1

A Fusion Protein Platform for Analyzing Tethered Agonism in the Adhesion Family of G

Protein-Coupled Receptors

Andrew N. Dates[1], Daniel T.D. Jones[1], Meredith A. Skiba[1], Maggie M. Burruss[1], Andrew C.

Kruse[1], and Stephen C. Blacklow[1,2,*]

[1]Department of Biological Chemistry and Molecular Pharmacology, Blavatnik Institute, Harvard

Medical School, Boston, MA 02115, USA

[2]Department of Cancer Biology, Dana Farber Cancer Institute, Boston, MA 02215, USA

*Correspondence: stephen_blacklow@hms.harvard.edu

Abstract

Adhesion G Protein-Coupled Receptors (aGPCRs) are a distinct family of transmembrane proteins that carry out numerous cellular functions in multicellular organisms. During maturation, aGPCRs undergo autoproteolysis to generate non-covalently associated N-terminal and C-terminal fragments (NTF and CTF, respectively). Signaling is induced when a tethered agonist at the N-terminus of the CTF is released from or unmasked by the NTF to access its binding site within the seven transmembrane domain of the CTF. Here, we show that the tethered sequence does not require a free N-terminus to signal and exploit this finding to create a fusion protein platform for interrogation of structure-function relationships in aGPCRs. We analyze tethered agonism in CD97, a protein implicated in immune cell function and cancer cell invasiveness, and demonstrate the platform's generality by creating constitutively active forms of several other aGPCRs. This platform should find wide applicability in analyzing signals transmitted by tethered agonists in aGPCRs.

Introduction

As the largest class of membrane proteins in the human genome, G protein-coupled receptors (GPCRs) are central modulators of physiology.[1] The 33 Adhesion G Protein-Coupled Receptors (aGPCRs) comprise the second-largest family of GPCRs (behind the rhodopsin family) and influence development, cell maturation, and tissue maintenance through roles in myelination, synapse formation, immune cell migration, auditory perception, and other critical processes.[2–9] Despite their abundance in the human genome, aGPCRs are relatively understudied compared to other GPCR classes,[10] and not currently targeted by therapeutics.

The aGPCRs have a unique modular domain organization, with three functional units (Fig. 1). The adhesion module typically consists of a series of protein interaction domains responsible for ligand binding.[11] Between the adhesion module and the membrane is a regulatory region for signaling, called the GPCR Autoproteolysis-Inducing (GAIN) domain. The third module consists of the seven-transmembrane (7TM) GPCR domain, which couples to intracellular effectors to execute biological outputs. A distinctive feature of the GAIN domain is that it undergoes autocatalytic cleavage during maturation. This auto-processing event divides the aGPCR into two subunits: an N-terminal fragment (NTF) containing the adhesion module and most of the GAIN domain and a C-terminal fragment (CTF) containing a highly conserved residual "stalk" from the GAIN domain followed by the 7TM domain and cytoplasmic tail.[12,13] The stalk is comprised of ~15-25 residues, with the first seven amino acids being highly conserved across family members and largely hydrophobic in nature. The first seven stalk residues are a sequestered β-strand in the GAIN domain and this sequence functions as a tethered agonist (TA) upon removal of the NTF, which has been postulated to be triggered upon the delivery of force by bound ligand to the adhesion module.[14–22]

An atomic-level view of tethered agonism in aGPCRs has recently emerged from cryo-electron microscopic (cryo-EM) studies of seven CTFs from different aGPCRs.[23–26] These studies define a binding pose of the TA in the orthosteric site and the conformational changes imposed on the GPCR to facilitate transition to the active, G protein-bound state. The TA adopts a conserved U-shaped conformation when buried within the orthosteric pocket and is connected to the first transmembrane domain (TM1) by a poorly conserved linker which comprises the remainder of the stalk.

Recombinant CTF proteins expressed with an added N-terminal methionine residue (M-

CTF) have been used to evaluate tethered agonism and the effects of specific TA residues on

aGPCR activity.[14] However, the M-CTF approach for use in comparing the signaling activity of

different mutants of a given receptor can be challenging, in part because the dynamic range of

the M-CTF-induced signal is variable among receptors, and in part because M-CTF constructs

lack N-terminal signal sequences, and as a result the amount of protein expressed (and the extent

to which the N-terminal methionine is retained) can vary among the mutants being tested.[14,16,20]

One facet of aGPCR activation that remains unclear is whether liberation of a native, free

N-terminus is strictly required for tethered agonism, or whether intramolecular agonism can

occur when the TA is preceded by protein sequences or domains. The recent cryo-EM structures

of aGPCR CTFs suggest the possibility of accommodating an N-terminal extension.[23–26] The

addition of short polypeptide sequences to the N-terminal end of the TA however, has had

opposing effects on signaling depending on the identities of receptor and fusion sequence tested,

suggesting that there may be variation in the capacity of individual receptors to accommodate an

N-terminal extension. [15,16,22,27,28] The divergence in these observations might be reconciled with a

simple, modular, and generalizable assay system to evaluate structure-function relationships in

aGPCRs.

Here, we assessed whether a free N-terminus is required for tethered agonism using

CD97 (ADGRE5) as a biologically important, prototypical example of an aGPCR.[29,30] CD97 is a

critical regulator of dendritic cell homeostasis in the spleen, where it senses its natural ligand

CD55 on red blood cells;[31–35] it is also highly expressed in many carcinomas, localizing to the

invasion front of tumors, and mediating invasion and migration phenotypes in multiple cancer

cell lines.[36–39] In constructing fusion proteins that place an unmasked TA of CD97 between an

inert N-terminal domain and the 7TM domain of CD97, we found that these fusion proteins are

potent, autonomously active molecules that require an intact TA for their signaling activity. We

then used this fusion protein approach to investigate structure-function relationships in CD97 and

adapted this strategy to create a platform for the study of aGPCR activity with high signal-to-

noise across a variety of aGPCRs and coupled G proteins. This work provides a broadly

applicable platform for the evaluation of molecular mechanisms of aGPCR agonism and

autoinhibition as well as the identification of G protein coupling profiles for this class of GPCRs.

Results

Development of a robust assay for the signaling activity of the CD97 CTF. To establish the signaling activity of CD97 using a reference method, we used reporter gene assays,[40] comparing the basal activity of full-length CD97 (CD97 FL) with that of the CD97 M-CTF and that of the CD97 M-CTF with the stalk sequence deleted (designated M-SL, for "stalkless"). Each of these proteins was expressed with a C-terminal FLAG epitope tag to facilitate detection by Western blot or flow cytometry using an anti-FLAG antibody (Fig. 1). Assays were performed using a serum response element (SRE) reporter, a cAMP Response Element (CRE) reporter, a Nuclear Factor of Activated T Cells Response Element (NFAT-RE) reporter, and a Serum Response Factor Response Element (SRF-RE) reporter (Extended Data Fig. 1); each reporter was validated using ligand-induced stimulation of the human β2 adrenergic (β2AR) and muscarinic M3 receptors (M3R) as controls (Extended Data Fig. 2a-d). When CD97 FL is expressed and tested in the reporter assay, it results in low basal activity with the SRE and SRF-RE reporters (Fig. 2 a,b). In contrast, expression of the CD97 M-CTF gives rise to constitutive activity with the SRE and SRF-RE reporters (Fig. 2 a,b), and no detectable difference with the NFAT-RE and CRE reporters (Fig. 2 c,d). The M-SL protein, which lacks the stalk sequence, shows negligible basal signaling activity with all reporters, confirming that the activity of M-CTF in the SRE and SRF-RE reporter assays depends on the presence of the TA. Because the steady-state amount of the M-SL protein was lower than that of its M-CTF counterpart (Extended Data Fig. 2 e,f), the amount of transfected DNA was titrated to achieve similar total protein amounts as judged by Western blot (Extended Data Fig. 2 g), and a dose-response analysis for the two constructs was also performed in the reporter assay (Extended Data Fig. 2 h). The results show that the M-SL signaling activity remains silent by comparison to M-CTF across a range of comparable protein amounts. These data are in agreement with previous experiments using the SRE assay and establish that CD97 also exhibits a stalk-dependent signal in the SRF-RE assay.[41,42]

We next sought to assemble a quiescent form of the CTF in which signaling could be induced by proteolysis, conceptually analogous to the PAR1 fusion of ADGRL3 that can be switched on with thrombin cleavage.[16] In our design, we created a fusion protein in which the tobacco etch virus protease (TEV) cleavage sequence directly links the C-terminus of an inert, thermostable protein domain to the N-terminus of the CD97 CTF. For initial studies, we chose maltose-binding protein (MBP) as a fusion headpiece, creating a construct we refer to as MBP-

CTF (Fig. 1). This design positions the N-terminal serine of the CD97 stalk sequence as the P1'

residue of the TEV cleavage site, enabling the removal of the fusion sequence to generate an

intact CTF identical to that produced by autocatalytic cleavage of the full-length protein *in vivo*.

We expressed this construct behind the IL2 signal sequence to facilitate consistent expression

and surface delivery among different CD97 variants. We also included an HA epitope tag at the

N-terminal end for easy cell-surface detection by flow cytometry and western blot.

Remarkably, we found that the MBP-CTF fusion protein and M-CTF possess similar

maximal constitutive signaling activity in the SRE assay, and that the DNA encoding the MBP-

CTF fusion was ~30-fold more potent than M-CTF, even *without* addition of TEV to catalyze

release of the N-terminal fusion domain (Fig. 3 a). As judged by Western blot, the MBP-CTF

fusion was produced in greater amounts than the M-CTF when equivalent quantities of DNA

were transfected, but this difference in abundance does not fully account for its increased

potency (Fig. 3 b). When the stalk sequence was deleted, the robust constitutive signaling

activity was abrogated, confirming the dependence of the reporter assay signal on the presence of

the stalk sequence in the fusion protein (Extended Data Fig. 3 a). Flow cytometry also showed

that the surface abundance of the MBP-CTF protein was much lower than that of CD97 FL or

the MBP-SL fusion, even though the total amount of MBP-CTF was comparable to that of the

two other proteins (Extended Data Fig. 3 b,c), suggesting that the active species of CD97 is

rapidly internalized in these cells (or conceivably is signaling from an intracellular

compartment[43,44]).

We tested whether a range of different N-terminal fusions resulted in comparable

constitutive signaling by the CD97 CTF. Indeed, sequences ranging from thermostable protein

domains such as T4-lysozyme and BRIL to entities as short as a 3xFLAG epitope sequence all

produce robust signals when tested in the SRE reporter assay (Fig. 3 c, Extended Data Fig. 4 a).

We then co-expressed the MBP-CTF with a TEV protease variant optimized for activity within

the secretory pathway to determine whether cleavage to generate the CTF would increase the

reporter signal.[45] At all amounts of transfected TEV-expressing plasmid tested, the MBP-CTF

fusion was efficiently cleaved to populate a ~34 kDa band corresponding to the native CTF. The

reporter assay signal remained comparable to the uncleaved MBP-CTF after cleavage and did

not increase after MBP removal, suggesting that the MBP fusion does not influence GPCR

signaling by the TA (Extended Data Fig. 4 b,c). The intrinsic signaling activity of our N-terminal

fusion proteins nicely complements recent structural studies, which show that the TA sequence

in other aGPCRs occupies the orthosteric site of the 7TM domain and appears to have an exit

vector that will accommodate additional N-terminal sequences extending into the extracellular

space.[23–26]

CD97 couples selectively to G12/13 heterotrimers. Reporter assay signals are a signaling readout

downstream of heterotrimeric G protein activity and do not unambiguously identify the G protein

relied upon for the signal (Extended Data Fig. 1). We used our CD97 MBP-CTF fusion protein

to define the G protein coupling profile of CD97 more precisely by testing its activity inducing

the SRE reporter in an engineered HEK293T cell line lacking six $G\alpha$ subunits: $G\alpha_s$ short, $G\alpha_s$

long, $G\alpha_q$, $G\alpha_{11}$, $G\alpha_{12}$, and $G\alpha_{13}$, referred to as HEKΔ6.[46] After dose optimization for each of the

$G\alpha$ subunits and control studies with a receptor of known coupling profile (Extended Data Fig.

5),[47,48] we then transfected these cells with the CD97 MBP-CTF, the SRE reporter, and each of a

series of six $G\alpha$ proteins, and recorded the signaling activity of the CD97 MBP-CTF associated

with each co-transfected $G\alpha$ subunit. Signaling occurred only when $G\alpha_{12}$ or $G\alpha_{13}$ were co-

transfected with the CD97 MBP-CTF in the HEKΔ6 cells (Fig. 4), giving a CD97 coupling

profile in agreement with other recent studies.[31,41] These data also indicate that MBP-CTF

molecules can be used generally to identify $G\alpha$ coupling profiles of receptors for which the

coupling partners are not known.

Using the MBP-CTF platform to probe structure-function relationships in tethered agonism and

TA masking. The TA lies laterally across the orthosteric pocket in recent cryo-EM structures of

other CTFs, [23–26] features which are preserved in a homology model of the CD97 CTF (Fig. 5

a,b). To probe structure-function relationships in the stalk of CD97, we systematically mutated

stalk residues in the CD97 MBP-CTF context and evaluated the effects on activity, mutating

residues in positions P1'-9' (531-539) individually to either alanine or lysine. For residues that

are alanine in the wild-type protein, we substituted glycine for alanine in the mutational scan

instead. Only F533[P3'] and L536[P6'] were intolerant of either substitution in the activity assay,

suggestive that both the size of the amino acid and hydrophobicity are necessary for activation at

these positions (Fig. 5 c). The M537A[P7'] mutation exhibited activity like that of the wild-type

protein, as did the A534G[P4'] and A538G[P8'] substitutions; however, signaling was abrogated by

the M537K$^{P7'}$, A534K$^{P4'}$, and A538K$^{P8'}$ mutations, indicating that either a small or non-polar

side chain is required at each of these positions. S531$^{P1'}$, S532$^{P2'}$, and H539$^{P9'}$ each tolerated

substitution to either alanine or lysine, as did I535$^{P5'}$, a highly conserved residue in the center of

the TA, which remarkably retained ~40% of wild-type activity even when mutated to lysine. We

also tested distal stalk residues in positions P10'-15' in triple alanine mutants and saw no effects

on activity (Extended Data Fig. 6 a). Note that the amount of mutant protein expressed was

comparable to that of the wild-type protein for all mutations tested except for F533A$^{P3'}$ and

L536A$^{P6'}$, where the steady state protein amount was lower (Extended Data Figure 6 b,c). The

observed effects of mutations to the stalk sequence align nicely with the homology model of the

CD97 CTF, in that buried hydrophobic residues are indispensable for agonism whereas residues

less important for agonism are tolerant of alanine/glycine mutation and tolerant of lysine

substitutions only when the position is exposed to solvent (Fig. 5 d).

202   We also determined the length of flexible tether needed between MBP and the TA to

permit constitutive agonism. All proteins tested in this experiment were present in similar

amounts, as judged by Western blot (Extended Data Fig. 7 a). Excision of the TEV cleavage

sequence and S531$^{P1'}$ had no effect on the activity of the MBP-fused CTF, whereas the

additional removal of S532$^{P2'}$ resulted in a reduction in activity to that of the full-length protein

with the TA sequestered by the GAIN domain (Fig. 6 a–c). This decrease in activity resulted

from steric interference by the MBP fusion, because reinsertion of the TEV cleavage site

between MBP and F533$^{P3'}$ restored activity to that of the parental MBP-TEV-CTF protein (Fig. 6

b,c).

211   In addition, we used our MBP-fusion strategy to determine the minimal portion of the

GAIN domain sufficient for TA sequestration in CD97. We inserted a series of increasing-length

segments of the GAIN domain between the TA and the TEV cleavage site of the MBP-CTF

fusion protein (Extended Data Fig. 7 b). Only the full-length GAIN insertion spontaneously

underwent autoproteolysis upon synthesis (Extended Data Fig. 7 c), and only this sequence

restrained the constitutive signaling activity of the MBP-CTF molecule (Fig. 6 d,e), showing that

a natively folded GAIN domain is the minimal requirement both for aGPCR processing and for

autoinhibition of TA-mediated aGPCR signaling.

The MBP-CTF fusion platform is a generalizable strategy for robust analysis of aGPCR

signaling. Lastly, we evaluated the generalizability of the MBP-fusion platform as a tool to study

aGPCR signaling properties with a panel of diverse aGPCRs. First, we compared FL, M-CTF,

and MBP-CTF forms of GPR56 (ADGRG1), an aGPCR which signals through $G_{12/13}$ and which

is evolutionarily and structurally distant from CD97 (Fig. 7 a,b) [14,23]. When 30 ng of plasmid

DNA expressing GPR56 was used, the MBP-CTF protein strongly stimulated the SRE and SRF-

RE reporters and modestly activated the NFAT-RE reporter, whereas neither the M-CTF protein

nor the FL protein produced a reporter signal substantially above background (Fig. 7 c–f,

Extended Data Fig. 8 a). Increasing the dose of transfected DNA for the M-CTF protein did

induce a reporter signal (Extended data Fig. 8 b). The amount of M-CTF protein produced was

also much lower than that of the MBP-CTF even at higher DNA doses, again highlighting the

advantage of using the MBP-CTF platform for reliable protein expression (Extended Data Fig. 8

c).

We curated a panel of additional MBP-CTFs corresponding to eight previously studied

aGPCRs (Fig. 7 g).[14,15,17,22,49–51] Our platform replicated cAMP-stimulation (as CRE reporter

activity) by ADGRF1, ADGRG2, ADGRG5, ADGRG6, and ADGRD1 and showed improved

signal amplitude by as much as 50-fold above previous analyses. We also confirmed stimulation

of the SRE reporter by ADGRF1, ADGRL3, ADGRG3, and ADGRV1. Every receptor construct

tested was reliably expressed at the amount of DNA used for transfection in our reporter assays

(Extended Data Fig. 8 c). ADGRL3 and ADGRD1, both of which were inactivated by fusion to

the PAR1 N-terminus, were highly active in at least one of the two reporters. In addition, our

scan uncovered three novel signals which have not been reported to date: induction of the CRE

by ADGRG3 and induction of the SRE by ADGRG5 and ADGRD1. These findings highlight the

utility of our platform to identify the downstream pathways accessible to individual receptors,

and demonstrate its general utility as a modular assay system to probe the activity and coupling

biology of diverse receptors in the aGPCR family.

<u>Discussion</u>

Here, we report the development of a modular and highly robust platform for evaluating

the signaling profiles of aGPCR family proteins. In our system, we fused MBP with a linker

containing a TEV cleavage site to the TA sequence at the amino terminal end of the aGPCR

signaling subunit (CTF). Engineered receptors with this domain organization are highly potent

and autonomously active in reporter gene assays. The signal rapidly reaches saturation, which is

maintained over a wide dose range, buffering the effects of small variations in protein abundance

when comparing mutants. This approach also provides advantages over the use of synthetic

peptides to activate 7TM domains *in trans* because it is more cost-effective and is insensitive to

variation in the solubility of the peptide sequence, which can confound interpretation of results

with synthetic peptide addition.[52] The signaling platform has a simple, modular design, is easy to

use, is amenable to implementation in engineered cells for determining G protein subtype

specificity, and both nicely complements and can be used in conjunction with other methods for

probing aGPCR activity, such as [^{35}S]GTPγS binding assays and proximity based recruitment

assays.[14,16]

We demonstrated the utility of our signaling platform by probing structure-function

relationships with the CD97 MBP-CTF. We defined its G protein subtype-specificity, probed the

functional roles of individual stalk residues, identified the closest allowable N-terminal fusion to

the TA, and determined the minimal sequence derived from the GAIN domain sufficient for

autoproteolysis and autoinhibition. We also showed that the structural requirements for

autoinhibition are more rigorous than simply blocking the N-terminal end of the stalk peptide

with a preceding polypeptide sequence. While other fusion constructs can inhibit autonomous

CTF activity,[16,27,28] our findings suggest that there are features of these fusion sequences which

interfere with autonomous activation, likely through some form of TA sequestration. In line with

this logic, we showed that, for the case of the GAIN domain, only an intact domain sequence

possesses such properties. These conclusions support the notion that two discrete states of the

TA exist, both of which involve the deep sequestration of its hydrophobic sequence within

another protein domain.

Across the wide variety of mutational variants tested in our studies of CD97, our platform

generated consistent amounts of expressed protein except for the F533A$^{P3'}$ and L536A$^{P6'}$ loss-of-

function mutations, which showed a modest decrease in total protein amounts. Mutation of these

residues to lysine had no detectable effect on protein expression (Extended Data Fig. 6 a,b). We

speculate that disabling nonpolar mutations expose the TA as a hydrophobic neoepitope, which

targets the protein for degradation by quality control machinery in the secretory pathway,

whereas substitution of a charged residue such as lysine is not detected as an exposed

hydrophobic surface and thus avoids a degradative fate.

We also showed that our platform should be applicable to signaling across other proteins

in the aGPCR family by testing representative examples across six aGPCR subfamilies. In doing

so, we uncovered three coupling activities not previously reported, indicating the utility of our

platform in identifying all accessible downstream pathways of a given receptor. The expression

advantages conferred by our system may prove particularly useful for studying aGPCRs that are

poorly expressed or less biochemically tractable in other assays. Of the 33 human aGPCRs,

several receptors do not possess canonical autoproteolytic cleavage sequences, raising the

question of whether they signal through intramolecular ligation of the TA.[53] Use of our fusion

protein platform to interrogate signaling of the CTF can determine whether aGPCRs that lack a

canonical autoproteolytic sequence are able to signal when the TA is unmasked.

Though our data show that TA unmasking is sufficient to activate aGPCRs, it is

important to recognize that our findings do not distinguish between existing models that favor

either complete dissociation of the TA from the rest of the GAIN domain or some form of de-

sequestration or unmasking of the TA as the key step in activation. Without a completely intact

GAIN domain, the default is a receptor molecule in an autonomously active state (Fig. 6 d). This

finding is also in line with previously reported clinical and *in vitro* observations that destabilizing

mutations of the GAIN domain lead to either autonomous activation or reduced force thresholds

of activation.[54,55] Additional data will be required to fully define how these disease-associated

mutations affect receptor function in molecular detail.

Methods

Materials

DMEM, PBS, and 0.25% trypsin were from Corning. Opti-MEM, 0.1mg/mL poly-D-lysine, and penicillin-streptomycin were from ThermoFisher Scientific. FBS was from GeminiBio. Lipofectamine 2000 was from Invitrogen. Carbachol was from Santa Cruz Biotechnology and (-)-isoproterenol hydrochloride was from Sigma-Aldrich. Rabbit anti-FLAG (D6W5B), Mouse anti-HA (6E2), Mouse anti-Beta Tubulin (D3U1W), Rabbit anti-GAPDH (14C10), and HRP-conjugated Rabbit anti-GAPDH (D16H11) XP antibodies were from Cell Signaling Technology. IRDye 800-conjugated Donkey Anti-Rabbit and IRDye 680-conjugated Goat Anti-Mouse secondary antibodies were from LI-COR Biosciences. To generate 647-conjugated Mouse anti-HA antibody, the antibody was purified from hybridoma supernatant and chemically conjugated to Alexa Fluor 647 NHS Ester from ThermoFisher Scientific. The Dual-Luciferase Reporter Assay System kit was from Promega.

Plasmid DNA Constructs

All proteins were assembled using human aGPCR and G protein sequences retrieved from the Uniprot database. Synthetic gene fragments (Integrated DNA Technologies) were inserted into vectors using Gibson assembly, and plasmid sequences were confirmed by DNA sequencing. Receptor constructs with a 5' IL2-signal sequence were subcloned into the pFuse-hIgG1-Fc2 vector using EcoRI and BglII restriction sites. M-CTF constructs (which do not contain an IL2-signal sequence) were inserted into the pFuse-hIgG1-Fc1 vector using EcoRI and BglII restriction sites. All receptor constructs included a 3' stop codon preceding the hIgG1-Fc coding sequence. An "empty vector" construct (referred to as "EV"; see Table S1) was generated by placement of a stop codon immediately after the IL2-signal sequence of the pFuse-hIgG1-Fc2 vector. A single glycine-serine linker was inserted between any epitope tag and the coding sequence of the protein of interest. Detailed construct features and calculated molecular weights for proteins used in this study are listed in Table S1.

Gα proteins were encoded using synthetic DNA sequences and subcloned into pcDNA3.1 using HindIII and NotI restriction sites. The secTEV construct was subcloned into pcDNA3.1 with HindIII and XhoI restriction sites. Point mutants were generated using a Q5 Site-Directed Mutagenesis Kit (New England Biolabs).

We generated luciferase reporter plasmids containing individual response elements, a minimal promoter (minP), and *luc2* firefly luciferase. These plasmids are analogous to the pGL4.29[CRE/minP/*luc2P*], pGL4.30[NFAT-RE/minP/*luc2P*], pGL4.33[SRE/*luc2P*], pGL4.34[SRF-RE/*luc2P*] plasmids with the exceptions that *luc2* is used in place of *luc2P* in our system and that minP is included in our SRE and SRF-RE plasmids. To assemble these plasmids, synthetic gene fragments containing response elements were inserted into a pGL4.23[*luc2*/minP] vector obtained from Promega using KpnI and HindIII (for SRE, SRF-RE, and CRE) or KpnI and NcoI (for NFAT-RE, which included the minP in the synthetic gene fragment) restriction sites. The pRL-TK plasmid (Promega) was used to express *Renilla* luciferase.

Cell Culture

Parental HEK293T and HEKΔ6[46] cells were cultured in DMEM supplemented with 10% FBS
and 1% penicillin-streptomycin at 37°C in a 5% CO_2, humidified incubator.

Reporter Gene Assays

Cells were seeded onto 96-well clear bottom, white walled, tissue culture-treated culture plates at
a density of 3 x 10^5 per well in 100 μL complete media (DMEM/10% FBS/1% penicillin-
streptomycin) and incubated overnight. Prior to seeding, plates were treated with 40 μL of 10
μg/mL aqueous poly-D-lysine solution for 30 minutes at room temperature. 16-20 hours after
seeding, media was replaced with 50 μL complete media and cells were transfected by adding a
solution volume of 20 μL containing plasmid DNA and Lipofectamine 2000 at a ratio of 3 μL
per 1 μg DNA in Opti-MEM.

For standard single-DNA dose assays in the CRE and NFAT-RE systems, each well was co-
transfected with 50 ng firefly reporter plasmid, 1 ng pRL-TK plasmid, and 10 ng receptor
plasmid. For standard single-DNA dose assays in the SRE and SRF-RE systems, each well was
co-transfected with 30 ng firefly reporter plasmid, 0.6 ng pRL-TK plasmid, and 30 ng receptor
plasmid. For receptor DNA titration experiments (Fig. 3 a, Extended Data Fig. 1 g,h, Extended
Data Fig. 3 a), total DNA was balanced with EV (see Table S1). For secTEV co-transfection
experiments (Fig. 3 c), a constant receptor DNA load of 30 ng was used and total DNA was
balanced with pcDNA3.1.

After 4-6 hours, the transfection media was replaced with 100 μL complete media and incubated
overnight. After 16-20 hours, media was replaced with 100 μL complete media (CRE and
NFAT-RE) or serum-free DMEM supplemented with 1% penicillin-streptomycin (SRE and
SRF-RE) and cells were incubated for 6-8 hours. Cells were then washed with 100 μL PBS and
lysed by addition of 20 μL 1 x Passive Lysis Buffer (Promega) to each well and nutated for 15
minutes at room temperature. Lysates were pipetted up and down, and the assay plate was spun
at 400 x g for 60 seconds to remove air bubbles. The ratio of firefly to *renilla* luciferase activity
was measured using automated reagent injection with a GloMax luminometer system (Promega),
in which 50 μL LAR II and 25 μL Stop & Glo reagent (Promega) were added to each well.

For Gα rescue optimization experiments in HEKΔ6 cells (Extended Data Fig. 5 a,d,e), varying
amounts of Gα subunits were transfected alongside 50 ng firefly reporter plasmid and 1 ng pRL-
TK plasmid (CRE, NFAT-RE) or 30 ng firefly reporter plasmid and 0.6 ng pRL-TK plasmid
(SRE), and total DNA was balanced with pcDNA3.1. For Gα rescue panels in the SRE assay
(Fig. 4 a, Extended Data Fig. 5 c), 30 ng receptor DNA was co-transfected with respective Gα
DNA loads (20 ng of $Gα_s$, $Gα_q$, $Gα_{11}$, and $Gα_{16}$; 0.3 ng of $Gα_{12}$ and $Gα_{13}$), 30 ng firefly reporter
plasmid, and 0.6 ng pRL-TK plasmid, and total DNA was balanced with pcDNA3.1. For $Gα_{12/13}$
titration experiments alongside CD97 receptor constructs (Fig. 4 b,c), total DNA was balanced
with pcDNA3.1.

Bioinformatics and Homology Model Generation

Sequence conservation of tethered agonist residues (Fig. 5 b) was derived from BLOSUM62
scores based upon a multiple sequence alignment beginning at the predicted autoproteolytic

392 cleavage sites of the 33 human aGPCR CTFs. For receptors without canonical cleavage sites, the
393 P1'-equivalent residue was determined using a multiple sequence alignment of the 32 human
394 GAIN domains. As ADGRA1 possesses neither a predicted cleavage motif nor GAIN domain,
395 the P1' residue was determined by alignment of its full-length sequence with the CTFs of the
396 other 32 human aGPCRs.
397
398 To generate a homology model of the CD97 CTF in the active signaling state (Fig. 5 a,d), the
399 sequences of human CD97 residues 531-835 and human $G\alpha_{13}$ residues 1-377 were submitted to
400 the ColabFold software server,[56] which predicted a structure of the TA-agonized CTF in complex
401 with $G\alpha_{13}$. The model of the CD97 GAIN domain (Extended Data Fig. 7 b) was extracted from
402 the alphafold2 structure prediction for full-length human CD97.[57,58]
403
Western Blot Analysis

Cells for western blots were prepared in poly-D-lysine-treated 24-well plates in parallel with
cells for reporter assays, using the same transfection mixtures (which included both reporter
plasmids and receptor and/or secTEV plasmids), media change timelines, and time of harvest.
Cell density, media volumes, and DNA loads were increased five-fold relative to the amounts
used in 96-well format reporter assays to account for the larger well size. At the time of harvest,
protein samples for blots were reduced and denatured in Laemmli buffer, loaded into 4 –20%
Tris-Glycine gels (Bio-Rad) for SDS-PAGE, and transferred onto nitrocellulose membranes
(Cytiva). Blots were blocked with 5% milk in TBST and incubated with primary antibodies
overnight at 4°C. Simultaneous analysis of proteins of interest and loading controls were carried
out for each blot (with the exception of those shown in Extended Data Fig. 2 e,f) through the use
of two imaging channels on a LI-COR Odyssey CLx Fluorescent Imaging System. Rabbit anti-
GAPDH antibody was used as a loading control for blots using Mouse anti-HA antibody, and
Mouse anti-Beta Tubulin antibody was used as a loading control for blots using Rabbit anti-
FLAG antibody. Blots were incubated with secondary antibodies for 30-60 minutes prior to
imaging. For blots simultaneously detecting FLAG, HA, and GAPDH (Extended Data Fig. 7 c,
Extended Data Fig. 8 b), FLAG and HA were detected using the LI-COR system and HRP-
conjugated GAPDH primary antibody was detected with Western Lightning Plus-ECL
(PerkinElmer) using a Bio-Rad Chemidoc System.
423
Cell Surface Staining and Flow Cytometry

Quantification of cell surface staining by flow cytometry was performed using cells from non-
coated 24-well plates prepared in parallel with cells for reporter assays, using the same
transfection mixtures (which included both reporter plasmids and receptor), media change
timelines, and time of harvest. Cell density, media volumes, and DNA loads were increased five-
fold relative to the amounts used in 96-well format reporter assays to account for the larger well
size. At the time of harvest, cells were resuspended in ice-cold 20 mM HEPES buffer, pH 7.4,
containing 150 mM NaCl and 0.1% bovine serum albumin (HBS/BSA), transferred to a 96-well
V-bottom plate (Corning), and washed once with HBS/BSA. Cells were incubated at 4°C with
647-conjugated mouse anti-HA antibody at 1 μg/mL in HBS/BSA for 20 minutes, washed twice
with HBS/BSA, and analyzed using a Beckman Coulter Cytoflex flow cytometer. Data were
analyzed with CytExpert software. Single cells were gated using forward and side scattering and
staining was quantified using the mean fluorescence intensity in the APC channel.

References

1. Schiöth HB, Lagerström MC. Structural diversity of G protein-coupled receptors and significance for drug discovery. *Nature Reviews Drug Discovery*. 2008;7(4):339-357. doi:10.1038/nrd2518

2. Ushkaryov YA, Lelianova V, Vysokov N v. Catching Latrophilin With Lasso: A Universal Mechanism for Axonal Attraction and Synapse Formation. *Frontiers in Neuroscience*. 2019;13. doi:10.3389/fnins.2019.00257

3. Arcos-Burgos M, Jain M, Acosta MT, et al. A common variant of the latrophilin 3 gene, LPHN3, confers susceptibility to ADHD and predicts effectiveness of stimulant medication. *Molecular Psychiatry*. 2010;15(11):1053-1066. doi:10.1038/mp.2010.6

4. Yona S, Lin H, Dri P, et al. Ligation of the adhesion-GPCR EMR2 regulates human neutrophil function. *The FASEB Journal*. 2008;22(3):741-751. doi:10.1096/fj.07-9435com

5. Bolliger MF, Martinelli DC, Südhof TC. The cell-adhesion G protein-coupled receptor BAI3 is a high-affinity receptor for C1q-like proteins. *Proc Natl Acad Sci U S A*. 2011;108(6):2534-2539. doi:10.1073/pnas.1019577108

6. Langenhan T, Piao X, Monk KR. Adhesion G protein-coupled receptors in nervous system development and disease. *Nature Reviews Neuroscience*. 2016;17(9):550-561. doi:10.1038/nrn.2016.86

7. McGee JA, Goodyear RJ, McMillan DR, et al. The very large G-protein-coupled receptor VLGR1: A component of the ankle link complex required for the normal development of auditory hair bundles. *Journal of Neuroscience*. 2006;26(24):6543-6553. doi:10.1523/JNEUROSCI.0693-06.2006

8. Monk KR, Oshima K, Jörs S, Heller S, Talbot WS. Gpr126 is essential for peripheral nerve development and myelination in mammals. *Development*. 2011;138(13):2673-2680. doi:10.1242/dev.062224

9. Piao X, Hill SS, Bodell A, et al. G protein-coupled receptor-dependent development of human frontal cortex. *Science*. 2004;303(5666):2033-2036. doi:10.1126/SCIENCE.1092780

10. Fredriksson R, Lagerström MC, Lundin LG, Schiöth HB. The G-protein-coupled receptors in the human genome form five main families. Phylogenetic analysis, paralogon groups, and fingerprints. *Mol Pharmacol*. 2003;63(6):1256-1272. doi:10.1124/MOL.63.6.1256

11. Purcell RH, Hall RA. Adhesion G Protein-Coupled Receptors as Drug Targets. *Annu Rev Pharmacol Toxicol*. 2018;58:429-449. doi:10.1146/ANNUREV-PHARMTOX-010617-052933

12. Araç D, Boucard AA, Bolliger MF, et al. A novel evolutionarily conserved domain of cell-adhesion GPCRs mediates autoproteolysis. *EMBO Journal*. 2012;31(6):1364-1378. doi:10.1038/emboj.2012.26

475 13. Prömel S, Langenhan T, Araç D. Matching structure with function: The GAIN domain of
476   Adhesion-GPCR and PKD1-like proteins. *Trends in Pharmacological Sciences*.
477   2013;34(8):470-478. doi:10.1016/j.tips.2013.06.002

478 14. Stoveken HM, Hajduczok AG, Xu L, Tall GG. Adhesion G protein-coupled receptors are
479   activated by exposure of a cryptic tethered agonist. *Proc Natl Acad Sci U S A*.
480   2015;112(19):6194-6199. doi:10.1073/pnas.1421785112

481 15. Liebscher I, Schön J, Petersen SC, et al. A Tethered Agonist within the Ectodomain
482   Activates the Adhesion G Protein-Coupled Receptors GPR126 and GPR133. *Cell Reports*.
483   2014;9(6):2018-2026. doi:10.1016/j.celrep.2014.11.036

484 16. Mathiasen S, Palmisano T, Perry NA, et al. G12/13 is activated by acute tethered agonist
485   exposure in the adhesion GPCR ADGRL3. *Nature Chemical Biology*. 2020;16(12):1343-
486   1350. doi:10.1038/s41589-020-0617-7

487 17. Wilde C, Fischer L, Lede V, et al. The constitutive activity of the adhesion GPCR
488   GPR114/ADGRG5 is mediated by its tethered agonist. *FASEB Journal*. 2016;30(2):666-
489   673. doi:10.1096/fj.15-276220

490 18. Demberg LM, Winkler J, Wilde C, et al. Activation of adhesion G protein-coupled
491   receptors: Agonist specificity of Stachel sequence-derived peptides. *Journal of Biological*
492   *Chemistry*. 2017;292(11):4383-4394. doi:10.1074/jbc.M116.763656

493 19. Peeters MC, Fokkelman M, Boogaard B, et al. The adhesion G protein-coupled receptor
494   G2 (ADGRG2/GPR64) constitutively activates SRE and NFκB and is involved in cell
495   adhesion and migration. *Cellular Signalling*. 2015;27(12):2579-2588.
496   doi:10.1016/j.cellsig.2015.08.015

497 20. Kishore A, Purcell RH, Nassiri-Toosi Z, Hall RA. Stalk-dependent and stalk-independent
498   signaling by the adhesion G protein-coupled receptors GPR56 (ADGRG1) and BAI1
499   (ADGRB1). *Journal of Biological Chemistry*. 2016;291(7):3385-3394.
500   doi:10.1074/jbc.M115.689349

501 21. Vizurraga A, Adhikari R, Yeung J, Yu M, Tall GG. Mechanisms of adhesion G protein–
502   coupled receptor activation. *Journal of Biological Chemistry*. 2020;295(41):14065-14083.
503   doi:10.1074/jbc.REV120.007423

504 22. Demberg LM, Rothemund S, Schöneberg T, Liebscher I. Identification of the tethered
505   peptide agonist of the adhesion G protein-coupled receptor GPR64/ADGRG2. *Biochem*
506   *Biophys Res Commun*. 2015;464(3):743-747. doi:10.1016/J.BBRC.2015.07.020

507 23. Barros-Álvarez X, Nwokonko RM, Vizurraga A, et al. The tethered peptide activation
508   mechanism of adhesion GPCRs. *Nature 2022 604:7907*. 2022;604(7907):757-762.
509   doi:10.1038/s41586-022-04575-7

510 24. Xiao P, Guo S, Wen X, et al. Tethered peptide activation mechanism of the adhesion
511   GPCRs ADGRG2 and ADGRG4. *Nature 2022 604:7907*. 2022;604(7907):771-778.
512   doi:10.1038/s41586-022-04590-8

25.   Ping YQ, Xiao P, Yang F, et al. Structural basis for the tethered peptide activation of
adhesion GPCRs. *Nature 2022 604:7907*. 2022;604(7907):763-770. doi:10.1038/s41586-
022-04619-y

26.   Qu X, Qiu N, Wang M, et al. Structural basis of tethered agonism of the adhesion GPCRs
ADGRD1 and ADGRF1. *Nature 2022 604:7907*. 2022;604(7907):779-785.
doi:10.1038/s41586-022-04580-w

27.   Frenster JD, Stephan G, Ravn-Boess N, et al. Functional impact of intramolecular
cleavage and dissociation of adhesion G protein-coupled receptor GPR133 (ADGRD1) on
canonical signaling. *Journal of Biological Chemistry*. 2021;296.
doi:10.1016/j.jbc.2021.100798

28.   Lizano E, Hayes JL, Willard FS. A synthetic method to assay adhesion-family G-protein
coupled receptors. Determination of the G-protein coupling profile of
ADGRG6(GPR126). *Biochemical and Biophysical Research Communications*.
2021;534:317-322. doi:10.1016/j.bbrc.2020.11.086

29.   Gray JX, Haino M, Roth MJ, et al. *CD97 Is a Processed, Seven-Transmembrane,*
Heterodimeric Receptor Associated with Inflammation. Vol 157.; 1996.
http://www.jimmunol.org/

30.   EICHLER W, AUST G, HAMANN D. Characterization of an early activation-dependent
antigen on lymphocytes defined by the monoclonal antibody BL-Ac(F2). *Scand J*
Immunol. 1994;39(1):111-115. doi:10.1111/J.1365-3083.1994.TB03348.X

31.   Liu D, Duan L, Rodda LB, et al. CD97 promotes spleen dendritic cell homeostasis
through the mechanosensing of red blood cells. *Science (1979)*. 2022;375(6581).
doi:10.1126/science.abi5965

32.   Liu D, Duan L, Cyster JG. Chemo- and mechanosensing by dendritic cells facilitate
antigen surveillance in the spleen. *Immunol Rev*. 2022;306(1):25-42.
doi:10.1111/IMR.13055

33.   Karpus ON, Veninga H, Hoek RM, et al. Shear stress-dependent downregulation of the
adhesion-G protein-coupled receptor CD97 on circulating leukocytes upon contact with its
ligand CD55. *J Immunol*. 2013;190(7):3740-3748. doi:10.4049/JIMMUNOL.1202192

34.   Wang T, Tian L, Haino M, et al. Improved antibacterial host defense and altered
peripheral granulocyte homeostasis in mice lacking the adhesion class G protein receptor
CD97. *Infection and Immunity*. 2007;75(3):1144-1153. doi:10.1128/IAI.00869-06

35.   Hamann J, Vogel B, van Schijndel GMW, van Lier RAW. The seven-span transmembrane
receptor CD97 has a cellular ligand (CD55, DAF). *J Exp Med*. 1996;184(3):1185-1189.
doi:10.1084/JEM.184.3.1185

36.   Ward Y, Lake R, Faraji F, et al. Platelets Promote Metastasis via Binding Tumor CD97
Leading to Bidirectional Signaling that Coordinates Transendothelial Migration. *Cell*
Reports. 2018;23(3):808-822. doi:10.1016/j.celrep.2018.03.092

37. Safaee M, Clark AJ, Ivan ME, et al. CD97 is a multifunctional leukocyte receptor with
distinct roles in human cancers (Review). *International Journal of Oncology*.
2013;43(5):1343-1350. doi:10.3892/ijo.2013.2075

38. Tjong WY, Lin HH. The role of the RGD motif in CD97/ADGRE5-and EMR2/ADGRE2-
modulated tumor angiogenesis. *Biochemical and Biophysical Research Communications*.
2019;520(2):243-249. doi:10.1016/j.bbrc.2019.09.113

39. Aust G, Zheng L, Quaas M. To Detach, Migrate, Adhere, and Metastasize:
CD97/ADGRE5 in Cancer. *Cells 2022, Vol 11, Page 1538*. 2022;11(9):1538.
doi:10.3390/CELLS11091538

40. Cheng Z, Garvin D, Paguio A, Stecha P, Wood K, Fan F. Luciferase Reporter Assay
System for Deciphering GPCR Pathways. *Current Chemical Genomics*. 2010;4:84.
doi:10.2174/1875397301004010084

41. Ward Y, Lake R, Yin JJ, et al. LPA receptor heterodimerizes with CD97 to amplify LPA-
initiated RHO-dependent signaling and invasion in prostate cancer cells. *Cancer
Research*. 2011;71(23):7301-7311. doi:10.1158/0008-5472.CAN-11-2381

42. Hilbig D, Sittig D, Hoffmann F, et al. Mechano-Dependent Phosphorylation of the PDZ-
Binding Motif of CD97/ADGRE5 Modulates Cellular Detachment. *Cell Reports*.
2018;24(8):1986-1995. doi:10.1016/j.celrep.2018.07.071

43. Irannejad R, Tomshine JC, Tomshine JR, et al. Conformational biosensors reveal GPCR
signalling from endosomes. *Nature*. 2013;495(7442):534-538.
doi:10.1038/NATURE12000

44. Irannejad R, Pessino V, Mika D, et al. Functional selectivity of GPCR-directed drug
action through location bias. *Nat Chem Biol*. 2017;13(7):799-806.
doi:10.1038/NCHEMBIO.2389

45. Cesaratto F, López-Requena A, Burrone OR, Petris G. Engineered tobacco etch virus
(TEV) protease active in the secretory pathway of mammalian cells. *J Biotechnol*.
2015;212:159-166. doi:10.1016/J.JBIOTEC.2015.08.026

46. Grundmann M, Merten N, Malfacini D, et al. Lack of beta-arrestin signaling in the
absence of active G proteins. *Nature Communications 2018 9:1*. 2018;9(1):1-16.
doi:10.1038/s41467-017-02661-3

47. Inoue A, Raimondi F, Kadji FMN, et al. Illuminating G-Protein-Coupling Selectivity of
GPCRs. *Cell*. 2019;177(7):1933-1947.e25. doi:10.1016/J.CELL.2019.04.044

48. Avet C, Mancini A, Breton B, et al. Effector membrane translocation biosensors reveal G
protein and βarrestin coupling profiles of 100 therapeutically relevant GPCRs. *Elife*.
2022;11. doi:10.7554/ELIFE.74101

49. Hu QX, Dong JH, Du HB, et al. Constitutive Gαi Coupling Activity of Very Large G
Protein-coupled Receptor 1 (VLGR1) and Its Regulation by PDZD7 Protein. *The Journal
of Biological Chemistry*. 2014;289(35):24215. doi:10.1074/JBC.M114.549816

50.   Nazarko O, Kibrom A, Winkler J, et al. A Comprehensive Mutagenesis Screen of the
590   Adhesion GPCR Latrophilin-1/ADGRL1. *iScience*. 2018;3:264-278.
591   doi:10.1016/J.ISCI.2018.04.019

592   51.   Gupte J, Swaminath G, Danao J, Tian H, Li Y, Wu X. Signaling property study of
593   adhesion G-protein-coupled receptors. *FEBS Letters*. 2012;586(8):1214-1219.
594   doi:10.1016/j.febslet.2012.03.014

595   52.   Shoichet BK. Screening in a spirit haunted world. *Drug Discovery Today*. 2006;11(13-
596   14):607-615. doi:10.1016/J.DRUDIS.2006.05.014

597   53.   Langenhan T. Adhesion G protein–coupled receptors—Candidate metabotropic
598   mechanosensors and novel drug targets. *Basic & Clinical Pharmacology & Toxicology*.
599   2020;126(S6):5-16. doi:10.1111/BCPT.13223

600   54.   Boyden SE, Desai A, Cruse G, et al. Vibratory Urticaria Associated with a Missense
601   Variant in ADGRE2. *N Engl J Med*. 2016;374(7):656-663.
602   doi:10.1056/NEJMOA1500611

603   55.   Piao X, Hill SS, Bodell A, et al. G protein-coupled receptor-dependent development of
604   human frontal cortex. *Science*. 2004;303(5666):2033-2036.
605   doi:10.1126/SCIENCE.1092780

606   56.   Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. ColabFold:
607   making protein folding accessible to all. *Nature Methods 2022 19:6*. 2022;19(6):679-682.
608   doi:10.1038/s41592-022-01488-1

609   57.   Jumper J, Evans R, Pritzel A, et al. Highly accurate protein structure prediction with
610   AlphaFold. *Nature 2021 596:7873*. 2021;596(7873):583-589. doi:10.1038/s41586-021-
611   03819-2

612   58.   Varadi M, Anyango S, Deshpande M, et al. AlphaFold Protein Structure Database:
613   massively expanding the structural coverage of protein-sequence space with high-
614   accuracy models. *Nucleic Acids Research*. 2022;50(D1):D439-D444.
615   doi:10.1093/NAR/GKAB1061

616   59.   Treisman R, Alberts AS, Sahai E. Regulation of SRF activity by Rho family GTPases.
617   *Cold Spring Harb Symp Quant Biol*. 1998;63:643-651. doi:10.1101/SQB.1998.63.643

618   60.   Hill CS, Wynne J, Treisman R. The Rho family GTPases RhoA, Rac1, and CDC42Hs
619   regulate transcriptional activation by SRF. *Cell*. 1995;81(7):1159-1170.
620   doi:10.1016/S0092-8674(05)80020-0

Acknowledgments

This work was supported by a grant from the Warren Alpert foundation (to S.C.B.) and award 1 R35-CA220340 (to S.C.B). A.N.D. was supported by a fellowship from the Fujifilm Corporation. M.A.S. was supported by a Merck Postdoctoral Fellowship from the Helen Hay Whitney Foundation. We thank Asuka Inoue (Tohoku University) for generously providing the HEKΔ6 knockout cell line. We also thank members of the Blacklow and Kruse labs for helpful discussions.

Author Contributions

A.N.D. and S.C.B. conceived of the research. A.N.D., D.T.D.J., M.A.S, A.C.K, and S.C.B designed experiments. A.N.D. and M.M.B. performed experiments. A.N.D., D.T.D.J., M.A.S, A.C.K., and S.C.B. interpreted data. A.N.D. and S.C.B. wrote the manuscript; D.T.D.J., M.A.S, A.C.K., and S.C.B. edited the manuscript.

Competing Interests

S.C.B. is on the scientific advisory board for and receives funding from Erasca, Inc., is an advisor to MPM Capital, and is a consultant for Scorpion Therapeutics, Odyssey Therapeutics, Droia Ventures, and Ayala Pharmaceuticals for unrelated projects. A.C.K. is a co-founder and consultant for Tectonic Therapeutic and Seismic Therapeutic and for the Institute for Protein Innovation, a non-profit research institute.

Figure 1 – Overview of CD97 constructs used in this study. (a) Schematic diagrams of
constructs. Full-length CD97 (CD97 FL) protein and Maltose-Binding Protein (MBP)-fused C-
terminal fragment (CTF) and "Stalkless" (SL) molecules include the interleukin 2 signal
sequence (IL2ss) in place of the native signal peptide. Methionine-capped C-terminal fragment
(M-CTF) and stalkless (M-SL) molecules have no signal sequence. In the MBP-fusion proteins,
a tobacco etch virus protease (TEV) cleavage sequence is located between the C-terminus of
MBP and the N-terminus of the CTF and SL molecules, utilizing serine 531 as its P1' residue.
(b) Schematic depicting the domain architecture of constructs in **(a)**. CD97 FL is a heterodimer
at the cell surface comprised of N-terminal and C-terminal fragments (NTF, CTF). CD97
contains five tandem EGF repeat domains followed by a GPCR autoproteolysis-inducing
(GAIN) domain, a 7TM domain, and a cytoplasmic tail. Autocatalytic cleavage occurs within the
GAIN domain to create the NTF and CTF. The stalk peptide is at the N-terminus of the CTF and
is sequestered within the GAIN domain.

Figure 2 – CD97 signaling assays using SRE, SRF-RE, NFAT-RE, and CRE reporter genes.
Signaling response of the indicated CD97-derived molecules in SRE **(a)**, SRF-RE **(b)**, NFAT-RE
(c), and CRE **(d)** reporter assays. Cells were transfected with 30 ng receptor DNA/well **(a,b)** or
10 ng receptor DNA/well **(c,d)**. Data are normalized to the mean value for empty vector. In all
panels, data are presented as mean ± s.d. of three technical replicates and representative of three
independent biological replicates. One-way analysis of variance (ANOVA) was used with
Tukey's multiple-comparison post-hoc test to compare the difference between samples
(****P<0.0001).

Figure 3 – Dose-response analysis of signaling by the MBP-CD97 fusion and analysis of other N-terminal fusions. (a) Comparison of HA-FL-FLAG, M-CTF-FLAG, and HA-MBP-CTF-FLAG in the SRE assay across a range of DNA transfection amounts (0.25, 0.5, 1, 2, 10, 30, or 60 ng per well). All data points are normalized to the mean value for empty vector (60 ng). Sigmoidal four parameter logistic curves were fit to each titration curve with nonlinear regression. Data are presented as mean ± s.d. of three technical replicates and representative of three independent biological replicates. Two-way ANOVA with Tukey's multiple-comparison post-hoc test was used to compare MBP-CTF and M-CTF data points at each DNA load, and one-way ANOVA with Tukey's multiple-comparison post-hoc test was used to compare 60 ng DNA loads with empty vector (ns, P>0.05, **P<0.01, ***P<0.001, ****P<0.0001). **(b)** α-FLAG Western blot comparison of protein abundance for the molecules evaluated in **(a)**. Individual lanes correspond to DNA amounts used in **(a)**. Blotting for β-tubulin was used to control for protein loading. **(c)** Signal response of the CD97 MBP-CTF when co-expressed with varying amounts (0-30 ng) of secreted TEV protease. Cells were transfected with 30 ng receptor DNA/well. Data are normalized to the mean value for empty vector and are presented as mean ± s.d. of three technical replicates and representative of three independent biological replicates. **(d)** Signaling data for a panel of N-terminal sequences fused to the CD97 CTF. Cells were transfected with 30 ng DNA/well. Data are normalized to the mean value for empty vector and are presented as mean ± s.d. of three technical replicates and representative of three independent biological replicates.

Figure 4 – Use of the MBP fusion platform to determine the G protein coupling profile of CD97. Engineered HEK293T cells lacking Gα$_s$, Gα$_{olf}$, Gα$_q$, Gα$_{11}$, Gα$_{12}$, and Gα$_{13}$ (HEKΔ6), as well as Gα$_{16}$ (which is not expressed in parental cells) were used to identify the G protein coupling profile of CD97. **(a)** Gα subunits (or empty vector for Gα-encoding plasmids, EV$_{Gα}$) were individually co-transfected with empty vector, CD97 FL, or CD97 MBP-CTF. Signals for each form of CD97 were evaluated in the SRE assay. Values are reported as the ratio of the raw luminescence values of firefly and renilla luciferases. **(b,c)** Signaling response of the CD97 MBP-CTF when co-transfected with increasing amounts of Gα$_{12}$ **(b)** or Gα$_{13}$ **(c)**. Data are normalized to the mean reporter signal generated by co-transfection of empty vector with 0 ng Gα plasmid (the leftmost bar on each graph). For all panels, cells were transfected with 30 ng receptor DNA/well. Data are presented as mean ± s.d. of three technical replicates and representative of three independent biological replicates. One-way ANOVA with Tukey's multiple-comparison post-hoc test was used to compare EV and MBP-CTF samples in **(b,c)** (ns, P>0.05,*P<0.05, ****P<0.0001).

710
Figure 5 – Scanning mutagenesis of the CD97 stalk sequence using the N-terminal fusion
platform. (a) ColabFold-generated structural prediction of the stalk-bound 7TM domain of
CD97 in complex with Gα13[56] A slice through the orthosteric pocket of the receptor is depicted,
with the bound stalk sequence colored on a blue-to-red rainbow scale from N- to C-terminus. **(b)**
Sequence conservation of the first nine stalk residues of CD97 based on alignment of the 33
human aGPCRs. **(c)** SRE signaling data for scanning alanine (or glycine for A534 and A538) and
lysine mutagenesis of the stalk sequence in CD97. All mutations were made in the context of
parental IL2ss-HA-MBP-TEV-CD97 CTF. Cells were transfected with 30 ng DNA/well. Data
are normalized as a percentage of the mean value for WT and presented as mean ± s.d. of three
technical replicates and representative of three independent biological replicates. **(d)** Mapping of
mutagenesis results onto the structural model of the CD97 CTF.

Figure 6 – Probing mechanisms of synthetic and natural autoinhibition in CD97. (a) Schematic diagrams depicting constructs in which MBP is fused at decreasing distances from the TA. **(b)** Signaling response of MBP fusion constructs shown in **(a)**. **(c)** Model illustrating the variation in steric interference depending on the fusion position of MBP. **(d)** Signaling response of proteins with insertion segments of varying lengths from the CD97 GAIN domain between MBP and the CD97 CTF (see Extended Data Fig. 7 a,b for construct design). **(e)** Schematic depiction of TA sequestration or exposure depending on whether part (left) or all of the GAIN domain is included (right). Data in **(b)** and **(d)** are normalized to the mean value for empty vector and are presented as mean ± s.d. of three technical replicates and representative of three independent biological replicates. One-way ANOVA with Tukey's multiple-comparison post-hoc test was used to compare MBP-CTF with all other receptor constructs in **(b)** and with the FL and MBP-GAIN-CTF constructs in **(d)** (ns, $P>0.05$, **$P<0.01$, ****$P<0.0001$).

Figure 7 – Signaling assays for MBP-fused CTFs from a panel of aGPCRs. (a) Schematic diagrams of GPR56 constructs evaluated. **(b)** Schematic depicting the domain architecture of GPR56 constructs. The GPR56 headpiece is comprised of a Pentraxin/Laminin/neurexin/sex-hormone-binding-globulin-Like (PLL) domain followed by a conserved GAIN domain. **(c–f)** Signaling response of the indicated GPR56-derived molecules in SRE **(c)**, SRF-RE **(d)**, NFAT-RE **(e)**, and CRE **(f)** reporter assays. Cells were transfected with 30 ng DNA/well **(c,d)** or 10 ng DNA/well **(e,f)**. **(g)** MBP-CTF constructs for a panel of aGPCRs were evaluated in the SRE and CRE reporter assays. Cells were transfected with 30 ng DNA/well (SRE) or 10 ng DNA/well (CRE). Each receptor is expressed using the HA-MBP-TEV-CTF-FLAG protein design. The long C-terminal tail regions of ADGRL3 and ADGRG2 were truncated for ease of cloning and expression (see Table S1). Data in **(c–g)** are normalized to the mean value for empty vector (EV). Data are reported as mean ± s.d. of three technical replicates and representative of two **(c–f)** or three **(g)** independent biological replicates.

Extended Data Figures

Extended Data Figure 1 – Effector dependence of genetic reporter assays utilized in this study. G_s and G_i heterotrimers modulate cAMP production, which is reported by the cAMP response element (CRE). The Serum Response Element (SRE) is dependent on signals from both $ERK_{1/2}$ and RhoA proteins.[59] $ERK_{1/2}$ is stimulated by G_i-derived $G\beta\gamma$ subunits as well as a signaling cascade downstream of phospholipase C (PLC) activity, which is controlled by G_q heterotrimer stimulation. The small GTPase RhoA is activated by GTP-bound $G\alpha_{12}$. The Serum Response Factor Response Element (SRF-RE) is an engineered form of the SRE reporter to respond exclusively to RhoA activity.[60] Finally, the Nuclear Factor of Activated T Cells Response Element (NFAT-RE) is downstream of the canonically G_q-modulated intracellular calcium signaling cascade. These pathways are not meant to be exhaustive, but reflect the generally-accepted signaling cascades downstream of the four primary G protein subfamilies.

Extended Data Figure 2 – Validation of genetic reporter assays and aGPCR construct
design. (a–d) Validation of reporters with muscarinic M3 receptors (M3R) stimulated with
10μM carbachol **(a–c)** or endogenous β-adrenergic receptors (β-ARs) stimulated with 100nM
isoproterenol **(d)**. Cells were treated for 6 hours with drug (induced) or DMSO (uninduced) prior
to readout. **(e,f)** Western blot detection of C-terminal FLAG epitope tags to analyze total protein
amounts for proteins evaluated in Fig. 2 a,b **(e)** and Fig. 2 c,d **(f)**. Blotting for β-tubulin was used
to control for protein loading. **(g,h)** Titration of M-CTF and M-SL DNA in Western blot against
C-terminal FLAG epitope tags **(g)** and the SRE assay **(h)**. **(i)** Effects of the addition of C-
terminal FLAG epitope tags on signaling constructs used in Fig. 2. Data in **(a–d,h,i)** are
normalized to the mean value for empty vector. Cells were transfected with 30 ng DNA/well.
Data are presented as mean ± s.d. of three technical replicates and representative of two **(h)** or
three **(a–d,i)** independent biological replicates.

Extended Data Figure 3 – Effects of removing the stalk sequence from the MBP-CTF framework. (a) Comparison of the indicated CD97-derived proteins in the SRE assay across a range of transfected DNA amounts (0.25, 0.5, 1, 2, 10, 30, or 60 ng per well). All data points are normalized to the mean value for empty vector (60 ng). Sigmoidal four parameter logistic curves were fit to each titration curve with nonlinear regression. Data are presented as mean ± s.d. of three technical replicates and representative of three independent biological replicates. Two-way ANOVA with Tukey's multiple-comparison post-hoc test was used to compare MBP-CTF and M-CTF data points at each DNA load, and one-way ANOVA with Tukey's multiple-comparison post-hoc test was used to compare 60 ng DNA loads with empty vector (ns, P>0.05,**P<0.01, ****P<0.0001). **(b)** Western blot detection of N-terminal HA tags on the proteins evaluated in **(a)**. Individual lanes correspond to the amounts of transfected DNA used in **(a).** Note that CD97 FL is predicted to be heavily glycosylated. Blotting for GAPDH was used to control for protein loading. **(c)** Flow cytometry to assess cell surface protein abundance. Proteins were detected using an antibody against the N-terminal HA epitope tag. Data are presented as mean fluorescence intensity values. Data are presented as mean ± s.d. of three technical replicates and representative of three independent biological replicates. One-way ANOVA with Tukey's multiple-comparison post-hoc test was used to compare samples (****P<0.0001).

Extended Data Figure 4 – Western blots probing a panel of N-terminal CTF fusions TEV
protease-catalyzed release of the MBP tag from MBP-CTF. (a) Western blot detection of N-
terminal HA epitope tags for fusion proteins evaluated in Fig. 3 c. Note that CD97 FL is
predicted to be heavily glycosylated. Blotting for GAPDH was used to control for protein
loading. **(b)** Western blot detection of N-terminal HA epitope tags of the CD97 MBP-CTF when
co-expressed with varying amounts (2-30 ng) of secreted TEV protease. Expected weights of and
pre- and post-processed protein are 76.6 kDa and 42.3 kDa, respectively. Blotting for GAPDH
was used to control for protein loading.

839
840
841
842

Extended Data Figure 5 – Calibration of G protein knockout cells and Gα rescue system to deconvolute G protein subtype specificity. (a) Reporter plasmids were co-transfected into HEKΔ6 cells with DNA encoding each Gα subtype (titrated up to 50 ng) and the relative luciferase activity was measured 30 hours after transfection. Gα DNA transfection amounts selected for rescue experiments are shown by dashed lines intersecting the x-axis: 20 ng for $G\alpha_s$, $G\alpha_q$, $G\alpha_{11}$, and $G\alpha_{16}$; 0.3 ng for $G\alpha_{12}$ and $G\alpha_{13}$. Data are normalized to the mean value for empty vector (50 ng). Data are presented as mean ± s.d. of three technical replicates and representative of two independent biological replicates. **(b)** Basal signaling activity induced by the amounts of Gα DNA transfected in rescue experiments. Values are reported as the ratio of the raw luminescence values of firefly and renilla luciferases. Data are presented as mean ± s.d. of three technical replicates and representative of three independent biological replicates. **(c)** Rescue experiment with 10 μM carbachol-stimulated M3R to confirm its known coupling profile of $G\alpha_q$ and $G\alpha_{11}$. Values are reported as the ratio of the raw luminescence values of firefly and renilla luciferases. Cells were transfected with 30 ng receptor DNA/well. Data are presented as mean ± s.d. of three technical replicates and representative of three independent biological replicates. **(d,e)** DNA titration with $G\alpha_s$ **(d)** and $G\alpha_{16}$ **(e)** to confirm their ability to induce signals in HEKΔ6 cells. Data are normalized to the mean value for empty vector (50 ng for **(d)** and 100 ng for **(e)**). Data are presented as mean ± s.d. of three technical replicates and representative of three independent biological replicates.

Extended Data Figure 6 – Effect of triple alanine mutations in the stalk linker and Western blot evaluation of the expression of all mutant proteins tested. (a) Signaling responses of MBP-CTF constructs harboring triple alanine mutations of residues in the P10-15 positions of the CD97 stalk sequence. Data are normalized to the mean value for empty vector. Data are presented as mean ± s.d. of three technical replicates and representative of three independent biological replicates. One-way ANOVA was used with Tukey's multiple-comparison post-hoc test to compare the difference between samples (ns, P>0.05). **(b,c).** Western blot detection of N-terminal HA epitope tags to evaluate total protein expression for constructs evaluated in Fig. 5 c and Extended Data Fig. 6 a. Blotting for GAPDH was used to control for protein loading.

Extended Data Figure 7 – Construct design and validation of molecules used to test synthetic and natural autoinhibition in CD97. (a) Western blot detection of N-terminal HA epitope tags to evaluate total protein expression for GAIN domain insertions evaluated in Fig. 6 b. Blotting for GAPDH was used to control for protein loading. **(b)** Alphafold2 prediction of the CD97 GAIN domain.[57,58] GAIN fragments tested in reporter assays are colored on the model as indicated on the schematics at right. **(c)** Western blot detection of N-terminal HA and C-terminal FLAG epitope tags on proteins tested in Fig. 6 d. Blotting for GAPDH was used to control for protein loading

Extended Data Figure 8 – Evaluation of expression and processing of a panel of aGPCR MBP-CTF proteins. (a) Western blot detection of C-terminal FLAG epitope tags on GPR56-derived proteins tested in Fig. 7 c,d. Blotting for β-tubulin was used to control for protein loading. **(b,c)** Titration of M-CTF and MBP-CTF DNA in the SRE assay **(b)** and Western blot against C-terminal FLAG epitope tags **(c)**. Data in **(b)** are normalized to the mean value for empty vector. Data are presented as mean ± s.d. of three technical replicates and representative of two independent biological replicates. **(d)** Western blot detection of N-terminal HA and C-terminal FLAG epitope tags for proteins evaluated in Fig. 7 g. Blotting for GAPDH was used to control for protein loading.

Table S1 – Description of constructs utilized in this study

Name	Mutations/ Modifications	Vector	Description	Calculated MW (if blotted)	Ref
SRE/MinP/*luc2*		pGL4.23	Luciferase reporter vector using the serum response element, a minimal promoter, and luc2 firefly luciferase		
SRF-RE/MinP/*luc2*		pGL4.23	Luciferase reporter vector using the serum response factor response element, a minimal promoter, and luc2 firefly luciferase		
NFAT-RE/MinP/*luc2*		pGL4.23	Luciferase reporter vector using the nuclear factor of activated T cells response element, a minimal promoter, and luc2 firefly luciferase		
CRE/MinP/*luc2*		pGL4.23	Luciferase reporter vector using the cAMP response element, a minimal promoter, and luc2 firefly luciferase		
EV		pFuse-hIgG1-Fc2	Empty pFuse-hIgG1-Fc2 plasmid with a stop codon inserted between the IL2 signal peptide and the multiple cloning site		
HA-CD97 FL-FLAG		pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, full-length CD97 isoform 1, and a FLAG tag	56.7 kDa (processed NTF); 35.4 kDa (processed CTF)	
CD97 M-CTF-FLAG	Residues 1-530 of CD97 isoform 1 removed	pFuse-hIgG1-Fc1	Methionine residue followed by a CD97 construct N-terminally truncated at the autoproteolytic cleavage site (CD97 CTF) and a FLAG tag	35.5 kDa	
CD97 M-SL-FLAG	Residues 1-545 of CD97 isoform 1 removed	pFuse-hIgG1-Fc1	Methionine residue followed by a CD97 construct N-terminally truncated at the start of TM1 (CD97 SL) and a FLAG tag	33.8 kDa	
HA-MBP-CD97 CTF-FLAG	Residues 1-530 of CD97 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, the CD97 CTF, and a FLAG tag	77.7 kDa	
HA-MBP-CD97 CTF		pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa	
secTEV	N23Q, C130S, and T173G substitutions	pcDNA3.1	IL2 signal peptide followed by the coding sequence of secTEV protease		45
HA-CD97 FL		pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, full-length CD97 isoform 1	90.9 kDa (unprocessed); 56.7 kDa (processed NTF)	
HA-BRIL-CD97 CTF		pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, bone restricted ifitm-like protein (BRIL), TEV cleavage sequence residues P6-P1, and the CD97 CTF	48.0 kDa	
HA-PGS-CD97 CTF		pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, a glycogen synthase domain from Pyrococcus abyssi (PGS), TEV cleavage sequence residues P6-P1, and the CD97 CTF	58.1 kDa	
HA-T4L-CD97 CTF		pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, T4-lysozyme, TEV cleavage sequence residues P6-P1, and the CD97 CTF	54.4 kDa	
HA-3xFLAG-CD97 CTF		pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, a 3xFLAG tag, TEV cleavage sequence residues P6-P1, and the CD97 CTF	39.0 kDa	
HA-MBP-CD97 CTF S531A	Alanine substitution S531A	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa	
HA-MBP-CD97 CTF S532A	Alanine substitution S532A	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa	
HA-MBP-CD97 CTF F533A	Alanine substitution F533A	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa	
HA-MBP-CD97 CTF A534G	Glycine substitution A534G	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa	

HA-MBP-CD97 CTF I535A	Alanine substitution I535A	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa
HA-MBP-CD97 CTF L536A	Alanine substitution L536A	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa
HA-MBP-CD97 CTF M537A	Alanine substitution M537A	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa
HA-MBP-CD97 CTF A538G	Glycine substitution A538G	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa
HA-MBP-CD97 CTF H539A	Alanine substitution H539A	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa
HA-MBP-CD97 CTF S531K	Lysine substitution S531K	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa
HA-MBP-CD97 CTF S532K	Lysine substitution S532K	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa
HA-MBP-CD97 CTF F533K	Lysine substitution F533K	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa
HA-MBP-CD97 CTF A534K	Lysine substitution A534K	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa
HA-MBP-CD97 CTF I535K	Lysine substitution I535K	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa
HA-MBP-CD97 CTF L536K	Lysine substitution L536K	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa
HA-MBP-CD97 CTF M537K	Lysine substitution M537K	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa
HA-MBP-CD97 CTF A538K	Lysine substitution A538K	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa
HA-MBP-CD97 CTF H539K	Lysine substitution H539K	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa
HA-MBP-S531		pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, the CD97 CTF, and a FLAG tag	75.8 kDa
HA-MBP-S532	Residues 1-531 of CD97 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, an N-terminally-truncated CD97 CTF, and a FLAG tag	75.7 kDa
HA-MBP-F533	Residues 1-532 of CD97 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, an N-terminally-truncated CD97 CTF, and a FLAG tag	75.6 kDa
HA-MBP-TEV-F533	Residues 1-532 of CD97 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, an N-terminally truncated CD97 CTF, and a FLAG tag	76.4 kDa
HA-MBP-GAIN-CTF-FLAG	Residues 1-256 of CD97 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, the CD97 GAIN domain, the CD97 CTF, and a FLAG tag	107.5 kDa (unprocessed); 72.2 kDa (processed NTF);
HA-MBP-Fragment 5-CTF-FLAG	Residues 1-347 of CD97 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, a fragment of the CD97 GAIN domain, the CD97 CTF, and a FLAG tag	97.5 kDa
HA-MBP-Fragment 4-CTF-FLAG	Residues 1-434 of CD97 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, a fragment of the CD97 GAIN domain, the CD97 CTF, and a FLAG tag	88.2 kDa

HA-MBP-Fragment 3-CTF-FLAG	Residues 1-489 of CD97 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, a fragment of the CD97 GAIN domain, the CD97 CTF, and a FLAG tag	82.2 kDa	
HA-MBP-Fragment 2-CTF-FLAG	Residues 1-510 of CD97 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, a fragment of the CD97 GAIN domain, the CD97 CTF, and a FLAG tag	79.8 kDa	
HA-MBP-Fragment 1-CTF-FLAG	Residues 1-520 of CD97 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, a fragment of the CD97 GAIN domain, the CD97 CTF, and a FLAG tag	78.8 kDa	
HA-GPR56 FL-FLAG		pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, full-length GPR56 isoform 1, and a FLAG tag	77.4 kDa	
GPR56 M-CTF-FLAG	Residues 1-382 of GPR56 isoform 1 removed	pFuse-hIgG1-Fc1	Methionine residue followed by a GPR56 construct N-terminally truncated at the autoproteolytic cleavage site (GPR56 CTF) and a FLAG tag	35.8 kDa	
HA-MBP-GPR56 CTF-FLAG	Residues 1-382 of GPR56 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, the GPR56 CTF, and a FLAG tag	78.0 kDa	
ADGRF1	Residues 1-566 of ADGRF1 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, the ADGRF1 CTF, and a FLAG tag	81.8 kDa	
ADGRL3	Residues 1-841 and 1134-1447 of ADGRL3 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, the ADGRL3 CTF with a truncated C-terminal tail, and a FLAG tag	76.6 kDa	
ADGRG3	Residues 1-249 of ADGRG3 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, the ADGRG3 CTF, and a FLAG tag	76.9 kDa	
ADGRV1	Residues 1-5883 of ADGRV1 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, the ADGRV1 CTF, and a FLAG tag	89.5 kDa	
ADGRG2	Residues 1-606 and 912-1017 of ADGRG2 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, the ADGRG2 CTF with a truncated C-terminal tail, and a FLAG tag	78.3 kDa	
ADGRG5	Residues 1-226 of ADGRG5 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, the ADGRG5 CTF, and a FLAG tag	76.8 kDa	
ADGRD1	Residues 1-544 of ADGRD1 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, the ADGRD1 CTF, and a FLAG tag	79.5 kDa	
ADGRG6	Residues 1-840 of ADGRG6 isoform 1 removed	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, the ADGRG6 CTF, and a FLAG tag	86.7 kDa	
HA-CHRM3-FLAG		pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, the CHRM3 coding sequence, and a FLAG tag		
CD97 M-CTF		pFuse-hIgG1-Fc1	Methionine residue followed by the CD97 CTF		
CD97 M-SL		pFuse-hIgG1-Fc1	Methionine residue followed by CD97 SL		
HA-MBP-CD97 SL		pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and CD97 SL	74.8 kDa	

HA-MBP-CD97 CTF YDV->A	Alanine substitutions Y540A, D541A, and V542A	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa	
HA-MBP-CD97 CTF EDW->A	Alanine substitutions E543A, D544A, and W545A	pFuse-hIgG1-Fc2	IL2 signal peptide followed by an HA tag, maltose binding protein, TEV cleavage sequence residues P6-P1, and the CD97 CTF	76.6 kDa	