**Short Title:** Lipid complement of *Arabidopsis* leaf mitochondria

**Defining the lipidome of Arabidopsis leaf mitochondria: Specific lipid complement and lipid**

**biosynthesis capacity**

Yi-Tse Liu[1,†], Jennifer Senkler[2,†], Cornelia Herrfurth[1,3], Hans-Peter Braun[2,*], Ivo Feussner[1,3,4,*]

[1]University of Goettingen, Albrecht-von-Haller-Institute for Plant Sciences, Department of Plant

Biochemistry, 37077 Goettingen, Germany. [2]Leibniz Universität Hannover, Institute of Plant

Genetics, Department of Plant Proteomics, 30419 Hannover, Germany. [3]University of

Goettingen, Goettingen Center for Molecular Biosciences (GZMB), Service Unit for

Metabolomics and Lipidomics, 37077 Goettingen, Germany. [4]University of Goettingen,

Goettingen Center for Molecular Biosciences (GZMB), Department of Plant Biochemistry, 37077

Goettingen, Germany

**\*Authors for correspondence:** Hans-Peter Braun, Herrenhäuser Str. 2, 30419 Hannover,

Germany; Fax. +49 511 76214351; Email: braun@genetik.uni-hannover.de; Ivo Feussner, Justus-

von-Liebig-Weg 11, 37077 Goettingen, Germany; Fax: +49-551-39-25749; Email: ifeussn@uni-

goettingen.de

†These authors contributed equally to this work.

The authors responsible for distribution of materials integral to the findings presented in this

article in accordance with the policy described in the Instructions for Authors

(https://academic.oup.com/plphys/pages/General-Instructions) are Hans-Peter Braun,

braun@genetik.uni-hannover.de and Ivo Feussner, ifeussn@uni-goettingen.de.

**Author contributions:** YL performed experiments including lipid analysis and pathway

construction; JS performed proteomic analysis. YL, JS, CH, HB and IF analyzed the data. YL wrote

the first draft of the manuscript. CH, HB and IF edited the manuscript. HB and IF designed and

supervised the study. IF agrees to serve as the author responsible for contact and ensures

communication.

**Competing interests:** Authors declare no competing interest.

**Key words:** *Arabidopsis thaliana*, lipidome, mitochondrium, proteome, respiration

**One sentence summary:** The lipid landscape of plant mitochondria suggests that they are

capable in generating several phospholipid classes with the assistant of membrane contact site-

localized proteins.

**Summary**

Mitochondria are often considered the power stations of the cell, playing critical roles in various

biological processes such as cellular respiration, photosynthesis, stress responses and

programmed cell death. To maintain the structural and functional integrities of mitochondria, it

is crucial to achieve a defined membrane lipid composition between different lipid classes

wherein specific proportions of individual lipid species are present. Although mitochondria are

capable of self-synthesizing a few lipid classes, many phospholipids are synthesized in the

endoplasmic reticulum and transferred to mitochondria via membrane contact sites, as

mitochondria are excluded from the vesicular transportation pathway. However, knowledge on

the capability of lipid biosynthesis in mitochondria and the precise mechanism of maintaining

the homeostasis of mitochondrial lipids is still scarce. Here we describe the lipidome of

mitochondria isolated from *Arabidopsis* leaves, including the molecular species of glycerolipids,

sphingolipids and sterols to depict the lipid landscape of mitochondrial membranes. In addition,

we define proteins involved in lipid metabolism by proteomic analysis and compare our data

with mitochondria from cell cultures since they still serve as model system. Proteins putatively

localized to the membrane contact sites are proposed based on the proteomic results and

online databases. Collectively, our results suggest that leaf mitochondria are capable - with the

assistance of membrane contact site-localised proteins - of generating several lipid classes

including phosphatidylethanolamines, cardiolipins, diacylgalactosylglycerols and free sterols. We

anticipate our work to be a foundation to further investigate the functional roles of lipids and

their involvement in biochemical reactions in plant mitochondria.

**Introduction**

Mitochondria are considered as semiautonomous organelles. According to a widely accepted

hypothesis, they descend from proteobacteria that have been engulfed to form an eukaryotic

cell. The two involved cells established an endosymbiosis (Gray et al., 1999). Mitochondria play

crucial roles in various cellular processes, including ATP generation, metabolite biosynthesis,

stress responses like phosphate starvation and initiation of programmed cell death (Jacoby et al.,

2012; Michaud et al., 2016). The mitochondria of plant cells even have extended functions,

many of which are related to photosynthesis (Braun, 2020). Mitochondria are enclosed by two

membranes, the outer (OM) and the inner (IM) mitochondrial membranes, that both require

defined protein and lipid compositions to maintain their functional integrity (Horvath and Daum,

2013; Michaud et al., 2017). The majority of the mitochondrial proteins is encoded by the

nuclear genome, synthesized in the cytoplasm and post-translationally imported into the

organelle. At the same time, a few proteins are encoded by the mitochondrial genome and

synthesized within the organelle. Recent studies have broadened the knowledge of protein and

lipid import in mitochondria, to the OM, the intermembrane space, the IM and the matrix

(Michaud et al., 2017).

Biological membranes are composed of a wide variety of lipid classes and molecular species.

According to the backbones of the lipid molecules, they are classified into three main lipid

categories – glycerolipids, sphingolipids and sterols. To delineate the molecular species in this

study, we later use the taxonomy containing two colon-separated units for the numbers of

carbons and double bonds in the fatty acyl moiety, and an additional unit after a semicolon for

the number of hydroxyl groups (when present) (Liebisch et al., 2020). Plant mitochondrial

membranes contain high amounts of glycerophospholipids, such as phosphatidylcholine (PC)

and phosphatidylethanolamine (PE), but contain less than 2 % of sterols (Moreau et al., 1974;

Bligny and Douce, 1980). Sphingolipids have so far not been detected in mitochondrial

membranes form plants. Most of these lipids are produced in the ER and are sorted to

mitochondria and other organelles afterwards. Nevertheless, mitochondria have also their own

capacity to generate some specific lipid classes. For instance, cardiolipin (CL) is synthesized in

the IM and is present exclusively in mitochondria (Douce et al., 1972; Babiychuk et al., 2003). CL

plays an essential role establishing the cristae of the IM and in maintaining the mitochondrial

ultrastructure. CL is formed through the condensation of phosphatidylglycerol (PG) and

diacylglycerol (DAG) and finally consists of four acyl chains. The inner envelope of the plastid is

the major site for generating PG molecules (Müller and Frentzen, 2001). However, enzymes

involved in PG biosynthesis are identified in mitochondria as well, suggesting that mitochondria

are capable of self-synthesizing PG and thus CL (Babiychuk et al., 2003). In plants, the PG

synthesizing enzymes, phosphatidylglycerolphosphate (PGP) synthase and PGP phosphatase

(PGPP), are associated with mitochondria, and plastids; whereas CL synthase (CLS) localizes

exclusively in the IM (Xu et al., 2002; Katayama et al., 2004). Notably, although mitochondrial

lipid biosynthesis is not the major lipid source in plant cells, it is critical for certain organisms. In

yeast, mitochondria are the major supplier of PE (Horvath and Daum, 2013). They generate PE

from phosphatidylserine (PS) by phosphatidylserine decarboxylase (PSD). In *Arabidopsis*, three

PSD enzymes have been identified and PSD1 localizes in mitochondria, providing PE molecules *in*

*situ* (Nerlich et al., 2007). Alternatively, some PS molecules are converted from PE through the

base-exchange pathway that substitutes the head groups of PE with serine molecules by PS

synthase1, PSS1 (Yamaoka et al., 2011). While knowing mitochondria are capable of synthesizing

PG, CL, PS and PE, its competence toward other lipid classes is still poorly understood (Li-Beisson

et al., 2013).

Lipid trafficking between ER, mitochondria and plastids is essential for mitochondrial membrane

biogenesis (Horvath and Daum, 2013; Michaud et al., 2017). Based on studies in yeast and

mammals, glycerophospholipid biosynthesis takes place at a distinct membrane stretch of the

ER, the mitochondria-associated membrane (MAM), wherein both ER and mitochondrial

proteins have been identified (Vance, 1990). Elevated activities of PE, PC and PS synthesizing

enzymes have been detected in purified yeast MAM. Moreover, this distinct membrane domain

seems to occur ubiquitously among plants and animal cells, suggesting its critical role during

evolution and in mediating lipid transfer between ER and mitochondria (Morré et al., 1971;

Staehelin, 1997; Achleitner et al., 1999; Michaud et al., 2016). Although mitochondria and

plastids work closely together in numerous pathways in plants, the lipid transport mechanism

between these two organelles is largely unknown. Nevertheless, a few studies have suggested

the relevance of lipid trafficking between these two organelles for survival, especially under

environmental stresses. During phosphate starvation, higher numbers of mitochondria – plastid

junctions are established (Jouhet et al., 2004). At the same time, drastic lipid remodeling of

mitochondria, plastids and plasma membrane arises. PC and PE are degraded to release the

phosphate residues for essential biological processes and the remaining molecules are recycled

to generate the typical plastidial glyceroglycolipid, digalactosyldiacylglycerol (DGDG). This

coincides with increased levels of CL in mitochondria isolated from *Arabidopsis* suspension cells

and calli (Michaud et al., 2017). In *Arabidopsis*, the protein complex involved in lipid trafficking

and tethering of the two mitochondrial membranes, the mitochondrial transmembrane

lipoprotein (MTL) complex, has been identified recently (Michaud et al., 2016). MTL is composed

of more than 200 subunits and it has been demonstrated that MTL promotes the translocation

of PE from IM to OM and the import of DGDG from plastids to mitochondria during phosphate

starvation. Furthermore, mutation of a newly identified MTL subunit, digalactosyldiacylglycerol

synthase suppressor 1 (DGS1), leads to alteration of plastidial and mitochondrial lipid

composition and deficiency in mitochondrial biogenesis (Li et al., 2019). The characterization of

the MTL complex provides an initial insight in understanding the mechanism of lipid trafficking

between mitochondria and plastids.

In this study, we aimed to characterize the lipid metabolism of *Arabidopsis* leaf mitochondria in

depth and to compare these data with published and own data from suspension cell cultures

since they serve as model system. Therefore, we conducted an in-depth lipidomic analysis,

providing the molecular species information of all lipid categories including glycerolipids,

sphingolipids and sterols to illustrate the lipid landscape of mitochondria in *Arabidopsis* leaves.

In combination with a proteomic approach, we intended to specify the capacity of lipid

biosynthesis and modification in plant mitochondria, defining lipid species and classes that may

be generated by mitochondria. We additionally propose putative membrane contact site-

localizing proteins and their roles in interorganelle communication. Our results suggest that leaf

mitochondria possess a defined lipid composition wherein specific lipid molecular species

appear. In addition, lipid trafficking between mitochondria, ER and plastids via membrane

contact sites provides assistance in maintaining the homeostasis of the lipid composition in

mitochondria.

**Results**

**Purity of mitochondrial fractions**

A combinatorial approach by lipidomics, proteomics and mining of online databases was applied

to investigate the lipid composition as well as the capacity of lipid metabolism of *Arabidopsis*

leaf mitochondria. Mitochondria were isolated from three independent experiments by

differential centrifugation combined with Percoll density gradient centrifugation. To ensure and

evaluate the purity of the mitochondrial fractions, all samples were investigated by two-

dimensional (2D) blue native (BN)/SDS PAGE and by shotgun proteome analyses. Our

mitochondrial fractions proved to be of high purity. The two photosystems were not detectable

on our 2D gels. Only trace amounts of the large subunit from Rubisco, the most abundant

protein in leaves, were visible in the leaf mitochondrial fractions (L-mito) on BN/SDS gels, and

the small subunit of Rubisco was barely visible (Fig. 1A, Supplemental Fig. S1). Since most of the

work on plant mitochondria was performed with suspension cell cultures so far, we isolated

mitochondria from dark-grown *Arabidopsis* cell culture (C-mito) for comparison. They were

investigated by the same procedure and used as an established control for contamination from

chloroplasts. The L-mito and C-mito fractions were highly similar on BN/SDS PAGE (Fig. 1,

Supplemental Fig. S1). In addition, the purity of the mitochondrial fractions was investigated by

label-free quantitative shotgun proteomics. Building on the proteome data (Supplemental Tab.

S1), the proportion of mitochondria-localized proteins was calculated to be in the range of 87 to

94 % (Fig. 1B, Supplemental Fig. S2). These results indicate that the mitochondrial fractions are

of high purity.

**Figure 1**. Purity of mitochondrial fractions. The purity of mitochondrial fractions was determined

by 2D blue native / SDS PAGE **(A)** and by summed-up peptide intensities of subcellular

compartments based on protein assignments as given by the Subcellular localization database

for Arabidopsis proteins (SUBAcon; www.suba.live) **(B)**.

**A**: Mitochondria were isolated from *Arabidopsis* leaves (L-mito) and cell cultures (C-mito).

Proteins were separated by 2D Blue native PAGE and Coommassie-stained. Numbers on top and

to the left of the 2D gels refer to the masses of standard protein complexes / proteins (in kDa),

the roman numbers above the gels to the identity of OXPHOS complexes. I+III$_2$: supercomplex

consisting of complex I and dimeric complex III; I: complex I; V: complex V; III$_2$: dimeric complex

III; IV: complex IV. The small (S; 14.5 kDa) and the large (L; 53.5 kDa) subunit of Rubisco are

indicated by green arrows. Mitochondrial preparations derive from three independent

experiments, and the corresponding 2D gels of the two remaining biological replicates as well as

one reference gel each for mitochondrial and chloroplast fractions from *Arabidopsis* cell culture

(Mito ref) or leaves (Cp rev) are shown in Supplemental Fig. S1. These three preparations were

used for lipidomics with LC-MS/MS (Fig. 3, 5, 6, and S3) and proteomics **(B)**.

**B:** Mitochondrial fractions from *Arabidopsis* leaves and cell cultures were analyzed by label-free

quantitative shotgun proteomics. Peptide intensities assigned to subcellular compartments

were summed-up and averaged results for L-mitos and C-mitos from the three independent

experiments were visualized by pie charts (for detailed results see Supplemental Fig. S2 and

Supplemental Tab. S1). Blue: mitochondria; green: plastids; gray: others; numbers in %.

**PC, PE and CL are strongly enriched glycerolipids in plant leaf mitochondria**

Glycerolipids are the most abundant lipids not only in mitochondria but also generally in plant

tissues, accounting for more than 90 % of the overall lipids in leaves (Li-Beisson et al., 2013;

Michaud et al., 2017). Glycerophospholipids are fundamental to all biological membranes, while

glyceroglycolipids are critical for photosynthetic membranes and localize mainly in plastids of

vegetative tissues (Hölzl and Dörmann, 2019). Therefore, lipids were extracted from leaves and

purified mitochondria therefrom to determine the proportion of the major glycerolipid classes

in a quantitative approach based on thin layer chromatography followed by gas chromatography

coupled to flame ionization detection (TLC-GC/FID, Fig. 2). In L-TE, glyceroglycolipids including

monogalactosyldiacylglycerols (MGDG; 38.9 ± 1.1 %) and DGDG (22.3 ± 3.8 %) contribute to the

majority of the overall lipids; while glycerophospholipids, PC (35.7 ± 6.0 %) and PE (37.2 ± 0.3 %),

are the most abundant lipids in L-mito. CL accounts for 4.0 ± 0.9 % in the lipidome of L-mito;

however, it was not detectable in L-TE because of its low abundance in total cellular lipids. PG,

as the precursor of CL, consists 5.6 ± 2.1 % of the overall lipids in L-mito.

**Figure 2.** Lipid class profiles of purified mitochondria and total leaf extracts. Glycerolipids of leaf

total extract (L-TE) and mitochondria isolated from leaves (L-mito) were analyzed quantitatively

by a TLC-GC/FID approach. Data of L-TE represent mean values in mol % from three

independent experiments ± SD; data of L-mito represent mean values in mol % from two of the

three independent experiments ± SEM. PC, phosphatidylcholine; PE, phosphatidylethanolamine;

CL, cardiolipin; PG, phosphatidylglycerol; MGDG, monogalactosyldiacylglycerol; DGDG,

digalactosyldiacylglycerol.

**Specific molecular glycerolipid species are enriched in plant leaf mitochondria**

Glycerophospholipids are the most abundant lipids in mitochondria (Fig. 2) (Horvath and Daum,

2013; Michaud et al., 2017). Therefore, the molecular species of glycerophospholipids, as well as

those of glyceroglycolipids and diacylglycerols, were analyzed from the same samples

originating from three independent experiments as for proteomics (Fig. 1) by LC-MS/MS (Fig. 3a).

Data are expressed in mol %. The most abundant PC and PE molecular species in mitochondria

with the acyl chains 18:2/18:3, 18:2/18:2 and 18:3/18:3 account for more than 60 % in both lipid

classes (Fig. 3a; Supplemental Tab. S2). The species 16:0/18:2, 16:0/18:3 of both lipid classes

however were depleted by about 50 % in this organelle in comparison to L-TE. PS (18:0/18:2)

composes up to 38.3 % in L-mito with an increase of $2^{4.3}$ fold comparing to L-TE. The

accumulation of this lipid species was the most remarkable one that was even visible when the

fatty acid profiles of the different lipid classes were calculated from the molecular species data,

showing an enrichment of 18:0 in mitochondria on the expense of 16:0 (Supplemental Fig. S3).

Together the main molecular species of these three lipid classes have a C18/C18 acyl chain

composition in common.

**Figure 3.** Profiles of the distribution and fold changes of the molecular glycerolipid species

between L-mito and L-TE. Heat map visualizations of (a) glycerophospholipids, (b)

glyceroglycolipids and (c) diacylglycerols illustrate the difference of species distribution based on

LC-MS/MS analyses. Each lipid class is represented by one set of joined columns, only DAG is

divided into two sets due to space constraints. Identity of columns in each set from left to right:

Column 1 lists the individual lipid class and the identity of the detected molecular lipid species.

Column 2 (L-TE) lists the respective distribution of each molecular species in L-TE, expressed as

the mean of its relative values in mol % in the three independent experiments also used for

proteomics and for sphingolipid and sterol analysis (Figs. 1, 5 and 6, Supplemental Tab. S1).

Column 3 (L-mito) lists the respective distribution of each molecular species in L-mito, expressed

as the mean of its relative values (mol %) in the three independent experiments also used for

proteomics. Column 4 lists the mean distribution of both sample types, and column 5 ($Log_2(FC)$)

lists the binary logarithm of fold change. Binary logarithm was applied when the mean values

are higher than 0.5 to inspect the fold changes (FC) between L-mito and L-TE. The heat map

colors represent mean intensity values according to the color map on the low right-hand side.

One and two asterisks (*, **) indicate p values < 0.05 and < 0.01, respectively, by Student's t-

test. PC, phosphatidylcholine; PE, phosphatidylethanolamine; PS: phosphatidylserine; PI:

phosphatidylinositol; PA phosphatidic acid; DAG: Diacylglycerol; CL, cardiolipin; PG,

phosphatidylglycerol; MGDG, monogalactosyldiacylglycerol; DGDG, digalactosyldiacylglycerol.

An important intermediate in glycerolipid metabolism is phosphatidic acid (PA), serving as

precursor or product for numerous glycerolipids including PC, PE, PG, phosphatidylinositol (PI)

and DAG. PA (16:0/18:2), PA (16:0/18:3) and PA (18:2/18:3) are the most abundant PA species in

both fractions, L-mito and L-TE. Notably, only a minor molecular PA species (16:0/16:0; 1.3 mol

%) accumulates preferentially in mitochondria with an enrichment of $2^{5.3}$ fold comparing to L-TE.

DAG as the other important lipid intermediate displays a higher complexity in its species profile.

DAG (16:0/18:2), DAG (16:0/18:3), DAG (18:2/18:3) and DAG (18:3/18:3) build a substantial

amount (>65 mol %) of the DAG profile in L-mito (Fig. 3c). As in case of PA, only the minor DAG

species (16:0/16:0) shows a $2^{2.2}$ fold increase in L-mito compared to L-TE.

PG is formed from PA in the envelopes of plastids and mitochondria and serves as precursor for

the mitochondrial lipid CL (Babiychuk et al., 2003). Hydrolysis and condensation of PG lead to

the formation of DAG and CL correspondingly. PG (16:0/18:2) and PG (16:0/18:3) are the major

species in L-mito, but only the former species is enriched in mitochondria ($2^{1.1}$ fold). In addition,

unsaturated PG species are significantly enriched in L-mito, which contribute to the structures of

the down-stream CL molecules. CL (72:10) and CL (72:11), composed of 18:2 and 18:3 acyl

chains, are the most abundant CL (Fig. 3a).

Glycerophospholipids serve not only as membrane building blocks but are also important

precursors for signaling molecules in the cell. PI, phosphatidylinositol monophosphate (PIP) and

phosphatidylinositol bisphosphate ($PIP_2$) exert regulatory functions in cell development and

polarity determination (Heilmann, 2016). In L-mito, molecular species with 16:0/18:2 and

16:0/18:3 acyl chains compose up to 80 % of PI and $PIP_2$; yet PIP was not detectable in L-mito.

PI (16:0/16:1) and PI (18:0/18:1) were only detectable in the mitochondrial extracts, and

PI (18:0/18:2) is significantly enriched in L-mito for $2^{1.2}$ fold in comparison to L-TE (Fig. 3a). On

the other hand, we obtained higher signals of $PIP_2$ (18:1/18:2) in L-mito.

Glyceroglycolipids carry carbohydrate residues as their head groups; for instance, the galactose-

containing lipids, MGDG and DGDG contain one and two galactoses, respectively (Hölzl and

Dörmann, 2019). MGDG (16:3/18:3) and DGDG (18:3/18:3) are the major components of the

overall glyceroglycolipids in both L-TE and L-mito (Fig. 3b). The amount of MGDG (16:0/18:3)

and MGDG (18:2/18:2) are specifically elevated in L-mito for $2^{1.5}$ and $2^{1.4}$ folds, respectively,

compared to L-TE. In contrast, the species MGDG (16:1/18:1) and MGDG (16:3/18:2) are

significantly decreased in L-mito. In summary, the major mitochondrial lipids PC, PE, PS and CL

consist primarily of C18/C18 species, while PI, PIP2 and PG consist primarily of C16/C18 species

and the lipid precursors DAG and PA resemble a mixture of both backbones.

**Proteomic analyses provides insights into the leaf mitochondrial glycerolipid metabolism**

**capacity**

To further analyze the proteome of the mitochondrial fractions listed in Supplemental Tab. S1,

their protein composition was investigated for enzymes from lipid metabolism. Subsequently,

their biological function and the subcellular localization were retrieved from The Arabidopsis

Information Resource (TAIR) and the Subcellular localization database for Arabidopsis proteins

(SUBAcon) databases (Supplemental Fig. S4).

About 40 proteins involved in the biosynthesis and modification of fatty acids and more complex

lipids were identified in mitochondrial extracts (Supplemental Tab. S3). Four enzymes from

glycerophospholipid metabolism were detected in all mitochondrial fractions (Fig. 4). Two

glycerol-3-phosphate dehydrogenases were identified (SDP6 and GPDHp), being involved in an

early step of glycerophospholipid biosynthesis. Phosphoethanolamine cytidylyltransferase (PECT)

was detected in our study by similar abundances in the L-mito and C-mito samples. CL is

synthesized after the condensation of PG and CDP-DAG by CL synthase (CLS) or after the

transacylation of monolyso-CL by monolysocardiolipin acyltransferase (LCLAT or Tafazzin)(Xu et

al., 2006). CLS and Tafazzin were identified in the shotgun proteomic analysis in this study with a

higher abundance in C-mito compared to L-mito. PG is synthesized from PA via CDP-DAG by

phosphatidylglycerophosphate synthase 1 (PGP1) (Xu et al., 2002) and it was only detected in C-

mito samples. Phosphatidylserine decarboxylase 1 (PSD1) synthesizes PE by decarboxylating the

headgroup of PS (Birner et al., 2001; Nerlich et al., 2007). Again in our study PSD1 was only

detected in C-mito samples. Interestingly PE and PC can be interconverted into each other via

PA and the detected phospholipase Dα1 (PLDα1) enzyme activity. PLDα1 was preferentially

found in C-mito samples. PIP and $PIP_2$ can be degraded by *myo*-inositol polyphosphate 1-

phosphatase (SAL1). Although many enzymes involved in phosphoinositide metabolism have

been annotated to localize in mitochondria, only SAL1 was identified in our approach in the C-

mito samples.

## a Glycerophospholipid metabolism

| L-mito | | | C-mito | | | Locus | Name | Function |
|---|---|---|---|---|---|---|---|---|
| | | | | | | AT2G38670 | PECT1 | Ethanolamine-phosphatecytidylyltransferase |
| | | | | | | AT2G39290 | *PGP1* | CDP-DAG-G3P3-phosphatidyltransferase1 |
| | | | | | | AT3G10370 | SDP6 | Glycerol-3-phosphatedehydrogenase |
| | | | | | | AT3G15730 | PLDα1 | PhospholipaseDalpha1 |
| | | | | | | AT4G04870 | CLS | Cardiolipinsynthase |
| | | | | | | AT4G16700 | PSD1 | PSdecarboxylaseproenzyme1 |
| | | | | | | AT1G78690 | LCLAT | Monolysocardiolipinacyltransferase |
| | | | | | | AT5G40610 | GPDHp | Glycerol-3-phosphatedehydrogenase |

## b Glyceroglycolipid metabolism

| L-mito | | | C-mito | | | Locus | Name | Function |
|---|---|---|---|---|---|---|---|---|
| | | | | | | AT4G33030 | *SQD1* | Sulfoquinovosyldiacylglycerol1 |

## c Phosphoinositide metabolism

| L-mito | | | C-mito | | | Locus | Name | Function |
|---|---|---|---|---|---|---|---|---|
| | | | | | | AT5G63980 | SAL1 | Inositolpolyphosphate1-phosphatase |

**Figure 4.** Mitochondrial localized proteins that are related to glycerolipid biosynthesis. (a) Glycerophospholipid, (b) glyceroglycolipid and (c) phosphoinositide metabolism. Proteins identified and/or localized in mitochondria were labeled (i) blue: proteins identified in the proteomic analysis of this study and also predicted to localize in mitochondria, (ii) green: proteins identified in the proteomic analysis of this study in mitochondria but predicted to localize in other organelles, (iii) bold font: exclusively localized in mitochondria and (iv) italic font: only identified in one of the mitochondrial populations. Heat maps visualize the protein abundance of three independent experiments of mitochondria purified from leaves and cell cultures from Fig. 1 and listed in Supplemental Tabs. S1 and S3. The heat map colors represent mean intensity values according to the color map on the top right-hand side. Predicted protein localization was based on The Arabidopsis Information Resource (TAIR; www.arabidopsis.org) and the Subcellular localization database for Arabidopsis proteins (SUBAcon; www.suba.live).

The biosynthesis of glyceroglycolipids takes place in the plastids envelopes by MGDG and DGDG

synthases (MGDs and DGDs), respectively (Hölzl and Dörmann, 2019). MGDG is generated by

adding a galactose head group to DAG; subsequent addition of another galactose by DGD1

generates DGDG. The detected specific species profiles of mitochondrial glyceroglycolipids may

suggest at least for a specific transport from plastids to mitochondria (Fig. 3b). The sulfolipid

SQDG is synthesized in two steps. Uridine diphosphate (UDP)-glucose is first combined with a

sulfite by UDP-sulfoquinovose synthase 1 (SQD1), followed by transferring the sulfoquinovose

head group to DAG and thus forms SQDG. Although SQD1 was identified in the C-mito samples

in this study, we did not detect SQDG lipids in any of the mitochondrial samples. Together we

could exclusively localize three biosynthetic steps to mitochondria: (i) CLS and catalyzing the

formation of CL as well as (ii) PSD1 and (iii) PECT1 facilitating PE formation.

**Specific molecular sphingolipid and sterol species suggest a function in mitochondria**

Sphingolipids and sterols are important modulators of membrane microdomains and play

critical roles in regulating the balance between cell survival and apoptosis. Therefore, we

profiled next their mitochondrial compositions in comparison to the total lipidome of leaves in

the same samples that were used for proteomics and glycerolipid analysis (Figs. 1 and 3) by LC-

MS/MS (Figs. 5 and 6) and expressed the lipid profiles in mol %. For sphingolipids both simple

and complex sphingolipids were measured, including long-chain bases (LCB), phosphorylated

LCB (LCB-P), ceramides (Cer), glucosylceramides (GlcCer) and glycosyl inositol

phosphoceramides (GIPC) (Fig. 5). All sphingolipids have LCBs, 18-carbon amino-alcohols, as

their backbones. Phytosphingosine (18:0;3), hosting three hydroxyl groups, is the most

abundant free LCB in both L-mito and L-TE (Fig. 5a). However, dihydrosphingosine (18:0;2) is

highly enriched in L-mito as well as phosphorylated dihydrosphingosine (18:0;2-P), which is the

major component in the LCB-P pool of L-mito (64.7 %). LCBs can be further *N*-acylated to

generate Cer, the basic structure of complex sphingolipid classes. Addition of glucoses to Cer

generates GlcCer, following sequential extension of phosphoinositol, hexose and/or hexose

derivatives generate series of GIPCs. Series 0, A and B GIPCs carry one, two and three additional

hexoses on the head groups of inositol phosphoceramides, respectively (Cacas et al., 2012;

Haslam and Feussner, 2022). Similar to LCB and LCB-P, only specific species were enriched in L-

mito (Fig. 5b): Cer (18:0;3/24:0;1), GlcCer (18:1;2/24:1;1) and series A hexose-carrying GIPC (H-

GIPC) (18:1;3/24:1;1). Remarkably, GIPCs were only detectable in mitochondrial samples (both

L-mito and C-mito) in our approach. However, proteins related to sphingolipid metabolism were

neither identified in plant mitochondria via proteomic analyses, nor retrieved from online

databases.

**Figure 5.** Profiles of the distribution and fold changes of the molecular sphingolipid species between L-mito and L-TE. Heat map visualizations of (a) long chain bases, long chain base-phosphates and (b) complex sphingolipids illustrate the difference of species distribution based on LC-MS/MS analyses. Each lipid class is represented by one set of joined columns. Identity of

columns in each set from left to right: Column 1 lists the individual lipid class and the identity of

the detected molecular lipid species. Column 2 (L-TE) lists the respective distribution of each

molecular species in L-TE, expressed as the mean of its relative values (mol %) in the three

independent experiments also used for proteomics and for glycerolipid and sterol analysis

(Figs. 1, 3 and 6, Supplemental Tab. S1). Column 3 (L-mito) lists the respective distribution of

each molecular species in L-mito, expressed as the mean of its relative values (mol %) in the

three independent experiments also used for proteomics. Column 4 lists the mean distribution

of both sample types, and column 5 (Log$_2$(FC)) lists the binary logarithm of fold change. Binary

logarithm was applied when the mean values are higher than 0.5 to inspect the fold changes (FC)

between L-mito and L-TE. The heat map colors represent mean intensity values according to the

color map on the low left-hand side. One and two asterisks (*, **) indicate p values < 0.05 and <

0.01, respectively, by Student's t-test. LCB, long-chain base; LCB-P, long chain base-phosphate;

Cer, ceramide; GlcCer, glycosylceramide; GIPC, glycosyl inositol phosphoceramide.

The common structure of sterols is a four-ring system, cyclopentanoperhydrophenanthrene,

with possible conjugation of hydroxyl groups and acyl chains. In plants, a complex mixture of

sterols can be found, including brassicasterol, campesterol, cholesterol, sitosterol and

stigmasterol (Cacas et al., 2012). Campesterol was identified as major free sterol with 61.7 % in

L-mito (Fig. 6). From the group of steryl glycosides cholesteryl and isofucosteryl glycoside

accumulated preferentially in L-mito. Considering SE, 16:0 and 18:0 containing cholesteryl esters

were enriched $2^{3.4}$ and $2^{5.3}$ folds, respectively, in L-mito. For ASG however the situation was

blurred. DWF1 was the only protein related to sterol metabolism that we identified in plant

mitochondria via proteomic analyses. In summary, we observed an accumulation of GIPCs and

free campersterol in leaf mitochondria.

**Figure 6.** Profiles of the distribution and fold changes of the molecular sterol species between L-mito and L-TE. Heat map visualizations of (a) free sterols (b) steryl glycosides (SG) and (c) acylated steryl glycosides (ASG) and steryl esters (SE) illustrate the difference of species distribution based on LC-MS/MS analyses. Each lipid class is represented by one set of joined

columns. Identity of columns in each set from left to right: Column 1 lists the individual lipid

class and the identity of the detected molecular lipid species. Column 2 (L-TE) lists the respective

distribution of each molecular species in L-TE, expressed as the mean of its relative values (mol

%) in the three independent experiments also used for proteomics and for glycerolipid and

sphingolipid analysis (Figs. 1, 3 and 5, Supplemental Tab. S1). Column 3 (L-mito) lists the

respective distribution of each molecular species in L-mito, expressed as the mean of its relative

values (mol %) in the three independent experiments also used for proteomics. Column 4 lists

the mean distribution of both sample types, and column 5 (Log$_2$(FC)) lists the binary logarithm of

fold change. The heat map colors represent mean intensity values according to the color map on

the low left-hand side. Binary logarithm was applied when the mean values are higher than 0.5

to inspect the fold changes (FC) between L-mito and L-TE. One and two asterisks (*, **) indicate

p values < 0.05 and < 0.01, respectively, by Student's t-test.

Mitochondria harbor various additional metabolic pathways for the synthesis of lipophilic

molecules. More than 20 fatty acid biosynthesis-related proteins were identified in our

mitochondrial proteome (Supplemental Fig. S5). They contribute to the synthesis of lipoic acid,

ubiquinone and other terpenoid-quinones. Abundance, function and predicted localization of

these proteins are specified in Supplemental Tab. S1 and S3.

**Lipid molecules can be imported by mitochondria**

Transport of DGDG from plastids to mitochondria and reallocation of PE from IM to OM via the

MTL complex have been demonstrated under phosphate-depleting conditions (Michaud et al.,

2016). The MTL complex has more than 200 subunits and 11 have been verified to associate

physically with the core subunits, Mic60 and Tom40, as previously shown by immunoblotting

(Michaud et al., 2016; Li et al., 2019). In our mitochondrial proteome listed in Supplemental Tab.

S1, 186 of the subunits (87 %) were identified including all the 11 verified ones (Fig. 7,

Supplemental Tab. S4). The newly identified MTL subunit, DGS1, which links mitochondrial

protein to lipid transport, is present in all mitochondrial extracts.

| L-mito | C-mito | Locus | Name | Description |
|---|---|---|---|---|
| | | AT1G06530 | PMD2 | Peroxisomal and mitochondrial division factor 2 |
| | | AT1G27390 | TOM20-2 | Translocase of outer membrane 20 kDa subunit 2 |
| | | AT2G37410 | TIM17-2 | Translocase inner membrane subunit 17-2 |
| | | AT2G42210 | B14.7 | Mitochondrial import inner membrane translocase subunit family protein |
| | | AT3G20000 | TOM40 | Translocase of the outer mitochondrial membrane 40 kDa |
| | | AT3G27080 | TOM20-3 | Translocase of outer membrane 20 kDa subunit 3 |
| | | AT4G39690 | Mic60 | Mitochondrial contact site and cristae organizing system subunit |
| | | AT5G12290 | DGS1 | dgd1 suppressor 1 |
| | | AT5G13430 | RISP | Ubiquinol-cytochrome c reductase iron-sulfur subunit |
| | | AT5G40930 | TOM20-4 | Translocase of outer membrane 20 kDa subunit 4 |
| | | ATMG00160 | COX2 | Cytochrome oxidase 2 |

| L-mito | C-mito | Locus | Name | Function |
|---|---|---|---|---|
| | | AT3G19820 | DWF1 | Delta(24)-sterolreductase |

Low High

**Figure 7.** MTL complex-associated proteins in mitochondria. Our proteomic analyses covered 186 of the 214 hypothetical subunits in the MTL complex (listed in Supplemental Tab. S4); 11 have been investigated by immunoblotting approaches in previous studies. Heat maps visualize the protein abundance of three independent experiments of mitochondria purified from leaves and cell cultures. The heat map colors represent mean intensity values according to the color map on the low right-hand side.

In addition, at least 10 outer envelope (OE)-localized proteins from plastids were co-purified with mitochondria, with higher abundance in C-mito (Fig. 8, Supplemental Tab. S5). They account for 2.1 % (10/471) of the identified plastidial proteins, which are enriched in comparison to the proportion of OE-localized proteins in plastids (46/3002, 1.5 %) (Inoue, 2007, 2011; Simm et al., 2013; Kim et al., 2019). The enrichment of OE-localized proteins in mitochondrial extracts suggest a close physical interaction between mitochondria and plastidial OE membrane. For instance, the membrane contact sites, which were not disrupted during the

432    isolation procedure, and therefore are present in the mitochondrial samples as membrane

433    patches.

| L-mito | | | C-mito | | | Locus | Name | Function |
|---|---|---|---|---|---|---|---|---|
| | | | | | | AT2G16640 | TOC132 | Translocon complex in the outer envelope membrane 132 |
| | | | | | | AT2G28900 | OEP16-1 | Outer plastid envelope protein 16-1 |
| | | | | | | AT2G43950 | OEP37 | Outer envelope protein 37 |
| | | | | | | AT3G06510 | GGGT | Galactolipid:galactolipid galactosyltransferase |
| | | | | | | AT3G17970 | TOC64-III | Translocon at the outer membrane of chloroplasts 64-III |
| | | | | | | AT3G46740 | TOC75-III | Translocon at the outer envelope membrane of chloroplasts 75-III |
| | | | | | | AT3G52420 | OEP7 | Outer envelope membrane protein 7 |
| | | | | | | AT4G02510 | TOC159 | Translocon at the outer envelope membrane of chloroplasts 159 |
| | | | | | | AT5G05000 | TOC34 | Translocon at the outer envelope membrane of chloroplasts 34 |
| | | | | | | AT5G19620 | OEP80 | Outer envelope protein of 80 kDa |

Low            High

**Figure 8.** Identified plastid outer envelope proteins in mitochondrial extracts. Our proteomic

analyses identified 10 outer envelope-localized proteins (listed in Supplemental Tab. S5). Heat

maps visualize the protein abundance of three independent experiments of mitochondria

purified from leaves and cell cultures. The heat map colors represent mean intensity values

according to the color map on the left-hand side.

**Discussion**

In this study, we present an in-depth dataset of lipid molecular species from glycerolipids,

sphingolipids and sterols of Arabidopsis leaf mitochondria. With the assistance of online

resources, we assigned and confirmed central lipid biosynthesis steps within these

mitochondrial fractions. Furthermore, we confirmed and expanded our knowledge on the

existence of a protein complex for lipid trafficking between mitochondria and other organelles.

In the past, the most abundant plant mitochondrial glycerolipid classes, such as PC, PE, PI, PA, CL,

MGDG and DGDG, from *Arabidopsis* cell cultures and calli under phosphate-depleted conditions

have been quantified with TLC-GC (Jouhet et al., 2004; Michaud et al., 2016). However, a TLC-

GC-based approach lacks the molecular species information concerning acyl chain length and

unsaturation degree of the lipids, which are critical parameters to determine the membranes

physical properties. In addition, neither sphingolipids nor sterols of mitochondria were profiled

to our very best knowledge formerly in *Arabidopsis*.

In our study, mitochondria were purified from leaves by differential and Percoll density gradient

centrifugations and compared with the well established mitochondrial preparations from

suspension cell culture (Jouhet et al., 2004; Klodmann et al., 2011; Behrens et al., 2013;

Michaud et al., 2016). Much effort was made to document purity of the resulting organelle

fraction: (i) visual inspection of the protein complex composition by 2D BN / SDS PAGE and (ii)

label-free quantitative shotgun proteins in combination with SUBAcon evaluation. The SUBAcon

database integrates worldwide knowledge on subcellular localization information of Arabidopsis

proteins based on *in vitro* or *in vivo* protein targeting experiments, mass spectrometry-based

analyses of organellar fractions, protein-protein interaction data and bioinformatics tools for

subcellular localization prediction. Based on the latter approach, purity of our mitochondrial

fractions can be estimated to be in the range of 90% (87 to 94%, Supplemental Fig. S2). Traces of

chloroplasts were present in our fractions. However, even Rubisco, which is considered to be

the most abundant chloroplast protein, was detectable only as a very faint spot on the 2D

BN/SDS gel of the L-mito fraction (Fig. 1). Finally, quantitative lipid analysis revealed the purity

of our mitochondrial fraction. MGDG, DGDG, PG and SQDG are the main components of

plastidial membranes (Hölzl and Dörmann, 2019). In *Arabidopsis* chloroplasts, the ratio of

MGDG : DGDG : PG : SQDG is about 1:0.5:0.1:0.3 (Awai et al., 2006). In contrast, a distinct ratio

of 1:0.3:0.4:0 was detected in L-mito in this study (Fig. 2). Moreover, only a few specific MGDG

and DGDG molecular species are enriched in L-mito compared to L-TE (Fig. 3B), suggesting that

the cross-contamination from intact plastids or bulk plastidial membranes are insignificant.

Interestingly a similar observation was recently made when tagged mitochondria were isolated

by affinity purification (Niehaus et al., 2020). However, in previous studies on mitochondria

isolated from cell cultures the amount of MGDG was much lower (~1.4 mol % vs. ~15 mol % in

Fig. 2) than in the mitochondrial fraction analyzed here from leaves (Jouhet et al., 2004;

Michaud et al., 2016). This can only be explained by a higher number of membrane contacts

between plastids and mitochondria in leaves as has been observed before in case of phosphate

starved cell cultures (Jouhet et al., 2004). On the other hand, more than 85 % of the subunits of

the MTL complex (deduced from its composition in yeast mitochondria, Fig. 7) and many OE-

localized proteins (Fig. 8) were identified in the mitochondrial samples via our proteomics

approach, strongly suggesting that a close contact between mitochondria, the ER and

chloroplasts exists in our preparations as has been observed before in *Arabidopsis* cell cultures

(Jouhet et al., 2004; Michaud et al., 2016). We conclude that the identified chloroplast lipids and

proteins rather originate from small pieces of plastidial membranes and, importantly, the

mitochondria-plastid contact sites, which are present in our mitochondrial fraction and

contribute to the measured lipid composition. While the protein complexes connecting ER and

mitochondria have been well described in yeast, their roles in mediating and/or facilitating lipid

translocation are less defined in plants. In contrast to yeast, to our knowledge only one

tethering protein has been described in flowering plants up to now (Michaud et al., 2016).

With the proteomic and lipidomic datasets as well as the online resources in hand, we assigned

central lipid biosynthesis reactions and refined models for a possible exchange of lipid

molecules between mitochondria, ER and plastids in plants in Fig. 9. Previous studies in yeast

and mammalian cells have shown that mitochondria are capable of synthesizing PE, PA, PG and

CL (Flis and Daum, 2013; Horvath and Daum, 2013; Tatsuta et al., 2014). Here, we expand the

knowledge that at least the glycerolipids PE and CL can be generated and/or modified in

Arabidopsis leaf mitochondria. Considering PE biosynthesis, the rate-limiting enzymes in CDP-

ethanolamine and PS decarboxylation pathways, PECT1 and PSD1, respectively, were identified

in our mitochondrial samples with higher abundance in C-mito (Fig. 4). It suggests that the

generation of mitochondrial PE in the cell culture may be more active in comparison to the

leaves. Moreover our observation was supported by earlier studies that localized PECT1 at the

mitochondrial periphery and PSD1 in mitochondria in *Arabidopsis* (Mizoi et al., 2006; Nerlich et

al., 2007). PE is one of the most abundant glycerophospholipids in biological membranes.

Therefore, a high demand is expected to supply mitochondria for their biogenesis in actively

dividing cell cultures. The other major component of biological membranes is PC. It is

considered to be synthesized in the ER and then transferred through protein complexes

connecting ER and mitochondria to mitochondria. The structural information of the enriched

lipid species provides evidences for the biosynthesis in the ER as well. Many PE and PC species in

mitochondrial samples have longer acyl moieties with 22 to 26 carbons, which can only be

synthesized in the ER (Haslam and Kunst, 2013). Conventionally, the majority of PS biosynthesis

was assumed to take place in the ER and before being transferred to mitochondria as well. PS

and PE, although they are able to interconvert, have distinct lipid profiles from each other,

suggesting that only selected lipid species are the substrates of PSD1. In addition, inositol

polyphosphate phosphatase 1 (SAL1), which is an enzyme that is involved in phosphoinositide

degradation and generates inositol phosphates, was identified in our mitochondrial samples

with higher abundance in C-mito comparing to L-mito. This finding confirmed a previous report

that localized it in both mitochondria and plastids (Estavillo et al., 2011). Inositol phosphates

play crucial roles in many biological processes including gene expression and regulation of cell

death through sphingolipids (Alcázar-Román and Wente, 2008; Donahue et al., 2010). Removal

of head groups from glycerophospholipids, mostly PC and PE, by phospholipase Dα1 (PLDα1)

results in PA, serving as important precursor in other glycerophospholipid biosynthesis

pathways (Fig. 9). Comparing to L-TE, the most enriched PA species in L-mito carry 16:0 and 18:0

fatty acyl moieties. This corresponds to the enriched PI, PG and DAG species in L-mito,

suggesting that PA may serve as an interconverting hinge between these glycerolipids in plant

mitochondria. While our study detected PLDα1 preferentially in C-mito samples, previous

studies localized it in the cytosol or at the plasmamembrane (Du et al., 2013). This may suggest

that PLDα1 localizes to contact sites either between plasmamembrane and mitochondria or

between ER and mitochondria. In addition, two glycerol-3-phosphate dehydrogenases were

identified (SDP6 and GPDHp). Whereas SDP6 was detected in mitochondria before, GPDHp was

localized in the plastid (Nandi et al., 2004; Quettier et al., 2008).

**Figure 9.** Model of lipid biosynthesis and trafficking within and between mitochondria, ER and

plastids in *Arabidopsis*. Lipid synthesis and transfer between membranes are indicated by black

and blue arrows, respectively. Proteins identified in L-mito or C-mito with additional ER or

plastidic localization based on The Arabidopsis Information Resource (TAIR;

www.arabidopsis.org) and the Subcellular localization database for Arabidopsis proteins

(SUBAcon; www.suba.live) are considered as putative contact-site localized proteins (dashed

frame). Lipid biosynthesis-related proteins indicated in other studies are depicted in grey. Full

names and functions of involved proteins are itemized in Supplemental Tab. S1. OM,

mitochondrial outer membrane; IM, mitochondrial inner membrane; OE, plastid outer envelope;

IE, plastid inner envelope; MTL, mitochondrial transmembrane lipoprotein complex. DHAP,

dihydroxyacetone phosphate; G3P, glyceraldehyde 3-phosphate; MAG, monoacylglycerol; DAG,

diacylglycerol; Etn, ethanolamine; TAG, triacylglycerol; IP, inositol phosphate; PGP,

phosphatidylglycerol phosphate; SQDG, sulfoquinovosyl diacylglycerol.

An enzyme being involved in synthesizing PG from PA is PGP1. It was found in our proteomic

analysis in C-mito samples (Fig. 4). CL can be synthesized through condensation of PG and DAG

by CLS, or addition of acyl chains to monolyso-CL by monolysocardiolipin acyltransferase (LCLAT

or Tafazzin). All identified enzymes involving in PG and CL biosynthesis, PGP1, CLS and LCLAT,

are more abundant in C-mito compared to L-mito. PGP1 was previously found to localize to both

mitochondria and plastids (Babiychuk et al., 2003), while CLS was localized to mitochondria

(Katayama et al., 2004). This suggests an intensive interaction between mitochondria and

plastids in cell cultures because (1) high amounts of proteins localized in contact sites imply a

closer connection between the two organelles; (2) more CLS is required to convert the plastid-

derived PG which is transferred to mitochondria presumably via membrane contact sites; (3)

active energy generation is vital for actively dividing cells and thus a high amount of CL is

required to assemble the respiratory chain complexes in mitochondria. This is further supported

by the specific PG species found in L-TE. L-TE contains substantial amounts of 16:1 (probably

$trans$-$\Delta^3$-16:1)/18:X PG species (~35 mol %) (Fig. 3), which are molecular species synthesized

entirely within plastids and probably exclusively localized there (Hölzl and Dörmann, 2019).

Therefore, it is very likely that the high amounts of these PG species in L-TE can be attributed to

the abundance of plastid membranes particularly thylakoids. Interestingly, the L-mito fraction

also contained significant levels (~8 mol %) of 16:1/18:X PG species. These PG species may be

not in mitochondria but be derived from the mitochondrion-plastid contact sites or

contaminations of plastid membranes (Fig. 9). Nevertheless PG which is directly after its uptake

converted into CL in mitochondria (Figs. 2 and 9).

Typically, glyceroglycolipids such as MGDG, DGDG and SQDG are considered to be synthesized in

plastids and are transferred to other organelles upon stress. For instance, DGDG is transferred

from plastidial membranes to other compartments including mitochondrial membranes and the

plasma membrane during phosphate starvation to compensate the loss of PC and PE (Jouhet et

al., 2004). In addition to vesicular transportation and lipid trafficking via contact sites, emerging

evidences support the hypothesis of lipid synthesis in $trans$ in yeast and plants (Mehrshahi et al.,

2013; Tavassoli et al., 2013; Michaud et al., 2017). That is, enzymes located at one membrane

might be capable of catalyzing a reaction on another membrane when in close proximity. In this

way, neither tethering proteins nor massive lipid remoling under stress conditions is required,

if the mitochondria can acquire lipids without setting up the junctions to other organelles.

Therefore, we suspect that the DGDG biosynthesis enzymes, DGD1/2, catalyze the reactions in

$trans$ or at the mitochondria-plastid contact sites, providing DGDG molecules to compensate the

loss of glycerophospholipids in real time (Fig. 4b). In addition, several subunits of the MTL

complex were identified in our proteomic approach including Tom40, Mic60, DGS1 etc. (Fig. 7).

It establishes a lipid trafficking system removing DGDG from plastids and allocating PE between

the IM and OM. Rapid remodeling of the mitochondrial membranes is supported by in *trans*

lipid biosynthesis and the MTL complex, likely in close cooperation with other complexes.

However, the underlying mechanism of this lipid transportation machinery is still largely

unknown. SQDG synthase (SQD1) was identified exclusively in C-mito, but no SQDG molecules

were detected (Fig. 4b). Earlier reports localized SQD1 in chloroplasts (Essigmann et al., 1998),

suggesting that SQD1 is either co-purified with plastid envelope membranes from cell cultures

or is localized in .

Sphingolipid biosynthesis takes place at the ER membrane and subsequently in the Golgi

apparatus for the addition of carbohydrate residues. Interestingly, in the last decade, several

sphingolipid-metabolizing enzymes have been identified in mitochondria purified from yeast

and mammalian systems, including Cer synthase and ceramidase (Bionda et al., 2004; Kitagaki et

al., 2007; Novgorodov et al., 2011). In animal models, mitochondria-synthesized Cer plays a

crucial role in cerebral ischemia-induced mitochondrial dysfunction. The connection between

sphingolipids and mitochondria-promoted apoptosis has been proposed in plants as well.

Recent studies in *Arabidopsis* have demonstrated that the ratio between LCB-P and Cer is

involved in maintaining the balance of cell survival and apoptosis (Watanabe et al., 2013; Bi et

al., 2014). However, no sphingolipid biosynthesis enzyme has been identified in plant

mitochondria hereto. GIPCs are the most abundant sphingolipids in mitochondria, although still

minor comparing to glycerophospholipids. It is known that PM-localized GIPCs are important in

signal transduction and intercellular recognition (Lenarčič et al., 2017; Ali et al., 2018). GIPCs

may as well establish the communication between mitochondria and other organelles, although

further analysis is required to understand the functions of sphingolipids in plant mitochondria.

Sterols, although taking part in both biotic and abiotic stress responses, are at low abundance in

mitochondrial samples. Sterol biosynthesis primarily takes place in the ER (Schaller, 2003).

Among them, sterol C-24 reductase (DWF1) was identified both in L-mito and C-mito, with

higher abundance in C-mito. DWF1 has been proposed to mediate the biosynthesis of all

phytosterols with higher specificity towards campesterol in Arabidopsis, but this protein was

localized to ER membranes (Klahre et al., 1998; Youn et al., 2018). Therefore, we suggest DWF1

as an additional contact site localized protein (Fig. 7). High amount of campesterol was detected

in L-mito, suggesting the existence of an onsite biosynthesis and/or sterol transporter to

facilitate the import of campesterol from the ER. However, unlike mammalian cells wherein

cholesterol transport proteins such as steroidogenic acute regulatory protein, StAR (Clark et al.,

1995), and MLN64 (Charman et al., 2010) have been identified, little is known about sterol

transporters in plants.

**Conclusions**

In summary, we expanded the knowledge regarding lipid biosynthesis and modification in plant

mitochondria by performing a global lipidome analysis of Arabidopsis leaf mitochondria to

provide their in-depth lipid molecular species profile including glycerolipids, sphingolipids and

sterols. Our proteome data confirmed that PE and CL can be synthesized in these organelles

partially by the assistance of putative contact-site localized proteins and / or in *trans* lipid

biosynthesis. Based on the proteomic results, we propose and confirm the existence of

membrane contact site-localized proteins and their aspects in lipid biosynthesis pathways. This

study serves as a foundation for additional researches in unveiling the functional roles of

mitochondrial lipids and the mechanisms of mitochondria-dependent signaling pathways.

**Materials and methods**

**Plant materials and growth conditions**

Rosette leaves of wild-type *Arabidopsis thaliana* (L.) Heynh Columbia-0 were used for both

extracting total lipid extract and purifying mitochondria. After sowing the seeds in pots with

three-day cold stratification at 4 °C, seedlings were grown under 16 h-day length at 24 °C with

60 % relative humidity and 150 µmol photons $m^{-2}$ $sec^{-1}$ for one week. Young seedlings were

transferred to large trays and grown with equal spacing for another three weeks before further

experimental procedures.

**Suspension culture of *Arabidopsis thaliana***

Suspension cultures were established starting from sterilized seeds of *Arabidopsis thaliana* wild-

type Columbia-0, grown on MS medium plates containing 0.8 % agar. Plant pieces were

transferred to B5 medium agar plates and cultivated for several weeks in the dark for callus

induction. Callus was finally transferred into liquid B5 medium including 3 % (w/v) sucrose,

0.01 % (w/v) 2,4-dichlorophenoxyacetic acid and 0.001 % (w/v) kinetin. Cultures were incubated

on a shaker at 24 °C in the dark. Callus was transferred weekly into new liquid medium (3 g /

100 ml).

**Mitochondria isolation**

*From Arabidopsis rosettes*

The mitochondria isolation procedure was described previously (Schikowsky et al., 2018). About

200 g of four-week-old *Arabidopsis* rosettes were collected and homogenized at 4 °C with a

waring blender in 1 liter disruption buffer (0.3 M sucrose, 60 mM TES, 25 mM tetrasodium

pyrophosphate, 10 mM potassium dihydrogen phosphate, 2 mM EDTA, 1 mM glycine, 1 %

PVP40, 1 % BSA, 50 mM sodium ascorbate, 20 mM cysteine; pH 8.0) by three times for 10 sec

with 30 sec intervals. The following procedures were performed on ice or at 4 °C. Two layers of

miracloth with supporting gauze were used to filter the homogenate into a beaker. The

remaining plant debris was first grinded with additional sea sand for 10 min by mortar and

pestle, and then filtered again through miracloth. The filtrates were combined and centrifuged

at 2,500 g for 5 min to eliminate the cell debris. Centrifugation with higher speed at 15,250 g for

15 min was applied on the supernatant to pellet mitochondria and other organelles. The

resulting pellets were resuspended with a paintbrush in wash buffer (0.3 M sucrose, 10 mM TES,

10 mM potassium dihydrogen phosphate; pH 7.5). The samples were adjusted to the final

volume of 12 ml with wash buffer and transferred to a Dounce homogenizer. Two strokes of

pestle were performed to disrupt large organelles like chloroplasts. Aliquots of 1 ml samples

were transferred carefully to Percoll gradients which had been established beforehand by

69,400 g centrifugation for 40 min in one-to-one ratio of Percoll and Percoll medium (0.6 M

sucrose, 20 mM TES, 2 mM EDTA, 20 mM potassium dihydrogen phosphate, 2 mM glycine;

pH 7.5). Mitochondria were separated from other components by centrifuging in the gradients

at 17,400 g for 20 min. The resulting mitochondrial fractions formed white clouds at the bottom

half of the gradients and were collected by Pasteur pipettes to clean centrifuge tubes. The

clean-up procedures were performed three to five times by filling up wash buffer in the

centrifuge tubes and pelleting the mitochondria with 17,200 g for 20 min, until the resulting

pellet was firm. After each washing steps, two to three pellets were combined in one tube until

all mitochondria from one biological replicate were pooled together. The mitochondrial pellets

were weighed, resuspended with wash buffer and aliquoted at the concentration of 0.1 g/ml.

*From cell cultures*

Mitochondria isolation from *Arabidopsis thaliana* suspension cell culture was carried out as

described before (Farhat et al., 2019). About 200 g fresh cells were harvested and homogenized

using disruption buffer (450 mM sucrose, 15 mM MOPS, 1.5 mM EGTA, 0.6 % (w/v) PVP40, 2 %

(w/v) BSA, 10 mM sodium ascorbate, 10 mM cysteine; pH 7.4) and a waring blender. During

several washing steps, cell fragments were removed (centrifugation twice for 5 minutes at

2,700 g and once for 5 minutes at 8,300 g). Crude mitochondria were pelleted at 17,000 g for

10 minutes, resuspended in washing buffer (0.3 M sucrose, 10 mM MOPS, 1 mM EGTA; pH 7.2),

homogenized using a Dounce homogenisator and loaded onto discontinuous Percoll gradients

(phases of 18 %, 23 % and 40 % Percoll in gradient buffer (0.3 M sucrose, 10 mM MOPS, 0.2 mM

EGTA; pH 7.2)). After ultracentrifugation (90 minutes, 70,000 g), purified mitochondria were

collected from the 23 %-40 % interphase. For Percoll removal, several washing steps (10 minutes,

14,500 g) were performed using resuspension buffer (0.4 M mannitol, 1 mM EGTA, 10 mM

tricine; pH 7.2) to gain a firm pellet of purified mitochondria.

Each of the three independently purified mitochondria populations from 4-week old rosette

leaves (L-mito) and 200 g cell cultures (C-mito), respectively were used for all experiments.

**BN/SDS-PAGE**

Gel electrophoresis procedures (blue-native (BN) and SDS PAGE) were performed as described

previously (Senkler et al., 2018), based on the published protocol given in (Wittig et al., 2006).

**Label-free quantitative shotgun proteomics**

*Protein sample preparation*

Sample preparation for shotgun proteome analysis was performed as described before (Thal et

al., 2018). The protein content of the mitochondrial fractions was determined using a Bradford

assay kit (Thermo Scientific, Rockford, USA). 50 µg protein of each sample were loaded onto a

SDS gel for sample purification (Thal et al., 2018). Electrophoresis was stopped when the

proteins reached the border between the stacking and the separating gel. Gels were

subsequently incubated in fixation solution (15% (v.v) ethanol; 10% (v.v) acetic acid) for 30 min,

stained for 1 h with Coomassie Brilliant Blue G250, and finally the protein band at the border of

the two gel phases was cut out into cubes with edge lengths of approximately 1 mm.

Trypsination of the proteins was carried out as described previously (Fromm et al., 2016).

*Shotgun proteomic LC-MS analysis*

704 Label-free quantitative mass spectrometric analyses of whole mitochondrial protein samples

705 from cell culture and long day *Arabidopsis thaliana* leaves were performed as outlined before

706 (Thal et al., 2018) using an Ultimate 3000 UPLC coupled to a Q Exactive Orbitrap mass

707 spectrometer (Thermo Scientific, Dreieich, Germany).

*Data processing*

In a first step, the resulting MS data were processed using the Proteome Discoverer Software

(Thermo Fisher Scientific, Dreieich, Germany) and searched with the Mascot search engine

(www.matrixscience.com) against the tair10 protein database (downloaded from

www.arabidopsis.org). For quantitative analyses, MS data were further processed as outlined

before (Rugen et al., 2019) using the MaxQuant software package (version 1.6.4.0), the

Andromeda search engine (Cox and Mann, 2008) and the tair10 protein database. For

determination of sample purity, peptide intensities were used, combined with the subcellular

locations of assigned proteins as given by SUBAcon from the SUBA platform (www.suba.live)

(Hooper et al., 2017). A proteomic heatmap was generated using the NOVA software

(www.bioinformatik.uni-frankfurt.de/tools/nova/index.php)(Giese et al., 2015). Identified

proteins of all six datasets (three biological replicates of leaf total mitochondrial protein (L-mito

1,2,3) and cell culture total mitochondrial protein (C-mito 1,2,3)) were hierarchically clustered in

a heatmap based on iBAQ (intensity Based Absolute Quantification) values (for primary results

see Supplemental Tab. S1).

**Lipid extraction**

Total lipid extract from leaf (L-TE) was obtained as described (Tarazona et al., 2015). Briefly, mixture of 150 mg frozen leaf powder or 100 mg mitochondria and 6 ml extraction buffer (propan-2-ol : hexane : water, 60:26:14 (v.v.v)) was incubated at 60 °C with shaking for 30 min. After centrifugation at 800 g for 20 min, the clear supernatant was transferred to clean tubes and evaporated under stream of nitrogen gas until dryness. Samples were reconstituted in 800 µl of TMW (tetrahydrofuran (THF) : methanol : water, 4:4:1 (v.v.v)). Procedure for extracting the mitochondrial lipids was adjusted by substituting the water fraction of the extraction buffer by equal volume of the mitochondrial aliquot. Further process was continued as described above.

**Lipid quantification**

*TLC separation of lipid classes*

Lipid extracts from 500 mg mitochondria and 50 mg leaves were spotted onto TLC 60 plates (20 × 20 cm$^2$, Merck KGaA, Darmstadt, Germany) in parallel with the following standards: PC (Larodan, Solna, Sweden), PE, CL, and PG (Avanti Polar Lipids, Birmingham, AL, USA), MGDG, and DGDG (purified from plant lipid extracts). Extracts were developed by a solvent mixture of chloroform : methanol : acetic acid (65:25:8 (v.v.v)). After visualizing the lipid spots under 528 nm UV light, the bands were scrapped out and converted to fatty acid methyl esters (FAME) before GC/FID analysis.

*Acidic transesterification*

Glycerolipids were transesterified by acidic methanolysis (Miquel and Browse, 1992) and converted to fatty acid methyl esters (FAME). The lipid-bound silica powder from corresponding spots on the TLC plate were scrapped out and added to 1 ml FAME solution (methanol : toluene : sulfuric acid : dimethoxypropane, 33:17:1.4:1 (v.v.v.v)) with 5 µg tripentadecanoin as an internal standard. After 1 h incubation at 80 °C, 1.5 ml saturated NaCl solution and 1.2 ml hexane were added subsequently. The resulting FAME was collected from the hexane phase, dried, and reconstituted in 10 µl acetonitrile.

*GC/FID analysis*

Lipid-bound fatty acids were analyzed after converting to FAMEs by a 6890N Network GC/FID

System with a medium polar cyanopropyl DB-23 column (30 m × 250 μm × 25 nm; Agilent

Technologies, Waldbronn, Germany) using helium as the carrier gas at 1 ml min$^{-1}$. Samples were

injected at 220 °C with an Agilent 7683 Series injector in split mode. After 1 min at 150 °C, the

oven temperature raised to 200 °C at the rate of 8 °C min$^{-1}$, increased to 250 °C in 2 min, and

held at 250 °C for 6 min. Peak integration was performed using the GC ChemStation (Agilent

Technologies).

**Lipid derivatization**

Phosphate-containing lipids - phosphatidic acids (PA), phosphoinositides (PIPs) and long-chain

base phosphates (LCB-P) - were derivatized to enhance their chromatographic separation and

mass spectrometric detection. Methylation procedure was applied on PA and PIPs as followed.

Aliquots of 100 μl lipid extracts were first brought to dryness under stream of nitrogen gas and

reconstituted with 200 μl methanol. Methylation reaction took place after the supply of 3.3 μl

trimethylsilyldiazomethane. After 30 min incubation at room temperature, the reaction was

terminated by neutralizing with 1 μl of 1.7 M acetic acid. Samples were dried under nitrogen gas

and redissolved in 100 μl TMW. Acetylation procedure was applied on LCB-P. Aliquots of 100 μl

lipid extracts were brought to dryness and reconstituted with 100 μl pyridine and 50 μl acetic

anhydride. After 30 min incubation at 50 °C, samples were dried under stream of nitrogen gas

with 50 °C water bath. To redissolve the samples, 100 μl TMW was used as the final solvent to

proceed with lipid analysis.

**Global lipidomic analysis with LC-MS**

Analysis conditions and system setup were as described (Tarazona et al., 2015). Samples were

separated by an ACQUITY UPLC system (Waters Crop., Milford, MA, USA) with a HSS T3 column

(100 mm x 1 mm, 1.8 μl; Waters Crop.), ionized by a chip-based nanoelectrospray using TriVersa

Nanomate (Advion BioScience, Ithaca, NY, USA) and analyzed by a 6500 QTRAP tandem mass

spectrometer (AB Sciex, Framingham, MA, USA). Aliquots of 2 μl were injected and separated

with a flow rate of 0.1 ml min$^{-1}$. The solvent system composed of methanol : 20 mM ammonium

acetate (3:7 (v.v)) with 0.1 % (v.v) acetic acid (solvent A) and THF : methanol : 20 mM

ammonium acetate (6:3:1 (v.v.v)) with 0.1 % (v.v) acetic acid (solvent B). According to the lipid

classes, different linear gradients were applied: start from 40 %, 65 %, 80 % or 90 % B for 2 min;

increase to 100 % B in 8 min; hold for 2 min and re-equilibrate to the initial conditions in 4 min.

Starting condition of 40 % solvent B were utilized for long-chain bases (LCB) and phosphorylated

long-chain bases (LCB-P); 80 % for diacylglycerol (DAG); 90 % for steryl esters (SE); 65 % for the

remaining lipid classes. Retention time alignment and peak integration were performed with

MultiQuant (AB Sciex). Quantitative results were calculated according to the amount of internal

standards.

**Biosynthesis pathways construction**

The Arabidopsis Information Resource (TAIR; www.arabidopsis.org), the Subcellular localization

database for Arabidopsis proteins (SUBAcon; www.suba.live), Kyoto Encyclopedia of Genes and

Genomes (KEGG; www.genome.jp/kegg) and the shotgun proteomic analyses of L-mito and C-

mito were combined to construct the lipid biosynthesis pathways in plant mitochondria.

Additionally, lipidomics data are depicted in the pathways to illustrate the biosynthetic fluxes.

**Supplemental Data**

**Supplemental Figure S1.** Purity inspection of mitochondrial protein complexes and

supercomplexes by two-dimensional blue-native/SDS PAGE. Mitochondrial fractions isolated

from Arabidopsis leaves (L-mito 1 and L-mito 3) and from Arabidopsis cell cultures (C-mito1 and

C-mito 3) were separated by 2D PAGE and Coomassie-stained (corresponding gels of fractions L-

mito 2 and C-mito 2 see Fig. 1). Numbers on top and to the left of the 2D gels refer to the

masses of standard protein complexes / proteins (in kDa), the roman numbers above the gels to

the identity of OXPHOS complexes (see Fig. 1 for detailed information). The arrows indicate the

large (L; 53,5 kDa) and the small (S; 14.5 kDa) subunit of Rubisco. To further confirm the

reproducibility of the preparations, 2D blue-native/SDS reference gels for mitochondrial and

chloroplast fractions (Mito-ref, Cp-ref) from *A. thaliana* are given to the bottom of the figure

(gels were taken from (Klodmann et al., 2011) and (Behrens et al., 2013)). Identity of the protein

complexes visible on the chloroplast reference gel: PSI – photosystem I; PSII – photosystem II;

Rub – Rubisco; LHCII – light harvesting complex II.

**Supplemental Figure S2.** Purity of mitochondrial fractions as determined by label-free quantitative shot gun proteomics. Three mitochondrial fractions isolated from Arabidopsis leaves (L-mitos) and cell cultures (C-mitos) were analysed. Summed-up peptide intensities were calculated for subcellular compartments based on protein assignments as given by SUBAcon (The Subcellular localization database for Arabidopsis proteins; www.suba.live**).** Blue: mitochondria; green: plastids; gray: others; numbers in %.

**Supplemental Figure S3.** Fatty acid profiles of the glycerolipids from L-mito, L-TE, C-mito and C-TE. Heat maps illustrate the difference of the fatty acid distribution based on LC-MS/MS analyses. The heat map colors represent mean intensity values according to the color map on the low right-hand side. Each column represents one lipid class wherein the acyl moieties of all species are listed and summarized to 100 %. Data represent mean values in mol % from three independent experiments which were also used for proteomics (Supplemental Tab. S1).

**Supplemental Figure S4.** Workflow for the construction of the biosynthesis pathways. Multiple databases were combined to build the lipid biosynthesis pathways in mitochondria, The Arabidopsis Information Resource (TAIR; www.arabidopsis.org), the Subcellular localization database for Arabidopsis proteins (SUBAcon; www.suba.live), Kyoto Encyclopedia of Genes and Genomes (KEGG; www.genome.jp/kegg) and the proteomic datasets of isolated mitochondria in this study. Information of protein localizations and backbones of the biosynthesis pathways were obtained from TAIR, SUBAcon and KEGG, respectively.

**Supplemental Figure S5.** Additional lipid biosynthesis enzymes which do not match with analyzed lipid classes in this study. Enzymes of (a) fatty acid biosynthesis, (b) unsaturated fatty acid biosynthesis, (c) lipoic acid metabolism and (d) ubiquinone and other terpenoid-quinone biosynthesis illustrate the general lipid biosynthesis in Arabidopsis mitochondria. Heat maps visualize the protein abundance based on shotgun proteomic analyses of three independent experiments of purified mitochondria from leaves and cell culture. The heat map colors represent mean intensity values according to the color map on the low right-hand side. Blue: proteins identified in the proteomic analysis of this study and also predicted to localize in mitochondria; green: proteins identified in the proteomic analysis of this study but predicted to localize in other organelles; bold font: exclusively localized in mitochondria; italic font: only identified in one of the mitochondrial populations. Further full names and functions are itemized in Supplemental Tab. S3. In the heat maps, proteins with high and low abundance are depicted in yellow and black, respectively. Predicted protein localization was based on The Arabidopsis Information Resource (TAIR; www.arabidopsis.org) and the Subcellular localization database for Arabidopsis proteins (SUBAcon; www.suba.live).

**Supplemental Table S1.** Overall proteins identified in the proteomic approach.

**Supplemental Table S2.** Overall lipids identified in the lipidomic approach.

**Supplemental Table S3.** Lipid biosynthesis-related proteins identified in the proteomic approach.

**Supplemental Table S4.** List of subunits of the MTL identified in the proteomic approach.

**Supplemental Table S5.** Proteins assigned to plastids regarding to SUBAcon.

**Acknowledgements**

We are grateful for the technical assistance from Sabine Freitag. YL has been a doctoral student

of the Ph.D. program "Molecular Biology" – International Max Planck Research School and the

Göttingen Graduate School for Neurosciences, Biophysics, and Molecular Biosciences (GGNB)

(DFG grant GSC 226) at the Georg August University Göttingen.

**Funding information**

IF and HPB acknowledge funding through the German Research Foundation (DFG: INST 186/822-

1, INST 186/1167-1, INST 187/503-1 and ZUK 45/2010).

**References**

**Achleitner G, Gaigg B, Krasser A, Kainersdorfer E, Kohlwein SD, Perktold A, Zellnig G, Daum G** (1999)

Association between the endoplasmic reticulum and mitochondria of yeast facilitates interorganelle

transport of phospholipids through membrane contact. Eur J Biochem **264:** 545-553

**Alcázar-Román AR, Wente SR** (2008) Inositol polyphosphates: a new frontier for regulating gene

expression. Chromosoma **117:** 1-13

**Ali U, Li H, Wang X, Guo L** (2018) Emerging roles of sphingolipid signaling in plant response to biotic and

abiotic stresses. Mol Plant **11:** 1328-1343

**Awai K, Xu C, Tamot B, Benning C** (2006) A phosphatidic acid-binding protein of the chloroplast inner

envelope membrane involved in lipid trafficking. Proc Natl Acad Sci USA **103:** 10817-10822

**Babiychuk E, Müller F, Eubel H, Braun H-P, Frentzen M, Kushnir S** (2003) *Arabidopsis*

phosphatidylglycerophosphate synthase 1 is essential for chloroplast differentiation, but is

dispensable for mitochondrial function. Plant J **33:** 899 - 909

**Behrens C, Blume C, Senkler M, Eubel H, Peterhänsel C, Braun H-P** (2013) The 'protein complex

proteome' of chloroplasts in *Arabidopsis thaliana*. Journal of Proteomics **91:** 73-83

**Bi F-C, Liu Z, Wu J-X, Liang H, Xi X-L, Fang C, Sun T-J, Yin J, Dai G-Y, Rong C, Greenberg JT, Su W-W, Yao**

**N** (2014) Loss of ceramide kinase in *Arabidopsis* impairs defenses and promotes ceramide

accumulation and mitochondrial $H_2O_2$ bursts. Plant Cell **26:** 3449-3467

**Bionda C, Portoukalian J, Schmitt D, Rodriguez-Lafrasse C, Ardail D** (2004) Subcellular

compartmentalization of ceramide metabolism: MAM (mitochondria-associated membrane) and/or

mitochondria? Biochem J **382:** 527-533

**Birner R, Burgermeister M, Schneiter R, Daum G** (2001) Roles of phosphatidylethanolamine and of its

several biosynthetic pathways in *Saccharomyces cerevisiae*. Molecular Biology of the Cell **12:** 997-

1007.

**Bligny R, Douce R** (1980) A precise localization of cardiolipin in plant cells. Biochim Biophys Acta **617:**

254-263

**Braun H-P** (2020) The Oxidative phosphorylation system of the mitochondria in plants. Mitochondrion **53:**

66-75

**Cacas J-L, Furt F, Le Guédard M, Schmitter J-M, Buré C, Gerbeau-Pissot P, Moreau P, Bessoule J-J,**

**Simon-Plas F, Mongrand S** (2012) Lipids of plant membrane rafts. Prog Lipid Res **51:** 272-299

**Charman M, Kennedy BE, Osborne N, Karten B** (2010) MLN64 mediates egress of cholesterol from

endosomes to mitochondria in the absence of functional Niemann-Pick Type C1 protein. J Lipid Res

**51:** 1023-1034

**Clark BJ, Pezzi V, Stocco DM, Rainey WE** (1995) The steroidogenic acute regulatory protein is induced by

angiotensin II and $K^+$ in H295R adrenocortical cells. Molecular and Cellular Endocrinology **115:** 215-

219

**Cox J, Mann M** (2008) MaxQuant enables high peptide identification rates, individualized p.p.b.-range

mass accuracies and proteome-wide protein quantification. Nature Biotechnol **26:** 1367-1372

**Donahue JL, Alford SR, Torabinejad J, Kerwin RE, Nourbakhsh A, Ray WK, Hernick M, Huang X, Lyons**

**BM, Hein PP, Gillaspy GE** (2010) The *Arabidopsis thaliana myo*-inositol 1-phosphate synthase1 gene

is required for *myo*-inositol synthesis and suppression of cell death. Plant Cell **22:** 888-903

**Douce R, Mannella CA, Bonner WD** (1972) Site of the biosynthesis of CDP-diglyceride in plant

mitochondria. Biochem Biophys Res Commun **49:** 1504-1509

**Du Z-Y, Chen M-X, Chen Q-F, Xiao S, Chye M-L** (2013) Arabidopsis acyl-CoA-binding protein ACBP1
participates in the regulation of seed germination and seedling development. Plant J **74:** 294-309
**Essigmann B, Guler S, Narang RA, Linke D, Benning C** (1998) Phosphate availability affects the thylakoid
lipid composition and the expression of *SQD1*, a gene required for sulfolipid biosynthesis in
*Arabidopsis thaliana*. Proc Natl Acad Sci USA **95:** 1950-1955
**Estavillo GM, Crisp PA, Pornsiriwong W, Wirtz M, Collinge D, Carrie C, Giraud E, Whelan J, David P,**
**Javot H, Brearley C, Hell R, Marin E, Pogson BJ** (2011) Evidence for a SAL1-PAP chloroplast
retrograde pathway that functions in drought and high light signaling in *Arabidopsis*    Plant Cell **23:**
3992-4012
**Farhat N, Hichri S, Hildebrandt TM, Debez A, Braun H-P** (2019) Composition and stability of the oxidative
phosphorylation system in the halophile plant *Cakile Maritima*. Front Plant Sci **10:** 1010
**Flis VV, Daum G** (2013) Lipid transport between the endoplasmic reticulum and mitochondria. Cold
Spring Harbor Perspectives in Biology **5:** a013235
**Fromm S, Senkler J, Eubel H, Peterhänsel C, Braun H-P** (2016) Life without complex I: proteome analyses
of an Arabidopsis mutant lacking the mitochondrial NADH dehydrogenase complex. J Exp Bot **67:**
3079-3093
**Giese H, Ackermann Jr, Heide H, Bleier L, Dröse S, Wittig I, Brandt U, Koch I** (2015) NOVA: a software
to analyze complexome profiling data. Bioinformatics **31:** 440-441
**Gray MW, Burger G, Lang BF** (1999) Mitochondrial evolution. Science **283:** 1476-1481
**Haslam TM, Feussner I** (2022) Diversity in sphingolipid metabolism across land plants. J Exp Bot **73:**
2785-2798
**Haslam TM, Kunst L** (2013) Extending the story of very-long-chain fatty acid elongation. Plant Sci **210:** 93-
107
**Heilmann I** (2016) Phosphoinositide signaling in plant development. Development **143:** 2044-2055
**Hölzl G, Dörmann P** (2019) Chloroplast Lipids and Their Biosynthesis. Annu Rev Plant Biol **70:** null
**Hölzl G, Dörmann P** (2019) Chloroplast lipids and their biosynthesis. Annu Rev Plant Biol **70:** 51-81
**Hooper CM, Castleden IR, Tanz SK, Aryamanesh N, Millar AH** (2017) SUBA4: the interactive data analysis
centre for Arabidopsis subcellular protein locations. Nucleic Acids Res **45:** D1064-D1074
**Horvath SE, Daum G** (2013) Lipids of Mitochondria. Prog Lipid Res **52:** 590-614
**Inoue K** (2007) The chloroplast outer envelope membrane: the edge of light and excitement. Journal of
Integrative Plant Biology **49:** 1100-1111
**Inoue K** (2011) Emerging roles of the chloroplast outer envelope membrane. Trends Plant Sci **16:** 550-
557

**Jacoby RP, Li L, Huang S, Pong Lee C, Millar AH, Taylor NL** (2012) Mitochondrial composition, function

and stress response in plants. Journal of Integrative Plant Biology **54:** 887-906

**Jouhet J, Marechal E, Baldan B, Bligny R, Joyard J, Block MA** (2004) Phosphate deprivation induces

transfer of DGDG galactolipid from chloroplast to mitochondria. Journal of Cell Biology **167:** 863-874

**Katayama K, Sakurai I, Wada H** (2004) Identification of an *Arabidopsis thaliana* gene for cardiolipin

synthase located in mitochondria. FEBS Lett **577:** 193-198

**Kim J, Na YJ, Park SJ, Baek S-H, Kim DH** (2019) Biogenesis of chloroplast outer envelope membrane

proteins. Plant Cell Rep **38:** 783-792

**Kitagaki H, Cowart LA, Matmati N, Vaena de Avalos S, Novgorodov SA, Zeidan YH, Bielawski J, Obeid**

**LM, Hannun YA** (2007) Isc1 regulates sphingolipid metabolism in yeast mitochondria. Biochim

Biophys Acta **1768:** 2849-2861

**Klahre U, Noguchi T, Fujioka S, Takatsuto S, Yokota T, Nomura T, Yoshida S, Chua NH** (1998) The

Arabidopsis *DIMINUTO/DWARF1* gene encodes a protein involved in steroid synthesis. Plant Cell **10:**

1677-1690

**Klodmann J, Senkler M, Rode C, Braun H-P** (2011) Defining the *protein complex proteome* of plant

mitochondria. Plant Physiol **157:** 587-598

**Lenarčič T, Albert I, Böhm H, Hodnik V, Pirc K, Zavec AB, Podobnik M, Pahovnik D, Žagar E, Pruitt R,**

**Greimel P, Yamaji-Hasegawa A, Kobayashi T, Zienkiewicz A, Gömann J, Mortimer JC, Fang L,**

**Mamode-Cassim A, Deleu M, Lins L, Oecking C, Feussner I, Mongrand S, Anderluh G, Nürnberger T**

(2017) Eudicot plant-specific sphingolipids determine host selectivity of microbial NLP cytolysins.

Science **358:** 1431-1434

**Li-Beisson Y, Shorrosh B, Beisson F, Andersson MX, Arondel V, Bates PD, Baud S, Bird D, DeBono A,**

**Durrett TP, Franke RB, Graham IA, Katayama K, Kelly AA, Larson T, Markham JE, Miquel M, Molina**

**I, Nishida I, Rowland O, Samuels L, Schmid KM, Wada H, Welti R, Xu C, Zallot R, Ohlrogge J** (2013)

Acyl-lipid metabolism. *In* C Somerville, E Meyerowitz, eds, The Arabidopsis Book. The American

Society of Plant Biologists, Rockville, MD, USA, p e0161

**Li L, Lavell A, Meng X, Berkowitz O, Selinski J, van de Meene A, Carrie C, Benning C, Whelan J, De Clercq**

**I, Wang Y** (2019) Arabidopsis DGD1 SUPPRESSOR1 is a subunit of the mitochondrial contact site and

cristae organizing system and affects mitochondrial biogenesis. Plant Cell **31:** 1856-1878

**Liebisch G, Fahy E, Aoki J, Dennis EA, Durand T, Ejsing C, Fedorova M, Feussner I, Griffiths WJ, Koefeler**

**H, Merrill AH, Murphy RC, O'Donnell VB, Oskolkova OV, Subramaniam S, Wakelam M, Spener F**

(2020) Update on LIPID MAPS classification, nomenclature and shorthand notation for MS-derived

lipid structures. J Lipid Res **61:** 1539–1555

**Mehrshahi P, Stefano G, Andaloro JM, Brandizzi F, Froehlich JE, DellaPenna D** (2013) Transorganellar

complementation redefines the biochemical continuity of endoplasmic reticulum and chloroplasts.

Proc Natl Acad Sci USA **110:** 12126-12131

**Michaud M, Gros V, Tardif M, Brugiere S, Ferro M, Prinz WA, Toulmay A, Mathur J, Wozny M, Falconet**

**D, Marechal E, Block MA, Jouhet J** (2016) AtMic60 is involved in plant mitochondria lipid trafficking

and is part of a large complex. Curr Biol **26:** 627-639

**Michaud M, Prinz WA, Jouhet J** (2017) Glycerolipid synthesis and lipid trafficking in plant mitochondria.

The FEBS Journal **284:** 376-390

**Miquel M, Browse J** (1992) *Arabidopsis* mutants deficient in polyunsaturated fatty acid synthesis. J Biol

Chem **267:** 1502-1509

**Mizoi J, Nakamura M, Nishida I** (2006) Defects in CTP:PHOSPHORYLETHANOLAMINE

CYTIDYLYLTRANSFERASE affect embryonic and postembryonic development in *Arabidopsis*. Plant

Cell **18:** 3370-3385

**Moreau F, Dupont J, Lance C** (1974) Phospholipid and fatty acid composition of outer and inner

membranes of plant mitochondria. Biochim Biophys Acta **345:** 294-304

**Morré DJ, Merritt WD, Lembi CA** (1971) Connections between mitochondria and endoplasmic reticulum

in rat liver and onion stem. Protoplasma **73:** 43-49

**Müller F, Frentzen M** (2001) Phosphatidylglycerophosphate synthases from *Arabidopsis thaliana*. FEBS

Lett **509:** 298-302

**Nandi A, Welti R, Shah J** (2004) The *Arabidopsis thaliana* dihydroxyacetone phosphate reductase gene

*SUPPRESSOR OF FATTY ACID DESATURASE DEFICIENCY1* is required for glycerolipid metabolism and

for the activation of systemic acquired resistance. Plant Cell **16:** 465–477

**Nerlich A, von Orlow M, Rontein D, Hanson AD, Dörmann P** (2007) Deficiency in phosphatidylserine

decarboxylase activity in the *psd*1 *psd*2 *psd*3 triple mutant of Arabidopsis affects

phosphatidylethanolamine accumulation in mitochondria. Plant Physiol **144:** 904-914

**Niehaus M, Straube H, Künzler P, Rugen N, Hegermann J, Giavalisco P, Eubel H, Witte C-P, Herde M**

(2020) Rapid affinity purification of tagged plant mitochondria (Mito-AP) for metabolome and

proteome analyses. Plant Physiol **182:** 1194–1210

**Novgorodov SA, Wu BX, Gudz TI, Bielawski J, Ovchinnikova TV, Hannun YA, Obeid LM** (2011) Novel

pathway of ceramide production in mitochondria. J Biol Chem **286:** 25352-25362

**Quettier A-L, Shaw E, Eastmond PJ** (2008) SUGAR-DEPENDENT6 Encodes a Mitochondrial Flavin Adenine

Dinucleotide-Dependent Glycerol-3-P Dehydrogenase, Which Is Required for Glycerol Catabolism

and Postgerminative Seedling Growth in Arabidopsis. Plant Physiol **148:** 519-528

**Rugen N, Straube H, Franken LE, Braun H-P, Eubel H** (2019) Complexome profiling reveals association of

PPR proteins with ribosomes in the mitochondria of plants. Molecular & Cellular Proteomics **18:**

1345-1362

**Schaller H** (2003) The role of sterols in plant growth and development. Prog Lipid Res **42:** 163-175

**Schikowsky C, Thal B, Braun H-P, Eubel H** (2018) Sample preparation for analysis of the plant

mitochondrial membrane proteome. *In* H-P Mock, A Matros, K Witzel, eds, Plant Membrane

Proteomics: Methods and Protocols. Springer New York, New York, NY, pp 163-183

**Senkler J, Rugen N, Eubel H, Hegermann J, Braun H-P** (2018) Absence of complex I implicates

rearrangement of the respiratory chain in european mistletoe. Curr Biol **28:** 1606-1613.e1604

**Simm S, Papasotiriou D, Ibrahim M, Leisegang M, Müller B, Schorge T, Karas M, Mirus O, Sommer M,**

**Schleiff E** (2013) Defining the core proteome of the chloroplast envelope membranes. Front Plant

Sci **4**

**Staehelin LA** (1997) The plant ER: a dynamic organelle composed of a large number of discrete functional

domains. Plant J **11:** 1151-1165

**Tarazona P, Feussner K, Feussner I** (2015) An enhanced plant lipidomics method based on multiplexed

liquid chromatography–mass spectrometry reveals additional insights into cold- and drought-

induced membrane remodeling. Plant J **84:** 621-633

**Tatsuta T, Scharwey M, Langer T** (2014) Mitochondrial lipid trafficking. Trends Cell Biol **24:** 44-52

**Tavassoli S, Chao JT, Young BP, Cox RC, Prinz WA, de Kroon AIPM, Loewen CJR** (2013) Plasma

membrane - endoplasmic reticulum contact sites regulate phosphatidylcholine synthesis. EMBO rep

**14:** 434-440

**Thal B, Braun H-P, Eubel H** (2018) Proteomic analysis dissects the impact of nodulation and biological

nitrogen fixation on *Vicia faba* root nodule physiology. Plant Mol Biol **97:** 233-251

**Vance JE** (1990) Phospholipid synthesis in a membrane fraction associated with mitochondria. J Biol

Chem **265:** 7248-7256

**Watanabe M, Balazadeh S, Tohge T, Erban A, Giavalisco P, Kopka J, Mueller-Roeber B, Fernie AR,**

**Hoefgen R** (2013) Comprehensive Dissection of Spatiotemporal Metabolic Shifts in Primary,

Secondary, and Lipid Metabolism during Developmental Senescence in Arabidopsis. Plant Physiol

**162:** 1290-1310

**Wittig I, Braun H-P, Schägger H** (2006) Blue native PAGE. Nat Protoc **1:** 418-428

**Xu C, Hartel H, Wada H, Hagio M, Yu B, Eakin C, Benning C** (2002) The *pgp1* mutant locus of Arabidopsis

encodes a phosphatidylglycerolphosphate synthase with impaired activity. Plant Physiol **129:** 594-

604

**Xu Y, Malhotra A, Ren M, Schlame M** (2006) The enzymatic function of Tafazzin. J Biol Chem **281:** 39217-
39224

**Yamaoka Y, Yu Y, Mizoi J, Fujiki Y, Saito K, Nishijima M, Lee Y, Nishida I** (2011) *PHOSPHATIDYLSERINE*
*SYNTHASE1* is required for microspore development in *Arabidopsis thaliana*. Plant J **67:** 648-661

**Youn JH, Kim T-W, Joo S-H, Son S-H, Roh J, Kim S, Kim T-W, Kim S-K** (2018) Function and molecular
regulation of DWARF1 as a C-24 reductase in brassinosteroid biosynthesis in Arabidopsis. J Exp Bot
**69:** 1873-1886