1 **Melanization of _Candida auris_ is Associated with Alteration of Extracellular pH**
2

3 Daniel F. Q. Smith[‡1], Nathan J. Mudrak[‡1,2], Daniel Zamith-Miranda[‡3,4], Leandro Honorato[5], Leonardo
4 Nimrichter [5], Barbara Smith[6], Gary Gerfen[7], Joshua D. Nosanchuk[#3,4], Arturo Casadevall[#1]
5

[1] W. Harry Feinstone Department of Molecular Microbiology and Immunology, The Johns Hopkins
Bloomberg School of Public Health, Baltimore, MD, USA
[2] Krieger School of Arts & Science, Johns Hopkins University, Baltimore, MD, USA
[3] Department of Microbiology and Immunology, Albert Einstein College of Medicine, Bronx, New York,
USA
[4] Division of Infectious Diseases, Department of Medicine, Albert Einstein College of Medicine, Bronx,
New York, USA
[5] Laboratório de Glicobiologia de Eucariotos, Departamento de Microbiologia Geral, Instituto de
Microbiologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro,
Brazil
[6] Institute for Basic Biomedical Sciences Microscope Facility, The Johns Hopkins School of Medicine,
Baltimore, MD, USA
[7] Department of Biochemistry, Albert Einstein College of Medicine, Bronx, New York, USA
19
20
21 [‡] DFQS, NJM, and DZM contributed equally to this work
22 [#] JDN and AC contributed equally to this work
23

Abstract

   Candida auris is a recently emerged global fungal pathogen, which causes life threatening infections, often in healthcare settings and with immunocompromised patients. *C. auris* infections are particularly worrisome because the fungus is often resistant to multiple classes of antifungal drugs. Furthermore, the fungus forms durable biofilms that are difficult to remove from surfaces. Due to the relatively recent, resilient, and resistant nature of this fungal pathogen, we investigated whether *C. auris* made common fungal virulence factor melanin. Melanin is a black-brown pigment that enables microbes to evade the mammalian immune response due to its strong antioxidant properties, ability to inactivate antimicrobial peptides, shielding of pathogen-associated molecular patterns, and inactivation of antifungal drugs. We found that certain strains of *C. auris,* predominantly from Clade I and IV, oxidized catecholamines into melanin pigment. The formation of this melanin occurred extracellularly, resulting in granule-like melanin structures that are deposited on the surface of the fungus. This melanization was primarily chemically driven and mediated by alkalinization of the extracellular environment. The *C. auris* strains had relatively high cell surface hydrophobicity, but we did not find a correlation between hydrophobicity and ability to melanize. Melanin protected the fungus from oxidative damage but we did not observe a protective role *in vitro* or *in vivo* during infection of macrophages or *Galleria mellonella*, respectively. In summary, *C. auris* alkalinizes the extracellular medium, which promotes the chemical autooxidation of the L-dopa to melanin that then attaches to its surface, thus illustrating a different mechanism for fungal melanization.

Introduction

Candida auris is an emerging fungal pathogen that is believed to original in marine wetlands and marshes (1,2). *C. auris* was first described in a clinical setting in 2009, but patient samples have been retrospectively determined to be *C. auris* as early as 1996 (3–5). Since its discovery, *C. auris* has caused outbreaks in at least 47 countries (3). *C. auris* is often acquired in healthcare settings and is believed to cause infections via biofilms formed on medical equipment and intravenous catheters leading to bloodstream infection (candidemia) and dissemination to organs (6). *C. auris* is further notable because it has a remarkably high resistance to known antifungal therapies, with exceedingly high resistances to azole and echinocandin classes of antifungal drugs (3,7). This poses a major concern for clinicians treating infections, as the fungus may be resistant to many of the commonly used antifungal therapeutics. Hence, a deeper understanding is needed about the physical properties of *C. auris* and how they contribute to virulence.

One unexplored aspect of *C. auris* physiology is whether it can produce the virulence factor melanin. Other *Candida* species, such as *C. albicans* and *C. glabrata* produce this virulence factor (8–11). Melanin is a black-brown, insoluble, acid resistant pigment found throughout a plethora of life forms, and has particular importance in the context of fungal virulence in mammals (12,13). In fungi, melanin is typically produced through either the oxidation of catecholamines (DOPA melanin), the tyrosine degradation pathway (pyomelanin), or through polyketide synthase pathways (DHN melanin) (12). Genes controlling melanization are affected by temperature, nutrient stress and copper/metal ion concentration (14–16), and the biochemical processes can be altered by the environmental antioxidant balance and pH (17,18). Melanin is formed either within the cell in organelles termed "melanosomes" and then exported to the cell wall, or in the cell wall itself (19,20). Once located within the cell wall the pigment is typically anchored through interactions with chitin, chitosan, and other cell wall components (9,21,22). Melanin strongly interacts with polysaccharides, lipids, and proteins, many of which are still intact following boiling in acid, lipid extraction, and enzymatic degradation, and fungi are thought to use these molecules as a scaffold for the melanin deposition in the cell wall (19,23,24). This melanin could also be secreted, as seen in *C. neoformans,* where melanin granules are shed from the cell wall and can be recovered from the supernatant (19,25).

Functionally, fungal melanin pigments have strong antioxidant properties that allow them to resist oxidative damage caused by the host immune cells, such as macrophage and neutrophil oxidative bursts (26). Additionally, melanin can bind and sequester antimicrobial peptides and antimicrobial enzymes that the host typically use to degrade and kill fungi during infection, as well as antifungal drugs used to treat infections (27–29). Fungal melanins located in the cell wall can alter cell wall composition and affect recognition of pathogen-associated molecular patterns that would otherwise be exposed. This may lead to diminished recognition by host immune cells. Conversely, DHN melanin can be recognized by the melanin sensing C-type lectin (MelLec) receptor, which is part of the human innate immune response, and can enhance fungal clearance (30). Melanin could also be presented to lymphocytes in a stimulatory context, which results in melanin-specific antibody production, which can inevitably lead to fungal opsonization, phagocytosis, and complement deposition (27,31,32).

In this study, we investigated the ability of 18 *C. auris* strains to produce melanin. Like other fungi, we found that *C. auris* produces a black melanin pigment, which was located on the surface of the cell wall. However, we found that, unlike other fungi, the melanization occurs extracellularly, in the supernatant. In liquid culture, the cell-free melanin can adhere to the surface of the cell and cause pigmentation of the fungus. We found this process to be mediated by the *C. auris'* ability to neutralize the pH of the media it is grown in, resulting in enhanced autoxidation of the catecholamine melanin substrate. This method of melanin production differs greatly from what is reported in other fungi. *C. auris* melanin

90   protected the fungus from oxidative stress but was not protective *in vitro* in the context of interactions with
91   macrophages, or in an *in vivo* challenge of *Galleria mellonella* larvae.
92
Results
Temperature Dependence of *Candida auris* Strain Melanization
To evaluate the ideal temperature for *Candida auris* melanization, we tested the degree of
melanization at both 30° and 37 °C. We found that the strains melanized to a greater extent at the higher
temperature compared to the lower temperature (Figure 1A-C). The melanin-capable strains of *C. auris*
were CDC 385, CDC 386, CDC 387, CDC 388, CDC 389, CDC 390, CDC 931, CDC 1097, CDC 1104,
and MMC1. Strain CDC 382 exhibited an intermediate melanin phenotype. Additionally, we note that
certain strains, namely CDC 388, CDC 390, CDC 1097, and MMC1 melanized significantly more at 37 °C
and not at the lower temperature. This has interesting implications for understanding regulation of
melanization and the role it has in warm environmental niches, as well as during infections of humans,
who have a normal core body temperature of ~37 ˚C.
Further, the capacity of individual strains to melanize was associated with their evolutionary
clade, with strains belonging to clades I and IV – associated with South Asia and South America,
respectively – demonstrating melanization activity, while those strains belonging to clades II and II –
associated with East Asia and Africa, respectively – did not. In addition, the single representative of clade
V – associated with Iran – melanized, but the lack of additional clade V isolates for testing hinders the
generalization of this finding.
110
Electron paramagnetic resonance (EPR)
Melanin generally contains a stable free-radical, which lends to its antioxidant properties. Stable
free radical species can be detected by electron paramagnetic resonance (EPR). Melanin from 7-days
cultures was extracted from 12 strains of *C. auris* and analyzed by EPR, with melanin extracted from *C.
neoformans* H99 used as a standard (Figure 1D-F). The EPR spectra from the non-melanizing strains did
not display the melanin-distinctive EPR peak (Figure 1E), but the EPR spectra from the pigment extracted
from the melanin-producing strains (Figure 1F) were similar to the spectrum obtained from *C. neoformans*
melanin (Figure 1D).
119
Time and Cell Density
To understand the optimal conditions for melanin production by *C. auris*, yeast cells were
incubated with L-DOPA starting with different cell densities, and the production of pigment was analyzed
for 13 days and compared to cells incubated in the absence of L-DOPA (Figure 2). *C. neoformans* was
also cultivated with the same conditions and compared to the *C. auris* melanization profile. Melanin
production was optimal by cells in high density (10^7 cells/mL). In this condition, melanin was visible from
day 3, and intensified along the days, reaching its peak between days 7 and 10. Some strains melanized
in a delayed fashion when in medium cell density (10^6 cells/mL), whereas none of the low density (10^5
cell/mL) cultures produced visible pigment. The cell density-associated phenotype presented by *C.
neoformans* was the opposite of that presented by *C. auris*, as melanin production by low cell density *C.
neoformans* cells was equally or more effective than in the high-density conditions.
131
***Candida auris* Strains Melanize Using Substrates Associated with DOPA melanin**
We examined various known melanin precursors to evaluate which the *C. auris* strains could use
as substrates for melanization, and if there are any strain-specific differences between the substrates
used. We found that at a starting inoculum of 10^7 cells/ml, melanin-capable *C. auris* strains were able to
use L-3,4-dihydroxyphenylalanine (L-DOPA), D-3,4-dihydroxyphenylalanine (D-DOPA), L-methyl-3,4-

dihydroxyphenylalanine (Methyl-DOPA/M-DOPA), dopamine, norepinephrine, epinephrine, and a mixture
of catecholamine neurotransmitters (dopamine, epinephrine, and norepinephrine) at the ratios found in
the mammalian brain (33). We found that in general, all melanin-capable *C. auris* strains were able to use
the same substrates (Figure 3A). Some strains, namely CDC 385 and CDC 386 were less able to
produce pigment when grown with dopamine. While the L-DOPA, D-DOPA, M-DOPA, dopamine, brain
mix, and norepinephrine produced dark brown-black pigments, the epinephrine produced a reddish-brown
amber color of melanization (Figure 3B).
Notably, the *C. auris* strains were unable to use L-tyrosine, homogentisic acid (HGA), and 4-
hydroxyphenylpyruvic acid (4-HPP) as precursors to melanization, which indicates that *C. auris* cannot
produce pyomelanin, a process typically mediated by laccases (34), and does not have a tyrosinase
enzyme that can convert L-tyrosine into suitable catecholamine melanin precursors. Additionally, the
strains are unable to oxidize ABTS, a laccase-specific substrate, further indicating that the melanization
agent is not a laccase (Figure 3C).
150
Melanin is Primarily Found in the Supernatant
During collection of the melanized cells, we noted that the supernatants of the cultures were
notably darker than expected given our work with another melanizing fungus, *C. neoformans* (Figure 4A,
B). Similarly, when cells were grown on agar plates with L-DOPA as a substrate for melanization, there
was a distinct halo of pigment surrounding the melanin-producing strains of *C. auris (*Figure 4C*)*. Notably,
there was little to no pigmentation of the yeast colony itself. This strongly indicated that the melanization
of the cells was primarily extracellular and did not originate within the cells themselves.
To evaluate whether the supernatant of the cells had melanin-producing enzymatic activity, we
added L-DOPA to the isolated supernatant. We found that the supernatants of melanizing strains, namely
the supernatant of MMC1, CDC 387 (B8441), and CDC 388 (B11098), were capable of oxidizing L-
DOPA, whereas the supernatant of non-melanizing strain CDC 381 (B11220) and minimal media alone
were not. This indicates, in part, that the melanin-producing components of the *C. auris* were secreted
(Figure 4D).
Using ultracentrifugation, we collected small particles and extracellular vesicles found within the
melanized supernatant of the liquid cultures. We found that a substantial amount of the pigment was in
this supernatant pellet. Using negative staining transmission electron microscopy (TEM), we found that
the pelleted melanin was within granule structures, comparable to what was seen in the cell wall
periphery of the melanized culture (Figure 4E, F). These structures measured 20-40 nm (Figure 4E), and
were similar in appearance to melanin granules secreted by *C. neoformans* (19). The granules from CDC
381 had the smallest mean diameter (~20 nm) and CDC 387 had the largest at ~40 nm (Figure 4F).
171
Cell-bound Melanin is localized to the Periphery of the Cell Wall
To visualize the localization of melanin on the *C. auris* cells grown in liquid media*,* we used light
microscopy, TEM, and scanning electron microscopy (SEM). Under light microscopy, dark pigmentation
of the cells appeared to be located primarily in the cell wall, with infrequent additional intracellular
pigmentation, possibly due to oxidation of L-DOPA within a large vacuole (Figure 5A). Comparing TEM
micrographs of melanized and non-melanized cultures, we saw electron-dense structures found
exclusively in the melanized cultures that primarily appeared on the periphery of the cell wall. These
electron-dense structures are likely melanin granules and appeared to be rounded and granular. The
extracellular melanin granules also appeared unbound to the cell wall (Figure 5B). These extracellular
melanin granules were similar to the secreted melanin granules that were roughly 20-40 nm in diameter
(Figure 5B, 4E-F). Similarly, samples imaged with SEM showed that melanized cells had raised structures

on their surface, indicative of the structures observed in TEM (Figure 5C). Further, in the case of CDC
387, 388, and MMC1, the melanin granules appear to hold the cells together in large, aggregated clumps.
To determine whether the directionality of the cell-bound melanin deposition (i.e., if melanization
occurred in the cell wall and then was released into media, or if melanization occurred primarily in the
media and then attached to the cell wall periphery), we performed a series of add-back experiments, in
which we added melanized supernatant to the cells from the same *C. auris* strain grown without L-DOPA.
We found that the cells become pigmented by three hours of incubation with the melanized supernatant
(Figure 6A-D). Additionally, when we added melanized supernatant from the CDC 388 strain to the non-
melanizing CDC 381 strain, the CDC 381 strain accumulated pigment (Figure 6D). First, since CDC 381
cannot readily produce pigment, this showed that the pigment accumulation in these add-back
experiments was due to adhesion of the already synthesized pigment in the supernatant rather than the
production of new pigment from unreacted L-DOPA in the supernatant. Second, it indicated that the
melanin-deficient cells did not have a cell wall difference that made them intrinsically unable to bind
extracellular melanin. These add-back experiment data, along with the secretion data in Figure 4 and
Figure 5, point to the conclusion that *C. auris* cell wall melanization occurs extracellularly and then sticks
to the outside of the cell wall.
We grew CDC 387 and CDC 388 with L-DOPA in the presence of compounds known to enhance
or block the proper formation of cell wall components (9,35), either through direct supplementation of the
cell wall components, by binding and blocking proper structure formation, or by inhibiting the enzyme
responsible for producing that cell wall component. This allowed us, in theory, to evaluate which
components of the cell wall were important for extracellular melanin adherence. Cultures conditions that
resulted in a darker supernatant than the control condition indicated the lack of melanin adherence to the
cell wall, and those with lighter supernatant indicated that the compound enhanced melanin adherence to
the cell wall.
Addition of N-acetylglucosamine (GlcNAc) enhances the cell wall's layer of chitin and also
increases cell wall melanization in *C. neoformans* and *C. albicans* (9,36). In CDC 387 and CDC 388,
addition of 5 mM GlcNAc did not enhance cell wall melanin adhesion, and resulted in darker supernatant,
perhaps indicating less adhesion (Figure 6E, F). In TEM micrographs, the melanin located in the cell wall
between the control and GlcNAc treated did not appear appreciably different (Figure 6G, H).
Conversely, we added 100 μM Calcofluor White (CFW), which forms hydrogen bonds with chitin
polymers as they grow and thus disrupts proper formation of chitin microfibrils (37,38). Adding this dye led
to markedly decreased supernatant pigmentation (Figure 6E, F). Intriguingly, under microscopic analysis,
the CFW appeared to precipitate and form crystals that bound the melanin. The adhesion of melanin to
the CFW crystals was the likely cause of the lack of pigmented supernatant. This resulted in black
crystals that had the fluorescent properties of the CFW, along with the expected CFW-stained cell walls
(Supplementary Figure 1A). The CFW crystals were also visible under electron microscopy (Figure 6H,
Supplementary Figure 1B). The crystallization persisted despite filter sterilization. In an assay with cell-
free supernatant with CFW and with/without L-DOPA, we confirmed that the melanin formed bound to the
CFW crystals and precipitated out of solution (Supplementary Figure 1C). We also noticed that the CDC
387 and CDC 388 strain supernatants without L-DOPA form a smaller population of CFW crystals
compared to the CDC 381 non-melanizing strains (Supplementary Figure 1D). This finding may indicate
that there is a correlation between the melanization factor and formation of CFW crystal fragments with a
smaller size, which could indicate better solubility.
Further, we grew the CDC 387 and CDC 388 strains under the presence of the antifungal
compound caspofungin at half of the inhibitory concentration reported by the Centers for Disease Control
and Prevention (CDC) (37), which inhibits the enzymes responsible for cell wall β-glucan synthesis.
Caspofungin treatment resulted in high levels of supernatant melanization (Figure 6E, F). Under light

microscopic and electron microscopic analysis, the treatment appeared to reduce the amount of melanin
present on the cell wall, with nearly no melanin visible on the cell wall via TEM (Figure 6G, H).
Additionally, in the CDC 388 strain, caspofungin treatment resulted in what appeared to be pseudohyphal
growth (Supplementary Figure 1E). Although the cells were able to grow in culture under this
concentration of caspofungin, in the electron micrograph, the cells appeared to have a deformed
morphology, and their cytoplasmic contents looked condensed and abnormal, potentially due to the
antifungal stress and cell wall defect (Figure 6G, H).
237
Melanization Affects the Hydrophobicity of Some *C. auris* Strains
Since melanin is a hydrophobic molecule, we investigated whether melanization had a correlation
with the cell surface hydrophobicity (CSH) of the *C. auris* strains. With *C. auris* strains grown in the
nutrient rich Yeast Peptone Dextrose (YPD) media and minimal media, 7 of 12 strains had CSH that was
greater than 80% (Figure 6I, J). There was no association between the strain hydrophobicity and its ability
to melanize, as the strongly melanizing strain CDC 387 and the weak/no melanizing strains CDC 381 and
MMC2 all had low hydrophobicity. Conversely, the non-melanizing strains CDC 383 and melanizing
strains CDC 388 had high hydrophobicity. Additionally, since hydrophobicity is known to play roles in
host-pathogen interactions (38–40), we investigated whether melanization affected the CSH of the *C.*
auris strains. Surprisingly, we found in the CDC 387 strain (and to a lesser extent in the non-melanizing
strain CDC 381), growth in L-DOPA for 7 days resulted in decreased CSH compared to the controls
grown without L-DOPA (Figure 6K). In CDC 388 and MMC1, we did not see a change in CSH following
melanization, although these strains had notably high levels of hydrophobicity at ~90%.
251
Supernatant Melanization is Resistant to Denaturing Conditions
To further elucidate the identity of the melanin-producing components, we sought to characterize
properties and constraints on supernatant melanization. We first assessed the supernatant's melanization
activity after exposure to high temperatures. We observed a modest reduction in the supernatant's
capacity to melanize after a 1 h incubation at 100˚C, and no significant difference between the
melanization activity of the precipitate or supernatant that formed after boiling (Figure 7A). After treating
samples with the serine proteinases Trypsin or Proteinase K, we assessed the supernatant melanization
activity and observed no reduction in the capacity for melanization (Figure 7B-D). Further, melanization
activity was unaffected by treatment with 1% w/v SDS or methanol; however, melanization was increased
after treatment with 6 M Urea in all conditions, indicating the urea affected the melanization reaction
irrespective of the melanizing strain-specific factor in the supernatant (Figure 7E). These observations
argue against the melanization factor being a protein.
We sought to characterize the size of the melanin-producing components. After passing the
supernatant with a 3 kDa protein cut-off filter, we found that melanization activity was partially conserved
in the flowthrough of the concentrator, suggesting that the melanization factor was not larger than 3 kDa.
To check whether a melanin-producing component was being protected from denaturation within
EVs, we treated supernatant with Triton X-100 and SDS in combination with proteinases. If the
melanization factor was protected within EVs, adding detergent will disturb the EV integrity and allow the
protease to degrade the protein contents within (41–43). We did not find that EVs protect a melanizing
factor from proteolytic degradation (Figure 7G). This was further supported by a lack of enriched
melanization activity observed in an EV-enriched ultracentrifuged pellet (Supplementary Figure 2).
273
Melanization Genes in the *C. auris* Genome
We searched the C. auris genome of enzymes that have been associated with melanin synthesis
in other fungi such as laccases, phenol oxidases and tyrosinases. We found a number of hypothetical

multicopper oxidase and ferroxidase genes reported in the *C. auris* genome. However, we have no
evidence that these have laccase or tyrosinase-like functions and appear similar to metal ion transporters,
peroxisomal membrane components, and ergosterol synthesis (Supplementary Table 1 and 2). Hence,
the absence of tyrosinase and laccase activity in *C. auris* corresponds with the absence of a clearly
identifiable enzyme in the genome.
282
Supernatant Neutral pH Correlated with *C. auris* Strains' Abilities to Melanize
Due to the smaller crystal size of the CFW in the supernatants from the melanizing strain, we
measured the pH of the supernatants since alkaline pH can enhance the solubility of CFW solutions
(44,45). Additionally, the urea treatment, which produced an alkaline environment, increased supernatant
melanization in all the strains. We found a strong correlation between the pH of the conditioned media
and the degree to which the cultures melanized (Figure 8A). We found that the melanizing strains have a
supernatant of around 6.5, whereas the non-melanizing strains tended to have a supernatant closer to pH
5.75. This change in pH was in the range in which L-DOPA and other catecholamines auto-oxidize to
melanin (Figure 8B).
We altered the supernatants' pHs to be either pH 5.5 or 7 by adding hydrochloric acid or
potassium hydroxide. We found that melanization activity was halted at pH 5.5, including the
supernatants of CDC 387 and CDC 388, but was enhanced at pH 7 (Figure 8C, D). Most notably, the
supernatant from CDC 381 – a non-melanizing strain – gained melanization activity at higher pH to an
extent statistically indistinguishable from that of the melanizing strains – CDC 387 and CDC 388 at the
same pH. This strongly suggests that the pH of the supernatant was a major contributor to the
extracellular melanization activity observed.
As endogenous production of ammonia has been reported in *C. albicans* under stress conditions
(46), we assessed the ammonia concentrations of supernatants using a commercially available
colorimetric assay that specifically detects ammonia (47). We observed higher ammonia concentrations in
supernatants from *C. auris* strains that melanized and had higher pH (Figure 8E,F).
We collected the oxidized L-DOPA from the supernatants of CDC 381, CDC 387, and CDC 388,
and minimal media alone at their baseline pH (~5.75, 6.5, 6.5, and 5.5, respectively) and at pH 7. We
imaged these melanized particles using negative staining TEM. Interestingly, we found that the melanized
particles from CDC 387 and CDC 388 had structural differences compared with the oxidized L-DOPA in
the minimal media at pH 7 (Figure 8G). The particles collected from the supernatant with oxidized L-
DOPA were similar to those collected from cultures, whereas the oxidized L-DOPA in minimal media
alone had smaller particles, rougher clumps of electron dense material (Figure 8G). This suggested that
there is some component in the supernatant that may help in structuring the auto-polymerized L-DOPA,
by acting as a scaffold, which allows it to form spherical and linear structures like beads on a string.
Melanin is known to strongly bind to carbohydrates, lipids, and proteins, which may be factors (19,23).
313
Melanin Protects *C. auris* from Oxidative Damage
C. auris cultures incubated, or not, with L-DOPA for 7 days were treated with hydrogen peroxide
(H_2O_2) and yeast viability was evaluated (Figure 9A). The incubation with H_2O_2 reduced the viability of all
evaluated strains of *C. auris*. The melanin-producing strains (CDC 387 and CDC 388) incubated with L-
DOPA were partially protected from the H_2O_2, when compared to cells from the same strain grown in the
absence of L-DOPA. Strain CDC 381 showed similar viability in the presence and in the absence of L-
DOPA, indicating that melanin protects *C. auris* yeast cells against oxidative attack by H_2O_2.
321
Melanin Does Not Protect *C. auris* Against Macrophage Killing

Melanized and control *C. auris* yeast cells were incubated with bone marrow-derived
macrophages (BMDM) for 2 and 24 hours and yeast killing was addressed. In the tested conditions
melanin did not confer protection to the yeast cells against macrophage killing (Figure 9B). It was
interesting to observe that whereas the killing in 2 hours was similar among the strains, in 24 hours CDC
381 was shown to be partially resistant against macrophage killing, as opposed to the susceptible
phenotype showed by strains CDC 387 and CDC 388.
329
330   **Melanin Does Not Affect *C. auris* Virulence During *Galleria mellonella in vivo* Infection**
331   Larvae of *G. mellonella* were infected with yeast cells from melanized and control *C. auris* and
332   survival was evaluated. As seen for the killing experiment with mouse macrophages, the presence of
333   melanin did not impact the survival of *Galleria mellonella* larvae (Figure 9C), suggesting that melanization
334   of *C. auris* prior to infection did not confer protection to the fungus in this invertebrate model.
335
Discussion:
In this paper, we have investigated the ability of *C. auris* to produce melanin, a multifunctional
pigment found across biology and in many pathogenic fungi. Some of the functions of melanin include the
ability to neutralize reactive oxygen species during immune activation, inactivate antimicrobial peptides,
and inactivate antifungal drugs (12,13,48). Due to these immune and treatment evasive properties, our
findings that *C. auris* can melanize may have clinical relevance, particularly in regard to our
understanding of how this fungus is treated.
We evaluated the ability of sixteen *C. auris* strains to melanize when grown in the presence of L-
DOPA, a common melanin substrate. We found that most strains melanized, manifested by darkening of
cultures, and that the ability to melanize was enhanced at higher temperatures (37˚C compared to 30˚C).
We had anticipated that melanization would occur similarly in all strains, so it was surprising that only
some strains had the ability to melanize. Further, based on our understanding of fungal melanization*,* we
expected melanization to occur more at 30˚C rather than 37˚C, as it does in other fungi such as
Cryptococcus neoformans (49). This also has relevancy for pathogenesis in humans, as human body
temperature is 37˚C and thus more conducive to ideal *C. auris* melanization conditions. Interestingly, the
pattern of melanization correlated with the clade in which the strain belonged to; *C. auris* strains from
Clade I and IV, which are associated with South Asia and South America, respectively, are the melanizing
strains, while the non-melanizing strains are from Clade II and III, which are typically associated with East
Asia and South Africa, respectively. This could point to some lineage-specific genetic or epigenetic
differences that are responsible for the melanization phenotype or regulating the melanization process.
Another aspect that differed from the pigmentation observed in other fungi was the impact of cell density
on melanization by *C. auris*. We observed that melanization in *C. auris* required high cell density (10^7
cells/mL). However, at an intermediate cell density (10^6 cells/mL) only some strains of *C. auris* (CDC 382,
385, 386, and 389) were able to produce melanin in a delayed fashion. This observation could indicate
the need of a quorum sensing molecule to trigger effective melanization in *C. auris*.
In addition, we tested the substrate specificity of the melanizing abilities of *C. auris.* We found that
C. auris produces melanins consistent with typical DOPA-melanins using catecholamine substrates such
as L-DOPA, D-DOPA, Methyl-DOPA, dopamine, epinephrine, and norepinephrine. *C. auris* was unable to
produce melanin from L-Tyrosine indicating there is no tyrosinase present, nor are there enzymes
capable of producing pyomelanin from a tyrosine degradation pathway. The *C. auris* genome has
hypothetical genes related to tyrosinase, but these genes appear more related to ergosterol synthesis or
magnesium/zinc ion transport. Similarly, *C. auris* was unable to form pyomelanin from homogentisic acid
(HGA) and 4-hydroxyphenylpyruvic acid (4-HPP). Laccases, such as those in *C. neoformans*, are able to
produce pyomelanin from HGA (34). These results in *C. auris* would suggest that a laccase does not exist

as it does in *C. neoformans,* or at least not one with as much of a wide-ranging substrate range.
Melanization in other DOPA-melanin producing fungi is often catalyzed by laccases (50) or multicopper
ferroxidases (11). Current annotation of the *C. auris* genome indicate there are hypothetical laccases,
oxidoreductase, and ferroxidases (Ontology Groups OG6_100257 and OG6_100380), however these
appear to be related to metal ion homeostasis and part of the peroxisomal membrane. Since we do not
know the function or expression pattern of these hypothetical genes, we cannot rule out that they are
capable of oxidizing catecholamines under certain conditions. However, our evidence points to a non-
laccase, non-tyrosinase, and general non-enzymatic mechanism as the responsible melanin-producing
factor.
We investigated the cellular localization of the melanin in three selected *C. auris* strains: CDC
381 which is a non-melanizing strain, CDC 387 which is a strongly melanizing strain, and CDC 388 which
is a moderately melanizing strain that mostly produces pigment at 37˚C. For some experiments, we
included the MMC1 strain that, like CDC 388, is a moderately melanizing strain. Using light microscopy,
we observed numerous dark aggregates on the outside of the melanized CDC 388, MMC, and particularly
CDC 387 cells, which tended to have a fluffy appearance and appeared in between areas where cells
were clumped together. In addition, the cells themselves appeared darker, and some had large
intracellular melanized spots which would appear to correspond to a vacuole. Based on our evidence that
melanization occurs extracellularly due to the alkalinization of the supernatant, we believe these dark
spots within the vacuole were due to L-DOPA autoxidation within the vacuole following storage under
nutrient-deprived conditions. In the CDC 381 strain, these features were minimal to non-existent. To
further gain understanding of the melanin on an ultrastructural level, we used TEM and SEM. In essence
we found that there were electron dense spherical structures on the surface of the melanized *C. auris,*
which were bound to the outermost layer of the cell wall or were associated with the cell wall. These
structures were only present in melanized cells, and were consistent with previously reported melanin
granules (9,19). Interestingly, these structures were not reliably seen intracellularly, and were significantly
smaller than the extracellular melanin structures seen during *C. albicans* melanization (9). The CDC 387
cells had a distinct phenotype where there were many multivesicular bodies (MVBs) within the cell and
merging with the plasma membrane, resulting in secretion of extracellular vesicles. Using SEM, we also
saw the presence of these spherical particles on the surface of the melanized *C. auris* strains, especially
the CDC 387 strains where the surface was heavily decorated in these structures.
One notable observed feature of the heavily melanized *C. auris* CDC 387 strain is that it tended
to form large aggregates of cells, which were melded together by extracellular melanin, as indicated by
light and electron microscopy. This aggregation is notable, as aggregation is a known physical property in
some strains of *C. auris,* outside of the context of melanin. The aggregate phenotype has been
hypothesized to function as a way for the fungus to remain within tissues and evade immune clearance,
and may play a role in biofilm formation and maintenance (51–53). If melanization is inducing a sort of
aggregative phenotype as our data suggests, this could have similar implications in understanding the
role of *C. auris* melanin in pathogenesis.
We that the supernatant of the melanized cultures were quite dark, particularly in the CDC 388
and MMC1 strains, which indicated there was secreted melanin. We also see this on solid agar, where all
the melanizing strains had a halo of melanin around the colonies and the non-melanizing strains do not.
The strongly melanizing strains CDC 385, CDC 386, CDC 387, and CDC 389 have the darkest halos of
melanin surrounding the colonies. The fungal colonies themselves remain white with no change in
pigmentation by 7 days of growth. The notion that the melanin is extracellular and secreted is supported
by the TEM and SEM which show melanized structures on the surface of the cells. Using negative
staining TEM of the isolated melanin from the supernatant, we see melanin granules similar to those seen
in TEM of whole cells, where the CDC 387 granules are the largest, followed by MMC1 and CDC 388.

The CDC 381 granules were the smallest and fewest. When we added the melanized supernatant to non-
melanized cells, including to CDC 381, we found that the melanin adhered to the cells and made the
appearance of a melanized cell pellet. These findings indicated that the extracellular melanin could
adhere to the outside of the cell wall and suggested that melanization occurred externally and aggregates
of the polymer adhere to the cell wall. It also shows that the CDC 381 cell wall is conducive to
melanization and its lack of melanization activity is not due to an inherent deficiency of the cell wall. When
L-DOPA was added to the cell-free supernatant of the cultures, we see that the supernatant from CDC
388 and CDC 387 had melanizing activity whereas the supernatant from CDC 381 did not, again,
indicating that the component responsible for melanization was extracellular.
To determine the extracellular factor responsible for melanization, we did a series of assays on
the cell-free supernatant. First, we determined that the melanization factor was stable at 100˚C, was
resistant to proteolytic degradation by trypsin and Proteinase K and was smaller than 3 kDa. These
observations together indicated that the melanization component was not a protein, but was likely a small
molecule. Further, the melanizing ability of the supernatant is not enriched in the extracellular vesicles
collected through ultracentrifugation, nor is it protected by EVs. We cannot exclude the possibility that
EVs are used as scaffolding for the melanization, as previously found in *C. neoformans* (54). However,
melanization activity is not lost after treatment with methanol, urea, nor SDS detergents.
Interestingly, we found that the ability for a strain to melanize correlated directly to the pH of the
supernatant within minimal media, where non-melanizing strains had a supernatant ~5.75 and melanizing
strains had a pH ~6.5. This strong correlation also corresponds to the steep increase in autoxidation of L-
DOPA between those pH ranges. We additionally found that artificially adjusting the pH of the
supernatant to pH 7 makes all the supernatants melanize, whereas adjusting the pH to 5.5 prevented all
the supernatants from melanizing, even the supernatants from the melanizing strains. These data
suggest that the melanization is due to the ability of some strains to alter the acidity of surrounding
environment more than others. We find strong evidence that this ability to neutralize the media was due to
the production of ammonia by some of the strains. Fungi, including *C. albicans* and *C. neoformans,* are
known to produce ammonia to help persist under pH stress (46,55,56). *C. auris* has been found in
environmental reservoirs within marine saltwater marshes (2). Microbes found in marine wetland
environmental niche are faced with alkaline stress and, as a result, may be naturally more alkaliphilic
(57).
L-DOPA autoxidation is highly dependent on pH (17,18). L-DOPA has an isoelectric point of pH
6.0, meaning that at pH 6 or above, the L-DOPA is more likely to have a deprotonated amine group,
which results in more energetically favorable oxidation, cyclization, and dopachrome formation (18). While
autoxidation of catecholamines could occur in the absence of any fungus, we found that the oxidized L-
DOPA from the supernatant of the *C. auris* strains had a different structure than the autoxidized L-DOPA
from the minimal media alone at a neutral pH. This indicated that there are some properties of the
supernatant that encourage the autoxidized L-DOPA to organize into specific spherical structures. Based
on previous understanding of how melanin interacts with other biological molecules, we hypothesize this
melanin scaffold to be polysaccharide, protein aggregates, or perhaps even lipids. The structure of the
melanin polymer around this scaffolding could change its biophysical properties. This could account for
one reason why there is significantly more melanin associated with the CDC 387 strain – the strain
changes the pH of the supernatant significantly, but the electron micrographs also show the cells have a
large amount of multivesicular bodies and organelles fusing with the plasma membrane, which would
indicate a relative hypersecretion of polysaccharides, proteins, and extracellular vesicles. Interestingly,
the supernatant of the non-melanized CDC 387 strain has a yellow tint to it, which indicates that some
molecules are being released in higher amounts compared with the CDC 381 and CDC 388 strains.

This pH-based mechanism of fungal melanization differs greatly from other ways in which fungi
produce melanin, including *C. albicans* (9–11)*.* In *C. albicans,* melanization is mediated by ferroxidases, a
class of enzymes that are genetically similar to the melanin-producing enzyme laccase in *C. neoformans.*
The knockout of some ferroxidase genes (*FET* genes) results in a loss of melanization phenotype.
We found that the *C. auris* strains, generally, were remarkably hydrophobic, with most strains' cell
surface hydrophobicity (CSH) being in the 90-100% range. Comparatively, the CSH of clinical isolates of
C. albicans strains ranges from about 2 to 41% (38). In *C. albicans,* CSH is reportedly associated with
increased adhesion to epithelial cells, biofilm formation, resistance to neutrophil-mediated killing, and
increased overall virulence (39,58–60). These findings have implications for studying how *C. auris*
interacts with hydrophobic surfaces, and how hydrophobicity may contribute to durable biofilm formations
within hospital environments and medical equipment. We do not find an association between CSH and
ability of the strain to melanize. Since melanin is a hydrophobic polymer, we anticipated that melanization
of cultures would enhance CSH. Surprisingly, we found that growth in minimal media with L-DOPA
reduced the hydrophobicity of the non-melanizing strain CDC 381 and the strong melanizing strain CDC
387, while the CSH of CDC 388 and MMC1 were virtually unchanged. We hypothesize that the
hydrophobic melanin is comparatively less hydrophobic than the hydrophobicity-mediators on the surface
of the *C. auris* cells, and when the melanin binds to the cell wall, it results in the relative decrease of CSH.
In *C. albicans,* CSH is believed to be mediated by exposure of hydrophobic cell wall mannoproteins
(40,61), which could correspond to areas in which melanin is deposited (62).
The antioxidant properties of melanin are partially responsible for its protective properties (63).
Melanized *C. auris* yeast cells were partially protected against hydrogen peroxide, and this protection was
stronger in the strain where melanization is more efficient (CDC 387). This observation is in accordance
with the expected properties of melanin pigments (12,48). However, melanin did not confer any protection
to *C. auris* in an *in vitro* challenge with BMDM. The *in vitro* killing of *C. auris* by BMDM was tested at
distinct times and, in the evaluated conditions, melanin played no role on protecting the fungus from the
phagocytes. The previously described degree of protection from killing, conferred by melanin to
pathogenic fungi varies considerably (18 % to > 36 %) (64), so it is possible that the pattern observed for
C. auris is similar to the one of *C. albicans*, where there is not a clear association between melanin and
virulence (9). Supporting the *in vitro* findings, *in vivo* experiments using the invertebrate *G. mellonella*,
showed that melanin produced by *C. auris* might not protect the fungus against innate immunity, but we
cannot rule out a potential action of melanin in interfering in processes that lead to acquired immunity.
In summary, we found that Clade I, IV, and V strains of *C. auris* grown with catecholamine
substrates produce melanin extracellularly by alkalinizing the media with ammonia, which promoted non-
enzymatic catecholamine oxidation. This extracellular melanin aggregates and binds to the outside of the
C. auris cell wall (Summarized in Figure 12). While we did not find evidence that the melanin formed by *C.*
auris interferes with the effector mechanisms of innate immune cells, nor is it an active melanization
process, our findings leave open the possibility that environment alkalinization could be a new
mechanism by which fungi can drive the production of melanin. It is also possible that melanization may
play a role in the environmental survival of *C. auris*, where the fungus is likely to encounter oxidative
stressors. Not much is known about the environmental niche of *C. auris,* however, the first environmental
isolates have been uncovered and described in the marine marshlands off the Andaman Islands in India
(2).
505
506   **Materials and Methods**
507
***Candida auris* strains and media**

All isolates of *Candida auris* strains were received from the Centers of Disease Control and Prevention

Food and Drug Administration Antimicrobial Resistance Isolate Bank

(https://wwwn.cdc.gov/arisolatebank/Panel/PanelDetail?ID=2), with the exceptions of MMC1 and MMC2,

which were previously described clinical isolates (65). CDC 381 is also known as B11220, CDC 387 is

also known as B8441, and CDC 388 is also known as B11098. All *C. auris* strains were first grown in

Yeast Peptone Dextrose (YPD) broth at 30°C until they reached stationary phase. Cultures were washed

twice and put into Minimal Media (15.0 mM glucose, 10.0 mM $MgSO_4$, 29.4 mM KH_2PO_4, 13.0 mM

glycine, 3.0 M vitamin B_1, pH 5.5) at 37°C at 10^7 cells/ml unless otherwise noted. Cells were grown for 7

days under continuous shaking.

***C. auris* Melanization in Liquid Media**

All *C. auris* strains were grown in minimal media as described above, with the addition of melanin

precursors to be tested. All compounds used for the melanization substrate assay and ABTS (2,2'-azino-

bis(3-ethylbenzothiazoline-6-sulfonic acid) laccase assay were prepared in concentrated stock solutions

and added to the culture at a final concentration of 1 mM except for caspofungin, which was added at 0.5

µg/ml final concentration. For the experiments evaluating the temperature dependence of the *C. auris*

strain melanization, cultures were also grown at 30°C for 7 days in minimal media. After 7 d, cultures were

scanned with a CanoScan9000F scanner at 600 dpi. Mean Gray Value for each well of culture was

determined using the Measure tool on FIJI image processing software (66).

Experiments where cultures were grown on solid agar petri dishes, 20 µl of stationary washed culture

were added to minimal media agar with 1 mM of 3,4, Dihydroxyl-L-phenylalanine (L-DOPA). Plates were

incubated at 37°C for 7 days and imaged with a scanner CanoScan9000F at 600 dpi.

Melanin extraction and electron paramagnetic resonance (EPR)

Melanin was extracted from selected strains of *C. auris* and one strain of *C. neoformans* (H99) as

described (67). Briefly, after growing for 7 days in minimal medium containing L-DOPA (1 mM), cells were

washed with sorbitol/sodium citrate solution, and incubated with Novozyme 234 for 1 hour at 30°C.

Samples were incubated with guanidine thiocyanate for 1 hour, followed by an incubation with HCl 6 M.

After boiling the solution for 1 hour, the pellets were washed and suspended in PBS. Extracted melanin

was submitted to EPR. The EPR spectra of the collected dark particles were registered using a Varian

E112 X-Band model spectrometer with a TE102 resonator and a liquid nitrogen finger Dewar. The EPR

runs were performed under the following parameters that are similar to our fungal melanin studies as

previously described (8): modulation frequency of 100 kHz, modulation amplitude of 1.6 G, center field of

3,310 G, sweep width of 100.0 G, microwave frequency of 9.2 GHz, microwave power of 1.0 mW, time

constant of 0.5 s, and temperature of 77 K .

Extracellular Melanin Isolation

Cultures grown for 7 days with or without L-DOPA as a substrate for melanization were centrifuged at

4,000 x *g* for 5 minutes. The supernatant was removed and filter sterilized through a 0.8 µm syringe filter

(Corning, Corning, NY). To measure the melanization of the supernatant, absorbance measurements

were performed at 492 nm using a SpectraMax iD5 spectrophotometer. For experiments collecting and

analyzing the extracellular melanin particles, the supernatants were then ultracentrifuged at 100,000 x *g*

for 1 hour at 4°C in a Beckman Coulter Optima L-90K UltraCentrifuge. Supernatants were decanted and

the melanized pellet was suspended in PBS.

Supernatant Melanization Activity

To assess phenoloxidase activity of the supernatant, L-DOPA was added to cell-free supernatant from non-melanized cultures. Subsequently, the supernatant was left to incubate for 72 hours at 37° C in darkness. Activity was determined through SpectraMax iD5 spectrophotometer readings at 492 nm taken at 0 and 72 hours and images were taken using the CanoScan9000F flatbed scanner at a resolution of 600 dpi. For the proteinase assays, supernatants were pre-treated with 1:10 Trypsin (Corning) at 37°C, or with approximately 200 μg/ml Proteinase K (New England BioLabs) at 60°C prior to the addition of L-DOPA. To measure the degree of protection conferred by extracellular vesicles against degradation of potential melanin-producing components, supernatants were pre-treated with 0.01% w/v SDS or 0.1% v/v Triton X-100 (Sigma) with or without Proteinase K. Supernatants were then incubated for one hour at 60°C as described. Heat degradation assays were performed by heating supernatant samples to 100°C for one hour prior to testing for melanization activity.

Light Microscopy

C. auris strains were imaged using light microscopy using an Olympus AX70 microscope and 100x objective.

Transmission Electron Microscopy (TEM)

C. auris melanized and non-melanized cells were imaged using TEM as described (25). Briefly, samples 2% (w/v) glutaraldehyde in 0.1 M cacodylate at room temperature for 2 h, followed by overnight incubation in 4% (w/v) formaldehyde, 1% (w/v) glutaraldehyde, and 0.1% PBS overnight at 4 °C. Samples were washed, fixed with 1% osmium tetroxide for 90 minutes, washed with dH$_2$O, serially dehydrated in ethanol, and embedded in SPURRS resin. Thin sections, 60 to 90 nm, were cut with a diamond knife on the Leica Ultracut E Ultramicrotome and picked up with 2x1 mm formvar coated copper slot grids. Grids were stained with 2% uranyl acetate (aq) and 0.4% lead citrate before imaging on a Hitachi 7600 TEM at 80 kV. Images were captured with an AMT XR80 CCD (8 megapixels, 16 bit). Secreted melanin was visualized using negative staining, in which 8 μl of sample was placed on a negative glow discharged 400 mesh ultra-thin carbon-coated grids (EMS CF400-CU-UL) for 30 s, followed by three quick rinses of Tris-buffered Saline (TBS) and staining with 2.5% uranyl acetate. Samples were imaged using a Hitachi 7600 TEM Electron Microscope at 80 kV. Images were captured with an AMT XR80 CCD (8 megapixels, 16 bit). Following image acquisition, diameter measurements of the secreted melanin particles were performed using the measurement tool of FIJI image processing software (66).

Scanning Electron Microscopy (SEM)

Briefly, samples were fixed in 2.5% (v/v) glutaraldehyde in 0.1 M sodium phosphate buffer (pH 7.3) overnight at 4 °C. Samples were placed on a poly-L-lysine coated coverslip (0.01 mg/ml , coated for 5 minutes, and rised twice in dH2O) for 1 hour, then washed, serially dehydrated in ethanol, chemically dried using hexamethyldisilazane (HMDS) and dried overnight in a desiccator. Samples were then placed on a metal sample stub (EMS aluminum 6mm pin, 12.7 mm diameter) with double sided carbon tape (EMS standard carbon adhesive tabs, 12 mm Diameter) and the underside of the coverslip coated with silver paint (EMS silver conductive coating). Samples were coated with 15 nm gold palladium (AuPd), on a Denton Desk III sputter coater before imaging on a ThermoFisher Helios FIB-SEM at 5 kV using an Ion Conversion and Electron (ICE) detector.

Extracellular Melanin Add-back

601 Cell-free melanized supernatant was added to a pellet of non-melanized cells from the same
602 corresponding strain of *C. auris.* Samples were mixed at 37°C for 3 h and pelleted at 4,000 x *g* for 5
603 minutes. Pelleted cells were image compared to control cells treated for 3 hours with the non-melanized
604 supernatant.
605
Cell Wall Disruption Assays
All cultures were grown in minimal media for 7 days with 1 mM L-DOPA. In addition to the L-DOPA, either
5 mM N-acetylglucosamine (GlcNAc) (9,35), 100 μg/ml Calcofluor White (9,35), or 50% MIC of
Caspofungin based on the CDC resistance profile (37) (Antibiotic Resistance Isolate Bank, Centers for
Disease Control and Prevention, Atlanta, GA, USA) was added to the culture at the beginning of the 7 d
incubation. Cultures were collected after 7 d, and supernatant and cells were examined under light and
electron microscopy as described above.
613
614 **pH Measurements and Supernatant pH Alteration**
615 Cultures were grown for 7 days in minimal media at 37°C. Cultures were centrifuged at 4,000 *x* g for 4
616 minutes, and supernatant was filter sterilized with a 0.22 μm PES Filter (Millipore-Sigma). Supernatant pH
617 was determined using a calibrated Fisher Scientific Accumet AB150 pH meter.
618
Cell-free supernatants were isolated, and their pH measured using a calibrated Fisher Scientific Accumet
AB150 pH meter. Samples were split into three groups: unaltered supernatant, supernatant manually
adjusted to pH 5.5 with hydrochloric acid (HCl), and supernatant manually adjusted to pH 7 with
potassium hydroxide (KOH). Titrations with HCl and KOH were performed using the same pH meter set to
continuously read pH. Each sample's three groups were then treated with 1 mM L-DOPA and incubated
at 37°C for 72 hours in darkness. Melanization activity was determined through SpectraMax iD5
spectrophotometer readings at 492 nm and images were taken using the CanoScan9000F flatbed
scanner at a resolution of 600 dpi.
627
Ammonia Quantification
Supernatant ammonia concentration was quantified using the commercially available API Ammonia Test
Kit (API) according to manufacturer's protocol, with test solution volumes scaled down in proportion to the
small volumes tested to maintain the established test solution concentrations. A standard curve of
ammonia concentrations from 16 ppm to 0.25 ppm was constructed by serial diluting 28% ammonium
hydroxide (Sigma) in PBS to enable validation of this modification and colorimetric correlation to known
ppm values. In a 48-well plate, 1 drop of both solutions 1 and 2 were added to 625 μl of cell-free
supernatant, after which the plate was agitated and left to sit for 5 minutes with a lid on. The color
change, from yellow to dark blue-green during the formation of indophenol blue dye, was quantified by
reading 680 nm absorbance using a SpectraMax iD5 spectrophotometer. In addition, plates were imaged
at 600 dpi using the CanoScan9000F flatbed scanner.
639
Cell Surface Hydrophobicity
Cell surface hydrophobicity was measured by Microbial Adhesion to Hexadecane (MATH) assay as
described (68). Briefly, cells were resuspended in PBS to an optical density of 0.2 - 0.4 at 600 nm using
SpectraMax iD5 spectrophotometer. This was measured in triplicate and recorded as the initial value. 3
mL of cells in PBS were added to a glass test tube followed by 400 μl of n-hexadecane. Tubes were
parafilmed and vortexed on high for 45 seconds each and left to settle for 2 minutes. Aliquots of the
aqueous (bottom) layer were carefully removed, and the absorbance at 600 nm was measured as the
final value. Hydrophobicity was calculated as: 100*(Initial Value – Final Value)/(Initial Value).

Oxidative stress

Yeast cells grown in the presence, or absence, of L-DOPA for 7 days at 37 °C were incubated in RPMI medium buffered with MOPS, containing or not, hydrogen peroxide (H_2O_2) 5mM for 3 hours at 37 °C under shaking. Cell suspensions were diluted and plated onto Sabouraud-agar plates for colony-forming units (CFU) counting. The cytotoxic effect of H_2O_2 in each strain was calculated as the number of yeast cells in the presence of H_2O_2, divided by the yeast count in the absence of H_2O_2, multiplied by 100 to be expressed as percentage.

Killing by bone marrow-derived macrophages

Bone marrow cells isolated from C57BL/6 mice were cultivated in RPMI containing 10% of FBS and 20% of L929 supernatant for 7 days with media addition on day 3. At the end of the differentiation time, cells were plated in 96-well plates (10^5 cells per well) and incubated at 37 °C for adherence. Macrophages were challenged with *C. auris* (0.5×10^5 yeast cells per well), grown in the presence or the absence of L-DOPA (1mM). In parallel, *C. auris* was added to wells in the same conditions but without macrophages. After 2 or 24 h, the plates were centrifuged, the supernatant discarded, and the pellets were suspended in sterile distilled H_2O. The suspensions were diluted and plated onto Sabouraud-agar plates and incubated at 30 °C for 24 h for CFU counting. For each experimental group, the percentage of yeast killing was calculated as: yeast + macrophage group divided by yeast in the absence of macrophages.

***Galleria mellonella* infection**

Groups of 10 insects (250-300 mg) in the final instar larval stage were used. Larvae were injected with melanized or non-melanized yeasts of *C. auris* (2×10^6 cells in 10 μL) into the haemocoel through the last left pro-leg strains using a Hamilton syringe. For these experiments the strains CDC 381, CDC 387 and CDC 388 were selected. PBS was used as a control. The insects were then placed in sterile Petri dishes and maintained in the dark at 37 °C. The numbers of living larvae were monitored twice daily and recorded for a period of 7 days. Larvae were considered dead if no response to physical stimulus was observed.

Bioinformatics Search of Laccase and Tyrosinase Genes in *C. auris*

The keywords "laccase" and "tyrosinase" were searched separately in FungiDB.org. Results were filtered and narrowed down to genes, and to those found in the taxonomic subset "[Candida] auris". Results were then exported and downloaded as a comma-delineated file (.csv) including the following table columns: Organism, GeneID, Product Description, Ortholog Group, EC numbers from OrthoMCL, Computed GO Component IDs, Computed GO Components, Computed GO Function IDs, Computed GO Functions, Computed GO Process IDs, Computed GO Processes, Interpro ID, Interpro Description, PFam ID, PFam Description, Superfamily ID, and Superfamily Description.

Acknowledgements:

We would like to thank the Johns Hopkins University School of Medicine Microscope Facility for their assistance with the electron microscopy experiments, and Rosanna Baker for her suggestions and assistance with the ammonia experiments. We would also like to thank Dr. Christine Chrissian for her helpful feedback during the writing process. The illustration in Figure 10 was made using Biorender.com. DFQS, NM, and AC were supported by National Institute of Allergy and Infection Disease R01 AI052733, R01 AI152078, and R01 AI052733. and the National Heart, Lung, and Blood Institute R01 HL059842. D.F.Q.S. was funded by National Institutes of Health 5T32GM008752-18 and 1T32AI138953-01A1. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the

695 manuscript. DZM and JDN were supported by National Institutes of Health (NIH)–National Institute of
696 Allergy and Infectious Diseases grant R21 AI124797. LN and LH were supported by grants from the
697 Brazilian agency Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq; grants
698 311179/2017-7 and 408711/2017-7) and FAPERJ (E-26/202.809/2018).
699

References
1. Casadevall A, Kontoyiannis DP, Robert V. On the Emergence of Candida auris: Climate Change,
Azoles, Swamps, and Birds. mBio [Internet]. 2019 Aug 27 [cited 2020 Sep 29];10(4). Available from:
https://mbio.asm.org/content/10/4/e01397-19

2. Arora P, Singh P, Wang Y, Yadav A, Pawar K, Singh A, et al. Environmental Isolation of Candida
auris from the Coastal Wetlands of Andaman Islands, India. mBio [Internet]. 2021 Mar 16 [cited
2022 Jan 23]; Available from: https://journals.asm.org/doi/abs/10.1128/mBio.03181-20

3. Ahmad S, Alfouzan W. Candida auris: Epidemiology, Diagnosis, Pathogenesis, Antifungal
Susceptibility, and Infection Control Measures to Combat the Spread of Infections in Healthcare
Facilities. Microorganisms. 2021 Apr;9(4):807.

4. Lee WG, Shin JH, Uh Y, Kang MG, Kim SH, Park KH, et al. First Three Reported Cases of
Nosocomial Fungemia Caused by Candida auris. Journal of Clinical Microbiology [Internet]. 2011
Sep [cited 2022 Jan 24]; Available from: https://journals.asm.org/doi/abs/10.1128/JCM.00319-11

5. Satoh K, Makimura K, Hasumi Y, Nishiyama Y, Uchida K, Yamaguchi H. Candida auris sp. nov., a
novel ascomycetous yeast isolated from the external ear canal of an inpatient in a Japanese
hospital. Microbiol Immunol. 2009 Jan;53(1):41–4.

6. Horton MV, Nett JE. Candida auris infection and biofilm formation: going beyond the surface. Curr
Clin Microbiol Rep. 2020 Sep;7(3):51–6.

7. Garcia-Bustos V, Cabanero-Navalon MD, Ruiz-Saurí A, Ruiz-Gaitán AC, Salavert M, Tormo MÁ, et
al. What Do We Know about Candida auris? State of the Art, Knowledge Gaps, and Future
Directions. Microorganisms. 2021 Oct 19;9(10):2177.

8. Almeida-Paes R, Figueiredo-Carvalho MH, da Silva LB, Gerfen G, S Araújo GR de, Frases S, et al.
Candida glabrata produces a melanin-like pigment that protects against stress conditions
encountered during parasitism. Future Microbiology. 2021 May 1;16(7):509–20.

9. Walker CA, Gómez BL, Mora-Montes HM, Mackenzie KS, Munro CA, Brown AJP, et al. Melanin
Externalization in Candida albicans Depends on Cell Wall Chitin Structures. Eukaryotic Cell
[Internet]. 2010 Sep [cited 2022 Jan 19]; Available from:
https://journals.asm.org/doi/abs/10.1128/EC.00051-10

10. Morris-Jones R, Gomez BL, Diez S, Uran M, Morris-Jones SD, Casadevall A, et al. Synthesis of
Melanin Pigment by Candida albicans In Vitro and during Infection. Infection and Immunity. 2005
Sep 1;73(9):6147–50.

11. Dai B, Xu Y, Gao N, Chen J. Wor1-regulated ferroxidases contribute to pigment formation in opaque
cells of Candida albicans. FEBS Open Bio. 2021;11(3):598–621.

12. Smith DFQ, Casadevall A. The Role of Melanin in Fungal Pathogenesis for Animal Hosts. In:
Rodrigues ML, editor. Fungal Physiology and Immunopathogenesis [Internet]. Cham: Springer
International Publishing; 2019 [cited 2020 Sep 24]. p. 1–30. (Current Topics in Microbiology and
Immunology). Available from: https://doi.org/10.1007/82_2019_173

13. Nosanchuk JD, Casadevall A. The contribution of melanin to microbial pathogenesis. Cellular
Microbiology. 2003;5(4):203–23.

14. Jiang N, Sun N, Xiao D, Pan J, Wang Y, Zhu X. A copper-responsive factor gene CUF1 is required
for copper induction of laccase in Cryptococcus neoformans. FEMS Microbiology Letters. 2009 Jul
1;296(1):84–90.

15. Mauch RM, Cunha V de O, Dias ALT. THE COPPER INTERFERENCE WITH THE
MELANOGENESIS OF *Cryptococcus neoformans*. Rev Inst Med trop S Paulo. 2013 Apr;55:117–
20.

16. Cordero RJB, Camacho E, Casadevall A. Melanization in Cryptococcus neoformans Requires
Complex Regulation. mBio [Internet]. 2020 Feb 4 [cited 2022 Jan 25]; Available from:
https://journals.asm.org/doi/abs/10.1128/mBio.03313-19

17. Chen Y chieh, Kendall T, Yip P, Davy A, Sefcik J, Sutter JU. Influence of Ions and pH on the
Formation of Solid- and Liquid-like Melanin. ACS Omega. 2020 Oct 6;5(39):25059–68.

18. Eslami M, Zare HR, Namazian M. Thermodynamic Parameters of Electrochemical Oxidation of l-
DOPA: Experimental and Theoretical Studies. J Phys Chem B. 2012 Oct 18;116(41):12552–7.

19. Camacho E, Vij R, Chrissian C, Prados-Rosales R, Gil D, O'Meally RN, et al. The structural unit of
melanin in the cell wall of the fungal pathogen Cryptococcus neoformans. Journal of Biological
Chemistry. 2019 Jul 5;294(27):10471–89.

20. Franzen AJ, Cunha MML, Miranda K, Hentschel J, Plattner H, da Silva MB, et al. Ultrastructural
characterization of melanosomes of the human pathogenic fungus Fonsecaea pedrosoi. Journal of
Structural Biology. 2008 Apr 1;162(1):75–84.

21. Baker LG, Specht CA, Donlin MJ, Lodge JK. Chitosan, the Deacetylated Form of Chitin, Is Necessary
for Cell Wall Integrity in Cryptococcus neoformans. Eukaryotic Cell [Internet]. 2007 May [cited 2022
Jan 24]; Available from: https://journals.asm.org/doi/abs/10.1128/EC.00399-06

22. Walton FJ, Idnurm A, Heitman J. Novel gene functions required for melanization of the human
pathogen Cryptococcus neoformans. Molecular Microbiology. 2005;57(5):1381–96.

23. Chrissian C, Camacho E, Kelly JE, Wang H, Casadevall A, Stark RE. Solid-state NMR spectroscopy
identifies three classes of lipids in Cryptococcus neoformans melanized cell walls and whole fungal
cells. Journal of Biological Chemistry. 2020 Oct 30;295(44):15083–96.

24. Chrissian C, Lin CPC, Camacho E, Casadevall A, Neiman AM, Stark RE. Unconventional
Constituents and Shared Molecular Architecture of the Melanized Cell Wall of C. neoformans and
Spore Wall of S. cerevisiae. Journal of Fungi. 2020 Dec;6(4):329.

25. Eisenman HC, Nosanchuk JD, Webber JBW, Emerson RJ, Camesano TA, Casadevall A.
Microstructure of Cell Wall-Associated Melanin in the Human Pathogenic Fungus Cryptococcus
neoformans. Biochemistry. 2005 Mar 1;44(10):3683–93.

26. Jacobson ES, Tinnell SB. Antioxidant function of fungal melanin. Journal of Bacteriology. 1993 Nov
1;175(21):7102–4.

27. Alviano DS, Franzen AJ, Travassos LR, Holandino C, Rozental S, Ejzemberg R, et al. Melanin from
Fonsecaea pedrosoi Induces Production of Human Antifungal Antibodies and Enhances the

Antimicrobial Efficacy of Phagocytes. Infection and Immunity [Internet]. 2004 Jan [cited 2022 Jan
24]; Available from: https://journals.asm.org/doi/abs/10.1128/IAI.72.1.229-237.2004

28. Doering TL, Nosanchuk JD, Roberts WK, Casadevall A. Melanin as a potential cryptococcal defence
against microbicidal proteins. Med Mycol. 1999 Jun;37(3):175–81.

29. Rosas ÁL, Casadevall A. Melanization decreases the susceptibility of Cryptococcus neoformans to
enzymatic degradation. Mycopathologia. 2001 Aug 1;151(2):53–6.

30. Stappers MHT, Clark AE, Aimanianda V, Bidula S, Reid DM, Asamaphan P, et al. Recognition of
DHN-melanin by MelLec, is required for protective immunity to Aspergillus. Nature. 2018 Mar
15;555(7696):382–6.

31. Nosanchuk JD, Rosas AL, Casadevall A. The Antibody Response to Fungal Melanin in Mice. The
Journal of Immunology. 1998 Jun 15;160(12):6026–31.

32. Rosas ÁL, Nosanchuk JD, Casadevall A. Passive Immunization with Melanin-Binding Monoclonal
Antibodies Prolongs Survival of Mice with Lethal Cryptococcus neoformans Infection. Infection and
Immunity [Internet]. 2001 May 1 [cited 2022 Jan 24]; Available from:
https://journals.asm.org/doi/abs/10.1128/IAI.69.5.3410-3412.2001

33. Baker RP, Chrissian C, Stark RE, Casadevall A. Cryptococcus neoformans melanization incorporates
multiple catecholamines to produce polytypic melanin. Journal of Biological Chemistry [Internet].
2022 Jan 1 [cited 2022 Jan 25];298(1). Available from: https://www.jbc.org/article/S0021-
9258(21)01329-6/abstract

34. Frases S, Salazar A, Dadachova E, Casadevall A. Cryptococcus neoformans Can Utilize the
Bacterial Melanin Precursor Homogentisic Acid for Fungal Melanogenesis. Applied and
Environmental Microbiology [Internet]. 2007 Jan [cited 2022 Jan 20]; Available from:
https://journals.asm.org/doi/abs/10.1128/AEM.01947-06

35. Perez-Dulzaides R, Camacho E, Cordero RJB, Casadevall A. Cell-wall dyes interfere with
Cryptococcus neoformans melanin deposition. Microbiology (Reading). 2018 Aug;164(8):1012–22.

36. Camacho E, Chrissian C, Cordero RJB, Liporagi-Lopes L, Stark RE, Casadevall A. N-
acetylglucosamine affects Cryptococcus neoformans cell-wall composition and melanin
architecture. Microbiology (Reading). 2017 Nov;163(11):1540–56.

37. Panel Details | Antimicrobial Resistance Isolate Bank | Antibiotic/Antimicrobial Resistance | CDC
[Internet]. [cited 2022 Jan 19]. Available from:
https://wwwn.cdc.gov/ARIsolateBank/Panel/PanelDetail?ID=2

38. Raut J, Rathod V, Karuppayil SM. Cell Surface Hydrophobicity and Adhesion: A Study on Fifty
Clinical Isolates of Candida albicans. Nippon Ishinkin Gakkai Zasshi. 2010;51(3):131–6.

39. Danchik C, Casadevall A. Role of Cell Surface Hydrophobicity in the Pathogenesis of Medically-
Significant Fungi. Frontiers in Cellular and Infection Microbiology [Internet]. 2021 [cited 2022 Jan
24];10. Available from: https://www.frontiersin.org/article/10.3389/fcimb.2020.594973

40. Krasowska A, Sigler K. How microorganisms use hydrophobicity and what does this mean for human
needs? Frontiers in Cellular and Infection Microbiology [Internet]. 2014 [cited 2022 Jan 24];4.
Available from: https://www.frontiersin.org/article/10.3389/fcimb.2014.00112

41. Wolf JM, Rivera J, Casadevall A. Serum albumin disrupts Cryptococcus neoformans and Bacillus
anthracis extracellular vesicles. Cell Microbiol. 2012 May;14(5):762–73.

42. Coelho C, Vij R, Smith DQ, Brady NR, Hamacher-Brady A, Casadevall A. Study of Microbial
Extracellular Vesicles:Separation by Density Gradients, Protection Assays and Labelling for Live
Tracking. Bio Protoc. 2020 Jan 20;10(2):e3502.

43. Coelho C, Brown L, Maryam M, Vij R, Smith DFQ, Burnet MC, et al. Listeria monocytogenes
virulence factors, including listeriolysin O, are secreted in biologically active extracellular vesicles. J
Biol Chem. 2019 Jan 25;294(4):1202–17.

44. Kim S, Inglett GE. Molecular weight and ionic strength dependence of fluorescence intensity of the
Calcofluor/β-glucan complex in flow-injection analysis. Journal of Food Composition and Analysis.
2006 Aug 1;19(5):466–72.

45. Wu J, Deng X, Tian B, Wang L, Xie B. Interactions between Oat β-Glucan and Calcofluor
Characterized by Spectroscopic Method. J Agric Food Chem. 2008 Feb 1;56(3):1131–7.

46. Vylkova S, Carman AJ, Danhof HA, Collette JR, Zhou H, Lorenz MC. The Fungal Pathogen Candida
albicans Autoinduces Hyphal Morphogenesis by Raising Extracellular pH. mBio [Internet]. 2011 Jul
1 [cited 2020 Sep 26];2(3). Available from: https://mbio.asm.org/content/2/3/e00055-11

47. Zadorojny C, Saxton S, Finger R. Spectrophotometric Determination of Ammonia. Journal (Water
Pollution Control Federation). 1973;45(5):905–12.

48. Liu S, Youngchim S, Zamith-Miranda D, Nosanchuk JD. Fungal Melanin and the Mammalian Immune
System. Journal of Fungi. 2021 Apr;7(4):264.

49. Jacobson ES, Emery HS. Temperature regulation of the cryptococcal phenoloxidase. Journal of
Medical and Veterinary Mycology. 1991 Jan 1;29(2):121–4.

50. Salas SD, Bennett JE, Kwon-Chung KJ, Perfect JR, Williamson PR. Effect of the laccase gene
CNLAC1, on virulence of Cryptococcus neoformans. J Exp Med. 1996 Aug 1;184(2):377–86.

51. Ben-Ami R, Berman J, Novikov A, Bash E, Shachor-Meyouhas Y, Zakin S, et al. Multidrug-Resistant
Candida haemulonii and C. auris, Tel Aviv, Israel. Emerg Infect Dis. 2017 Feb;23(2):195–203.

52. Rossato L, Colombo AL. Candida auris: What Have We Learned About Its Mechanisms of
Pathogenicity? Frontiers in Microbiology [Internet]. 2018 [cited 2022 Jan 20];9. Available from:
https://www.frontiersin.org/article/10.3389/fmicb.2018.03081

53. Bravo Ruiz G, Lorenz A. What do we know about the biology of the emerging fungal pathogen of
humans Candida auris? Microbiological Research. 2021 Jan 1;242:126621.

54. Eisenman HC, Frases S, Nicola AM, Rodrigues ML, Casadevall A. Vesicle-associated melanization
in Cryptococcus neoformans. Microbiology (Reading). 2009 Dec;155(Pt 12):3860–7.

55. Fu MS, Coelho C, Leon-Rodriguez CMD, Rossi DCP, Camacho E, Jung EH, et al. Cryptococcus
neoformans urease affects the outcome of intracellular pathogenesis by modulating
phagolysosomal pH. PLOS Pathogens. 2018 Jun 15;14(6):e1007144.

56. Vylkova S, Lorenz MC. Modulation of Phagosomal pH by Candida albicans Promotes Hyphal
Morphogenesis and Requires Stp2p, a Regulator of Amino Acid Transport. PLOS Pathogens. 2014
Mar 13;10(3):e1003995.

57. Meena B, Rajan LA, Vinithkumar NV, Kirubagaran R. Novel marine actinobacteria from emerald
Andaman & Nicobar Islands: a prospective source for industrial and pharmaceutical byproducts.
BMC Microbiol. 2013 Jun 22;13:145.

58. Singleton DR, Fidel PL Jr, Wozniak KL, Hazen KC. Contribution of cell surface hydrophobicity protein
1 (Csh1p) to virulence of hydrophobic Candida albicans serotype A cells. FEMS Microbiology
Letters. 2005 Mar 1;244(2):373–7.

59. Hazen KC, Brawner DL, Riesselman MH, Jutila MA, Cutler JE. Differential adherence of hydrophobic
and hydrophilic Candida albicans yeast cells to mouse tissues. Infection and Immunity [Internet].
1991 Mar [cited 2022 Jan 24]; Available from: https://journals.asm.org/doi/abs/10.1128/iai.59.3.907-
912.1991

60. Silva-Dias A, Miranda IM, Branco J, Monteiro-Soares M, Pina-Vaz C, Rodrigues AG. Adhesion,
biofilm formation, cell surface hydrophobicity, and antifungal planktonic susceptibility: relationship
among Candida spp. Front Microbiol [Internet]. 2015 [cited 2020 Sep 28];6. Available from:
https://www.frontiersin.org/articles/10.3389/fmicb.2015.00205/full?utm_source=newsletter&utm_me
dium=web&utm_campaign=Microbiology-w15-2015

61. Fukazawa Y, Kagaya K. Molecular bases of adhesion of Candida albicans. Journal of Medical and
Veterinary Mycology. 1997 Jan 1;35(2):87–99.

62. Tronchin G, Pihet M, Lopes-Bezerra LM, Bouchara JP. Adherence mechanisms in human pathogenic
fungi. Med Mycol. 2008 Dec 1;46(8):749–72.

63. Mattoon ER, Cordero RJB, Casadevall A. Fungal Melanins and Applications in Healthcare,
Bioremediation and Industry. J Fungi (Basel). 2021 Jun 18;7(6):488.

64. Nosanchuk JD, Casadevall A. Impact of melanin on microbial virulence and clinical resistance to
antimicrobial compounds. Antimicrob Agents Chemother. 2006 Nov;50(11):3519–28.

65. Zamith-Miranda D, Heyman HM, Cleare LG, Couvillion SP, Clair GC, Bredeweg EL, et al. Multi-omics
Signature of Candida auris, an Emerging and Multidrug-Resistant Pathogen. mSystems [Internet].
2019 Jun 11 [cited 2022 Jan 19]; Available from:
https://journals.asm.org/doi/abs/10.1128/mSystems.00257-19

66. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source
platform for biological-image analysis. Nature Methods. 2012 Jul;9(7):676–82.

67. Chatterjee S, Prados-Rosales R, Itin B, Casadevall A, Stark RE. Solid-state NMR Reveals the
Carbon-based Molecular Architecture of Cryptococcus neoformans Fungal Eumelanins in the Cell
Wall. J Biol Chem. 2015 May 29;290(22):13779–90.

68. Vij R, Danchik C, Crawford C, Dragotakes Q, Casadevall A. Variation in Cell Surface Hydrophobicity
among Cryptococcus neoformans Strains Influences Interactions with Amoebas. mSphere
[Internet]. 2020 Apr 29 [cited 2020 Sep 21];5(2). Available from:
https://msphere.asm.org/content/5/2/e00310-20

Figure Legends:

Figure 1. Effect of Temperature on Melanization. Melanization variably occurs in *C. auris* strains. This melanization can be quantified (A). Melanization occurs less at 30˚C (B) than at 37˚C (C) particularly for the melanizing strains CDC 388, CDC 390, and MMC1. Experiments performed in biological replicates. Panels (B) and (C) are representative images. (D-F) Electronic paramagnetic resonance (EPR) of melanin isolated from *Cryptococcus neoformans* (D), non-melanizing *C. auris* strains (E) and melanizing *C. auris* strains (F) grown in minimal media with L-DOPA. Error bars represent standard deviation from three independent experiments.

Figure 2. Cell density and time course. *C. auris* yeast cells were incubated at increasing cell densities in minimal media in the presence, or absence, of L-DOPA. Melanin formation was monitored through the course of 13 days and quantified using ImageJ. The graphs show means and standard deviation from 3 independent experiments.

Figure 3. Melanization by Substrate. (A) *C. auris* melanizing strains use catecholamines such as L-DOPA, D-DOPA, Methyl-DOPA, dopamine, epinephrine, and norepinephrine as substrates for melanization. Overall, there are few differences between the melanizing strains' abilities to oxidize these substrates. **(B)** The color of the pigments formed were typically brown-black, with the exception of the orange produced with epinephrine. Red dotted line represents the baseline mean gray value of the background. Values above the line (closer to 0) signify wells darker than the background. **(C)** Strains of *C. auris,* CDC 381, 387, and 388 do not have laccase activity when grown with the laccase-specific substrate ABTS, while the positive control – H99 strain of *Cryptococcus neoformans* – does have laccase activity, as indicated by the blue color formed in culture. Data represents biological triplicate, with Panel **(B,C)** showing a representative image of the melanized cultures. Abbreviations: L-Tyrosine (TYR), Methyl-DOPA (M-DOPA), dopamine (DA), epinephrine (EPI), norepinephrine (NOR), Brain mixture (BRAIN), homogentisic acid (HGA), 4-hydroxyphenylpyruvic acid (4-HPP), Caspofungin (CASPO). Error bars indicate standard deviation.

Figure 2. Melanization primarily occurs extracellularly in C. auris. (A-B) The supernatant of melanized cultures have darker supernatants than their non-melanized counterparts, particularly the supernatant from the CDC 388 strain. **(C)** When grown on solid agar, all melanizing strains have a dark halo surrounding their colony, with darker color around strong melanizing strains. The colonies of the strains themselves are notably white. **(D)** The cell-free supernatant from CDC 387 and CDC 388 (melanin-capable strains) have the ability to melanize, indicating a melanization factor is present. **(E-F)** Cultures grown with L-DOPA have melanin particles in the supernatant that can be collected and visualized via electron microscopy. The strongly melanizing CDC 387 strain has the largest population of these particles, the moderate melanizing strains CDC 388 and MMC1 have moderately sized population of these particles, and the weak melanizing strain CDC 381 has the smallest population. All experiments done in biological triplicate. Panels **(A), (C)** and **(F)** are representative images. Scale Bar indicates 500 nm. Error bars indicate standard deviation.

Figure 5. Melanin is Primarily Deposited on the Cell Wall Surface. (A) Light Microscopy shows that the 387, 388, and MMC1 cells have darker color when grown in minimal media with L-DOPA, while there is little color change in the CDC 381 cell color. Some of the pigmented cells have dark pigmentation within the cell. CDC 387 in particular appears to have large pigmented extracellular structures. Scale bars represent 10 μm. Images representative of three biological replicates. **(B)** Transmission electron

microscopy shows electron dense particles on the outside of only the 387, 388, and MMC1 cells grown in
the presence of L-DOPA, which is indicative of melanin pigment. These dark particles are found on the
external cell wall and in the extracellular space, and are especially visible in the CDC 387 and MMC1
micrographs. Scale bars represent 500 nm. **(C)** Similar external granule structures on the surface of CDC
387, 388, and MMC1 cells grown with L-DOPA can be seen in SEM micrographs. Scale bars represent 5
μm.

Figure 6. Extracellular Melanin Adheres to β-glucans. **(A-C)** Adding non-melanized cells to the
melanized supernatant from its corresponding strain resulted in pigmented cells after 3 hours of
incubation. **(D)** The melanized supernatant from the CDC 388 strain causes the non-melanizing CDC 381
strain to become pigmented. Adding the building block for chitin, GlcNAc, to the CDC 387 (**E**) and CDC
388 (**F**) resulted in increased supernatant melanization, blocking chitin synthesis with CFW resulted in
less melanin in supernatant but also caused melanin coated CFW crystals on the fungal surface. Blocking
β-Glucan synthesis with caspofungin resulted in increased supernatant melanization in both strains. **(G-H)**
Electron microscopy revealed no major differences in melanin adherence to the cell wall in either the
CDC 387 **(G)** and CDC 388 **(H)** strains treated with GlcNAc. There were clear electron-dense crystal
structures on the cell walls of CFW conditions, and the caspofungin-treated conditions had little to no
melanin on the cell surface. While caspofungin-treated cells grew in culture, they look atypical in TEM
with a condensed cytoplasm. All scale bars represent 500 nm. **(I,J)** There is no association between the
hydrophobicity of a *C. auris* strain and its ability to melanize. **(K)** Although, the melanization of certain
strains, namely CDC 387, results in a significant reduction of CSH compared to its non-melanized control.
No statistically reduction that happens in CDC 381, 388, or MMC1 strains. All experiments are
representative of three biological replicates. Statistical differences tested with a Two-Way ANOVA with
multiple comparisons using Prism GraphPad, with $p < 0.0001$ = ****. Error bars indicate standard deviation.

Figure 7. C. auris Melanization Factor is Resistant to Denaturing Conditions. **(A)** Heating the
supernatant to 100°C slightly reduces the melanizing ability of the supernatant but does not completely
abolish melanizing ability of the supernatant nor the precipitate that forms following boiling (inset). Adding
proteinase K (**B**) or Trypsin (**C**) does not reduce the melanizing ability of the supernatant. **(D)** Images of
the oxidation after 72 h within the samples treated with proteinase K and trypsin (E) Similarly, 33%
methanol and 1% SDS did not reduce melanization activity, while 6M urea uniformly enhanced DOPA
oxidation in all groups. **(F)** Filtration through a 3 kDa filter indicates that the melanizing factor is smaller
than 3 kDa. **(G)** The melanizing factor is not protected from proteolytic degradation by being shielded
within extracellular vesicles (EV), as addition of detergent to disturb EVs does not render the melanization
factor more suscept to proteolytic degradation. Error bars indicate standard deviation.

Figure 3. Supernatant pH Correlates to Melanization Ability. **(A)** The pH of the supernatant from the
C. auris cultures corresponds to the ability of that strain to melanize, with two distinct clusters that
correspond to melanizing versus non-melanizing strains. **(B)** The correlation between pH and
melanization from the *C. auris* strains corresponds to a similar increase in autoxidation of L-DOPA in the
same range of pH. **(C)** Reducing the pH of the supernatants to 5.5 abolished melanization ability in the
CDC 387 and 388 strains, while raising the pH of the supernatants to 7 enhanced melanization in all the
supernatants, including CDC 381. **(D)** Images of the L-DOPA oxidation in the supernatant following pH
alteration. **(E,F)** *C. auris* strains that melanize are associated with increased ammonia concentration in
the supernatant, with a concentration of approximately 16 ppm, while non-melanizing strains have an
ammonia concentration between 8 and 5 ppm. **(G)** Aggregates from pH 7 melanized cell-free supernatant
from CDC 387 and CDC 388 look similar to those produced in cultures (Figure 4F), whereas the pigment

collected from the pH 7 minimal media looks more amorphous and lacks structure. The Scale Bar
represents 500 nm. Error bars indicate standard deviation.

Figure 9. Melanin from *C. auris* during interaction with the host. (**A**) Melanized and control yeast cells
were incubated with, or without, hydrogen peroxide (H_2O_2) 5 mM for 3 hours at 37° C. Yeast viability was
addressed by CFU counting and values of H_2O_2-treated *C. auris* were expressed as a percentage of cells
in the absence of H_2O_2. Each symbol in the graph represents one experiment performed in triplicates.
Data was treated by One-way ANOVA followed by Sidak's multiple comparison test. (**B**) Bone marrow-
derived macrophages (BMDM) were challenged with melanized or control *C. auris* yeast cells for 2 or 24
hours. After macrophage lysis, fungal viability was addressed by CFU counting. The killing percentage
was expressed as the amount of viable yeast in the presence of macrophages, divided by the amount of
yeast in the absence of macrophages for each strain and condition. Each symbol represents one
experiment performed in triplicates. (**C**) Melanized and control *C. auris* yeast cells were inoculated into
the haemocoel of *G. mellonella* larvae (n = 10/group). The number of living larvae was monitored for 7
days after inoculation.

Figure 10. Proposed Model for Candida auris Melanization. *C. auris* from Clades I and IV can
neutralize the pH of their surrounding environment by secreting either ammonia, an alkalinizing enzyme,
or some other pH modulator. The neutralized pH encourages oxidation of catecholamines present, which
begin to form a pigment. The oxidized catecholamines may aggregate on secreted proteins,
polysaccharides, or lipids to form melanin granules. These melanin granules are deposited on the surface
of the fungi. In cases of high degrees of melanization, nearly all the extracellular melanin ends up
adhering to the surface of the fungus, leaving a relatively light-colored supernatant.

Supplementary Figure 1. Interactions of Calcofluor White and Melanin. (*A*) Melanizing *C. auris*
strains (CDC 387) grown with L-DOPA and Calcofluor white (CFW) have visible crystal formation under
light microscopy. These crystals are dark in color, suggesting the presence of melanin or polymerized L-
DOPA, and they are fluorescent under a 405 nm wavelength laser, indicating they also contain CFW. (B)
These crystals can also be seen under transmission electron microscopy. (C) Adding L-DOPA and CFW
to non-melanized cell-free supernatant resulted in the formation of black crystals, which under the
fluorescent microscope have signal, indicating that the crystals are a combination of both CFW and
melanin/autopolymerized L-DOPA. (D) Additionally, when CFW is added to supernatant, the crystals that
form in CDC 381 strain are large and settle quickly, leaving few crystals in suspension, whereas CDC 387
and 388 form smaller crystals that do not settle rapidly and remain in the suspension. Inset is an image of
the falcon tube containing the supernatant with CFW, and micrographs show a sample taken from the
non-settled suspension. (E) Pseudohyphal morphology seen in the melanized CDC 388 culture treated
with 50% MIC caspofungin for 1 week. Scale bars in A, C, D, and E represent 10 μm, and scale bars in B
represent 500 nm.

1015

1016   ***Supplementary Figure 2. Melanization activity is not associated with Extracellular Vesicles.***
1017   Following ultracentrifugation of the *C. auris*-conditioned media, there is equal melanization activity in both
1018   the supernatant and the pellet (containing extracellular vesicles) resuspended in 200 μl of supernatant,
1019   which represents a ~200x concentration.

1020

1021   ***Supplementary Table 1.*** **List of laccase homologues in *C. auris* from FungiDB**

1022

1023   ***Supplementary Table 2.*** **List of tyrosinase homologues in *C. auris* from FungiDB**

Figure 1. Effect of Temperature on Melanization. Melanization occurs in only some of the *C. auris* strains. This melanization can be quantified (A). Melanization occurs less at 30˚C (B) than at 37˚C (C) particularly for the melanizing strains CDC 388, CDC 390, and MMC1. Experiments performed in biological replicates. Panels (B) and (C) are representative images. (D-F) Electronic paramagnetic resonance (EPR) of melanin isolated from *Cryptococcus neoformans* (D), non-melanizing *C. auris* strains (E) and melanizing *C. auris* strains (F) grown in minimal media with L-DOPA. Error bars represent standard deviation from three independent experiments.

Figure 2. Cell density and time course. *C. auris* yeast cells were incubated at increasing cell densities in minimal media in the presence, or absence, of L-DOPA. Melanin formation was monitored through the course of 13 days and quantified using ImageJ. The graphs show means and standard deviation from 3 independent experiments.

Figure 3. Melanization by Substrate. **(A)** *C. auris* melanizing strains use catecholamines such as L-DOPA, D-DOPA, Methyl-DOPA, dopamine, epinephrine, and norepinephrine as substrates for melanization. Overall, there are few differences between the melanizing strains' abilities to oxidize these substrates. **(B)** The color of the pigments formed were typically brown-black, with the exception of the orange produced with epinephrine. Red dotted line represents the baseline mean gray value of the background. Values above the line (closer to 0) signify wells darker than the background. **(C)** Strains of *C. auris,* CDC 381, 387, and 388 do not have laccase activity when grown with the laccase-specific substrate ABTS, while the positive control – H99 strain of *Cryptococcus neoformans* – does have laccase activity, as indicated by the blue color formed in culture. Data represents biological triplicate, with Panel **(B,C)** showing a representative image of the melanized cultures. Abbreviations: L-Tyrosine (TYR), Methyl-DOPA (M-DOPA), dopamine (DA), epinephrine (EPI), norepinephrine (NOR), Brain mixture (BRAIN), homogentisic acid (HGA), 4-hydroxyphenylpyruvic acid (4-HPP), Caspofungin (CASPO). Error bars indicate standard deviation.

Figure 2. Melanization primarily occurs extracellularly in *C. auris*. **(A-B)** The supernatant of melanized cultures have darker supernatants than their non-melanized counterparts, particularly the supernatant from the CDC 388 strain. **(C)** When grown on solid agar, all melanizing strains have a dark halo surrounding their colony, with darker color around strong melanizing strains. The colonies of the strains themselves are notably white. **(D)** The cell-free supernatant from CDC 387 and CDC 388 (melanin-capable strains) have the ability to melanize, indicating a melanization factor is present. **(E-F)** Cultures grown with L-DOPA have melanin particles in the supernatant that can be collected and visualized via electron microscopy. The strongly melanizing CDC 387 strain has the largest population of these particles, the moderate melanizing strains CDC 388 and MMC1 have moderately sized population of these particles, and the weak melanizing strain CDC 381 has the smallest population. All experiments done in biological triplicate. Panels **(A)**, **(C)** and **(F)** are representative images. Scale Bar indicates 500 nm. Error bars indicate standard deviation.

Figure 5. Melanin is Primarily Deposited on the Cell Wall Surface. (A) Light Microscopy shows that the 387, 388, and MMC1 cells have darker color when grown in minimal media with L-DOPA, while there is little color change in the CDC 381 cell color. Some of the pigmented cells have dark pigmentation within the cell. CDC 387 in particular appears to have large pigmented extracellular structures. Scale bars represent 10 µm. Images representative of three biological replicates. **(B)** Transmission electron microscopy shows electron dense particles on the outside of only the 387, 388, and MMC1 cells grown in the presence of L-DOPA, which is indicative of melanin pigment. These dark particles are found on the external cell wall and in the extracellular space, and are especially visible in the CDC 387 and MMC1 micrographs. Scale bars represent 500 nm. **(C)** Similar external granule structures on the surface of CDC 387, 388, and MMC1 cells grown with L-DOPA can be seen in scanning electron micrographs. Scale bars represent 5 µm.

Figure 6. Extracellular Melanin Adheres to β-glucans. **(A-C)** Adding non-melanized cells to the melanized supernatant from its corresponding strain resulted in pigmented cells after 3 hours of incubation. **(D)** The melanized supernatant from the CDC 388 strain causes the non-melanizing CDC 381 strain to become pigmented. Adding the building block for chitin, GlcNAc, to the CDC 387 **(E)** and CDC 388 **(F)** resulted in increased supernatant melanization, blocking chitin synthesis with CFW resulted in less melanin in supernatant but also caused melanin-coated CFW crystals on the fungal surface. Blocking beta-Glucan synthesis with caspofungin resulted in increased supernatant melanization in both strains. **(G-H)** Electron microscopy revealed no major differences in melanin adherence to the cell wall in either the CDC 387 **(G)** and CDC 388 **(H)** strains treated with GlcNAc. There were clear electron-dense crystal structures on the cell walls of CFW conditions, and the caspofungin-treated conditions had little to no melanin on the cell surface. While caspofungin-treated cells grew in culture, they look atypical in TEM with a condensed cytoplasm. All scale bars represent 500 nm. **(I,J)** There is no association between the hydrophobicity of a *C. auris* strain and its ability to melanize. **(K)** Although, the melanization of certain strains, namely CDC 387, results in a significant reduction of CSH compared to its non-melanized control. No statistically reduction reduction that happens in CDC 381, 388, or MMC1 strains. All experiments are representative of three biological replicates. Statistical differences tested with a Two-Way ANOVA with multiple comparisons using Prism GraphPad, with p<0.0001 = ****. Error bars indicate standard deviation.

Figure 7. C. auris Melanization Factor is Resistant to Denaturing Conditions. (A) Heating the supernatant to 100˚C slightly reduces the melanizing ability of the supernatant but does not completely abolish melanizing ability of the supernatant nor the precipitate that forms following boiling (inset). Adding proteinase K (**B**) or Trypsin (**C**) does not reduce the melanizing ability of the supernatant. **(D)** Images of the oxidation after 72 h within the samples treated with proteinase K and trypsin (E) Similarly, 33% methanol and 1% SDS did not reduce melanization activity, while 6M urea uniformly enhanced DOPA oxidation in all groups. **(F)** Filtration through a 3 kDa filter indicates that the melanizing factor is smaller than 3 kDa. **(G)** The melanizing factor is not protected from proteolytic degradation by being shielded within extracellular vesicles (EV), as addition of detergent to disturb EVs does not render the melanization factor more suscept to proteolytic degradation. Error bars indicate standard deviation.

Figure 3. Supernatant pH Correlates to Melanization Ability. (A) The pH of the supernatant from the *C. auris* cultures corresponds to the ability of that strain to melanize, with two distinct clusters that correspond to melanizing versus non-melanizing strains. **(B)** The correlation between pH and melanization from the *C. auris* strains corresponds to a similar increase in autoxidation of L-DOPA in the same range of pH. **(C)** Reducing the pH of the supernatants to 5.5 abolished melanization ability in the CDC 387 and 388 strains, while raising the pH of the supernatants to 7 enhanced melanization in all the supernatants, including CDC 381. **(D)** Images of the L-DOPA oxidation in the supernatant following pH alteration. **(E,F)** *C. auris* strains that melanize are associated with increased ammonia concentration in the supernatant, with a concentration of approximately 16 ppm, while non-melanizing strains have an ammonia concentration between 8 and 5 ppm. **(G)** Aggregates from pH 7 melanized cell-free supernatant from CDC 387 and CDC 388 look similar to those produced in cultures (Figure 4F), whereas the pigment collected from the pH 7 minimal media looks more amorphous and lacks structure. The Scale Bar represents 500 nm. Error bars indicate standard deviation.

Figure 9. Melanin from *C. auris* during interaction with the host. (**A**) Melanized and control yeast cells were incubated with, or without, hydrogen peroxide (H_2O_2) 5 mM for 3 hours at 37° C. Yeast viability was addressed by CFU counting and values of H_2O_2-treated *C. auris* were expressed as a percentage of cells in the absence of H_2O_2. Each symbol in the graph represents one experiment performed in triplicates. Data was treated by One-way ANOVA followed by Sidak's multiple comparison test. (**B**) Bone marrow-derived macrophages (BMDM) were challenged with melanized or control *C. auris* yeast cells for 2 or 24 hours. After macrophage lysis, fungal viability was addressed by CFU counting. The killing percentage was expressed as the amount of viable yeast in the presence of macrophages, divided by the amount of yeast in the absence of macrophages for each strain and condition. Each symbol represents one experiment performed in triplicates. (**C**) Melanized and control *C. auris* yeast cells were inoculated into the haemocoel of *G. mellonella* larvae (n = 10/group). The number of living larvae was monitored for 7 days after inoculation.

Proposed Model of *Candida auris* Melanization

① *C. auris* from South Asian and South American clades begins to neutralize the pH of the media, either enzymatically or through the production of small molecules when grown at 37°C in minimal media.

② In neutral pH, catecholamines autooxidize in the media and begins to aggregate on polysaccharides or protein aggregates in the supernatant, which act as scaffolding. These aggregates - Melanin granules - begin to adhere to the peripheral cell wall, while the supernatant remains melanized.

③ Cells with abundant adherent melanin begin to clump and form cell aggregates. The supernatant becomes relatively melanin-free, while less melanin-adherent strains retain melanized supernatant.

Figure 10. Proposed Model for Candida auris Melanization. *C. auris* from Clades I and IV are able to neutralize the pH of their surrounding environment by secreting either ammonia, an alkalinizing enzyme, or some other pH modulator. The neutralized pH encourages oxidation of catecholamines present, which begin to form a pigment. The oxidized catecholamines may aggregate on secreted proteins, polysaccharides, or lipids to form melanin granules. These melanin granules are deposited on the surface of the fungi. In cases of high degrees of melanization, nearly all of the extracellular melanin ends up adhering to the surface of the fungus, leaving a relatively light-colored supernatant.

Supplementary Figure 1. Interactions of Calcofluor White and Melanin. (A)Melanizing *C. auris* strains (CDC 387) grown with L-DOPA and Calcofluor white (CFW) have visible crystal formation under light microscopy. These crystals are dark in color, suggesting the presence of melanin or polymerized L-DOPA, and they are fluorescent under a 405 nm wavelength laser, indicating they also contain CFW. (B) These crystals can also be seen under transmission electron microscopy. (C) Adding L-DOPA and CFW to non-melanized cell-free supernatant resulted in the formation of black crystals, which under the fluorescent microscope have signal, indicating that the crystals are a combination of both CFW and melanin/autopolymerized L-DOPA. (D) Additionally, when CFW is added to supernatant, the crystals that form in CDC 381 strain are large and settle quickly, leaving few crystals in suspension, whereas CDC 387 and 388 form smaller crystals that do not settle rapidly and remain in the suspension. Inset is an image of the falcon tube containing the supernatant with CFW, and micrographs show a sample taken from the non-settled suspension. Scale bars in A, C, D, and E represent 10 μm, and scale bars in B represent 500 nm.

Supplementary Figure 2. Melanization activity is not associated with Extracellular Vesicles. Following ultracentrifugation of the *C. auris*-conditioned media, there is equal melanization activity in both the supernatant and the pellet (containing extracellular vesicles) resuspended in 200 μl of supernatant, which represents a ~200x concentration.

Organism	CDC Strain	Gene ID	Product Description	Ortholog Group	EC numbers from OrthoMCL	Computed GO Component IDs	Computed GO Components	Computed GO Function IDs	Computed GO Functions	Computed GO Process IDs	Computed GO Processes	Interpro ID	Interpro Description	PFam ID	PFam Description	Superfamily ID	Superfamily Description
Candida auris strain B11220	381	CJI96_0000827	hypothetical_protein	OG6_100257	1.10.3.2 (Laccase);1.10.3.3 (L-ascorbate oxidase);1.16.3.1 (Ferroxidase);1.7.2.1 (Nitrite reductase (NO-forming));2.7.-.- (Transferring phosphorus-containing groups.);3.2.1.- (Glycosidases, i.e. enzymes hydrolyzing O- and S-glycosyl compounds.);3.2.1.3 (Glucan 1,4-alpha-glucosidase);3.2.1.97 (Endo-a	GO:0016021	integral component of membrane	GO:0005507;GO:0016491	copper ion binding;oxid oreductase activity	GO:000681 1;GO:00550 72;GO:0055 114	ion transport;iron ion homeostasis;obsolete oxidation-reduction process	IPR001117;IPR002355;IPR008972;IPR011706;IPR011707;IPR033138	Multicopper oxidase, type 1;Multicopper oxidase, copper-binding site;Cupredoxin;Multicopper oxidase, type 2;Multicopper oxidase, type 3;Multicopper oxidases, conserved site	PF00394;PF07731;PF07732	Multicopper oxidase, type 1;Multicopper oxidase, type 2;Multicopper oxidase, type 3	SSF49503	Cupredoxin
Candida auris strain B11220	381	CJI96_0001509	hypothetical_protein	OG6_100380	1.10.3.2 (Laccase)	GO:0016021	integral component of membrane	N/A	N/A	N/A	N/A	IPR007248	Mpv17/PMP22	PF04117	Mpv17/PMP22	N/A	N/A
Candida auris strain B11220	381	CJI96_0001510	hypothetical_protein	OG6_100380	1.10.3.2 (Laccase)	GO:0016021	integral component of membrane	N/A	N/A	N/A	N/A	IPR007248	Mpv17/PMP22	PF04117	Mpv17/PMP22	N/A	N/A
Candida auris strain B11220	381	CJI96_0003577	hypothetical_protein	OG6_100257	1.10.3.2 (Laccase);1.10.3.3 (L-ascorbate oxidase);1.16.3.1 (Ferroxidase);1.7.2.1 (Nitrite reductase (NO-forming));2.7.-.- (Transferring phosphorus-containing groups.);3.2.1.- (Glycosidases, i.e. enzymes hydrolyzing O- and S-glycosyl compounds.);3.2.1.3 (Glucan 1,4-alpha-glucosidase);3.2.1.97 (Endo-a	GO:0016021	integral component of membrane	GO:0005507;GO:0016491	copper ion binding;oxid oreductase activity	GO:000681 1;GO:00550 72;GO:0055 114	ion transport;iron ion homeostasis;obsolete oxidation-reduction process	IPR001117;IPR002355;IPR008972;IPR011706;IPR011707;IPR033138	Multicopper oxidase, type 1;Multicopper oxidase, copper-binding site;Cupredoxin;Multicopper oxidase, type 2;Multicopper oxidase, type 3;Multicopper oxidases, conserved site	PF00394;PF07731;PF07732	Multicopper oxidase, type 1;Multicopper oxidase, type 2;Multicopper oxidase, type 3	SSF49503	Cupredoxin
Candida auris strain B11221	383	CJI97_000078	unspecified product	OG6_100380	1.10.3.2 (Laccase)	GO:0016021	integral component of membrane	N/A	N/A	N/A	N/A	IPR007248	Mpv17/PMP22	PF04117	Mpv17/PMP22	N/A	N/A
Candida auris strain B11221	383	CJI97_000519	unspecified product	OG6_100257	1.10.3.2 (Laccase);1.10.3.3 (L-ascorbate oxidase);1.16.3.1 (Ferroxidase);1.7.2.1 (Nitrite reductase (NO-forming));2.7.-.- (Transferring phosphorus-containing groups.);3.2.1.- (Glycosidases, i.e. enzymes hydrolyzing O- and S-glycosyl compounds.);3.2.1.3 (Glucan 1,4-alpha-glucosidase);3.2.1.97 (Endo-a	N/A	N/A	GO:0005507;GO:0016491	copper ion binding;oxid oreductase activity	GO:000681 1;GO:00550 72;GO:0055 114	ion transport;iron ion homeostasis;obsolete oxidation-reduction process	IPR001117;IPR002355;IPR008972;IPR011706;IPR011707;IPR033138	Multicopper oxidase, type 1;Multicopper oxidase, copper-binding site;Cupredoxin;Multicopper oxidase, type 2;Multicopper oxidase, type 3;Multicopper oxidases, conserved site	PF00394;PF07731;PF07732	Multicopper oxidase, type 1;Multicopper oxidase, type 2;Multicopper oxidase, type 3	SSF49503	Cupredoxin
Candida auris strain B11221	383	CJI97_003068	unspecified product	OG6_100257	1.10.3.2 (Laccase);1.10.3.3 (L-ascorbate oxidase);1.16.3.1 (Ferroxidase);1.7.2.1 (Nitrite reductase (NO-forming));2.7.-.- (Transferring phosphorus-containing groups.);3.2.1.- (Glycosidases, i.e. enzymes hydrolyzing O- and S-glycosyl compounds.);3.2.1.3 (Glucan 1,4-alpha-glucosidase);3.2.1.97 (Endo-a	N/A	N/A	GO:0005507;GO:0016491	copper ion binding;oxid oreductase activity	GO:005511 4	obsolete oxidation-reduction process	IPR001117;IPR002355;IPR008972;IPR011706;IPR011707;IPR033138	Multicopper oxidase, type 1;Multicopper oxidase, copper-binding site;Cupredoxin;Multicopper oxidase, type 2;Multicopper oxidase, type 3;Multicopper oxidases, conserved site	PF00394;PF07731;PF07732	Multicopper oxidase, type 1;Multicopper oxidase, type 2;Multicopper oxidase, type 3	SSF49503	Cupredoxin
Candida auris strain B11245	386	CJJ09_001108	unspecified product	OG6_100257	1.10.3.2 (Laccase);1.10.3.3 (L-ascorbate oxidase);1.16.3.1 (Ferroxidase);1.7.2.1 (Nitrite reductase (NO-forming));2.7.-.- (Transferring phosphorus-containing groups.);3.2.1.- (Glycosidases, i.e. enzymes hydrolyzing O- and S-glycosyl compounds.);3.2.1.3 (Glucan 1,4-alpha-glucosidase);3.2.1.97 (Endo-a	GO:0016021	integral component of membrane	GO:0005507;GO:0016491	copper ion binding;oxid oreductase activity	GO:000681 1;GO:00550 72;GO:0055 114	ion transport;iron ion homeostasis;obsolete oxidation-reduction process	IPR001117;IPR002355;IPR008972;IPR011706;IPR011707;IPR033138	Multicopper oxidase, type 1;Multicopper oxidase, copper-binding site;Cupredoxin;Multicopper oxidase, type 2;Multicopper oxidase, type 3;Multicopper oxidases, conserved site	PF00394;PF07731;PF07732	Multicopper oxidase, type 1;Multicopper oxidase, type 2;Multicopper oxidase, type 3	SSF49503	Cupredoxin
Candida auris strain B11245	386	CJJ09_002041	unspecified product	OG6_100380	1.10.3.2 (Laccase)	GO:0016021	integral component of membrane	N/A	N/A	N/A	N/A	IPR007248	Mpv17/PMP22	PF04117	Mpv17/PMP22	N/A	N/A
Candida auris strain B11245	386	CJJ09_002042	unspecified product	OG6_100380	1.10.3.2 (Laccase)	GO:0016021	integral component of membrane	N/A	N/A	N/A	N/A	IPR007248	Mpv17/PMP22	PF04117	Mpv17/PMP22	N/A	N/A
Candida auris strain B11245	386	CJJ09_002483	unspecified product	OG6_100257	1.10.3.2 (Laccase);1.10.3.3 (L-ascorbate oxidase);1.16.3.1 (Ferroxidase);1.7.2.1 (Nitrite reductase (NO-forming));2.7.-.- (Transferring phosphorus-containing groups.);3.2.1.- (Glycosidases, i.e. enzymes hydrolyzing O- and S-glycosyl compounds.);3.2.1.3 (Glucan 1,4-alpha-glucosidase);3.2.1.97 (Endo-a	N/A	N/A	GO:0005507;GO:0016491	copper ion binding;oxid oreductase activity	GO:000681 1;GO:00550 72;GO:0055 114	ion transport;iron ion homeostasis;obsolete oxidation-reduction process	IPR001117;IPR002355;IPR008972;IPR011706;IPR011707;IPR033138	Multicopper oxidase, type 1;Multicopper oxidase, copper-binding site;Cupredoxin;Multicopper oxidase, type 2;Multicopper oxidase, type 3;Multicopper oxidases, conserved site	PF00394;PF07731;PF07732	Multicopper oxidase, type 1;Multicopper oxidase, type 2;Multicopper oxidase, type 3	SSF49503	Cupredoxin
Candida auris strain B8441	387	B9J08_000072	hypothetical protein	OG6_100380	1.10.3.2 (Laccase)	GO:0016021	integral component of membrane	N/A	N/A	N/A	N/A	IPR007248	Mpv17/PMP22	PF04117	Mpv17/PMP22	N/A	N/A
Candida auris strain B8441	387	B9J08_000073	hypothetical protein	OG6_100380	1.10.3.2 (Laccase)	GO:0016021;GO:0005743	integral component of membrane;mitochondrial inner membrane	N/A	N/A	GO:000606 7	ethanol metabolic process	IPR007248	Mpv17/PMP22	PF04117	Mpv17/PMP22	N/A	N/A
Candida auris strain B8441	387	B9J08_000517	hypothetical protein	OG6_100257	1.10.3.2 (Laccase);1.10.3.3 (L-ascorbate oxidase);1.16.3.1 (Ferroxidase);1.7.2.1 (Nitrite reductase (NO-forming));2.7.-.- (Transferring phosphorus-containing groups.);3.2.1.- (Glycosidases, i.e. enzymes hydrolyzing O- and S-glycosyl compounds.);3.2.1.3 (Glucan 1,4-alpha-glucosidase);3.2.1.97 (Endo-a	GO:0033573	high-affinity iron permease complex	GO:0005507;GO:0004322;GO:0016491	copper ion binding;ferr oxidase activity;oxid oreductase activity	GO:000681 1;GO:00550 72;GO:0055 114;GO:003 3215	ion transport;iron ion homeostasis;obsolete oxidation-reduction process;reductive iron assimilation	IPR001117;IPR002355;IPR008972;IPR011706;IPR011707;IPR033138	Multicopper oxidase, type 1;Multicopper oxidase, copper-binding site;Cupredoxin;Multicopper oxidase, type 2;Multicopper oxidase, type 3;Multicopper oxidases, conserved site	PF00394;PF07731;PF07732	Multicopper oxidase, type 1;Multicopper oxidase, type 2;Multicopper oxidase, type 3	SSF49503	Cupredoxin
Candida auris strain B8441	387	B9J08_002997	hypothetical protein	OG6_100257	1.10.3.2 (Laccase);1.10.3.3 (L-ascorbate oxidase);1.16.3.1 (Ferroxidase);1.7.2.1 (Nitrite reductase (NO-forming));2.7.-.- (Transferring phosphorus-containing groups.);3.2.1.- (Glycosidases, i.e. enzymes hydrolyzing O- and S-glycosyl compounds.);3.2.1.3 (Glucan 1,4-alpha-glucosidase);3.2.1.97 (Endo-a	GO:0016021	integral component of membrane	GO:0005507;GO:0016491	copper ion binding;oxid oreductase activity	GO:000681 1;GO:00550 321;GO:005 5114;GO:00 70193	ion transport;iron ion homeostasis;meiotic cell cycle;obsolete oxidation-reduction process;synaptonemal complex organization	IPR001117;IPR002355;IPR008972;IPR011706;IPR011707;IPR033138	Multicopper oxidase, type 1;Multicopper oxidase, copper-binding site;Cupredoxin;Multicopper oxidase, type 2;Multicopper oxidase, type 3;Multicopper oxidases, conserved site	PF00394;PF07731;PF07732	Multicopper oxidase, type 1;Multicopper oxidase, type 2;Multicopper oxidase, type 3	SSF49503	Cupredoxin
Candida auris strain B11243	931	CJJ07_002884	unspecified product	OG6_100380	1.10.3.2 (Laccase)	GO:0016021	integral component of membrane	N/A	N/A	N/A	N/A	IPR007248	Mpv17/PMP22	PF04117	Mpv17/PMP22	N/A	N/A
Candida auris strain B11243	931	CJJ07_002885	unspecified product	OG6_100380	1.10.3.2 (Laccase)	GO:0016021	integral component of membrane	N/A	N/A	N/A	N/A	IPR007248	Mpv17/PMP22	PF04117	Mpv17/PMP22	N/A	N/A
Candida auris strain B11243	931	CJJ07_003625	unspecified product	OG6_100257	1.10.3.2 (Laccase);1.10.3.3 (L-ascorbate oxidase);1.16.3.1 (Ferroxidase);1.7.2.1 (Nitrite reductase (NO-forming));2.7.-.- (Transferring phosphorus-containing groups.);3.2.1.- (Glycosidases, i.e. enzymes hydrolyzing O- and S-glycosyl compounds.);3.2.1.3 (Glucan 1,4-alpha-glucosidase);3.2.1.97 (Endo-a	GO:0016021	integral component of membrane	GO:0005507;GO:0016491	copper ion binding;oxid oreductase activity	GO:000681 1;GO:00550 72;GO:0055 114	ion transport;iron ion homeostasis;obsolete oxidation-reduction process	IPR001117;IPR002355;IPR008972;IPR011706;IPR011707;IPR033138	Multicopper oxidase, type 1;Multicopper oxidase, copper-binding site;Cupredoxin;Multicopper oxidase, type 2;Multicopper oxidase, type 3;Multicopper oxidases, conserved site	PF00394;PF07731;PF07732	Multicopper oxidase, type 1;Multicopper oxidase, type 2;Multicopper oxidase, type 3	SSF49503	Cupredoxin
Candida auris strain B11243	931	CJJ07_003902	unspecified product	OG6_100257	1.10.3.2 (Laccase);1.10.3.3 (L-ascorbate oxidase);1.16.3.1 (Ferroxidase);1.7.2.1 (Nitrite reductase (NO-forming));2.7.-.- (Transferring phosphorus-containing groups.);3.2.1.- (Glycosidases, i.e. enzymes hydrolyzing O- and S-glycosyl compounds.);3.2.1.3 (Glucan 1,4-alpha-glucosidase);3.2.1.97 (Endo-a	N/A	N/A	GO:0005507;GO:0016491	copper ion binding;oxid oreductase activity	GO:000681 1;GO:00550 72;GO:0055 114	ion transport;iron ion homeostasis;obsolete oxidation-reduction process	IPR001117;IPR002355;IPR008972;IPR011706;IPR011707;IPR033138	Multicopper oxidase, type 1;Multicopper oxidase, copper-binding site;Cupredoxin;Multicopper oxidase, type 2;Multicopper oxidase, type 3;Multicopper oxidases, conserved site	PF00394;PF07731;PF07732	Multicopper oxidase, type 1;Multicopper oxidase, type 2;Multicopper oxidase, type 3	SSF49503	Cupredoxin

Supplementary Table 1. List of laccase homologues in C. auris from FungiDB

Organism	CDC. Strain	Gene ID	Product Description	Ortholog Group	EC numbers from OrthoMCL	Computed GO Component IDs	Computed GO Components	Computed GO Function IDs	Computed GO Functions	Computed GO Process IDs	Computed GO Processes	Interpro ID	Interpro Description	PFam ID	PFam Description	Superfamily ID	Superfamily Description
Candida auris strain B11220	381	CJI96_0000655	hypothetical_protein	OG6_105162	1.14.18.1 (Tyrosinase)	GO:0005789;GO:0016021	endoplasmic reticulum membrane;integral component of membrane	N/A	N/A	GO:0006629	lipid metabolic process	IPR005352	Erg28	PF03694	Erg28	N/A	N/A
Candida auris strain B11220	381	CJI96_0000674	hypothetical_protein	OG6_101780	N/A	GO:0016021	integral component of membrane	N/A	N/A	N/A	N/A	N/A	N/A	PF01544	N/A	SSF143865; SSF144083	CorA soluble domain-like; Magnesium transport protein CorA, transmembrane region
Candida auris strain B11220	381	CJI96_0003008	hypothetical_protein	OG6_101780	N/A	GO:0016021	integral component of membrane	N/A	N/A	N/A	N/A	N/A	N/A	PF01544	N/A	SSF143865; SSF144083	CorA soluble domain-like; Magnesium transport protein CorA, transmembrane region
Candida auris strain B11221	383	CJI97_001444	unspecified product	OG6_101780	N/A	GO:0016020	membrane	GO:0046873	metal ion transmembrane transporter activity	GO:0030001;GO:0055085	metal ion transport;transmembrane transport	IPR002523	Mg2+ transporter protein, CorA-like/Zinc transport protein ZntB	PF01544	Mg2+ transporter protein, CorA-like/Zinc transport protein ZntB	SSF143865; SSF144083	CorA soluble domain-like; Magnesium transport protein CorA, transmembrane region
Candida auris strain B11221	383	CJI97_002873	unspecified product	OG6_105162	1.14.18.1 (Tyrosinase)	GO:0005789;GO:0016021	endoplasmic reticulum membrane;integral component of membrane	N/A	N/A	GO:0006629	lipid metabolic process	IPR005352	Erg28	PF03694	Erg28	N/A	N/A
Candida auris strain B11221	383	CJI97_002891	unspecified product	OG6_101780	N/A	GO:0016021;GO:0016020	integral component of membrane;membrane	GO:0046873	metal ion transmembrane transporter activity	GO:0030001;GO:0055085	metal ion transport;transmembrane transport	IPR002523	Mg2+ transporter protein, CorA-like/Zinc transport protein ZntB	PF01544	Mg2+ transporter protein, CorA-like/Zinc transport protein ZntB	SSF143865; SSF144083	CorA soluble domain-like; Magnesium transport protein CorA, transmembrane region
Candida auris strain B11245	386	CJJ09_001261	unspecified product	OG6_101780	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	PF01544	N/A	SSF143865; SSF144083	CorA soluble domain-like; Magnesium transport protein CorA, transmembrane region
Candida auris strain B11245	386	CJJ09_001280	unspecified product	OG6_105162	1.14.18.1 (Tyrosinase)	GO:0005789;GO:0016021	endoplasmic reticulum membrane;integral component of membrane	N/A	N/A	GO:0006629	lipid metabolic process	IPR005352	Erg28	PF03694	Erg28	N/A	N/A
Candida auris strain B11245	386	CJJ09_003418	unspecified product	OG6_101780	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	PF01544	N/A	SSF143865; SSF144083	CorA soluble domain-like; Magnesium transport protein CorA, transmembrane region
Candida auris strain B8441	387	B9J08_001162	hypothetical protein	OG6_101780	N/A	GO:0005886	plasma membrane	GO:0022890;GO:0046873	inorganic cation transmembrane transporter activity;metal ion transmembrane transporter activity	GO:0006812;GO:0015693;GO:0055085	cation transport;magnesium ion transport;transmembrane transport	N/A	N/A	PF01544	N/A	SSF143865; SSF144083	CorA soluble domain-like; Magnesium transport protein CorA, transmembrane region
Candida auris strain B8441	387	B9J08_002817	hypothetical protein	OG6_105162	1.14.18.1 (Tyrosinase)	GO:0005789;GO:0016021	endoplasmic reticulum membrane;integral component of membrane	GO:0030674	protein-macromolecule adaptor activity	GO:0070988;GO:0006696;GO:0006629	demethylation;ergosterol biosynthetic process;lipid metabolic process	IPR005352	Erg28	PF03694	Erg28	N/A	N/A
Candida auris strain B8441	387	B9J08_002836	hypothetical protein	OG6_101780	N/A	GO:0000329;GO:0016021;GO:1990816	fungal-type vacuole membrane;integral component of membrane;vacuole-mitochondrion membrane contact site	GO:0015095	magnesium ion transmembrane transporter activity	GO:0010961;GO:0030001;GO:0055085	cellular magnesium ion homeostasis;metal ion transport;transmembrane transport	N/A	N/A	PF01544	N/A	SSF143865; SSF144083	CorA soluble domain-like; Magnesium transport protein CorA, transmembrane region
Candida auris strain B11243	931	CJI07_003398	unspecified product	OG6_105162	1.14.18.1 (Tyrosinase)	GO:0005789;GO:0016021	endoplasmic reticulum membrane;integral component of membrane	N/A	N/A	GO:0006629	lipid metabolic process	IPR005352	Erg28	PF03694	Erg28	N/A	N/A
Candida auris strain B11243	931	CJI07_004279	unspecified product	OG6_101780	N/A	GO:0016020	membrane	GO:0046873	metal ion transmembrane transporter activity	GO:0030001;GO:0055085	metal ion transport;transmembrane transport	IPR002523	Mg2+ transporter protein, CorA-like/Zinc transport protein ZntB	PF01544	Mg2+ transporter protein, CorA-like/Zinc transport protein ZntB	SSF143865; SSF144083	CorA soluble domain-like; Magnesium transport protein CorA, transmembrane region
Candida auris strain B11243	931	CJI07_004614	unspecified product	OG6_101780	N/A	GO:0016020	membrane	GO:0046873	metal ion transmembrane transporter activity	GO:0030001;GO:0055085	metal ion transport;transmembrane transport	IPR002523	Mg2+ transporter protein, CorA-like/Zinc transport protein ZntB	PF01544	Mg2+ transporter protein, CorA-like/Zinc transport protein ZntB	SSF143865; SSF144083	CorA soluble domain-like; Magnesium transport protein CorA, transmembrane region

Supplementary Table 2. List of tyrosinase homologues in C. auris from FungiDB