TITLE:

Multifaceted effects of bycatch mitigation measures on target/non-target species for pelagic longline

fisheries and consideration for bycatch management

AUTHORS:

Daisuke Ochi[#], Kei Okamoto and Shintaro Ueno

AFFILIATION:

Fisheries Resources Institute, Japan Fisheries Research and Education Agency

Number of tables: 10

Number of figures: 9

[#]Corresponding author,

Address: Fisheries Resources Institute, Japan Fisheries Research and Education Agency, 2-12-4,

Fukuura, Kanazawa-ku, Yokohama-city, Kanagawa 2368648, Japan

e-mail: otthii80s@gmail.com

____

SUMMARY

The pelagic longline fishery, in an effort to reduce bycatch of sea turtles, have developed and deployed fisheries bycatch mitigation techniques such as replacing J/tuna hooks and squid bait with circle hooks and whole fish bait. However, little emphasis has been placed on the side effects of bycatch mitigation measures on endangered species other than target bycatch species. Several previous studies of the side effects have been marred by lack of control for the covariates. Here, based on long-term data obtained from research cruises by a pelagic longline vessel, we examined the effects of using circle hooks and whole fish bait to replace squid bait on the fishing mortality of target and non-target fishes, and also bycatch species. A quantitative evaluation analysis of our results, based on a Bayesian approach, showed the use of circle hooks to increase mouth hooking in target and bycatch species, and their size to be proportional to the magnitude of the effect. Although deploying circle hooks increased fishing mortality per unit effort (MPUE) for shortfin mako shark, the magnitude of changing the bait species from squid to fish clearly had a far greater impact on MPUE than the use of circle hooks. Because the impact of the introduction of bycatch mitigation measures on species other than the focused bycatch species is non-negligible, a quantitative assessment of bycatch mitigation-related fishing mortality is critical before introducing such measures.

Keywords: billfishes, circle hook, finfish bait, fisheries management, sea turtle, sharks, tuna, longline fishery

INTRODUCTION

Unintentional catch by fisheries is referred to as bycatch, and bycatch of especially

endangered species, can have devastating effects on such populations. Thus, efforts to

minimize bycatch of endangered species are strongly encouraged at all layers from local to

international in terms of species conservation. For example, in the tuna longline fishery,

concerns about the increased conservation risk for seabirds and sea turtles by unintentional

and fatal catch have been a major issue among many regional fisheries management

organizations (RFMOs) since the 1990s (Wallace et al. 2013; Dias et al. 2019). In recent years,

some elasmobranchs, whose populations are declining further, are also treated as bycatch

species. The decline of these species has focused attention at both the national and the

international levels. Even for target fish that are not bycatch species, addressing the

deterioration of stock status caused by overfishing requires reductions in unintentional fishing

mortality (for instance, billfishes in the North Atlantic; Kerstetter & Graves 2006; Diaz 2008).

Several studies have weighed the sustainability of tuna longline fisheries against the

conservation of species vulnerable to bycatch (Hall et al. 2000; Melvin et al. 2014; Clarke et al.

2015)—particularly for seabirds and marine turtles—with the development of several effective

bycatch mitigation measures (Melvin et al. 2014; Swimmer et al. 2017). Bycatch mitigation

measures in longline fisheries target specific animal groups and are evaluated based on their

success in reducing mortality due to bycatch of specific vulnerable species. While the impact

on the catch of the target fish is the primary consideration when evaluating bycatch mitigation

techniques, few studies have examined the impacts and tradeoffs on species not targeted by

mitigation techniques (Pacheco et al. 2011; Gilman et al. 2016). However, the introduction of

bycatch mitigation measures requires an assessment of the impact not only on the target

bycatch species on the other ecologically related species prior to their introduction (Reinhardt

et al. 2018).

Use of circle hook and whole finfish bait are typical sea turtle bycatch mitigation

measures for pelagic longline fisheries (Watson et al. 2005; Gilman et al. 2006; Yokota et al.

2009; Stokes et al. 2011). The tip of the circle hook bends inward, and when a fish or sea turtle

swallows the hooked bait, the circle hook cannot hook inside the digestive tract; instead, as the

hook exits the mouth, a torque causes it to hook through the edge of the mouth. This property

allows for easy hook removal and has been reported to reduce the mortality rate of bycatch sea

turtles on board and after release (Kiyota et al. 2004; Cooke & Suski 2004; Kerstetter &

Graves 2006). Reports of positive effects of circle hooks include those for other species—such

as reduced haulback and post-release mortality in sharks, reduced post-release mortality in

swordfish, and increased catch rates in tuna, the target fish. The use of fish bait instead of

squid bait potentially reduces bycatch (Watson et al. 2005; Yokota et al. 2009, 2011) and

mortality rates (Stokes et al. 2011; Parga et al. 2015) of sea turtles. These sea turtle mitigation

measures, however, exact other costs. Several meta-analytic studies have reported the circle

hook-related reduction of sea turtle mortality rate but increase in billfish and shark catch rates

(Gilman et al. 2016; Reinhardt et al. 2018; Santos et al. 2020). The use of whole fish bait also

reportedly increases the catch of sharks and other species (Foster et al. 2012). However, few

studies have allowed for quantitative evaluation of the effects of these mitigation measures

experimentally on species beyond sea turtles. This limited data concern is due in part to the

reliance on observer data from commercial vessels, and small sample sizes, small comparison

groups, and lack of experimental rigor from research vessels. In addition, the impact

assessment for sharks underestimates catch rates and mortality associated with missed catches

due to "bite-off" branchline (Reinhardt et al. 2018) and several other issues. In addition, many

experimental studies and meta-analyses (e.g., Diaz 2008; Godin et al. 2012; Huang et al. 2016;

Pacheco et al. 2011; Yokota et al. 2006a) only evaluate gear impacts at significance levels

without evaluating the magnitude of the effect. Small significant differences may be judged

not to matter much when assessing overall risk in bycatch species. Studies that controlled for

these conditions would allow for an evaluation of adverse effects without the confounding

problems described above. Also, many studies use catch/bycatch rates (Andraka et al. 2013;

Foster et al. 2012; Gilman et al. 2007; Watson et al. 2005; Yokota et al. 2006a) and mortality

rates (at haulback or after release; Carruthers et al. 2009; Gallagher et al. 2014; Horodysky et

al. 2005; Kerstetter et al. 2003) as important impact indicators, without considering

irreversible impacts, such as the number of organisms killed at the time of catch. Additionally,

hooking location itself (mouth, swallow, or external) is believed to have a strong influence on

mortality rate, which demands the estimation of risk under specific hooking conditions, taking

causal relationships into account.

Here, we used data from controlled experiments to analyze these confounding factors

and developed a Bayesian statistical model to evaluate the effects of changing hook and bait

species on fish species other than turtles—particularly, tuna, swordfish, and sharks—and then

verified the contribution of circle hooks and fish bait to mortality rate, catch rate, and fatal

catch rate, respectively. We also discussed the appropriate assessment of bycatch mitigation

measures in fisheries management.

METHOD

Experimental Operations

We analyzed data from the R/V Taikei No. 2 longline research operation conducted in the

Northwest Pacific Ocean between 2002 and 2010—a typical Japanese shallow-setting

operation targeting mainly swordfish and sharks (set depth: shallowest 47.44m±14.22SD,

deepest 72.41m±10.87SD (n=98 ops.); based on the time depth recorder data [SBT500,

Murayama Denki Ltd.]), using four hooks per basket, a wire leader, and a night soaking

(Yokota et al. 2006a). A total of 286,363 hooks from 306 operations (range: 400 – 964 hooks

per operation) were deployed in the experiment (Table 1). The area of operation ranged from around the Izu Islands in Japan to off the east coast of northeastern Honshu—typical fishing ground for Japanese shallow-setting longliners (Fig. 1; Hiraoka et al. 2016). We deployed 11 different hooks (Table 2) and we describe hook shapes and other details of these hooks in Appendix 1 following the measurement method of Yokota et al. (2006b). From these hooks, we regarded the 4.0-sun tuna hooks as a control group. Since the remaining ten different circle hooks varied greatly in size (Table 2) and sample sizes were too small to analyze each hook type separately, we divided them into two groups ("small-C" and "large-C" hook) according to size for convenience (threshold: straight total length = 68 mm AND maximum total width = 80 mm; approximately equivalent to 5.0-sun or 18/0). Because degree of hook offset has been reported to affect catch and hooking location (Cooke & Suski 2004), most hooks were <10°but some were nearly 15°. The bait comprised chub mackerel (*Scomber japonicus*) and Japanese common squid (*Todarodes pacificus*) in the range of 20–30 cm in fork length or dorsal mantle length. These were frozen and stored, then completely thawed before being hooked. The sequence of line setting was divided into several experimental segments, and a different combination of hook and bait type was applied for each segment, with an alternate order of segments at each operation.

The researcher recorded catch for each operation for all catch/bycatch, and the species caught, fate of catch (alive or dead), hooking location (mouth, swallow, and external hooking), time of catch, and float ID. The researchers determined if the catch was alive or dead based on the movement of the animals and the degree of injuries before being hauled. Float ID was recorded when a float was dropped during line setting and when it was retrieved to the deck during line hauling in order to calculate soak time. Hooking locations were recorded for catches caught using squid bait. At the start of longline operations, the researcher also collected sea surface temperature with a water thermometer (DS-1; Murayama Denki Ltd.)

equipped on the vessel, which we subsequently used in the analysis.

Since the experiment was conducted using multiple sizes (degrees) of circle hooks in

combination with whole fish bait and squid bait, we categorized hook shapes into three main

types:

1.   control (tuna hook 4.0-sun; tuna)

2.   smaller circle hook (smaller than the threshold; small-C)

3.   larger circle hook (the threshold or larger; large-C)

In Table 1, we show longline effort separated by bait and hook type, tabulated

according to the above categorization. We selected the following species for analysis: blue

shark *Prionace glauca*, shortfin mako shark *Isurus oxyrinchus*, bigeye tuna *Thunnus obesus*,

swordfish *Xiphias gladius*, striped marlin *Kajikia audax*, common dolphinfish *Coryphaena*

hippurus, escolar *Lepidocybium flavobrunneum* and, longnosed lancetfish *Alepisaurus ferox*,

and loggerhead turtle *Caretta caretta*.

Statistical Analysis

We conducted all analyses using a Bayesian approach to estimate parameters. We adopted

haulback mortality rate, catch per unit effort (CPUE; /1000 hooks), and mortality per unit

effort (MPUE; /1000 hooks) (Afonso et al. 2011) as indices to evaluate the impact of hook and

bait type on fishing mortality within the analyses. We used the following data as inputs to the

model: number of caught (individuals), longline effort (number of hooks), fate at hauling,

hooking location, year and location of operation, water temperature at operation, and soaking

time.

Because of the missing data due to not recording the hooking location when fish bait

was used, we split the analysis to evaluate the impact of hook type and bait type on fishing

mortality into two models. Model 1 evaluated only the effect of hook type based on capture

events with squid bait and assumed that the use of circle hooks would change to more mouth

hooking of each species, resulting in improved mortality rate. Model 2 evaluated both hook

and bait types, and assumed that the combination of the two would result in large fluctuations

in mortality rate, CPUE and MPUE for each species.

We based MODEL 1 on a logit regression using a Bernoulli distribution. We express

the observed hooking location H and haulback mortality rate M in MODEL 1 by the following

equations:

$$M \sim Bernoulli(p_{dead}) \tag{1}$$

$$logit(p_{1,dead}) = \beta_{1,hkloc} + \beta_{1,sst} + \beta_{1,soaktime} \tag{2}$$

$$H \sim Categorical(softmax(\theta_{hkloc \cdot hook})) \tag{3}$$

where β_1 is the parameter in each explanatory variable (*hkloc:* hooking location, *sst:* sea

surface temperature and *soaktime:* soak time), $p_{1,dead}$ is the expected haulback mortality rate,

and $\theta_{hkloc \cdot hook}$ is the expected probability of hooking location in each hook type.

We structured MODEL 2 to calculate the expected number of mortalities per effort

(MPUE) based on the parameters estimated in both the mortality and CPUE estimation subsets.

In MODEL 2, due to lack of hooking location data, we calculate $p_{2,dead}$ from a modified

equation (2) as follows;

$$logit(p_{2,dead}) = \beta_{2,hook} + \beta_{2,bait} + \beta_{2,sst} + \beta_{2,soaktime} \tag{4}$$

181   where β_2 is the parameter in each explanatory variable (*hook:* hook type, *bait:* bait type).

182   The CPUE subset is based on a log regression using a Poisson distribution as the error

183   structure. We express the number of catches C per operation using the expected CPUE λ as

184   follows:

$$C \sim Poisson(\lambda + \log(E)) \tag{5}$$

$$\lambda = \log\left(\gamma_{hook} + \gamma_{bait} + \gamma_{lat} + \gamma_{sst} + r_{year}\right) \tag{6}$$

$$r_{year} \sim N(0, \ \sigma^2) \tag{7}$$

where E is the longline effort γ is a parameter in each explanatory variable (*lat:* latitude where

the longline set), r_{year} is a random effect of annual fluctuation on CPUE, and σ is a standard

deviation.

We obtained the expected values of hook-bait-specific MPUE ζ by multiplying CPUE

by at-haulback mortality rate as follows:

$$\zeta = \hat{p}_{2,dead} \times \hat{\lambda}$$

$$\tag{8}$$

where $\hat{p}_{2,dead}$ denotes the expected mortality rate standardized for hook and bait type, and $\hat{\lambda}$

denotes the estimated CPUE standardized for hook and bait type. We used the standardization

method for abundance indices used in fisheries stock assessment (Maunder & Punt 2004). In

this method, explanatory variables other than the factor subject to standardization are averaged

to predict the objective variable, which in stock assessment is a time scale such as years or

months, but in our case, we modified the standardization scale to reference hook type and bait

type.

We calculated each parameter based on a Bayesian approach with Markov chain

Monte Carlo (MCMC) sampling. For the MCMC sampling, we used cmdstan 2.28.2 (Stan

Development Team 2021). As a prior distribution for σ, we used a half student-t distribution

with 2 degrees of freedom, mean 0, and variance 2.5, and we used a uniform distribution for

the other parameters (β_1, β_2, γ, θ). We computed the posterior distribution using Stan with

15000 sampling iterations including 10000 warmup iterations, number of chains as 4, and no

sinning. We calculated Bayesian credible intervals based on the highest density interval (HDI)

for the estimates. Although the Bayesian approach for the estimates precluded significance

testing, we determined the difference between the estimates of the experimental group and

those of the control group ("swallowing" for hooking location, "tuna" hook for hook type, and

"squid" for bait type), and if the lower and upper limits of HDI for the difference did not

exceed 0, we considered the difference as a difference for convenience (assuming region of

practical equivalence [ROPE] as 0; Kruschke 2015). In Appendix 2 we show the Stan code

used to estimate each parameter in MODEL 1 and 2. For other data handling, statistical

analysis, and plotting, we used R4.3.0 (R Core Team 2023) and packages "cmdstanr 0.6.1,"

"ggalluvial 0.12.5," "ggthemes 4.2.4," "mapdata 2.3.1," "maps 3.4.1," "sf 1.0–14," "tidybayes

3.0.6," and "tidyverse 2.0.0" (Pebesma 2018; Wickham et al. 2019; Becker et al. 2020;

Brunson 2020; Arnold 2021; Gabry & Češnovar 2021; Becker et al. 2022; Kay 2023).

RESULTS

Summary Statistics

Sufficient catches/bycatches of blue shark, longnosed lancetfish, common dolphinfish, and

shortfin mako shark swordfish, bigeye tuna, loggerhead turtle, escolar and striped marlin were

recorded for the later analysis (Table 3). The main species listed in Table 3 as "other species"

are listed as follows; salmon shark *Lamna ditropis* (N = 229), pelagic stingray

Pteroplatytrygon violacea (N = 199), pomflets *Brama spp.* (N = 135), bigeye thresher shark

Alopias superciliosus (N = 89), and albacore *Thunnus alalunga* (N = 69). The sample sizes of

these "other species" were too skewed among experimental groups to converge the later

analysis.

In Table 3 we also show the number of fish caught by hook type and bait type. Bigeye

tuna had extremely low catches on large-C hooks, and loggerheads had low catches on fish

bait. The most common hooking location at the time of catch was mouth hooking for all nine

species, with extremely few hook locations other than mouth hooking, especially for bigeye

tuna and escolar (Table 4). The proportion of mortality of captured species at haulback varied

greatly by species. The haulback mortality rate was low for blue sharks, common dolphinfish,

escolar, and shortfin mako sharks and, although higher for bigeye tuna, longnose lancetfish,

striped marlin, and swordfish, was generally extremely low, especially in loggerhead turtles.

The length-based size composition of the nine species (precaudal length for blue

sharks and shortfin mako sharks, straight-line carapace length for loggerhead turtles,

eye-to-fork length for striped marlin and swordfish, and fork length for all other species) used

in the analysis was not statistically compared in this study and was not included in the model

because it did not contribute to haulback mortality rate, but in Appendix 3 we include a

histogram.

Almost all parameters in the models for the nine species were successfully converged

(Rhat < 1.1) but some of the parameters for loggerhead turtle were not converged, as we

describe below.

Output of MODEL 1

In Table 5 we show occurrence estimates of hooking locations by hook type throughout the

model. We observed differences in hooking location by hook type for blue shark, common

dolphinfish, loggerhead turtle, and swordfish, with an increase in mouth hooking and a

decrease in hook swallowing for large-C for loggerhead turtle and a clear increase in mouth

hooking for small-C for shortfin mako shark and swordfish (Fig. 2). In blue sharks, the

frequency of hook swallowing decreased in both small-C and large-C.

In Table 6 we show haulback mortality rates by hooking location. We observed clear

differences in haulback mortality rate by hooking location for blue shark, shortfin mako shark,

striped marlin, and swordfish (Fig. 3). Haulback mortality rates after hook swallowing for blue

sharks were lower than those for external hooking, and higher than those for mouth hooking.

We observed lower haulback mortality rates for shortfin mako shark, striped marlin, and

swordfish from hook swallowing than from mouth hooking.

Output of MODEL 2

Haulback mortality rate was higher for large-C than for tuna hook types in bigeye tuna, but did

not differ among hook types in other species (Table 7, Fig. 4). On the other hand, the mortality

rate for two bait types differed in bigeye tuna and blue shark, with a rate reduction and

increase with the use of fish bait in bigeye tuna and shortfin mako shark, respectively.

In Fig. 5 and Fig. 6 we show the effects of two covariates—sea surface temperature

(SST) and soak time—on haulback mortality rate. The response to SST differed by species,

with haulback mortality rate increasing with higher SST for shortfin mako sharks, swordfish,

common dolphinfish, and escolar, and conversely increasing with lower SST for bigeye tuna

and longnose lancetfish. We observed little fluctuation in haulback mortality rate with SST in

blue shark and striped marlin. In general, haulback mortality rate increased with increasing

soak time. However, for bigeye tuna, haulback mortality rate decreased with soak time, but the

trend was not clear, and for blue sharks, haulback mortality rate increased only slightly with

increased soak time.

In Table 8 we show standardized CPUE by hook and bait type. Differences in CPUE

among hook type were found for all species except striped marlin (Fig. 7). We observed higher

CPUE for small-C in bigeye tuna and escolar and higher CPUE for both small and large -C in

blue shark, common dolphinfish, longnose lancetfish, shortfin mako shark, and swordfish.

Especially in blue sharks, the difference in CPUE could be over 30 catch/1000 hooks

depending on the hook type.

We observed differences in standardized CPUE by bait type in bigeye tuna, blue

shark, shortfin mako shark, common dolphinfish, and escolar. CPUE decreased with whole

281 fish bait in bigeye tuna and blue sharks, but increased in shortfin mako shark, common

282 dolphinfish, and escolar. In the case of blue shark, escolar, shortfin mako shark, and swordfish,

283 the bait effect could be varied with circle hooks, suppressing the CPUE-increasing effect by

284 fish bait in bigeye tuna, blue shark, and swordfish with the use of circle hooks, while

285 conversely this combination boosted increase of CPUE in common dolphinfish, escolar, and

286 shortfin mako shark.

287   In Table 9 we show MPUE values by hook and bait type. Compared to differences in

288 MPUE among hook and bait type, those in CPUE were relatively small. We observed

289 differences in MPUE by hook types, with higher MPUE for small-C in bigeye tuna, escolar

290 and shortfin mako shark, and higher MPUE for both small and large-C in blue shark, common

291 dolphinfish, longnose lancetfish and swordfish (Fig. 8). We confirmed decreases in MPUE by

292 fish bait in bigeye tuna and blue shark, and blue sharks, and conversely, increases in MPUE in

293 common dolphinfish, shortfin mako shark, and swordfish. In escolar, however, MPUE

294 increased only when small-C was combined with whole fish bait. The effect of the

295 combination of wholefish bait and circle hook varied in bigeye tuna, blue shark, common

296 dolphinfish, shortfin mako shark, and swordfish. In bigeye tuna, blue shark, and swordfish, the

297 effect of the circle hook on MPUE was suppressed by fish bait, while in common dolphinfish

298 and shortfin mako shark, MPUE increased significantly when whole fish bait and circle hook

299 were used together.

301 ***Hook and bait effect for sea turtle bycatch***

302 We failed to complete our analysis for loggerhead turtles throughout the models because the

303 bias in frequency of capture events among the experimental groups was too large and did not

304 converge except for only a part of MODEL 1. Instead, we show the nominal CPUE, haulback

305 mortality rate, and MPUE for each experimental group in Table 10, and hooking location and

haulback mortality rate by each hook type with squid bait in Fig. 9. Most individuals survived

regardless of hooking location. When whole fish bait was used, haulback mortality rate and

associated MPUE were zero. For squid bait, large-C had the smallest CPUE, mortality, and

MPUE.

DISCUSSION

Our experimental comparisons showed that the hook shape and bait type—both considered as

effective bycatch mitigation measures for sea turtles—have extremely multifaceted effects for

teleost fishes and sharks and, in some species, the direction of the effects was conflicted. The

results provide significant insight into two aspects of the management of vulnerable bycatch

species in tuna fisheries: how the confrontational effect of bycatch mitigation measures should

be managed, and in which processes of fishing mortality intervention in the management of

vulnerable species should occur. As a specific concern regarding the former, when considering

shortfin mako shark, which are experiencing significant stock depletion in the North Atlantic

(Sims et al. 2018; ICCAT 2019), the implementation of bycatch mitigation measures for sea

turtles, which are also required to reduce fishery bycatch, could conversely increase fishing

mortality and become a conservation risk. Previously, most of bycatch mitigation measure

assessments have focused on whether they reduce the impact of vulnerable bycatch species of

concern being bycaught, with the secondary impact being from an economic perspective—in

other words, whether the catch rate of commercial species is reduced or not. Here we show

that widely adopted bycatch mitigation measures may have a negative impact on some

endangered species. Where both loggerhead turtle and shortfin mako with opposing effects are

at low abundance, management measures should be based on a thorough discussion

identifying the optimal combination of mitigation measures, accompanied by scientific

evidence. With regard to the latter issue, the mortality reduction expected from circle hooks is

not very promising, especially for species with high catch mortality, since the main effect of

circle hooks is to minimize internal organ damage, which is of little use for species that have

died from other causes, such as heat stress or suffocation. For such species, consideration of

methods to reduce the catch itself rather than the mortality rate will be of greater benefit.

Increased mouth hooking of loggerhead turtle by large circle hooks is consistent

with existing studies. The circle hook prevents internal organ damage and improves the

probability of live release (Cooke & Suski 2004) while the impact of circle hooks on haulback

mortality in this study could not be evaluated due to skewed data about mortality events. For

the same reason, the MODEL 2 analysis could not evaluate the effects of circle hook and bait

type on CPUE, mortality rate, and MPUE of loggerhead turtle. However, since mortality event

of loggerhead turtle did not occur at all when fish bait was used, it may be assumed that there

is an effect of mortality reduction by using fish bait. This result is also consistent with existing

studies, and is related to the lower attractant effect of whole fish bait on marine turtles and the

increased probability of swallowing caused by the difficulty to bite off the bait (Stokes et al.

2011; Parga et al. 2015).

Our study indicated that reduced hook swallowing with circle hooks and increased

haulback mortality after hook swallowing were found for sharks and swordfish. Hook

swallowing has been reported to increase the likelihood of fatal damage to internal organs.

Although previous studies have reported that studies to attach satellite tags to white marlin

Kajikia albida caught by recreational fishing using circle hooks and subsequently released

have reduced the post-release mortality rate (Horodysky & Graves 2005), unfortunately, the

present study did not corroborate this information. Our results also presented different effects

of hooking location by hook types, with more frequent mouth hooking by small circle hooks in

many species. Few studies have examined the relationship among size of circle hook, hooking

location and haulback mortality of non-turtle species. However, two studies discussed the possibility that relative differences in mouth and hook size, and differences in feeding behavior toward prey (swallowing the prey whole or biting it off) may affect the hooking location (Epperly et al. 2012; Gilman et al. 2020).

In some species, an increase in MPUE was observed only with the use of small-C hooks among the two types of circle hooks. Except for the case of shortfin mako shark, this indicates that the increase in CPUE due to small-C hooks had a greater impact on MPUE than the haulback mortality rate. There have been many previous findings on the effects of circle hook use on CPUE, with elevated CPUE for tunas and no consistent trend for other teleosts and sharks. However, although few previous studies have focused on hook size and made comparisons, catch rates for skipjack, shortbill spearfish, escolar, and lancetfish are reported to have decreased when larger hooks were used (Curran & Beverly 2012). Considering the effect of hook size in terms of the catch process, it is unlikely that catch rates increased due to swallowing, as the results of MODEL 1 indicate an increase in mouth hooking for many species. In the case of shortfin mako shark, the increase was observed in CPUE for both small and large-C hooks, but this increase was not observed in MPUE for large-C hooks. Since the increase in CPUE with large-C was small, the effect on MPUE may have been masked by the uncertain haulback mortality rate. The effects of the bait type varied by species, unlike the pattern for hook types, with the use of whole fish bait decreasing CPUE and MPUE for blue sharks, while we observed the exact opposite effect for shortfin mako sharks, in that fish bait increased CPUE and MPUE—and even haulback mortality rate. Whole fish bait also reduced MPUE in bigeye tuna, while in common dolphinfish and escolar, it tended to increase MPUE and CPUE. We observed few effects of bait type in billfishes. Several studies have examined the effects of fish bait without circle hook and have reported reduced catch rates for tropical tunas, blue sharks, and escolar and increased catch rates for shortfin mako, porbeagle shark,

and white marlin (Fernandez-Carvalho et al. 2015; Foster et al. 2012; Watson et al. 2005;

Yokota et al. 2009). In swordfish, some previous studies evaluating the effect of switching to

whole fish bait from squid bait have reported conflicting effects (increase: Santos et al. 2012;

Foster et al. 2012; decrease: Fernandez-Carvalho et al. 2015). Very few studies have assessed

haulback mortality by bait types other than those on sea turtles. The catch rates of the target

species, as previously noted, were affected by bait texture, but a mechanistic explanation for

mortality effects is lacking. Since the likelihood that differences in feeding behavior among

species have an effect is high, this issue could be resolved through comparative studies based

on observations of feeding behavior, as in the case of circle hooks.

When the effects of hook and bait types were considered simultaneously, it was found

that the bait type had a more significant impact on CPUE, mortality, and MPUE than the hook

type. However, whether this impact was beneficial or detrimental varied greatly depending on

the species. Although the combination of circle hooks and fish bait is effective for avoiding

sea turtle bycatch, this combination may pose a high mortality risk for endangered species like

shortfin mako shark, and even in the case of target fishes like bigeye tuna, may counteract the

expected positive effect of the circle hook on catch rate. In the case of shortfin mako shark, for

example, changing from tuna hooks to small-C hooks increases MPUE by about 1.8 times, and

changing the bait from squid to fish increases MPUE by over 7 times (Table 9), and in the case

of bigeye tuna, the CPUE estimate, which increased by 2.3 times with small-C hooks, returned

to the same level as tuna hooks by changing from squid to fish bait (Table 8). Such substantial

changes in CPUE and MPUE would not be ignored when managing fisheries for those species.

Although very limited studies have simultaneously examined the interrelationship between

hook and bait types, all studies support the conclusion that the combination of hook type and

bait type causes fluctuations in catch rates and that the direction of response varies among

species (Coelho et al. 2012; Foster et al. 2012; Fernandez-Carvalho et al. 2015).

Water temperature and soak time emerged as significant factors affecting haulback

mortality rate, which had been reported in sharks (Carruthers et al. 2009; Gallasher et al. 2014)

and sea turtles (Watson et al. 2005). This indicates these covariates need to be controlled

statistically or experimentally when assessing the effects of hook and bait type on mortality

rate. The effect of water temperature—particularly during the depth and time of day when

hooked—and changes in water temperature up to the time the fish is landed, are considered to

be influential. In addition, in high water temperature environments, studies have identified an

increased risk of suffocation due to decreased dissolved oxygen in water and increased

physiological metabolic rate (Skomal & Bernal 2010). Gallagher et al. (2014) reported an

increase in haulback mortality rate for four shark species when caught during high water

temperatures. In addition, for the species that adopt rum ventilation, prolonged soak time

inevitably increases the risk of suffocation due to the restriction of swimming behavior by

being hooked. Mortality rates of tuna, swordfish, and sharks reportedly increased with

increasing soak time (Epperly et al. 2012; Gallagher et al. 2014).

Here, we quantified our data through experimental operations that standardized the

various conditions, but not all aspects were completely controlled. For example, while

previous studies on hook size have examined the correspondence with actual measurements

(Gilman et al. 2016), several shapes of circle hook were used in the experiment in this study,

precluding examination of effects of individual hook types due to sample size issues. We were

also unable to examine hooking location of the catch when fish bait was used. These

omissions, while having a limited impact on the present conclusions, are probably variables

that should be considered for a deeper examination of the effects of terminal gear on catch and

bycatch. In this experiment, wire leaders were used on all branchlines to minimize the effects

of sharks' bite-off. While some studies have described concerns that wire leaders may increase

catch rates, especially for rare sharks, they are considered essential for at least experimentally

verifying accurate catch and mortality rates for shark species. We know from this and previous studies that haulback mortality rates for sharks are much lower than those for teleosts (Reinhardt et al. 2018), and the implementation of appropriate releases, even with wire leaders, allow for the reduction of risk for vulnerable shark species.

Here, based on a Bayesian approach, we succeeded in presenting a quantitative impact assessment of terminal gear on teleosts, sharks, and sea turtles by directly calculating the expected values for mortality rate, CPUE, and MPUE with each terminal gear. Calculating MPUE using this model can be a very useful tool because it provides a more direct estimate than does CPUE or mortality rate alone of catch/bycatch risk to populations of those species. Although we did not include post-release mortality rate in the model due to lack of data, it would be possible to estimate overall fishing mortality in the model by designing additional experiments so that mark–recapture is conducted at the same time. Even if it is not possible to use wire leaders for the proportion of "cryptic catch" due to bite-off, it is possible to extrapolate this proportion into the model to make predictions regarding mortality—a development we anticipate.

In this study, we successfully conducted a quantitative impact assessment of bycatch integrating catch rate and mortality using a Bayesian approach that directly predicts mortality and CPUE. Although the data used in the analysis relied solely on the results of an Asian-style longline experiment in the Pacific Ocean and may therefore contain inherent biases, the same analysis method can be used in conjunction with data from other experiments conducted in other areas and fishing styles to provide a more integrated assessment.

REFERENCES

Afonso AS, Hazin FHV, Carvalho F, Pacheco JC, Hazin H, Kerstetter DW, Murie D, Burgess

GH (2011) Fishing gear modifications to reduce elasmobranch mortality in pelagic and

bottom longline fisheries off Northeast Brazil. Fisheries Research **108**:336–343.

Andraka S, Mug M, Hall M, Pons M, Pacheco L, Parrales M, Rendón L, Parga ML, Mituhasi

T, Segura Á, et al. (2013) Circle hooks: Developing better fishing practices in the

artisanal longline fisheries of the Eastern Pacific Ocean. *Biological Conservation* **160**,

214–224. https://doi.org/10.1016/j.biocon.2013.01.019

Arnold JB (2021) ggthemes: Extra Themes, Scales and Geoms for "ggplot2." Available at

https://cran.r-project.org/package=ggthemes

Becker RA, Wilks AR, Brownrigg R (2020) mapdata: Extra Map Databases. Available at

https://cran.r-project.org/package=mapdata

Becker RA, Wilks AR, Brownrigg R, Minka TP, Deckmyn A (2022) maps: Draw

geographical maps. Available at https://cran.r-project.org/package=maps

Brunson JC (2020) ggalluvial: Layered Grammar for Alluvial Plots. *Journal of Open Source

Software* **5**:2017.

Carruthers EH, Schneider DC, Neilson, JD (2009) Estimating the odds of survival and

identifying mitigation opportunities for common bycatch in pelagic longline fisheries.

Biological Conservation **142**, 2620–2630. https://doi.org/10.1016/j.biocon.2009.06.010

Clarke S, Sato M, Small C, Sullivan B, Inoue Y, Ochi D (2015) Bycatch in longline fisheries

for tuna and tuna-like species; A global review of status and mitigation measures. *FAO

Fisheries and Aquaculture Technical Paper* **588**. Available at

https://www.fao.org/3/a-i4017e.pdf.

Coelho R, Santos MN, Amorim S (2012) Effects of hook and bait on targeted and bycatch

fishes in an equatorial Atlantic pelagic longline fishery. *Bulletin of Marine Science* **88**, 449–467.

Cooke SJ, Suski CD (2004) Are circle hooks an effective tool for conserving marine and freshwater recreational catch-and-release fisheries? *Aquatic Conservation: Marine and Freshwater Ecosystems* **14**, 299–326.

Curran D, Beverly S (2012) Effects of 16/0 circle hooks on pelagic fish catches in three south pacific albacore longline fisheries. *Bulletin of Marine Science* **88**, 485–497. https://doi.org/10.5343/BMS.2011.1060

Dias MP, Martin R, Pearmain EJ, Burfield IJ, Small C, Phillips RA, Yates O, Lascelles B, Borboroglu PG, Croxall JP (2019) Threats to seabirds: A global assessment. *Biological Conservation* **237**, 525–537.

Diaz GA (2008) The effect of circle hooks and straight (J) hooks on the catch rates and numbers of white marlin and blue marlin released alive by the U.S. pelagic longline fleet in the Gulf of Mexico. *North American Journal of Fisheries Management* **28**, 500–506. https://doi.org/10.1577/m07-089.1

Epperly SP, Watson JW, Foster DG, Shah AK (2012) Anatomical hooking location and condition of animals captured with pelagic longlines: The grand banks experiments 2002-2003. *Bulletin of Marine Science* **88**, 513–527.

Fernandez-Carvalho J, Coelho R, Santos MN, Amorim S (2015) Effects of hook and bait in a tropical northeast Atlantic pelagic longline fishery: Part II—Target, bycatch and discard fishes. *Fisheries Research* **164**, 312–321.

Foster DG, Epperly SP, Shah AK, Watson JW (2012) Evaluation of hook and bait type on the catch rates in the western North Atlantic Ocean pelagic longline fishery. *Bulletin of Marine Science* **88**, 529–545.

Gabry J, Češnovar R (2021) cmdstanr: R Interface to "CmdStan." Available at

502 https://mc-stan.org/cmdstanr/reference/cmdstanr-package.html

503 Gallagher AJ, Orbesen ES, Hammerschlag N, Serafy JE (2014) Vulnerability of oceanic

504  sharks as pelagic longline bycatch. *Global Ecology and Conservation* **1**, 50–59.

505 Gilman E, Chaloupka M, Bach P, Fennell H, Hall M, Musyl M, Piovano S, Poisson F, Song L

506  (2020) Effect of pelagic longline bait type on species selectivity: a global synthesis of

507  evidence. *Reviews in Fish Biology and Fisheries* **30**, 535–551.

508  https://doi.org/10.1007/S11160-020-09612-0/TABLES/2

509 Gilman E, Chaloupka M, Swimmer Y, Piovano S (2016) A cross-taxa assessment of pelagic

510  longline by-catch mitigation measures: conflicts and mutual benefits to elasmobranchs.

511  *Fish and Fisheries* **17**, 748–784. https://doi.org/10.1111/faf.12143

512 Gilman E, Kobayashi D, Swenarton T, Brothers N, Dalzell P, Kinan-Kelly I (2007) Reducing

513  sea turtle interactions in the Hawaii-based longline swordfish fishery. *Biological*

514  *Conservation* **139**, 19–28. https://doi.org/10.1016/j.biocon.2007.06.002

515 Gilman E, Zollett E, Beverly S, Nakano H, Davis K, Shiode D, Dalzell P, Kinan I (2006)

516  Reducing sea turtle by-catch in pelagic longline fisheries. *Fish and Fisheries* **7**, 2–23.

517  https://doi.org/10.1111/j.1467-2979.2006.00196.x

518 Godin AC, Carlson JK, Burgener V (2012) The effect of circle hooks on shark catchability

519  and at-vessel mortality rates in longlines fisheries. *Bulletin of Marine Science* **88**,

520  469–483. https://doi.org/10.5343/bms.2011.1054

521 Hall MA, Alverson DL, Metuzals KI (2000) By-catch: problems and solutions. *Marine*

522  *Pollution Bulletin* **41**, 204–219. https://doi.org/10.1016/S0025-326X(00)00111-9

523 Hiraoka Y, Kanaiwa M, Ohshimo S, Takahashi N, Kai M, Yokawa K (2016) Relative

524  abundance trend of the blue shark *Prionace glauca* based on Japanese distant-water and

525  offshore longliner activity in the North Pacific. *Fisheries Science* **82**, 687–699.

526  https://doi.org/10.1007/s12562-016-1007-7

Horodysky AZ, Graves JE (2005) Application of pop-up satellite archival tag technology to

estimate postrelease survival of white marlin (*Tetrapturus albidus*) caught on circle and

straight-shank ("J") hooks in the western North Atlantic recreational fishery. *Fishery*

Bulletin **103**, 84.

Huang HW, Swimmer Y, Bigelow K, Gutierrez A, Foster DG (2016) Influence of hook type

on catch of commercial and bycatch species in an Atlantic tuna fishery. *Marine Policy* **65**,

68–75. https://doi.org/10.1016/j.marpol.2015.12.016

ICCAT (2019) Report of the 2019 Shortfin mako shark stock assessment update meeting.

Collective Volume of Scientific Papers, ICCAT **76**, 1–77.

Kay M (2023) tidybayes: Tidy data and geoms for Bayesian models. Available at

https://doi.org/10.5281/zenodo.1308151

Kerstetter DW, Graves JE. (2006) Effects of circle versus J-style hooks on target and

non-target species in a pelagic longline fishery. *Fisheries Research* **80**, 239–250.

https://doi.org/10.1016/j.fishres.2006.03.032

Kerstetter DW, Luckhurst BE, Prince ED, Graves JE (2003) Use of pop-up satellite archival

tags to demonstrate survival of blue marlin (*Makaira nigricans*) released from pelagic

longline gear. *Fishery Bulletin* **101**, 939–941.

Kiyota M, Yokota K, Nobetsu T, Minami H, Nakano H (2004) Assessment of mitigation

measures to reduce interactions between sea turtles and longline fishery. *Proceedings of*

the International Symposium on SEASTAR 2000 and Bio-Logging Science (The 5th

SEASTAR 2000 Workshop), 24–29.

Kruschke JK (2015) Doing bayesian data analysis: a tutorial with R, JAGS, and Stan. 2nd eds.

(Academic Press, New York, USA)

Maunder MN, Punt AE (2004) Standardizing catch and effort data: a review of recent

approaches. *Fisheries Research* **70**, 141–159.

552 https://doi.org/10.1016/j.fishres.2004.08.002

553 Melvin EF, Guy TJ, Read LB (2014) Best practice seabird bycatch mitigation for pelagic

554 longline fisheries targeting tuna and related species. *Fisheries Research* **149**, 5–18.

555 https://doi.org/10.1016/j.fishres.2013.07.012

556 Pacheco JC, Kerstetter DW, Hazin FH, Hazin H, Segundo RSSL, Graves JE, Carvalho F,

557 Travassos PE (2011) A comparison of circle hook and J hook performance in a western

558 equatorial Atlantic Ocean pelagic longline fishery. *Fisheries Research* **107**, 39–45.

559 https://doi.org/10.1016/j.fishres.2010.10.003

560 Parga ML, Pons M, Andraka S, Rendón L, Mituhasi T, Hall M, Pacheco L, Segura A, Osmond

561 M, Vogel N (2015) Hooking locations in sea turtles incidentally captured by artisanal

562 longline fisheries in the Eastern Pacific Ocean. *Fisheries Research* **164**, 231–237.

563 https://doi.org/10.1016/j.fishres.2014.11.012

564 Pebesma E (2018) Simple features for R: Standardized support for spatial vector data. *The R*

565 *Journal* **10**, 439–446. https://doi.org/10.32614/RJ-2018-009

566 R Core Team (2023) R: A Language and Environment for Statistical Computing. Available at

567 https://www.r-project.org/

568 Reinhardt JF, Weaver J, Latham PJ, Dell'Apa A, Serafy JE, Browder JA, Christman M, Foster

569 DG, Blankinship DR (2018) Catch rate and at-vessel mortality of circle hooks versus

570 J-hooks in pelagic longline fisheries: A global meta-analysis. *Fish and Fisheries* **19**,

571 413–430. https://doi.org/10.1111/faf.12260

572 Santos MN, Coelho R, Fernandez-Carvalho J, Amorim S (2012) Effects of Hook and bait on

573 sea turtle catches in an equatorial Atlantic pelagic longline fishery. *Bulletin of Marine*

574 *Science* **88**, 683–701. https://doi.org/10.5343/bms.2011.1065

575 Santos CC, Rosa D, Coelho R (2020) Progress on a meta-analysis for comparing hook, bait

576 and leader effects on target, bycatch and vulnerable fauna interactions. *Collective Volume*

of Scientific Papers ICCAT **77**, 182–217.

Sims DW, Mucientes G, Queiroz N (2018) Shortfin mako sharks threatened by inaction. *Science* **359**, 1342. https://doi.org/ 10.1126/science.aat0315

Skomal G, Bernal D (2010) Physiological responses to stress in sharks. In 'Sharks and Their Relatives II.' (Eds JC Carrier, JA Musick, MR Heithaus), pp. 475–506. (CRC Press, Boca Raton, USA)

Stokes L, Hataway D, Epperly S, Shah A, Bergmann C, Watson J, Higgins B (2011) Hook ingestion rates in loggerhead sea turtles *Caretta caretta* as a function of animal size, hook size, and bait. *Endangered Species Research* **14**, 1–11. https://doi.org/10.3354/esr00339

Swimmer Y, Gutierrez A, Bigelow K, Barceló C, Schroeder B, Keene K, Shattenkirk K, Foster DG (2017) Sea turtle bycatch mitigation in U.S. longline fisheries. *Frontiers in Marine Science* **4**, 260. https://doi.org/10.3389/FMARS.2017.00260

Stan Development Team (2021) Stan modeling language users guide and reference manual, 2.28.2. Available at https://mc-stan.org

Wallace BP, Kot CY, Dimatteo AD, Lee T, Crowder LB, Lewison RL (2013) Impacts of fisheries bycatch on marine turtle populations worldwide: toward conservation and research priorities. *Ecosphere* **4**, 1–49. https://doi.org/10.1890/ES12-00388.1

Watson JW, Epperly SP, Shah AK, Foster DG (2005) Fishing methods to reduce sea turtle mortality associated with pelagic longlines. *Canadian Journal of Fisheries and Aquatic Sciences* **62**, 965–981. https://doi.org/10.1139/f05-004

Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Grolemund G, Hayes A, Henry L, Hester J et al. (2019) Welcome to the tidyverse. *Journal of Open Source Software* **4**, 1686. https://doi.org/10.21105/joss.01686

Yokota K, Kiyota M, Minami H (2006a) Shark catch in a pelagic longline fishery: Comparison of circle and tuna hooks. *Fisheries Research* **81**, 337–341.

https://doi.org/10.1016/j.fishres.2006.08.006

Yokota K, Kiyota M, Okamura H (2009) Effect of bait species and color on sea turtle bycatch

and fish catch in a pelagic longline fishery. *Fisheries Research* **97**, 53–58.

https://doi.org/10.1016/J.FISHRES.2009.01.003

Yokota K, Minami H, Kiyota M (2006b) Measurement-points examination of circle hooks for

pelagic longline fishery to evaluate effects of hook design. *Bulletin of Fishery Research*

Agency **17**, 83–102

Yokota K, Minami H, Kiyota M (2011) Effectiveness of tori-lines for further reduction of

incidental catch of seabirds in pelagic longline fisheries. *Fisheries Science* **77**, 479–485.

https://doi.org/10.1007/s12562-011-0357-4

Table 1 Fishing effort (longline hooks) in experimental operations used in the analysis.

hook type	bait type		total
	squid	fish	effort
tuna	97,834	71,146	168,980
small-C	70,882	7,658	78,540
large-C	31,872	6,976	38,848
total effort	200,588	85,780	286,368

Table 2 List of hook types used in the experiment.hook name	size	category	straight total length(mm)	maximum total width(mm)
Komatu Keisaku tunahook	4.0 sun	tuna	63	63
Doitomi Tunamutsu	4.0 sun	small-C (circle hook)	58	69
Komatsu Keisaku modified	4.0 sun	small-C (circle hook)	60	68
Komatsu Keisaku modified	4.5 sun	small-C (circle hook)	63	71
Komatsu Keisaku modified	4.8 sun	small-C (circle hook)	N/A	N/A
Komatsu Keisaku type Koshina	4.5 sun	small-C (circle hook)	62	76
Komatsu Keisaku type North America	4.3 sun	small-C (circle hook)	57	63
Tankichi Uruwa	3.8 sun	small-C (circle hook)	56	64
Komatsu Keisaku modified	5.2 sun	large-C (circle hook)	74	81
Komatsu Keisaku type North America	5.2 sun	large-C (circle hook)	76	85
Pacific Fishing Tackle circle hook	18/0	large-C (circle hook)	68	80

Table 3 Number of fish caught by hook and bait type in the experimental operation

species	total catch	hook type			bait type	
		tuna	small-C	large-C	squid	fish
blue shark	13,018	8,084	3,562	1,372	8,903	4,115
longnose lancetfish	1,297	945	257	95	692	605
common dolphinfish	505	206	181	118	363	142
shortfin mako shark	485	298	134	53	262	223
swordfish	288	129	112	47	249	39
bigeye tuna	269	114	146	9	201	68
loggerhead turtle	268	128	113	27	259	9
escolar	163	79	56	28	108	55
striped marlin	145	70	53	22	126	19
other species	1,578	-	-	-	-	-

Table 4 Composition of hooking location and fate at hauling.

species	hooking location				fate at haulback		
	swallowed	mouth	external	unknown (*)	alive	dead	unknown (*)
blue shark	2,270	2,608	66	8,074	11,701	1,066	251
longnose lancetfish	25	288	16	968	165	1,010	122
common dolphinfish	39	159	4	303	398	87	20
shortfin mako shark	66	68	18	333	363	118	4
swordfish	59	118	18	93	55	226	7
bigeye tuna	7	128	2	132	83	183	3
loggerhead turtle	109	116	14	29	256	5	7
escolar	1	62	1	99	108	40	15
striped marlin	12	80	8	45	68	76	1

(*) Includes catches that dropped off before reseachers checked or lack of survey.

Table 5 Estimates of the posterior distribution of the proportion of hooking location by hook when squid bait is used (median). Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses.

species	hook type		
	tuna	small-C	large-C
hooking place: external			
blue shark	0.020 (0.015 - 0.027)	0.008 (0.005 - 0.012)	0.005 (0.001 - 0.014)
loggerhead turtle	0.025 (0.006 - 0.066)	0.079 (0.037 - 0.143)	0.031 (0.001 - 0.154)
longnose lancetfish	0.049 (0.020 - 0.092)	0.031 (0.011 - 0.067)	0.074 (0.018 - 0.185)
shortfin mako	0.102 (0.041 - 0.199)	0.107 (0.052 - 0.185)	0.136 (0.021 - 0.381)
striped marlin	0.083 (0.026 - 0.186)	0.068 (0.016 - 0.173)	0.048 (0.002 - 0.235)
swordfish	0.075 (0.030 - 0.148)	0.109 (0.057 - 0.184)	0.031 (0.001 - 0.154)
hooking place: mouth			
blue shark	0.497 (0.475 - 0.518)	0.548 (0.528 - 0.568)	0.560 (0.517 - 0.603)
loggerhead turtle	0.439 (0.346 - 0.534)	0.480 (0.381 - 0.577)	0.746 (0.544 - 0.892)
longnose lancetfish	0.857 (0.792 - 0.908)	0.903 (0.848 - 0.943)	0.872 (0.740 - 0.953)
shortfin mako	0.356 (0.238 - 0.486)	0.524 (0.418 - 0.630)	0.379 (0.154 - 0.650)
striped marlin	0.759 (0.623 - 0.869)	0.831 (0.696 - 0.924)	0.883 (0.660 - 0.982)
swordfish	0.528 (0.416 - 0.637)	0.676 (0.574 - 0.767)	0.611 (0.405 - 0.790)
hooking place: swallowed			
blue shark	0.483 (0.461 - 0.505)	0.444 (0.424 - 0.464)	0.434 (0.391 - 0.478)
loggerhead turtle	0.533 (0.438 - 0.626)	0.438 (0.343 - 0.537)	0.210 (0.078 - 0.405)
longnose lancetfish	0.092 (0.052 - 0.148)	0.064 (0.032 - 0.111)	0.046 (0.007 - 0.145)
shortfin mako	0.536 (0.405 - 0.665)	0.365 (0.266 - 0.472)	0.460 (0.213 - 0.724)
striped marlin	0.150 (0.066 - 0.273)	0.093 (0.030 - 0.207)	0.048 (0.002 - 0.234)
swordfish	0.394 (0.289 - 0.505)	0.211 (0.136 - 0.303)	0.343 (0.174 - 0.550)

Table 6 Estimated haulback mortality rates (median of posterior distribution) by hooking location when squid bait is used. Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses. Loggerhead turtles were excluded because there were no mortalities and the calculation had not been converged.

species	external	mouth	swallowed
blue shark	0.295 (0.185 - 0.411)	0.052 (0.044 - 0.061)	0.157 (0.141 - 0.172)
shortfin mako shark	0.496 (0.262 - 0.729)	0.134 (0.061 - 0.219)	0.279 (0.176 - 0.394)
longnose lancetfish	0.956 (0.810 - 1.000)	0.860 (0.816 - 0.899)	0.798 (0.621 - 0.934)
swordfish	0.802 (0.595 - 0.959)	0.776 (0.694 - 0.853)	0.919 (0.842 - 0.979)
striped marlin	0.798 (0.488 - 0.990)	0.431 (0.315 - 0.547)	0.968 (0.839 - 1.000)

Table 7 Haulback mortality rate by hook and bait type (median of posterior distribution). Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses.

Species	Hook type		
	tuna	small-C	large-C
Bait type: squid			
bigeye tuna	0.735 (0.638 - 0.824)	0.757 (0.680 - 0.830)	0.960 (0.812 - 1.000)
blue shark	0.084 (0.077 - 0.092)	0.081 (0.072 - 0.091)	0.077 (0.062 - 0.093)
common dolphinfish	0.154 (0.099 - 0.214)	0.128 (0.081 - 0.181)	0.180 (0.108 - 0.264)
escolar	0.190 (0.087 - 0.313)	0.300 (0.164 - 0.453)	0.182 (0.061 - 0.341)
longnose lancetfish	0.901 (0.870 - 0.930)	0.925 (0.893 - 0.954)	0.890 (0.827 - 0.941)
shortfin mako	0.161 (0.105 - 0.221)	0.187 (0.122 - 0.257)	0.182 (0.081 - 0.297)
striped marlin	0.531 (0.399 - 0.663)	0.472 (0.335 - 0.612)	0.530 (0.314 - 0.736)
swordfish	0.829 (0.755 - 0.895)	0.783 (0.700 - 0.860)	0.833 (0.713 - 0.933)
Bait type: fish			
bigeye tuna	0.473 (0.324 - 0.626)	0.503 (0.337 - 0.665)	0.886 (0.569 - 1.000)
blue shark	0.078 (0.068 - 0.088)	0.075 (0.063 - 0.088)	0.071 (0.056 - 0.088)
common dolphinfish	0.238 (0.156 - 0.325)	0.200 (0.103 - 0.311)	0.274 (0.163 - 0.398)
escolar	0.242 (0.114 - 0.390)	0.370 (0.172 - 0.588)	0.233 (0.056 - 0.468)
longnose lancetfish	0.909 (0.879 - 0.936)	0.931 (0.895 - 0.961)	0.898 (0.840 - 0.946)
shortfin mako	0.307 (0.235 - 0.379)	0.347 (0.227 - 0.477)	0.340 (0.196 - 0.493)
striped marlin	0.691 (0.459 - 0.887)	0.638 (0.373 - 0.869)	0.690 (0.387 - 0.923)
swordfish	0.853 (0.728 - 0.948)	0.812 (0.652 - 0.939)	0.857 (0.698 - 0.967)

Table 8 Standardized catch per unit effort (CPUE) by hook and bait type (median of posterior distribution). Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses.

Species	Hook type		
	tuna	small-C	large-C
Bait type: squid			
bigeye tuna	0.420 (0.046 - 1.195)	1.095 (0.140 - 3.127)	0.882 (0.075 - 2.969)
blue shark	42.648 (34.036 - 52.508)	71.133 (56.592 - 87.324)	60.571 (47.909 - 74.668)
common dolphinfish	0.756 (0.355 - 1.307)	1.839 (0.850 - 3.179)	1.349 (0.588 - 2.339)
escolar	0.235 (0.083 - 0.452)	0.702 (0.257 - 1.358)	0.371 (0.118 - 0.764)
longnose lancetfish	3.143 (2.555 - 3.783)	5.023 (4.036 - 6.073)	4.194 (3.171 - 5.292)
shortfin mako	1.030 (0.626 - 1.483)	1.953 (1.202 - 2.855)	1.597 (0.902 - 2.423)
striped marlin	0.378 (0.204 - 0.582)	0.465 (0.247 - 0.728)	0.428 (0.201 - 0.745)
swordfish	0.785 (0.539 - 1.065)	1.186 (0.795 - 1.618)	1.268 (0.799 - 1.831)
Bait type: fish			
bigeye tuna	0.158 (0.016 - 0.473)	0.410 (0.036 - 1.227)	0.330 (0.025 - 1.177)
blue shark	35.877 (28.719 - 44.397)	59.875 (47.685 - 73.886)	50.932 (40.260 - 62.990)
common dolphinfish	1.853 (0.845 - 3.252)	4.502 (2.013 - 7.853)	3.298 (1.493 - 5.894)
escolar	0.499 (0.171 - 1.000)	1.488 (0.484 - 2.986)	0.787 (0.224 - 1.710)
longnose lancetfish	3.592 (2.863 - 4.399)	5.745 (4.546 - 7.137)	4.802 (3.532 - 6.226)
shortfin mako	2.262 (1.365 - 3.302)	4.291 (2.570 - 6.365)	3.524 (1.960 - 5.451)
striped marlin	0.369 (0.163 - 0.639)	0.452 (0.185 - 0.800)	0.417 (0.153 - 0.806)
swordfish	0.809 (0.495 - 1.188)	1.222 (0.725 - 1.824)	1.310 (0.724 - 2.059)

Table 9 Estimated MPUE by hook and bait type (median of posterior distribution). Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses.

Species	Hook type		
	tuna	small-C	large-C
Bait type: squid			
bigeye tuna	0.307 (0.032 - 0.881)	0.826 (0.082 - 2.339)	0.828 (0.062 - 2.790)
blue shark	3.589 (2.802 - 4.495)	5.761 (4.420 - 7.264)	4.668 (3.390 - 6.176)
common dolphinfish	0.116 (0.046 - 0.217)	0.233 (0.092 - 0.440)	0.241 (0.086 - 0.466)
escolar	0.044 (0.010 - 0.098)	0.207 (0.056 - 0.444)	0.066 (0.010 - 0.170)
longnose lancetfish	2.829 (2.277 - 3.399)	4.639 (3.709 - 5.612)	3.718 (2.804 - 4.751)
shortfin mako	0.165 (0.087 - 0.269)	0.362 (0.187 - 0.588)	0.288 (0.102 - 0.544)
striped marlin	0.199 (0.099 - 0.322)	0.218 (0.103 - 0.362)	0.223 (0.081 - 0.423)
swordfish	0.648 (0.438 - 0.889)	0.925 (0.606 - 1.274)	1.048 (0.634 - 1.534)
Bait type: fish			
bigeye tuna	0.074 (0.006 - 0.225)	0.203 (0.019 - 0.633)	0.277 (0.018 - 1.005)
blue shark	2.797 (2.134 - 3.535)	4.482 (3.347 - 5.808)	3.627 (2.578 - 4.877)
common dolphinfish	0.437 (0.163 - 0.809)	0.895 (0.299 - 1.814)	0.893 (0.325 - 1.741)
escolar	0.119 (0.028 - 0.272)	0.540 (0.123 - 1.243)	0.180 (0.022 - 0.506)
longnose lancetfish	3.263 (2.595 - 4.005)	5.337 (4.219 - 6.662)	4.297 (3.145 - 5.603)
shortfin mako	0.693 (0.390 - 1.063)	1.480 (0.727 - 2.425)	1.182 (0.511 - 2.087)
striped marlin	0.248 (0.092 - 0.455)	0.280 (0.089 - 0.541)	0.277 (0.078 - 0.586)
swordfish	0.683 (0.397 - 1.014)	0.977 (0.535 - 1.488)	1.103 (0.581 - 1.770)

Table 10 Nominal CPUE, Haulback mortality rate and MPUE of Loggerhead turtle and all figures are based on aggregated operational data, not estimates.

hook type	CPUE		haulback mortality		MPUE	
	squid	fish	squid	fish	squid	fish
tuna	1.247	0.084	0.0385	0.000	0.0480	0.000
large-C	0.941	0.287	0.0090	0.000	0.0085	0.000
small-C	2.257	0.261	0.0246	0.000	0.0555	0.000

Figure 1 Locations where the longline operation experiment was conducted.

Figure 2 Differences in the estimated probability of the "swallowed" hooking location of each circle hook type from tuna hook when squid bait is used. The red dotted line indicates that the difference is zero.

Figure 3　Differences in the estimated haulback mortality rate of each target hooking location from "swallowed" hooking location when squid bait is used. The red dotted line indicates that the difference is zero.

Figure 4 Differences in estimated haulback mortality between each experimental group and the control group (squid x tuna hook). The red dotted line indicates that the difference is zero.

Figure 5 Relationship between sea surface temperature (SST) variability and haulback mortality rate at longline operations. Solid lines indicate median; masked areas indicate 95% Bayesian credible interval.

Figure 6 Relationship between soak time (time from setting the branch line to hauling) variability and haulback mortality rate at longline operations. Solid lines indicate median; masked areas indicate 95% Bayesian credible interval.

Figure 7　Differences in the standardized catch per unit effort (CPUE) for each experimental group from those for the control group (squid x tuna hook). The red dotted line indicates that the difference is zero.

Figure 8   Differences in the estimated MPUE (mortality per unit effort) between those for each experimental group and those for the control group (squid x tuna hook). The red dotted line indicates that the difference is zero.

Figure 9 Alluvial plot of hooking locations and associated haulback mortality rates of loggerhead turtles by hook when squid bait is used.

Appendix 1 Detailed measurements of hooks used in the experiment and a figure explaining measurement points of hooks (copied from Yokota et al. 2006b).

manufacture	Komatsu Keisaku	Hisamatsu Tankichi	Doitomi	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Pacific Fishing Tackle MFG., CO.
hook name	tuna hook	Uruwa hook BKN	tuna circle hook SS-170	modified circle hook	modified circle hook	modified circle hook	cirlce hook type Koshina	cirlce hook type North America	cirlce hook type North America	modified circle hook	circle hook
standardized size	4.0 sun	3.8 sun	#4	4.0 sun	4.5 sun	4.8 sun	4.5 sun	4.3 sun	5.2 sun	5.2 sun	18/0
material	stainless steel	hard steel	stainless steel	stainless steel	stainless steel	stainless steel	stainless steel	hard steel	hard steel	stainless steel	stainless steel
hook eye	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring
shank thickness(mm)	5.3	4.0	4.1	5.3	5.2	N/A	4.9	5.3	5.7	5.3	5.1
straight total length(mm)	63	56	58	60	63	N/A	62	57	74	76	68
straight total width(mm)	38	44	49	47	49	N/A	56	45	56	54	59
minimum total width(mm)	38	36	39	41	45	N/A	51	41	52	48	51
maximum total width(mm)	63	64	69	68	71	N/A	76	63	81	85	80
front length(mm)	41	33	38	35	44	N/A	47	39	49	47	45
minimum inner width(mm)	27	20	15	24	25	N/A	26	20	27	26	27
L-W ratio	1.7	1.3	1.2	1.3	1.3	N/A	1.1	1.3	1.3	1.4	1.2
max-min ratio	1.7	1.8	1.8	1.7	1.6	N/A	1.5	1.5	1.6	1.8	1.6
incurved point angle		80	70	90	70	N/A	70	65	75	80	80
offset angle	5°≦θ＜10°	θ≈0°	θ≈0°	θ≈10°	5°≦θ＜10°	θ＜10°	θ≈10°	5°≦θ＜10°	10°≦θ＜15°	θ≈5°	10°≦θ＜15°
offset width(mm)	0.9	≈0	≈0	1.8	2.6	N/A	2.1	1.5	3.7	1.0	5.4
weight (g)	19.9	12.2	15.0	19.7	21.6	N/A	21.4	19.4	30.3	25.5	23.2

Appendix 2 Stan code used for MCMC sampling of (a) Model 1 and (b) Model 2

(a)   MODEL 1

```
data {
 int HK_cat;
 int<lower=1> N1;
 Int<lower=1> LOC_cat;
 int<lower=1, upper=LOC_cat> LOC[N1];
 int<lower=1, upper=HK_cat> HK[N1];
 int<lower=0, upper=1> M[N1];
 int L;
 matrix[N1, L] X1;
 Int N2;
 Int Catch[N2];
 Int L2;
 matrix[N2, L2] X2;
 vector[N2] O;
 Int YRID;
 int YR[N2];
}

transformed data{
 vector[HK_cat] Zeros;
 Zeros = rep_vector(0, HK_cat);
}

parameters {
 matrix[HK_cat, LOC_cat - 1] theta_raw;
 vector[L] beta;
 vector[L2] beta2;
 vector[YRID] r;
 real<lower= 0> sigma;

}

transformed parameters {
 vector[N2] mu;
 vector[N1] phi;
 matrix[HK_cat, LOC_cat] theta;
 phi = inv_logit(X1 * beta);
 theta = append_col(Zeros, theta_raw);
 for(i in 1:N2)
 mu[i] = X2[i,] * beta2 + r[YR[i]];

model {
 sigma ~ student_t(4, 0, 2.5);
 r ~ normal(0, sigma);
 for(n in 1:N1){
 M[n] ~ bernoulli(phi[n]);
 target += categorical_lpmf(LOC[n]|softmax(theta[HK[n],]'));
 }
 Catch ~ poisson(exp(mu + log(O)));
}
```

(b)   MODEL 2

```
data {
 int HK_BAIT_comb;
 int<lower=1> N1;
 int<lower=0, upper=1> M[N1];
 int L;
 matrix[N1, L] X1;
 int N2;
 int Catch[N2];
 int L2;
 matrix[N2, L2] X2;
 vector[N2] O;
 int YRID;
 int YR[N2];
}

parameters {
 vector[L] beta;
 vector[L2] beta2;
 vector[YRID] r;
 real<lower= 0> sigma;
}

transformed parameters {
 vector[N2] mu;
 vector[N1] phi;
 phi = inv_logit(X1 * beta);
 for(i in 1:N2)
 mu[i] = X2[i,] * beta2 + r[YR[i]];
}

model {
 sigma ~ student_t(4, 0, 2.5);
 r ~ normal(0, sigma);
 for(n in 1:N1){
 M[n] ~ bernoulli(phi[n]);
 }
 Catch ~ poisson(exp(mu + log(O)));
}
```

Appendix 3 Size distributions by species for the major species captured in the study, with body

length as an index of precaudal length for blue sharks and shortfin mako sharks, straight-line

carapace length for loggerhead turtles, eye-to-fork length for striped marlin and swordfish, and fork

length for all other species.