TITLE:

Multifaceted effects of bycatch mitigation measures on target/non-target species for pelagic longline

fisheries and consideration for bycatch management

AUTHORS:

Daisuke Ochi[#], Kei Okamoto and Shintaro Ueno

AFFILIATION:

Fisheries Resources Institute, Japan Fisheries Research and Education Agency

Number of tables: 11

Number of figures: 9

[#]Corresponding author,

Address: Fisheries Resources Institute, Japan Fisheries Research and Education Agency, 2-12-4,

Fukuura, Kanazawa-ku, Yokohama-city, Kanagawa 2368648, Japan

e-mail: otthii80s@gmail.com

SUMMARY

The pelagic longline fishery, in an effort to reduce bycatch of sea turtles, have developed and deployed fisheries bycatch mitigation techniques such as replacing J/tuna hooks and squid bait with circle hooks and whole fish bait. However, little emphasis has been placed on the side effects of bycatch mitigation measures on endangered species other than target bycatch species. Several previous studies of the side effects have been marred by lack of control for the covariates. Here, based on long-term data obtained from research cruises by a pelagic longline vessel, we examined the effects of using circle hooks and whole fish bait to replace squid bait on the fishing mortality of target and non-target fishes, and also bycatch species. A quantitative evaluation analysis of our results, based on a Bayesian approach, showed the use of circle hooks to increase mouth hooking in target and bycatch species, and their size to be proportional to the magnitude of the effect. Although deploying circle hooks increased catch per unit effort (CPUE) and fishing mortality per unit effort (MPUE), changing the bait species from squid to fish clearly had a far greater impact on shark mortality than the use of circle hooks. Because the impact of the introduction of bycatch mitigation measures on species other than the focused bycatch species is non-negligible, a quantitative assessment of bycatch mitigation-related fishing mortality is critical before introducing such measures.

Keywords: billfishes, bycatch, bycatch mitigation measure, fisheries management, sea turtle, sharks, tuna, longline fishery

INTRODUCTION

Unintentional catch by fisheries is referred to as bycatch, and calls for its reduction are global, especially because the bycatch of endangered species can have devastating effects on such populations. For example, in the tuna longline fishery, concerns about the increased conservation risk for seabirds and sea turtles by unintentional and fatal catch have been a major issue among many regional fisheries management organizations (RFMOs) since the 1990s (Wallace et al. 2013; Dias et al. 2019), and in recent years, some elasmobranchs, whose populations are declining further, are also treated as bycatch species. The decline of these species has focused attention at both the national and the RFMO levels. Even for target fish that are not bycatch species, addressing the deterioration of stock status caused by overfishing requires reductions in unintentional fishing mortality (for instance, billfishes in the North Atlantic; Kerstetter & Graves 2006; Diaz 2008).

Several studies have weighed the sustainability of tuna longline fisheries against the conservation of species vulnerable to bycatch (Hall et al. 2000; Melvin et al. 2014; Clarke et al. 2015)—particularly for seabirds and marine turtles—with the development of several effective bycatch mitigation measures (Melvin et al. 2014; Swimmer et al. 2017). Bycatch mitigation measures in longline fisheries target specific animal groups and are evaluated based on their success in reducing mortality due to bycatch of specific vulnerable species. While the impact on the catch of the target fish is the primary consideration when evaluating bycatch mitigation techniques, few studies have examined the impacts and tradeoffs on species not targeted by mitigation techniques (Pacheco et al. 2011; Gilman et al. 2016). However, an ecosystem-based fisheries management approach to the introduction of bycatch reduction measures demands the assessment of the impact not only on the target bycatch species but also on the ecosystem itself in which the target species lives (Reinhardt et al. 2018).

The circle hook and finfish bait are typical sea turtle bycatch mitigation measures for pelagic longline fisheries (Watson et al. 2005; Gilman et al. 2006; Yokota et al. 2009; Stokes et al. 2011). The tip of the circle hook bends inward, and when a fish or sea turtle swallows the hooked bait, the circle hook cannot hook inside the digestive tract; instead, as the hook exits the mouth, a torque force causes it to hook through the edge of the mouth. This property allows for easy hook removal and has been reported to reduce the mortality rate of bycatch sea turtles on board and after release (Kiyota et al. 2004; Cooke & Suski 2004; Kerstetter & Graves 2006). Reports of positive effects of circle hooks include those for other

species—such as reduced haulback and post-release mortality in sharks, reduced post-release mortality

in swordfish, and increased catch rates in tuna, the target fish. The use of fish bait instead of squid bait

potentially reduces bycatch (Watson et al. 2005; Yokota et al. 2009, 2011) and mortality rates (Stokes et

al. 2011; Parga et al. 2015) of sea turtles.

These sea turtle mitigation measures, however, exact other costs. Several meta-analytic studies

have reported the circle hook-related reduction of sea turtle mortality but increase in billfish and shark

catch rates (Reinhardt et al. 2018). The use of whole fish bait also reportedly increases the catch of

sharks and other species (Foster et al. 2012). However, few studies have allowed for quantitative

evaluation of the effects of these mitigation measures experimentally on species beyond sea turtles. This

is due in part to the reliance on observer data from commercial vessels, and small sample sizes, small

comparison groups, and lack of experimental rigor from research vessels. In addition, the impact

assessment for sharks underestimates catch rates and mortality associated with missed catches due to

"bite-off" branchline (Reinhardt et al. 2018) and several other issues. In addition, many experimental

studies and meta-analyses only evaluate gear impacts at significance levels without evaluating the

magnitude of the effect. Small significant differences may be judged not to matter much when assessing

overall risk in bycatch species. Studies that controlled for these conditions would allow for an evaluation

of adverse effects without the confounding problems described above. Also, many studies use

catch/bycatch rates and mortality rates (at haulback or after release) as important impact indicators,

without considering irreversible impacts, such as the number of organisms killed at the time of catch.

Additionally, hooking position itself is believed to have a strong influence on mortality rate, which

demands the estimation of risk under specific hooking conditions, taking causal relationships into

account.

Here, we used data from controlled experiments to analyze these confounding factors and

developed a Bayesian statistical model to evaluate the effects of changing hook and bait species on fish

species other than turtles—particularly, tuna, swordfish, and sharks—and then verified the contribution of

circle hooks and fish bait to mortality rate, catch rate, and fatal catch rate, respectively. We also discussed

the appropriate assessment of bycatch mitigation measures in fisheries management.

METHOD

Experimental Operations

We analyzed data from the R/V Taikei No. 2 longline research operation conducted in the Northwest

Pacific Ocean between 2002 and 2010—a typical Japanese shallow-setting operation targeting mainly

swordfish and shark, using four hooks per basket, a wire leader, and a night soaking (Yokota et al. 2006a).

We regarded the 4.0-sun tuna hooks as a control group against which we assessed 10 different sized

circle hooks (Table 1). In Appendix 1 we describe hook shapes and other details of these hooks following

the measurement method of Yokota et al. (2006b). Since the circle hooks varied in size and sample

numbers were too small to analyze each hook separately, we divided them into two groups according to

size. Because degree of hook offset has been reported to affect catch and hooking position (Cooke &

Suski 2004), most hooks were circle hooks <10°but some were nearly 15°. The bait comprised chub

mackerel *Scomber japonicus* and Japanese common squid *Todarodes pacificus* in the range of 20–30 cm

in fork length or dorsal mantle length. These were frozen and stored, then completely thawed before

being hooked. The sequence of line setting was divided into several experimental segments, and a

different combination of hook and bait type was applied for each segment, with an alternate order of

segments at each operation.

The researcher recorded catch for each operation for all catch/bycatch, and the species caught,

status of haulback (alive or dead), hooking location (mouth, swallow, and body hooking), time of catch,

and basket number. Hook location was recorded only for squid bait. At the start of longline operations, the

researcher also collected sea surface temperature, which we subsequently used in the analysis.

Since the experiment was conducted using multiple sizes (degrees) of circle hooks in

combination with whole fish bait and squid bait, we categorized hook shapes into three main types:

1. control (tuna hook 4.0-sun; tuna)

2. smaller circle hook (smaller than 5.0-sun; small-C)

3. larger circle hook (18/0 and 5.0-sun or more; large-C)

In Table 1 we show annual longline effort separated by bait and hook type, tabulated according to

the above categorization. We selected the following species for analysis: blue shark *Prionace glauca*,

shortfin mako shark *Isurus oxyrinchus*, bigeye tuna *Thunnus obesus*, swordfish *Xiphias gladius*, striped

marlin *Kajikia audax*, common dolphinfish *Coryphaena hippurus*, escolar *Lepidocybium*

*flavobrunneum*and, longnosed lancetfish *Alepisaurus ferox*, and loggerhead turtles *Caretta caretta* (to

confirm the effect on sea turtle bycatch).

Statistical Analysis

We conducted all analyses using a Bayesian approach to estimate parameters. We adopted

haulback mortality rate, catch per unit effort (CPUE), and mortality per unit effort (MPUE) (Afonso et al.

2011) as indices to evaluate the impact of hook and bait type on fishing mortality within the analyses. We

used the following data as inputs to the model: number of (units) caught, longline effort (number of hooks),

fate at hauling, hooking location, year and location of operation, water temperature at operation, and

soaking time.

Because of the heterogeneity of the data set due to some missing data, we split the analysis into

two models. MODEL 1 relates only to relationships between hooking location and haulback mortality rate,

while MODEL 2 comprises two subsets for estimating haulback mortality rate, CPUE and MPUE.

We based MODEL 1 on a logit regression using a Bernoulli distribution. The model assumes that

hooking position is most directly related to mortality rate and attempts to express the indirect effect of

hook type by estimating probability of hooking location for each hook type. We express the observed

hooking location *H* and haulback mortality rate *M* in MODEL 1 by the following

equations:

$$M \sim Bernoulli(p_{dead}) \#(1)$$

$$logit(p_{dead}) = \beta_{1,hook} + \beta_{1,sst} + \beta_{1,soaktime} \#(2)$$

$$H \sim Categorical\left(softmax(\theta_{hook})\right) \#(3)$$

where β_1 is the parameter in each explanatory variable, p_{dead} is the expected haulback mortality rate, and

θ_{hook} is the expected probability of hooking location.

We structured MODEL 2 to calculate the expected number of mortalities per effort (MPUE)

based on the parameters estimated in both the mortality and CPUE estimation subsets. In MODEL 2, due

to lack of hooking location data, we calculate p_{dead} from a modified equation (2) as follows;

$$logit(p_{dead}) = \beta_{2,hook} + \beta_{2,bait} + \beta_{2,sst} + \beta_{2,soaktime} \#(4)$$

152   where β_2 is the parameter in each explanatory variable.

153   The CPUE subset is almost same as a GLMM analysis in assuming the Poisson distribution as

154   the error structure. We express the number of catches C per operation using the expected CPUE λ as

155   follows:

$$C \sim Poisson(\lambda + \log(E)) \#(5)$$

$$\lambda = \log(\gamma_{hook} + \gamma_{bait} + \gamma_{latitude} + \gamma_{sst} + r_{year}) \#(6)$$

$$r_{year} \sim N(0, \ \sigma^2) \#(7)$$

157   where E is the longline effort, γ is a parameter in each explanatory variable, r_{year} is a random effect of

158   annual fluctuation on CPUE, and σ is a standard deviation.

159   We obtained the expected values of hook-bait-specific MPUE ζ by multiplying CPUE by

160   at-haulback mortality rate as follows:

$$\zeta = \widehat{p_{dead}} \times \hat{\lambda} \#(9)$$

where $\widehat{p_{dead}}$ denotes the expected mortality rate standardized for hook and bait type, and $\hat{\lambda}$ denotes the

estimated CPUE standardized for hook and bait type. We used the standardization method for

abundance indices used in fisheries stock assessment (Maunder & Punt 2004). In this method,

explanatory variables other than the factor subject to standardization are averaged to predict the objective

variable, which in stock assessment is a time scale such as years or months, but in our case we modified

the standardization scale to reference hook type and bait type.

We calculated each parameter based on a Bayesian approach with Markov chain Monte Carlo

(MCMC) sampling. For the MCMC sampling, we used cmdstan 2.28.2 (Stan Development Team 2021).

As a prior distribution for each parameter, we used a half student-t distribution with 2 degrees of freedom,

mean 0, and variance 2.5 for σ, and we used a uniform distribution for the other parameters. We

computed the posterior distribution using Stan with 15000 sampling iterations including 10000 warmup

iterations, number of chains as 4, and no sinning. We calculated Bayesian credible intervals based on the

highest density interval (HDI) for the estimates. Although the Bayesian approach for the estimates

precluded significance testing, we determined the difference between the estimates of the experimental

group and those of the control group ("swallowing" for hooking location, "tuna" hook for hook type, and

"squid" for bait type), and if the lower and upper limits of HDI for the difference did not exceed 0, we

considered the difference as a difference for convenience (assuming region of practical equivalence

[ROPE] as 0; Kruschke 2015). In Appendix 2 we show the Stan code used to estimate each parameter in

MODEL 1 and 2. For other data handling, statistical analysis, and plotting, we used R4.1.1 (R Core Team

2021) and packages "cmdstanr 0.4.0," "ggalluvial 0.12.3," "ggthemes 4.2.4," "mapdata 2.3.0," "maps

3.4.0," "sf 1.0–2," "tidybayes 3.0.1," and "tidyverse 1.3.1" (Becker & Brownrigg. 2018; Pebesma 2018;

Wickham et al. 2019; Brunson 2020; Arnold 2021; Richard et al. 2018, 2021; Gabry & Češnovar 2021;

Kay 2021).

RESULTS

A total of 286,363 hooks from 612 operations were deployed in the experiment (Table 2). The area of

operation ranged from around the Izu Islands in Japan to off the east coast of northeastern

Honshu—typical fishing ground for Japanese shallow-setting longliners (Fig. 1; Hiraoka et al. 2016).

Summary Statistics

The most common species caught was blue shark, followed by longnosed lancetfish, common dolphinfish,

and shortfin mako shark (Table 3). Major miscellaneous fish species included, in descending order of

abundance, salmon shark *Lamna ditropis* (N = 229), pelagic stingray *Pteroplatytrygon violacea* (N = 199),

pomflets *Brama spp.* (N = 135), bigeye thresher shark *Alopias superciliosus* (N = 89), and albacore

Thunnus alalunga (N = 69), but for these species the sample sizes were too skewed among experimental

groups to converge the MCMC sampling.

In Table 4 we show the number of fish caught by hook type and bait type. Bigeye tuna had

extremely low catches on large-C hooks, and loggerheads had low catches on fish bait. The most

common hooking position at the time of catch was mouth hooking for all nine species, with extremely few

hook locations other than mouth hooking, especially for bigeye tuna and esclolar (Table 4). The proportion

of fatalities of captured species at haulback varied greatly by species (Table 5). The haulback mortality

rate was low for blue sharks, common dolphinfish, escolar, and shortfin mako sharks and, although higher

for bigeye tuna, longnose lancetfish, striped marlin, and swordfish, was generally extremely low,

especially in loggerhead turtles.

The size composition of the nine species used in the analysis was not statistically compared in

this study and was not included in the model because it did not contribute to haulback mortality rate, but in

Appendix 3 we include a histogram.

Almost all parameters in the models for the nine species were successfully converged (Rhat <

1.1) but some of the parameters for loggerhead turtle were not converged, as we describe below.

Output of MODEL 1

In Table 6 we show percentage estimates of hooking locations by hook type throughout the model. We

observed differences in hooking location by hook type for blue shark, common dolfinfish, loggerhead

turtle, and swordfish, with an increase in mouth hooking and a decrease in swallowing for large-C for

loggerhead turtle and a clear increase in mouth hooking for small-C for shortfin mako shark and swordfish

(Fig. 2). In blue sharks, the frequency of swallowing decreased in both small-C and large-C.

In Table 7 we show haulback mortality rates by hooking location. We observed clear differences

in haulback mortality rate by hooking location for blue shark, shortfin mako shark, striped marlin, and

swordfish (Fig. 3). Haulback mortality rates after hook swallowing for blue sharks were lower than those

for body hooking, and higher than those for mouth hooking. We observed lower haulback mortality rates

for shortfin mako shark, striped marine, and swordfish from hook swallowing than from mouth hooking.

Output of MODEL 2

Haulback mortality was higher for large-C than for tuna hook types in bigeye tuna, but did not differ

among hook types in other species (Table 8, Fig. 4). On the other hand, the haulback mortality for two bait

types differed in bigeye tuna and blue shark, with a rate reduction and increase with the use of fish bait in

bigeye tuna and shortfin mako shark, respectively.

In Fig. 5 and Fig. 6 we show the effects of two environmental factors—sea surface temperature

(SST) and soak time—on haulback mortality rate. The response to SST differed by species, with haulback

mortality rate increasing with higher SST for shortfin mako sharks, swordfish, common dolphinfish, and

escolar, and conversely increasing with lower SST for bigeye and longnose lancetfish. We observed little

fluctuation in haulback mortality rate with SST in blue shark and striped marlin. In general, haulback

mortality rate increased with increasing soak time. However, for bigeye tuna, haulback mortality rate

decreased with soak time, but the trend was not clear, and for blue sharks, haulback mortality rate

increased only slightly with increased soak time.

In Table 9 we show standardized CPUE by hook and bait type. Differences in CPUE among

hook type were found for all species except striped marlin (Fig. 7). We observed higher CPUE for small-C

in bigeye tuna, escolar, longnose lancetfish, and swordfish, and higher CPUE for both large and small-C

in blue shark, shortfin mako shark, and common dolphinfish. Especially in blue sharks, the difference in

CPUE ranged from 10 to 20 depending on the hook type.

We observed differences in standardized CPUE by bait type in bigeye tuna, blue shark, shortfin

mako shark, common dolphinfish, and escolar. CPUE decreased with whole fish bait in bigeye tuna and

blue sharks, but increased in shortfin mako shark, common dolphinfish, and escolar. In the case of blue

shark, escolar, shortfin mako shark, and swordfish, the bait effect could be varied with circle hooks,

suppressing the CPUE-increasing effect by fish bait in bigeye tuna, blue shark, and swordfish with the use

of circle hooks, while conversely this combination boosted increase of CPUE in common dolphinfish,

escolar, and shortfin mako shark.

In Table 10 we show calculated MPUE values by hook and bait type. Compared to differences in

MPUE among hook and bait type, those in CPUE were relatively small. We observed differences in

MPUE by all hook types except for those in common dolphinfish and striped marlin, with higher MPUE for

small-C in bigeye tuna, blue shark, escolar, longnosed lancetfish, and swordfish (Fig. 8). We confirmed

decreases in MPUE by fish bait in bigeye tuna and blue shark, and blue sharks, and conversely,

increases in MPUE in common dolphinfish, shortfin mako shark, and swordfish. In escolar, however,

MPUE increased only when small-C was combined with whole fish bait. The effect of the combination of

whole fish bait and circle hook varied in bigeye tuna, blue shark, common dolphinfish, shortfin mako shark,

and swordfish. In bigeye tuna, blue shark, and swordfish, the effect of the circle hook on MPUE was

suppressed by fish bait, while in common dolphinfish and shortfin mako shark, MPUE increased

significantly when whole fish bait and circle hook were used together.

Hook and bait effect for sea turtle bycatch

We failed to complete our analysis for loggerhead turtles throughout the models because the bias in frequency of capture events among the experimental groups was too large and did not converge except for only a part of MODEL 1. Instead, in Table 11 we show the nominal CPUE, haulback mortality rate, and MPUE for each experimental group, and in Fig. 9, hooking location and haulback mortality rate by each hook type with squid bait. Most individuals survived regardless of hooking location. When whole fish bait was used, haulback mortality rate and associated MPUE were zero. For squid bait, large-C had the smallest CPUE, mortality, and MPUE.

DISCUSSION

Our experimental comparisons showed that the hook shape and bait type—both considered as effective bycatch mitigation measures for sea turtles—have extremely multifaceted effects for teleost fishes and sharks and, in some species, the direction of the effects was conflicted. The results provide significant insight into two aspects of the management of vulnerable bycatch species in tuna fisheries: how the confrontational effect of bycatch mitigation measures should be managed, and in which processes of fishing mortality intervention in the management of vulnerable species should occur. As a specific concern regarding the former, when considering shortfin mako shark, which are experiencing significant stock depletion in the North Atlantic (Sims et al. 2018; ICCAT 2019), the implementation of bycatch mitigation measures for sea turtles, which are also required to reduce fishery bycatch, could conversely increase fishing mortality and become a conservation risk. Previously, most of bycatch mitigation measure assessments have focused on whether they reduce the impact of vulnerable bycatch species of concern being bycaught, with the secondary impact being from an economic perspective—in other words, whether the catch rate of commercial species is reduced or not. Here we show for the first time that widely adopted bycatch mitigation measures may have a negative impact on some endangered species. Where both species with opposing effects are at low abundance, management measures should be based on a thorough discussion identifying the optimal combination of mitigation measures, accompanied by scientific evidence. With regard to the latter issue, the mortality reduction expected from circle hooks is not very promising, especially for species with high catch mortality, since the main effect of circle hooks is to minimize internal organ damage, which is of little use for species that have died from other causes, such as heat stress or suffocation. For such species, consideration of methods to reduce the catch itself rather than the mortality rate will be of greater benefit.

Because the sea turtle bycatch events in this study were highly skewed toward some terminal gear combinations, we were unable to make many statistical comparisons regarding haulback mortality rate, CPUE, or MPUE, while the size of circle hooks was positively related to increased mouth hooking. This result is consistent with existing studies, which show that the circle hook prevents internal organ damage and improves the probability of live release (Cooke & Suski 2004). Also, although we were unable to make statistical comparisons, CPUE was low, and some combinations did not cause bycatch at all. This result is also consistent with existing studies, and is related to the lower attractant effect of whole

fish bait on marine turtles and the increased probability of swallowing caused by the difficulty to bite off the

bait (Stokes et al. 2011; Parga et al. 2015).

The results of MODEL 1 indicate that circle hook use, rather than tuna hook use; clearly reduced

hook swallowing for blue sharks, shortfin mako sharks, and swordfish, and in these three species,

haulback mortality rate after mouth hooking was also lower than that after hook swallowing. Hook

swallowing has been reported to increase the likelihood of fatal damage to internal organs. Although

previous studies have reported that studies to attach satellite tags to white marlin *Tetrapturus albidus*

caught by recreational fishing using circle hooks and subsequently released have reduced the

post-release mortality rate (Horodysky & Graves 2005), unfortunately, the present study did not

corroborate this information. Since no data regarding post-release mortality were collected in our

research, evaluating the overall impact of hook shape and bait type on bycatch mortality was not possible.

Our results showed different effects of hooking location by hook shapes, with more frequent mouth

hooking by small circle hooks in many species. Few studies have examined the relationship between

hook size and hook hooking position, on the one side, and mortality of non-turtle species by circle hook,

on the other, the possibility that relative differences in mouth and hook size, and differences in feeding

behavior toward prey (swallowing the prey whole or biting it off) may affect the hooking position. Blue

shark studies have reported the negative and long-lasting effects of gut hooking (Epperly et al. 2012;

Gilman et al. 2020).

The results of MODEL 2, which examined the effects of hooks and bait, showed increases in

MPUE due to the use of small-C hooks of all species except striped marlin. This result suggests that the

increased CPUE by small-C hook contributed more to the increase in the MPUE than haulback mortality

rate. Interestingly, we did not observe an increased MPUE with large-C hooks. There have been many

previous findings on the effects of circle hook use on CPUE, with elevated CPUE for tunas and no

consistent trend for sharks and other teleosts. However, although few previous studies have focused on

hook size and made comparisons, catch rates for skipjack, shortbill spearfish, escolar, and lancetfish are

reported to have decreased when larger hooks were used (Curran & Beverly 2012). Considering the

effect of hook size in terms of the catch process, it is unlikely that catch rates increased due to swallowing,

as the results of MODEL 1 indicate an increase in mouth hooking for many species. The results of

MODEL 2 showed that the effects of the bait type varied by species, unlike the pattern for hook types, with

the use of whole fish bait decreasing CPUE and MPUE for blue sharks, while we observed the exact

opposite effect for shortfin mako sharks, in that fish bait increased CPUE and MPUE—and even haulback

mortality rate. Whole fish bait also reduced MPUE in bigeye tuna, while in common dolphinfish and

escolar, it tended to increase MPUE and CPUE. We observed few effects of bait type in billfishes.

Although few studies have examined the effects of bait type alone, an increase in catch rate by fish bait

has been noted for shortfin mako sharks (Foster et al. 2012). In swordfish, studies evaluating the effect of

switching to whole fish bait from squid bait have reported conflicting effects (increase: Santos et al. 2012;

decrease: Foster et al. 2012). Very few studies have assessed haulback mortality by bait types other than

those on sea turtles, but in blue sharks, the use of fish bait reduced haulback mortality (Epperly et al.

2012). The catch rates of the target species, as previously noted, were affected by bait texture, but a

mechanistic explanation for mortality effects is lacking. Since the likelihood that differences in feeding

behavior among species have an effect is high, this issue could be resolved through comparative studies

based on observations of feeding behavior, as in the case of circle hooks.

When we considered the effects of hook and bait type simultaneously, bait types were shown to

have a relatively greater impact than hook types. For example, in shortfin mako sharks and dolphinfish,

the mortality-increasing effect of fish bait was further amplified by that of the circle hook, while in blue

sharks the mortality-increasing effect of the circle hook was counteracted by the bait effect. Although the

combination of circle hooks and fish bait is considered to be very useful for avoiding sea turtle bycatch,

this very combination may pose a high mortality risk for some species, and even in the case of target

fishes, may counteract the expected positive effect of the circle hook on catch rate. Although very limited

studies have simultaneously examined the interrelationship between hook and bait types, all studies

support the conclusion that the combination of hook type and bait type causes fluctuations in catch rates

and that the direction of response varies among species (Coelho et al. 2012; Foster et al. 2012;

Fernandez-Carvalho et al. 2015).

Water temperature and soak time emerged as significant factors affecting haulback mortality

rate. This indicates the significance of these environmental factors, and the need to control them

statistically or experimentally when assessing the effects of hook and bait type on mortality rate. The

effect of water temperature—particularly during the depth and time of day when hooked—and changes in

water temperature up to the time the fish is landed, are considered to be influential. In addition, in high

water temperature environments, studies have identified an increased risk of suffocation due to

decreased dissolved oxygen in water and increased physiological metabolic rate (Skomal & Bernal 2010);

Gallagher et al. (2014) reported an increase in haulback mortality rate for four shark species when caught

during high water temperatures. In addition, for the species that adopt rum ventilation, prolonged soak

time inevitably increases the risk of suffocation due to the restriction of swimming behavior by being

hooked. Mortality rates of tuna, swordfish, and sharks reportedly increased with increasing soak time

(Epperly et al. 2012; Gallagher et al. 2014).

Here, we quantified our data through experimental operations that standardized the various

conditions, but not all aspects were completely controlled. For example, while previous studies on hook

size have examined the correspondence with actual measurements (Gilman et al. 2016), several shapes

of circle hook were used in the experiment in this study, precluding examination of effects of individual

hook types due to sample size issues. We were also unable to examine hooking location of the catch

when fish bait was used. These omissions, while having a limited impact on the present conclusions, are

probably variables that should be considered for a deeper examination of the effects of terminal gear on

catch and bycatch. In this experiment, wire leaders were used on all branchlines to minimize the effects of

sharks' bite-off. While some studies have described concerns that wire leaders may increase catch rates,

especially for rare sharks, they are considered essential for at least experimentally verifying accurate

catch and mortality rates for shark species. We know from this and previous studies that haulback

mortality rates for sharks are much lower than those for teleosts (Reinhardt et al. 2018), and the

implementation of appropriate releases, even with wire leaders, allow for the reduction of risk for

vulnerable shark species.

Here, based on a Bayesian approach, we succeeded in presenting a quantitative impact

assessment of terminal gear on teleosts, sharks, and sea turtles by directly calculating the expected

values for mortality rate, CPUE, and MPUE with each terminal gear. Calculating MPUE using this model

can be a very useful tool because it provides a more direct estimate than does CPUE or mortality rate

alone of catch/bycatch risk to populations of those species. Although we did not include post-release

mortality rate in the model due to lack of data, it would be possible to estimate overall fishing mortality in

the model by designing additional experiments so that mark–recapture is conducted at the same time.

Even if it is not possible to use wire leaders for the proportion of "cryptic catch" due to bite-off, it is

possible to extrapolate this proportion into the model to make predictions regarding mortality—a

development we anticipate.

In this study, we successfully conducted a quantitative impact assessment of bycatch integrating

catch rate and mortality using a Bayesian approach that directly predicts mortality and CPUE. Although

the data used in the analysis relied solely on the results of an Asian-style longline experiment in the

Pacific Ocean and may therefore contain inherent biases, the same analysis method can be used in

conjunction with data from other experiments conducted in other areas and fishing styles to provide a

more integrated assessment.

ACKNOWLEDGMENT

We greatly appreciate for the crew of the R/V Taikei No. 2, especially Fishmaster O. Asano, who was very

helpful in coordinating the research operations, and members of the Fisheries Resources Institute

(formerly known as the National Research Institute of Far Seas Fisheries), especially Y. Inoue, Y. Semba,

A. Kurashima, Y. Fujinami, N. Katsumata, K. Yokota, K. Oshima, H. Minami, who also assisted with the

research and analysis, and Y. Takeuchi who assisted model running using the computer cluster. The

authors would like to than Enago (www.enago.jp) for English language review.

DECLARATION OF INTEREST STATEMENT

REFERENCES

Afonso AS, Hazin FHV, Carvalho F, Pacheco JC, Hazin H, Kerstetter DW, Murie D, Burgess GH. 2011. Fishing

gear modifications to reduce elasmobranch mortality in pelagic and bottom longline fisheries off

Northeast Brazil. Fisheries Research **108**:336–343.

Arnold JB. 2021. ggthemes: Extra Themes, Scales and Geoms for "ggplot2." Available from

https://cran.r-project.org/package=ggthemes.

Brunson JC. 2020. {*ggalluvial*}: Layered grammar for alluvial plots. Journal of Open Source Software **5**:2017.

Clarke S, Sato M, Small C, Sullivan B, Inoue Y, Ochi D. 2015. Bycatch in longline fisheries for tuna and

tuna-like species; A global review of status and mitigation measures. FAO FISHERIES AND

AQUACULTURE TECHNICAL PAPER **588**. Available from www.fao.org/3/a-i4017e.pdf.

Coelho R, Santos MN, Amorim S. 2012. Effects of hook and bait on targeted and bycatch fishes in an

equatorial Atlantic pelagic longline fishery. Bulletin of Marine Science **88**:449–467.

Cooke SJ, Suski CD. 2004. Are circle hooks an effective tool for conserving marine and freshwater recreational

catch-and-release fisheries? Aquatic Conservation: Marine and Freshwater Ecosystems **14**:299–326.

Curran D, Beverly S. 2012. Effects of 16/0 circle hooks on pelagic fish catches in three south pacific albacore

longline fisheries. Bulletin of Marine Science **88**:485–497.

Dias MP, Martin R, Pearmain EJ, Burfield IJ, Small C, Phillips RA, Yates O, Lascelles B, Borboroglu PG,

Croxall JP. 2019. Threats to seabirds: A global assessment. Biological Conservation **237**:525–537.

Diaz GA. 2008. The Effect of Circle Hooks and Straight (J) Hooks on the catch rates and numbers of white

marlin and blue marlin released alive by the U.S. pelagic longline fleet in the Gulf of Mexico. North

American Journal of Fisheries Management **28**:500–506.

Epperly SP, Watson JW, Foster DG, Shah AK. 2012. Anatomical hooking location and condition of animals

captured with pelagic longlines: The grand banks experiments 2002-2003. Bulletin of Marine Science

88:513–527.

Fernandez-Carvalho J, Coelho R, Santos MN, Amorim S. 2015. Effects of hook and bait in a tropical northeast

Atlantic pelagic longline fishery: Part II—Target, bycatch and discard fishes. Fisheries Research

164:312–321.

Foster DG, Epperly SP, Shah AK, Watson JW. 2012. Evaluation of hook and bait type on the catch rates in the

western North Atlantic Ocean pelagic longline fishery. Bulletin of Marine Science **88**:529–545.

Gabry J, Češnovar R. 2021. *cmdstanr*: R Interface to "CmdStan."

Gallagher AJ, Orbesen ES, Hammerschlag N, Serafy JE. 2014. Vulnerability of oceanic sharks as pelagic

longline bycatch. Global Ecology and Conservation **1**:50–59. Elsevier.

Gilman E, Chaloupka M, Bach P, Fennell H, Hall M, Musyl M, Piovano S, Poisson F, Song L. 2020. Effect of

pelagic longline bait type on species selectivity: a global synthesis of evidence. Reviews in Fish Biology

and Fisheries **30**:535–551.

Gilman E, Chaloupka M, Swimmer Y, Piovano S. 2016. A cross-taxa assessment of pelagic longline by-catch

mitigation measures: conflicts and mutual benefits to elasmobranchs. Fish and Fisheries **17**:748–784.

Gilman E, Zollett E, Beverly S, Nakano H, Davis K, Shiode D, Dalzell P, Kinan I. 2006. Reducing sea turtle

by-catch in pelagic longline fisheries. Fish and Fisheries **7**:2–23.

Hall MA, Alverson DL, Metuzals KI. 2000. By-catch: problems and solutions. Marine Pollution Bulletin

41:204–219.

Hiraoka Y, Kanaiwa M, Ohshimo S, Takahashi N, Kai M, Yokawa K. 2016. Relative abundance trend of the

blue shark *Prionace glauca* based on Japanese distant-water and offshore longliner activity in the North

Pacific. Fisheries Science **82**:687–699.

Horodysky AZ, Graves JE. 2005. Application of pop-up satellite archival tag technology to estimate postrelease

survival of white marlin (*Tetrapturus albidus*) caught on circle and straight-shank ("J") hooks in the

western North Atlantic recreational fishery. Fishery Bulletin **103**:84.

ICCAT. 2019. Report of the 2019 shortfin mako shark stock assessment update meeting. Collective Volume of

Scientific Papers, ICCAT **76**:1–77.

Kay M. 2021. {*tidybayes*}: Tidy Data and Geoms for {*Bayesian*} Models. Available from

http://mjskay.github.io/tidybayes/.

Kerstetter DW, Graves JE. 2006. Effects of circle versus J-style hooks on target and non-target species in a

pelagic longline fishery. Fisheries Research **80**:239–250.

Kiyota M, Yokota K, Nobetsu T, Minami H, Nakano H. 2004. Assessment of mitigation measures to reduce

interactions between sea turtles and longline fishery. Pages 24–29. Proceedings of the International

Symposium on SEASTAR 2000 and Bio-logging Science (The 5th SEASTAR 2000 Workshop).

Kruschke JK. 2015. Doing Bayesian data analysis: A tutorial with R, JAGS, and Stan. Second eds. Academic

Press, New York.

Maunder MN, Punt AE. 2004. Standardizing catch and effort data: a review of recent approaches. Fisheries

Research **70**:141–159.

Melvin EF, Guy TJ, Read LB. 2014. Best practice seabird bycatch mitigation for pelagic longline fisheries

targeting tuna and related species. Fisheries Research **149**:5–18.

Pacheco JC, Kerstetter DW, Hazin FH, Hazin H, Segundo RSSL, Graves JE, Carvalho F, Travassos PE. 2011.

A comparison of circle hook and J hook performance in a western equatorial Atlantic Ocean pelagic

longline fishery. Fisheries Research **107**:39–45.

Parga ML, Pons M, Andraka S, Rendón L, Mituhasi T, Hall M, Pacheco L, Segura A, Osmond M, Vogel N.

2015. Hooking locations in sea turtles incidentally captured by artisanal longline fisheries in the Eastern

Pacific Ocean. Fisheries Research **164**:231–237.

Pebesma E. 2018. Simple features for R: Standardized support for spatial vector data. The R Journal

10:439–446.

R Core Team. 2021. R: A Language and environment for statistical computing. Vienna, Austria. Available from

https://www.r-project.org/.

Reinhardt JF, Weaver J, Latham PJ, Dell'Apa A, Serafy JE, Browder JA, Christman M, Foster DG, Blankinship

DR. 2018. Catch rate and at-vessel mortality of circle hooks versus J-hooks in pelagic longline fisheries:

A global meta-analysis. Fish and Fisheries **19**:413–430.

Richard A, Becker OS, Brownrigg, R, ARWR. 2018. *Mapdata*: Extra map databases. Available from

https://cran.r-project.org/package=mapdata.

Richard A, Becker OS, Brownrigg R, Minka TP, ARWR, Deckmyn. A. 2021. *Maps*: Draw geographical maps.

Available from https://cran.r-project.org/package=maps.

Santos MN, Coelho R, Fernandez-Carvalho J, Amorim S. 2012. Effects of hook and bait on sea turtle catches

in an equatorial Atlantic pelagic longline fishery. Bulletin of Marine Science **88**:683–701.

Sims DW, Mucientes G, Queiroz N. 2018. Shortfin mako sharks threatened by inaction. Science **359**:1342.

Skomal G, Bernal D. 2010. Physiological Responses to Stress in Sharks. Pages 475–506 in Carrier JC, Musick

JA, and Heithaus MR (eds) Sharks and Their Relatives II. CRC Press, Boca Raton.

Stokes L, Hataway D, Epperly S, Shah A, Bergmann C, Watson J, Higgins B. 2011. Hook ingestion rates in

loggerhead sea turtles *Caretta caretta* as a function of animal size, hook size, and bait. Endangered

Species Research **14**:1–11.

Swimmer Y, Gutierrez A, Bigelow K, Barceló C, Schroeder B, Keene K, Shattenkirk K, Foster DG. 2017. Sea

turtle bycatch mitigation in U.S. longline fisheries. Frontiers in Marine Science **4**:260.

Stan Development Team. 2021. Stan modeling language users guide and reference manual, 2.28.2. Available

from https://mc-stan.org.

Wallace BP, Kot CY, Dimatteo AD, Lee T, Crowder LB, Lewison RL. 2013. Impacts of fisheries bycatch on

marine turtle populations worldwide: toward conservation and research priorities. Ecosphere **4**:1–49.

Watson JW, Epperly SP, Shah AK, Foster DG. 2005. Fishing methods to reduce sea turtle mortality associated

with pelagic longlines. Canadian Journal of Fisheries and Aquatic Sciences **62**:965–981.

Wickham H et al. 2019. Welcome to the {*tidyverse*}. Journal of Open Source Software **4**:1686.

Yokota K, Kiyota M, Minami H. 2006a. Shark catch in a pelagic longline fishery: Comparison of circle and tuna

hooks. Fisheries Research **81**:337–341.

Yokota K, Kiyota M, Okamura H. 2009. Effect of bait species and color on sea turtle bycatch and fish catch in a

pelagic longline fishery. Fisheries Research **97**:53–58.

Yokota K, Minami H, Kiyota M. 2006b. Measurement-points examination of circle hooks for pelagic longline

fishery to evaluate effects of hook design. Bulletin of Fishery Research Agency **17**:83–102.

Yokota K, Minami H, Kiyota M. 2011. Effectiveness of tori-lines for further reduction of incidental catch of

seabirds in pelagic longline fisheries. Fisheries Science **77**:479–485.

Table 1 List of hook types used in the experiment

hook name	size	category
Komatu Keisaku tunahook	4.0 sun	tuna
Doitomi Tunamutsu	4.0 sun	small-C (circle hook)
Komatsu Keisaku modified	4.0 sun	small-C (circle hook)
Komatsu Keisaku modified	4.5 sun	small-C (circle hook)
Komatsu Keisaku modified	4.8 sun	small-C (circle hook)
Komatsu Keisaku type Etsuna	4.5 sun	small-C (circle hook)
Komatsu Keisaku type North America	4.3 sun	small-C (circle hook)
Tankichi Uruwa	3.8 sun	small-C (circle hook)
Komatsu Keisaku modified	5.2 sun	large-C (circle hook)
Komatsu Keisaku type North America	5.2 sun	large-C (circle hook)
Pacific Fishing Tackle circle hook	18/0	large-C (circle hook)

521   Table 2 Fishing effort (longline hooks) in experimental operations used in the analysis.

hook type	bait type		total effort
	squid	fish	
tuna	97,834	71,146	168,980
small-C	70,882	7,658	78,540
large-C	31,872	6,976	38,848
total effort	200,588	85,780	286,368

Table 3 Number of fish caught by species in the experimental operation

species	total catch
blue shark	13,018
longnose lancetfish	1,297
common dolphinfish	505
shortfin mako shark	485
swordfish	288
bigeye tuna	269
loggerhead turtle	268
escolar	163
striped marine	145
other species	1,578

Table 4 Number of catch by hook and bait type.

species	hook type			bait type	
	tuna	small-C	large-C	squid	fish
blue shark	8,084	3,562	1,372	8,903	4,115
longnose lancetfish	945	257	95	692	605
common dolfinfish	206	181	118	363	142
shortfin mako shark	298	134	53	262	223
swordfish	129	112	47	249	39
bigeye tuna	114	146	9	201	68
loggerhead turtle	128	113	27	259	9
escolar	79	56	28	108	55
striped marine	70	53	22	126	19

Table 5 Composition of hooking location and fate at hauling.

species	hooking location				fate at haulback		
	swallowed	mouth	body	unknown(*)	alive	dead	unknown(*)
blue shark	2,270	2,608	66	8,074	11,701	1,066	251
longnose lancetfish	25	288	16	968	165	1,010	122
common dolphinfish	39	159	4	303	398	87	20
shortfin mako shark	66	68	18	333	363	118	4
swordfish	59	118	18	93	55	226	7
bigeye tuna	7	128	2	132	83	183	3
loggerhead turtle	109	116	14	29	256	5	7
escolar	1	62	1	99	108	40	15
striped marine	12	80	8	45	68	76	1

(*) Includes catches that dropped off before reseachers checked or lack of survey.

Table 6 Estimates of the posterior distribution of the proportion of hooking location by hook when squid bait is used (median). Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses.

species	hook type		
	tuna	small-C	large-C
hooking place: body			
blue shark	0.020 (0.015 - 0.027)	0.008 (0.005 - 0.012)	0.005 (0.001 - 0.014)
loggerhead turtle	0.025 (0.006 - 0.066)	0.079 (0.037 - 0.143)	0.031 (0.001 - 0.154)
longnose lancetfish	0.049 (0.020 - 0.092)	0.031 (0.011 - 0.067)	0.074 (0.018 - 0.185)
shortfin mako	0.102 (0.041 - 0.199)	0.107 (0.052 - 0.185)	0.136 (0.021 - 0.381)
striped marine	0.083 (0.026 - 0.186)	0.068 (0.016 - 0.173)	0.048 (0.002 - 0.235)
swordfish	0.075 (0.030 - 0.148)	0.109 (0.057 - 0.184)	0.031 (0.001 - 0.154)
hooking place: mouth			
blue shark	0.497 (0.475 - 0.518)	0.548 (0.528 - 0.568)	0.560 (0.517 - 0.603)
loggerhead turtle	0.439 (0.346 - 0.534)	0.480 (0.381 - 0.577)	0.746 (0.544 - 0.892)
longnose lancetfish	0.857 (0.792 - 0.908)	0.903 (0.848 - 0.943)	0.872 (0.740 - 0.953)
shortfin mako	0.356 (0.238 - 0.486)	0.524 (0.418 - 0.630)	0.379 (0.154 - 0.650)
striped marine	0.759 (0.623 - 0.869)	0.831 (0.696 - 0.924)	0.883 (0.660 - 0.982)
swordfish	0.528 (0.416 - 0.637)	0.676 (0.574 - 0.767)	0.611 (0.405 - 0.790)
hooking place: swallowed			
blue shark	0.483 (0.461 - 0.505)	0.444 (0.424 - 0.464)	0.434 (0.391 - 0.478)
loggerhead turtle	0.533 (0.438 - 0.626)	0.438 (0.343 - 0.537)	0.210 (0.078 - 0.405)
longnose lancetfish	0.092 (0.052 - 0.148)	0.064 (0.032 - 0.111)	0.046 (0.007 - 0.145)
shortfin mako	0.536 (0.405 - 0.665)	0.365 (0.266 - 0.472)	0.460 (0.213 - 0.724)
striped marine	0.150 (0.066 - 0.273)	0.093 (0.030 - 0.207)	0.048 (0.002 - 0.234)
swordfish	0.394 (0.289 - 0.505)	0.211 (0.136 - 0.303)	0.343 (0.174 - 0.550)

Table 7 Estimated haulback mortality rates (median of posterior distribution) by hooking location when squid bait is used. Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses. Loggerhead turtles were excluded because there were no mortalities and the calculation had not been converged.

species	body	mouth	swallowed
blue shark	0.295 (0.185 - 0.411)	0.052 (0.044 - 0.061)	0.157 (0.141 - 0.172)
shortfin mako shark	0.496 (0.262 - 0.729)	0.134 (0.061 - 0.219)	0.279 (0.176 - 0.394)
longnose lancetfish	0.956 (0.810 - 1.000)	0.860 (0.816 - 0.899)	0.798 (0.621 - 0.934)
swordfish	0.802 (0.595 - 0.959)	0.776 (0.694 - 0.853)	0.919 (0.842 - 0.979)
striped marine	0.798 (0.488 - 0.990)	0.431 (0.315 - 0.547)	0.968 (0.839 - 1.000)

Table 8 Haulback mortality rate by hook and bait type (median of posterior distribution). Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses.

species	hook type		
	tuna	small-C	large-C
bait type : squid			
bigeye tuna	0.436 (0.036 - 1.231)	1.033 (0.100 - 2.938)	0.896 (0.067 - 2.980)
blue shark	44.457 (35.464 - 54.243)	66.395 (53.022 - 81.268)	56.701 (45.075 - 69.726)
common dolphinfish	0.709 (0.336 - 1.182)	1.676 (0.775 - 2.811)	1.193 (0.534 - 2.020)
escolar	0.213 (0.074 - 0.408)	0.630 (0.225 - 1.216)	0.353 (0.108 - 0.724)
longnose lancetfish	3.326 (2.742 - 3.993)	4.677 (3.757 - 5.683)	4.109 (3.074 - 5.261)
shortfin mako	1.051 (0.686 - 1.458)	1.723 (1.084 - 2.431)	1.547 (0.939 - 2.364)
striped marine	0.320 (0.169 - 0.512)	0.369 (0.181 - 0.603)	0.356 (0.155 - 0.640)
swordfish	0.731 (0.496 - 1.004)	1.079 (0.706 - 1.504)	1.115 (0.673 - 1.661)
bait type : fish			
bigeye tuna	0.185 (0.014 - 0.550)	0.441 (0.050 - 1.323)	0.383 (0.016 - 1.342)
blue shark	37.299 (29.425 - 45.517)	55.736 (44.428 - 68.588)	47.620 (37.760 - 58.788)
common dolphinfish	1.781 (0.858 - 3.031)	4.210 (2.019 - 7.284)	3.000 (1.404 - 5.264)
escolar	0.476 (0.153 - 0.941)	1.410 (0.443 - 2.826)	0.787 (0.221 - 1.712)
longnose lancetfish	3.544 (2.849 - 4.346)	4.982 (3.904 - 6.216)	4.379 (3.172 - 5.761)
shortfin mako	2.327 (1.518 - 3.322)	3.813 (2.453 - 5.622)	3.418 (1.913 - 5.221)
striped marine	0.371 (0.163 - 0.663)	0.428 (0.170 - 0.777)	0.411 (0.144 - 0.822)
swordfish	0.751 (0.456 - 1.116)	1.107 (0.636 - 1.672)	1.142 (0.618 - 1.848)

Table 9 Standardized catch per unit effort (CPUE) by hook and bait type (median of posterior distribution). Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses.

species	hook type		
	tuna	small-C	large-C
bait type : squid			
bigeye tuna	0.734 (0.638 - 0.823)	0.757 (0.679 - 0.830)	0.960 (0.812 - 1.000)
blue shark	0.084 (0.076 - 0.092)	0.081 (0.072 - 0.091)	0.077 (0.062 - 0.093)
common dolphinfish	0.154 (0.097 - 0.214)	0.128 (0.081 - 0.180)	0.180 (0.107 - 0.262)
escolar	0.189 (0.085 - 0.311)	0.300 (0.163 - 0.451)	0.181 (0.058 - 0.336)
longnose lancetfish	0.901 (0.869 - 0.929)	0.925 (0.893 - 0.954)	0.890 (0.827 - 0.941)
shortfin mako	0.161 (0.107 - 0.223)	0.186 (0.120 - 0.253)	0.181 (0.084 - 0.300)
striped marine	0.531 (0.401 - 0.665)	0.472 (0.334 - 0.610)	0.530 (0.321 - 0.740)
swordfish	0.829 (0.755 - 0.894)	0.783 (0.700 - 0.859)	0.832 (0.711 - 0.933)
bait type : fish			
bigeye tuna	0.473 (0.326 - 0.627)	0.503 (0.339 - 0.666)	0.886 (0.570 - 1.000)
blue shark	0.078 (0.068 - 0.088)	0.075 (0.062 - 0.088)	0.071 (0.055 - 0.088)
common dolphinfish	0.237 (0.158 - 0.326)	0.200 (0.103 - 0.312)	0.273 (0.164 - 0.398)
escolar	0.242 (0.114 - 0.389)	0.370 (0.174 - 0.586)	0.231 (0.061 - 0.472)
longnose lancetfish	0.909 (0.880 - 0.937)	0.931 (0.895 - 0.960)	0.898 (0.839 - 0.945)
shortfin mako	0.308 (0.242 - 0.387)	0.347 (0.227 - 0.479)	0.342 (0.197 - 0.488)
striped marine	0.691 (0.457 - 0.883)	0.640 (0.368 - 0.865)	0.692 (0.384 - 0.920)
swordfish	0.853 (0.728 - 0.947)	0.811 (0.651 - 0.935)	0.856 (0.700 - 0.967)

Table 10 Estimated MPUE by hook and bait type (median of posterior distribution). Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses.

species	hook type		
	tuna	small-C	large-C
bait type: squid			
bigeye tuna	0.318 (0.026 - 0.902)	0.780 (0.064 - 2.214)	0.841 (0.072 - 2.828)
blue shark	3.744 (2.910 - 4.646)	5.388 (4.152 - 6.790)	4.370 (3.163 - 5.756)
common dolphinfish	0.108 (0.043 - 0.196)	0.213 (0.091 - 0.399)	0.214 (0.080 - 0.405)
escolar	0.040 (0.009 - 0.089)	0.187 (0.052 - 0.403)	0.063 (0.009 - 0.162)
longnose lancetfish	2.994 (2.443 - 3.586)	4.321 (3.440 - 5.239)	3.641 (2.706 - 4.706)
shortfin mako	0.169 (0.088 - 0.263)	0.321 (0.156 - 0.498)	0.278 (0.102 - 0.535)
striped marine	0.169 (0.076 - 0.279)	0.173 (0.075 - 0.299)	0.185 (0.065 - 0.360)
swordfish	0.604 (0.403 - 0.838)	0.840 (0.538 - 1.186)	0.919 (0.533 - 1.385)
bait type: fish			
bigeye tuna	0.087 (0.007 - 0.267)	0.219 (0.016 - 0.675)	0.319 (0.016 - 1.158)
blue shark	2.909 (2.229 - 3.684)	4.185 (3.112 - 5.415)	3.396 (2.408 - 4.571)
common dolphinfish	0.420 (0.168 - 0.765)	0.836 (0.288 - 1.664)	0.812 (0.286 - 1.558)
escolar	0.113 (0.026 - 0.261)	0.511 (0.113 - 1.177)	0.178 (0.022 - 0.512)
longnose lancetfish	3.217 (2.582 - 3.953)	4.629 (3.668 - 5.846)	3.919 (2.800 - 5.169)
shortfin mako	0.712 (0.432 - 1.063)	1.311 (0.679 - 2.117)	1.153 (0.527 - 2.046)
striped marine	0.250 (0.089 - 0.469)	0.264 (0.081 - 0.525)	0.273 (0.072 - 0.589)
swordfish	0.633 (0.375 - 0.959)	0.883 (0.466 - 1.364)	0.961 (0.497 - 1.590)

Table 11 Nominal CPUE, Haulback mortality rate and MPUE of Loggerhead turtle。 All figures are based on aggregated operational data, not estimates.

hook type	CPUE		haulback mortality		MPUE	
	squid	fish	squid	fish	squid	fish
tuna	1.247	0.084	0.0385	0.000	0.0480	0.000
large-C	1.034	0.277	0.0090	0.000	0.0093	0.000
small-C	2.148	0.261	0.0246	0.000	0.0528	0.000

Figure 1 Locations where the longline operation experiment was conducted.

Figure 2 Differences in the estimated probability of the "swallowed" hooking position of each circle hook type from tuna hook when squid bait is used. The red dotted line indicates that the difference is zero.

Figure 3　Differences in the estimated haulback mortality rate of each target hooking position from "swallowed" hooking position when squid bait is used. The red dotted line indicates that the difference is zero.

Figure 4 Differences in estimated haulback mortality between each experimental group and the control group (squid x tuna hook). The red dotted line indicates that the difference is zero.

Figure 5 Relationship between sea surface temperature (SST) variability and haulback mortality rate at longline operations. Solid lines indicate median; masked areas indicate 95% Bayesian credible interval.

Figure 6 Relationship between soak time (time from setting the branch line to hauling) variability and haulback mortality rate at longline operations. Solid lines indicate median; masked areas indicate 95% Bayesian credible interval.

Figure 7　Differences in the standardized catch per unit effort (CPUE) for each experimental group from those for the control group (squid x tuna hook). The red dotted line indicates that the difference is zero.

Figure 8   Differences in the estimated MPUE (mortality per unit effort) between those for each experimental group and those for the control group (squid x tuna hook). The red dotted line indicates that the difference is zero.

Figure 9 Alluvial plot of hooking locations and associated haulback mortality rates of loggerhead turtles by hook when squid bait is used.

Appendix 1 Detailed measurements of hooks used in the experiment and a figure explaining measurement points of hooks (copied from Yokota et al. 2006b).

manufacture	Komatsu Keisaku	Hisamatsu Tankichi	Doitomi	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Pacific Fishing Tackle MFG., CO.
hook name	tuna hook	Uruwa hook BKN	tuna circle hook SS-170	modified circle hook	modified circle hook	modified circle hook	cirlce hook type Etsuna	cirlce hook type North America	cirlce hook type North America	modified circle hook	circle hook
standardized size	4.0 sun	3.8 sun	#4	4.0 sun	4.5 sun	4.8 sun	4.5 sun	4.3 sun	5.2 sun	5.2 sun	18/0
material	stainless steel	hard steel	stainless steel	stainless steel	stainless steel	stainless steel	stainless steel	hard steel	hard steel	stainless steel	stainless steel
hook eye	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring
shank thickness(mm)	5.3	4.0	4.1	5.3	5.2	N/A	4.9	5.3	5.7	5.3	5.1
straight total length(mm)	63	56	58	60	63	N/A	62	57	74	76	68
straight total width(mm)	38	44	49	47	49	N/A	56	45	56	54	59
minimum total width(mm)	38	36	39	41	45	N/A	51	41	52	48	51
maximum total width(mm)	63	64	69	68	71	N/A	76	63	81	85	80
front length(mm)	41	33	38	35	44	N/A	47	39	49	47	45
minimum inner width(mm)	27	20	15	24	25	N/A	26	20	27	26	27
L-W ratio	1.7	1.3	1.2	1.3	1.3	N/A	1.1	1.3	1.3	1.4	1.2
max-min ratio	1.7	1.8	1.8	1.7	1.6	N/A	1.5	1.5	1.6	1.8	1.6
incurved point angle		80	70	90	70	N/A	70	65	75	80	80
offset angle	$5° \leqq θ < 10°$	$θ ≈ 0°$	$θ ≈ 0°$	$θ ≈ 10°$	$5° \leqq θ < 10°$	$θ < 10°$	$θ ≈ 10°$	$5° \leqq θ < 10°$	$10° \leqq θ < 15°$	$θ ≈ 5°$	$10° \leqq θ < 15°$
offset width(mm)	0.9	≈ 0	≈ 0	1.8	2.6	N/A	2.1	1.5	3.7	1.0	5.4
weight (g)	19.9	12.2	15.0	19.7	21.6	N/A	21.4	19.4	30.3	25.5	23.2

Appendix 2 Stan code used for MCMC sampling of (a) Model 1 and (b) Model 2

(a)   MODEL 1

```
data {
    int hook_cat;                           // Hook type category
    int<lower=1> N1;                        // Number of hooked individual
    int<lower=1> H_cat;                     // Hooking Hation category
    int<lower=1, upper=H_cat> H[N1];        // Hooking location in the nth individual
    int<lower=1, upper=hook_cat> hook[N1];  // Hook type that captured the nth individual.
    int<lower=0, upper=1> M[N1];            // Fate at haulback for the nth individual.
    int L;                                  // Number of estimation parameters explaining mortality rate
    matrix[N1, L] X1;                       // Parameter-variable matrix for parameter estimation
    // for prediction //
    matrix[H_cat, L] X1_pred;               // Parameter-variable matrix for prediction
    int N3;                                 // Number of data records for prediciton
    matrix[N3, L] X1_pred2;                 // Parameter-variable matrix for prediction of haulback mortality relating to SST and soaktime
}

transformed data{
    vector[hook_cat] Zeros;                 //Intercepts of probability of hooking location for categorical distribution

    Zeros = rep_vector(0, hook_cat);
}

parameters {
    matrix[hook_cat, H_cat - 1] theta_raw;  // Probability of hooking location (before transformation)
    vector[L] beta;                         // Vector of parameters for explanatory variables of haulback mortality rate
}

transformed parameters {
    vector[N1] p_dead;                      // Haulback mortality rate
    matrix[hook_cat, H_cat] theta;          // Probability of hooking location (transformed for categorical distribution)

    p_dead = inv_logit(X1 * beta);
    theta = append_col(Zeros, theta_raw);
}

model {
    for(n in 1:N1){
    M[n] ~ bernoulli(p_dead[n]);
    target += categorical_lpmf(H[n]|softmax(theta[hook[n],]'));
    }
}

generated quantities{
    matrix[hook_cat, H_cat] theta_pred;     // Predicted probability of hooking location
    vector[H_cat] p_dead_pred;              // Predicted haulback mortality rate
    vector[N3] M_pred;                      // Predicted haulback mortality rate relating to SST and soaktime

    for(l in 1:H_cat)
        p_dead_pred[l] = inv_logit(X1_pred[l,] * beta);
    for(k in 1:hook_cat){
        theta_pred[k,] = softmax(theta[k,]')';
    }
    for(l in 1:N3)
    M_pred[l] = inv_logit(X1_pred2[l,] * beta);
}
```

(b)   MODEL 2

```
data {
    // Estimation of haulback mortality rate //
    int hook_bait_comb;                    // Category combining hook and bait type
    int<lower=1> N1;                        // Number of individuals determined to its fate at haulback
    int<lower=0, upper=1> M[N1];           // Fate at haulback for the nth individual.
    int L;                                  // Number of estimation parameters explaining mortality rate
    matrix[N1, L] X1;                       // Parameter-variable matrix for parameter estimation of haulback mortality rate

    // Estimation and standardization of CPUE //
    int N2;                                 // Number of capture records
    int C[N2];                              // Catch number in the nth records
    int L2;                                 // Number of estimation parameters explaining CPUE
    matrix[N2, L2] X2;                      // Parameter-variable matrix for parameter estimation of CPUE
    vector[N2] O;                           // Offset (number of hooks)
    int YRID;                               // Random variable: Year category
    int YR[N2];                             // Year when the nth capture was recorded

    // for prediction //
    matrix[hook_bait_comb, L] X1_pred;     // Parameter-variable matrix for parameter estimation of haulback mortality rate
    matrix[hook_bait_comb, L2] X2_pred;    // Parameter-variable matrix for parameter estimation of CPUE
    int N3;                                 // Number of data records for prediction of MPUE
    matrix[N3, L] X1_pred2;                 // Parameter-variable matrix for parameter estimation of MPUE
}

parameters {
    vector[L] beta;                         // Vector of parameters for explanatory variables of haulback mortality rate
    vector[L2] beta2;                       // Vector of parameters for explanatory variables of CPUE
    vector[YRID] r_year;                    // Parameter of random variable (YEAR)
    real<lower= 0> sigma;                   // Hyperperameter for random variable
}

transformed parameters {
    vector[N2] lambda;                      // CPUE
    vector[N1] p_dead;                      // Haulback mortality rate

    p_dead = inv_logit(X1 * beta);
    for(i in 1:N2)
    lambda[i] = X2[i,] * beta2 + r_year[YR[i]];
}

model {
    sigma ~ student_t(4, 0, 2.5);
    r_year ~ normal(0, sigma);
    for(n in 1:N1){
        M[n] ~ bernoulli(p_dead[n]);
    }
    C ~ poisson(exp(lambda + log(O)));
}

generated quantities{
    vector[hook_bait_comb] p_dead_pred;    // Predicted probability of haulback mortality
    vector[hook_bait_comb] lambda_pred;    // Standardized CPUE
    vector[hook_bait_comb] zeta_pred;      // Predicted MPUE

    for(l in 1:hook_bait_comb){
        p_dead_pred[l] = inv_logit(X1_pred[l,] * beta);
        lambda_pred[l] = exp(X2_pred[l, ] * beta2);
        zeta_pred[l] = lambda_pred[l] * p_dead_pred[l];
    }
}
```

Appendix 3 Size distributions by species for the major species captured in the study, with body size as an index of precaudal length for blue sharks and shortfin mako sharks, straight-line carapace length for loggerhead turtles, eye-to-fork length for striped marine and swordfish, and fork length for all other species.