TITLE:

Multifaceted effects of bycatch mitigation measures on target/non-target species for pelagic longline

fisheries and consideration for bycatch management

AUTHORS:

Daisuke Ochi [#], Kei Okamoto and Shintaro Ueno

AFFILIATION:

Fisheries Resources Institute, Japan Fisheries Research and Education Agency

Number of tables: 10

Number of figures: 9

[#]Corresponding author,

Address: Fisheries Resources Institute, Japan Fisheries Research and Education Agency, 2-12-4,

Fukuura, Kanazawa-ku, Yokohama-city, Kanagawa 2368648, Japan

e-mail: otthii80s@gmail.com

____

SUMMARY

The pelagic longline fishery, in an effort to reduce bycatch of sea turtles, have developed and

deployed fisheries bycatch mitigation techniques such as replacing J/tuna hooks and squid bait

with circle hooks and whole fish bait. However, little emphasis has been placed on the side

effects of bycatch mitigation measures on endangered species other than target bycatch

species. Several previous studies of the side effects have been marred by lack of control for the

covariates. Here, based on long-term data obtained from research cruises by a pelagic longline

vessel, we examined the effects of using circle hooks and whole fish bait to replace squid bait

on the fishing mortality of target and non-target fishes, and also bycatch species. A

quantitative evaluation analysis of our results, based on a Bayesian approach, showed the use

of circle hooks to increase mouth hooking in target and bycatch species, and their size to be

proportional to the magnitude of the effect. While deploying circle hooks did not increase

fishing mortality per unit effort (MPUE) for shortfin mako sharks, combining to whole fish

bait had a significant increase on MPUE. Because the impact of the introduction of bycatch

mitigation measures on species other than the focused bycatch species is non-negligible, a

quantitative assessment of bycatch mitigation-related fishing mortality is critical before

introducing such measures.

Keywords: billfishes, circle hook, finfish bait, fisheries management, sea turtle, sharks, tuna,

longline fishery

INTRODUCTION

Unintentional catch by fisheries is referred to as bycatch, and bycatch of especially endangered species, can have devastating effects on such populations. Thus, efforts to minimize bycatch of endangered species are strongly encouraged at all layers from local to international in terms of species conservation. For example, in the tuna longline fishery, concerns about the increased conservation risk for seabirds and sea turtles by unintentional and fatal catch have been a major issue among many regional fisheries management organizations (RFMOs) since the 1990s (Wallace et al. 2013; Dias et al. 2019). In recent years, some elasmobranchs, whose populations are declining further, are also treated as bycatch species. The decline of these species has focused attention at both the national and the international levels. Even for target fish that are not bycatch species, addressing the deterioration of stock status caused by overfishing requires reductions in unintentional fishing mortality (for instance, billfishes in the North Atlantic; Kerstetter & Graves 2006; Diaz 2008).

Several studies have weighed the sustainability of tuna longline fisheries against the conservation of species vulnerable to bycatch (Hall et al. 2000; Melvin et al. 2014; Clarke et al. 2015)—particularly for seabirds and marine turtles—with the development of several effective bycatch mitigation measures (Melvin et al. 2014; Swimmer et al. 2017). Bycatch mitigation measures in longline fisheries target specific animal groups and are evaluated based on their success in reducing mortality due to bycatch of specific vulnerable species. While the impact on the catch of the target fish is the primary consideration when evaluating bycatch mitigation techniques, few studies have examined the impacts and tradeoffs on species not targeted by mitigation techniques (Pacheco et al. 2011; Gilman et al. 2016). However, the introduction of bycatch mitigation measures requires an assessment of the impact not only on the target bycatch species on the other ecologically related species prior to their introduction (Reinhardt et al. 2018).

Use of circle hook and whole finfish bait are typical sea turtle bycatch mitigation

measures for pelagic longline fisheries (Watson et al. 2005; Gilman et al. 2006; Yokota et al.

2009; Stokes et al. 2011). The tip of the circle hook bends inward, and when a fish or sea turtle

swallows the hooked bait, the circle hook cannot hook inside the digestive tract; instead, as the

hook exits the mouth, a torque causes it to hook through the edge of the mouth. This property

allows for easy hook removal and has been reported to reduce the mortality rate of bycatch sea

turtles on board and after release (Kiyota et al. 2004; Cooke & Suski 2004; Kerstetter &

Graves 2006). Reports of positive effects of circle hooks include those for other species—such

as reduced haulback and post-release mortality in sharks, reduced post-release mortality in

swordfish, and increased catch rates in tuna, the target fish. The use of fish bait instead of

squid bait potentially reduces bycatch (Watson et al. 2005; Yokota et al. 2009, 2011) and

mortality rates (Stokes et al. 2011; Parga et al. 2015) of sea turtles. These sea turtle mitigation

measures, however, exact other costs. Several meta-analytic studies have reported the circle

hook-related reduction of sea turtle mortality rate but increase in billfish and shark catch rates

(Gilman et al. 2016; Reinhardt et al. 2018; Santos et al. 2020). The use of whole fish bait also

reportedly increases the catch of sharks and other species (Foster et al. 2012). However, few

studies have allowed for quantitative evaluation of the effects of these mitigation measures

experimentally on species beyond sea turtles. This limited data concern is due in part to the

reliance on observer data from commercial vessels, and small sample sizes, small comparison

groups, and lack of experimental rigor from research vessels. In addition, the impact

assessment for sharks underestimates catch rates and mortality associated with missed catches

due to "bite-off" branchline (Reinhardt et al. 2018). In addition, many experimental studies

and meta-analyses (e.g., Diaz 2008; Godin et al. 2012; Huang et al. 2016; Pacheco et al. 2011;

Yokota et al. 2006a) only evaluate gear impacts at significance levels without evaluating the

magnitude of the effect. Small significant differences may be judged not to matter much when

assessing overall risk in bycatch species. Studies that controlled for these conditions would

allow for an evaluation of adverse effects without the confounding problems described above.

Also, many studies use catch/bycatch rates (Andraka et al. 2013; Foster et al. 2012; Gilman et

al. 2007; Watson et al. 2005; Yokota et al. 2006a) and mortality rates (at haulback or after

release; Carruthers et al. 2009; Gallagher et al. 2014; Horodysky et al. 2005; Kerstetter et al.

2003) as important impact indicators, without considering irreversible impacts, such as the

number of organisms killed at the time of catch. Additionally, hooking location itself (mouth,

swallow, or external) is believed to have a strong influence on mortality rate, which demands

the estimation of risk under specific hooking conditions, taking causal relationships into

account.

Here, we used data from controlled experiments to analyze these confounding factors

and developed a Bayesian statistical model to evaluate the effects of changing hook and bait

species on fish species other than turtles—particularly, tuna, swordfish, and sharks—and then

verified the contribution of circle hooks and fish bait to mortality rate, catch rate, and fatal

catch rate, respectively. We also discussed the appropriate assessment of bycatch mitigation

measures in fisheries management.

METHOD

Experimental Operations

We analyzed data from the R/V Taikei No. 2 longline research operation conducted in the

Northwest Pacific Ocean between 2002 and 2010—a typical Japanese shallow-setting

operation targeting mainly swordfish and sharks (set depth: shallowest 47.44m±14.22SD,

deepest 72.41m±10.87SD (n=98 ops.); based on the time depth recorder data [SBT500,

Murayama Denki Ltd.]), using four hooks per basket, a wire leader, and a night soaking

(Yokota et al. 2006a). A total of 286,363 hooks from 306 operations (range: 400 – 964 hooks

per operation) were deployed in the experiment (Table 1). The area of operation ranged from

around the Izu Islands in Japan to off the east coast of northeastern Honshu—typical fishing

ground for Japanese shallow-setting longliners (Fig. 1; Hiraoka et al. 2016). We deployed 11

different hooks (Table 2) and we describe hook shapes and other details of these hooks in

Appendix 1 following the measurement method of Yokota et al. (2006b). Because degree of

hook offset has been reported to affect catch and hooking location (Cooke & Suski 2004),

most hooks were <10° but some were nearly 15°. The bait comprised chub mackerel (*Scomber*

japonicus) and Japanese common squid (*Todarodes pacificus*) in the range of 20–30 cm in

fork length or dorsal mantle length. These were frozen and stored, then completely thawed

before being hooked. The sequence of line setting was divided into several experimental

segments, and a different combination of hook and bait type was applied for each segment,

with an alternate order of segments at each operation.

The researcher recorded catch for each operation for all catch/bycatch, and the species

caught, fate of catch (alive or dead), hooking location (mouth, swallow, and external hooking),

time of catch, and float ID. The researchers determined if the catch was alive or dead based on

the movement of the animals and the degree of injuries before being hauled. Float ID was

recorded when a float was dropped during line setting and when it was retrieved to the deck

during line hauling in order to calculate soak time. Hooking locations were recorded for

catches caught using squid bait. At the start of longline operations, the researcher also

collected sea surface temperature with a water thermometer (DS-1; Murayama Denki Ltd.)

equipped on the vessel, which we subsequently used in the analysis.

The experiment was conducted using multiple sizes of circle hooks, which varied

greatly in size. However, those sample sizes were too small to analyze each hook type

separately. For convenience, we have classified the hook shapes based on size (threshold:

straight total length = 68 mm AND maximum total width = 80 mm; approximately equivalent

to 5.0-sun or 18/0). This led to the following three main hook types:

1. control (tuna hook 4.0-sun; tuna)

2. smaller circle hook (smaller than the threshold; small-C)

3. larger circle hook (the threshold or larger; large-C)

In Table 1, we show longline effort separated by bait and hook type, tabulated

according to the above categorization. We selected the following species for analysis: blue

shark *Prionace glauca*, shortfin mako shark *Isurus oxyrinchus*, bigeye tuna *Thunnus obesus*,

swordfish *Xiphias gladius*, striped marlin *Kajikia audax*, common dolphinfish *Coryphaena*

hippurus, escolar *Lepidocybium flavobrunneum* and, longnosed lancetfish *Alepisaurus ferox*,

and loggerhead turtle *Caretta caretta*.

Statistical Analysis

We conducted all analyses using a Bayesian approach to estimate parameters. We adopted

haulback mortality rate, catch per unit effort (CPUE; /1000 hooks), and mortality per unit

effort (MPUE; /1000 hooks) (Afonso et al. 2011) as indices to evaluate the impact of hook and

bait type on fishing mortality within the analyses. We used the following data as inputs to the

model: number of caught (individuals), longline effort (number of hooks), fate at hauling,

hooking location, year and location of operation, water temperature at operation, and soaking

time.

Because of the missing data due to not recording the hooking location when fish bait

was used, we split the analysis to evaluate the impact of hook type and bait type on fishing

mortality into two models. Model 1 evaluated only the effect of hook type based on capture

events with squid bait and assumed that the use of circle hooks would change to more mouth

hooking of each species, resulting in improved mortality rate. Model 2 evaluated both hook

and bait types, and assumed that the combination of the two would result in large fluctuations

in mortality rate, CPUE and MPUE for each species.

We based MODEL 1 on a logit regression using a Bernoulli distribution. We express

the observed hooking location H and haulback mortality rate M in MODEL 1 by the following

equations:

$$M \sim Bernoulli(p_{dead}) \tag{1}$$

$$logit(p_{1,dead}) = \beta_{1,hkloc} + \beta_{1,sst} + \beta_{1,soaktime} \tag{2}$$

$$H \sim Categorical\left(softmax(\theta_{hkloc \cdot hook})\right) \tag{3}$$

where β_1 is the parameter in each explanatory variable (*hkloc:* hooking location, *sst:* sea

surface temperature and *soaktime:* soak time), $p_{1,dead}$ is the expected haulback mortality rate,

and $\theta_{hkloc \cdot hook}$ is the expected probability of hooking location in each hook type.

We structured MODEL 2 to calculate the expected number of mortalities per effort

(MPUE) based on the parameters estimated in both the mortality and CPUE estimation

subsets. In MODEL 2, due to lack of hooking location data, we calculate $p_{2,dead}$ from a

modified equation (2) as follows;

$$logit(p_{2,dead}) = \beta_{2,hook} + \beta_{2,bait} + \beta_{2,sst} + \beta_{2,soaktime} \tag{4}$$

178   where β_2 is the parameter in each explanatory variable (*hook:* hook type, *bait:* bait type).

179   The CPUE subset is based on a log regression using a Poisson distribution as the error

180   structure. We express the number of catches C per operation using the expected CPUE λ as

181   follows:

$$C \sim Poisson(\lambda + \log(E)) \tag{5}$$

$$\lambda = \log\left(\gamma_{hook} + \gamma_{bait} + \gamma_{lat} + \gamma_{sst} + r_{year}\right) \tag{6}$$

$$r_{year} \sim N(0, \sigma^2) \tag{7}$$

182   where E is the longline effort, γ is a parameter in each explanatory variable (*lat:* latitude where

183   the longline set), r_{year} is a random effect of annual fluctuation on CPUE, and σ is a standard

184   deviation.

185   We obtained the expected values of hook-bait-specific MPUE ζ by multiplying CPUE

186   by at-haulback mortality rate as follows:

$$\zeta = \hat{p}_{2,dead} \times \hat{\lambda} \qquad (8)$$

where $\hat{p}_{2,dead}$ denotes the expected mortality rate standardized for hook and bait type, and $\hat{\lambda}$

denotes the estimated CPUE standardized for hook and bait type. We used the standardization

method for abundance indices used in fisheries stock assessment (Maunder & Punt 2004). In

this method, explanatory variables other than the factor subject to standardization are averaged

to predict the objective variable, which in stock assessment is a time scale such as years or

months, but in our case, we modified the standardization scale to reference hook type and bait

type.

We calculated each parameter based on a Bayesian approach with Markov chain

Monte Carlo (MCMC) sampling. For the MCMC sampling, we used cmdstan 2.28.2 (Stan

Development Team 2021). As a prior distribution for σ, we used a half student-t distribution

with 2 degrees of freedom, mean 0, and variance 2.5, and we used a uniform distribution for

the other parameters ($\beta_1, \beta_2, \gamma, \theta$). We computed the posterior distribution using Stan with

15000 sampling iterations including 10000 warmup iterations, number of chains as 4, and no

sinning. We calculated Bayesian credible intervals based on the highest density interval (HDI)

for the estimates. Although the Bayesian approach for the estimates precluded significance

testing, we determined the difference between the estimates of the experimental group and

those of the control group ("swallowing" for hooking location, "tuna" hook for hook type, and

"squid" for bait type), and if the lower and upper limits of HDI for the difference did not

exceed 0, we considered the difference as a difference for convenience (assuming region of

practical equivalence [ROPE] as 0; Kruschke 2015). In Appendix 2 we show the Stan code

used to estimate each parameter in MODEL 1 and 2. For other data handling, statistical

analysis, and plotting, we used R4.3.0 (R Core Team 2023) and packages "cmdstanr 0.6.1,"

"ggalluvial 0.12.5," "ggthemes 4.2.4," "mapdata 2.3.1," "maps 3.4.1," "sf 1.0–14," "tidybayes

3.0.6," and "tidyverse 2.0.0" (Pebesma 2018; Wickham et al. 2019; Becker et al. 2020;

Brunson 2020; Arnold 2021; Gabry & Češnovar 2021; Becker et al. 2022; Kay 2023).

RESULTS

Summary Statistics

Sufficient catches/bycatches of blue shark, longnose lancetfish, common dolphinfish, and

shortfin mako shark swordfish, bigeye tuna, loggerhead turtle, escolar and striped marlin were

recorded for the later analysis (Table 3). The main species listed in Table 3 as "other species"

are listed as follows; salmon shark *Lamna ditropis* (N = 229), pelagic stingray

Pteroplatytrygon violacea (N = 199), pomflets *Brama spp.* (N = 135), bigeye thresher shark

Alopias superciliosus (N = 89), and albacore *Thunnus alalunga* (N = 69). The sample sizes of

these "other species" were too skewed among experimental groups to converge the later

analysis.

In Table 3 we also show the number of fish caught by hook type and bait type. Bigeye

tuna had extremely low catches on large-C hooks, and loggerheads had low catches on fish

bait. The most common hooking location at the time of catch was mouth hooking for all nine

species, with extremely few hook locations other than mouth hooking, especially for bigeye

tuna and escolar (Table 4). The proportion of mortality of captured species at haulback varied

greatly by species. The haulback mortality rate was low for blue sharks, common dolphinfish,

escolar, and shortfin mako sharks and, higher for bigeye tuna, longnose lancetfish, striped

marlin, and swordfish. In the case of loggerhead turtles, the mortality rate was extremely low.

The length-based size composition of the nine species (precaudal length for blue

sharks and shortfin mako sharks, straight-line carapace length for loggerhead turtles, eye-to-

fork length for striped marlin and swordfish, and fork length for all other species) used in the

analysis was not statistically compared in this study and was not included in the model

because it did not contribute to haulback mortality rate, but in Appendix 3 we included

histograms.

Almost all parameters in the models for the nine species were successfully converged

(Rhat < 1.1) but the whole models did not converge for bigeye tuna, common dolphinfish,

escolar in the MODEL 1, and some of the parameters in the MOEDL 1 and the whole model in

the MODEL 2 for loggerhead turtle did not converge, as we describe below.

Output of MODEL 1

In Table 5 we show occurrence estimates of hooking locations by hook type throughout the

model. We observed differences in hooking location with an increase in mouth hooking and a

decrease in hook swallowing for large-C for loggerhead turtle and a clear increase in mouth

hooking for small-C for shortfin mako shark and swordfish (Fig. 2). In blue sharks, the

frequency of hook swallowing decreased in both small-C and large-C.

In Table 6 we show haulback mortality rates by hooking location. We observed clear

differences in haulback mortality rate by hooking location for blue shark, shortfin mako shark,

striped marlin, and swordfish (Fig. 3). Haulback mortality rates after hook swallowing for blue

sharks were lower than those for external hooking, and higher than those for mouth hooking.

We observed lower haulback mortality rates for shortfin mako shark, striped marlin, and

swordfish from hook swallowing than from mouth hooking.

 Output of MODEL 2

Haulback mortality rate was higher for large-C than for tuna hook types in bigeye tuna, but did

not differ among hook types in other species (Table 7, Fig. 4). On the other hand, the mortality

rate for two bait types differed in bigeye tuna and blue shark, with a rate reduction and

increase with the use of fish bait in bigeye tuna and shortfin mako shark, respectively.

In Fig. 5 and Fig. 6 we show the effects of two covariates—sea surface temperature

(SST) and soak time—on haulback mortality rate. The response to SST differed by species,

with haulback mortality rate increasing with higher SST for shortfin mako sharks, swordfish,

common dolphinfish, and escolar, and conversely increasing with lower SST for bigeye tuna

and longnose lancetfish. We observed little fluctuation in haulback mortality rate with SST in

blue shark and striped marlin. In general, haulback mortality rate increased with increasing

soak time. However, for bigeye tuna, haulback mortality rate decreased with soak time, but the

trend was not clear, and for blue sharks, haulback mortality rate increased only slightly with

increased soak time.

We observed higher CPUE for only small-C in blue shark, bigeye tuna, common

dolphinfish and escolar (Fig. 7, Table 8). We observed differences in standardized CPUE by

bait type in bigeye tuna, blue shark, common dolphinfish, escolar, and shortfin mako shark.

CPUE decreased with whole fish bait in bigeye tuna and blue sharks, but increased in common

dolphinfish, escolar, and shortfin mako shark. In the case of blue shark, escolar, and shortfin

mako shark, the bait effect could be varied with circle hooks, suppressing the CPUE-

increasing effect by fish bait in bigeye tuna, and blue shark with the use of circle hooks, while

conversely this combination boosted increase of CPUE in common dolphinfish, escolar, and

shortfin mako shark.

Compared to differences in CPUE among hook and bait type, those in MPUE were

relatively small (Table 9). We observed differences with higher MPUE for only small-C in

bigeye tuna and escolar, (Fig. 8). We confirmed decreases in MPUE by fish bait in bigeye tuna

and blue shark, and conversely, increases in MPUE in common dolphinfish, shortfin mako

shark. The effect of the combination of whole fish bait and circle hook varied in bigeye tuna,

blue shark, common dolphinfish escolar and shortfin mako shark. In bigeye tuna and blue

shark, the effect of the circle hook on MPUE was suppressed by fish bait, while in common

dolphinfish, escolar and shortfin mako shark, MPUE significantly increased when whole fish

bait and circle hook were used together.

Hook and bait effect for sea turtle bycatch

We failed to complete our analysis for loggerhead turtles throughout the models because the

bias in frequency of capture events among the experimental groups was too large and did not

converge except for only a part of MODEL 1. Instead, we show the nominal CPUE, haulback

mortality rate, and MPUE for each experimental group in Table 10, and hooking location and

haulback mortality rate by each hook type with squid bait in Fig. 9. Most individuals survived

regardless of hooking location. When whole fish bait was used, haulback mortality rate and

associated MPUE were zero. For squid bait, large-C had the smallest CPUE, mortality, and

MPUE.

DISCUSSION

Our experimental comparisons showed that the hook and bait type—both considered as

effective bycatch mitigation measures for sea turtles—have extremely multifaceted effects for

teleost fishes and sharks and, in some species, the direction of the effects was conflicted. The

results provide significant insight into two aspects of the management of vulnerable bycatch

species in tuna fisheries: how the confrontational effect of bycatch mitigation measures should

be managed, and in which processes of fishing mortality intervention in the management of

vulnerable species should occur. As a specific concern regarding the former, when considering

shortfin mako shark, which are experiencing significant stock depletion in the North Atlantic

(Sims et al. 2018; ICCAT 2019), the implementation of bycatch mitigation measures for sea

turtles, which are also required to reduce fishery bycatch, could conversely increase fishing

mortality and become a conservation risk. Previously, most of bycatch mitigation measure

assessments have focused on whether they reduce the impact of vulnerable bycatch species of

concern being bycaught, with the secondary impact being from an economic perspective—in

other words, whether the catch rate of commercial species is reduced or not. Where both

loggerhead turtle and shortfin mako with opposing effects are at low abundance, management

measures should be based on a thorough discussion identifying the optimal combination of

mitigation measures, accompanied by scientific evidence. With regard to the latter issue, the

mortality reduction expected from circle hooks is not very promising, especially for species

with high catch mortality, since the main effect of circle hooks is to minimize internal organ

damage, which is of little use for species that have died from other causes, such as heat stress

or suffocation. For such species, consideration of methods to reduce the catch itself rather than

the mortality rate will be of greater benefit.

Increased mouth hooking of loggerhead turtle by large circle hooks is consistent

with existing studies. The circle hook prevents internal organ damage and improves the

probability of live release (Cooke & Suski 2004) while the impact of circle hooks on haulback

mortality in this study could not be evaluated due to skewed data about mortality events. For

the same reason, the MODEL 2 analysis could not evaluate the effects of circle hook and bait

type on CPUE, mortality rate, and MPUE of loggerhead turtle. However, since mortality event

of loggerhead turtle did not occur at all when fish bait was used, it may be assumed that there

is an effect of mortality reduction by using fish bait. This result is also consistent with existing

studies, and is related to the lower attractant effect of whole fish bait on marine turtles and the

increased probability of swallowing caused by the difficulty to bite off the bait (Stokes et al.

2011; Parga et al. 2015).

Our study indicated that reduced hook swallowing with circle hooks and increased

haulback mortality after hook swallowing were found for sharks and swordfish. Hook

swallowing has been reported to increase the likelihood of fatal damage to internal organs.

Although previous studies have reported that studies to attach satellite tags to white marlin

Kajikia albida caught by recreational fishing using circle hooks and subsequently released

have reduced the post-release mortality rate (Horodysky & Graves 2005), unfortunately, the

present study did not corroborate this information. Our results also presented different effects

of hooking location by hook types, with more frequent mouth hooking by small circle hooks in

many species. Few studies have examined the relationship among size of circle hook, hooking

location and haulback mortality of non-turtle species. However, two studies discussed the

possibility that relative differences in mouth and hook size, and differences in feeding

behavior toward prey (swallowing the prey whole or biting it off) may affect the hooking

location (Epperly et al. 2012; Gilman et al. 2020).

An increase in CPUE and MPUE was observed with the use of small-C hooks among

the two types of circle hooks. This indicates that the increase in CPUE due to small-C hooks

had a greater impact on MPUE than the haulback mortality rate. There have been many

previous findings on the effects of circle hook use on CPUE, with elevated CPUE for tunas

and no consistent trend for other teleosts and sharks. Interestingly, we did not observe an

increased CPUE and MPUE with large-C hooks. Although few previous studies have focused

on hook size and made comparisons, catch rates for skipjack, shortbill spearfish, escolar, and

lancetfish are reported to have decreased when larger hooks were used (Curran & Beverly

2012; Gilman et al. 2018). Considering the effect of hook size in terms of the catch process, it

is unlikely that catch rates increased due to swallowing, as the results of MODEL 1 indicate an

increase in mouth hooking for many species. In the case of blue shark and shortfin mako

shark, the increase was observed in CPUE for both small-C, but this increase was not observed

in MPUE. It may be explained by that the effect of the circle hook on MPUE may have been

masked by the uncertainty of haulback mortality rate. Several studies have examined the

effects of fish bait without circle hook and have reported reduced catch rates for tropical tunas,

blue sharks, and escolar and increased catch rates for shortfin mako, porbeagle shark, and

white marlin (Fernandez-Carvalho et al. 2015; Foster et al. 2012; Watson et al. 2005; Yokota et

al. 2009). In swordfish, some previous studies evaluating the effect of switching to whole fish

bait from squid bait have reported conflicting effects (increase: Santos et al. 2012; Foster et al.

2012; decrease: Fernandez-Carvalho et al. 2015). Very few studies have assessed haulback

mortality by bait types other than those on sea turtles. The catch rates of the target species, as

previously noted, were affected by bait texture, but a mechanistic explanation for mortality

effects is lacking. Since the likelihood that differences in feeding behavior among species have

an effect is high, this issue could be resolved through comparative studies based on

observations of feeding behavior, as in the case of circle hooks.

When the effects of hook and bait types were considered simultaneously, it was found

that the bait type had a more significant impact on CPUE, mortality rate, and MPUE than the

hook type. However, whether this impact was beneficial or detrimental varied greatly

depending on the species. Although the combination of circle hooks and fish bait is effective

for avoiding sea turtle bycatch, this combination may pose a high mortality risk for

endangered species like shortfin mako shark, and even in the case of target fishes like bigeye

tuna, may counteract the expected positive effect of the circle hook on catch rate. In the case

of shortfin mako, for example, changing the bait from squid to fish from tuna hooks to small-C

hooks increases MPUE by about 4.0 times, and changing the bait from squid to fish from tuna

hooks to small-C hooks further increases MPUE by about 5.9 times (Table 9), and in the case

of bigeye tuna, the CPUE estimate, which increased by 1.9 times with small-C hooks, returned

to the same level as tuna hooks by changing from squid to fish bait (Table 8). Such substantial

changes in CPUE and MPUE would not be ignored when managing fisheries for those species.

Although very limited studies have simultaneously examined the interrelationship between

hook and bait types, all studies support the conclusion that the combination of hook type and

bait type causes fluctuations in catch rates and that the direction of response varies among

species (Coelho et al. 2012; Foster et al. 2012; Fernandez-Carvalho et al. 2015).

Water temperature and soak time emerged as significant factors affecting haulback

mortality rate, which had been reported in sharks (Carruthers et al. 2009; Gallasher et al. 2014)

and sea turtles (Watson et al. 2005). This indicates these covariates need to be controlled

statistically or experimentally when assessing the effects of hook and bait type on mortality

rate. The effect of water temperature—particularly during the depth and time of day when

hooked—and changes in water temperature up to the time the fish is landed, are considered to

be influential. In addition, in high water temperature environments, studies have identified an

increased risk of suffocation due to decreased dissolved oxygen in water and increased

physiological metabolic rate (Skomal & Bernal 2010). Gallagher et al. (2014) reported an

increase in haulback mortality rate for four shark species when caught during high water

temperatures. In addition, for the species that adopt rum ventilation, prolonged soak time

inevitably increases the risk of suffocation due to the restriction of swimming behavior by

being hooked. Mortality rates of tuna, swordfish, and sharks reportedly increased with

increasing soak time (Epperly et al. 2012; Gallagher et al. 2014).

We quantified our data through experimental operations that standardized the various

conditions, but not all aspects were completely controlled. For example, while previous

studies on hook size have examined the correspondence with actual measurements (Gilman et

al. 2016), several shapes of circle hook were used in the experiment in this study, precluding

examination of effects of individual hook types due to sample size issues. We were also unable

to examine hooking location of the catch when fish bait was used. These omissions, while

having a limited impact on the present conclusions, are probably variables that should be

considered for a deeper examination of the effects of terminal gear on catch and bycatch. In

this experiment, wire leaders were used on all branchlines to minimize the effects of sharks'

bite-off. While some studies have described concerns that wire leaders may increase catch

rates, especially for rare sharks, they are considered essential for at least experimentally

verifying accurate catch and mortality rates for shark species. We know from this and previous

studies that haulback mortality rates for sharks are much lower than those for teleosts (Afonso

et al. 2012; Reinhardt et al. 2018), and the implementation of appropriate releases, even with

wire leaders, allow for the reduction of risk for vulnerable shark species.

Here, based on a Bayesian approach, we succeeded in presenting a quantitative

impact assessment of terminal gear on teleosts, sharks, and sea turtles by directly calculating

the expected values for mortality rate, CPUE, and MPUE with each terminal gear. Calculating

MPUE using this model can be a very useful tool because it provides a more direct estimate

than does CPUE or mortality rate alone of catch/bycatch risk to populations of those species.

Although we did not include post-release mortality rate in the model due to lack of data, it

would be possible to estimate overall fishing mortality in the model by designing additional

experiments so that mark–recapture is conducted at the same time. Even if it is not possible to

use wire leaders for the proportion of "cryptic catch" due to bite-off, it is possible to

extrapolate this proportion into the model to make predictions regarding mortality—a

development we anticipate. Although the data used in the analysis relied solely on the results

of an Asian-style longline experiment in the Pacific Ocean and may therefore contain inherent

biases, the same analysis method can be used in conjunction with data from other experiments

conducted in other areas and fishing styles to provide a more integrated assessment.

REFERENCES

Afonso AS, Hazin FHV, Carvalho F, Pacheco JC, Hazin H, Kerstetter DW, Murie D, Burgess

GH (2011) Fishing gear modifications to reduce elasmobranch mortality in pelagic and

bottom longline fisheries off Northeast Brazil. Fisheries Research **108**:336–343.

Afonso AS, Santiago R, Hazin H, Hazin FHV (2012) Shark bycatch and mortality and hook

bite-offs in pelagic longlines: Interactions between hook types and leader materials. Fish

Res. 131-133, 9–14. https://doi:10.1016/j.fishres.2012.07.001.

Andraka S, Mug M, Hall M, Pons M, Pacheco L, Parrales M, Rendón L, Parga ML, Mituhasi

T, Segura Á, et al. (2013) Circle hooks: Developing better fishing practices in the

artisanal longline fisheries of the Eastern Pacific Ocean. *Biological Conservation* **160**,

214–224. https://doi.org/10.1016/j.biocon.2013.01.019

Arnold JB (2021) ggthemes: Extra Themes, Scales and Geoms for "ggplot2." Available at

https://cran.r-project.org/package=ggthemes

Becker RA, Wilks AR, Brownrigg R (2020) mapdata: Extra Map Databases. Available at

https://cran.r-project.org/package=mapdata

Becker RA, Wilks AR, Brownrigg R, Minka TP, Deckmyn A (2022) maps: Draw

geographical maps. Available at https://cran.r-project.org/package=maps

Brunson JC (2020) ggalluvial: Layered Grammar for Alluvial Plots. *Journal of Open Source*

Software **5**:2017.

Carruthers EH, Schneider DC, Neilson, JD (2009) Estimating the odds of survival and

identifying mitigation opportunities for common bycatch in pelagic longline fisheries.

Biological Conservation **142**, 2620–2630. https://doi.org/10.1016/j.biocon.2009.06.010

Clarke S, Sato M, Small C, Sullivan B, Inoue Y, Ochi D (2015) Bycatch in longline fisheries

for tuna and tuna-like species; A global review of status and mitigation measures. *FAO*

Fisheries and Aquaculture Technical Paper **588**. Available at https://www.fao.org/3/a-
i4017e.pdf.

Coelho R, Santos MN, Amorim S (2012) Effects of hook and bait on targeted and bycatch
fishes in an equatorial Atlantic pelagic longline fishery. *Bulletin of Marine Science* **88**,
449–467.

Cooke SJ, Suski CD (2004) Are circle hooks an effective tool for conserving marine and
freshwater recreational catch-and-release fisheries? *Aquatic Conservation: Marine and*
Freshwater Ecosystems **14**, 299–326.

Curran D, Beverly S (2012) Effects of 16/0 circle hooks on pelagic fish catches in three south
pacific albacore longline fisheries. *Bulletin of Marine Science* **88**, 485–497.
https://doi.org/10.5343/BMS.2011.1060

Dias MP, Martin R, Pearmain EJ, Burfield IJ, Small C, Phillips RA, Yates O, Lascelles B,
Borboroglu PG, Croxall JP (2019) Threats to seabirds: A global assessment. *Biological*
Conservation **237**, 525–537.

Diaz GA (2008) The effect of circle hooks and straight (J) hooks on the catch rates and
numbers of white marlin and blue marlin released alive by the U.S. pelagic longline fleet
in the Gulf of Mexico. *North American Journal of Fisheries Management* **28**, 500–506.
https://doi.org/10.1577/m07-089.1

Epperly SP, Watson JW, Foster DG, Shah AK (2012) Anatomical hooking location and
condition of animals captured with pelagic longlines: The grand banks experiments 2002-
2003. *Bulletin of Marine Science* **88**, 513–527.

Fernandez-Carvalho J, Coelho R, Santos MN, Amorim S (2015) Effects of hook and bait in a
tropical northeast Atlantic pelagic longline fishery: Part II—Target, bycatch and discard
fishes. *Fisheries Research* **164**, 312–321.

Foster DG, Epperly SP, Shah AK, Watson JW (2012) Evaluation of hook and bait type on the

catch rates in the western North Atlantic Ocean pelagic longline fishery. *Bulletin of*

Marine Science **88**, 529–545.

Gabry J, Češnovar R (2021) cmdstanr: R Interface to "CmdStan." Available at https://mc-

stan.org/cmdstanr/reference/cmdstanr-package.html

Gallagher AJ, Orbesen ES, Hammerschlag N, Serafy JE (2014) Vulnerability of oceanic

sharks as pelagic longline bycatch. *Global Ecology and Conservation* **1**, 50–59.

Gilman E, Chaloupka M, Bach P, Fennell H, Hall M, Musyl M, Piovano S, Poisson F, Song L

(2020) Effect of pelagic longline bait type on species selectivity: a global synthesis of

evidence. *Reviews in Fish Biology and Fisheries* **30**, 535–551.

https://doi.org/10.1007/S11160-020-09612-0/TABLES/2

Gilman E, Chaloupka M, Swimmer Y, Piovano S (2016) A cross-taxa assessment of pelagic

longline by-catch mitigation measures: conflicts and mutual benefits to elasmobranchs.

Fish and Fisheries **17**, 748–784. https://doi.org/10.1111/faf.12143

Gilman E, Kobayashi D, Swenarton T, Brothers N, Dalzell P, Kinan-Kelly I (2007) Reducing

sea turtle interactions in the Hawaii-based longline swordfish fishery. *Biological*

Conservation **139**, 19–28. https://doi.org/10.1016/j.biocon.2007.06.002

Gilman E, Zollett E, Beverly S, Nakano H, Davis K, Shiode D, Dalzell P, Kinan I (2006)

Reducing sea turtle by-catch in pelagic longline fisheries. *Fish and Fisheries* **7**, 2–23.

https://doi.org/10.1111/j.1467-2979.2006.00196.x

Godin AC, Carlson JK, Burgener V (2012) The effect of circle hooks on shark catchability

and at-vessel mortality rates in longlines fisheries. *Bulletin of Marine Science* **88**, 469–

483. https://doi.org/10.5343/bms.2011.1054

Hall MA, Alverson DL, Metuzals KI (2000) By-catch: problems and solutions. *Marine*

Pollution Bulletin **41**, 204–219. https://doi.org/10.1016/S0025-326X(00)00111-9

Hiraoka Y, Kanaiwa M, Ohshimo S, Takahashi N, Kai M, Yokawa K (2016) Relative

abundance trend of the blue shark *Prionace glauca* based on Japanese distant-water and

offshore longliner activity in the North Pacific. *Fisheries Science* **82**, 687–699.

https://doi.org/10.1007/s12562-016-1007-7

Horodysky AZ, Graves JE (2005) Application of pop-up satellite archival tag technology to

estimate postrelease survival of white marlin (*Tetrapturus albidus*) caught on circle and

straight-shank ("J") hooks in the western North Atlantic recreational fishery. *Fishery*

Bulletin **103**, 84.

Huang HW, Swimmer Y, Bigelow K, Gutierrez A, Foster DG (2016) Influence of hook type

on catch of commercial and bycatch species in an Atlantic tuna fishery. *Marine Policy*

65, 68–75. https://doi.org/10.1016/j.marpol.2015.12.016

ICCAT (2019) Report of the 2019 Shortfin mako shark stock assessment update meeting.

Collective Volume of Scientific Papers, ICCAT **76**, 1–77.

Kay M (2023) tidybayes: Tidy data and geoms for Bayesian models. Available at

https://doi.org/10.5281/zenodo.1308151

Kerstetter DW, Graves JE. (2006) Effects of circle versus J-style hooks on target and non-

target species in a pelagic longline fishery. *Fisheries Research* **80**, 239–250.

https://doi.org/10.1016/j.fishres.2006.03.032

Kerstetter DW, Luckhurst BE, Prince ED, Graves JE (2003) Use of pop-up satellite archival

tags to demonstrate survival of blue marlin (*Makaira nigricans*) released from pelagic

longline gear. *Fishery Bulletin* **101**, 939–941.

Kiyota M, Yokota K, Nobetsu T, Minami H, Nakano H (2004) Assessment of mitigation

measures to reduce interactions between sea turtles and longline fishery. *Proceedings of*

the International Symposium on SEASTAR 2000 and Bio-Logging Science (The 5th

SEASTAR 2000 Workshop), 24–29.

Kruschke JK (2015) Doing bayesian data analysis: a tutorial with R, JAGS, and Stan. 2nd eds.

(Academic Press, New York, USA)

Maunder MN, Punt AE (2004) Standardizing catch and effort data: a review of recent

approaches. *Fisheries Research* **70**, 141–159.

https://doi.org/10.1016/j.fishres.2004.08.002

Melvin EF, Guy TJ, Read LB (2014) Best practice seabird bycatch mitigation for pelagic

longline fisheries targeting tuna and related species. *Fisheries Research* **149**, 5–18.

https://doi.org/10.1016/j.fishres.2013.07.012

Pacheco JC, Kerstetter DW, Hazin FH, Hazin H, Segundo RSSL, Graves JE, Carvalho F,

Travassos PE (2011) A comparison of circle hook and J hook performance in a western

equatorial Atlantic Ocean pelagic longline fishery. *Fisheries Research* **107**, 39–45.

https://doi.org/10.1016/j.fishres.2010.10.003

Parga ML, Pons M, Andraka S, Rendón L, Mituhasi T, Hall M, Pacheco L, Segura A, Osmond

M, Vogel N (2015) Hooking locations in sea turtles incidentally captured by artisanal

longline fisheries in the Eastern Pacific Ocean. *Fisheries Research* **164**, 231–237.

https://doi.org/10.1016/j.fishres.2014.11.012

Pebesma E (2018) Simple features for R: Standardized support for spatial vector data. *The R*

Journal **10**, 439–446. https://doi.org/10.32614/RJ-2018-009

R Core Team (2023) R: A Language and Environment for Statistical Computing. Available at

https://www.r-project.org/

Reinhardt JF, Weaver J, Latham PJ, Dell'Apa A, Serafy JE, Browder JA, Christman M, Foster

DG, Blankinship DR (2018) Catch rate and at-vessel mortality of circle hooks versus J-

hooks in pelagic longline fisheries: A global meta-analysis. *Fish and Fisheries* **19**, 413–

430. https://doi.org/10.1111/faf.12260

Santos MN, Coelho R, Fernandez-Carvalho J, Amorim S (2012) Effects of Hook and bait on

sea turtle catches in an equatorial Atlantic pelagic longline fishery. *Bulletin of Marine*

Science **88**, 683–701. https://doi.org/10.5343/bms.2011.1065

Santos CC, Rosa D, Coelho R (2020) Progress on a meta-analysis for comparing hook, bait

and leader effects on target, bycatch and vulnerable fauna interactions. *Collective Volume*

of Scientific Papers ICCAT **77**, 182–217.

Sims DW, Mucientes G, Queiroz N (2018) Shortfin mako sharks threatened by inaction.

Science **359**, 1342. https://doi.org/ 10.1126/science.aat0315

Skomal G, Bernal D (2010) Physiological responses to stress in sharks. In 'Sharks and Their

Relatives II.' (Eds JC Carrier, JA Musick, MR Heithaus), pp. 475–506. (CRC Press, Boca

Raton, USA)

Stokes L, Hataway D, Epperly S, Shah A, Bergmann C, Watson J, Higgins B (2011) Hook

ingestion rates in loggerhead sea turtles *Caretta caretta* as a function of animal size, hook

size, and bait. *Endangered Species Research* **14**, 1–11. https://doi.org/10.3354/esr00339

Swimmer Y, Gutierrez A, Bigelow K, Barceló C, Schroeder B, Keene K, Shattenkirk K,

Foster DG (2017) Sea turtle bycatch mitigation in U.S. longline fisheries. *Frontiers in*

Marine Science **4**, 260. https://doi.org/10.3389/FMARS.2017.00260

Stan Development Team (2021) Stan modeling language users guide and reference manual,

2.28.2. Available at https://mc-stan.org

Wallace BP, Kot CY, Dimatteo AD, Lee T, Crowder LB, Lewison RL (2013) Impacts of

fisheries bycatch on marine turtle populations worldwide: toward conservation and

research priorities. *Ecosphere* **4**, 1–49. https://doi.org/10.1890/ES12-00388.1

Watson JW, Epperly SP, Shah AK, Foster DG (2005) Fishing methods to reduce sea turtle

mortality associated with pelagic longlines. *Canadian Journal of Fisheries and Aquatic*

Sciences **62**, 965–981. https://doi.org/10.1139/f05-004

Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Grolemund G, Hayes

A, Henry L, Hester J et al. (2019) Welcome to the tidyverse. *Journal of Open Source*

Software **4**, 1686. https://doi.org/10.21105/joss.01686

Yokota K, Kiyota M, Minami H (2006a) Shark catch in a pelagic longline fishery:

Comparison of circle and tuna hooks. *Fisheries Research* **81**, 337–341.

https://doi.org/10.1016/j.fishres.2006.08.006

Yokota K, Kiyota M, Okamura H (2009) Effect of bait species and color on sea turtle bycatch

and fish catch in a pelagic longline fishery. *Fisheries Research* **97**, 53–58.

https://doi.org/10.1016/J.FISHRES.2009.01.003

Yokota K, Minami H, Kiyota M (2006b) Measurement-points examination of circle hooks for

pelagic longline fishery to evaluate effects of hook design. *Bulletin of Fishery Research*

Agency **17**, 83–102

Yokota K, Minami H, Kiyota M (2011) Effectiveness of tori-lines for further reduction of

incidental catch of seabirds in pelagic longline fisheries. *Fisheries Science* **77**, 479–485.

https://doi.org/10.1007/s12562-011-0357-4

Table 1 Fishing effort (longline hooks) in experimental operations used in the analysis.

hook type	bait type		total effort
	squid	fish	
tuna	97,834	71,146	168,980
small-C	70,882	7,658	78,540
large-C	31,872	6,976	38,848
total effort	200,588	85,780	286,368

Table 2 List of hook types used in the experiment.hook name	size	category	straight total length(mm)	maximum total width(mm)
Komatu Keisaku tunahook	4.0 sun	tuna	63	63
Doitomi Tunamutsu	4.0 sun	small-C (circle hook)	58	69
Komatsu Keisaku modified	4.0 sun	small-C (circle hook)	60	68
Komatsu Keisaku modified	4.5 sun	small-C (circle hook)	63	71
Komatsu Keisaku modified	4.8 sun	small-C (circle hook)	N/A	N/A
Komatsu Keisaku type Koshina	4.5 sun	small-C (circle hook)	62	76
Komatsu Keisaku type North America	4.3 sun	small-C (circle hook)	57	63
Tankichi Uruwa	3.8 sun	small-C (circle hook)	56	64
Komatsu Keisaku modified	5.2 sun	large-C (circle hook)	74	81
Komatsu Keisaku type North America	5.2 sun	large-C (circle hook)	76	85
Pacific Fishing Tackle circle hook	18/0	large-C (circle hook)	68	80

Table 3 Number of fish caught by hook and bait type in the experimental operation

species	total catch	hook type			bait type	
		tuna	small-C	large-C	squid	fish
blue shark	13,018	8,084	3,562	1,372	8,903	4,115
longnose lancetfish	1,297	945	257	95	692	605
common dolphinfish	505	206	181	118	363	142
shortfin mako shark	485	298	134	53	262	223
swordfish	288	129	112	47	249	39
bigeye tuna	269	114	146	9	201	68
loggerhead turtle	268	128	113	27	259	9
escolar	163	79	56	28	108	55
striped marlin	145	70	53	22	126	19
other species	1,578	-	-	-	-	-

Table 4 Composition of hooking location and fate at hauling.

species	hooking location				fate at haulback		
	swallowed	mouth	external	unknown (*)	alive	dead	unknown (*)
blue shark	2,270	2,608	66	8,074	11,701	1,066	251
longnose lancetfish	25	288	16	968	165	1,010	122
common dolphinfish	39	159	4	303	398	87	20
shortfin mako shark	66	68	18	333	363	118	4
swordfish	59	118	18	93	55	226	7
bigeye tuna	7	128	2	132	83	183	3
loggerhead turtle	109	116	14	29	256	5	7
escolar	1	62	1	99	108	40	15
striped marlin	12	80	8	45	68	76	1

(*) Includes catches that dropped off before reseachers checked or lack of survey.

Table 5 Estimates of the posterior distribution of the proportion of hooking location by hook when squid bait is used (median). Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses.

species	hook type		
	tuna	small-C	large-C
hooking place: external			
blue shark	0.020 (0.015 - 0.027)	0.008 (0.005 - 0.012)	0.005 (0.001 - 0.014)
loggerhead turtle	0.025 (0.006 - 0.066)	0.079 (0.037 - 0.143)	0.031 (0.001 - 0.154)
longnose lancetfish	0.049 (0.020 - 0.092)	0.031 (0.011 - 0.067)	0.074 (0.018 - 0.185)
shortfin mako	0.102 (0.041 - 0.199)	0.107 (0.052 - 0.185)	0.136 (0.021 - 0.381)
striped marlin	0.083 (0.026 - 0.186)	0.068 (0.016 - 0.173)	0.048 (0.002 - 0.235)
swordfish	0.075 (0.030 - 0.148)	0.109 (0.057 - 0.184)	0.031 (0.001 - 0.154)
hooking place: mouth			
blue shark	0.497 (0.475 - 0.518)	0.548 (0.528 - 0.568)	0.560 (0.517 - 0.603)
loggerhead turtle	0.439 (0.346 - 0.534)	0.480 (0.381 - 0.577)	0.746 (0.544 - 0.892)
longnose lancetfish	0.857 (0.792 - 0.908)	0.903 (0.848 - 0.943)	0.872 (0.740 - 0.953)
shortfin mako	0.356 (0.238 - 0.486)	0.524 (0.418 - 0.630)	0.379 (0.154 - 0.650)
striped marlin	0.759 (0.623 - 0.869)	0.831 (0.696 - 0.924)	0.883 (0.660 - 0.982)
swordfish	0.528 (0.416 - 0.637)	0.676 (0.574 - 0.767)	0.611 (0.405 - 0.790)
hooking place: swallowed			
blue shark	0.483 (0.461 - 0.505)	0.444 (0.424 - 0.464)	0.434 (0.391 - 0.478)
loggerhead turtle	0.533 (0.438 - 0.626)	0.438 (0.343 - 0.537)	0.210 (0.078 - 0.405)
longnose lancetfish	0.092 (0.052 - 0.148)	0.064 (0.032 - 0.111)	0.046 (0.007 - 0.145)
shortfin mako	0.536 (0.405 - 0.665)	0.365 (0.266 - 0.472)	0.460 (0.213 - 0.724)
striped marlin	0.150 (0.066 - 0.273)	0.093 (0.030 - 0.207)	0.048 (0.002 - 0.234)
swordfish	0.394 (0.289 - 0.505)	0.211 (0.136 - 0.303)	0.343 (0.174 - 0.550)

Table 6 Estimated haulback mortality rates (median of posterior distribution) by hooking location when squid bait is used. Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses. Loggerhead turtles were excluded because there were no mortalities and the calculation had not been converged.

species	external	mouth	swallowed
blue shark	0.295 (0.185 - 0.411)	0.052 (0.044 - 0.061)	0.157 (0.141 - 0.172)
shortfin mako shark	0.496 (0.262 - 0.729)	0.134 (0.061 - 0.219)	0.279 (0.176 - 0.394)
longnose lancetfish	0.956 (0.810 - 1.000)	0.860 (0.816 - 0.899)	0.798 (0.621 - 0.934)
swordfish	0.802 (0.595 - 0.959)	0.776 (0.694 - 0.853)	0.919 (0.842 - 0.979)
striped marlin	0.798 (0.488 - 0.990)	0.431 (0.315 - 0.547)	0.968 (0.839 - 1.000)

Table 7 Haulback mortality rate by hook and bait type (median of posterior distribution). Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses.

Species	Hook type		
	tuna	small-C	large-C
Bait type: squid			
bigeye tuna	0.734 (0.636 - 0.823)	0.757 (0.679 - 0.830)	0.959 (0.813 - 1.000)
blue shark	0.084 (0.077 - 0.093)	0.081 (0.072 - 0.091)	0.077 (0.062 - 0.093)
common dolphinfish	0.154 (0.099 - 0.215)	0.128 (0.081 - 0.180)	0.180 (0.110 - 0.265)
escolar	0.188 (0.082 - 0.309)	0.301 (0.165 - 0.454)	0.182 (0.060 - 0.340)
longnose lancetfish	0.901 (0.870 - 0.930)	0.925 (0.892 - 0.954)	0.889 (0.826 - 0.940)
shortfin mako	0.161 (0.106 - 0.223)	0.187 (0.123 - 0.258)	0.181 (0.085 - 0.300)
striped marlin	0.532 (0.400 - 0.665)	0.472 (0.334 - 0.609)	0.530 (0.318 - 0.740)
swordfish	0.829 (0.754 - 0.895)	0.783 (0.698 - 0.859)	0.832 (0.711 - 0.934)
Bait type: fish			
bigeye tuna	0.473 (0.319 - 0.622)	0.503 (0.337 - 0.663)	0.884 (0.573 - 1.000)
blue shark	0.078 (0.069 - 0.088)	0.075 (0.063 - 0.088)	0.071 (0.056 - 0.088)
common dolphinfish	0.238 (0.159 - 0.326)	0.201 (0.102 - 0.311)	0.274 (0.162 - 0.400)
escolar	0.242 (0.114 - 0.390)	0.371 (0.179 - 0.596)	0.233 (0.054 - 0.469)
longnose lancetfish	0.909 (0.879 - 0.936)	0.931 (0.894 - 0.961)	0.898 (0.841 - 0.947)
shortfin mako	0.307 (0.237 - 0.381)	0.348 (0.226 - 0.478)	0.339 (0.196 - 0.493)
striped marlin	0.691 (0.457 - 0.885)	0.638 (0.375 - 0.870)	0.691 (0.386 - 0.925)
swordfish	0.853 (0.727 - 0.950)	0.811 (0.648 - 0.934)	0.856 (0.700 - 0.967)

Table 8 Standardized catch per unit effort (CPUE) by hook and bait type (median of posterior distribution). Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses.

Species	Hook type		
	tuna	small-C	large-C
Bait type: squid			
bigeye tuna	0.433 (0.057 - 1.230)	0.825 (0.102 - 2.340)	0.680 (0.051 - 2.300)
blue shark	45.141 (36.893 - 53.976)	51.084 (41.940 - 61.337)	43.953 (35.882 - 53.072)
common dolphinfish	0.755 (0.384 - 1.282)	1.280 (0.622 - 2.172)	0.823 (0.383 - 1.403)
escolar	0.199 (0.060 - 0.409)	0.511 (0.157 - 1.054)	0.289 (0.079 - 0.631)
longnose lancetfish	3.287 (2.514 - 4.144)	3.598 (2.700 - 4.604)	3.204 (2.243 - 4.304)
shortfin mako	1.087 (0.771 - 1.433)	1.402 (0.969 - 1.878)	0.982 (0.615 - 1.436)
striped marlin	0.312 (0.140 - 0.537)	0.282 (0.118 - 0.493)	0.294 (0.112 - 0.564)
swordfish	0.721 (0.441 - 1.047)	0.825 (0.500 - 1.231)	0.840 (0.467 - 1.333)
Bait type: fish			
bigeye tuna	0.184 (0.020 - 0.550)	0.352 (0.033 - 1.047)	0.289 (0.020 - 1.059)
blue shark	36.417 (29.748 - 43.741)	41.224 (33.670 - 49.667)	35.459 (28.602 - 42.824)
common dolphinfish	1.855 (0.897 - 3.150)	3.146 (1.523 - 5.431)	2.023 (0.917 - 3.514)
escolar	0.437 (0.127 - 0.918)	1.123 (0.321 - 2.419)	0.634 (0.159 - 1.456)
longnose lancetfish	3.499 (2.643 - 4.507)	3.829 (2.829 - 5.003)	3.414 (2.372 - 4.724)
shortfin mako	2.294 (1.598 - 3.089)	2.953 (1.986 - 4.073)	2.071 (1.258 - 3.104)
striped marlin	0.350 (0.131 - 0.660)	0.316 (0.108 - 0.617)	0.331 (0.095 - 0.689)
swordfish	0.714 (0.395 - 1.124)	0.817 (0.434 - 1.308)	0.834 (0.389 - 1.394)

Table 9 Estimated MPUE by hook and bait type (median of posterior distribution). Lower and upper limits of Bayesian credible interval (95% highest density interval [HDI]) are shown in parentheses.

Species	Hook type		
	tuna	small-C	large-C
Bait type: squid			
bigeye tuna	0.316 (0.035 - 0.899)	0.623 (0.078 - 1.774)	0.637 (0.051 - 2.173)
blue shark	3.802 (3.031 - 4.628)	4.143 (3.263 - 5.120)	3.383 (2.494 - 4.401)
common dolphinfish	0.115 (0.048 - 0.211)	0.163 (0.067 - 0.301)	0.148 (0.056 - 0.279)
escolar	0.037 (0.008 - 0.088)	0.152 (0.035 - 0.345)	0.052 (0.007 - 0.140)
longnose lancetfish	2.960 (2.255 - 3.734)	3.322 (2.457 - 4.225)	2.839 (1.986 - 3.849)
shortfin mako	0.174 (0.096 - 0.261)	0.260 (0.144 - 0.397)	0.177 (0.066 - 0.328)
striped marlin	0.165 (0.068 - 0.294)	0.132 (0.051 - 0.243)	0.153 (0.048 - 0.320)
swordfish	0.595 (0.353 - 0.866)	0.643 (0.388 - 0.971)	0.694 (0.369 - 1.109)
Bait type: fish			
bigeye tuna	0.087 (0.008 - 0.266)	0.174 (0.017 - 0.540)	0.241 (0.014 - 0.906)
blue shark	2.844 (2.232 - 3.519)	3.094 (2.359 - 3.935)	2.527 (1.816 - 3.348)
common dolphinfish	0.438 (0.176 - 0.800)	0.627 (0.213 - 1.238)	0.550 (0.197 - 1.057)
escolar	0.104 (0.019 - 0.250)	0.407 (0.086 - 0.996)	0.144 (0.015 - 0.439)
longnose lancetfish	3.177 (2.389 - 4.094)	3.558 (2.611 - 4.651)	3.055 (2.075 - 4.207)
shortfin mako	0.703 (0.444 - 1.004)	1.020 (0.547 - 1.586)	0.697 (0.312 - 1.196)
striped marlin	0.237 (0.075 - 0.469)	0.195 (0.054 - 0.413)	0.219 (0.051 - 0.497)
swordfish	0.601 (0.323 - 0.960)	0.653 (0.324 - 1.065)	0.703 (0.320 - 1.201)

Table 10 Nominal CPUE, Haulback mortality rate and MPUE of Loggerhead turtle and all figures are based on aggregated operational data, not estimates.

hook type	CPUE		haulback mortality		MPUE	
	squid	fish	squid	fish	squid	fish
tuna	1.247	0.084	0.0385	0.000	0.0480	0.000
large-C	0.816	0.143	0.0090	0.000	0.0073	0.000
small-C	1.566	0.261	0.0246	0.000	0.0385	0.000

Figure 1 Locations where the longline operation experiment was conducted.

Figure 2 Differences in the estimated probability of the "swallowed" hooking location of each circle hook type from tuna hook when squid bait is used. The red dotted line indicates that the difference is zero.

Figure 3   Differences in the estimated haulback mortality rate of each target hooking location from "swallowed" hooking location when squid bait is used. The red dotted line indicates that the difference is zero.

Figure 4 Differences in estimated haulback mortality between each experimental group and the control group (squid x tuna hook). The red dotted line indicates that the difference is zero.

Figure 5 Relationship between sea surface temperature (SST) variability and haulback mortality rate at longline operations. Solid lines indicate median; masked areas indicate 95% Bayesian credible interval.

Figure 6 Relationship between soak time (time from setting the branch line to hauling) variability and haulback mortality rate at longline operations. Solid lines indicate median; masked areas indicate 95% Bayesian credible interval.

Figure 7　Differences in the standardized catch per unit effort (CPUE) for each experimental group from those for the control group (squid x tuna hook). The red dotted line indicates that the difference is zero.

Figure 8   Differences in the estimated MPUE (mortality per unit effort) between those for each experimental group and those for the control group (squid x tuna hook). The red dotted line indicates that the difference is zero.

Figure 9 Alluvial plot of hooking locations and associated haulback mortality rates of loggerhead turtles by hook when squid bait is used.

Appendix 1 Detailed measurements of hooks used in the experiment and a figure explaining measurement points of hooks (copied from Yokota et al. 2006b).

manufacture	Komatsu Keisaku	Hisamatsu Tankichi	Doitomi	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Komatsu Keisaku	Pacific Fishing Tackle MFG., CO.
hook name	tuna hook	Uruwa hook BKN	tuna circle hook SS-170	modified circle hook	modified circle hook	modified circle hook	cirlce hook type Koshina	cirlce hook type North America	cirlce hook type North America	modified circle hook	circle hook
standardized size	4.0 sun	3.8 sun	#4	4.0 sun	4.5 sun	4.8 sun	4.5 sun	4.3 sun	5.2 sun	5.2 sun	18/0
material	stainless steel	hard steel	stainless steel	stainless steel	stainless steel	stainless steel	stainless steel	hard steel	hard steel	stainless steel	stainless steel
hook eye	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring	yes, with ring
shank thickness(mm)	5.3	4.0	4.1	5.3	5.2	N/A	4.9	5.3	5.7	5.3	5.1
straight total length(mm)	63	56	58	60	63	N/A	62	57	74	76	68
straight total width(mm)	38	44	49	47	49	N/A	56	45	56	54	59
minimum total width(mm)	38	36	39	41	45	N/A	51	41	52	48	51
maximum total width(mm)	63	64	69	68	71	N/A	76	63	81	85	80
front length(mm)	41	33	38	35	44	N/A	47	39	49	47	45
minimum inner width(mm)	27	20	15	24	25	N/A	26	20	27	26	27
L-W ratio	1.7	1.3	1.2	1.3	1.3	N/A	1.1	1.3	1.3	1.4	1.2
max-min ratio	1.7	1.8	1.8	1.7	1.6	N/A	1.5	1.5	1.6	1.8	1.6
incurved point angle		80	70	90	70	N/A	70	65	75	80	80
offset angle	$5° \leqq \theta < 10°$	$\theta \approx 0°$	$\theta \approx 0°$	$\theta \approx 10°$	$5° \leqq \theta < 10°$	$\theta < 10°$	$\theta \approx 10°$	$5° \leqq \theta < 10°$	$10° \leqq \theta < 15°$	$\theta \approx 5°$	$10° \leqq \theta < 15°$
offset width(mm)	0.9	≈ 0	≈ 0	1.8	2.6	N/A	2.1	1.5	3.7	1.0	5.4
weight (g)	19.9	12.2	15.0	19.7	21.6	N/A	21.4	19.4	30.3	25.5	23.2

Appendix 2 Stan code used for MCMC sampling of (a) Model 1 and (b) Model 2

(a)   MODEL 1

```
data {
 int HK_cat;
 int<lower=1> N1;
 int<lower=1> LOC_cat;
 int<lower=1, upper=LOC_cat> LOC[N1];
 int<lower=1, upper=HK_cat> HK[N1];
 int<lower=0, upper=1> M[N1];
 int L;
 matrix[N1, L] X1;
 int N2;
 int  Catch[N2];
 int L2;
 matrix[N2, L2] X2;
 vector[N2] O;
 int YRID;
 int YR[N2];
}

transformed data{
 vector[HK_cat] Zeros;
 Zeros = rep_vector(0, HK_cat);
}

parameters {
 matrix[HK_cat, LOC_cat - 1] theta_raw;
 vector[L] beta;
 vector[L2] beta2;
 vector[YRID] r;
 real<lower= 0> sigma;

}

transformed parameters {
 vector[N2] mu;
 vector[N1] phi;
 matrix[HK_cat, LOC_cat] theta;
 phi = inv_logit(X1 * beta);
 theta = append_col(Zeros, theta_raw);
 for(i in 1:N2)
 mu[i] = X2[i,] * beta2 + r[YR[i]];

model {
 sigma ~ student_t(4, 0, 2.5);
 r ~ normal(0, sigma);
 for(n in 1:N1){
 M[n] ~ bernoulli(phi[n]);
 target += categorical_lpmf(LOC[n]|softmax(theta[HK[n],]'));
 }
 Catch ~ poisson(exp(mu + log(O)));
}
```

(b)   MODEL 2

```
data {
  int HK_BAIT_comb;
  int<lower=1> N1;
  int<lower=0, upper=1> M[N1];
  int L;
  matrix[N1, L] X1;
  int N2;
  int Catch[N2];
  int L2;
  matrix[N2, L2] X2;
  vector[N2] O;
  int YRID;
  int YR[N2];
}

parameters {
  vector[L] beta;
  vector[L2] beta2;
  vector[YRID] r;
  real<lower= 0> sigma;
}

transformed parameters {
  vector[N2] mu;
  vector[N1] phi;
  phi = inv_logit(X1 * beta);
  for(i in 1:N2)
  mu[i] = X2[i,] * beta2 + r[YR[i]];
}

model {
  sigma ~ student_t(4, 0, 2.5);
  r ~ normal(0, sigma);
  for(n in 1:N1){
  M[n] ~ bernoulli(phi[n]);
  }
  Catch ~ poisson(exp(mu + log(O)));
}
```

2 Appendix 3 Size distributions by species for the major species captured in the study, with body

3 length as an index of precaudal length for blue sharks and shortfin mako sharks, straight-line

4 carapace length for loggerhead turtles, eye-to-fork length for striped marlin and swordfish, and fork

5 length for all other species.

6