A Multi-Tissue Comparison and Molecular Characterization of Canine Organoids

Authors: Christopher Zdyrski[1,2,3]*, Vojtech Gabriel[1]*, Oscar Ospina[3], Hannah Wickham[1], Dipak K. Sahoo[5], Kimberly Dao[3], Leeann S. Aguilar Meza[1], Abigail Ralston[3], Leila Bedos[5], William Bastian[6], Sydney Honold[1], Pablo Piñeyro[7], Eugene F. Douglass[6], Jonathan P. Mochel[#1,2,3], Karin Allenspach[#1,2,3,5]

[1] SMART Pharmacology, Department of Biomedical Sciences, Iowa State University, Ames, IA, USA
[2] Present Address: SMART Pharmacology, Precision One Health Initiative, University of Georgia, Athens, GA, USA
[3] 3D Health Solutions Inc., Ames, IA, USA
[4] Department of Biostatistics and Bioinformatics, Moffitt Cancer Center, Tampa, FL, USA
[5] Department of Veterinary Clinical Sciences, Iowa State University, Ames, IA, USA
[6] Pharmaceutical & Biomedical Sciences, Institute of Bioinformatics, University of Georgia, Athens, GA
[7] Veterinary Diagnostic Laboratory, Iowa State University, Ames, IA, USA

*Authors contributed equally to the manuscript.
co-senior authors.

Corresponding authors: Christopher Zdyrski (czdyrski@uga.edu) Ph.D.; K. Allenspach (Karin.Allenspach@uga.edu) DVM, Ph.D.; Jonathan P. Mochel (jpmochel@uga.edu) DVM, Ph.D.

DECLARATION OF INTERESTS

K. Allenspach is a co-founder of LifEngine Animal Health and 3D Health Solutions. She serves as a

consultant for Ceva Animal Health, Bioiberica, LifeDiagnostics, Antech Diagnostics, Deerland Probiotics,

and Mars.

J.P. Mochel is a co-founder of LifEngine Animal Health and 3D Health Solutions. Dr. Mochel serves as a

consultant for Ceva Animal Health, Boehringer Ingelheim, Dechra Ltd. and Ethos Animal Health.

C. Zdyrski is the Director of Research and Product Development at 3D Health Solutions.

K. Dao was an employee of 3D Health Solutions.

A. Ralston was an employee of 3D Health Solutions.

Other authors do not have any conflict of interest to declare.

Multiple patents related to this work have been filed through the Iowa State University Research

Foundation.

13 **ABSTRACT**

Organoids are 3-dimensional (3D) stem cell-derived cell culture lines that offer a variety of

technical advantages compared to traditional 2-dimensional (2D) cell cultures. Although murine

models have proved useful in biomedical research, rodent models often fail to adequately mimic

human physiology and disease progression, resulting in poor preclinical prediction of

therapeutic drug efficacy and toxicity. With the advent of organoid technology, many of these

challenges can be overcome. Previously, the use of canine organoids in drug testing and

disease modeling was limited to organoids originating from the intestine, liver, kidney, lung, and

urinary bladder. Here, we report the cultivation, maintenance, and molecular characterization of

two novel adult-stem cell-derived canine organoid cell lines, including the endometrium and

pancreas, in addition to previously reported bladder, lung, and liver organoids from two

genetically related canines. Five tissues and organoid lines from each donor were characterized

using bulk RNA-seq, allowing for a unique, multi-organ comparison between these two

individuals and identification of specific cell types such as glandular epithelial cells in

endometrial organoids.

Keywords: canine; dog; organoids; reverse translational medicine; stem cell; endometrium;

lung; pancreas; bladder; liver

INTRODUCTION

Numerous *in vitro* models are used as preclinical biological and pharmacological research tools (Hickman et al., 2014). The most prevalent *in vivo* models for biomedical research include *Drosophila melanogaster* and *Caenorhabditis elegans*, both of which are widely used for genetic research, *Danio rerio*, a model used primarily for mutagenesis screening, and numerous mammalian models for which advanced genetic tools are available (Kim et al., 2020b). Mouse models are extensively used in biomedical research due to their cost-effectiveness, fast-growing nature, and availability of genetic mutants (Kim et al., 2020b). Differences in diet, living environment, circadian rhythm, and the short lifespan are among the issues limiting the translational relevance of rodent models (Perlman, 2016). Although the murine model has proven effective in a variety of biological research areas, rodents frequently fail to adequately mimic human physiology and disease progression, hence compromising their predictive performance in preclinical pharmaceutical research (Gordon et al., 2009; Wang et al., 2018a). Approximately ninety percent of experimental drugs fail to make the transition from discovery to successful clinical trials (Mochel et al., 2018; Van Norman, 2016; Shalev, 2006). Drug development is successful in about 5% of those ultimately approved, and average costs including post-approval R&D in the U.S. exceed $2 billion and it takes 10-15 years to reach clinical trials per drug (Brancato et al., 2020; DiMasi et al., 2016). The use of 3D organoids in the screening stage of drug discovery could drastically reduce the use of live animals for drug development (Mollaki, 2021). Ultimately, additional research is warranted to identify alternative *in vitro* models that can more accurately replicate human physiology and reduce animal use. Conventional pharmacology research involves using 2D cell culture and animal testing prior to human clinical trials (Brancato et al., 2020).

Organoids are 3D self-organized, miniature, and simplified versions of organs *in vitro*. Adult stem cells can self-renew, differentiate into multiple cell types, and are genomically stable over

multiple passages (Fatehullah et al., 2016; Huch and Koo, 2015; Huch et al., 2015). Unlike

traditional 2D cell lines, organoids grow in a 3D extracellular matrix, allowing for the recreation

of more realistic tissue architecture and physiological responses (Hynds and Giangreco, 2013).

Recently, donor-to-donor variability in human ileum- and colon-derived organoids has been

investigated; however, more research is needed across other biomedical models, including

dogs (Mohammadi et al., 2021). Furthermore, organoids can be used in both basic and applied

biomedical research, including the study of genetic disorders, cancers, and infectious diseases

(Kim et al., 2020b; Nantasanti et al., 2015; Usui et al., 2017; Zhou et al., 2018). Organoids can

be a valuable tool for personalized medicine where patient-specific organoids can be grown and

incubated with drug candidates to predict effectiveness prior to patient treatment (Bartfeld and

Clevers, 2017; Kaushik et al., 2018). Organoids may become useful for regenerative medicine;

for example, hepatic organoids can be transplanted to a patient and transdifferentiated to

various hepatocellular regional identities *in vivo* (Sampaziotis et al., 2021). Organoid cell lines

drastically reduce the number of animals needed for drug testing as they can be expanded

indefinitely in culture and cryopreserved for future use. Organoid technology has made it

possible to undertake pharmacological research and testing in a manner that is more

responsible from an ethical standpoint, while also simplifying the process of genetic

manipulation (Artegiani et al., 2020; Takeda et al., 2019). While human organoids are a valuable

research tool in the biomedical field, they come with limitations. Public concern plays a key role

in tissue sampling from human patients (Lehmann et al., 2019). Ethical concerns for the use of

human-derived organoids include chimeric research and genetic editing of organoids derived

from patients (Munsie et al., 2017). Although a growing number of studies have characterized

cell populations using transcriptomic data across human organs, there is a lack of similar

studies in non-human models (Jones et al., 2022).

The reverse translational paradigm, in which data from human clinical research might aid in the

development of veterinary therapeutics and vice versa, is garnering a growing amount of

interest (Schneider et al., 2018). Dogs share similar lifestyles and diets with their owners due to

the close relationship between dogs and humans, often including a sedentary lifestyle and an

increased risk of developing obesity (Chandler et al., 2017). The longer lifespan of dogs over

that of mice predisposes dogs to develop analogous chronic diseases to humans, including

diabetes mellitus (Adin and Gilor, 2017), ocular diseases (Sebbag et al., 2018, 2019),

inflammatory bowel disease (Chandra et al., 2019; Jergens and Simpson, 2012; Kopper et al.,

2021), congestive heart failure (Mochel and Danhof, 2015; Mochel et al., 2019; Silva and Emter,

2020), cancers (Knapp et al., 2020), and cognitive dysfunction (Ozawa et al., 2019), among

others (Wang et al., 2018a). Therefore, in addition to being used as a large animal model for

preclinical drug safety assessment, dogs are also emerging as a translatable model for

demonstrating proof-of-concept efficacy studies, particularly in the field of oncology (LeBlanc et

al., 2016; Schaefer et al., 2016). While dogs excel as a model in many applications compared to

rodents, they come with their own challenges in the form of expensive housing and ethical

concerns about using live dogs in research. Dogs are recognized as companion animals in

western countries, and there are ongoing worldwide initiatives to limit their use in research

through the 3Rs (*Reduce*, *Replace*, *Refine*) principles (Hasiwa, 2011; Russell and Burch, 1959).

A potential solution that provides more access to the canine model while decreasing reliance on

live animal use lies in organoid technology.

Organoids still represent a relatively novel technology and lack formal standardization in

isolation, maintenance, and downstream applications. We previously demonstrated the ability to

culture canine intestinal organoids from healthy and diseased tissues and demonstrated the

translational potential of these organoids for human medicine (Chandra et al., 2019). Our

research group has been working to standardize the culture and maintenance of organoid cell

lines to maximize the reproducibility of our findings across different experimental sites (Gabriel

et al., 2022a). This investment into the standardization of protocols includes downstream

applications such as the use of a permeable support system for canine organoids in drug testing

and discovery (Gabriel et al., 2022b), use of organoids in viral testing (García-Rodríguez et al.,

2020), and regenerative medicine (Sampaziotis et al., 2021).

Currently, there is a lack of canine organoid models compared to other major biomedical

species to accurately depict and study various diseases, drugs, and biological phenomena.

This report describes the successful cultivation of two novel canine organoid lines, including the

endometrium and pancreas, to the best of the authors' knowledge, there has not been any peer-

reviewed description of these organoids, in addition to previously described bladder (Elbadawy

et al., 2022), lung (Shiota (Sato) et al., 2023), and liver (Nantasanti et al., 2015) organoids from

two related dogs. By comparing five tissue-specific organoid lines obtained from two genetically

related donors (B816 and B818), this study aims to acquire insight into gene expression in

different organoids and their corresponding tissues. Preliminary analyses using RNA

sequencing, immunohistochemistry, and immunofluorescence were used to characterize the

relationships between individual organoid cell lines and their parent tissue. In addition to

characterization, we compared related individuals across organs in this preliminary analysis,

thus preparing these organoids to next be utilized and tested in a variety of biomedical

applications and functional assays.

RESULTS

Organoid Expansion

Organoid cell lines were successfully established from five organs, including the uterus, lung,

pancreas, urinary bladder, and liver, two of which are novel, originating from two female canine

individuals. All tissues were isolated on the same day, minced, and embedded in Matrigel.

Organoids were cultivated simultaneously, and growth progression, passage number, and

media are reported in **Supplemental file 1.** Samples were passaged between two and four

times to remove excess tissue fragments before being harvested for characterization and

freezing (**Figure 1-figure supplement 1**). Wells with any remaining tissue were excluded from

any characterization to ensure reliable results. The same media was used for all organoid lines

and the media composition is listed in **Supplemental file 2** and consists of growth factors which

encourage the growth of multiple tissues. Several of the canine organoids displayed a variety of

distinct morphological phenotypes characterized via light microscopy and hematoxylin & eosin

(H&E) staining (**Figure 1**).

Immunohistochemistry (IHC) for Pan cytokeratin (PanCK) was used to confirm the epithelial

origin of the canine organoids, while tissues were positive only in epithelial regions (**Figure 2**).

To better confirm and distinguish that the organoids were not remaining tissue, smooth muscle

actin (SMA) was stained (**Figure 2**) and vimentin (VIM) was used to identified mesenchymal

cells in the tissues and organoids (**Figure 2**).

Morphological and histological characterization of canine organoid cell lines

Uterus

A subset of endometrial organoids formed a tubular structure appreciated on brightfield

microscopy during culture (**Figure 1**). H&E staining suggested that the culture consists of

endometrial epithelial cells and glandular epithelial cells (**Figure 1**). RNA-seq data indicated that

SRY-box transcription factor 17 (SOX17) was upregulated in endometrium organoids (**Figure**

3B). This protein is expressed in the human endometrium, specifically in the luminal and

glandular epithelium (Kinnear et al., 2019). SOX17 is important for endometrial glandular

development and function in mice (Turco et al., 2017), and was also expressed in organoids

derived from human menstrual flow, consistent with their function in endometrial gland

development (Cindrova-Davies et al., 2021). Uroplakin Ib (UPK1B) was upregulated in

endometrium organoids compared to uterus tissues (**Figure 3A**). The top endometrium-specific

genes for organoids included Actin gamma 1 (ACTG1), Clusterin (CLU), and Prothymosin alpha

(PTMA), whereas uterine tissues showed high expression of multiple ribosomal proteins (**Figure**

3D). Regarding endometrium-specific genes, intra-organoid comparison revealed 1,039 unique

genes (**Figure 4A**), uterine intra-tissue comparisons revealed 2,930 unique genes (**Figure 4B**),

and 10,487 genes were expressed in both organoids and tissues (**Figure 4C**). For

immunofluorescence (IF), Vimentin (VIM) was positive in both tissues and organoids while

Paired Box 8 (PAX8) identified endometrial glands in tissues (**Figure 5**).

Pancreas

The pancreatic organoids displayed two phenotypes, one resembling spheroids and another of

a flowering organoid (**Figure 1**). H&E staining suggested that the culture mainly consisted of

cells derived from intercalated ducts, with a few cells potentially differentiating into endocrine

cells (**Figure 1**). Genes that were found to be upregulated in organoids compared to tissue

included Dual oxidase 2 (DUOX2), Pyroglutamylated Rfamide peptide (QRFP), Cadherin 17

(CDH17), and Early growth response 1 (EGR1) (**Figure 3A**). Noteworthy is that Somatostatin

(SST) expression was reduced in organoids suggesting our culture does not contain a

significant number of neuroendocrine delta cells (**Figure 3A**). However, maltase-glucoamylase

2 (MGAM2) and NK6 homeobox 1 (NKX6-1) were also upregulated in organoids (**Figure 3B**).

NKX6-1 serves as a marker of multipotent pancreatic progenitors, indicating their ability to

differentiate into ductal, acinar, and endocrine cells (Wiedenmann et al., 2021). Upregulated

genes in pancreas tissues included cells positive for insulin (INS), glucagon (GCG), and multiple

markers characteristic of pancreatic acinar cells (**Figure 3C**). One of the most highly expressed

pancreas-specific genes in the organoids included cytokeratin 7 (KRT7), indicating that most of

the cells in the organoids are of epithelial origin and likely represent pancreatic ductal cells

(Wiedenmann et al., 2021). Regarding pancreas-specific genes, intra-organoid comparison

revealed 482 unique genes (**Figure 4A**), intra-tissue comparison revealed 344 unique genes

(**Figure 4B**), and 10,173 genes were expressed in both organoids and tissues (**Figure 4C**). IF

staining on pancreatic tissues identified insulin and glucagon in the islets (**Figure 5**); however,

the organoids did not express measurable insulin nor glucagon.

Lung

The lung organoids displayed three distinct phenotypes with flowering differentiated organoids

and bulbous organoids constituting most of the culture, while a small proportion had a

morphology resembling alveolar structures (**Figure 1**). H&E staining suggests our culture

consists of alveolar type-2 cells (AT2) and bronchial epithelial cells (**Figure 1**). The lung marker,

NK2 homeobox 1 (NKX2-1) (Dost et al., 2020), was upregulated in organoids (**Figure 3B**), while

Surfactant Protein B (SFTPB) and Surfactant Protein C (SFTPC) gene expression were

detected and specific to both lung tissues and organoids (**Figure 3D**). Upregulated genes in

organoids compared to tissues included Pyroglutamylated RFamide peptide (QRFP) and

Peptidoglycan recognition protein 1 (PGLYRP1). For lung-specific genes, intra-organoid

comparison revealed 736 unique genes (**Figure 4A**), intra-tissue comparisons revealed 1,963

unique genes (**Figure 4B**), and 10,597 genes were expressed in both organoids and tissues

(**Figure 4C**). IF staining clearly identified alveolar type-1 cells (AT1) for Claudin 18 (CLDN18)

while Lysosome-associated membrane glycoprotein 3 (LAMP3) identified AT2 cells in the tissue

sections, however, organoids did not show IF signal for either (**Figure 5**).

Urinary Bladder

Bladder organoids displayed a single phenotype constituting round organoids without a visible

lumen or internal chamber typical of spheroids of other tissues such as intestine (**Figure 1**)

(Chandra et al., 2019; Gabriel et al., 2022a). H&E staining clearly showed the organoids present

morphological features consistent with transitional epithelium, consisting of characteristic basal

and umbrella cell layers (**Figure 1 and Figure 5**). Desmoglein 3 (DSG3), which is a basal cell

marker (Elbadawy et al., 2019), and Loricrin cornified envelope precursor protein (LORICRIN),

an intermediate cell marker (Lin et al., 2013), were upregulated in organoids compared to

tissues (**Figure 3A**). Uroplakin (UPK) proteins are specific to terminally differentiated urothelial

cells (Lu et al., 2022), and Uroplakin 2 (UPK2) were both upregulated in organoids (**Figure 3B**).

The bladder-specific marker EIF4A1 was expressed in both organoids and tissues. Regarding

bladder-specific genes, intra-organoid comparisons revealed 1,071 unique genes (**Figure 4A**),

intra-tissue comparisons revealed 949 unique genes (**Figure 4B**), and 10,499 genes were

expressed in both organoids and tissues (**Figure 4C**). The presence of umbrella cells and basal

cells was confirmed by positive IF staining with Uroplakin Ia (UPK1A) and Cytokeratin 5 (CK5)

(**Figure 5**) in tissues (Elbadawy et al., 2022; Kim et al., 2020a). UPK1A expression was present

in the "pores" of umbrella cells, while expression of CK5 was positive and limited to the outside

of the bladder organoids, which is characteristic of inverted epithelial growth in organoid cultures

(**Figure 5**).

Liver

Liver organoids morphologically resembled the pancreatic organoids, with one phenotype

resembling spheroids and another of a more flowering organoid. Cellular morphology observed

under H&E evaluation suggests that most of the cells were differentiated cholangiocytes (**Figure**

1). Trefoil factor 1 (TFF1) and Tripartite Motif Containing 71 (TRIM71) were upregulated in liver

organoids compared to liver tissue (**Figure 3A**). TFF1 encodes a protein critical in the

regeneration of the liver after injury by promoting biliary lineage differentiation and inhibiting

hepatic lineage (Hayashi et al., 2018). Single-cell RNA sequencing of the human liver described

a transcriptional profile of a cell population within cholangiocytes where the DE genes included

TFF1 (MacParland et al., 2018). TRIM71 was also upregulated in organoids (**Figure 3B**) and

has previously been hypothesized to be involved in promoting rapid self-renewal in

undifferentiated mouse embryonic stem cells (Chang et al., 2012). Liver-specific organoid

markers included multiple heterogeneous nuclear ribonucleoproteins (**Figure 3D**). Albumin

(ALB) was the most highly expressed liver-specific gene in tissue due to the large percentage of

hepatocytes (**Figure 3D**). Regarding liver-specific genes, intra-organoid comparisons revealed

1,858 unique genes (**Figure 4A**), tissues had 1,386 unique genes (**Figure 4B**), and 9,984

genes were expressed in both organoids and tissues (**Figure 4C**). Using IF, hepatocytes in the

tissue stained positively for Hydroxyacid Oxidase 1 (HAO1) (**Figure 5**) (Kampf et al., 2014). As

expected, tissues showed high expression of HAO1 as hepatocytes constitute the major cell

type in liver tissue while organoids were negative for this marker, suggesting they almost

entirely consist of differentiated cholangiocytes, consistent with previous descriptions in other

liver-derived organoid cultures (Aktas et al., 2022; Zdyrski et al., 2024).

Insights into organ-specific genes

The usage of multiple methods (**Figure 6A**) including RNA-seq, allowed for identification of

differentially expressed genes between tissues and organoids (**Figure 3A**), and assisted in the

determination of the major cell populations present and absent in the organoid cell lines. We

acknowledge that the organoids are exclusively composed of epithelial cells and lack other

populations present in intact tissue, such as immune cells and endothelial cells. Genes

expressed were identified for each tissue type (**Figure 4C**) to emphasize the similarity of

expression patterns of the organoid models compared with their tissue of origin. A comparison

of mRNA expression across tissues and organoids can be seen in **Figure 4.** Between 76% and

80% (**Figure 4C**) of all expressed genes overlapped for each organ between tissues and

organoids.

Principal component analysis (PCA) (**Figure 4D**) was used to visualize major sources of genetic

variance for the different samples, where principal components 1 and 2 (PC1, PC2) effectively

separated epithelial organoids and their tissues of origin. Furthermore, PCA across all

organoids and all tissues, separately, clearly demonstrated the strongest sample clustering

between genetically related animals (**Figure 4E**). Intra-organoid and intra-tissue comparisons

identified upregulated genes (**Figure 3B and 3C**) and the ten most highly expressed tissue-

specific genes (**Figure 3D**). The expression of unique and overlapping genes was further

compared for each sample type (**Figure 4A and 4B**). Gene set enrichment analysis was used to

characterize the global transcriptional programs that best characterized both organoids and

tissues samples (**Figure 6E**). Stouffer integration of transcriptional pathway z-scores yielded a

consensus scoring of the major global differences between paired organoids and source tissues

(**Figure 6B**, **Supplemental file 8)**. Overall, tissue and organoid samples were best separated

by transcriptional hallmarks associated with inflammation and proliferation, respectively (**Figure**

6D). This preliminary characterization will assist in determining potential applications for these

novel canine organoid models for research applications (**Figure 6C**).

DISCUSSION

Canine organoids as biomedical models

Canines can serve as a superior model to mice for translational research applications,

especially due to their tendency to develop analogous chronic diseases to those of humans and

their shared similarity in lifestyle. However, using canines for translational research presents

some obstacles. Their use in research can be ethically questionable and resource-intensive.

Organoids can overcome some of these challenges and could potentially represent an excellent

alternative to expanding the biomedical applications of the canine model. Developing novel

canine organoid models will accelerate research efforts toward advanced veterinary

therapeutics as well as for preclinical drug screening in human medicine. Furthermore,

advances in organoid technology are being made in areas including personalized drug testing

using patient-derived organoid cultures (Huch and Koo, 2015; Mullenders et al., 2019).

Typically, studies report the cultivation of one tissue-specific organoid cell line, while others

combine organoid lines from multiple individuals to make conclusions. By combining unrelated

donors' information, donor-to-donor variability can be neglected, thus ignoring relevant

differences in patient populations. We aim to broaden the applications of the dog model in

biomedicine while minimizing animal usage by developing new canine organoid lines and

studying gene expression profiles across different epithelial tissues. While our tissue samples

consist of various cell types (such as epithelial cells, vascular cells, and immune cells), our

organoids are epithelial in origin. Comparisons between the original tissues and the derived

organoids expose the constraints of current epithelial organoid models. Recently, protocols for

co-culturing immune cells with organoids have become more attainable, thus enhancing the

complexity and applicability of organoid models. (Chakrabarti et al., 2021).

Many laboratories utilize tissue-specific protocols and media supplemented with various growth

factors to expand their organoids which can be costly and laborious (Kaushik et al., 2018;

Tortorella et al., 2022). In the current study, all described cultures were grown under the same

conditions using the same organoid expansion media (*Complete media with growth factors with*

ROCK inhibitor and GSK3β inhibitor – CMGF+ R/G) and differentiation media (*Complete media*

with growth factors – CMGF+). The use of the same media composition lends itself to future

applications of co-culture or use in assembloid models where multiple organoid lines are

combined and continued growth in a shared media composition is ideal. While our media

composition allows for the growth and expansion of organoids and a variety of cell types, the

authors acknowledge that for differentiation of certain epithelial cell types, such as within the

pancreas, specific growth factors and culture conditions may need to be optimized per tissue.

Additionally, many protocols emphasize tissue removal to create a suspension of stem cells

during organoid isolations. Here, we utilize mechanical dissociation during isolation for the

inclusion of small tissue pieces during the initial growth that we believe can benefit canine

organoid expansion, assisting the most in liver cultures. The tissue is then dissociated

enzymatically during organoid passages and hence removed from the culture prior to any

analysis. This phenomenon we observed may be due to intercellular signaling from the stem

cells still attached to damaged tissue, resulting in the release of damage-associated signals,

increasing the initial growth of the stem cells, or potentially due to the availability of the

extracellular matrix from the tissue at the beginning of the culture.

We report the cultivation, characterization, and comparison of five organoid lines (endometrium,

pancreas, lung, bladder, and liver), two of them novel, derived from the same animal, from two

genetically related canine donors of the same litter, sex, and age. Additionally, the isolation,

cultivation, and media composition were identical for the five organoid cell lines, eliminating the

need for tissue-specific growth factors. To the best of the authors' knowledge, the data in this

study, therefore, constitutes the most comprehensive comparison of tissue-specific expression

across canine organoids available to date. These newly available canine organoids could be

applied for more rapid translational applications, such as the identification of new therapeutics,

the study of genetic editing technologies, and the development of better disease models. Due to

the nature of the samples being derived from the same donors, these lines have the potential to

be used in downstream experiments including organ-on-a-chip (Dongeun et al., 2010) and

assembloid cultures (Birey et al., 2017).

Organoid characterization and biomedical applications

Uterus

Previously, it has been shown that UPK1B, which was upregulated in our canine endometrial

organoids, was upregulated after an endometrial biopsy, and the protein was found in glandular-

epithelial cells (Kalma et al., 2009). Human endometrium organoids typically resemble a cystic-

shaped organoid unlike the canine endometrial organoids, which contained large tubular

structures (Boretto et al., 2017; Turco et al., 2017). One study administered hormones to the

culture and noticed columnar epithelial morphology with the formation of larger vacuoles

(Cindrova-Davies et al., 2021). Previously, cultures of 3D uterine glands explants and stromal

cells had limited viability surviving only for four days with the resemblance of spheroids

beginning to form (Stadler et al., 2009). Furthermore, 3D organotypic canine endometrium

cultures have been previously described (Bartel et al., 2013), however this study simply isolated

differentiated endometrial glands and stromal cells from tissue and co-cultured them for 48

hours, not attempting to proliferate or expand the cells. Endometrium organoids may also be

useful in the future for investigating diseases such as endometriosis and endometrial cancers

(Turco et al., 2017). Additionally, they may be used in studies related to embryo implantation

into the endometrium (Rawlings et al., 2021). Finally, co-culturing endometrium organoids on

three-dimensional scaffolds may provide insight into implantation studies (Cindrova-Davies et

al., 2021).

Pancreas

The characterization of our canine pancreas-derived organoids suggests that they primarily

contain intercalated pancreatic ductular cells. In addition, we believe these stem cells should

have the capacity to differentiate into neuroendocrine cells but optimization of growth factors in

the media may be required. In dogs, pancreatitis is by far the most common disease of the

exocrine pancreas (Lim et al., 2014; Xenoulis, 2015), therefore, a healthy pancreatic canine

organoid model could assist in studying the pathophysiology of pancreatitis in this species.

Murine *in vivo* models have long been known to have limited translatability for modeling

pancreatic cancer in humans (Bailey and Carlson, 2019). Canine pancreatic ductal organoids

could potentially be used for disease modeling of pancreatic ductal adenocarcinoma (PDAC),

which is one of the most lethal types of cancer in humans (Wiedenmann et al., 2021).

Furthermore, differentiation of our canine pancreatic cultures can be attempted in the future

since recently described methods successfully differentiated 2D canine pancreatic ductal cells

into insulin- and glucagon-producing beta-like cells (Gao et al., 2022). Such applications could

then be used for pancreatic hormone production and studying drug target screening and

toxicological effects on the endocrine pancreas.

Lung

Canine lung organoids have previously been described (Shiota (Sato) et al., 2023); however,

the organoid cultures we describe here contained a variety of distinct morphologies in addition

to what has previously been reported. Due to the differences in the abundance of various

morphologies, it was difficult to determine the expression of the least populated phenotypes

using bulk RNA-sequencing. Nonetheless, phenotypic and genotypic characterization suggest

that our culture may contain alveolar type 2 (AT2) cells. AT2 cells are responsible for expression

of surfactant proteins in the lungs and differentiation into AT1 cells which cover more than 95%

of the alveolar surface area and are crucial for gas-exchange (Barkauskas et al., 2013; Wang et

al., 2018b). Lung organoid models have recently helped uncover cell pathways critical during

lung repair and regeneration and to identify damage-associated transient progenitors (DATPs)

which represent a distinct population of AT2-lineage cells (Choi et al., 2020). Having both

bronchial epithelial cells and AT2 cell types present in our lung organoids increases the number

of future potential applications of the organoids. It has been shown that the lung arises from

cells expressing the NKX2-1 transcription factor (Kaushik et al., 2018), which is upregulated in

our lung organoids. For example, lung organoids have previously been used in the Transwell

system for studying viral uptake into cells (Zhou et al., 2018). The use of a lung organoid model

derived from human pluripotent stem cells showed that AT2-like cells are susceptible to SARS-

CoV-2 infection, and infection of organoids resulted in the upregulation of chemokines similar to

that reported in patients with COVID-19 (Han et al., 2021). Similarly, canine lung organoids

could be used for in-depth pathophysiology studies of viruses causing canine infectious

respiratory disease (CIRD) complex, including canine parainfluenza virus (CPIV), canine

adenovirus (CAV) type 2, and canine herpesvirus (Reagan and Sykes, 2020).

Urinary Bladder

Canine bladder cancer organoids were previously described and exposed to anticancer drugs to

describe their potential role in research and precision medicine (Elbadawy et al., 2019). Since

canine bladder cancer is a well-established model for human muscle-invasive bladder cancer

(Knapp et al., 2020), canine bladder cancer organoids represent a valuable model for

translational preclinical research (Minkler et al., 2021). In addition, Elbadawy et al. has recently

described healthy canine bladder organoids (Elbadawy et al., 2022). This report expands the

knowledge and accessibility of healthy canine bladder organoids, with those described here

displaying a similar morphological phenotype to those previously described (Elbadawy et al.,

2022). Using 3D patient-derived tumor organoids to predict the response to chemotherapeutic

protocols has great potential in oncological precision medicine. Therefore, there is a need for

healthy canine bladder organoids to serve as controls when attempting to identify novel

therapeutic strategies (Yu et al., 2021).

Liver

Canine liver-derived organoids have been previously described from both normal and

COMMD1-deficient dogs and were cultured to model copper storage disease, which is also

known as Wilson's disease in humans (Nantasanti et al., 2015). Our group has recently further

standardized the protocol for canine hepatic organoid culture (Gabriel et al., 2022a). Based on

the characterization outlined in that publication, our canine liver organoid culture is mainly

comprised of differentiated cholangiocytes. The previous study describing canine hepatic

organoids in expansion media showed that the organoids minimally expressed the mature

hepatocyte (CYP3A12) marker while stably expressing the following markers: stem cell (CD133

and LGR5), cholangiocyte (KRT19 and SOX9), and early hepatocyte (FOXA1 and HNF4α)

(Nantasanti et al., 2015). Future efforts could involve the development of media compositions,

including growth factors that will enhance the differentiation of hepatic stem cells into mature,

differentiated hepatocytes rather than cholangiocytes, which were first described in murine liver

organoids (Huch et al., 2013), then first attempted in dogs (Nantasanti et al., 2015), further

refined in dogs (Kruitwagen et al., 2020). Our group has recently investigated the ability of

canine liver-derived organoid cultures to differentiate into mature hepatocytes by comparing six

different media compositions (Gabriel et al., 2024). Optimization of such differentiation media

could open avenues to explore their usefulness for hepatic toxicity assays in drug research, in

addition to modeling various analogous cholangiopathies and hepatocellular diseases in

canines.

CONCLUSION

Applications of organoid technologies are rapidly expanding and now encompass protocols to

develop reliable *in vitro* models of various diseases. Further differentiation or enrichment of

certain cell populations within the organoids characterized here is warranted to expand the

current scope of applications for canine organoids. We report the successful isolation, culture,

and characterization of two novel canine organoid lines. These novel organoid lines will

enhance future use of the technology in fields including drug development, clinical applications,

and personalized medicine applications. Furthermore, a multi-tissue comparison of five canine

organoid lines derived from two genetically related individuals allowed for direct evaluation of

inter-organ and inter-individual variance in both in vivo and in vitro gene expression. The

antibodies optimized in this study for the canine model can be used in differentiation

experiments attempting to enrich specific cell types such as differentiated hepatocytes and

pancreatic beta cells. Future directions, including derivation of organoids from adult and

diseased canines and further characterization utilizing single-cell RNA sequencing, will help

identify unique subpopulations of cells within the organoids, further increasing the applicability of

these new translational *in vitro* models.

MATERIALS AND METHODS

Tissue collection

Dogs were used under permit (ref. IACUC-18-065), and proper Institutional Animal Care and

Use Committee (IACUC) protocols were followed. For this study, two 4-week-old intact female

canines were euthanized via intravenous sodium pentobarbital overdose due to unrelated

reasons, and tissues were quickly harvested (donor details in **Supplemental file 3**).

Approximately 2 cm x 2 cm tissue biopsies were obtained and then rinsed three times in 10 mL

of 1X Complete Chelating Solution (1X CCS, composition and further details can be found in

Gabriel et al. 2022) and transferred to 6 mL of Dulbecco's Modified Eagle Medium/Nutrient

Mixture F-12 (Advanced DMEM/F12) with the addition of Pen Strep (Gabriel et al., 2022a).

Organoid isolation and cultivation

Organoid isolation and maintenance were based on a modified protocol described by Saxena et

al. in 2016, which was optimized to include the standardized culture, expansion, and harvesting

of canine intestinal and hepatic organoids in Gabriel et al. 2022 (Gabriel et al., 2022a; Saxena

et al., 2016). A subset of the tissue pieces was minced with a scalpel until a consistency was

achieved that would fit into a 10 mL pipette, at which point the samples underwent the typical

canine organoid isolation protocol. Samples were washed with 5 mL of 1X complete chelating

solution (CCS) then vortexed. After the tissue settled, supernatant was removed down to the 5

mL mark and a total of five washes were done. During the last two washes, supernatant was

removed down to the 3 mL mark. Next, 3 mL of 1X CCS containing the tissue sample was

transferred to a 6 well plate, the sample tube was rinsed with an additional 3 mL of 1X CCS and

transferred to the 6 well plate. Then, 150 µL of 0.5 M EDTA (Invitrogen, ref. 15575-038) was

added and the plate was incubated at 4°C while rocking for 10 min. The sample was then

transferred to a tube containing 5 mL of CCS and 2 mL of FBS and inverted. Next, the

supernatant and ~100-200 µL of tissue was transferred to an empty tube and centrifuged at 700

g for 5 min at 4°C. Supernatant was discarded and the sample was rinsed with 6 mL of DMEM,

before again spinning and removing supernatant. Samples were then mixed in either of two

Matrigel compositions Phenol red-free (Corning ref. 356231) or Phenol red (Corning ref.

356230) and plated in 30 µL drops in 24 well plates. Plates were incubated for ~20 minutes at

37°C to solidify the Matrigel. Samples were expanded in our growth media (CMGF+ R/G), which

is supplemented with Y-27632 ROCK inhibitor (Biogems, ref. 1293823) and a GSK3β inhibitor,

Stemolecule CHIR99021 (Stemgent, ref. 04-0004). Total media volumes typically consisted of

500 µL on Monday and Wednesday, and 750 µL on Friday, any deviations are listed in

Supplemental file 1. Passaging was done as previously described, through the addition of 500

µL TrypLE Express to 500 µL of DMEM and organoids which was then incubated in a 37°C heat

bath for 10 minutes. Dissociation was stopped by dilution of the TrypLE Express with 6 mL of

DMEM which was then centrifuged and removed. Cleaning of the organoids was used to

replace Matrigel or change the density of the organoids in the culture. Before harvesting for

characterization, both Y-27632 ROCK inhibitor and GSK3β inhibitor were retracted (CMGF+) for

five days to discourage the culture from stem cell expansion and allow time for potential

differentiation of cell lines (see **Supplemental file 2** for complete media details). Samples used

for paraffin embedding were ensured to be plated in Phenol red-free Matrigel.

Cryopreservation of Organoids

Two freezing medias were used to cryopreserve organoids which consisted of (1) 50% CMGF+

R/G, 40% FBS, and 10% DMSO as well as (2) Cryostor CS10 (BioLife Solutions; 210102).

Recovery after Cryostor CS10 was more reliable and thus was favored. Prior to freezing,

organoids were recovered from Matrigel and resuspended in an appropriate freezing media.

After being placed in a 1 mL cryovial, samples were placed in the fridge for 10 minutes, then

moved to a -80°C freezer overnight in a Mr. Frosty container (Nalgene; 5100-0001) filled with

isopropanol, and finally stored in liquid nitrogen (-196°C) indefinitely.

RNA extractions and sequencing

After isolation, expansion, and differentiation (between 17 and 31 days), organoids were

pelleted and resuspended in 100 µL of Phosphate Buffered Saline (PBS) and transferred to a

cryovial. The sample tube was flushed with 900 µL of RNAlater and subsequently added to the

cryovial before being stored in liquid nitrogen (-196°C). Tissue biopsies were directly placed into

cryovials containing 1 mL of RNAlater and stored in liquid nitrogen. Upon thawing, tissue

samples were quickly rinsed in PBS to remove excess salts from the RNAlater solution and

were immediately transferred to 800 µL of Trizol and homogenized with a pestle. Organoid

samples were thawed and transferred to a 15 mL tube with 2 mL of PBS, then centrifuged at

1,200 g at 4°C for 5 min to pellet the organoids. RNAlater was removed, and 1 mL of Trizol was

added to the organoids and homogenized via brief vortexing.

After homogenizing, samples were stored at room temperature for 5 minutes and then

centrifuged at 12,000 g at 4°C for 10 min to eliminate debris and polysaccharides. The

supernatant was transferred to a new tube, and chloroform (0.2 mL chloroform per mL Trizol)

was added. Samples were shaken vigorously for 20 sec and stored at room temperature for 2-3

minutes before being centrifuged at 10,000 g at 4°C for 18 minutes. The aqueous phase was

transferred to a sterile 1.5 mL Rnase-free tube. Then an equal volume of 100% RNA-free EtOH

was slowly added and mixed before being transferred to a Qiagen Rneasy column (Rneasy Mini

kit) seated in a collection tube which was centrifuged for 30 seconds at 8,000 g. Flow-through

was discarded, and the Qiagen Dnase treatment protocol was followed. Next, 500 µL of buffer

RPE was added and centrifuged for 30 seconds at 8,000 g. Flow-through was again discarded,

and 500 µL of buffer RPE was added and centrifuged for 2 minutes at 8,000 g. Flow-through

was discarded, and columns were centrifuged for 1 minute at 8,000 g to remove the remaining

buffer. RNA was eluted in 50 μL of Rnase-free water and allowed to sit for 2 minutes before

being centrifuged for 1 minute at 8,000 g. Samples were centrifuged again at 8,000 g,

immediately analyzed on a Nanodrop, and frozen at -80°C.

Prior to library preparation, RNA samples were quantified with an Agilent 2100 Bioanalyzer

(Eukaryotic Total RNA Nano). Further quantification was done by GENEWIZ using a Qubit 2.0

Fluorometer (ThermoFisher Scientific) and a 4200 Tapestation (Agilent). An ERCC RNA Spike-

In Mix kit (ThermoFisher Scientific cat. 4456740) was used to normalize total RNA prior to

library preparation. A NEBNext Ultra II RNA Library Prep Kit for Illumina (New England Biolabs,

Ipswich, MA, USA) was used for library preparation. mRNAs were initially enriched with

Oligod(T) beads and then fragmented for 15 minutes at 94°C. Next, first and second-strand

cDNA was synthesized, end-repaired, and adenylated at 3'ends, and universal adapters were

ligated to cDNA fragments. This was followed by index addition and library enrichment by PCR

with limited cycles. Libraries were validated on the Agilent TapeStation (Agilent Technologies,

Palo Alto, CA, USA) and quantified using a Qubit 2.0 Fluorometer (ThermoFisher Scientific,

Waltham, MA, USA) as well as by quantitative PCR (KAPA Biosystems, Wilmington, MA, USA).

The libraries were multiplexed and clustered onto two flowcells and were loaded onto an

Illumina HiSeq 4000 instrument. The samples were sequenced using a 2x150bp Paired-End

(PE) configuration. The HiSeq Control Software (HCS) conducted image analysis and base

calling. Raw sequence data (.bcl files) generated from Illumina HiSeq was converted into fastq

files and de-multiplexed using Illumina bcl2fastq 2.20 software with one mismatch allowed for

index sequence identification.

RNA sequencing

The total number of reads from tissues and organoids ranged from ~18×10^6 to 27×10^6

(**Supplemental file 4**). The average Phred quality score was 35 before quality control

procedures (see *Processing, mapping, and quantification of RNA-seq libraries*). Comparisons

were made between organoids and their native tissues, across organoids, and across tissues,

for both B816 and B818 individuals.

Processing, mapping, and quantification of RNA-seq libraries

Raw sequence files were inspected in FastQC v0.11 and MultiQC v1.7 (Kim et al., 2015; Li et

al., 2009) to verify their quality. Barcodes were trimmed from reads and reads with a quality

score < 20 were discarded from downstream analysis cutadapt v3.5 (Martin, 2011). The data

set was de-duplicated with BBDuK v38.94 (https://sourceforge.net/projects/bbmap/) with a

search k-mer size of 18bp. The resulting reads were passed to SortMeRNA v2.1 (Kopylova et

al., 2012) to filter out rRNA sequences based on similarity with the SILVA v111 and Rfam v11.0

databases (Gardner et al., 2009; Quast et al., 2013). After each step, reads were inspected with

FastQC and MultiQC to ensure the quality of the data. Prior to the alignment of reads to a dog

genome with STAR v2.5, an index was created from ROS_Cfam_1.0 (RefSeq:

GCF_014441545). In average for all samples, 90.5% of the reads mapped to unique targets

within the reference genome (**Supplemental file 4**). Sequences from the ERCC spike-in

controls were included in this index to quantify their abundances in the samples. The resulting

BAM files were passed to Subread v1.6 to obtain gene-level counts via the featureCounts

algorithm.

Differential gene expression analysis

Gene counts mapped to ERCC spike-in controls by featureCounts were extracted. Then, we

calculated library size scaling factors based solely on ERCC counts using edgeR v3.36

(Robinson et al., 2009) as implemented in R v4.1 (Team, 2013) and using the trimmed mean of

M-values (TMM) method (Robinson and Oshlack, 2010) to normalize the ERCC counts. The

scaling factors were used to normalize the gene counts and calculate log2-transformed counts

per million (CPM) after adding a 0.5 as a constant to all the values.

Our goal was to detect differences in gene expression between extracted tissues and the

corresponding organoids. Prior to differential gene expression analysis, visualization of

transcriptional variance was explored with principal component analysis and multidimensional

scaling. This unsupervised analysis revealed similarity between samples as expected by tissue-

of-origin (endometrium, lung, pancreas, bladder, or liver) and type (extracted or organoid). No

obvious outliers were detected during this exploratory analysis. The model under testing was

expr=β_1+β_2 x organoid, with type indicating if the sample was an organoid or not. Gene-wise

dispersions were estimated, and outlier effects were reduced with the estimateDisp function in

edgeR (using the robust=T option). Negative binomial generalized linear models (GLM) were

fitted for each gene, and statistical significance for the difference in mean expression was

obtained by performing Bayes quasi-likelihood F-tests (glmQLFTest function in edgeR).

Visualization of the results via heatmaps and Venn diagrams were generated via the

ComplexHeatmap (Gu et al., 2016) and VennDiagram (Chen and Boutros, 2011) R packages.

Genes unique to each organ are listed in **Supplemental file 5** and **Supplemental file 6**.

Transcriptional Pathways for both tissues and organoids were analyzed by conducting single-

sample Gene Set Enrichment Analysis (ssGSEA) using the VIPER algorithm (Alvarez et al.,

2016). This analysis focused on 50 Hallmark gene sets which were obtained from the Molecular

Signatures Database (MsigDB) (Liberzon et al., 2015). Scaled pathway enrichment scores were

converted to z-scores and was visualized and clustered using the gplots R package. All data

and analysis code has been made publicly available on a GitHub repository.

Paraffin embedding and immunohistochemistry

After organoids were expanded, they were then allowed to grow in CMGF+ for five days, media

was removed, and 500 µL of Formalin-acetic acid-alcohol (FAA, composition in Gabriel et al.

2022) was added to each well (Gabriel et al., 2022a). After 24 hours, FAA was replaced with

70% ethanol and samples were paraffin-embedded and mounted on slides at the Iowa State

University Histopathology laboratory. Tissues were fixed in paraformaldehyde and paraffin-

embedded according to standard histology procedures. Tissues and organoids were stained

with hematoxylin and eosin (H&E).

For immunohistochemistry, samples were deparaffinized and rehydrated through a series of

alcohol changes to deionized water. Endogenous peroxidase within the samples was then

quenched using a hydrogen peroxide bath. Heat induced epitope retrieval was performed using

either a tris-EDTA or citrate buffer. Immunohistochemistry (IHC) antibodies for Pan cytokeratin

(Agilent, M0821), smooth actin (BioGenex, MU128-UC) and vimentin (Agilent, M0725) were

used on both tissues and organoids. An indirect method of IHC staining was then carried out

using a biotinylated secondary antibody followed by a streptavidin. The samples were then

incubated with NovaRED™ (Vector, SK-4800) chromogen, counterstained with hematoxylin,

and dehydrated. Light microscopy images were taken on a Leica Aperio GT 450 Scanner and

analyzed with ImageScope (v12.4.3.5008) or on an Olympus BX40 light microscope.

Immunofluorescence

For deparaffinization, slides were placed in xylene twice for ten minutes, then transferred to

100% ethanol twice for one minute with regular agitation. After the last alcohol wash, slides

were laid on tissue paper for five minutes to dry. After deparaffinization, tissues and organoids

underwent Heat Induced Epitope Retrieval (HIER) with either Citrate buffer (pH 6) or a

Tris/EDTA buffer (pH 9) using a HybEZ II Oven at 75°C for two hours. After two hours, the tray

was taken out of the oven, and the slides were allowed to cool with the lid off for 15 minutes.

Once cool, the slides were rinsed in PBS twice for two minutes each, then rinsed in PBS for ten

minutes. The tissues and organoids were permeabilized by incubation in 0.25% Triton in PBS

twice for ten minutes each. After three PBS rinses, both tissues and organoids were blocked in

Casein in PBS for one hour at room temperature. Tissues and organoids were incubated in a

humidity chamber with their primary antibody overnight at 4°C at the appropriate concentration.

The next day, the slides were again rinsed in PBS prior to the secondary being added at 1:1000

in PBS for one hour at room temperature, with slides being rinsed again. The slides were then

incubated with DAPI (Sigma, D9542-1MG) at 1:500 in PBS for twenty minutes, and washed

three times for ten minutes in PBS, then switched to distilled water. Fluoroshield (Sigma, F6182-

20ML) was used to mount the slides, and after drying overnight, the slides were imaged on a

Stellaris confocal microscope. Antibodies, dilutions, and antigen retrieval techniques for each

tissue type can be seen in **Supplemental file 7**. Scale bars were added to immunofluorescent

images using Leica LAS AF Lite (v. 2.6.0 build 7266).

DATA AVAILABILITY

The RNA-seq raw reads generated in this study are available in the Sequence Read Archive

(NCBI-SRA BioProject PRJNA847879) as well as the aligned files being available on the NIH

ICDC (https://caninecommons.cancer.gov/#/study/ORGANOIDS01). The bioinformatic scripts

are available on Github (https://github.com/chris-zdyrski/Novel_Canine_Organoids).

ACKNOWLEDGEMENTS

We are grateful for startup funds received from Iowa State University as well as a generous

donation from the Armbrust family through the Iowa State University Foundation. We would like

to thank Dr. Jodi Smith for the use of her colony. We are also appreciative of Dr. Adrien

Aertsens and Sichao Mao for their help with tissue collection. We thank Dr. Nicole Valenzuela

for lending access to Iowa State University's computing resources. We thank Akhil

Vinithakumari for assistance with graphing of data, Ahona Mukherjee for assisting with

immunofluorescence, and Dr. David Meyerholz for marker recommendations. We appreciate the

timely processing of samples by the Iowa State University Pathology Department and the Iowa

State University Histopathology Department.

FIGURE LEGENDS

624
Figure 1. Morphological and histological characterization of canine organoid lines
derived from a single donor.
Bright field and hematoxylin & eosin (H&E) images and approximate proportions of the cultures
for the five organoid cell lines derived from the uterus (red=80%, yellow=20%), lung (red=60%,
yellow=35%, orange=5%), pancreas (red=50%, yellow=50%), bladder (red=100%), and liver
(red=80%, yellow=20%) of canine individual B816. Red, yellow, and orange arrows indicate
distinct morphologies in each organoid line while blue, green, and black arrows indicate similar
histological areas of the organoids and tissues. Structures identified in tissues with histological
similarities in organoids were found in endometrium (blue=glandular epithelial cells,
green=endometrial epithelial cells), lung (blue=alveolar type-1 cells, green=bronchial epithelial
cells, black=alveolar type-2 cells), pancreas (blue=endocrine cells, green=intercalated ducts),
bladder (blue=transitional epithelium), and liver (blue=cholangiocyte) organoids. Images of
organoid cultures were captured using a Leica Dmi1 microscope. Scale bars are provided at 5X
(500 μm), 10X (200 μm), and 40X (60 μm) magnifications.

639
Figure 2. Immunohistochemistry comparison of tissues and organoids.
Immunohistochemistry staining of organoids and tissues for both donors, B816 and B818. Scale
bars for each image are displayed in µm. Antibodies for pan cytokeratin, smooth muscle actin,
and vimentin were used. Images of negative control slides are provided in **Figure 2-figure**
supplement 1.

Figure 3. Expression of RNA and identification of tissue-specific markers for both organoids and tissues.
(A) RNA heatmaps of the differentially expressed (DE) genes (FWER < 0.05) between tissues and organoids of the same organs. Tissue-specific markers were identified across the five tissues for both (B) organoids (FWER < 0.05) and (C) tissues (FWER < 0.05). Upregulated expression is red, white is neutral, and blue represents suppressed expression. (D) The ten most highly expressed tissue-specific genes from two genetically related donors for both organoids and tissues, as well as genes in common between organoids and tissues are denoted in **bold**.

655
Figure 4. Comparison of mRNA expression similarity between organoids and tissue
samples for each organ.
Venn diagrams of genes expressed from both donors (B816 and B818) comparing (A)
organoids and (B) tissues from the same organs. (C) Venn diagrams showing the comparison of
mRNA expression between organoids and tissues from each organ. (D) Principal component
analysis (PCA) plots of mRNA expression across all organoid and tissue samples. (E) PCA
plots for either organoid or tissue samples. Tissue types are color coded in the legend.
663

665
Figure 5. Protein characterization across tissues and organoids.
Immunofluorescent staining of organoids and tissues for both donors, B816 and B818.
Antibodies are labeled in the channel they appear in, and images are merged with DAPI (blue).
Images were taken at either 20X or 60X magnification, scale bars are 100 µm. A green
background represents positive expression of the marker while a red background indicates the
lack of expression in the current field. Immunofluorescent details are listed in **Supplemental file
7**, and negative control images taken under the same conditions are shown in **Figure 5-figure
supplement 1**.

674
Figure 6. Characterization of organoids, pathway enrichment analysis, and potential
biomedical applications.
(A) Initial characterization and preservation methods used for the intra-donor derived canine
organoid cell lines. (B) A list of the top pathways when comparing organoids to tissues. (C) A list
of potential biomedical applications for canine organoid models. (D) Analysis of proliferation and
inflammation pathways shown across all organoids and tissues. (E) Enriched pathways for
either organoids or tissues (red = upregulated, blue = downregulated).

682
Figure 1-figure supplement 1. Morphogenesis of organoids over time.
Light microscopy images of organoids derived from B816 tissues over time. Images spanned
from two days after isolation, through a passage, to the earliest harvesting, day 17. Images
were taken at 5X magnification, scale bars are 500 μm.

687
Figure 2-figure supplement 1. Immunohistochemistry comparison of tissues and organoids.
Negative control images for immunohistochemistry staining of organoids and tissues for both donors, B816 and B818. Images were taken at 20X magnification, scale bars are in μm.

692
Figure 5-figure supplement 1. Protein characterization across tissues and organoids.
Immunofluorescent staining of the negative control slides for organoids and tissues for both
donors, B816 and B818. Images were merged with DAPI (blue) and were taken at either 20X or
60X magnification, scale bars are 100 μm.

697 **Supplemental file 1.** Growth chart for all organoid samples derived from two dogs. Descriptions
698 include media changes, passages attempted, and total days in culture.
699
700 **Supplemental file 2.** Composition of CMGF+ culture media without the addition of RG.
701
702 **Supplemental file 3.** Donor information including age, sex, and breed from the two donors,
703 B816 and B818.
704
705 **Supplemental file 4.** Average sequence length, total reads, quality scores, and percent of
706 mapped reads obtained from the RNA-seq samples of organoids and tissues. Numbers are
707 shown for the data set before and after quality control procedures (read trimming, deduplication,
708 and filtering).
709
710 **Supplemental file 5.** The gene symbol, logCPM, and logFC values for each combination of
711 organoids grown regarding the RNA-seq analysis. The logCPM represent the log-counts per
712 million which are normalized expression counts. The logFC=Log fold change which compares
713 the level of expression between two treatments.
714
715 **Supplemental file 6.** The gene symbol, logCPM, and logFC values for each combination of
716 tissues harvested regarding the RNA-seq analysis. The logCPM represent the log-counts per
717 million which are normalized expression counts. The logFC=Log fold change which compares
718 the level of expression between two treatments.
719
720 **Supplemental file 7.** Antibody details, concentrations of primary and secondary antibodies, and
721 antigen retrieval techniques used.
722
723 **Supplemental file 8.** Pathway analysis results for organoid and tissue samples from both B816
724 and B818. Red indicates upregulated while blue indicates downregulated pathways.
725

LITERATURE CITED

Adin, C.A., and Gilor, C. (2017). The Diabetic Dog as a Translational Model for Human Islet

Transplantation.

Aktas, R.G., Karski, M., Issac, B., Sun, L., Rockowitz, S., Sliz, P., and Vakili, K. (2022). Long-

Term Characteristics of Human-Derived Biliary Organoids under a Single Continuous Culture

Condition. Cells *11*. https://doi.org/10.3390/cells11233797.

Alvarez, M.J., Shen, Y., Giorgi, F.M., Lachmann, A., Ding, B.B., Hilda Ye, B., and Califano, A.

(2016). Functional characterization of somatic mutations in cancer using network-based

inference of protein activity. Nat Genet *48*, 838–847. https://doi.org/10.1038/ng.3593.

Artegiani, B., Hendriks, D., Beumer, J., Kok, R., Zheng, X., Joore, I., Chuva de Sousa Lopes, S.,

van Zon, J., Tans, S., and Clevers, H. (2020). Fast and efficient generation of knock-in human

organoids using homology-independent CRISPR–Cas9 precision genome editing. Nat Cell Biol

22, 321–331. https://doi.org/10.1038/s41556-020-0472-5.

Bailey, K.L., and Carlson, M.A. (2019). Porcine models of pancreatic cancer. Front Oncol *9*.

https://doi.org/10.3389/fonc.2019.00144.

Barkauskas, C.E., Cronce, M.J., Rackley, C.R., Bowie, E.J., Keene, D.R., Stripp, B.R., Randell,

S.H., Noble, P.W., and Hogan, B.L.M. (2013). Type 2 alveolar cells are stem cells in adult lung.

Journal of Clinical Investigation *123*, 3025–3036. https://doi.org/10.1172/JCI68782.

Bartel, C., Tichy, A., Schoenkypl, S., Aurich, C., and Walter, I. (2013). Effects of steroid

hormones on differentiated glandular epithelial and stromal cells in a three dimensional cell

culture model of the canine endometrium. BMC Vet Res *9*. https://doi.org/10.1186/1746-6148-9-

86.

Bartfeld, S., and Clevers, H. (2017). Stem cell-derived organoids and their application for

medical research and patient treatment. J Mol Med *95*, 729–738.

https://doi.org/10.1007/s00109-017-1531-7.

Birey, F., Andersen, J., Makinson, C.D., Islam, S., Wei, W., Huber, N., Fan, H.C., Metzler,

K.R.C., Panagiotakos, G., Thom, N., et al. (2017). Assembly of functionally integrated human

forebrain spheroids. Nature *545*, 54–59. https://doi.org/10.1038/nature22330.

Boretto, M., Cox, B., Noben, M., Hendriks, N., Fassbender, A., Roose, H., Amant, F.,

Timmerman, D., Tomassetti, C., Vanhie, A., et al. (2017). Development of organoids from

mouse and human endometrium showing endometrial epithelium physiology and long-term

expandability. Development (Cambridge) *144*, 1775–1786. https://doi.org/10.1242/dev.148478.

Brancato, V., Oliveira, J.M., Correlo, V.M., Reis, R.L., and Kundu, S.C. (2020). Could 3D

models of cancer enhance drug screening? Biomaterials *232*.

https://doi.org/10.1016/j.biomaterials.2019.119744.

Chakrabarti, J., Koh, V., So, J.B.Y., Yong, W.P., and Zavros, Y. (2021). A Preclinical Human-

Derived Autologous Gastric Cancer Organoid/Immune Cell Co-Culture Model to Predict the

Efficacy of Targeted Therapies. Journal of Visualized Experiments *2021*.

https://doi.org/10.3791/61443.

Chandler, M., Cunningham, S., Lund, E.M., Khanna, C., Naramore, R., Patel, A., and Day, M.J.

(2017). Obesity and Associated Comorbidities in People and Companion Animals: A One Health

Perspective. J Comp Pathol *156*, 296–309. https://doi.org/10.1016/j.jcpa.2017.03.006.

Chandra, L., Borcherding, D.C., Kingsbury, D., Atherly, T., Ambrosini, Y.M., Bourgois-Mochel,

A., Yuan, W., Kimber, M., Qi, Y., Wang, Q., et al. (2019). Derivation of adult canine intestinal

organoids for translational research in gastroenterology. BMC Biol *17*, 1–21.

https://doi.org/10.1186/s12915-019-0652-6.

Chang, H.M., Martinez, N.J., Thornton, J.E., Hagan, J.P., Nguyen, K.D., and Gregory, R.I.

(2012). Trim71 cooperates with microRNAs to repress Cdkn1a expression and promote

embryonic stem cell proliferation. Nat Commun *3*. https://doi.org/10.1038/ncomms1909.

Chen, H., and Boutros, P.C. (2011). VennDiagram: A package for the generation of highly-

customizable Venn and Euler diagrams in R. BMC Bioinformatics *12*.

https://doi.org/10.1186/1471-2105-12-35.

Choi, J., Park, J.E., Tsagkogeorga, G., Yanagita, M., Koo, B.K., Han, N., and Lee, J.H. (2020).

Inflammatory Signals Induce AT2 Cell-Derived Damage-Associated Transient Progenitors that

Mediate Alveolar Regeneration. Cell Stem Cell *27*, 366-382.e7.

https://doi.org/10.1016/j.stem.2020.06.020.

Cindrova-Davies, T., Zhao, X., Elder, K., Jones, C.J.P., Moffett, A., Burton, G.J., and Turco,

M.Y. (2021). Menstrual flow as a non-invasive source of endometrial organoids. Commun Biol *4*.

https://doi.org/10.1038/s42003-021-02194-y.

DiMasi, J.A., Grabowski, H.G., and Hansen, R.W. (2016). Innovation in the pharmaceutical

industry: New estimates of R&D costs. J Health Econ *47*, 20–33.

https://doi.org/10.1016/j.jhealeco.2016.01.012.

Dongeun, H., D, M.B., Akiko, M., Martín, M.-Z., Yuan, H.H., and E, I.D. (2010). Reconstituting

Organ-Level Lung Functions on a Chip. Science (1979) *328*, 1662–1668.

https://doi.org/10.1126/science.1188302.

Dost, A.F.M., Moye, A.L., Vedaie, M., Tran, L.M., Fung, E., Heinze, D., Villacorta-Martin, C.,

Huang, J., Hekman, R., Kwan, J.H., et al. (2020). Organoids Model Transcriptional Hallmarks of

Oncogenic KRAS Activation in Lung Epithelial Progenitor Cells. Cell Stem Cell *27*, 663-678.e8.

https://doi.org/10.1016/j.stem.2020.07.022.

Elbadawy, M., Usui, T., Mori, T., Tsunedomi, R., Hazama, S., Nabeta, R., Uchide, T.,

Fukushima, R., Yoshida, T., Shibutani, M., et al. (2019). Establishment of a novel experimental

model for muscle-invasive bladder cancer using a dog bladder cancer organoid culture. Cancer

Sci *110*, 2806–2821. https://doi.org/10.1111/cas.14118.

Elbadawy, M., Fujisaka, K., Yamamoto, H., Tsunedomi, R., Nagano, H., Ayame, H., Ishihara, Y.,

Mori, T., Azakami, D., Uchide, T., et al. (2022). Establishment of an experimental model of

normal dog bladder organoid using a three-dimensional culture method. Biomedicine &

Pharmacotherapy *151*, 113105. https://doi.org/10.1016/j.biopha.2022.113105.

Fatehullah, A., Tan, S.H., and Barker, N. (2016). Organoids as an in vitro model of human

development and disease. Nat Cell Biol *18*, 246–254. https://doi.org/10.1038/ncb3312.

Gabriel, V., Zdyrski, C., Sahoo, D.K., Dao, K., Bourgois-Mochel, A., Kopper, J., Zeng, X.-L.,

Estes, M.K., Mochel, J.P., and Allenspach, K. (2022a). Standardization and Maintenance of 3D

Canine Hepatic and Intestinal Organoid Cultures for Use in Biomedical Research. Journal of

Visualized Experiments 1–28. https://doi.org/10.3791/63515.

Gabriel, V., Zdyrski, C., Sahoo, D.K., Dao, K., Bourgois-Mochel, A., Atherly, T., Martinez, M.N.,

Volpe, D.A., Kopper, J., Allenspach, K., et al. (2022b). Canine Intestinal Organoids in a Dual-

Chamber Permeable Support System. Journal of Visualized Experiments

https://doi.org/10.3791/63612.

Gabriel, V., Lincoln, A., Zdyrski, C., Ralston, A., Wickham, H., Honold, S., Ahmed, B.H.,

Paukner, K., Feauto, R., Merodio, M.M., et al. (2024). Evaluation of different media

compositions promoting hepatocyte differentiation in the canine liver organoid model. Heliyon

10. https://doi.org/10.1016/j.heliyon.2024.e28420.

Gao, Y., Guan, W., and Bai, C. (2022). Pancreatic Duct Cells Isolated From Canines

Differentiate Into Beta-Like Pancreatic Islet Cells. Front Vet Sci *8*.

https://doi.org/10.3389/fvets.2021.771196.

García-Rodríguez, I., Sridhar, A., Pajkrt, D., and Wolthers, K.C. (2020). Put Some Guts into It:

Intestinal Organoid Models to Study Viral Infection. Viruses *12*, 1288.

https://doi.org/10.3390/v12111288.

Gardner, P.P., Daub, J., Tate, J.G., Nawrocki, E.P., Kolbe, D.L., Lindgreen, S., Wilkinson, A.C.,

Finn, R.D., Griffiths-Jones, S., Eddy, S.R., et al. (2009). Rfam: Updates to the RNA families

database. Nucleic Acids Res *37*. https://doi.org/10.1093/nar/gkn766.

Gordon, I., Paoloni, M., Mazcko, C., and Khanna, C. (2009). The comparative oncology trials

consortium: Using spontaneously occurring cancers in dogs to inform the cancer drug

development pathway. PLoS Med *6*. https://doi.org/10.1371/journal.pmed.1000161.

Gu, Z., Eils, R., and Schlesner, M. (2016). Complex heatmaps reveal patterns and correlations

in multidimensional genomic data. Bioinformatics *32*, 2847–2849.

https://doi.org/10.1093/bioinformatics/btw313.

Han, Y., Duan, X., Yang, L., Nilsson-Payant, B.E., Wang, P., Duan, F., Tang, X., Yaron, T.M.,

Zhang, T., Uhl, S., et al. (2021). Identification of SARS-CoV-2 inhibitors using lung and colonic

organoids. Nature *589*, 270–275. https://doi.org/10.1038/s41586-020-2901-9.

Hasiwa, N. (2011). Critical evaluation of the use of dogs in biomedical research and testing in

Europe. ALTEX *28*, 326–340. https://doi.org/10.14573/altex.2011.4.326.

Hayashi, Y., Yamaguchi, J., Kokuryo, T., Ebata, T., Yokoyama, Y., and Nagino, M. (2018). Loss

of trefoil factor 1 inhibits biliary regeneration but accelerates the hepatic differentiation of

progenitor cells in mice. Biochem Biophys Res Commun *506*, 12–19.

https://doi.org/10.1016/j.bbrc.2018.10.023.

Hickman, J.A., Graeser, R., de Hoogt, R., Vidic, S., Brito, C., Gutekunst, M., van der Kuip, H.,

and Imi Predect consortium (2014). Three-dimensional models of cancer for pharmacology and

cancer cell biology: Capturing tumor complexity in vitro/ex vivo. Biotechnol J *9*, 1115–1128.

https://doi.org/10.1002/biot.201300492.

Huch, M., and Koo, B.K. (2015). Modeling mouse and human development using organoid

cultures. Development (Cambridge) *142*, 3113–3125. https://doi.org/10.1242/dev.118570.

Huch, M., Dorrell, C., Boj, S.F., Van Es, J.H., Li, V.S.W., Van De Wetering, M., Sato, T., Hamer,

K., Sasaki, N., Finegold, M.J., et al. (2013). In vitro expansion of single Lgr5 + liver stem cells

induced by Wnt-driven regeneration. Nature *494*, 247–250. https://doi.org/10.1038/nature11826.

Huch, M., Gehart, H., Van Boxtel, R., Hamer, K., Blokzijl, F., Verstegen, M.M.A., Ellis, E., Van

Wenum, M., Fuchs, S.A., De Ligt, J., et al. (2015). Long-term culture of genome-stable bipotent

stem cells from adult human liver. Cell *160*, 299–312. https://doi.org/10.1016/j.cell.2014.11.050.

Hynds, R.E., and Giangreco, A. (2013). Concise review: The relevance of human stem cell-

derived organoid models for epithelial translational medicine. Stem Cells *31*, 417–422.

https://doi.org/10.1002/stem.1290.

Jergens, A.E., and Simpson, K.W. (2012). Inflammatory bowel disease in veterinary medicine.

Front Biosci (Elite Ed) *4*, 1404–1419. https://doi.org/10.2741/e470.

Jones, R.C., Karkanias, J., Krasnow, M.A., Pisco, A.O., Quake, S.R., Salzman, J., Yosef, N.,

Bulthaup, B., Brown, P., Harper, W., et al. (2022). The Tabula Sapiens: A multiple-organ, single-

cell transcriptomic atlas of humans. Science (1979) *376*.

https://doi.org/10.1126/science.abl4896.

Kalma, Y., Granot, I., Gnainsky, Y., Or, Y., Czernobilsky, B., Dekel, N., and Barash, A. (2009).

Endometrial biopsy-induced gene modulation: first evidence for the expression of bladder-

transmembranal uroplakin Ib in human endometrium. Fertil Steril *91*.

https://doi.org/10.1016/j.fertnstert.2008.01.043.

Kampf, C., Mardinoglu, A., Fagerberg, L., Hallström, B.M., Edlund, K., Lundberg, E., Pontén, F.,

Nielsen, J., and Uhlen, M. (2014). The human liver-specific proteome defined by transcriptomics

and antibody-based profiling. FASEB Journal *28*, 2901–2914. https://doi.org/10.1096/fj.14-

250555.

Kaushik, G., Ponnusamy, M.P., and Batra, S.K. (2018). Concise Review: Current Status of

Three-Dimensional Organoids as Preclinical Models. Stem Cells *36*, 1329–1340.

https://doi.org/10.1002/stem.2852.

Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: A fast spliced aligner with low

memory requirements. Nat Methods *12*, 357–360. https://doi.org/10.1038/nmeth.3317.

Kim, E., Choi, S., Kang, B., Kong, J.H., Kim, Y., Yoon, W.H., Lee, H.R., Kim, S.E., Kim, H.M.,

Lee, H.S., et al. (2020a). Creation of bladder assembloids mimicking tissue regeneration and

cancer. Nature *588*, 664–669. https://doi.org/10.1038/s41586-020-3034-x.

Kim, J., Koo, B.K., and Knoblich, J.A. (2020b). Human organoids: model systems for human

biology and medicine. Nat Rev Mol Cell Biol *21*, 571–584. https://doi.org/10.1038/s41580-020-

0259-3.

Kinnear, S., Salamonsen, L.A., Francois, M., Harley, V., and Evans, J. (2019). Uterine SOX17:

a key player in human endometrial receptivity and embryo implantation. Sci Rep *9*.

https://doi.org/10.1038/s41598-019-51751-3.

Knapp, D.W., Dhawan, D., Ramos-Vara, J.A., Ratliff, T.L., Cresswell, G.M., Utturkar, S.,

Sommer, B.C., Fulkerson, C.M., and Hahn, N.M. (2020). Naturally-Occurring Invasive Urothelial

Carcinoma in Dogs, a Unique Model to Drive Advances in Managing Muscle Invasive Bladder

Cancer in Humans. Front Oncol *9*. https://doi.org/10.3389/fonc.2019.01493.

Kopper, J.J., Iennarella-Servantez, C., Jergens, A.E., Sahoo, D.K., Guillot, E., Bourgois-Mochel,

A., Martinez, M.N., Allenspach, K., and Mochel, J.P. (2021). Harnessing the Biology of Canine

Intestinal Organoids to Heighten Understanding of Inflammatory Bowel Disease Pathogenesis

and Accelerate Drug Discovery: A One Health Approach. Frontiers in Toxicology *3*.

https://doi.org/10.3389/ftox.2021.773953.

Kopylova, E., Noé, L., and Touzet, H. (2012). SortMeRNA: Fast and accurate filtering of

ribosomal RNAs in metatranscriptomic data. Bioinformatics *28*, 3211–3217.

https://doi.org/10.1093/bioinformatics/bts611.

Kruitwagen, H.S., Oosterhoff, L.A., van Wolferen, M.E., Chen, C., Nantasanti Assawarachan,

S., Schneeberger, K., Kummeling, A., van Straten, G., Akkerdaas, I.C., Vinke, C.R., et al.

(2020). Long-Term Survival of Transplanted Autologous Canine Liver Organoids in a COMMD1-

Deficient Dog Model of Metabolic Liver Disease. Cells *9*. https://doi.org/10.3390/cells9020410.

LeBlanc, A.K., Mazcko, C.N., and Khanna, C. (2016). Defining the Value of a Comparative

Approach to Cancer Drug Development. Clinical Cancer Research *22*, 2133–2138.

https://doi.org/10.1158/1078-0432.CCR-15-2347.

Lehmann, R., Lee, C.M., Shugart, E.C., Benedetti, M., Charo, R.A., Gartner, Z., Hogan, B.,

Knoblich, J., Nelson, C.M., and Wilson, K.M. (2019). Human organoids: A new dimension in cell

biology. Mol Biol Cell *30*, 1129–1137. https://doi.org/10.1091/mbc.E19-03-0135.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

and Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*,

2078–2079. https://doi.org/10.1093/bioinformatics/btp352.

Liberzon, A., Birger, C., Thorvaldsdóttir, H., Ghandi, M., Mesirov, J.P., and Tamayo, P. (2015).

The Molecular Signatures Database Hallmark Gene Set Collection. Cell Syst *1*, 417–425.

https://doi.org/10.1016/j.cels.2015.12.004.

Lim, S.Y., Nakamura, K., Morishita, K., Sasaki, N., Murakami, M., Osuga, T., Ohta, H.,

Yamasaki, M., and Takiguchi, M. (2014). Qualitative and quantitative contrast-enhanced

ultrasonographic assessment of cerulein-induced acute pancreatitis in dogs. J Vet Intern Med

28, 496–503. https://doi.org/10.1111/jvim.12319.

Lin, C., Yin, Y., Stemler, K., Humphrey, P., Kibel, A.S., Mysorekar, I.U., and Ma, L. (2013).

Constitutive β-catenin activation induces male-specific tumorigenesis in the bladder urothelium.

Cancer Res *73*, 5914–5925. https://doi.org/10.1158/0008-5472.CAN-12-4198.

Lu, J., Zhao, M., Wu, C., Chu, C., Zhang, C.Z., and Cao, Y. (2022). Comparison of RNAscope

and immunohistochemistry for evaluation of the UPK2 status in urothelial carcinoma tissues.

Diagn Pathol *17*. https://doi.org/10.1186/s13000-022-01191-x.

MacParland, S.A., Liu, J.C., Ma, X.Z., Innes, B.T., Bartczak, A.M., Gage, B.K., Manuel, J.,

Khuu, N., Echeverri, J., Linares, I., et al. (2018). Single cell RNA sequencing of human liver

reveals distinct intrahepatic macrophage populations. Nat Commun *9*.

https://doi.org/10.1038/s41467-018-06318-7.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing

reads. EMBnet J *17*, 10. https://doi.org/10.14806/ej.17.1.200.

Minkler, S., Lucien, F., Kimber, M.J., Sahoo, D.K., Bourgois-Mochel, A., Musser, M., Johannes,

C., Frank, I., Cheville, J., Allenspach, K., et al. (2021). Emerging roles of urine-derived

components for the management of bladder cancer: One man's trash is another man's treasure.

Cancers (Basel) *13*, 1–13. https://doi.org/10.3390/cancers13030422.

Mochel, J.P., and Danhof, M. (2015). Chronobiology and Pharmacologic Modulation of the

Renin-Angiotensin-Aldosterone System in Dogs: What Have We Learned? Rev Physiol

Biochem Pharmacol *169*, 43–69. https://doi.org/10.1007/112_2015_27.

Mochel, J.P., Jergens, A.E., Kingsbury, D., Kim, H.J., Martín, M.G., and Allenspach, K. (2018).

Intestinal Stem Cells to Advance Drug Development, Precision, and Regenerative Medicine: A

Paradigm Shift in Translational Research. AAPS J *20*, 17. https://doi.org/10.1208/s12248-017-

0178-1.

Mochel, J.P., Teng, C.H., Peyrou, M., Giraudel, J., Danhof, M., and Rigel, D.F. (2019).

Sacubitril/valsartan (LCZ696) significantly reduces aldosterone and increases cGMP circulating

levels in a canine model of RAAS activation. European Journal of Pharmaceutical Sciences

128, 103–111. https://doi.org/10.1016/j.ejps.2018.11.037.

Mohammadi, S., Morell-Perez, C., Wright, C.W., Wyche, T.P., White, C.H., Sana, T.R., and

Lieberman, L.A. (2021). Assessing donor-to-donor variability in human intestinal organoid

cultures. Stem Cell Reports *16*, 2364–2378.

https://doi.org/https://doi.org/10.1016/j.stemcr.2021.07.016.

Mollaki, V. (2021). Ethical challenges in organoid use. BioTech *10*.

https://doi.org/10.3390/biotech10030012.

Mullenders, J., de Jongh, E., Brousali, A., Roosen, M., Blom, J.P.A., Begthel, H., Korving, J.,

Jonges, T., Kranenburg, O., Meijer, R., et al. (2019). Mouse and human urothelial cancer

organoids: A tool for bladder cancer research. Proc Natl Acad Sci U S A *116*, 4567–4574.

https://doi.org/10.1073/pnas.1803595116.

Munsie, M., Hyun, I., and Sugarman, J. (2017). Ethical issues in human organoid and gastruloid

research. Development (Cambridge) *144*, 942–945. https://doi.org/10.1242/dev.140111.

Nantasanti, S., Spee, B., Kruitwagen, H.S., Chen, C., Geijsen, N., Oosterhoff, L.A., Van

Wolferen, M.E., Pelaez, N., Fieten, H., Wubbolts, R.W., et al. (2015). Disease modeling and

gene therapy of copper storage disease in canine hepatic organoids. Stem Cell Reports *5*, 895–

907. https://doi.org/10.1016/j.stemcr.2015.09.002.

Van Norman, G.A. (2016). Drugs, Devices, and the FDA: Part 1: An Overview of Approval

Processes for Drugs. JACC Basic Transl Sci *1*, 170–179.

https://doi.org/10.1016/j.jacbts.2016.03.002.

Ozawa, M., Inoue, M., Uchida, K., Chambers, J.K., Takeuch, Y., and Nakayama, H. (2019).

Physical signs of canine cognitive dysfunction. Journal of Veterinary Medical Science *81*, 1829–

1834. https://doi.org/10.1292/jvms.19-0458.

Perlman, R.L. (2016). Mouse Models of Human Disease: An Evolutionary Perspective. Evol

Med Public Health eow014. https://doi.org/10.1093/emph/eow014.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and

Glöckner, F.O. (2013). The SILVA ribosomal RNA gene database project: Improved data

processing and web-based tools. Nucleic Acids Res *41*. https://doi.org/10.1093/nar/gks1219.

Rawlings, T.M., Makwana, K., Tryfonos, M., and Lucas, E.S. (2021). Organoids to model the

endometrium: implantation and beyond. Reproduction and Fertility *2*, R85–R101.

https://doi.org/10.1530/raf-21-0023.

Reagan, K.L., and Sykes, J.E. (2020). Canine Infectious Respiratory Disease. Veterinary Clinics

of North America - Small Animal Practice *50*, 405–418.

https://doi.org/10.1016/j.cvsm.2019.10.009.

Robinson, M.D., and Oshlack, A. (2010). A scaling normalization method for differential

expression analysis of RNA-seq data. Genome Biol *11*. https://doi.org/10.1186/gb-2010-11-3-

r25.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2009). edgeR: A Bioconductor package for

differential expression analysis of digital gene expression data. Bioinformatics *26*, 139–140.

https://doi.org/10.1093/bioinformatics/btp616.

Russell, W.M.S., and Burch, R.L. (1959). The principles of humane experimental technique

(Methuen).

Sampaziotis, F., Muraro, D., Tysoe, O.C., Sawiak, S., Beach, T.E., Godfrey, E.M., Upponi, S.S.,

Brevini, T., Wesley, B.T., Garcia-Bernardo, J., et al. (2021). Cholangiocyte organoids can repair

bile ducts after transplantation in the human liver. Science (1979) *371*, 839–846.

https://doi.org/10.1126/science.aaz6964.

Saxena, K., Blutt, S.E., Ettayebi, K., Zeng, X.-L., Broughman, J.R., Crawford, S.E., Karandikar,

U.C., Sastri, N.P., Conner, M.E., Opekun, A.R., et al. (2016). Human Intestinal Enteroids: a New

Model To Study Human Rotavirus Infection, Host Restriction, and Pathophysiology. J Virol *90*,

43–56. https://doi.org/10.1128/jvi.01930-15.

Schaefer, K., Rensing, S., Hillen, H., Burkhardt, J.E., and Germann, P.G. (2016). Is Science the

only Driver in Species Selection? An Internal Study to Evaluate Compound Requirements in the

Minipig Compared to the Dog in Preclinical Studies. Toxicol Pathol *44*, 474–479.

https://doi.org/10.1177/0192623315624572.

Schneider, B., Balbas-Martinez, V., Jergens, A.E., Troconiz, I.F., Allenspach, K., and Mochel,

J.P. (2018). Model-based reverse translation between veterinary and human medicine: The one

health initiative. CPT Pharmacometrics Syst Pharmacol *7*, 65–68.

https://doi.org/10.1002/psp4.12262.

Sebbag, L., McDowell, E.M., Hepner, P.M., and Mochel, J.P. (2018). Effect of tear collection on

lacrimal total protein content in dogs and cats: A comparison between Schirmer strips and

ophthalmic sponges. BMC Vet Res *14*. https://doi.org/10.1186/s12917-018-1390-7.

Sebbag, L., Kirner, N.S., Allbaugh, R.A., Reis, A., and Mochel, J.P. (2019). Kinetics of

Fluorescein in Tear Film After Eye Drop Instillation in Beagle Dogs: Does Size Really Matter?

Front Vet Sci *6*. https://doi.org/10.3389/fvets.2019.00457.

Shalev, M. (2006). New FDA recommendations to speed drug development. Lab Anim (NY) *35*,

13. https://doi.org/10.1038/laban0306-13a.

Shiota (Sato), Y., Elbadawy, M., Suzuki, K., Tsunedomi, R., Nagano, H., Ishihara, Y.,

Yamamoto, H., Azakami, D., Uchide, T., Fukushima, R., et al. (2023). Derivation of a new model

of lung adenocarcinoma using canine lung cancer organoids for translational research in

pulmonary medicine. Biomedicine and Pharmacotherapy *165*.

https://doi.org/10.1016/j.biopha.2023.115079.

Silva, K.A.S., and Emter, C.A. (2020). Large Animal Models of Heart Failure: A Translational

Bridge to Clinical Success. JACC Basic Transl Sci *5*, 840–856.

https://doi.org/10.1016/j.jacbts.2020.04.011.

Stadler, K., Handler, J., Schoenkypl, S., and Walter, I. (2009). A three-dimensional culture

model of canine uterine glands. In Vitro Cell Dev Biol Anim *45*, 35–43.

https://doi.org/10.1007/s11626-008-9127-8.

Takeda, H., Kataoka, S., Nakayama, M., Ali, M.A.E., Oshima, H., Yamamoto, D., Park, J.W.,

Takegami, Y., An, T., Jenkins, N.A., et al. (2019). CRISPR-Cas9-mediated gene knockout in

intestinal tumor organoids provides functional validation for colorectal cancer driver genes. Proc

Natl Acad Sci U S A *116*, 15635–15644. https://doi.org/10.1073/pnas.1904714116.

Team, R.C. (2013). R: A language and environment for statistical computing.

Tortorella, I., Argentati, C., Emiliani, C., Martino, S., and Morena, F. (2022). The role of physical

cues in the development of stem cell-derived organoids. European Biophysics Journal *51*, 105–

117. https://doi.org/10.1007/s00249-021-01551-3.

Turco, M.Y., Gardner, L., Hughes, J., Cindrova-Davies, T., Gomez, M.J., Farrell, L.,

Hollinshead, M., Marsh, S.G.E., Brosens, J.J., Critchley, H.O., et al. (2017). Long-term,

hormone-responsive organoid cultures of human endometrium in a chemically defined medium.

Nat Cell Biol *19*, 568–577. https://doi.org/10.1038/ncb3516.

Usui, T., Sakurai, M., Nishikawa, S., Umata, K., Nemoto, Y., Haraguchi, T., Itamoto, K., Mizuno,

T., Noguchi, S., Mori, T., et al. (2017). Establishment of a dog primary prostate cancer organoid

using the urine cancer stem cells. Cancer Sci *108*, 2383–2392.

https://doi.org/10.1111/cas.13418.

Wang, J., Wang, T., Sun, Y., Feng, Y., Kisseberth, W.C., Henry, C.J., Mok, I., Lana, S.E.,

Dobbin, K., Northrup, N., et al. (2018a). Proliferative and invasive colorectal tumors in pet dogs

provide unique insights into human colorectal cancer. Cancers (Basel) *10*.

https://doi.org/10.3390/cancers10090330.

Wang, Y., Tang, Z., Huang, H., Li, J., Wang, Z., Yu, Y., Zhang, C., Li, J., Dai, H., Wang, F., et

al. (2018b). Pulmonary alveolar type I cell population consists of two distinct subtypes that differ

in cell fate. Proc Natl Acad Sci U S A *115*, 2407–2412.

https://doi.org/10.1073/pnas.1719474115.

Wiedenmann, S., Breunig, M., Merkle, J., von Toerne, C., Georgiev, T., Moussus, M., Schulte,

L., Seufferlein, T., Sterr, M., Lickert, H., et al. (2021). Single-cell-resolved differentiation of

human induced pluripotent stem cells into pancreatic duct-like organoids on a microwell chip.

Nat Biomed Eng *5*, 897–913. https://doi.org/10.1038/s41551-021-00757-2.

Xenoulis, P.G. (2015). Diagnosis of pancreatitis in dogs and cats. Journal of Small Animal

Practice *56*, 13–26. https://doi.org/10.1111/jsap.12274.

Yu, L., Li, Z., Mei, H., Li, W., Chen, D., Liu, L., Zhang, Z., Sun, Y., Song, F., Chen, W., et al.

(2021). Patient-derived organoids of bladder cancer recapitulate antigen expression profiles and

serve as a personal evaluation model for CAR-T cells in vitro. Clin Transl Immunology *10*.

https://doi.org/10.1002/cti2.1248.

Zdyrski, C., Gabriel, V., Gessler, T.B., Ralston, A., Sifuentes-Romero, I., Kundu, D., Honold, S.,

Wickham, H., Topping, N.E., Sahoo, D.K., et al. (2024). Establishment and characterization of

turtle liver organoids provides a potential model to decode their unique adaptations. Commun

Biol *7*. https://doi.org/10.1038/s42003-024-05818-1.

Zhou, J., Li, C., Sachs, N., Chiu, M.C., Wong, B.H.-Y., Chu, H., Poon, V.K.-M., Wang, D., Zhao,

X., Wen, L., et al. (2018). Differentiated human airway organoids to assess infectivity of

emerging influenza virus. Proceedings of the National Academy of Sciences *115*, 6822–6827.

https://doi.org/10.1073/pnas.1806308115.

1061