1  **Title: Interlimb coordination in Parkinson's Disease is minimally affected by a**
2  **visuospatial dual task**

Allen Hill[a] ORCID: 0000-0003-0939-0765
Julie Nantel[a]* ORCID: 0000-0001-7253-9222

[a] School of Human Kinetics, University of Ottawa, Ottawa, ON, Canada
* corresponding author, jnantel@uottawa.ca

# Abstract

Parkinson's disease (PD) leads to reduced spatial and temporal interlimb coordination during gait as well
as reduced coordination in the upper or lower limbs. Multi-tasking when walking is common during real-
world activities, and affects some gait characteristics, like gait speed and variability. However, the impact
of a dual task (DT) on intra and interlimb coordination of both lower and upper limbs when walking in
people with PD remains unknown. Seventeen volunteers with mild to moderate PD (11 males, $65 \pm 8$ years,
$173 \pm 8$ cm, $74 \pm 20$ kg, Unified Parkinson's Disease Rating Scale motor section $10 \pm 5$) participated in gait
trials in an Extended-CAREN system, which includes a treadmill, 12-camera Vicon motion capture system,
and a 180° field-of-view virtual reality projection screen. Participants completed a 3 min walking trial and
a 2 min visuospatial word recognition DT trial at their preferred walking pace. Single and DT were com-
pared with a paired t-test, and the less and more affected (LA, MA) sides were tested for equivalence in
sensitivity to the DT. During the DT, we found the LA shoulder ROM decreased by 1.5°, and the LA
shoulder peak flexion decreased by 1.1° (p<.028, $g_{av}$>.12). The LA and MA hip ROM were differently
affected by the dual task (p=.023), and intralimb coordination was affected by dual tasking equivalently
between sides (p=.004). These results suggest that during normal single-task gait, people with PD use at-
tentional resources to compensate for reduced arm swing. Furthermore, our results indicate that any effect
of DT on lower intralimb coordination is not meaningfully different between the LA and MA sides.
**Keywords**: Parkinson's disease, dual task, interlimb coordination, relative phase, more affected side

# Introduction

Parkinson's disease (PD) is a multisystem neurodegenerative disease characterized by the loss of dopaminergic neurons in the substantia nigra with cascading effects in other regions, including those involved in cholinergic systems (Yarnall et al., 2011; Poewe et al., 2017). The neurodegeneration in PD begins unilaterally and progresses bilaterally, and motor symptoms mirror this unilateral emergence and bilateral progression, commonly resulting in one side being more affected than the other (Djaldetti et al., 2006). In addition to the three cardinal motor symptoms of PD—bradykinesia, rigidity, and tremor—people with PD develop further motor deficits, such as hypokinesia (reduced movement amplitude), increased movement variability, gait asymmetry, and postural instability, all of which may contribute to or reflect impairments in coordinating the upper and lower limbs during locomotion (van Emmerik & Wagenaar, 1996; Yogev et al., 2007; Plotnik et al., 2007; Mirelman et al., 2019). Coordination of the upper and lower limbs is required to maintain dynamic gait stability in the presence of major and minor perturbations (Marigold & Misiaszek, 2009; Krasovsky et al., 2012). In addition, gait tasks involving speed modulation require adaptations in interlimb coordination. Such tasks have shown kinematic differences in older adult fallers compared to non-faller peers (Barak et al., 2006; Shishov et al., 2017).

Coordination is defined as the context- and phase-dependent control of spatial and temporal cyclical relationships between body segments (Krasovsky & Levin, 2010). One prominent feature of normal gait coordination is the frequency-matched anti-phase swing (180 deg phase offset) between the arms, legs, and ipsilateral arm-leg pairs, while contralateral arm-leg pairs swing in phase with each other (Wagenaar & van Emmerik, 2000). (One-to-one arm and leg swing frequencies are expected for gait speeds above 0.8 m/s, while slower gait sometimes exhibits 2:1 arm swing to stride frequencies (Wagenaar & van Emmerik, 2000).) These phase relationships result in gait which is generally temporally and spatially symmetric in healthy adults (Sadeghi et al., 2000; Killeen et al., 2018). In contrast, asymmetry in spatial or temporal gait characteristics is recognized as a disruption in the coordination of normal gait and occurs in several pathological populations, including those with PD (Yogev et al., 2007; Huang et al., 2012; Park et al., 2016).

One approach to measuring coordination is based on dynamical systems theory and uses the continuous relative phase (CRP) between coupled oscillators to characterize the state of the system (Haken et al., 1985). Previous studies using CRP have found that, compared to healthy peers, people with PD have reduced coordination stability and increased average phase error (from anti- or in-phase) for upper and lower interlimb coordination during gait and for upper limb coordination during bimanual tasks (van Emmerik & Wagenaar, 1996; Winogrodzka et al., 2005; Almeida & Brown, 2013). Other studies support the presence of reduced interlimb coordination stability in PD (Roemmich et al., 2013), and increased phase error in some upper and lower limb pairs compared to healthy peers and comparing PD freezers to non-freezers (Nanhoe-Mahabier et al., 2011).

Homologous limb coordination is decreased in PD as well. Plotnik and collaborators (2007) reported worse lower limb coordination in people with PD compared to healthy controls using the phase coordination index (PCI), a combined measure of accuracy and consistency in the anti-phase coordination of step timing during gait. Coordination between the upper limbs is also reduced or altered in people with PD compared to healthy controls (Huang et al., 2012; Sterling et al., 2015).

Although there is support for coordination deficits in PD, the more and less affected (MA and LA) sides may contribute differently to interlimb coordination amidst the predominantly unilateral symptoms of early PD. However, the relative nature of coordination makes it difficult to separately characterize coordination between the MA and LA sides. Some studies have found that the LA side maintains an ability to adapt to and compensate for task requirements, such as split-belt walking (Roemmich et al., 2014) or dual-tasking during gait (Siragy & Nantel, 2020); despite the potential benefits of adaptation, this may appear as "worse" coordination due to the decreased similarity between sides. Other studies have found that people with PD showed increased PCI (ie less accurate and/or consistent coordination) when asked to use non-preferred step lengths and times, or when asked to adapt to external mechanical constraints, such as walking on split-belt treadmills (Williams et al., 2013; Fasano et al., 2016); the authors suggest that the increased PCI indicates a generally reduced ability to adapt (lower limb) coordination in PD.

However, adaption to split-belt walking differed between MA and LA sides, depending on which side was
sped up (Fasano et al., 2016), and therapy focused on the LA side may be more effective than a balanced
or MA-side focus (Ricciardi et al., 2015). In addition, two studies have found that ipsilateral arm-leg
coordination is reduced on the MA side (Nanhoe-Mahabier et al., 2011; Roemmich et al., 2013). CRP can
be used to measure coordination stability within individual limbs (i.e. intralimb coordination), which
could contribute to a better understanding of coordination deficits in specific limbs (Barela et al., 2000;
Byrne et al., 2002). However, we are not aware of previous studies having investigated this in people with
PD.
Performing a secondary task during gait adds an attentional stressor that can reveal the loss of
automaticity for some aspects of gait (e.g. gait speed, gait variability), and this is measured as a change in
performance while dual tasking, often referred to as the dual-task cost (Yogev-Seligmann et al. 2008). In
addition, experimental studies have shown that coordination of bimanual movement patterns can be
stabilized by an attentional focus, but a dual task serving as an attentional distractor interferes with the
stabilization (Temprado et al., 1999). Since gait while dual tasking may be more similar to gait in the real
world than typical laboratory recordings of steady state gait (Hillel et al., 2019), dual tasking during gait
may offer an ecologically valid method of manipulating attention to highlight coordinative deficits. Studies
of coordination and dual tasking in people with PD show a larger decline in lower limb coordination,
measured by PCI, during a dual task compared to healthy controls, and PCI in PD fallers and freezers is
more affected by a dual task than PD non-fallers and non-freezers (Plotnik & Hausdorff, 2008; Plotnik et
al., 2009; Plotnik, Giladi, et al., 2011). In addition, studies have found reduced arm swing during gait in
people with PD during a dual task compared to single task gait (Mirelman et al., 2016; Baron et al., 2018).
These results suggest that loss of automaticity in coordination may be associated with the progression of
PD to more severe outcomes.
While effects of dual tasking on bilateral coordination and asymmetry have been found (Plotnik,
Dagan, et al., 2011), as well as differences in spatiotemporal and movement characteristics between the
MA and LA limbs in the upper and lower body (Mirelman et al., 2016; Siragy & Nantel, 2020), the effect of
motor symptom presentation (i.e. MA and LA sides) in coordination within and between the upper and
lower limbs has not been sufficiently explored.  Differential dual-task cost between sides in interlimb
coordination in people with PD could provide evidence of how the asymmetric presentation of the disease
affects coordination in the upper and lower limbs. Therefore, the purpose of this study is to investigate
MA/LA side differences in coordination deficits in people with PD by observing interlimb coordination
within and between the upper and lower limbs during single and dual-task gait.

## Methods

### Participants

Volunteers with mild to moderate PD (between I-III Hoehn & Yahr) were recruited from the Ottawa-
Gatineau area. Exclusion criteria included any additional neurological impairment, a recent orthopedic
injury or surgery that could interfere with gait, the use of a walking aid, or any discomfort with using a
projected virtual reality system. Participant characteristics for sex, age, height, weight, handedness,
Unified Parkinson's Disease Rating Scale (UPDRS) motor section (III), freezing and falling status, and
interval since diagnosis of PD were recorded. The MA side was defined as the side where PD motor
symptoms first occurred, as reported by participants. Participants were tested on their normal
medication. All participants provided written informed consent, and the study was approved by local
ethics review boards.

### Protocol

Data was collected using the CAREN system (CAREN-Extended, Motekforce Link, Amsterdam, NL). The
CAREN system consists of an instrumented split-belt treadmill (Bertec Corp., Columbus. OH) embedded
in a six degree-of-freedom motion platform, a 12-camera motion capture system (Vicon, Oxford, UK), and
a 180 deg field of view projection screen. The participants wore a safety harness attached to an overhead
frame on the motion platform to prevent falls without restricting movement.

Participants were allowed an initial familiarization period with the CAREN system for several minutes
until they reported being comfortable; the familiarization period was also used to determine their
preferred walking speed for the gait trials. The single task trial was 3 min long, while the dual task trial
was 2 min long. The dual task consisted of a visuospatial word recognition and acknowledgement task
where a word was shown at eye level at a random position between 20-70 deg to the left or right of center;
participants were asked to notice and acknowledge the word by reading it aloud. Twelve words were
randomly drawn from a standard list of 16 possible words in the native language of the participant
(English or French). The dual task began 20 s into the trial, and a new word was shown for 3 s every 2-4 s
for 80 s. The dual task was designed to be an ecologically valid recreation of common daily life situations
(e.g. public transportation terminal, etc.) requiring perception and comprehension of visual cues (Siragy
& Nantel, 2020; Ahmadi et al., 2021). The dual task trial occurred after the single task trial. Participants
were allowed to rest between trials when requested.

**Data reduction**
A set of 57 markers (Wilken et al., 2012) was used to capture full-body kinematics at 100 Hz; marker
data was then filtered using a 4th order dual-pass Butterworth low-pass filter with a cutoff frequency of
12 Hz. Low pass filter cutoff frequency was chosen based on a residual frequency analysis of marker data
using an RMS noise of 0.5 mm measured from static markers fixed to the motion platform (Winter, 2009).
Gait events were calculated using an algorithm based on the local extrema of the vertical position and
velocity of the heel marker (Roerdink et al., 2008). OpenSim was used with the Rajagopal et al. (2016)
model to perform inverse kinematics to extract bilateral knee, elbow, shoulder, and hip flexion/extension
angles (Delp et al., 2007). The wrist pronation/supination range of motion of the model was expanded
from 90 deg to 160 deg to better match normative biomechanical characteristics (Shaaban et al., 2008).
To detect changes in spatial coordination, range of motion (ROM), ROM coefficient of variation (COV),
and peak flexion were measured on the MA and LA sides for the shoulder and hip joints. Similarly, changes
in temporal coordination were assessed using average intercycle phase variability of CRP (interpreted as
coordination stability) to measure for ipsilateral shoulder and hip interlimb joint pairs and intralimb
(shoulder and elbow, hip and knee) joint pairs. Continuous phase was calculated as the angle of the
complex analytic signal produced by the Hilbert transform after centering the original signal (Lamb &
Stöckl, 2014).

**Statistics**
Effects of dual tasking on individual variables were tested using paired t-tests. Additionally, a TOST
procedure was used to test for equivalence between sides for changes due to dual tasking (DTC); this was
done for all lateral variables (e.g. shoulder ROM, which was measured on both LA and MA sides)  (Lakens,
2017). All tests were performed with $\alpha = 0.05$. Unless otherwise noted, equivalence bounds were set per
variable using unstandardized values (i.e. the original units/scale, without normalizing to group error) to
reduce bias (Lakens, 2017) and estimated from previously published results for healthy older adults. The
maximum bilateral difference in DTC for shoulder ROM and peak flexion was estimated to be 3.5° (Plate
et al., 2015, tbl. 1; Killeen et al., 2018, fig. 2). The maximum bilateral difference in DTC for shoulder ROM
COV was estimated at 6° (Mirelman et al., 2016, tbl. 2). Bounds for hip ROM and peak flexion were set as
0.5°, based on changes in DTC between left and right limbs in PD (Ribeiro et al., 2019, tbl. 2). Lower
intralimb phase variability bounds were set to 0.85° (Ghanavati et al., 2014, fig. 2A). Appropriate reference
data could not be found for the remaining variables (hip ROM COV, ipsilateral phase variability, and upper
intralimb phase variability), and the equivalence bound was set at Cohen's d=0.36, estimated from the
maximum bilateral difference in change in arm swing ROM due to DT in healthy older adults (Plate et al.,
2015, tbl. 1).
Effect sizes were calculated using Cohen's dav with a Hedge's g correction, noted as gav (Cumming,
2011; Lakens, 2013). A sensitivity power analysis for a two-tailed t-test was conducted in G*Power (Faul
et al., 2007) to find that the minimum detectable effect size with 80% power is $d_z = 0.72$ when $\alpha = .05$
and the sample size is 17. A sensitivity power analysis for a TOST equivalence test using the TOSTER
package in R showed that the minimum bounds that would be rejectable with 80% power is $d = .90$ when

$\alpha = .05$ and the sample size is 17. Arm swing for a given shoulder was treated as functionally absent when
ROM was less than 5 deg. No meaningful coordination was expected between a shoulder with functionally
absent arm swing and either the ipsilateral hip or the contralateral shoulder; therefore, the ipsilateral and
intralimb CRP variables for affected subjects/shoulders were removed as outliers—3 subjects at most,
depending on the variable. Circular statistics (circular mean and standard deviations) were used for the
variables with angular units (Fisher, 1993). All data reduction and statistical analyses were performed
with the Julia language using open-source libraries and code (Bezanson et al., 2017; Hill & Nantel, 2023).

# Results

Participant demographics are reported in Table 1. Twenty subjects who met the inclusion criteria
were recruited. Two participants with severe dyskinesia were excluded from this study, and a third
participant was excluded for moving their hands while talking during a significant portion of a trial; both
behaviors produce movements (Jankovic, 2005) which are disruptive to the normal coordination patterns
of steady state gait. Subjects were diagnosed with PD an average of 7.4 ± 4.5 years prior to study
participation.

**Table 1 Participant demographics**

| Characteristic | Mean ± SD | Range |
|---|---|---|
| Sex (n) | 11 M, 6 F | |
| Age (years) | 64.8 ± 7.7 | 48-79 |
| Height (cm) | 173.3 ± 7.6 | 165-188 |
| Weight (kg) | 74.2 ± 19.9 | 52-128 |
| Handedness | 14 R, 3 L | |
| More affected side | 9 R, 8 L | |
| UPDRS III | 10.2 ± 5.3 | 0-20 |
| Freezers (n) | 5 | |
| Fallers (n, <1yr) | 10 | |

166  Single and dual task results were calculated using an average of 139 ± 19 and 74 ± 6 steps, respectively
167  (dual task trial length was shorter than the single task trial). Preferred gait speed was 1.0 ± 0.2 m/s among
168  subjects.

**Unilateral effects of DT**

170  Spatial coordination results are reported in Table 2. The LA shoulder ROM and peak flexion decreased
171  by 1.5 deg 95% CI [−2.8,−0.2] and 1.1 deg [−2.1,−0.1], respectively, during dual task compared to single
172  task walking. Temporal coordination results are reported in Table 3; no significant differences were
173  detected (p>.061).

**Table 2 Spatial coordination in the more and less affected sides during single and dual task**

| Variable | | Single task | Dual task | t-test | p-value | $g_{av}$ |
|---|---|---|---|---|---|---|
| Shoulder ROM (deg) | LA | 21.2 ± 13.3 | 19.7 ± 12.4 | t(16)=−2.44 | 0.027 | −0.12 |
| | MA | 17.9 ± 12.6 | 19.7 ± 14.8 | t(16)=0.69 | 0.501 | 0.13 |
| Shoulder ROM COV (%) | LA | 17.2 ± 9.1 | 25.0 ± 38.5 | t(16)=1.01 | 0.329 | 0.32 |
| | MA | 22.1 ± 14.8 | 16.4 ± 6.6 | t(16)=−1.55 | 0.141 | −0.51 |
| Shoulder peak flexion (deg) | LA | 15.3 ± 8.6 | 14.2 ± 8.2 | t(16)=−2.41 | 0.028 | 0.47 |
| | MA | 13.3 ± 7.7 | 14.1 ± 9.3 | t(16)=0.48 | 0.64 | 0.05 |
| Hip ROM (deg) | LA | 38.5 ± 4.1 | 37.6 ± 3.9 | t(16)=−1.61 | 0.127 | −0.22 |
| | MA | 37.6 ± 5.1 | 37.6 ± 5.3 | t(16)=0.14 | 0.892 | 0.01 |
| Hip ROM COV (%) | LA | 4.2 ± 1.4 | 4.7 ± 1.8 | t(16)=1.82 | 0.087 | 0.27 |
| | MA | 4.5 ± 1.6 | 4.6 ± 1.8 | t(16)=1.12 | 0.278 | 0.10 |

| | | | | | | |
|---|---|---|---|---|---|---|
| Hip peak flexion (deg) | LA | 22.7 ± 3.9 | 22.4 ± 4.4 | t(16)=−0.92 | 0.374 | −0.01 |
| | MA | 20.4 ± 5.1 | 20.5 ± 5.0 | t(16)=0.10 | 0.923 | −0.18 |

175      ROM = range of motion, MA/LA = more/less affected side, $g_{av}$ = Cohen's $d_{av}$ effect size corrected with
176      Hedge's g.

178      **Table 3 Temporal coordination (PCI and phase variability) in and between the more and less**
179      **affected sides during single and dual task**

| Variable | | Single task | Dual task | t-test | p-value | $g_{av}$ |
|---|---|---|---|---|---|---|
| PCI (deg) | | 6.6 ± 3.0 | 7.6 ± 4.0 | t(16)=1.70 | 0.109 | 0.28 |
| Shoulder interlimb (deg) | | 20.1 ± 9.9 | 21.4 ± 14.3 | t(13)=0.41 | 0.688 | 0.1 |
| Hip interlimb (deg) | | 7.1 ± 2.7 | 7.1 ± 2.1 | t(16)=−0.13 | 0.895 | −0.02 |
| Ipsilateral shoulder-hip (deg) | LA | 18.1 ± 15.5 | 22.1 ± 17.5 | t(15)=1.49 | 0.157 | 0.24 |
| | MA | 16.6 ± 9.1 | 14.6 ± 6.0 | t(13)=−0.43 | 0.675 | −0.15 |
| Contralateral shoulder-hip (deg)* | LA | 17.9 ± 15.4 | 21.9 ± 17.7 | t(15)=1.48 | 0.16 | 0.24 |
| | MA | 16.3 ± 8.9 | 14.0 ± 6.0 | t(13)=−0.57 | 0.581 | −0.2 |
| Upper intralimb (deg) | LA | 22.5 ± 10.2 | 26.3 ± 16.0 | t(15)=2.03 | 0.061 | 0.28 |
| | MA | 31.5 ± 21.6 | 26.8 ± 13.5 | t(13)=−0.58 | 0.573 | −0.16 |
| Lower intralimb (deg) | LA | 6.6 ± 2.5 | 7.1 ± 2.5 | t(16)=1.62 | 0.126 | 0.21 |
| | MA | 6.8 ± 2.4 | 7.1 ± 1.9 | t(16)=0.87 | 0.396 | 0.11 |

PCI = phase coordination index (Plotnik et al., 2007), MA/LA = more/less affected side, $g_{av}$ = Cohen's
$d_{av}$ effect size corrected with Hedge's g. *LA/MA refers to the shoulder of the contralateral shoulder-
hip pair. Arm swing for three subjects was treated as functionally absent in one or both shoulders,
which reduced the degrees of freedom for the t-tests of some variables.

**Bilateral difference in DTC**
Shoulder ROM DTC was not significantly different between LA and MA sides, $t(16) = -1.26, p = .225$;
and the equivalence test was also not significant ($p = .467$), preventing a conclusion of equivalence to a
healthy older adult population. Shoulder ROM CoV DTC was not significantly different between sides,
$t(16) = 1.62, p = .125$; nor was the test of equivalence ($p = .808$). Shoulder peak flexion DTC was not
significantly different between sides, $t(16) = -1.13, p = .276$; nor was the test of equivalence ($p = .177$).
Hip ROM DTC was significantly different between sides, $M = -0.967°$ 95% CI $[-1.8, -0.15]$, $t(16) = -2.52$,
$p = .023$, and was not equivalent ($p = .88$). Hip ROM CoV DTC was not significantly different between
sides, $t(16) = 1.17, p = .26$; nor was the test of equivalence ($p = .378$). Hip peak flexion was not
significantly different between sides, $t(16) = -0.69, p = .502$; nor was the test of equivalence ($p = .223$).
Ipsilateral shoulder-hip phase variability DTC was not significantly different between sides, $t(13) = 1.64$,
$p = .124$; nor was the test of equivalence ($p = .614$). Upper intralimb phase variability DTC was not
significantly different between sides, $t(13) = 0.93, p = .369$; nor was the test of equivalence ($p = .342$).
Lower intralimb phase variability DTC was not significantly different between sides, $t(16) = 1.53, p =$
$.146$; and was statistically equivalent ($p = .004$).

# Discussion

In this study, we found that few aspects of coordination are unilaterally affected by a visuospatial dual
task in a cohort of mild to moderate PD. Furthermore, few aspects of coordination exhibit a significant
bilaterally different sensitivity to dual tasking, however, most variables do not support a conclusion of
equivalent sensitivity of coordination between the MA and LA sides. During the dual task, the LA shoulder
ROM and peak flexion decreased compared to single task performance (Table 2), but the change in
shoulder ROM and peak flexion due to dual tasking were not significantly different between the LA and
MA sides. A significant difference was noted in hip ROM DTC between MA and LA sides, but neither side

was affected by dual tasking. Lower intralimb phase variability was not affected by dual tasking and was
not different and equivalent between sides at $\pm0.85°$ bounds.
Our results are generally inconclusive, as null findings for both t-tests and equivalence tests prevent
meaningful conclusions about the presence or absence of meaningful (i.e. clinically relevant) effects.
Reduced arm swing during dual tasking on the LA side is similar to previous studies (Mirelman et al.,
2016; Baron et al., 2018). Additionally, gait asymmetry present in people with PD often results in LA ROM
and coordination that are more similar to—but not always matching—healthy peers, while the MA side is
less similar to healthy peers (Roggendorf et al., 2012; Roemmich et al., 2013). Qualitatively, the single task
shoulder ROM in our data showed that the arm swing was fairly symmetric (Table 2), and slightly reduced
on both sides compared to normative arm swing data for healthy older adults at similar gait speeds (Plate
et al., 2015; Killeen et al., 2018).
However, we note that the absence of a significant effect of dual tasking on the MA side should not be
interpreted as evidence for differing behaviors/responses to dual tasking between the MA and LA sides,
as Gelman and Stern (2006) have previously stated, "comparisons of the sort 'X is statistically significant
and Y is not' can be misleading". Indeed, we directly compared the unilateral DTC and found no difference
between sides. Since equivalence at the tested bounds was not reached, there remains the possibility of a
bilateral difference in DTC at an effect size that is smaller than we were powered to detect but too large
to reject at our chosen equivalence bounds.
Lower intralimb phase variability was not affected by the dual task used in this study, and the
difference in responses of each side to dual tasking was equivalent to estimated responses in healthy older
adults. Different dual tasks have been found to have population specific effects (i.e. task interference) (Al-
Yahya et al., 2011), and previous studies on gait and dual tasks in PD suggests that fallers and/or freezers
may be more sensitive to dual task interference (Plotnik, Giladi, et al., 2011; Bekkers et al., 2018). More
research is needed to determine whether such increased sensitivity for PD fallers or freezers would apply
to the novel visuospatial task—intended to mimic an ecologically realistic scenario—that was used in this
study.

**Limitations**
Due to the absence of a control group, we are only able to quantify the response of a mild PD group to
the DT, and are unable to discuss how responses to the DT in a group of healthy controls might differ from
this group of people with PD. In addition, the participants in this study had mild to moderate PD, and the
arm swing ROM and ROM asymmetry within our cohort is markedly different compared to previously
reported PD cohorts (Roggendorf et al., 2012; Isaias et al., 2012; Sterling et al., 2015; Mirelman et al.,
2016); it is unclear how coordination would respond in people with similarly mild PD—but with larger
levels of arm swing asymmetry, or in people with more severe PD—which have more symmetric arm
swing but reduced ROM (Roggendorf et al., 2012). Similarly, the severity of PD—in terms of UPDRS III
scores—may be correlated with dual task interference, however the sparse range of UPDRS III values in
our cohort prevented the use of statistical tests that might detect any such interactions.
Additionally, different tasks are known to have unique and specific effects on different aspects of gait
which may limit the generalizability of our results (Al-Yahya et al., 2011; Rochester et al., 2014), and the
visuospatial dual task in this study was simple (12/17 participants demonstrated perfect performance,
and the remaining 5 participants responded to 10 ± 1.5 words out of 12). However, Baron et al. (2018)
found that arm swing kinematics were sensitive to multiple common dual tasks.
Finally, treadmill walking is different from overground walking and may prevent some common
responses in PD, such as reducing gait speed when distracted. However, our results are still informative
about the nature of coordination deficits in PD and the methods they use to compensate for coordination
deficits when optimally medicated and walking in this specific environment.

# Conclusion

Our results show that in a group of mild PD, a visuospatial dual task during gait contributes to
decreased arm ROM on the LA side, but only hip ROM is differently affected by a dual task on the MA and

LA sides. Furthermore, with the exception of coordination within the lower limbs, tests of equivalent
sensitivity to dual tasking on the MA and LA sides were inconclusive. Despite the inconclusive tests, these
results support cumulative science by providing reference data useful in calculating effect sizes for power
analyses of future studies on the topic (Lakens, 2013). More research is needed to support or dispute the
existence of differences in coordination between the MA and LA sides.

# Declarations

## Funding
263     This work was supported by the Natural Sciences and Engineering Research Council of Canada
264 (https://www.nserc-crsng.gc.ca) [RGPIN-2016-04928 to J.N., RGPAS 493045-2016 to J.N.], and by the
265 Ontario Ministry of Research, Innovation and Science (https://www.ontario.ca/page/early-researcher-
266 awards) Early Researcher Award [ER 16-12-206 to J.N.]. The funders had no role in study design, data
267 collection and analysis, decision to publish, or preparation of the manuscript.

## Competing interests
269     The authors have no relevant financial or non-financial interests to disclose.

## Ethics approval
271     This study was performed in accordance with the Declaration of Helsinki. Approval was granted by
272 the Ottawa Health Science Network Research Ethics Board (No. 20170291-01H) and the University of
273 Ottawa Office of Research Ethics and Integrity (No. A06-17-03).

## Consent to participate
275     Written informed consent was obtained from all study participants.

## Data and code availability
277     The software and data produced and analyzed for this study are openly available from the Zenodo data
278 repository at https://doi.org/10.5281/zenodo.8364708 (Hill & Nantel, 2023)

## Author contributions
280     **Allen Hill:** Conceptualization, Methodology, Software, Validation, Formal Analysis, Investigation,
281 Resources, Data Curation, Writing – Original Draft. **Julie Nantel:** Investigation, Resources, Writing –
282 Review & Editing, Supervision, Project administration, Funding acquisition

**Bibliography**

Ahmadi S, Siragy T, Nantel J (2021) Regularity of kinematic data between single and dual-task treadmill walking in people with Parkinson's disease. *Journal of NeuroEngineering and Rehabilitation*, **18**, 20. https://doi.org/10.1186/s12984-021-00807-5

Almeida QJ, Brown MJN (2013) Is DOPA-Responsive Hypokinesia Responsible for Bimanual Coordination Deficits in Parkinson's Disease? *Frontiers in Neurology*, **4**. https://doi.org/10.3389/fneur.2013.00089

Al-Yahya E, Dawes H, Smith L, Dennis A, Howells K, Cockburn J (2011) Cognitive motor interference while walking: A systematic review and meta-analysis. *Neuroscience & Biobehavioral Reviews*, **35**, 715–728. https://doi.org/10.1016/j.neubiorev.2010.08.008

Barak Y, Wagenaar RC, Holt KG (2006) Gait Characteristics of Elderly People With a History of Falls: A Dynamic Approach. *Physical Therapy*, **86**, 1501–1510. https://doi.org/10.2522/ptj.20050387

Barela JA, Whitall J, Black P, Clark JE (2000) An examination of constraints affecting the intralimb coordination of hemiparetic gait. *Human Movement Science*, **19**, 251–273. https://doi.org/10.1016/S0167-9457(00)00014-2

Baron EI, Miller Koop M, Streicher MC, Rosenfeldt AB, Alberts JL (2018) Altered kinematics of arm swing in Parkinson's disease patients indicates declines in gait under dual-task conditions. *Parkinsonism & Related Disorders*, **48**, 61–67. https://doi.org/10.1016/j.parkreldis.2017.12.017

Bekkers EMJ, Dockx K, Devan S, Van Rossom S, Verschueren SMP, Bloem BR, Nieuwboer A (2018) The Impact of Dual-Tasking on Postural Stability in People With Parkinson's Disease With and Without Freezing of Gait. *Neurorehabilitation and Neural Repair*, **32**, 166–174. https://doi.org/10.1177/1545968318761121

Bezanson J, Edelman A, Karpinski S, Shah V (2017) Julia: A Fresh Approach to Numerical Computing. *SIAM Review*, **59**, 65–98. https://doi.org/10.1137/141000671

Byrne JE, Stergiou N, Blanke D, Houser JJ, Kurz MJ, Hageman PA (2002) Comparison of Gait Patterns between Young and Elderly Women: An Examination of Coordination. *Perceptual and Motor Skills*, **94**, 265–280. https://doi.org/10.2466/pms.2002.94.1.265

Cumming G (2011) *Understanding The New Statistics: Effect Sizes, Confidence Intervals, and Meta-Analysis*. Routledge, New York. https://doi.org/10.4324/9780203807002

Delp SL, Anderson FC, Arnold AS, Loan P, Habib A, John CT, Guendelman E, Thelen DG (2007) OpenSim: Open-Source Software to Create and Analyze Dynamic Simulations of Movement. *IEEE Transactions on Biomedical Engineering*, **54**, 1940–1950. https://doi.org/10.1109/TBME.2007.901024

Djaldetti R, Ziv I, Melamed E (2006) The mystery of motor asymmetry in Parkinson's disease. *The Lancet Neurology*, **5**, 796–802. https://doi.org/10.1016/S1474-4422(06)70549-X

van Emmerik REA, Wagenaar RC (1996) Dynamics of movement coordination and tremor during gait in Parkinson's disease. *Human Movement Science*, **15**, 203–235. https://doi.org/10.1016/0167-9457(95)00044-5

Fasano A, Schlenstedt C, Herzog J, Plotnik M, Rose FEM, Volkmann J, Deuschl G (2016) Split-belt locomotion in Parkinson's disease links asymmetry, dyscoordination and sequence effect. *Gait & Posture*, **48**, 6–12. https://doi.org/10.1016/j.gaitpost.2016.04.020

Faul F, Erdfelder E, Lang A-G, Buchner A (2007) G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, **39**, 175–191. https://doi.org/10.3758/BF03193146

Fisher NI (1993) *Statistical Analysis of Circular Data*. Cambridge University Press. https://doi.org/10.1017/CBO9780511564345

Gelman A, Stern H (2006) The Difference Between "Significant" and "Not Significant" is not Itself Statistically Significant. *The American Statistician*, **60**, 328–331. https://doi.org/10.1198/000313006X152649

Ghanavati T, Salavati M, Karimi N, Negahban H, Ebrahimi Takamjani I, Mehravar M, Hessam M (2014) Intra-limb coordination while walking is affected by cognitive load and walking speed. *Journal of Biomechanics*, **47**, 2300–2305. https://doi.org/10.1016/j.jbiomech.2014.04.038

Haken H, Kelso JAS, Bunz H (1985) A theoretical model of phase transitions in human hand movements. *Biological Cybernetics*, **51**, 347–356. https://doi.org/10.1007/BF00336922

Hill A, Nantel J (2023) Data and Code for: Interlimb coordination in Parkinson's Disease is minimally affected by a visuospatial dual task. https://doi.org/10.5281/zenodo.8364708

Hillel I, Gazit E, Nieuwboer A, Avanzino L, Rochester L, Cereatti A, Croce UD, Rikkert MO, Bloem BR, Pelosin E, Del Din S, Ginis P, Giladi N, Mirelman A, Hausdorff JM (2019) Is every-day walking in older adults more analogous to dual-task walking or to usual walking? Elucidating the gaps between gait performance in the lab and during 24/7 monitoring. *European Review of Aging and Physical Activity*, **16**, 6. https://doi.org/10.1186/s11556-019-0214-5

Huang X, Mahoney JM, Lewis MM, Guangwei Du, Piazza SJ, Cusumano JP (2012) Both coordination and symmetry of arm swing are reduced in Parkinson's disease. *Gait & Posture*, **35**, 373–377. https://doi.org/10.1016/j.gaitpost.2011.10.180

Isaias IU, Volkmann J, Marzegan A, Marotta G, Cavallari P, Pezzoli G (2012) The Influence of Dopaminergic Striatal Innervation on Upper Limb Locomotor Synergies. *PLOS ONE*, **7**, e51464. https://doi.org/10.1371/journal.pone.0051464

Jankovic J (2005) Motor fluctuations and dyskinesias in Parkinson's disease: Clinical manifestations. *Movement Disorders*, **20**, S11–S16. https://doi.org/10.1002/mds.20458

Killeen T, Elshehabi M, Filli L, Hobert MA, Hansen C, Rieger D, Brockmann K, Nussbaum S, Zörner B, Bolliger M, Curt A, Berg D, Maetzler W (2018) Arm swing asymmetry in overground walking. *Scientific Reports*, **8**, 12803. https://doi.org/10.1038/s41598-018-31151-9

Krasovsky T, Baniña MC, Hacmon R, Feldman AG, Lamontagne A, Levin MF (2012) Stability of gait and interlimb coordination in older adults. *Journal of Neurophysiology*, **107**, 2560–2569. https://doi.org/10.1152/jn.00950.2011

Krasovsky T, Levin MF (2010) Review: Toward a Better Understanding of Coordination in Healthy and Poststroke Gait. *Neurorehabilitation and Neural Repair*, **24**, 213–224. https://doi.org/10.1177/1545968309348509

Lakens D (2013) Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and ANOVAs. *Frontiers in Psychology*, **4**. https://doi.org/10.3389/fpsyg.2013.00863

Lakens D (2017) Equivalence Tests: A Practical Primer for t Tests, Correlations, and Meta-Analyses. *Social Psychological and Personality Science*, **8**, 355–362. https://doi.org/10.1177/1948550617697177

Lamb PF, Stöckl M (2014) On the use of continuous relative phase: Review of current approaches and outline for a new standard. *Clinical Biomechanics*, **29**, 484–493. https://doi.org/10.1016/j.clinbiomech.2014.03.008

Marigold DS, Misiaszek JE (2009) Whole-Body Responses: Neural Control and Implications for Rehabilitation and Fall Prevention. *The Neuroscientist*, **15**, 36–46. https://doi.org/10.1177/1073858408322674

Mirelman A, Bernad-Elazari H, Thaler A, Giladi-Yacobi E, Gurevich T, Gana-Weisz M, Saunders-Pullman R, Raymond D, Doan N, Bressman SB, Marder KS, Alcalay RN, Rao AK, Berg D, Brockmann K, Aasly J, Waro BJ, Tolosa E, Vilas D, Pont-Sunyer C, Orr-Urtreger A, Hausdorff JM, Giladi N (2016) Arm swing as a potential new prodromal marker of Parkinson's disease. *Movement Disorders*, **31**, 1527–1534. https://doi.org/10.1002/mds.26720

Mirelman A, Bonato P, Camicioli R, Ellis TD, Giladi N, Hamilton JL, Hass CJ, Hausdorff JM, Pelosin E, Almeida QJ (2019) Gait impairments in Parkinson's disease. *The Lancet Neurology*, **18**, 697–708. https://doi.org/10.1016/S1474-4422(19)30044-4

Nanhoe-Mahabier W, Snijders AH, Delval A, Weerdesteyn V, Duysens J, Overeem S, Bloem BR (2011) Walking patterns in Parkinson's disease with and without freezing of gait. *Neuroscience*, **182**, 217–224. https://doi.org/10.1016/j.neuroscience.2011.02.061

Park K, Roemmich RT, Elrod JM, Hass CJ, Hsiao-Wecksler ET (2016) Effects of aging and Parkinson's disease on joint coupling, symmetry, complexity and variability of lower limb movements during gait. *Clinical Biomechanics*, **33**, 92–97. https://doi.org/10.1016/j.clinbiomech.2016.02.012

385 Plate A, Sedunko D, Pelykh O, Schlick C, Ilmberger JR, Bötzel K (2015) Normative data for arm swing
386  asymmetry: How (a)symmetrical are we? *Gait & Posture*, **41**, 13–18.
387  https://doi.org/10.1016/j.gaitpost.2014.07.011
388 Plotnik M, Dagan Y, Gurevich T, Giladi N, Hausdorff JM (2011) Effects of cognitive function on gait and
389  dual tasking abilities in patients with Parkinson's disease suffering from motor response fluctuations.
390  *Experimental Brain Research*, **208**, 169–179. https://doi.org/10.1007/s00221-010-2469-y
391 Plotnik M, Giladi N, Dagan Y, Hausdorff JM (2011) Postural instability and fall risk in Parkinson's disease:
392  impaired dual tasking, pacing, and bilateral coordination of gait during the "ON" medication state.
393  *Experimental Brain Research*, **210**, 529–538. https://doi.org/10.1007/s00221-011-2551-0
394 Plotnik M, Giladi N, Hausdorff JM (2007) A new measure for quantifying the bilateral coordination of
395  human gait: effects of aging and Parkinson's disease. *Experimental Brain Research*, **181**, 561–570.
396  https://doi.org/10.1007/s00221-007-0955-7
397 Plotnik M, Giladi N, Hausdorff JM (2009) Bilateral coordination of gait and Parkinson's disease: the
398  effects of dual tasking. *Journal of Neurology, Neurosurgery & Psychiatry*, **80**, 347–350.
399  https://doi.org/10.1136/jnnp.2008.157362
400 Plotnik M, Hausdorff JM (2008) The role of gait rhythmicity and bilateral coordination of stepping in the
401  pathophysiology of freezing of gait in Parkinson's disease. *Movement Disorders*, **23**, S444–S450.
402  https://doi.org/10.1002/mds.21984
403 Poewe W, Seppi K, Tanner CM, Halliday GM, Brundin P, Volkmann J, Schrag A-E, Lang AE (2017)
404  Parkinson disease. *Nature Reviews Disease Primers*, **3**, 17013. https://doi.org/10.1038/nrdp.2017.13
405 Rajagopal A, Dembia CL, DeMers MS, Delp DD, Hicks JL, Delp SL (2016) Full-Body Musculoskeletal Model
406  for Muscle-Driven Simulation of Human Gait. *IEEE Transactions on Biomedical Engineering*, **63**, 2068–
407  2079. https://doi.org/10.1109/TBME.2016.2586891
408 Ribeiro TS, Sousa AC de, Lucena LC de, Santiago LMM, Lindquist ARR (2019) Does dual task walking
409  affect gait symmetry in individuals with Parkinson's disease? *European Journal of Physiotherapy*, **21**,
410  8–14. https://doi.org/10.1080/21679169.2018.1444086
411 Ricciardi L, Ricciardi D, Lena F, Plotnik M, Petracca M, Barricella S, Bentivoglio AR, Modugno N, Bernabei
412  R, Fasano A (2015) Working on asymmetry in Parkinson's disease: randomized, controlled pilot
413  study. *Neurological Sciences*, **36**, 1337–1343. https://doi.org/10.1007/s10072-015-2082-8
414 Rochester L, Galna B, Lord S, Burn D (2014) The nature of dual-task interference during gait in incident
415  Parkinson's disease. *Neuroscience*, **265**, 83–94. https://doi.org/10.1016/j.neuroscience.2014.01.041
416 Roemmich RT, Field AM, Elrod JM, Stegemöller EL, Okun MS, Hass CJ (2013) Interlimb coordination is
417  impaired during walking in persons with Parkinson's disease. *Clinical Biomechanics*, **28**, 93–97.
418  https://doi.org/10.1016/j.clinbiomech.2012.09.005
419 Roemmich RT, Nocera JR, Stegemöller EL, Hassan A, Okun MS, Hass CJ (2014) Locomotor adaptation and
420  locomotor adaptive learning in Parkinson's disease and normal aging. *Clinical Neurophysiology*, **125**,
421  313–319. https://doi.org/10.1016/j.clinph.2013.07.003
422 Roerdink M, Coolen B (H. ), Clairbois B (H. ). E, Lamoth CJC, Beek PJ (2008) Online gait event detection
423  using a large force platform embedded in a treadmill. *Journal of Biomechanics*, **41**, 2628–2632.
424  https://doi.org/10.1016/j.jbiomech.2008.06.023
425 Roggendorf J, Chen S, Baudrexel S, van de Loo S, Seifried C, Hilker R (2012) Arm swing asymmetry in
426  Parkinson's disease measured with ultrasound based motion analysis during treadmill gait. *Gait &
427  Posture*, **35**, 116–120. https://doi.org/10.1016/j.gaitpost.2011.08.020
428 Sadeghi H, Allard P, Prince F, Labelle H (2000) Symmetry and limb dominance in able-bodied gait: a
429  review. *Gait & Posture*, **12**, 34–45. https://doi.org/10.1016/S0966-6362(00)00070-9
430 Shaaban H, Pereira C, Williams R, Lees VC (2008) The Effect Of Elbow Position On The Range Of
431  Supination And Pronation Of The Forearm. *Journal of Hand Surgery (European Volume)*, **33**, 3–8.
432  https://doi.org/10.1177/1753193407087862
433 Shishov N, Gimmon Y, Rashed H, Kurz I, Riemer R, Shapiro A, Debi R, Melzer I (2017) Old adult fallers
434  display reduced flexibility of arm and trunk movements when challenged with different walking
435  speeds. *Gait & Posture*, **52**, 280–286. https://doi.org/10.1016/j.gaitpost.2016.12.004

Siragy T, Nantel J (2020) Absent Arm Swing and Dual-Tasking Decreases Trunk Postural Control and
Dynamic Balance in People with Parkinson's Disease. *Frontiers in Neurology*, **11**.
https://doi.org/10.3389/fneur.2020.00213
Sterling NW, Cusumano JP, Shaham N, Piazza SJ, Liu G, Kong L, Du G, Lewis MM, Huang X (2015)
Dopaminergic modulation of arm swing during gait among Parkinson's disease patients. *Journal of*
*Parkinson's disease*, **5**, 141–150. https://doi.org/10.3233/JPD-140447
Temprado J-J, Zanone P-G, Monno A, Laurent M (1999) Attentional load associated with performing and
stabilizing preferred bimanual patterns. *Journal of Experimental Psychology: Human Perception and*
*Performance*, **25**, 1579–1594. https://doi.org/10.1037/0096-1523.25.6.1579
Wagenaar RC, van Emmerik REA (2000) Resonant frequencies of arms and legs identify different
walking patterns. *Journal of Biomechanics*, **33**, 853–861. https://doi.org/10.1016/S0021-
9290(00)00020-8
Wilken JM, Rodriguez KM, Brawner M, Darter BJ (2012) Reliability and minimal detectible change values
for gait kinematics and kinetics in healthy adults. *Gait & Posture*, **35**, 301–307.
https://doi.org/10.1016/j.gaitpost.2011.09.105
Williams AJ, Peterson DS, Earhart GM (2013) Gait coordination in Parkinson disease: Effects of step
length and cadence manipulations. *Gait & Posture*, **38**, 340–344.
https://doi.org/10.1016/j.gaitpost.2012.12.009
Winogrodzka A, Wagenaar RC, Booij J, Wolters EC (2005) Rigidity and bradykinesia reduce interlimb
coordination in Parkinsonian gait. *Archives of Physical Medicine and Rehabilitation*, **86**, 183–189.
https://doi.org/10.1016/j.apmr.2004.09.010
Winter DA (2009) *Biomechanics and Motor Control of Human Movement*. John Wiley & Sons, Inc.,
Hoboken, New Jersey.
Yarnall A, Rochester L, Burn DJ (2011) The interplay of cholinergic function, attention, and falls in
Parkinson's disease. *Movement Disorders*, **26**, 2496–2503. https://doi.org/10.1002/mds.23932
Yogev G, Plotnik M, Peretz C, Giladi N, Hausdorff JM (2007) Gait asymmetry in patients with Parkinson's
disease and elderly fallers: when does the bilateral coordination of gait require attention?
*Experimental Brain Research*, **177**, 336–346. https://doi.org/10.1007/s00221-006-0676-3
Yogev-Seligmann G, Hausdorff JM, Giladi N (2008) The role of executive function and attention in gait.
*Movement Disorders*, **23**, 329–342. https://doi.org/10.1002/mds.21720
466