Title: Nicotine-Related Beliefs Induce Dose-Dependent Responses in the Human Brain

Authors: Ofer Perl*[1,2,3], Anastasia Shuster[1,2], Matthew Heflin[1,2], Soojung Na[1,2], Ambereen Kidwai[4], Natalie Booker[5], William C. Putnam[5], Vincenzo G. Fiore[1,2], Xiaosi Gu[1,2,3]*

Affiliations:

[1]Center for Computational Psychiatry, Icahn School of Medicine at Mount Sinai; New York, NY, 10029, USA

[2]Department of Psychiatry, Icahn School of Medicine at Mount Sinai; New York, NY 10029, USA,

[3]Nash Family Department of Neuroscience, Icahn School of Medicine at Mount Sinai; New York, NY, 10029, USA

[4]School of Behavioral and Brain Sciences, University of Texas at Dallas; Dallas, TX, 75080, USA

[5]Department of Pharmacy Practice, Jerry H. Hodge School of Pharmacy; Texas Tech University Health Sciences Center, Dallas, TX, 75235, USA

*Correspondence:
Ofer Perl (O.P) oferikoo@gmail.com, Xiaosi Gu (X.G) xiaosi.gu@mssm.edu.

Abstract (113 words):
Could non-pharmacological constructs, such as beliefs, impact brain activities in a dose-dependent manner as drugs do? Here, nicotine-addicted human participants were made to believe that an electronic cigarette they were about to vape contained either "low", "medium", or "high" levels of nicotine, while nicotine strength was held constant. We then measured their brain signals during decision-making with functional MRI. Across the entire brain, we found that only the thalamus, a brain region rich in nicotinic acetylcholine receptors, showed a dose-dependent response to beliefs in both neural activity and connectivity with the prefrontal cortex. These findings demonstrate a dose-response relationship between beliefs and neurobiology, a mechanism previously known to only apply to pharmacological agents.

Introduction

The concept of dose-response comes from pharmacology where the amount of active ingredients in a drug is known to modulate biological processes proportionally (*1*, *2*). In the case of neuropharmacology, establishing dose-response relationships has been fundamental for understanding the biochemical mechanisms mediating drug actions in the human brain. But is it possible that abstract cognitive constructs—such as beliefs—could modulate neurophysiological responses in a similar dose-dependent manner? Cognitive interventions that rely on changing one's beliefs, have been shown to affect brain and behavior (*3*). Still, these non-pharmacological, narrative-based interventions have traditionally been considered qualitative in nature and thus, lacking the measurable precision that is usually required for medicine or biological research.

We hypothesized that human beliefs could modulate neural activities in a dose-dependent fashion that is similar to pharmacologically-induced dose responses. This hypothesis is based on accumulating evidence suggesting that beliefs can interact with human biology. Notable examples include placebo and nocebo effects, in which one's symptoms improve or worsen due to positive ("placebo") or negative beliefs ("nocebo") about a treatment, while the drug contains no active ingredients (*3*). Even in the presence of a powerful neuroactive substance (e.g., nicotine), beliefs about nicotine can exert a binary all-or-none type of effect on neural responses in smokers (*4*). Collectively, these findings support the notion that beliefs can broadly affect neurobiological activities in humans. However, it is unclear whether beliefs can exert a dose-dependent-like effect on the brain. If so, such a finding would reveal a higher degree of sophistication and precision of the interaction between human beliefs and neurobiology than previously understood.

Results

We investigated this question by telling nicotine-addicted human participants that the strength of nicotine contained in an electronic cigarette (e-cig) they were about to vape was either "low", "medium", or "high". This made participants believe that nicotine content differed across e-cigs although the actual nicotine content was in fact kept constant across belief conditions. After vaping, participants (20 participants and a total of 60 scans) performed a monetary decision-making task during functional magnetic resonance imaging (fMRI; **Fig. 1a**). We chose to measure brain signals during reward processing because both nicotine itself and belief about nicotine have been shown to influence reward-related neural activities probed by similar decision-making paradigms (*4–7*).

Smokers' self-reported belief about nicotine content increased linearly with engineered belief

We took extensive measures to control for confounding factors. First, participants might vape less in "weaker" nicotine conditions due to a lack of interest. To control for this, we set vaping time to 20 minutes and quantified the amount of nicotine intake by measuring cartridge weight before and after vaping. Nicotine intake—indexed by change in cartridge weight (mg) —did not differ across belief conditions ($F(2,38) = 1.806$, $P = 0.178$; **Fig. 1b**). Second, nicotine metabolism might differ between belief conditions. To address this, we collected saliva samples

both before and after vaping for high-performance liquid chromatography tandem mass
spectrometric (LC-MS/MS) analytical quantification of cotinine—a nicotine metabolite
indicative of plasma nicotine levels (*8*). Changes in cotinine concentrations (ng/mL) were
comparable across conditions ($F(2,32) = 0.959$, $P = 0.393$; **Fig. 1c**). Third, we measured their
exhaled carbon monoxide (CO) before vaping as an index of participants' baseline nicotine
saturation level. We did not observe differences in CO levels across conditions (parts per million,
ppm; $F(2,32) = 0.364$, $P = 0.698$; **Fig. 1d**). Finally, we examined if participants subjectively
believed the instructions by asking them to rate perceived nicotine strength using a 10-point
scale. Participants' ratings significantly increased as a function of instructed beliefs (mean ± SD
(AU); 'low': = 3.52 ± 0.61, 'medium': = 4.52 ± 0.41, 'high': = 5.82 ± 0.47, $F(2,38) = 9.71$, $P =$
0.0004; **Fig. 1e**). Taken together, these analyses mitigate the concern that imbalanced nicotine
consumption, metabolism, or baseline deprivation might have contributed to any neural
differences across conditions, while confirming that our instruction successfully influenced
participants' beliefs about nicotine strength.
98
99   **Brain responses for monetary signals in bilateral posterior thalamus increased as a**
100   **function of belief in a dose-dependent fashion**
101
Next, we examined how instructed beliefs about nicotine might have modulated brain activities
during reward-based decision-making in our participants. In a whole-brain analysis of reward-
related neural activities, only the thalamus exhibited dose-dependent responses to instructed
beliefs about nicotine strength (peak at (-15, -19, -1); family-wise error (FWE) cluster-level
corrected $k = 50$, $P = 0.006$; **Fig. 2a**) Region of interest (ROI) analysis using an independent
anatomical mask further confirmed this dose-dependent responses in the posterior thalamus
(mean ± SD (AU); 'low': = 0.157 ± 1.047, 'medium': = 0.601 ± 0.714, 'high': = 2.914 ± 0.865;
standard rmANOVA test $F(2,38) = 3.62$, $P = 0.036$, **Fig. 2b;** permutation test $P = 0.002$, **Fig.**
2c). Several control analyses confirmed that the identified dose-response in thalamic activation
was not due to spatial smoothing (**Supplementary Fig. 1**) or sensorimotor effects (i.e., button
press or visual cues; **Supplementary Fig. 2**). Furthermore, in contrast to the thalamus, the
ventral striatum – a region also known to encode nicotine- and reward-related signals – tracked
reward activities across all conditions ($P_{FWE} < 0.05$, **Supplementary Fig. 3**), but its responses
were not modulated by beliefs (**Supplementary Fig. 4-5**). These results show that across the
entire brain, only the thalamus exhibited dose-dependent responses to instructed beliefs about
nicotine strength.
118
Given the individual variability in susceptibility to instructions, we next asked whether
participants' subjective beliefs, indexed by their self-reported perception about nicotine strength,
also modulated thalamic responses. We found that across all participants and all sessions,
subjective ratings of perceived nicotine strength correlated with reward-related activities in the
thalamus (Spearman correlation, $r = 0.27$, $P = 0.035$, **Fig. 2d**), suggesting that these brain signals
were linked to participants' perceptions about nicotine strength following belief instructions.
Taken together, these analyses further confirmed that experimental instructions about nicotine
strength shaped subjective perception in our participants and induced dose-dependent neural
responses in the thalamus, a brain region with one of the highest concentrations of nicotinic
acetylcholine receptors (nAChRs) and a main binding site for nicotine (*9, 10*).
129

130 **Belief about nicotine modulated connectivity between prefrontal cortex and thalamus in a**
131 **dose-dependent manner**
132
At the circuit level, the thalamus is heavily connected to various cortical regions and is known to
contribute to higher-order cognition via these connections (*11*). Thus, we hypothesized that
belief might also modulate the functional connectivity between the thalamus and higher cortical
regions such as prefrontal regions. To this end, we carried out a psychophysiological interaction
(PPI ; see Methods) analysis (*12*) to investigate how beliefs about nicotine affected task-related
functional connectivity between thalamus and rest of the brain. We found that across the whole
brain, belief about nicotine modulated functional connectivity between the thalamus and the
ventromedial prefrontal cortex only (vmPFC; peak at (-11, 50, -6), $P_{uncorrected} < 0.005$, $k = 5$) in a
dose-dependent fashion (**Fig. 3a-b**). The vmPFC is a brain region heavily implicated in the
computation of value and belief updating (*13*). Thus, this result suggests that in addition to
modulating thalamic activation itself, belief about nicotine was also capable of scaling circuit-
level interactions between the thalamus and prefrontal regions involving in higher-level
cognition and decision-making.
146
147 **Discussion**
148
Can beliefs impact brain activity in a dose-dependent manner? Here, we found that neural
responses in the thalamus varied as a function of belief about nicotinic strength. The thalamus —
especially its posterior portion—contains one of the highest densities of nAChRs in the human
brain, quantified by both autoradiography (*14*) and functional imaging (*15*). This anatomical
feature is hypothesized to account for the attention-enhancing effect of nicotine (*15*), as the
thalamus is also known to gate sensory information. Expanding previous work showing acute
dose-dependent responses induced by nicotine in the human thalamus (*16*), our finding
demonstrates that cognitive constructs such as beliefs and expectations can modulate fine-
grained biological mechanisms in the human brain in a way similar to pharmacological agents.
158
We also found that vmPFC-thalamus connectivity scaled as a function of belief about nicotine
strength. The vmPFC has been extensively studied in the context of value-based decision-making
processes, and particularly in computing a "common currency" of subjective value (*17*). In
serving this role, the vmPFC has been shown to receive input from both the ventral tegmental
area and the basal ganglia via the thalamus (*18*). Furthermore, vmPFC-thalamus functional
coupling has been shown to subserve expectations about incoming visual stimuli (*19*). Our
finding expands previous work by demonstrating that instead of a binary "switch", the vmPFC-
thalamus circuit encodes information related to beliefs and expectations in a dose-dependent
manner, highlighting the importance of this circuitry mechanism in high-order cognition.
168
169 In sum, this study provides insight into the dose-dependent impact of abstract concepts such as
170 beliefs, on the fronto-thalamic neural circuitry in a manner resembling pharmacological dose-
171 dependent effects. This finding opens up new avenues for systematically leveraging the impact
172 of narratives on the brain in mental health research and treatment.
173
174
175

Materials and Methods
Participants
The study was approved by the Institutional Review Board of the University of Texas at Dallas
and the University of the Texas Southwestern Medical Center where data collection was
conducted. The sample size was computed by G*Power 3.1.9.7. Assuming a three-measurements
repeated-measures ANOVA F-test with the effect size of 0.4, alpha of 0.05, and power of 0.95,
the minimally required sample size was 18 participants.
183
Nicotine-dependent adult participants were recruited from the Dallas-Fort Worth (DFW)
metropolitan area. Inclusion criteria include an age of 18 years and older, normal or adjusted to
normal vision, and smoking a minimum of 10 cigarettes a day for a period exceeding one year
but with no prior experience with vaping devices or current attempt to quit smoking. Exclusion
criteria included the use of illicit drugs in the past two months, a history of traumatic brain
injury, any current substance abuse (excluding nicotine and alcohol), any contraindication to
MRI, or previous or current psychiatric, neurological, or major medical conditions. Participants
were scanned in accordance with a protocol approved by the UT Dallas and UT Southwestern
institutional review boards after providing written consent.
193
Twenty-three participants enrolled in this study and underwent three fMRI sessions, spaced
about one week apart. Three participants were excluded from analysis for the following reasons:
one participant was excluded due to software malfunction, one due to falling asleep in the
scanner and one due to loss of behavioral data for one of the scanning sessions. The final sample
therefore comprised of 20 participants (6 females, age: 41.1± 11.97 years, age range: 24-61
years). Participants were all right-hand dominant.
200
Experimental design
Upon arrival at the laboratory, participants completed demographic, state (Positive and Negative
Affect Schedule, Beck's Depression Inventory, Drug Abuse Screening Test, Short Michigan
Alcohol Screening Test, EQ, Toronto Alexithymia Scale, Behavioral Inhibition System and
Domain-Specific Risk-Taking questionnaires) and smoking history surveys (Fagerström Test for
Nicotine Dependence, Wisconsin Withdrawal Scale, Shiffman-Jarvik Withdrawal scale) .
Participants provided saliva samples for measuring cotinine, the primary metabolite of nicotine.
Saliva samples were collected using a passive drool method until a volume of 1.8 – 2ml was
obtained. They were then coded and stored in designated freezers until sent for analysis.
Participants' exhaled carbon monoxide (CO) levels served as proxy for their satiety status. These
were acquired by a Smokerlyzer (coVita micro+basic, Santa Barbara, CA) prior to e-cig vaping
in each session. The measurement took place in a designated behavioral testing rooms adjacent
to the scanners. Participants continuously exhaled through a designated straw until a
measurement appeared on screen. For nicotine delivery, we used the Blu e-cig atomizer with
disposable cartridges in the 'classic tobacco' flavor. Following the fMRI scans, participants
repeated the state-based series of surveys and provided a second saliva sample.
217
Three participants' data were removed from cotinine analysis: two due to cotinine readings
exceeding 3 standard deviations from the mean of the cohort and one due to missing data. Data
for all three sessions (instructed belief conditions) were discarded from this analysis. Three
participants' data (non-overlapping with the previous omission) were removed from CO

analysis: two due to readings exceeding 3 standard deviations from times the mean of the cohort
and one due to missing data. Once again, data for all three sessions (instructed belief conditions)
were discarded from this analysis.
225
Prior to vaping, the e-cig cartridge was weighed three times on a high precision scale and the
average of the three measurements was logged as the baseline weight of the cartridge. A similar
procedure was done post-vaping and the change in cartridge weight represented the amount of
nicotine liquid consumed by the participant. Using a double-blind procedure, neither participants
nor the research assistants (M.H. and A.K.) responsible for data collection had prior knowledge
about the true nicotine content in the e-cigs. The order of belief conditions was randomly
assigned for each participant. The e-cig cartridges were carefully re-labelled as 'low', 'medium',
or 'high' by the PI (X.G.) herself to avoid un-blinding by either the participant or the research
assistants. Research assistants (M.H. and A.K.) who interacted with the participants adhered to a
fixed text during the manipulation stating: "The cartridge you will use today will contain: [mild-
to-no nicotine] / [a medium amount of nicotine] / [a high amount of nicotine]." These
experimenters also made sure that participants used the e-cig properly, the device was well
powered, and that vapor was visible. Participants were told they could vape as much as they wish
for the next 20 minutes and were left alone to vape. After 20 minutes they were questioned about
any issues with the e-cig. Participants were then prompted to reply how they would rate the
strength of the nicotine in the e-cig on a scale of 0 to 10, compared to a normal cigarette.
242
<u>Cotinine detection in saliva</u>
Chemicals and reagents: Optima LC-MS grade acetonitrile, water and methanol were purchased
from Fisher Scientific (New Jersey, USA). Reagent grade ammonium formate was purchased
from Sigma Aldrich (Missouri, USA). Cotinine was purchased from Sigma Aldrich. Rac-
cotinine-d3 was obtained from Toronto Research Chemicals (Ontario, CAN). All other
chemicals and reagents were of analytical grade and used without further purification. Blank
human saliva procured from primary investigator. (New York, USA).
250
Preparation of stock and working solutions of analyte and internal standard: Primary stock
solutions of cotinine for the calibration curve (CC) and quality control samples (QC) were
prepared from a 1 mg/mL stock solution in methanol. Stock solutions of cotinine were stored at -
20 °C, and subsequent dilutions were conducted using water. Primary stock solutions of the
bioanalytical method's internal standard (IS) d3-cotinine were prepared by accurately weighing
d3-cotinine and dissolving in methanol to yield a 1 mg/mL stock solution. Stock solutions of d3-
cotinine were stored at -20 °C, and subsequent dilutions were conducted using water. For spiking
of saliva samples with cotinine, a working stock solution of 1000 ng/mL cotinine in water was
prepared and stored at -20 °C. For spiking of saliva samples with d3-cotinine, a working stock
solution of 300 ng/mL d3-cotinine in water was prepared and stored at -20 °C.
261
Preparation of calibration curve and quality control samples for analysis of saliva samples:
Cotinine was validated over a calibration range that supports low concentration and high
concentration samples. The calibration curve ranged from 5 ng/mL to 1000 ng/mL. Calibration
curve samples were prepared by spiking 1µL of the working stock (1 mg/mL cotinine in water)
into 250 µL of saliva. This represented the top calibration curve point (i.e., the upper limit of
quantification or ULOQ). The remaining calibration curve samples were prepared by serial

dilution of the ULOQ standard in saliva. Quality control were prepared in a similar fashion by
spiking 80 µL of the working stock (1000 ng/mL cotinine in water) into 250 µL of plasma. This
represented the high-quality control standard (HQC). The medium-quality control standard
(MQC) and low-quality control standard (LQC) were prepared by serial dilution of the HQC
standard in saliva. Spiking volume of the working standard did not exceed 5% of the matrix
volume. QCs for the calibration curve were prepared at 5, 30, and 80 ng/mL.
274
275 *Saliva Collection:* Patients provided a saliva sample collected at various time-points throughout a
276 session. Samples were collected using a passive drool method (i.e., Salimetrics 'SalivaBio'
277 collection aid). Saliva samples were analyzed for cotinine concentrations using a validated LC-
278 MS/MS method.
279
280 *Saliva Sample Preparation:* Acetonitrile (350 µL) and 100 µL (300 ng/mL) d3-cotinine was
281 added to a 250 µL aliquot of saliva. The resultant mixture was centrifuged for 5 min at 5000 rpm
282 at a temperature of 4°C. Five hundred microliters (500µL) of the clear supernatant, was removed,
283 placed in a new Eppendorf tube, and dried using a SpeedVac. Samples were reconstituted using
284 250 µL of water.
285
HPLC operating conditions: A Shimadzu CBM-20A Nexera X2 series LC system (Shimadzu
Corporation, Kyoto, Japan) equipped with degasser (DGU-20A) and binary pump (LC-30AD)
along with auto-sampler (SIL-30AC) and (CTO-30A) column oven. The autosampler was
maintained at 15 °C. An injection volume of 1 µL was used and chromatographic separation was
achieved using a Kinetex Biphenyl (2.6 µm, 50 × 2.1mm) column. The mobile phase, consisting
of 2mM ammonium formate in water (pump A) and methanol:water (95:5) with 0.2% formic
acid (pump B) used for the method. The mobile phase pumped using a gradient program at a
flow rate of 0.8 mL/min into the mass spectrometer electrospray ionization chamber in positive
polarity. Gradient program initiated with 5% of B and maintained for 1.0min, then ramped to
75%B by 2.5min and maintained at 75%B until 3.5min, changed back to 5%B by 4.0min and
maintained until 6.01min at the end system controller stop command.
Mass spectrometry operating conditions: Quantitation was achieved employing electrospray
ionization in positive ion mode for the analytes using a SCIEX QTRAP 6500+ mass
spectrometer (Redwood, CA, USA) equipped with the Turbo V source operated at 550 °C. The
nebulizer gas, auxiliary gas, curtain gas, CAD gas were set at 45, 45, 30 psi and 'medium',
respectively. The declustering potential (DP), collision energy (CE), entrance potential (EP) and
collision cell exit potential (CXP) were 141, 31, 10, 10 V for (Cotinine-1); 141, 47, 10, 8 V for
(Cotinine-2); 141, 31, 10, 12 V for (d3-Cotinine-1); and 141, 29, 10, 8 V for (d3-Cotinine-3),
respectively. Detection of the ions was carried out in the multiple-reaction monitoring mode
(MRM), by monitoring the precursor > product transitions of 177.0 > 80 and 177.0 > 98.0 (sum
over 2 MRMs) for cotinine and 181.2 > 80.1 and 181.2> 101.1 (sum over 2 MRMs) for d3-
cotinine. The data obtained were processed by Analyst software™ (version 1.6.3).
308
309 *Method Validation:* The methods for analysis of cotinine in saliva were validated according to
310 the United States FDA's May 2018 Guidance for Industry on 'Bioanalytical Method Validation'.
311 The method was found to have acceptable sensitivity, selectivity, matrix effect, linearity,
312 accuracy, precision, recovery, dilution integrity and stability.
313

Decision-making task
This task was developed based on a previous investment task (*4*, *5*), but with the modification
that participants were allowed to place both positive ('invest') and negative ('short') bets.
Briefly, participants were allocated an initial sum of 100 monetary units (i.e., their portfolio) at
the beginning of the experiment which could be invested in stock markets. Participants were
informed that their final payment would be scaled according to their actual gains or losses in the
task. Each participant played a total of 10 markets per visit, each consisting of 20 trials. The
stock market prices in the task were chosen from true historical stock market prices. Each task
block commenced by a caption titled 'new market' followed by a graphic display of past market
dynamics. In each trial t, the participant observes the price history of a stock market (including
the trial before, p_{t-1}) and places a bet, b_t. Next, a new market price p_t is revealed, and portfolio
amounts are updated to reflect the recent outcome. The fractional market return r_t ,is defined as r_t
$= (p_t - p_{t-1}) / p_{t-1}$. In each of the 20 trials, participants had unconstrained time to decide on their
investment moves. Participants were able to choose to either invest normally (if they think the
price will go up) or short sell (if they think the price will go down). Notably, shorting the market
would result in gaining from market drops. Thus, people could benefit from either a positive or
negative price change and the absolute value of market return $|r_t|$ represents the magnitude of the
potential gain. Participants provided their choice using a slider bar and finalized their choice by a
button press. Following a 750 ms delay, the new market price was revealed and the fractional
change in market price was applied to the portfolio. In later analyses this event is termed 'market
reveal'. Each trial concluded in a 750 ms inter-trial interval in which the slider turned gray and
became unresponsive. A total of 30 different markets were used across all three visits. Mean
session duration in the stock market task was 14.91 ± 3.06 minutes and did not differ across
conditions ($F(2,59) = 0.28$, $P = 0.76$).
338
Image acquisition and preprocessing
Whole-brain anatomical and functional MRI data were acquired on a Philips Achieva scanner
with a 3T field strength. High-resolution T1-weighted scans ($1.0 \times 1.0 \times 1.0$ mm) were acquired
using a 3D magnetization prepared rapid gradient-echo (MPRAGE) sequence. Functional images
were acquired using echo-planar imaging (EPI) and tilted 30° from AC-PC axis. The detailed
settings for the functional imaging were repetition time (TR) = 2,000 ms, echo time (TE) = 25
ms, flip angle = 90°, voxel size = $3.4 \times 3.4 \times 4.0$ mm, 38 slices. The average number of
functional images acquired was 457.37 ± 91.67. All imaging data were preprocessed using
standard statistical parametric mapping (SPM12, Wellcome Department of Imaging
Neuroscience) algorithms *(fil.ion.ucl.ac.uk/spm)*. Functional images were applied a slice time
correction.
350
To account for large head movements often caused by participants' coughing during the scan, we
used the ArtRepair toolbox to examine and repair volumes with large motion artifacts. We used
the *art_motionregress* and *art_global* modules of the single subject pipeline. The ArtRepair
algorithm was further used to generate the motion parameters to be included in the GLM design
matrix. Volumes were examined for fast head movements using the automated defaults such that
volumes with movement of >0.5 mm/TR were tagged and interpolated with the nearest usable
volumes. Overall, 6 out of 60 scans were repaired. The mean functional images for each subject
were co-registered to the subject's high-resolution T1 structural scan, using a 12-parameter
affine transformation. The participant's T1 image was segmented into gray and white matter and

then normalized using nonlinear basis functions to the Montreal Neurological Institute (MNI)
space with the functional images normalized to the template and resampled into $3.4 \times 3.4 \times 3.4$-
mm functional voxels. Functional images were smoothed spatially using a 6 mm full-width at
half-maximum Gaussian kernel. A temporal high-pass filter of 128 Hz was applied to the fMRI
data, and temporal autocorrelation was modeled using a first-order autoregressive function.
365
Statistical analysis
Throughout this study, we used a within-participant repeated measures ANOVA implemented in
MATLAB (*anova_rm*) to assess differences between the three conditions of instructed belief.
Normality was assessed with Shapiro-Wilk tests wherever appropriate. During analysis of the
various controls if data for one of the sessions was missing, that participant was excluded from
this specific analysis.
372
373
374 Behavioral modeling
375 We carried out linear mixed-effects multiple regression analyses on the series of market returns
376 and investment decisions extracted from each subject. Multiple regression was carried out in R
377 (RStudio 1.1.463, 2018) using the 'lmer' function as follows:
378
379 $|b_{t+1}| \sim 1 + |b_t| + |r_t| + $ *instructed beliefs* $ + ($*instructed beliefs* $* |b_t| \,|\, $ *participant*$)$
380
381 The final return of each market was excluded from the regression, as there is no investment
382 decision following the final market segment. Similarly, the first trial was also removed since it
383 had no preceding investment decision. *P* values were approximated via Satterthwaite's degrees
384 of freedom method. Linear mixed effects were estimated using the full information maximum
385 likelihood (FIML) method.
386
General linear modeling (GLM) of fMRI data
Event-related analyses of the fMRI data were conducted using SPM12 (Wellcome Department of
Imaging Neuroscience). We conducted general linear modeling (GLM) of the functional scans of
each participant bey modeling the observed BOLD signals and regressors to identify the
relationship between the task events and the hemodynamic response. Regressors related to all
visual and motor events were created by convolving a train of delta functions representing the
sequence of individual events with the default basis function in SPM12, which consists of a
synthetic hemodynamic response function composed of two gamma functions. The GLM
included six separate regressors: (1) new market screen display; (2) market history display; (3)
all key presses; (4) market price reveal of trial 1; (5) market price reveal of trials 2–19; (6)
market price reveal of trial 20. Additionally, six parameters generated during motion correction
were entered as covariates. In the GLM, absolute market return $|r|$ was entered as a parametric
modulator of market reveal of trials 2–19. We carried out linear contrasts of the parameter
estimates to identify the effects in each participant. Statistical maps from all participants were
then entered into a second-level group analysis to implement a random-effects statistical model.
A within-subject repeated-measures ANOVA model was conducted for the factor of instructed
beliefs ('low', 'medium', 'high'). Statistical inference was made based on the F statistics derived
from whole-brain rmANOVA statistical maps. Significant effects were identified at the level of
$P < 0.005$, with a cluster size threshold of $k = 50$. Maps were rendered using MRIcroGL.

406
<u>Thalamic parcellation and ROI analyses</u>
We extracted parameter estimates from bilateral thalamus using an anatomical mask (WFU Pick
atlas (*20*). Thalamic parcellations were obtained from the Lead-DBS MATLAB toolbox. Each
segmented nucleus or region was transformed into the experimental dataset's functional space
using the MarsBaR toolbox. The THOMAS (Thalamus optimized multi-atlas segmentation) atlas
(*21*) contains 12 non-overlapping nuclei, two of which (MGN and MTT) were too small to be
meaningfully transformed to our functional space and were therefore not used. As before, we
modeled BOLD activity tracking of fluctuations in magnitude of market return, |r|, and carried
out a group analysis with a within-subject rmANOVA design per region of the thalamic
parcellation. To account for multiple comparisons, we applied the false-detection rate (FDR)
correction to the extracted ROIs at $q = 0.05$.
418
419
<u>Permutation analysis</u>
We iteratively shuffled labels for instructed beliefs ('low', 'medium', 'high') within each
participant while maintaining their original ratio (i.e., one of each per participant). For each
viable permutation (i.e., a permutation whose model estimation converged and yielded any
significant voxels), a within-subject rmANOVA was carried out, following which a parameter
estimate was derived from the same ROI as the original design matrix to generate a surrogate
distribution of beta estimates ($N = 1,000$).
427
<u>Psycho-physiological interaction (PPI) analysis</u>
PPI analysis provides a measure of change in functional connectivity between different brain
regions under a specific psychological context (*12*). We defined a seed region – the thalamus –
as defined by the WFU anatomical atlas and a psychological context ('market reveal' – the
presentation of the investment's return). We then conducted a whole-brain PPI analysis per
condition of instructed beliefs and compared those in a within-subject repeated-measures design.
The generated PPI model included the PPI term, the physiological regressor, the psychological
regressor, and nuisance regressors of six motion parameters. The threshold of significance for the
group-level rmANOVA from the PPI regressor was set to be the same as in the GLM ($P < 0.005$)
with a cluster size threshold of $k = 5$.
438
439 **Data availability**
440 Data supporting the findings of this study are deposited in: https://osf.io/3hq6s/
441
Code availability
The scripts used for data acquisition and analysis are available in: https://osf.io/3hq6s/
Analyses were conducted using open software and toolboxes available online as described in
Materials and Methods (SPM: www.fil.ion.ucl.ac.uk/spm/software/spm12; R Studio:
https://www.rstudio.com/products/rstudio/download/#download; Lead-DBS: https://www.lead-
dbs.org/download/; MRIcroGL: https://www.nitrc.org/projects/mricrogl/)
448
449
450
451

Acknowledgements
We thank staff members at the UT Southwestern Imaging Center for their assistance with
scanning, Jennifer Jung and Mark Labinski for their help with developing the fMRI task, and
Jessica Maclin for her help with participant recruitment. We also thank Dr. Laura Berner for her
advice on the statistical analysis, and Drs. Daniela Schiller and Paul Kenny for their helpful
discussions and comments on the manuscript.
459

Author Contributions
Conceptualization: XG.
Methodology: OP, AS, NB, WCP, XG
Investigation: MH, SN, AK
Visualization: OP
Funding acquisition: XG
Project administration: MH, XG
Supervision: XG
Writing – original draft: OP, WCP, VGF, XG
Writing – review and editing: OP, VGF, XG

Competing interests
The authors declare that they have no competing interests.
472

Funding
National Institute on Drug Abuse grant R01DA043695 (XG)
National Institute on Drug Abuse grant R21DA049243 (XG)
University of Texas, Dallas internal funding (XG)
Mental Illness Research, Education, and Clinical Center at the James J. Peter Veterans Affairs
Medical Center grant MIRECC VISN 2 (VGF)
479

480 **Additional information**
481 Supplementary Text
482 Figs. S1 to S7
483 Tables S1
484

References
1. N. D. Volkow, M. Morales, The Brain on Drugs: From Reward to Addiction. *Cell*. **162**,
712–725 (2015).
2. E. R. Korpi, B. den Hollander, U. Farooq, E. Vashchinkina, R. Rajkumar, D. J. Nutt, P.
Hyytiä, G. S. Dawe, *Pharmacol. Rev.*, in press, doi:10.1124/pr.115.010967.
3. T. D. Wager, J. K. Rilling, E. E. Smith, A. Sokolik, K. L. Casey, R. J. Davidson, S. M.
Kosslyn, R. M. Rose, J. D. Cohen, Placebo-induced changes in FMRI in the anticipation
and experience of pain. *Science*. **303**, 1162–1167 (2004).
4. X. Gu, T. Lohrenz, R. Salas, P. R. Baldwin, A. Soltani, U. Kirk, P. M. Cinciripini, P. R.
Montague, Belief about nicotine selectively modulates value and reward prediction error
signals in smokers. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 2539–2544 (2015).

5.   T. Lohrenz, K. McCabe, C. F. Camerer, P. R. Montague, Neural signature of fictive
learning signals in a sequential investment task. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 9493–
9498 (2007).
6.   P. H. Chiu, T. M. Lohrenz, P. R. Montague, Smokers' brains compute, but ignore, a
fictive error signal in a sequential investment task. *Nat. Neurosci.* **11**, 514–520 (2008).
7.   X. Gu, T. Lohrenz, R. Salas, P. R. Baldwin, A. Soltani, U. Kirk, P. M. Cinciripini, P. R.
Montague, Belief about nicotine modulates subjective craving and insula activity in
deprived smokers. *Front. Psychiatry.* **7**, 1–11 (2016).
8.   G. O. Petersen, C. E. Leite, J. M. Chatkin, F. V. Thiesen, Cotinine as a biomarker of
tobacco exposure: Development of a HPLC method and comparison of matrices. *J. Sep.*
Sci. **33**, 516–521 (2010).
9.   S. Wonnacott, Presynaptic nicotinic ACh receptors. *Trends Neurosci.* **20**, 92–98 (1997).
10. D. Paterson, A. Nordberg, Neuronal nicotinic receptors in the human brain. *Prog.*
Neurobiol. **61**, 75–111 (2000).
11. G. M. G. Shepherd, N. Yamawaki, Untangling the cortico-thalamo-cortical loop: cellular
pieces of a knotty circuit puzzle. *Nat. Rev. Neurosci.* **22**, 389–406 (2021).
12. K. J. Friston, C. Buechel, G. R. Fink, J. Morris, E. Rolls, R. J. Dolan, Psychophysiological
and Modulatory Interactions in Neuroimaging. *Neuroimage.* **6**, 218–229 (1997).
13. M. F. S. Rushworth, T. E. J. Behrens, Choice, uncertainty and value in prefrontal and
cingulate cortex. *Nat. Neurosci.* **11**, 389–397 (2008).
14. D. P. Spurden, J. A. Court, S. Lloyd, A. Oakley, R. Perry, C. Pearson, R. G. L. Pullen, E.
K. Perry, Nicotinic receptor distribution in the human thalamus: Autoradiographical
localization of [3H]nicotine and [125I]α-bungarotoxin binding. *J. Chem. Neuroanat.* **13**,
105–113 (1997).
15. N. S. Lawrence, T. J. Ross, E. A. Stein, Cognitive mechanisms of nicotine on visual
attention. *Neuron.* **36**, 539–548 (2002).
16. E. A. Stein, J. Pankiewicz, H. H. Harsch, J. Cho, S. A. Fuller, R. G. Hoffmann, M.
Hawkins, S. M. Rao, P. A. Bandettini, A. S. Bloom, Nicotine-Induced Limbic Cortical
Activation in the Human Brain : A Functional MRI Study. *Am. J. Psychiatry.* **155**, 1009–
1015 (1998).
17. P. W. Glimcher. D.J. Levy, Root of all value. *Changes.* **29**, 997–1003 (2012).
18. S. N. Haber, B. Knutson, The reward circuit: Linking primate anatomy and human
imaging. *Neuropsychopharmacology.* **35**, 4–26 (2010).
19. G. Barbalat, N. Bazargani, S.-J. Blakemore, The Influence of Prior Expectations on
Emotional Face Perception in Adolescence. *Cereb. Cortex.* **23**, 1542–1551 (2012).
20. J. A. Maldjian, P. J. Laurienti, R. A. Kraft, J. H. Burdette, An automated method for
neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets.
Neuroimage. **19**, 1233–1239 (2003).
21. J. H. Su, F. T. Thomas, W. S. Kasoff, T. Tourdias, E. Y. Choi, B. K. Rutt, M. Saranathan,
Thalamus Optimized Multi Atlas Segmentation (THOMAS): fast, fully automated
segmentation of thalamic nuclei from structural MRI. *Neuroimage.* **194**, 272–282 (2019).
22. W. Schultz, P. Dayan, P. R. Montague, A Neural Substrate of Prediction and Reward.
Science. **275**, 1593–1599 (1997).
539

Perl et al.

540
Fig. 1. Experimental paradigm and sanity check measures. (**a**) Participants completed three
visits. In each, we collected saliva samples for cotinine measurement, measured carbon
monoxide (CO) levels, instructed beliefs, and measured brain activities using fMRI as
participants engaged in a decision-making task. (**b**) Change in cartridge weight (pre – post) (P =
0.178), (**c**) cotinine concentration (P = 0.393), or (**d**) CO level (P = 0.698) did not differ between
conditions. (**e**) Effect of instructed beliefs on self-reported belief about nicotine strength (P =
0.0004). Bars depict group means and points represent participants. Error bars are SEM. ***P <
0.001.

549
Fig. 2. Belief about nicotine strength induced dose-dependent responses in the thalamus.
(**a**) Whole-brain effects of instructed beliefs about nicotine on value-tracking signals (cluster-
level P_{FWE} = 0.006, k = 50). (**b**) Parameter estimates representing reward-related activities
extracted from an independent thalamus mask (shown in purple) across belief conditions (P =
0.036). Bars depict group means, points represent participants. Error bars are SEM. *$P < 0.05$.
(**c**) Permutation analysis for instructed beliefs conditions (N = 1,000, P = 0.002). A histogram
comprised of surrogate distribution for beta estimates (black bars). Red line denotes mean of true

beta values. (**d**) Correlation between thalamic signals and subjective belief rating regarding perceived nicotine strength (r = 0.27, *P* = 0.035). Black dashed line is linear fit.

Fig. 3. Belief about nicotine strength modulated thalamus-vmPFC functional connectivity in a dose-dependent fashion. (**a**) Whole-brain effects of instructed beliefs on the psychophysiological interaction (PPI) between the thalamus and rest of the brain (*P* < 0.005, uncorrected, k = 5). (**b**) Parameter estimates extracted from (a) representing functional coupling strength between the thalamus and vmPFC. Bars depict group means, points represent participants. Error bars are SEM.