Urban birds' tolerance towards humans was largely unaffected by increased variation in

human levels due to COVID-19 shutdowns

Peter Mikula[1,2,3*], Martin Bulla[1,4*], Daniel T. Blumstein[5], Yanina Benedetti[1], Kristina Floigl[1],

Jukka Jokimäki[6], Marja-Liisa Kaisanlahti-Jokimäki[6], Gábor Markó[7], Federico Morelli[1,8],

Anders Pape Møller[9,10], Anastasiia Siretckaia[1], Sára Szakony[11], Michael A. Weston[12], Farah

Abou Zeid[1], Piotr Tryjanowski[2,3,13†], Tomáš Albrecht[14,15†]

[1]Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Kamýcká 129,

16500 Prague, Czech Republic

[2]TUM School of Life Sciences, Ecoclimatology, Technical University of Munich, 85354,

Freising, Germany

[3]Institute for Advanced Study, Technical University of Munich, 85748, Garching, Germany

[4]Department of Behavioural Ecology and Evolutionary Genetics, Max Planck Institute for

Biological Intelligence, Eberhard Gwinner Str. 7, 82319 Seewiesen, Germany

[5]Department of Ecology and Evolutionary Biology, University of California, 621 Young Drive,

South, Los Angeles CA 90095, USA

[6]Arctic Centre, University of Lapland, PO Box 122, 96101, Rovaniemi, Finland

[7]Department of Plant Pathology, Institute of Plant Protection, Hungarian University of

Agriculture and Life Sciences, Ménesi út 44, 1118, Budapest, Hungary

[8]Institute of Biological Sciences, University of Zielona Góra, Prof. Z. Szafrana St. 1, 65516,

Zielona Góra, Poland

[9]Ecologie Systématique Evolution, Université Paris-Sud, CNRS, AgroParisTech, Université

Paris-Saclay, 91405, Orsay Cedex, France

[10]Ministry of Education Key Laboratory for Biodiversity Sciences and Ecological Engineering,

College of Life Sciences, Beijing Normal University, 100875, Beijing, China

[11]Department of Ecology, Institute of Biology, University of Veterinary Medicine Budapest,

Rottenbiller u. 50., 1077, Budapest, Hungary

[12]Deakin Marine, School of Life and Environmental Sciences, Deakin University, Burwood

Campus, 221 Burwood Highway, VIC 3125, Burwood, Australia

[13]Institute of Zoology, Poznań University of Life Sciences, Wojska Polskiego 71C, 60625,

Poznań, Poland

[14]Institute of Vertebrate Biology, Czech Academy of Sciences, Květná 8, 60365, Brno, Czech

Republic

[15]Department of Zoology, Faculty of Science, Charles University, Viničná 7, 12844, Prague,

Czech Republic

Authors for the correspondence: Peter Mikula (petomikula158@gmail.com); Martin Bulla

(bulla.mar@gmail.com); Tomáš Albrecht (albrecht@ivb.cz)

*These authors contributed equally to this work

†These authors jointly supervised this work

ORCID:

Peter Mikula: https://orcid.org/0000-0002-2731-9105

Martin Bulla: https://orcid.org/0000-0002-1328-1927

Yanina Benedetti: https://orcid.org/0000-0003-1600-2310

Daniel T. Blumstein: https://orcid.org/0000-0001-5793-9244

Jukka Jokimäki: https://orcid.org/0000-0002-7903-4128

Gábor Markó: https://orcid.org/0000-0003-1351-4070

Federico Morelli: https://orcid.org/0000-0003-1099-1357

Piotr Tryjanowski: https://orcid.org/0000-0002-8358-0797

Michael A. Weston: https://orcid.org/0000-0002-8717-0410

Tomáš Albrecht: https://orcid.org/0000-0002-9213-0034

Abstract

The coronavirus disease 2019 (COVID-19) pandemic dramatically altered human activities, and, during shutdowns. Here, we evaluated whether urban birds from five countries changed their tolerance towards humans during the COVID-19 shutdowns. We collected 6369 flight initiation distance estimates for 147 bird species and found that human numbers in parks (at a given hour, day, week or year - before and during the COVID-19 shutdowns) had a little effect on birds' tolerance of approaching humans. Apart from the actual human numbers in the area (hourly scale), the effect of human activity at other temporal scales centered around zero. The results were similar across countries, for most species or when we restricted our analyses only to species sampled both before and during the COVID-19 shutdowns. As expected, the level of daily human presence in parks (measured by Google Mobility Reports) correlated negatively with the stringency of governmental restrictions (a weekly proxy for human presence) and was overall lower during COVID-19 shutdowns than during the post-shutdown year (2022). Our results highlight the resilience of birds to changes in human numbers on multiple temporal scales, the complexities of linking animal fear responses to human behavior, and the challenge of quantifying both simultaneously *in situ*.

Keywords: antipredator behavior; escape distance; flight initiation distance; human presence, lockdown; urbanization

Introduction

The actions taken to control the coronavirus disease 2019 (hereafter COVID-19) pandemic locked inhabitants in their dwellings and thus changed the pattern of human outdoor activities [1–9]. This situation created a quasi-experiment that offered a unique opportunity to study how rapid changes in human behavior affect wildlife [10–14]. The shifts in human presence and activity associated with the COVID-19 pandemic might have been particularly noticeable in urban areas [12,15]. Consequently, the COVID-19 shutdowns might have elicited complex effects on urban nature [10]. For example, some animals increased their abundances in human-dominated landscapes, moved to new areas, or shifted the timing and character of their activities; some of these effects were area- or species-specific [10,11,16–23]

To persist in urban habitats, wild animals must increase their tolerance to humans and human disturbance [24,25]. Indeed, on a gradient from natural to urban environments animals seem more tolerant of humans, where human densities are high and human–wildlife interactions are generally harmless [24,26–29]. Additionally, within cities animal tolerance to humans seems to increase with increasing level of human presence (hereafter "human presence") but the association is often weak [30–33]. Whether the COVID-19 shutdowns induced changes in human activities and subsequently altered the urban landscape of fear and animal responses to human presence is still poorly understood [22,34].

Here, we explored how the COVID-19 shutdowns altered avian tolerance towards humans across parks of five cities in five countries (Finland – Rovaniemi, Poland – Poznan, Czechia – Prague, Hungary – Budapest, Australia – Melbourne). During the breeding seasons before the COVID-19 shutdowns (before March 2020) and during the COVID-19 shutdowns (2020–2021), we measured avian tolerance towards humans by quantifying flight initiation distance, which is the distance from an approaching human at which a bird escapes [26,28,35–38]. Because the COVID-19 interventions increased the variation in human presence outdoors [39–

41] and the temporal scale at which birds evaluate the changes in the landscape of fear and adjust their response is unknown, we tested how changes in human presence across multiple temporal scales (hour-to-hour, day-to-day, week-to-week and year-to-year) were associated with avian escape distance. We used (1) actual human numbers recorded during escape distance observation as a measure of hourly changes in human presence, (hereafter "Number of humans"), Google Mobility Reports data (hereafter "Google Mobility", https://www.google.com/covid19/mobility) as a measure of daily changes in human presence in parks, the stringency of governmental restrictions (hereafter "Stringency index"; https://ourworldindata.org/covid-stringency-index; based on data originally published in [42]) as a proxy for weekly changes in human presence, and years before COVID-19 shutdowns and years during COVID-19 shutdowns as a coarse proxy for yearly changes in humans presence (hereafter "Period"), which we validated with the other three, above described proxies of human presence. Although the baseline escape distance (the intercept) and the strength of association (the slope) between flight initiation distance and human presence may vary across species, we expected that if individuals, regardless of species, responded to shutdowns, then this would induce a similar response. In other words, we expected a negative relationship between escape distance of birds and human presence. However, if shutdowns affected human presence in a country-specific manner, we would expect country-specific effects of shutdowns on avian tolerance. Moreover, if shutdowns had little effect on human behavior, we expected no change in avian tolerance.

Results

We found no major and clear differences in the overall avian tolerance towards approaching humans on all temporal scales (Fig. 1). Specifically, (1) according to the expectation, the actual

human numbers in the area when an escape trial was conducted (hourly temporal scale)

correlated negatively, albeit weakly, with escape distance. (2) The relative human numbers

within each day as measured by Google Mobility (daily temporal scale) and (3) weekly changes

in human numbers as proxied by Stringency index (weekly temporal scale) were unrelated to

avian escape distance. Finally, (4) avian escape distances were similar in the years before and

during COVID-19 shutdowns (hereafter referred to as "Period").

The results were similar across countries (Fig. 1; Table S1), and robust to the presence and

absence of the control for starting distance (i.e. the distance between a bird and an observer

when the escape distance trial started; Fig. 1), the changes in random effects structure of the

models, or, in the case of Period, restriction of the analyses only to species sampled during both

periods (Fig. S1; Table S2). Moreover, the effect sizes from the global model, where all data

were analyzed together, were similar to the meta-analytical effect sizes based on country-

specific estimates (Fig. 1).

Altogether, while the response of humans to shutdowns was country-specific and the presence

of humans in parks decreased in some countries, increased in others, or remained unchanged

[7,8,43], avian tolerance, as we measured it, did not reflect such major changes (Fig. 1). The

country-specific effect sizes are small and mostly center around zero (Fig. 1). The effects were

also inconsistent within countries and sites as well as within species (Fig. S3–S10). In other

words, urban birds were largely inflexible in their escape distance in relation to changes in

human presence.

When exploring the patterns of human presence, we found that the actual number of humans at

the time of an escape trial correlated weakly, and in a country specific manner, with relative

daily human presence (Google Mobility; Fig. 2a) and weekly human presence (Stringency

index; Fig. 2b). As expected, (i) daily human presence generally correlated negatively, albeit

weakly, with weekly human presence (Fig. 3), and (ii) relative daily human presence was

generally lower during years with COVID-19 shutdowns (2020, 2021) than in 2022, a post-

shutdown year (Fig. 4). However, day-to-day variation in human presence seems larger (Fig. 4,

middle plots) than changes induced by shutdowns (Fig. 4, left and right plots). In other words,

the COVID-19 shutdown's influence on between-year differences in human numbers might

have been negligible in studied countries. Yet, we cannot also exclude a possibility that the

shutdowns changed human activities; hence, our year-to-year findings may reflect resilience of

the birds to changes in human activities. Also, data on Google Mobility prior to the COVID-19

shutdowns were unavailable and thus we do not know the true between-year reduction of human

presence due to shutdowns.

Figure 1. Avian tolerance towards humans across four temporal scales. Avian tolerance according to human levels during (a) the escape distance trial (hourly scale) measured as number of humans within a 50-meter radius, (b) the day of the escape trial as proxied by Google Mobility (daily scale), (c) week of the escape trial as proxied by the stringency of governmental measures (weekly scale), and (d) the period of the escape trial, i.e. before vs during the COVID-19 shutdowns (yearly scale). (a-d) The dots with horizontal lines represent estimated standardized effect size and their 95% confidence intervals, the numbers represent sample sizes. For the **countries** (i.e. country-specific models) and **"All"** (i.e. global models containing all countries), the effect sizes and 95% confidence intervals come from the joint posterior distribution of 5000 simulated values generated by the *sim* function from the *arm* package [44] using the mixed model outputs controlled for starting distance of the observer (filled circles) or not (empty circles; Table S1). The models were further controlled for year (in case of Period fitted as a random intercept), flock size (ln-transformed), body size (ln-transformed), temperature (also a proxy for a day within the breeding season: $r_{Pearson} = 0.48$; Fig. S2), and time of a day, as well as for the non-independence of data points by fitting random intercepts of day of the week, genus, species, species at a given day and year, country (in "All"), site, and species within a site, while fitting, in case of "All" (global model on all countries) Number of humans, Google Mobility, Stringency index, or Period as a random slope within country (i.e. allowing for different effect in each country) and in case of Period models (country-specific or "All") also within site (i.e. allowing for different Period effect at each site).

Fitting Number of humans, Google Mobility, Stringency index or Period as random slope at other

random intercepts produces similar results (Fig. S1, Table S2). The multicollinearity was small as

correlations between predictors were weak (Fig. S2). For the **"Combined"**, the estimates and 95%

confidence intervals represent the meta-analytical means based on the country-specific estimates and

their standard deviation (from the country-specific models), and sample size per country. Note that the

effect sizes are small and estimates tend to center around zero.

Figure 2. Numbers of humans within 50 meters during bird escape trial in association with daily human

presence in parks (Google Mobility, left) and stringency of anti-COVID governmental restrictions

(Stringency index, right). Dots represent individual data points (top panels on original scale, bottom on

ln-scale), jittered to increase visibility. Lines with shaded areas represent predictions with 95% CIs from

mixed effect models that controlled for the year (in case of Finland and Hungary) and non-independence

of data points by including day of the week within the year as a random intercept and Stringency index

or Google Mobility as a random slope (Table S3a–d). Human numbers during shutdowns were missing

for Australia and Poland. Note the weak and country-specific associations.

Figure 3. Association between human presence in parks (Google Mobility) and the stringency of anti-COVID governmental restrictions (Stringency index). Lines with shaded areas represent predicted relationships from country-specific mixed effect models controlled for the year and non-independence of data points by including weekday within the year as a random intercept and Stringency index as a random slope (Table S3e). Dots represent raw data, jittered to increase visibility, for days when we collected escape distances in each city. Colors indicate country. Note the generally negative but weak association between the Google Mobility and the Stringency index.

Figure 4. Changes in human presence in parks within and between years and countries. Left plots represent distributions (histograms) of daily human presence (Google Mobility), middle plots the raw daily data across years, and right plots locally estimated scatterplot smoothing (Loess). Dotted vertical line in "Distribution" plots indicates baseline value of human presence, separating negative values that represent decreased human presence and positive values that indicate increased human presence when compared with the baseline. The baseline is country and weekday specific (for weekday-specific patterns, see Fig. S11). Note that Google Mobility data were unavailable for the years before the COVID-19 pandemic (i.e. before 2020) but the year 2022 was without shutdowns in the studied countries.

Discussion

We used (i) flight initiation distance as a measure of avian tolerance toward humans and capitalized on (ii) COVID-19 shutdowns that induced variation in human levels in parks to

reveal that, in general, urban bird populations exhibited no major shifts in their tolerance toward

humans in response to fluctuations in human levels across different temporal scales, i.e. hourly

(actual number of humans during the experimental trial), daily (proxied by Google Mobility),

weekly (proxied by Stringency index), and yearly (proxied by whether observation was

conducted in years before or during COVID-19 shutdowns). These findings were largely

consistent across countries, and independent of model specification. The trends for avian

tolerance towards humans to increase with the increasing hourly (and to lesser extent also daily)

human levels outdoors were small and weak.

Avian responses to human levels across different timescales

The weak trends for birds to increase their tolerance to humans (i.e. tolerate closer approaches

by humans) when human levels increased within short temporal scales, as well as the species-

and area-specific nature of this effect that we report, are in line with previous studies [32,33,45–

47]. Altogether, these results indicate that, in some situations, urban birds may be able to

flexibly adjust their tolerance to short-term changes in human levels. Behavioral flexibility

plays a role in short-term adjustments [31,46,48], although our large-scale and multi-taxonomic

comparison indicates that the room for behavioral adjustments may be limited for urban birds

(e.g. because environmental filtering has left urban areas with mainly bolder individuals; see

below).

Our findings that urban birds do not seem to react to the change in human levels on longer

temporal scales, such as between weeks or years, is novel as we are not aware of a single study

which directly tests for such effects. Rather, previous studies investigated whether animal

tolerance has changed over time regardless of actual human levels, e.g. with the time–since

urbanization [49,50]. However, a recent study using individually marked dark-eyed juncos

(*Junco hyemalis*) showed, similar to our findings, weak effects of the COVID-19 shutdowns on

flight initiation distance in urban birds [22]. The weak response of birds to long-term changes

in human levels may indicate that (i) birds react to long-term shifts in human behavior in a non-

linear manner (which has to our knowledge never been tested) and/or such (ii) shifts in human

behavior have to reach a threshold value to trigger observable changes in bird behavior. Thus,

the changes in human behavior induced, for example by the COVID-19 shutdowns, might have

been insufficient to reach such threshold, since human presence in parks greatly fluctuates over

the day, across weeks and seasonally (Fig. 4; e.g. [7,39–41]), perhaps explaining why the

occurrence of wildlife in the cities did not always increase during the COVID-19 shutdowns

[17,19]. To test whether the response of birds to changing human levels is non-linear or based

on threshold human levels, repeated escape distance trials of marked individuals during various

times of day and different days of the week, as well as under various natural or experimentally

induced levels of human presence and activity are needed. However, most studies that

investigated escape responses of animals towards humans and used marked individuals did not

expose these individuals to the various levels of human disturbance ([48,51–53], but see [22]).

Our findings also highlight the socio-ecological complexities of linking wildlife and human

behaviors. In other words, the within-day and day-to-day fluctuations in human presence and

outdoor activities raise questions of how and when to measure human presence and activity in

field studies on animal tolerance towards humans. For instance, we still do not understand

whether animals respond to the overall daily human presence and activity, or whether they

flexibly adjust their behavior to levels of human presence and activity at a given time. We also

acknowledge the substantial evidence that birds can discriminate between different forms of

human activity, meaning that metrics of human abundance may not directly reflect perceptions

of risk by birds [54–56]. Finally, we acknowledge that some of our data were not designed to

answer questions about short-term (mainly hourly) adjustments. Thus, it is possible that despite

high human densities at some days of the week or during specific times of the day, animals may

shift their activities accordingly (e.g. forage when human activity is low or rest and hide when

human presence is high).

Possible mechanisms behind the weak response

The general lack of response to the changes in human levels we described, has three possible

implications. (i) Most urban bird species in our sample may be relatively inflexible in their

escape responses because the species may be already adapted to human presence. In other

words, natural selection on escape behavior of sampled urban birds might be at an optimum,

and departing from this may be difficult. Moreover, bird species that colonized urbanized areas

relatively earlier are more tolerant toward humans and also their escape responses to human

approaches are less variable [49]. Our results might also indicate that the room for current

(ongoing) selection is limited even when human presence changes on relatively long temporal

scale such as in the case of the COVID-19 shutdowns. However, some studies reported changes

in space use by wildlife [18,20,21], and these could arise, as our results indicate, from fixed and

non-plastic animal responses to humans who changed their activities.

(ii) Urban environment might act as a filter for bold individuals [52,53,57]. Thus, the lack of

consistent changes in the escape behavior of urban birds may indicate an absence (or low influx)

of generally shy, less tolerant individuals and species from rural or less disturbed areas into the

cities. In other words, the urban environment has filtered birds based on their inherent levels of

tolerance in a similar way regardless of variation in human levels. Indeed, based on genetic

studies, in some species, rural populations are a source of individuals to urban populations

[58,59].

(iii) Urban birds might have been already habituated to or become tolerant of variation in the

levels of human presence, irrespective of the potential changes in human activity patterns [60–

62]. Urban animals generally habituate to human disturbance quickly and once habituated may

show only a small variation in their tolerance [48,63]. Indeed, we found that even corvids (Fig.

S3–S6), generally considered to have high levels of behavioral plasticity, do not deviate from

trends observed in other species. Moreover, tolerance of individually-marked dark-eyed juncos

changed little during COVID-19 shutdowns, although individuals became more tolerant

(shortened their escape distance) after shutdowns when the university campus study site was

repopulated with humans [22]. Such findings are puzzling, but indicate that habituation-like

processes [64] might have already reduced plasticity. That said, there were between-site

differences in the escape distance of the same species (Fig. S7–S10) and we thus need further

studies of marked individuals to better understand this phenomenon.

Conclusions

In sum, our findings highlight that urban birds might be most sensitive to immediate variation

in human activity when making escape decisions. Whether there are generally more people

during a week or year either does not seem especially influential, or alternatively such variation

in human levels (given the hourly and daily variation) is insufficient to trigger any further

response (beyond the immediate one).

Human population and disturbance are predicted to further increase during the 21st century [65].

Therefore, animals will be increasingly forced to occupy human-altered environments, and

altered environments will host more humans. Our results indicate that urban birds may not

flexibly and quickly change their escape behavior to a temporarily changed human presence in

cities. It is unclear whether such tolerance reflects a natural selection on urban tolerance,

differential settlement of individual birds in cities, or habituation-like processes that reduce

plasticity.

We need comprehensive genetic investigations of current populations along urban–

rural/wilderness gradients combined with detailed investigation of individually-marked birds

(*sensu* [52,53]) and other taxa (*sensu* [50,66]) that are repeatedly tested over time. To properly

disentangle the role of differential settlement and habituation, marked birds should be

investigated under experimentally manipulated human densities or when human densities

change such as those that follow natural or human-induced disasters, predicable seasonal

events, or rapid urban development.

Our study further highlights that we still know little about how to measure human outdoor

presence and which of its features are most relevant for behavioral adjustments in free-living

animals. Answering these questions are essential if we are to successfully manage biodiversity

in the Anthropocene.

Materials and Methods

Study areas

Flight initiation distances were collected in Finland (J.J. and field assistant), Poland (P.T.),

Czech Republic (P.M., F.M., Y.B., K.F., A.S. and F.A.Z.), Hungary (G.M. and S.S.), and

Australia (M.W. and field assistants). In the Czech Republic, P.M. sampled 46 sites out of 56.

In Australia, M.W. sampled 32 sites out of 37. Each site was sampled by the same observer

both before and during the shutdowns. All escape distances were collected during the breeding

season (Europe: 1 April – 1 August; Australia: 15 August – 15 March) when most birds are

territorial and active in the areas around their nests, and only in urban areas, i.e. areas with

continuous urban elements, including multi-story buildings, family houses, or roads, with built-

up area >50%, building density >10 buildings/ha, residential human density >10 buildings/ha

[67]. Most data were collected in urban green areas, particularly parks and cemeteries. Finnish

data were collected in Rovaniemi (66.500°N, 25.733°E; 64,000 inhabitants, 75–203 m a. s. l.),

Polish in Poznań (52.406°N, 16.925°E; 0.53 million inhabitants, 60–154 m a. s. l.), Czech data

in Prague (50.083°N, 14.417°E; 1.3 million inhabitants, 177–399 m a. s. l.), Hungarian in

Budapest (47.498°N, 19.041°E; 1.8 million inhabitants, 96–527 m a. s. l.), and Australian in

Melbourne (37.821°S, 144.961°E; 5.2 million inhabitants, 5–169 m a. s. l.). For each city, we

collected data for two breeding seasons before the pandemic, covering the 2018 and 2019

seasons immediately preceding the emergence of the COVID-19 (Finland, Hungary, Poland;

and until March 2020 in Australia) and for up to two breeding seasons during the COVID-19

shutdowns (seasons starting in 2020 and 2021; for the Czech Republic, only starting in 2021).

The 2019 data were unavailable for the Czech Republic, and thus we used data from 2014 and

2018.

Avian tolerance towards humans

Avian tolerance towards human approach was estimated by a simple but widely used method,

quantifying the flight initiation distance which is the distance at which birds escape when

approached by a human observer [26,35–37]. The flight initiation distance reflects a trade-off

between the fitness-related benefits of not escaping and the costs of fleeing [29,35,68,69]. The

flight initiation distance estimates are highly consistent for individuals, populations, and species

tested within similar contexts [26,47,49,51–53].

All data were collected by trained observers skilled in bird identification, and using a standard

procedure outlined previously [32,33,37,45,47,70]. Briefly, when a focal bird (typically an

adult individual) was spotted, a single observer moved at a normal walking speed ($\sim$1 ms^{-1})

directly towards the bird (with head and gaze oriented towards this bird). When the focal bird

first started to escape (i.e. hopped, walked, ran, or flew away), the distance of the observer to

the bird was noted. The escape distance was measured by counting the number of steps of

known approximate length and converting them to meters or using a rangefinder (with $\pm$1 m

resolution). Our previous validation showed that the between-observer variation in escape

distance estimates is low [71] and that distance measured by counting steps did not differ from

the distance measured by a rangefinder [31]. In 79% of the cases (n = 5032 out of 6369

observations), we approached birds located on the ground. The escape distance of birds

positioned above the ground (e.g. perching on vegetation) was estimated as the Euclidean

distance that equals the square-root of the sum of the squared horizontal distance and the

squared height above the ground. Previous studies typically revealed no effect of perch height

on birds' flight initiation distance [72–75]. We approached only individuals that were not on or

next to their nests and we focused on 'relaxed birds', which we defined as those without any

initial signs of distress.

Birds often occur in flocks. In these cases, we randomly selected a single individual from a

flock and measured its response. All fieldworkers wore outdoor clothes without any bright

colors. Within each city we collected data at many sites (at the level of park, cemetery, etc.); to

avoid repeated sampling of the same individuals, we did not re-sample the same location during

the same breeding season. Within a sampling event at a given site, individuals of the same

species were sampled only if it was obvious that they were different individuals (e.g. because

of their simultaneous presence or if morphological features, e.g. sex-specific coloration or age,

enabled us to distinguish between different individuals). The flight initiation distances were

collected during favorable weather conditions (i.e. no rain and no strong wind). In total, we

collected 6369 flight initiation distance estimates for 147 bird species representing 2693 before-

shutdown estimates for 68 species and 3676 during-shutdown estimates for 135 species.

Predictors and covariates

To investigate how hourly changes in the human levels in parks influence escape distance

(hourly number of humans), for a subset of escape distance observation we estimated the

number of humans within a 50 m radius during each escape distance trial (N = 3504; $N_{Finland}$

$_{(2018–2021)}$ = 761, $N_{Poland\ (2018)}$ = 263, $N_{Czechia\ (2018,\ 2021)}$ = 1024, $N_{Hungary\ (2018–2021)}$ = 1456).

To explore how daily and weekly changes in the levels of human presence influence escape

distance, we extracted two variables. First, for each city, we extracted Google Mobility Reports

for parks, covering 2020–2022 (https://www.google.com/covid19/mobility). Google Mobility

reports the change in human presence for each day of the week relative to its baseline value,

estimated as the median value from 3 January – 6 February 2020. Note that Google Mobility

provides the relative weekday change from the baseline and thus lacks the absolute values of

human mobility; this index also lacks freely available data before 2020. Nevertheless, the year

2022 was without governmental restrictions and hence might approximate the normal (before

COVID-19) human levels outdoors, albeit with a caveat as the pandemic has changed human

working habits, e.g. more people currently work from home [76,77]. Second, for each country

and day, we extracted data on the strength of governmental anti-pandemic measures

characterized by Stringency index (https://ourworldindata.org/covid-stringency-index; [42]).

This index is rescaled to values from 0 to 100 (0 = no restrictions; 100 = strictest restrictions)

and represents a composite measure based on nine response indicators, including school

closures, workplace closures, cancellation of public events, restrictions on public gatherings,

closures of public transport, stay-at-home requirements, public information campaigns,

restrictions on internal movements, and international travel controls. Note that Stringency index

was not habitat-specific and did not focus directly on parks and cemeteries – the source of most

of our data.

Finally, to explore how yearly changes in the levels of human presence (induced by COVID-

19 shutdowns) influenced escape distance, each observation was scored as collected before (0)

or during (1) the COVID-19 shutdowns (hereafter "Period").

Life-history, social, contextual, and environmental factors may influence escape responses of

birds [26,35,37,47,78–81] and potentially confound associations between avian escape

responses and changes in human outdoor activity before and during the COVID-19 shutdowns.

Hence, we extracted information on seven parameters. (1) The 'starting distance' was estimated

as the distance to the bird (in meters) when an observer started the escape distance trial. (2) The

'flock size' was calculated as the number of all conspecific individuals moving, feeding, or

perching together that were visually separated from other conspecific or mixed-species

individuals. Note that we avoided approaching mixed-species bird groups. (3) The species-

specific 'body size' was approximated as body mass (in grams) and obtained as the mean of

female and male values from EltonTraits 1.0 database [82]. (4) The ambient 'temperature' was

estimated as the air temperature (°C) at the site during data collection either by the thermometer

directly during the fieldwork (Finland) or retrospectively using online tools (Poland, Czechia,

Hungary, Australia). (5) The 'time' of data collection was rounded to the nearest hour. (6) The

date of data collection was noted as a 'day' since the start of the breeding season (Europe: Day

1 = 1 April; Australia: Day 1 = 15 August). (7) The 'site' represents a unique identifier of each

sampled park, cemetery, city district, etc.

Statistical analyses

Bird tolerance in relation to human levels outdoors

We explored whether changes in human levels from hour-to-hour (i.e. the actual number of

humans during escape distance trials), day-to-day (proxied by Google Mobility), week-to-week

(proxied by Stringency index), or year-to-year (proxied by Period) correlated with avian escape

distance. Using mixed effect models, we first fitted ln-transformed escape distance as a

response and the Number of humans, Google Mobility, Stringency index, or Period as a

predictor of interest, while controlling for starting distance of the observer (ln-transformed),

flock size (ln-transformed), body mass (ln-transformed), temperature (also a proxy for a day

within the breeding season: $r_{\text{Pearson}} = 0.48$; Fig. S2), time of day and year (in case of Period

fitted as a random intercept). To account for circular properties of time, time was transformed

into radians ($2 \times$ time $\times \pi/24$) and fitted as sine and cosine of radians [83]. Multicollinearity

among explanatory variables was checked by the correlation matrix, which suggested that

correlations between variables entered simultaneously in the same model were generally weak

(Fig. S2). To account for the non-independence of data points [84,85], we further fitted random

intercepts of weekday (day of the week), genus, species, species at a given day, country, site,

and species within a site, while fitting the predictor of interest (Number of humans, Google

Mobility, Stringency index or Period) as a random slope within country (i.e. allowing for

different effect in each country) and in case of Period models also within site (i.e. allowing for

different Period effect at each site). Fitting Number of humans, Google Mobility, Stringency

index or Period as random slope at other random intercepts, or simplifying the random effects

structures, produces similar results (Fig. S1, Table S1 and S2). To further investigate the

robustness of the findings, also because the number of sampled species (135 vs 68) and unique

sampling days (141 vs 161) differed between period before and during the COVID-19

shutdowns, apart from using all observation, we fitted the models to more conservative datasets.

In the case of Number of humans, Google Mobility, and Stringency index, we used a dataset

with at least five observations per species ($N_{\text{Number of humans}} = 3446$, $N_{\text{Google Mobility}} = 3545$, N

$_{\text{Stringency index}} = 3573$), and another one with at least 10 observations per species ($N_{\text{Number of humans}}$

$= 3408$, $N_{\text{Google Mobility}} = 3399$, $N_{\text{Stringency index}} = 3446$). In the case of Period, we used a dataset

with at least five observations per species and Period (i.e. at least five observations before and

five during the COVID-19 shutdowns; $n = 5260$), the other with at least ten observations per

species and each Period ($n = 5106$). Such alternative approaches gave nearly identical results

(Fig. S1; Table S2).

Although, the mixed effect models account for possible country-specific effects and are more

conservative than country-specific models, we also fitted country-specific mixed-models. The

country-specific models had a similar structure to the global model, excluding the random

slopes and also random intercept of country and in the case of Poland also a random intercept

of site because specific sites were not noted in Poland. We then verified the results from the

full mixed effect model by using the country-specific estimates, standard deviations and sample

sizes to estimate a meta-analytical mean using the *meta.summaries* function from the *rmeta* R-

package [86].

Moreover, because of ongoing debate about whether to control escape distance for starting

distance [87], we also fitted an alternative set of models without starting distance of the

observer.

Exploring human presence

We further checked, using a limited dataset, whether actual human numbers during escape

distance trials reflected Google Mobility and Stringency index. To do so we fitted mixed effect

models with the number of humans as a response variable (on original scale or on ln-scale after

adding 0.01 to number of humans), and Google Mobility or Stringency index as a predictor

while also controlling for the year and the non-independence of data points by including

weekday within the year as a random intercept, and Google Mobility or Stringency index as a

random slope.

We further investigated the extent of day-to-day variation in Google Mobility, as well as

whether Google Mobility reflected the stringency of governmental restrictions. We fitted a

mixed effect model with Google Mobility as a response variable, and Stringency index as a

predictor while also controlling for the year and the non-independence of data points by

including weekday within the year as a random intercept and Stringency index as a random

slope.

Finally, to estimate whether shutdowns changed human mobility in sampled cities, we visually

compared Google Mobility in parks during shutdowns (2020–2021) and after shutdowns (2022;

Fig. 4).

General statistical procedure

We used R version 4.3.0 for all statistical analyses. All mixed models were fitted with the *lme4*

v. 1.1-29 package [88]. We then used the sim function from the *arm* v. 1.12-2 package and a

non-informative prior distribution [89,90] to create a sample of 5000 simulated values for each

model parameter (i.e. posterior distribution). We report effect sizes and model predictions by

the medians, and the uncertainty of the estimates and predictions by the Bayesian 95% credible

intervals represented by the 2.5 and 97.5 percentiles (i.e. 95% CI) from the posterior distribution

of 5000 simulated or predicted values. We graphically inspected the goodness of fit, and the

distribution of the residuals (see plots of model assumptions in [91]).

All continuous variables were standardized by subtracting the mean and dividing by the

standard deviation.

Since the need for phylogenetic control depends on the phylogenetic signal in the model's

residuals [92], we tested whether the residual variance contained a phylogenetic signal. To do

this we extracted the residuals from the models on the full datasets (Fig. 1; models 1a from

Tables S1a–d), and fitted the residuals as a new response variable in an intercept-only Bayesian

linear regression fitted with STAN [93] using the *brm* function from the *brms* v. 2.17 package

[94,95], with species and their phylogenetic relatedness as random effects. The phylogenetic

relatedness was included as a phylogenetic covariance matrix calculated with the *inverseA*

function in the *MCMCglmm* v. 2.33 package [96] from the maximum credibility tree build using

maxCladeCred function in the *phangorn* v. 2.8.1 package [97] and 100 randomly sampled

species-level phylogenetic trees (Hackett backbone) from the BirdTree online tool

(http://birdtree.org) [98]. Priors were specified using the *get_prior* function from *brms*, which

uses the Student's t distribution for the intercept and the standard deviation [94]. The target

average proposal acceptance probability was increased to 0.99 to make the sampling more

conservative to posterior distributions with high curvature [94]. Five MCMC chains ran for

5000 iterations each while discarding the first 2500 iterations as burn-in, and sampling every

5^{th} iteration, which resulted in 5000 samples of model parameters. The independence of samples

in the Markov chain was assessed using graphical diagnostics, and the convergence was

estimated using the Gelman-Rubin diagnostics which was 1 for all parameters, indicating model

convergence [99]. Phylogeny explained zero or little variance in the residuals from a model on

the number of humans (95% CI: 0–0.6% of variance in random effects), Google Mobility (0–

0.4%), Stringency index (0–0.4%), or Period (0–0.2%) and the model without phylogeny fitted

the data on residuals better than the model with phylogeny (i.e. the estimated Bayes factor in

favor of non-phylogenetic model was 138 for the number of humans, 96 for Google Mobility,

115 for Stringency index, 99 for Period; their posterior probabilities were 0.99 for all; Table

S4), which justifies our use of non-phylogenetic comparative methods.

All results are reproducible with the open-access data and code available from [91], which also

provides visual representations of model assumptions. Visualizations were generated by the

ggplot2 v. 3.3.6 package [100] and other supporting packages (for details, see [91]).

Acknowledgments

We thank Wolfgang Forstmeier and Lotte Schlicht for an advice on statistical analyses. M.W.

thanks Eliza Anderson, Mickey Balzereit, Sammi Cunningham, Alisha Dabonde, Chloe Daws,

Megan Dennis, Zoe Kellett, Clement Masse, Aaron Moore, Bailey Raats, Max Radvan,

Rebecca Frost, Anthony Rendall, Natalie Searle, Mel Sheedy, Will Standish, Mia Stott, and

Millie Toomey for help with the collection of escape responses of birds in Melbourne. Clement

Masse helped us with data collection in Rovaniemi. J.J. was partly supported by the project

"Responsible Tourism Planning" (2019-2023; project number 326348) founded by the

Academy of Finland (PROFI 5 Competitive funding to strengthen universities' research

profiles). P.M. and P.T. gratefully acknowledges the support the TUM – Institute for Advanced

Study – Hans Fisher Senior Fellowship. P.M. is also thankful to Anna Milosavljevičová for

support during the research. M.B. is thankful to Barbora and Majlen for their patience and

support.

Ethics

The fieldwork protocols complied with the current laws of the countries. This kind of research

requires no special permits in Europe. In Australia, Animal ethics approvals (Deakin University

Animal Ethics Committee Permits B10-2018 and B08-2021) and permits (DEWLP, 10008731

and 10010123) were obtained. All fieldwork was conducted following the approved guidelines.

Data were collected in public places and private lands where no special permit was required.

The method used to estimate avian tolerance towards human disturbance was designed to cause

only brief and minimal disturbance to birds; in cities, this disturbance typically does not differ

from standard background disturbance caused by other site visitors.

Competing interests

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

Data availability

Data and computer code used to generate the results, as well as all supporting material including

the plots of model assumptions, are freely available from the Open Science Framework

repository https://martinbulla.github.io/avian_FID_covid/ [91].

References

1.   Fang H, Wang L, Yang Y. Human mobility restrictions and the spread of the Novel
Coronavirus (2019-nCoV) in China. J Public Econ. 2020;191: 104272.
doi:10.1016/j.jpubeco.2020.104272
2.   Gatto M, Bertuzzo E, Mari L, Miccoli S, Carraro L, Casagrandi R, et al. Spread and
dynamics of the COVID-19 epidemic in Italy: Effects of emergency containment
measures. Proc Natl Acad Sci U S A. 2020;117: 10484–10491.
doi:10.1073/pnas.2004978117
3.   Huang X, Li Z, Jiang Y, Li X, Porter D. Twitter reveals human mobility dynamics
during the COVID-19 pandemic. Gao S, editor. PLoS One. 2020;15: e0241957.
doi:10.1371/journal.pone.0241957

4.   Kraemer MUG, Yang CH, Gutierrez B, Wu CH, Klein B, Pigott DM, et al. The effect
581   of human mobility and control measures on the COVID-19 epidemic in China. Science
582   (1979). 2020;368: 493–497. doi:10.1126/science.abb4218
583   5.   Venter ZS, Aunan K, Chowdhury S, Lelieveld J. COVID-19 lockdowns cause global
584   air pollution declines. Proc Natl Acad Sci U S A. 2020;117: 18984–18990.
585   doi:10.1073/pnas.2006853117
586   6.   March D, Metcalfe K, Tintoré J, Godley BJ. Tracking the global reduction of marine
587   traffic during the COVID-19 pandemic. Nature Communications 2021 12:1. 2021;12:
588   1–12. doi:10.1038/s41467-021-22423-6
589   7.   Venter ZS, Barton DN, Gundersen V, Figari H, Nowell M. Urban nature in a time of
590   crisis: recreational use of green space increases during the COVID-19 outbreak in Oslo,
591   Norway. Environmental Research Letters. 2020;15: 104075. doi:10.1088/1748-
592   9326/ABB396
593   8.   Randler C, Tryjanowski P, Jokimäki J, Kaisanlahti-Jokimäki ML, Staller N. SARS-
594   CoV2 (COVID-19) Pandemic Lockdown Influences Nature-Based Recreational
595   Activity: The Case of Birders. International Journal of Environmental Research and
596   Public Health 2020, Vol 17, Page 7310. 2020;17: 7310. doi:10.3390/IJERPH17197310
597   9.   Liu Y, Wang Z, Rader B, Li B, Wu CH, Whittington JD, et al. Associations between
598   changes in population mobility in response to the COVID-19 pandemic and
599   socioeconomic factors at the city level in China and country level worldwide: a
600   retrospective, observational study. Lancet Digit Health. 2021;3: e349–e359.
601   doi:10.1016/S2589-7500(21)00059-5
602   10.   Bates AE, Primack RB, Biggar BS, Bird TJ, Clinton ME, Command RJ, et al. Global
603   COVID-19 lockdown highlights humans as both threats and custodians of the
604   environment. Biol Conserv. 2021;263: 109175. doi:10.1016/J.BIOCON.2021.109175
605   11.   Derryberry EP, Phillips JN, Derryberry GE, Blum MJ, Luther D. Singing in a silent
606   spring: Birds respond to a half-century soundscape reversion during the COVID-19
607   shutdown. Science (1979). 2020;370: 575–579. doi:10.1126/SCIENCE.ABD5777
608   12.   Rutz C, Loretto MC, Bates AE, Davidson SC, Duarte CM, Jetz W, et al. COVID-19
609   lockdown allows researchers to quantify the effects of human activity on wildlife. Nat
610   Ecol Evol. 2020;4: 1156–1159. doi:10.1038/s41559-020-1237-z
611   13.   Soto EH, Botero CM, Milanés CB, Rodríguez-Santiago A, Palacios-Moreno M, Díaz-
612   Ferguson E, et al. How does the beach ecosystem change without tourists during
613   COVID-19 lockdown? Biol Conserv. 2021;255: 108972.
614   doi:10.1016/J.BIOCON.2021.108972
615   14.   Zellmer AJ, Wood EM, Surasinghe T, Putman BJ, Pauly GB, Magle SB, et al. What
616   can we learn from wildlife sightings during the COVID-19 global shutdown?
617   Ecosphere. 2020;11: e03215. doi:10.1002/ecs2.3215
618   15.   Park M, Lim JT, Wang L, Cook AR, Dickens BL. Urban-Rural Disparities for COVID-
619   19: Evidence from 10 Countries and Areas in the Western Pacific. Health Data Science.
620   2021;2021: 1–9. doi:10.34133/2021/9790275
621   16.   Manenti R, Mori E, Di Canio V, Mercurio S, Picone M, Caffi M, et al. The good, the
622   bad and the ugly of COVID-19 lockdown effects on wildlife conservation: Insights
623   from the first European locked down country. Biol Conserv. 2020;249: 108728.
624   doi:10.1016/j.biocon.2020.108728
625   17.   Gordo O, Brotons L, Herrando S, Gargallo G. Rapid behavioural response of urban
626   birds to COVID-19 lockdown. Proceedings of the Royal Society B: Biological
627   Sciences. 2021;288: 20202513. doi:10.1098/rspb.2020.2513
628   18.   Schrimpf MB, des Brisay PG, Johnston A, Smith AC, Sánchez-Jasso J, Robinson BG,
629   et al. Reduced human activity during COVID-19 alters avian land use across North

America. Sci Adv. 2021;7: 5073–5095.
doi:10.1126/SCIADV.ABF5073/SUPPL_FILE/SCIADV.ABF5073_DATA_S1_TO_S
4

19. Vardi R, Berger-Tal O, Roll U. iNaturalist insights illuminate COVID-19 effects on
large mammals in urban centers. Biol Conserv. 2021;254: 108953.
doi:10.1016/J.BIOCON.2021.108953

20. Warrington MH, Schrimpf MB, Des Brisay P, Taylor ME, Koper N. Avian behaviour
changes in response to human activity during the COVID-19 lockdown in the United
Kingdom. Proceedings of the Royal Society B: Biological Sciences. 2022;289.
doi:10.1098/RSPB.2021.2740/

21. Wilmers CC, Nisi AC, Ranc N. COVID-19 suppression of human mobility releases
mountain lions from a landscape of fear. Current Biology. 2021;31: 3952-3955.e3.
doi:10.1016/J.CUB.2021.06.050

22. Diamant ES, MacGregor-Fors I, Blumstein DT, Yeh PJ. Urban birds become less
fearful following COVID-19 reopenings. Proceedings of the Royal Society B . 2023
[cited 7 May 2023]. doi:10.1101/2023.01.04.522762

23. Montgomery RA, Raupp J, Parkhurst M. Animal Behavioral Responses to the COVID-
19 Quietus. Trends Ecol Evol. 2021;36: 184–186. doi:10.1016/J.TREE.2020.12.008

24. Samia DSM, Nakagawa S, Nomura F, Rangel TF, Blumstein DT. Increased tolerance
to humans among disturbed wildlife. Nat Commun. 2015;6: 8877.
doi:10.1038/ncomms9877

25. Ducatez S, Sol D, Sayol F, Lefebvre L. Behavioural plasticity is associated with
reduced extinction risk in birds. Nat Ecol Evol. 2020; 1–6. doi:10.1038/s41559-020-
1168-8

26. Díaz M, Møller AP, Flensted-Jensen E, Grim T, Ibáñez-Álamo JD, Jokimäki J, et al.
The geography of fear: a latitudinal gradient in anti-predator escape distances of birds
across Europe. Mettke-Hofmann C, editor. PLoS One. 2013;8: e64634.
doi:10.1371/journal.pone.0064634

27. Samia DSM, Blumstein DT, Díaz M, Grim T, Ibáñez-Álamo JD, Jokimäki J, et al.
Rural-urban differences in escape behavior of European birds across a latitudinal
gradient. Front Ecol Evol. 2017;5: 66.

28. Mikula P, Tomášek O, Romportl D, Aikins TK, Avendaño JE, Braimoh-Azaki BDA, et
al. Bird tolerance to humans in open tropical ecosystems. Nat Commun. 2023;14: 2146.
doi:10.1038/s41467-023-37936-5

29. Samia DSM, Blumstein DT, Stankowich T, Cooper WE. Fifty years of chasing lizards:
new insights advance optimal escape theory. Biological Reviews. 2016;91: 349–366.
doi:10.1111/brv.12173

30. Webb N V, Blumstein DT. Variation in Human Disturbance Differentially Affects
Predation Risk Assessment in Western Gulls. Condor. 2005;107: 178–181.
doi:10.1093/condor/107.1.178

31. Mikula P. Pedestrian density influences flight distances of urban birds. Ardea.
2014;102: 53–60. doi:10.5253/078.102.0105

32. Mikula P, Jokimäki J, Kaisanlahti-Jokimäki ML, Markó G, Morelli F, Møller AP, et al.
Face mask-wear did not affect large-scale patterns in escape and alertness of urban and
rural birds during the COVID-19 pandemic. Science of The Total Environment.
2021;793: 148672. Available: 10.1016/j.scitotenv.2021.148672

33. Morelli F, Leveau LM, Mikula P, MacGregor-Fors I, Bocelli ML, Quesada-Acuña SG,
et al. Are birds more afraid in urban parks or cemeteries? A Latin American study
contrasts with results from Europe. Science of The Total Environment. 2023;861:
160534. doi:10.1016/J.SCITOTENV.2022.160534

34.   Díaz M, Møller AP. Lockdown effects on fear revealed direct and indirect effects of
human presence on perceived predation risk. Science of The Total Environment.
2023;872: 162122. doi:10.1016/J.SCITOTENV.2023.162122
35.   Stankowich T, Blumstein DT. Fear in animals: a meta-analysis and review of risk
assessment. Proceedings of the Royal Society of London B: Biological Sciences.
2005;272: 2627–2634. doi:10.1098/rspb.2005.3251
36.   Weston MA, McLeod EM, Blumstein DT, Guay P-J. A review of flight-initiation
distances and their application to managing disturbance to Australian birds. Emu.
2012;112: 269–286. doi:10.1071/MU12026
37.   Blumstein DT. Developing an evolutionary ecology of fear: how life history and
natural history traits affect disturbance tolerance in birds. Anim Behav. 2006;71: 389–
399. doi:10.1016/j.anbehav.2005.05.010
38.   Burger J. Effects of Motorboats and Personal Watercraft on Flight Behavior over a
Colony of Common Terns. Condor. 1998;100: 528–534. doi:10.2307/1369719
39.   Csomós G, Borza EM, Farkas JZ. Exploring park visitation trends during the Covid-19
pandemic in Hungary by using mobile device location data. Scientific Reports 2023
13:1. 2023;13: 1–12. doi:10.1038/s41598-023-38287-3
40.   Zhao H, Mailloux BJ, Cook EM, Culligan PJ. Change of urban park usage as a
response to the COVID-19 global pandemic. Scientific Reports 2023 13:1. 2023;13: 1–
11. doi:10.1038/s41598-023-46745-1
41.   Volenec ZM, Abraham JO, Becker AD, Dobson AP. Public parks and the pandemic:
How park usage has been affected by COVID-19 policies. PLoS One. 2021;16:
e0251799. doi:10.1371/JOURNAL.PONE.0251799
42.   Hale T, Angrist N, Goldszmidt R, Kira B, Petherick A, Phillips T, et al. A global panel
database of pandemic policies (Oxford COVID-19 Government Response Tracker).
Nature Human Behaviour 2021 5:4. 2021;5: 529–538. doi:10.1038/s41562-021-01079-
8
43.   Geng D (Christina), Innes J, Wu W, Wang G. Impacts of COVID-19 pandemic on
urban park visitation: a global analysis. J For Res (Harbin). 2021;32: 553–567.
doi:10.1007/S11676-020-01249-W/TABLES/5
44.   Gelman A, Su Y-S, Yajima M, Hill J, Pittau MG, Kerman J, et al. Data Analysis Using
Regression and Multilevel/Hierarchical Models [R package arm version 1.13-1]. 2022
[cited 7 May 2023]. Available: https://CRAN.R-project.org/package=arm
45.   Morelli F, Mikula P, Benedetti Y, Bussière R, Jerzak L, Tryjanowski P. Escape
behaviour of birds in urban parks and cemeteries across Europe: Evidence of
behavioural adaptation to human activity. Science of the Total Environment.
2018;631–632: 803–810. doi:10.1016/j.scitotenv.2018.03.118
46.   Rodriguez-Prieto I, Fernández-Juricic E, Martín J, Regis Y. Antipredator behavior in
blackbirds: habituation complements risk allocation. Behavioral Ecology. 2009;20:
371–377. doi:10.1093/beheco/arn151
47.   Mikula P, Díaz M, Albrecht T, Jokimäki J, Kaisanlahti-Jokimäki M-L, Kroitero G, et
al. Adjusting risk-taking to the annual cycle of long-distance migratory birds. Sci Rep.
2018;8: 13989. doi:10.1038/s41598-018-32252-1
48.   Cavalli M, Baladrón A V., Isacch JP, Biondi LM, Bó MS. The role of habituation in
the adjustment to urban life: An experimental approach with burrowing owls.
Behavioural Processes. 2018;157: 250–255. doi:10.1016/J.BEPROC.2018.10.011
49.   Symonds MRE, Weston MA, van Dongen WFD, Lill A, Robinson RW, Guay P-J.
Time since urbanization but not encephalisation is associated with increased tolerance
of human proximity in birds. Front Ecol Evol. 2016;4: 117.
doi:10.3389/fevo.2016.00117

50.   Uchida K, Blumstein DT. Habituation or sensitization? Long-term responses of yellow-
731   bellied marmots to human disturbance. Behavioral Ecology. 2021;32: 668–678.
732   doi:10.1093/BEHECO/ARAB016

733   51.   Carrete M, Martínez-Padilla J, Rodríguez-Martínez S, Rebolo-Ifrán N, Palma A, Tella
734   JL. Heritability of fear of humans in urban and rural populations of a bird species. Sci
735   Rep. 2016;6: 31060. doi:10.1038/srep31060

736   52.   Carrete M, Tella JL. High individual consistency in fear of humans throughout the
737   adult lifespan of rural and urban burrowing owls. Sci Rep. 2013;3: 1–7.
738   doi:10.1038/srep03524

739   53.   Carrete M, Tella JL. Individual consistency in flight initiation distances in burrowing
740   owls: a new hypothesis on disturbance-induced habitat selection. Biol Lett. 2010;23:
741   167–170.

742   54.   Slater C, Cam G, Qi Y, Liu Y, Guay PJ, Weston MA. Camera shy? Motivations,
743   attitudes and beliefs of bird photographers and species-specific avian responses to their
744   activities. Biol Conserv. 2019;237: 327–337. doi:10.1016/J.BIOCON.2019.07.016

745   55.   Lethlean H, Van Dongen WFD, Kostoglou K, Guay PJ, Weston MA. Joggers cause
746   greater avian disturbance than walkers. Landsc Urban Plan. 2017;159: 42–47.
747   doi:10.1016/J.LANDURBPLAN.2016.08.020

748   56.   Radkovic AZ, Van Dongen WFD, Kirao L, Guay P-J, Weston MA. Birdwatchers
749   evoke longer escape distances than pedestrians in some African birds. Journal of
750   Ecotourism. 2019;18: 100–106. doi:10.1080/14724049.2017.1372765

751   57.   Sprau P, Dingemanse NJ. An approach to distinguish between plasticity and non-
752   random distributions of behavioral types along urban gradients in a wild passerine bird.
753   Front Ecol Evol. 2017;5: 92. doi:10.3389/FEVO.2017.00092/BIBTEX

754   58.   Minias P, Włodarczyk R, Minias A, Dziadek J. How birds colonize cities: genetic
755   evidence from a common waterbird, the Eurasian coot. J Avian Biol. 2017;48: 1095–
756   1103. doi:10.1111/JAV.01334

757   59.   Indykiewicz P, Podlaszczuk P, Janiszewska A, Minias P. Extensive gene flow along
758   the urban–rural gradient in a migratory colonial bird. J Avian Biol. 2018;49.
759   doi:10.1111/JAV.01723

760   60.   Ellenberg U, Mattern T, Seddon PJ. Habituation potential of yellow-eyed penguins
761   depends on sex, character and previous experience with humans. Anim Behav.
762   2009;77: 289–296. doi:10.1016/J.ANBEHAV.2008.09.021

763   61.   Rankin CH, Abrams T, Barry RJ, Bhatnagar S, Clayton DF, Colombo J, et al.
764   Habituation revisited: An updated and revised description of the behavioral
765   characteristics of habituation. Neurobiol Learn Mem. 2009;92: 135–138.
766   doi:10.1016/J.NLM.2008.09.012

767   62.   Zaccaroni M, Ciuffreda M, Paganin M, Beani L. Does an early aversive experience to
768   humans modify antipredator behaviour in adult Rock partridges?
769   http://dx.doi.org/101080/0892701420079522561. 2010;19: 193–200.
770   doi:10.1080/08927014.2007.9522561

771   63.   Vincze E, Papp S, Preiszner B, Seress G, Bókony V, Liker A. Habituation to human
772   disturbance is faster in urban than rural house sparrows. Behavioral Ecology. 2016;27:
773   1304–1313. doi:10.1093/beheco/arw047

774   64.   Čapkun-Huot C, Blumstein DT, Garant D, Sol D, Réale D. Toward a unified
775   framework for studying behavioural tolerance. Trends Ecol Evol. 2024 [cited 16 Jan
776   2024]. doi:10.1016/J.TREE.2023.12.006

777   65.   Sanderson EW, Jaiteh M, Levy MA, Redford KH, Wannebo A V., Woolmer G. The
778   Human Footprint and the Last of the Wild: The human footprint is a global map of
779   human influence on the land surface, which suggests that human beings are stewards of

nature, whether we like it or not. Bioscience. 2002;52: 891–904. doi:10.1641/0006-
3568(2002)052[0891:thfatl]2.0.co;2

66. Blumstein DT, Sanchez M, Philson CS, Bliard L. Is flight initiation distance associated
with longer-term survival in yellow-bellied marmots, Marmota flaviventer? Anim
Behav. 2023;202: 21–28. doi:10.1016/J.ANBEHAV.2023.05.013

67. Marzluff JM, Bowman R, Donnelly R. A historical perspective on urban bird research:
trend, terms, and approaches. In: Marzluff JM, Bowman R, Donnelly R, editors. Avian
Ecology and Conservation in an Urbanizing World. New York: Kluwer; 2001. pp. 20–
47.

68. Ydenberg RC, Dill LM. The economics of fleeing from predators. Adv Study Behav.
1986;16: 229–249. doi:10.1016/S0065-3454(08)60192-8

69. Albrecht T, Klvaňa P. Nest Crypsis, Reproductive Value of a Clutch and Escape
Decisions in Incubating Female Mallards Anas platyrhynchos. Ethology. 2004;110:
603–613. doi:10.1111/J.1439-0310.2004.00992.X

70. Blumstein DT, Samia DSM, Stankowich T, Cooper WE. Best practice for the study of
escape behavior. In: Cooper WE, Blumstein DT, editors. Escaping from Predators: An
Integrative View of Escape Decisions. Cambridge: Cambridge University Press; 2015.
pp. 407–419. doi:10.1017/CBO9781107447189.017

71. Guay P-J, McLeod EM, Cross R, Formby AJ, Maldonado SP, Stafford-Bell RE, et al.
Observer effects occur when estimating alert but not flight-initiation distances. Wildlife
Research. 2013;40: 289. doi:10.1071/WR13013

72. Bjørvik LM, Dale S, Hermansen GH, Munishi PKT, Moe SR. Bird flight initiation
distances in relation to distance from human settlements in a Tanzanian floodplain
habitat. J Ornithol. 2015;156: 239–246. doi:10.1007/s10336-014-1121-1

73. Kalb N, Anger F, Randler C. Flight initiation distance and escape behavior in the black
redstart (Phoenicurus ochruros). Ethology. 2019;125: 430–438.
doi:10.1111/ETH.12867

74. Sreekar R, Goodale E, Harrison RD. Flight initiation distance as behavioral indicator of
hunting pressure: a case study of the sooty-headed bulbul (Pycnonotus aurigaster) in
Xishuangbanna, SW China. Trop Conserv Sci. 2020;8: 505–512.
doi:10.1177/194008291500800214

75. Braimoh B, Iwajomo S, Wilson M, Chaskda A, Ajang A, Cresswell W. Managing
human disturbance: factors influencing flight-initiation distance of birds in a West
African nature reserve. https://doi.org/102989/0030652520171388300. 2018;89: 59–
69. doi:10.2989/00306525.2017.1388300

76. de Palma A, Vosough S, Liao F. An overview of effects of COVID-19 on mobility and
lifestyle: 18 months since the outbreak. Transp Res Part A Policy Pract. 2022;159:
372–397. doi:10.1016/J.TRA.2022.03.024

77. Kong X, Zhang A, Xiao X, Das S, Zhang Y. Work from home in the post-COVID
world. Case Stud Transp Policy. 2022;10: 1118. doi:10.1016/J.CSTP.2022.04.002

78. Morelli F, Benedetti Y, Díaz M, Grim T, Ibáñez-Álamo JD, Jokimäki J, et al.
Contagious fear: Escape behavior increases with flock size in European gregarious
birds. Ecol Evol. 2019;9: 6096–6104. doi:10.1002/ece3.5193

79. Reynolds C, Henry DAW, Tye DRC, Tye ND. Defining separation zones for coastal
birds at a wetland of global importance. Wildlife Research. 2020;48: 134–141.
doi:10.1071/wr20098

80. Piratelli AJ, Favoretto GR, de Almeida Maximiano MF. Factors affecting escape
distance in birds. Zoologia (Curitiba). 2015;32: 438–444. doi:10.1590/S1984-
46702015000600002

81.   Legagneux P, Ducatez S. European birds adjust their flight initiation distance to road
speed limits. Biol Lett. 2013;9. doi:10.1098/RSBL.2013.0417
82.   Wilman H, Belmaker J, Simpson J, de la Rosa C, Rivadeneira MM, Jetz W. EltonTraits
1.0: Species-level foraging attributes of the world's birds and mammals. Ecology.
2014;95: 2027–2027. doi:10.1890/13-1917.1
83.   Bulla M, Valcu M, Dokter AM, Dondua AA, Kosztolányi A, Rutten A, et al.
Unexpected diversity in socially synchronized rhythms of shorebirds. Nature.
2016;540: 109–113. doi:doi:10.1038/nature20563
84.   Schielzeth H, Forstmeier W. Conclusions beyond support: overconfident estimates in
mixed models. Behavioral Ecology. 2009;20: 416–420.
doi:10.1093/BEHECO/ARN145
85.   Barr DJ, Levy R, Scheepers C, Tily HJ. Random effects structure for confirmatory
hypothesis testing: Keep it maximal. J Mem Lang. 2013;68: 255–278.
doi:10.1016/J.JML.2012.11.001
86.   Lumley T. rmeta: Meta-Analysis. 2018 [cited 7 May 2023]. Available: https://cran.r-
project.org/web/packages/rmeta/index.html
87.   Cooper WE, Samia DSM, Blumstein DT. FEAR, spontaneity, and artifact in economic
escape theory: A review and prospectus. Adv Study Behav. 2015;47: 147–179.
doi:10.1016/BS.ASB.2015.02.002
88.   Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models using
lme4. J Stat Softw. 2015;67: 1–48. Available: http://arxiv.org/abs/1406.5823
89.   Gelman A, Hill J. Data analysis using regression and multilevel/hierarchical models.
Cambridge: Cambridge University Press; 2007.
90.   Gelman A, Su Y-S. arm: Data analysis using regression and multilevel/hierarchical
models. 2018 [cited 18 May 2022]. Available: http://CRAN.R-
project.org/package=arm
91.   Bulla M, Mikula P, Blumstein DT, Benedetti Y, Floigl K, Jokimäki J, et al. Supporting
information for "Urban birds' flight responses were largely unaffected by the COVID-
19 shutdowns', Github. 2023 [cited 7 May 2023].
doi:https://martinbulla.github.io/avian_FID_covid/
92.   Uyeda JC, Zenil-Ferguson R, Pennell MW. Rethinking phylogenetic comparative
methods. Syst Biol. 2018;67: 1091–1109. doi:10.1093/SYSBIO/SYY031
93.   Stan Development Team. Stan Modeling Language Users Guide and Reference
Manual, l, Version 2.28. 2021.
94.   Bürkner PC. brms: An R package for Bayesian multilevel models using Stan. J Stat
Softw. 2017;80: 1–28. doi:10.18637/jss.v080.i01
95.   Bürkner PC. Advanced Bayesian multilevel modeling with the R package brms. R
Journal. 2018;10: 395–411. doi:10.32614/RJ-2018-017
96.   Hadfield JD. MCMC methods for multi-response generalised linear mixed models: the
MCMCglmm R package. J Stat Softw. 2010;33: 1–22. doi:10.18637/jss.v033.i02
97.   Schliep KP. phangorn: phylogenetic analysis in R. Bioinformatics. 2011;27: 592–593.
doi:10.1093/bioinformatics/btq706
98.   Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO. The global diversity of birds in
space and time. Nature. 2012;491: 444–448. doi:10.1038/nature11631
99.   Brooks SP, Gelman A. General methods for monitoring convergence of iterative
simulations)? Journal of Computational and Graphical Statistics. 1998;7: 434–455.
doi:10.1080/10618600.1998.10474787
100. Wickham H. Ggplot2: Elegant graphics for data analysis. 2nd ed. Use R! Cham,
Switzerland: Springer International Publishing; 2016.
878