Diverse strategies for tracking seasonal environmental niches at hemispheric scale

Jeremy Cohen*[1,2] & Walter Jetz[1,2]

[1]Department of Ecology and Evolutionary Biology, Yale University, 165 Prospect Street, New

Haven, CT, 06520 USA

[2]Center for Biodiversity and Global Change, Yale University, 165 Prospect Street, New Haven,

CT, 06520 USA

*Correspondence to: jeremy3cohen@gmail.com

Author emails (in order of authorship): jeremy3cohen@gmail.com, walter.jetz@yale.edu

Author statement: All authors agreed to submission of the manuscript and accept the

responsibility for the accuracy and integrity of the manuscript.

Author contributions: All authors devised ideas for the study. JMC compiled annotated GBIF

data. JMC calculated niche similarity metrics. JMC accessed functional trait databases and

harmonized phylogenetic information. JMC conducted statistical analyses and generated figures.

All authors helped write the manuscript.

Keywords: Environmental niche, climatic niche, hypervolume, seasonal niche tracking, birds,

big data, seasonality, migration, functional trait groups, phylogenetic signal, GBIF

Abstract:

Species depend upon a constrained set of environmental conditions, or niches, for survival and

reproduction that are increasingly lost under climatic change. Seasonal environments require

species to either track their niches via movement or undergo physiological or behavioral changes

to survive. Here we identify the tracking of both environmental niche position and breadth across

619 New World bird species and assess their phylogenetic and functional underpinning.

Partitioning niche position and breadth tracking can inform whether climatic means or extremes

limit seasonal distributions. We uncover diverse strategies, including the tracking of niche

position, breadth, both, or neither, suggesting highly variable sensitivity to ongoing climatic

change. There was limited phylogenetic determinism to this variation, but a strong association

with functional attributes that differed between niche position and breadth tracking. Our findings

imply significant functional consequences for communities and ecosystems as impending climate

change affects some niche tracking strategies more than others.

Introduction

Species survive and reproduce under a specific set of environmental conditions, known as the environmental niche or n-dimensional hypervolume (Blonder et al. 2014; Hutchinson 1957; Lu, Winner, and Jetz 2021). In seasonal environments, species must adjust to a constantly shifting window of available conditions through one of several strategies. When remaining stationary, species must either maintain tolerance to a wide range of conditions or undergo physiological and behavioral changes to survive seasonal variation, known as 'niche switching' (Nakazawa et al. 2004). Alternatively, seasonally mobile animals can occupy a dynamic niche that remains relatively narrow across the annual cycle, known as 'niche tracking' (Gómez et al. 2016; Somveille, Rodrigues, and Manica 2018; Winger et al. 2019). Seasonal niche tracking is central to the persistence of species with limited behavioral or physiological capacity to adjust their niches (Fandos et al. 2020; Zurell et al. 2018). Understanding the functional and phylogenetic drivers of niche tracking behavior across diverse species can allow researchers to predict how species mediate exposure to novel, potentially adverse conditions as climate change progresses (Tingley et al. 2009; La Sorte and Jetz 2012; Somveille, Rodrigues, and Manica 2015).

After the documentation of seasonal niche tracking behavior in single species (Fandos et al. 2020) and smaller clades (Gómez et al. 2016; Eyres et al. 2020), a more general understanding of the patterns, causes and consequences of seasonal niche tracking across diverse taxonomic groups remains missing. For example, the role of phylogeny and functional traits in driving seasonal niche tracking across a diverse species set remains largely unexplored (but see Zurell et al. 2018). Several studies have hypothesized that niche tracking may be phylogenetically conserved (Gómez et al. 2016; Martínez–Meyer, Townsend Peterson, and Navarro–Sigüenza 2004), as is typical of behavioral and migratory traits (Outlaw and Voelker 2006), but this has not been evaluated rigorously or broadly. Alternatively, niche tracking may have repeatedly evolved in tandem with species' functional traits. For example, obligate insectivores or small-bodied species may be most likely to closely track their niche over the annual cycle to satisfy narrow dietary or thermal requirements (Gómez et al. 2016; Huey et al. 2012). Separating these potential drivers will allow researchers to better predict niche flexibility and niche tracking, and thus climate change vulnerability, among rare species or those from under-sampled regions.

Previous work on niche tracking has focused on seasonal similarity in niche positions, or mean environmental conditions, without accounting for niche components such as niche breadth (the volume or range of tolerable conditions). However, niche breadth is a central additional dimension because climate change is altering both the means and variances of climatic conditions (Rahmstorf and Coumou 2011). For example, species may seasonally track niche breadth instead of changing the central niche position when extremes are more limiting to their survival and reproductive success than climatic means, as is the case for numerous species (Albright et al. 2010; Ummenhofer and Meehl 2017). Thus, species may use the tracking of niche breadth as an alternative strategy to niche position tracking to persist in the face of

seasonality, one unexplored by the existing niche tracking literature, which instead considers
niche breadth as an annually static variable (Gómez et al. 2016; Zurell et al. 2018). To better
identify complex variation in niche tracking, seasonal niche similarity should be partitioned into
constituent components, including *niche position similarity*, or the distance between niche
centroids, representing the difference between the average conditions a species experiences in
each season; and *niche breadth similarity*, or the proportional difference in niche breadths (Lu,
Winner, and Jetz 2021), representing the range of conditions a species can tolerate.
Hypervolumes quantified using non-parametric techniques, such as kernel-density estimates or
support vector machines (Blonder 2018; Brown, Holland, and Jordan 2020), are difficult to
partition into constituent components. However, recently developed parametric methods for
quantifying the niche allow for partitioning of these components and direct hypothesis testing
against predictions derived from theories (Lu, Winner, and Jetz 2021).

We use these new metrics to distinguish five primary strategies for seasonal niche tracking (Fig.
1; Fig. 2): i) "Non-trackers" retain neither the position nor the breadth of their niches over the
annual cycle; ii) "Breadth trackers", or "position shifters", track niche breadth but shift their
position, suggesting that their seasonal ranges may be limited by environmental variation; iii)
"Position trackers", or "breadth expanders", track only niche position and adjust niche breadth
between seasons and may be seasonally limited by environmental means; Finally, "complete
trackers" track both niche position and breadth, but may do so by either iv) migrating to track
weather conditions across the annual cycle or v) remaining stationary in an aseasonal
environment.

However, a simple comparison of niche space during the breeding and overwintering seasons
(henceforth, *seasonal* niche similarity) cannot fully distinguish these strategies (Fig.1b, columns
4-5), despite obviously divergent dispersal capability and potential for climate change adaptation
(Eyres et al. 2020). We therefore consider an additional metric, *locational* niche similarity, in
which the realized overwintering hypervolume is compared with that had the species remained at
its breeding range, its *stationary winter niche* (Fig. 1c). This measure, which emphasizes the
similarity of conditions at specific locations, is important for understanding seasonal niche
tracking because it accounts for the environmental distance covered by the species through
purposeful movement. Among temperate breeders, seasonal and locational niche similarity are
likely to have an inverse relationship; for example, a niche tracking species with little difference
between its breeding and overwintering niche is likely to experience highly distinct
overwintering conditions compared with those at its breeding sites.

114
Figure 1: A typology of seasonal niche tracking strategies. Seasonal niche similarity is decomposed
into two components, similarity in niche position and niche breadth. A given species may (1) be a non-
tracker, tracking neither niche position or breadth, (2) track niche breadth, but shift position, (3) track
niche position, but expand breadth, or (4-5), be a complete tracker, tracking both niche position and
breadth. a) shows the differing use of climate zones over space and time; b) compares seasonal niches
between the breeding (orange) and overwintering (purple) seasons (solid arrows represent positional
similarity, and brackets represent breadth similarity); c) compares the location of the winter conditions at
breeding locations (light purple) and the overwintering niche, revealing that for strategy 4 and 5 niche
tracking is achievable through multiple life history strategies. Dashed arrows represent locational
positional similarity.

Tracks niche breadth, but shifts position

Tracks both niche breadth and position, either by movement or stationarity in less seasonal environments

Tracks neither niche breadth nor position

Tracks niche position, but expands breadth

Niche breadth tracking

Niche position tracking

Figure 2. Five niche tracking strategies. Conceptual schematic outlining the contribution of niche breadth and position tracking to each of five possible niche tracking strategies.

Here, we use a hemispheric system to address patterns and drivers of these strategies; specifically, 619 bird species, representing nearly the full diversity of birds breeding in the US and Canada. This species set is ideally suited given its tremendous variation in migratory strategy, diversity of functional trait groups (Barnagaud et al. 2017) and uniquely comprehensive occurrence data. We also leverage a recently developed environmental annotation tool (Li et al. 2021) and a novel parametric hypervolume method (Lu, Winner, and Jetz 2021) to quantify similarity between the environmental niches occupied by each species during both the breeding and overwintering seasons. We use this system to address the following questions:

1) What is the prevalence of different niche tracking strategies while accounting for both niche position and breadth? To date, a quantitative multi-species assessment of strategies across a diverse taxonomic group has been missing.

2) What is the relationship between the tracking of niche position and the tracking of niche breadth? Do species that maximize niche positional similarity across seasons also tend to retain similar niche breadth, or do species largely track only one or the other?

3) How phylogenetically and functionally determined are niche tracking strategies? Across
functional trait groups, we hypothesize seasonal niche tracking to be most common in a) long-
distance migrants, because they can physically relocate to suitable locations (Laube, Graham,
and Böhning-Gaese 2015; Zurell et al. 2018); b) small-bodied species, because they have low
thermal inertia and generally narrow thermal breadths (Huey et al. 2012; Albright et al. 2017),
suggesting they cannot tolerate large seasonal variation in the niche; c) insectivores, because
abundant insect prey is only available under specific temperature, precipitation and productivity
levels (Winkler, Luo, and Rakhimberdiev 2013); and d) species occupying open or water
habitats, because they are not shielded from climate variability by forest structure (Jarzyna et al.
2016). Given that behavioral and migratory traits are often phylogenetically conserved (Outlaw
and Voelker 2006), we expect a strong phylogenetic signal in seasonal niche tracking behavior
(Gómez et al. 2016).

4) Do cross-species comparisons of *locational* niche similarity reveal important behavioral
strategies and functional or phylogenetic associations not apparent when quantifying only
seasonal niche similarity? We predict an inverse relationship between seasonal and locational
similarity across species and expect to observe functional trait relationships with locational
similarity that are not observed with temporal similarity.

We expect the emerging insights to not only address these questions but more generally offer an
assessment of seasonal niche dynamics as a system to understand realized strategies for
mitigating exposure to climatic change.
164
165 **Material and Methods**
166
Species selection and environmental data

We identified the 672 bird species that annually breed or overwinter in the United States and
Canada based on American Birding Association birding codes (Association 2008) updated to the
Clements bird taxonomy as of 2021 (Clements 2007). These codes are a widely-accepted
authority to distinguish regularly occurring species from irregularly occurring vagrants, or those
species which sporadically appear on the continent each year but whose occurrence is not
predictably tied to a given location. We excluded species not native to the US or Canada or those
that are primarily marine, for which weather data is unavailable.

In August 2021, we accessed the Spatiotemporal Observation Annotation Tool (STOAT) v1.0, a
novel cloud-based toolbox for flexible biodiversity annotations (Li et al. 2021), to download
annotated Global Biodiversity Information Facility data (https://www.gbif.org/) for all species
(data compilation, analyses and visualizations were all completed in R 4.1.0; R Core Team
2021). To minimize the potential for spatiotemporal sampling bias, we then thinned points by
selecting one point from each location (5x5km grid cell) per week.

We annotated observation points with three environmental dimensions: daily maximum
temperature (sourced from NASA-MODIS; https://lpdaac.usgs.gov/products/mod11a1v006/),
enhanced vegetation index (EVI; from MODIS), and precipitation (from CHELSA v2.1; Karger
et al. 2021), each summarized to a 1km buffer (0.5km radius) over 30 days prior to the
observation (imported using jsonlite and httr packages; Ooms 2014; Wickham and Wickham
2020). We did not assess *a priori* whether these variables are equally relevant across species; for
example, certain species may be limited by precipitation but not temperature within their range,
or *vice versa*. Alternatively, some species may be limited by environmental factors not
considered in our estimation of species niches. However, this variable set represents the
environmental factors that most commonly drive species distributions (Qian 2010), and these
variables have low collinearity, allowing each axis to remain independent. Estimating niches
with the same variable set for all species was necessary to ensure consistency in cross-species
evaluation of seasonal changes to both niche position and breadth.

For each species, we restricted observation points to those with environmental data available for
all dimensions. Further, we temporally cropped data to season (December-February for the
overwintering season and June-August for breeding season) and spatially cropped points to the
American continents ($< -30°$ longitude). Sufficient data for analysis (>20 points per season after
filtering) was available for 619 species. Database management was completed using *tidyverse*
packages (Wickham et al. 2019).

To quantify locational similarity, we conducted the same niche characterizations for breeding
locations during the overwintering season. For each species, we created a hypothetical set of
occurrence points corresponding to coordinate locations during the breeding season, each with
randomized winter calendar dates (December-February), annotating them with environmental
data as described above.

Ecological niche modeling and seasonal similarity

For every species, we calculated parametric measurements of the similarity between the seasonal
three-dimensional hypervolumes. The hypervolumes were characterized as multivariate normal
distributions, allowing us to derive analytical estimates for the breadth and position of each
hypervolume (MVNH package; Lu, Winner, and Jetz 2021). We partitioned niche similarity into
two metrics: i) *niche position similarity*, which quantifies the distance among hypervolume
centroids in each season based on the sign-flipped, log-transformed Mahalanobis distance, used
to assess changes in niche position between seasons; and ii) *niche breadth similarity*, which
represents the similarity in *niche breadth* (volume) between seasons and is measured as the sign-
flipped, log-transformed determinant ratio.

To quantify seasonal similarity, we calculated positional and breadth similarity between the
breeding and overwintering seasonal hypervolumes. To quantify locational similarity, we
calculated positional similarity between the overwintering hypervolume and available

hypervolume during winter at breeding locations. Thus, we had a total of three metrics of
seasonal niche similarity per species. We visualized pairwise two-dimensional hypervolumes for
each species using *ggplot2* (Wickham 2011).

Functional traits and phylogeny

We obtained species-level functional trait values and an avian phylogenetic tree to test
associations between trait groups, phylogeny, and seasonal niche tracking. We derived migration
distances from (La Sorte et al. 2022; La Sorte personal communication), body mass from the
Eltonian trait database (Wilman et al. 2014), and habitat preference and diet from Barnagaud et
al. (2017). We grouped several categories of each categorical predictor to avoid false positives
associated with small sample sizes and to keep our conclusions broad. Diet categories included
carnivore, invertebrate, omnivore, and herbivore (combining 'fruit', 'nectar', 'vegetation', and
'seed' categories). Habitat categories included water ('coastal', 'open_water',
'riparian_wetlands'), open ('semi-open', 'rock', 'arid'), generalist ('urban', 'developed'), and
forest.

We updated an avian phylogeny from Jetz et al. (2012) to account for recent taxonomic changes,
updating species names to the Clements bird taxonomy as of 2021 (Clements 2007) and treating
recently split species as having no phylogenetic distance (Appendix 2). We also harmonized
species names in all trait datasets to Clements.

Cross-species models

To assess the role of functional traits in seasonal niche tracking, we used weighted multivariate
phylogenetic generalized least squares (PGLS) models (caper package; Orme et al. 2013).
Phylogenetically correlated model errors in PGLS account for the non-independence of the
species due to their phylogenetic relatedness (Symonds and Blomberg 2014). The dependence of
the model errors arises from trait axes that we did not include in the analysis and that may be
subject to niche conservatism so that model errors reflect the unobserved trait and thus the
phylogenetic distance between species. We used species' similarity in niche position and breadth
as the response variables in the models and explained their variation with categorical trait
variables. Prior to use in models, continuous predictor variables (migration distance and log-
transformed body mass) were scaled to improve model fit and response variables were log-
transformed and sign-flipped (representing niche similarity rather than dissimilarity) to improve
interpretability. We weighted all species points in models by log-transformed minimum sample
size (number of points in the season with less data) to account for uncertainty in seasonal niche
dissimilarity estimates (true error estimates are unavailable when estimating niche dissimilarity).
We fit three models testing the relationships between the four functional traits and three metrics
of niche similarity: seasonal niche position similarity, seasonal niche breadth similarity, and
locational niche position similarity. To evaluate the reliability of multivariate models, we also fit
univariate models for each combination of functional trait predictor and response variable.

To measure the role of phylogeny in both positional and breadth similarity, we quantified
Blomberg's K (Blomberg, Garland Jr et al. 2003) and compared it to a null distribution of K after
randomizing species' responses 1,000 times (picante package; Kembel et al. 2010). $K < 1$
suggests greater than expected phylogenetically-correlated variance within clades, while $K > 1$
suggests variance among clades (i.e., phylogenetically-conserved trait). We also quantified
lambda to assess the extent of phylogenetic conservation of niche metrics via Brownian motion.
We visualized relationships between niche metrics and functional traits using *ggplot2* (Wickham
2011)*, ggExtra* (Attali and Baker 2019) and *RcolorBrewer* (Neuwirth and Neuwirth 2011)*.* We
visualized partial residuals using *visreg* (Breheny, Burchett, and Breheny 2020).

Results

We found all five niche-tracking strategies represented among North American birds (Fig 3). For
example, Red-bellied woodpeckers (*Melanerpes carolinus*; Fig. 3a) are largely stationary birds
occupying highly seasonal environments, with low seasonal similarity between their breeding
and overwintering niches, but high locational similarity between their overwintering and
stationary winter niches. In contrast, Pine grosbeaks (*Pinicola enucleator*; Fig. 3b) are short-
distance migrants and "breadth trackers", tracking niche breadth while shifting position across
seasons. Migratory Swainson's thrushes (*Catharus ustulatus*; Fig. 3c) are "position trackers",
tracking their niche position across seasons, but expanding their niche during the breeding
season. Prothonotary warblers (*Protonotaria citrea*; Fig. 3d) are long-distance migrants that
abandon breeding territories to closely track both niche components across seasons, resulting in
high seasonal and low locational similarity. Finally, Great kiskadees (*Pitangus sulphuratus*; Fig.
3e) are stationary tropical residents with conserved niches across the annual cycle, resulting in
both high seasonal and high locational similarity. We select these species for visualizations
because they represent each strategy well in both two-dimensional niche space (temperature and
EVI only; Fig. 3) and three-dimensional space (remainder of results).

The 619 bird species analyzed spanned the full diversity of niche tracking strategies (Fig. 4;
Appendix 1). Across seasons, the median positional similarity (-log[Mahalonobis distance]) was
-1.95 +/- 0.05 SE (i.e., centroids approximately 3 SDs apart), suggesting that many species are
not closely tracking this niche attribute over the annual cycle (Ponti et al. 2020). For 60 species
(9.7%), centroids were within 1 SD, suggesting highly similar niches in each season for this
subset. The median among-season similarity in niche breadth was 2.17 +/- 0.03, equivalent to a
4:1 ratio in niche breadth between seasons, suggesting that many species expand their niche
breadths as they traverse seasons. 129 species (22%) had a niche breadth similarity equivalent to
a < 2:1 ratio, suggesting similarly-sized seasonal niches.

Figure 3: Diverse niche tracking strategies. (**a**) Red-bellied woodpeckers (*Melanerpes carolinus*) are non-trackers, with low *seasonal* similarity (solid arrow) between their breeding (orange points and ellipsoid) and overwintering (dark purple) two-dimensional niche (presented as standardized ellipse area). Meanwhile, the difference between their overwintering niche and winter conditions at breeding locations (light purple, dotted ellipsoid), or *locational* similarity (dotted arrow), is high. (**b**) Pine grosbeaks (*Pinicola enucleator*) track their niche breadth but shift position across seasons, while (**c**) Swainson's thrushes (*Catharus ustulatus*) track their niche position but expand niche breadth. (**d**) Prothonotary warblers (*Protonotaria citrea*) closely track their niche across seasons via migration, resulting in high seasonal and low locational similarity. (**e**) Great kiskadees (*Pitangus sulphuratus*) track their niche by remaining stationary in less seasonal environments, leading to high seasonal and locational similarity. Position and breadth similarity values are provided in parentheses. Note that this figure displays two-dimensional niche space, though our statistical analyses consider three-dimensional niches, including precipitation as additional dimension.

Tracking niche position was loosely tied to tracking niche breadth; species that tracked niche position more often tracked niche breadth, and *vice versa* (Generalized linear model: β=0.179, R^2=0.073, p<0.001; Fig. 4). 170 (27.5%) species were "complete" niche trackers, tracking both position and breadth more than most species (positive values on both axes in Fig. 4), while 209 (33.8%) were non-trackers, tracking both less than others (negative values in Fig. 4). However, numerous species tracked one but not the other (Fig. 4), suggesting a complex diversity of approaches to seasonal niche tracking across bird species. Only 80 species (12.9%) tracked niche

316 position but expanded breadth, while double that number – 160 (25.8%) – tracked breadth but
317 shifted position.
318

319
Figure 4: Interspecific variation in tracking niche position and breadth across seasons. The figure
shows niche similarities between the breeding and overwintering seasons for the 619 bird species
analyzed based on three niche dimensions (for red highlights, see Fig. 3). Strategies of tracking niche
position, breadth, both, or neither are partitioned into four categories (boxes separated by dashed lines).
Dashed lines highlight the means. Marginal density plots illustrate that species tracking only one niche
component often track niche breadth more closely than niche position, with medians represented by
notches. The trendline represents a linear relationship between niche position and breadth similarity with
associated 95% confidence interval (shading). See Fig. S2 for equivalent patterns for locational similarity.
328
We found seasonal niche tracking strategies to be closely associated with functional traits, and
their role to be consistent across weighted multivariate models accounting for all traits and
phylogenetic structure simultaneously (Fig. 5; Tables S1-S2) as well as univariate models

(Tables S3-S4). As expected, both niche positional and breadth similarity were very strongly
linked to migration distance (Phylogenetic generalized least-squares models: position, β=0.43,
p<0.001; breadth, β=0.25, p<0.001), which was the strongest functional predictor; long-distance
migrants were most likely to maximize similarity, while residents minimized similarity and
others fell in the middle (Fig. 5a,d). Body mass emerged as another important functional
predictor of seasonal niche tracking, especially with regards to positional similarity. Body mass
negatively predicted niche position similarity (β=-0.80, p<0.001) and was positively associated
with breadth similarity (β=0.53, p<0.001). Small-bodied birds maximized seasonal similarity,
while large-bodied birds minimized it (Fig. 5b,e). Finally, both diet ($F_{1,3}$=5.70, p<0.005) and
habitat preference ($F_{1,3}$=9.68, p<0.001) categories described variation in positional similarity,
though only habitat preference drove breadth similarity ($F_{1,3}$=10.15, p<0.005). As predicted,
insectivores maximized positional and breadth similarity, herbivores minimized these, and
omnivores fell in the middle (Fig. 5c,g). Waterbirds tracked their niches more closely than other
species (Fig. 5d,h).
346

Figure 5: Functional traits predict variation in seasonal niche tracking, as shown by partial residual
plots based on phylogenetic least-squares models. Y-axis values represent (a) positional and (b) breadth
similarity. Silhouettes correspond to categorical functional trait values as follows: Dietary preference (left
to right: carnivore, herbivore, insectivore, omnivore); Habitat preference (forest, generalist,
open/grassland, water).

Surprisingly, we found little evidence of a phylogenetic signal in seasonal niche tracking. For
both niche position and breadth, Within-group phylogenetic variation was greater and among-
group variation lesser than expected by chance (Bloomberg's K always < 0.1; position: p<0.001;
breadth: p=0.10; Table S5) and did not support Ornstein-Uhlenbeck trait evolution.
Measurements of lambda indicated a moderate phylogenetic signal for species tracking niche
position (λ=0.66, p<0.001), suggesting possible Brownian-motion trait evolution, but little signal
for niche breadth (λ=0.28, p<0.001).

Species that maximized seasonal niche similarity typically minimized locational niche similarity,
and vice versa (β=-0.408, R^2=0.237, $p<0.001$; Fig. S1). For locational similarity, the link
between tracking niche position and niche breadth was stronger than for seasonal similarity, with
42% of species tracking both components and 29% neither (Fig. S2). Only 15% of species
tracked locational niche position but not breadth, and 13% tracked locational breadth but not
position. Likewise, relationships between functional traits and locational positional similarity
were generally opposite to those observed with seasonal positional similarity. Long-distance
migrants (β=-0.331, $p<0.001$), small-bodied birds (β=0.95, $p<0.001$), and waterbirds ($F_{1,3}$=7.75,
$p<0.001$) each minimized locational positional similarity, though we found no effect of diet. As
with seasonal similarity, we found no evidence for a phylogenetic signal in locational niche
similarity (Table S5).

373
374

Discussion

For mobile organisms, seasonal niche tracking is a central part of the behavioral repertoire for
mediating exposure to novel climate conditions. Here, we assessed the representation of distinct
seasonal niche tracking strategies across North American breeding birds and evaluated the
functional and phylogenetic drivers of these strategies. Our analyses revealed five distinct
seasonal niche tracking strategies and their key functional drivers. We find that only a bit more
than half of species can be categorized as either "complete trackers", tracking both niche position
and breadth across seasons, or "non-trackers", tracking neither. However, almost 40% of species
only track niche position or breadth, revealing complexity in niche tracking strategies that is
previously unexplored in niche-tracking studies. Twice as many species tracked only niche
breadth (26%) compared with those that tracked niche position and expanded niche breadth
(13%), suggesting that tracking breadth is often a prerequisite for tracking position and that birds
may be most specialized in terms of the range of environmental conditions they can tolerate
rather than means. Importantly, niche tracking studies that do not consider niche breadth would
likely have labeled the 26% of species that are niche breadth trackers as non-trackers, ignoring
the role that niche breadth tracking and expansion plays in allowing species to persist over
seasons. Species limited by extreme or highly variable weather conditions rather than climatic
means may be more apt to prioritize niche breadth tracking while allowing niche position to
seasonally fluctuate. Meanwhile, a species with inflexible thermal or dietary requirements that is
specialized only during certain parts of the year (e.g., due to strict breeding requirements) may
track niche position but not breadth. However, further work is needed to pinpoint the ecological
motivations behind selection of breadth tracking as a niche tracking strategy.

We found that migration distance was the most important functional driver of seasonal niche
tracking, despite this link previously receiving mixed support among cross-species niche tracking
studies (Zurell et al. 2018; Laube, Graham, and Böhning-Gaese 2015; Gómez et al. 2016).
Although several studies (Gómez et al. 2016; Laube, Graham, and Böhning-Gaese 2015) failed

to detect a link between migration and niche tracking, their approaches compared species within
single families, including residential species that are exclusively tropical. Because these residents
occupy relatively stable environments, they can track their niches despite being stationary (Eyres
et al. 2020). This example helps illustrate why migration distance and niche tracking are less
correlated than might be expected – tropical residents can track their niches well despite their
stationarity, while long-distance migrants may experience high seasonal variation despite their
migration if they annually move between arctic breeding grounds and boreal or temperate
wintering grounds. Indeed, although migration distance was the strongest correlate of seasonal
niche tracking in our study, the correlation was moderate (R^2=0.249).

Body size was revealed as the next most important functional trait associated with seasonal niche
tracking when controlling for other traits. Small-bodied birds may not be able to tolerate large
variation in their thermal niches because they have low thermal inertia (Huey et al. 2012;
Albright et al. 2017). Further, small-bodied animals have shown greater sensitivity to extreme
weather events (Cohen, Fink, and Zuckerberg 2021) and more rapidly alter phenological timing
in tune with interannual weather variability (Cohen, Lajeunesse, and Rohr 2018) compared with
large-bodied animals, suggesting that they must carefully maintain thermal limits. Additionally,
small-bodied birds may be more likely to evolve niche-tracking strategies dependent on seasonal
migration because long-distance flight carries lower energetic costs than it does for large birds
(Watanabe 2016).

Finally, diet and habitat preference were additional meaningful drivers of seasonal niche tracking
behavior, with insectivores and carnivores tracking their niches more than herbivores and
omnivores, and waterbirds more so than other species. Insectivores may prioritize niche-tracking
because they are dependent upon prey that is most commonly available under specific thermal
and productivity limits (Winkler, Luo, and Rakhimberdiev 2013). Surprisingly, carnivores
tracked their niche positions even more closely than insectivores, but many of these species are
predating on herpetofauna also active during warm weather. Though we predicted both open and
water habitat specialists to be the strongest niche trackers, as they are less shielded from the
external environment by habitat structure (Jarzyna et al. 2016), only water birds tracked their
niche more closely than other groups (Fig 4d,h). Bird species that occupy open habitats may
make greater use of microhabitat structure than water birds, which are typically exposed (e.g.,
(Shew, Nielsen, and Sparling 2019)).

Although many studies have assumed seasonal niche tracking to be phylogenetically conserved
(Gómez et al. 2016; Martínez–Meyer, Townsend Peterson, and Navarro–Sigüenza 2004), this
has not to our knowledge been evaluated over a large, diverse taxonomic group such as North
American breeding birds. There are several explanations as to why we did not detect a
phylogenetic signal. First, migratory behavior has arisen in numerous, diverse lineages of birds,
and is flexible even within species (Zink 2011). Second, closely related species often occupy

distinct climatic zones that vary greatly in seasonality, especially if speciation resulted from

character displacement (Newton 2003). Finally, species-level exposure to and adaptation

potential for climate change may be quite independent from phylogeny (Khaliq et al. 2015;

Davis et al. 2010). Based on our analyses, we conclude that functional traits likely play a larger

role than phylogeny in driving seasonal niche tracking in birds.

Complex variation among niche tracking strategies cannot fully be assessed via seasonal niche

comparisons; for example, a species that niche-tracks via movement and a stationary species in

an aseasonal environment have similar seasonal niches. To parse these strategies, we estimated

locational niche similarity, or the similarity between overwintering niches and the *stationary*

winter niche for each species. Species with high seasonal similarity often had low locational

similarity, and *vice versa*, reflecting a trade-off between conserving niches and ranges

throughout the annual cycle. While many species tracked only niche position or breadth between

seasons, species adhered to locationally tracking both or neither components much more often,

with few tracking just one. Therefore, species may be leaving breeding areas in winter for the

purposes of modifying both the mean and variance of environmental conditions. We observed

several key differences between functional trait relationships and either seasonal or locational

niche similarity. For instance, migration distance and body size were equally important as drivers

of locational niche similarity, possibly because large-bodied birds are especially reluctant to

energetically invest in movement (Watanabe 2016). Further, in contrast with seasonal niche

similarity, there was no effect of diet on locational similarity, perhaps because food availability

depends more on environmental conditions themselves rather than the distance birds travel to

track them. Thus, we demonstrate the importance of considering both seasonal and locational

similarity in species niches to understand variation in niche tracking strategies.

We estimated niche metrics at the species level to effectively compare niche tracking strategies

across a taxonomic group as broad and diverse as birds and to make use of the enormous

quantity of available species-level occurrence data. However, we did not explore niche variation

as experienced at lower levels of organization, including populations, demographic groups, and

individuals (Fandos et al. 2020; Fandos and Tellería 2020; Carlson et al. 2021). Populations of

species with wide geographic ranges might be adapted to highly distinct environmental

conditions (Broggi et al. 2005). Within a population, demographic groups can also occupy

different niches – sexes can occupy spatially distinct regions during the nonbreeding season, or

young can be shielded in microclimate during breeding season (Shipley et al. 2020). Finally,

individuals may track different niche components than populations (Fandos et al. 2020) or might

have strong variation in their ecological niches during different parts of the year and thus vary in

niche-tracking. For example, individual white storks (*Ciconia ciconia*) and sandhill cranes

(*Antigone canadensis*) range from subtropical residents to short-distance migrants to longer-

distance migrants moving between northern temperate and tropical zones each year (Krapu et al.

2014; Fandos et al. 2020; Carlson et al. 2021). Summarizing niche tracking strategies at the

species level may limit our inferential ability in two ways: 1) species-level niche breadth is likely greater than individual niche breadth given variability in individual niches (Carlson et al. 2021), and 2) the accuracy of species-level niche centroids for individuals likely depends on the variation in climate zones at which the species exists, so long as individuals and populations are locally adapted (Araújo and Costa-Pereira 2013). Although most occurrence data is still available at the species level, new technologies such as GPS tracking offer exciting opportunities for assessing individual niches over time and across hierarchies of organismal organization (Jetz et al. 2022; Costa-Pereira et al. 2022) and complement the presented species-level findings.

As climate change progresses, species are increasingly being exposed to conditions that are quite different from those to which they have adapted (Pacifici et al. 2015). Gradual shifts in range boundaries or phenology allow species to keep pace with long-term changes in climatic means (La Sorte and Jetz 2012; Rushing et al. 2020; Koleček, Adamík, and Reif 2020). However, niche tracking is an important tool for species to buffer both the changing mean environmental conditions and increasing interannual variability associated with climate change by selecting optimal environmental conditions over short time scales (Román-Palacios and Wiens 2020), especially for species with narrow thermal or habitat requirements. Our findings reveal a broad diversity in niche tracking strategies and uncover important functional trait associations with migratory behavior and body size. This has ecological consequences for the assemblages and ecosystems that will see different functional perturbations due to the different susceptibility of niche tracking strategies to climatic change (Barbet-Massin and Jetz 2015). For example, as niche tracking becomes more difficult with climate change, small-bodied and insectivorous birds may be especially likely to fail to maintain their niches, resulting in proportional declines of these species relative to the broader avian community assemblage.

Thus, we provide a framework for future studies to assess the degree of tracking both niche position and breadth and capability for climate change adaptation in many other species globally, including those from understudied regions with limited available data. However, additional work is needed to assess how species have already been keeping up with climate change that has occurred. Understanding how climate change is modifying species' ability to conserve their niches will be a critical step towards determining which species are most likely to adapt to increasing short-term variability in climate.

Acknowledgements

We thank J. Makinen and M. Lu for their modeling advice and thoughtful comments on the analyses and J. Wilshire, A. Ranipeta, R. Li, and the Map of Life team at the Yale Center for Biodiversity and Global Change for implementing the Spatiotemporal Observation Annotation Tool to annotate occurrence data. We thank F. La Sorte for providing species-level migration

521 distances. We acknowledge funding from NSF grant DEB-1441737 and NASA grants
522 80NSSC17K0282 and 80NSSC18K0435.
523
524 **Data Sharing and Accessibility**
525
526 GBIF data is available for public use online at https://www.gbif.org/. Code generated to conduct
527 the analyses will be made available in a public repository such as Dryad or Figshare.
528
529 **References**
530

Albright, Thomas P., Denis Mutiibwa, Alexander. R. Gerson, Eric Krabbe Smith, William A. Talbot,
Jacqueline J. O'Neill, Andrew E. McKechnie, and Blair O. Wolf. 2017. "Mapping Evaporative
Water Loss in Desert Passerines Reveals an Expanding Threat of Lethal Dehydration."
Proceedings of the National Academy of Sciences 114 (9): 2283–88.
https://doi.org/10.1073/pnas.1613625114.
Albright, Thomas P., Anna M. Pidgeon, Chadwick D. Rittenhouse, Murray K. Clayton, Brian D.
Wardlow, Curtis H. Flather, Patrick D. Culbert, and Volker C. Radeloff. 2010. "Combined
Effects of Heat Waves and Droughts on Avian Communities across the Conterminous United
States." *Ecosphere* 1 (5): art12. https://doi.org/10.1890/ES10-00057.1.
Araújo, Márcio S., and Raul Costa-Pereira. 2013. "Latitudinal Gradients in Intraspecific Ecological
Diversity." *Biology Letters* 9 (6): 20130778. https://doi.org/10.1098/rsbl.2013.0778.
Association, American Birding. 2008. "American Birding Association Checklist: Birds of the Continental
United States and Canada." American Birding Association, Colorado Springs, Colorado.
Barbet-Massin, Morgane, and Walter Jetz. 2015. "The Effect of Range Changes on the Functional
Turnover, Structure and Diversity of Bird Assemblages under Future Climate Scenarios." *Global
Change Biology* 21 (8): 2917–28. https://doi.org/10.1111/gcb.12905.
Barnagaud, Jean-Yves, Pierre Gaüzère, Benjamin Zuckerberg, Karine Princé, and Jens-Christian
Svenning. 2017. "Temporal Changes in Bird Functional Diversity across the United States."
Oecologia 185 (4): 737–48. https://doi.org/10.1007/s00442-017-3967-4.
Blonder, Benjamin. 2018. "Hypervolume Concepts in Niche- and Trait-Based Ecology." *Ecography* 41
(9): 1441–55. https://doi.org/10.1111/ecog.03187.
Blonder, Benjamin, Christine Lamanna, Cyrille Violle, and Brian J. Enquist. 2014. "The N-Dimensional
Hypervolume." *Global Ecology and Biogeography* 23 (5): 595–609.
Breheny, Patrick, Woodrow Burchett, and Maintainer Patrick Breheny. 2020. "Package 'Visreg.'"
Broggi, Juli, Esa Hohtola, Markku Orell, and Jan-Åke Nilsson. 2005. "Local Adaptation to Winter
Conditions in a Passerine Spreading North: A Common-Garden Approach." *Evolution* 59 (7):
1600–1603.
Brown, Matilda JM, Barbara R. Holland, and Greg J. Jordan. 2020. "Hyperoverlap: Detecting Biological
Overlap in n-Dimensional Space." *Methods in Ecology and Evolution* 11 (4): 513–23.
Carlson, Ben S., Shay Rotics, Ran Nathan, Martin Wikelski, and Walter Jetz. 2021. "Individual
Environmental Niches in Mobile Organisms." *Nature Communications* 12 (1): 1–10.
Clements, James F. 2007. *Clements Checklist of Birds of the World*. Comstock Pub. Associates/Cornell
University Press.
Cohen, Jeremy M., Daniel Fink, and Benjamin Zuckerberg. 2021. "Extreme Winter Weather Disrupts
Bird Occurrence and Abundance Patterns at Geographic Scales." *Ecography* 44 (8): 1143–55.
Cohen, Jeremy M., Marc J. Lajeunesse, and Jason R. Rohr. 2018. "A Global Synthesis of Animal
Phenological Responses to Climate Change." *Nature Climate Change* 8 (3): 224–28.
https://doi.org/10.1038/s41558-018-0067-3.

Costa-Pereira, Raul, Remington J. Moll, Brett R. Jesmer, and Walter Jetz. 2022. "Animal Tracking Moves Community Ecology: Opportunities and Challenges." *Journal of Animal Ecology*.

Davis, Charles C., Charles G. Willis, Richard B. Primack, and Abraham J. Miller-Rushing. 2010. "The Importance of Phylogeny to the Study of Phenological Response to Global Climate Change." *Philosophical Transactions of the Royal Society B: Biological Sciences* 365 (1555): 3201–13. https://doi.org/10.1098/rstb.2010.0130.

Eyres, Alison, Katrin Böhning-Gaese, C. David L. Orme, Carsten Rahbek, and Susanne A. Fritz. 2020. "A Tale of Two Seasons: The Link between Seasonal Migration and Climatic Niches in Passerine Birds." *Ecology and Evolution* 10 (21): 11983–97.

Fandos, Guillermo, Shay Rotics, Nir Sapir, Wolfgang Fiedler, Michael Kaatz, Martin Wikelski, Ran Nathan, and Damaris Zurell. 2020. "Seasonal Niche Tracking of Climate Emerges at the Population Level in a Migratory Bird." *Proceedings of the Royal Society B: Biological Sciences* 287 (1935): 20201799. https://doi.org/10.1098/rspb.2020.1799.

Fandos, Guillermo, and José Luis Tellería. 2020. "Seasonal Niche-Tracking Behaviour of Two Partially Migratory Passerines." *Ibis* 162 (2): 307–17.

Gómez, Camila, Elkin A. Tenorio, Paola Montoya, and Carlos Daniel Cadena. 2016. "Niche-Tracking Migrants and Niche-Switching Residents: Evolution of Climatic Niches in New World Warblers (Parulidae)." *Proceedings of the Royal Society B: Biological Sciences* 283 (1824): 20152458.

Huey, Raymond B., Michael R. Kearney, Andrew Krockenberger, Joseph A. M. Holtum, Mellissa Jess, and Stephen E. Williams. 2012. "Predicting Organismal Vulnerability to Climate Warming: Roles of Behaviour, Physiology and Adaptation." *Philosophical Transactions of the Royal Society B: Biological Sciences* 367 (1596): 1665–79. https://doi.org/10.1098/rstb.2012.0005.

Hutchinson, G. E. 1957. "The Multivariate Niche." In *Cold Spring Harbor Symposia on Quantitative Biology*, 22:415–21.

Jarzyna, Marta A., Benjamin Zuckerberg, Andrew O. Finley, and William F. Porter. 2016. "Synergistic Effects of Climate and Land Cover: Grassland Birds Are More Vulnerable to Climate Change." *Landscape Ecology* 31 (10): 2275–90. https://doi.org/10.1007/s10980-016-0399-1.

Jetz, Walter, Grigori Tertitski, Roland Kays, Uschi Mueller, Martin Wikelski, Susanne Åkesson, Yury Anisimov, et al. 2022. "Biological Earth Observation with Animal Sensors." *Trends in Ecology & Evolution* 37 (4): 293–98. https://doi.org/10.1016/j.tree.2021.11.011.

Karger, Dirk Nikolaus, Adam M Wilson, Colin Mahony, and Niklaus E Zimmermann. 2021. "Global Daily 1km Land Surface Precipitation Based on Cloud Cover-Informed Downscaling," 20.

Kembel, Steven W., Peter D. Cowan, Matthew R. Helmus, William K. Cornwell, Helene Morlon, David D. Ackerly, Simon P. Blomberg, and Campbell O. Webb. 2010. "Picante: R Tools for Integrating Phylogenies and Ecology." *Bioinformatics* 26 (11): 1463–64.

Khaliq, Imran, Susanne A. Fritz, Roland Prinzinger, Markus Pfenninger, Katrin Böhning-Gaese, and Christian Hof. 2015. "Global Variation in Thermal Physiology of Birds and Mammals: Evidence for Phylogenetic Niche Conservatism Only in the Tropics." *Journal of Biogeography* 42 (11): 2187–96. https://doi.org/10.1111/jbi.12573.

Koleček, Jaroslav, Peter Adamík, and Jiří Reif. 2020. "Shifts in Migration Phenology under Climate Change: Temperature vs. Abundance Effects in Birds." *Climatic Change* 159 (2): 177–94. https://doi.org/10.1007/s10584-020-02668-8.

Krapu, Gary L., David A. Brandt, Paul J. Kinzel, and Aaron T. Pearse. 2014. "Spring Migration Ecology of the Mid-Continent Sandhill Crane Population with an Emphasis on Use of the Central Platte River Valley, Nebraska." *Wildlife Monographs* 189 (1): 1–41.

La Sorte, Frank A., and Walter Jetz. 2012. "Tracking of Climatic Niche Boundaries under Recent Climate Change: Niche Tracking under Recent Climate Change." *Journal of Animal Ecology* 81 (4): 914–25. https://doi.org/10.1111/j.1365-2656.2012.01958.x.

La Sorte, Frank A., Alison Johnston, Amanda D. Rodewald, Daniel Fink, Andrew Farnsworth, Benjamin M. Van Doren, Tom Auer, and Matthew Strimas-Mackey. 2022. "The Role of Artificial Light at

Night and Road Density in Predicting the Seasonal Occurrence of Nocturnally Migrating Birds."
Diversity and Distributions.
Laube, Irina, Catherine H. Graham, and Katrin Böhning-Gaese. 2015. "Niche Availability in Space and
Time: Migration in Sylvia Warblers." *Journal of Biogeography* 42 (10): 1896–1906.
Li, Richard, Ajay Ranipeta, John Wilshire, Jeremy Malczyk, Michelle Duong, Robert Guralnick, Adam
Wilson, and Walter Jetz. 2021. "A Cloud-Based Toolbox for the Versatile Environmental
Annotation of Biodiversity Data." Edited by Andrew J. Tanentzap. *PLOS Biology* 19 (11):
e3001460. https://doi.org/10.1371/journal.pbio.3001460.
Lu, Muyang, Kevin Winner, and Walter Jetz. 2021. "A Unifying Framework for Quantifying and
Comparing N-Dimensional Hypervolumes." *Methods in Ecology and Evolution* 12 (10): 1953–
68.
Martínez–Meyer, Enrique, A. Townsend Peterson, and Adolfo G. Navarro–Sigüenza. 2004. "Evolution of
Seasonal Ecological Niches in the Passerina Buntings (Aves: Cardinalidae)." *Proceedings of the*
Royal Society of London. Series B: Biological Sciences 271 (1544): 1151–57.
Newton, Ian. 2003. *Speciation and Biogeography of Birds*. Academic Press.
Ooms, Jeroen. 2014. "The Jsonlite Package: A Practical and Consistent Mapping between Json Data and r
Objects." *ArXiv Preprint ArXiv:1403.2805*.
Orme, David, R. Freckleton, G. Thomas, T. Petzoldt, S. Fritz, N. Isaac, and W. Pearse. 2013. "The Caper
Package: Comparative Analysis of Phylogenetics and Evolution in R." *R Package Version* 5 (2):
1–36.
Outlaw, Diana C., and Gary Voelker. 2006. "Phylogenetic Tests of Hypotheses for the Evolution of
Avian Migration: A Case Study Using the Motacillidae." *The Auk* 123 (2): 455–66.
Pacifici, Michela, Wendy B. Foden, Piero Visconti, James E. M. Watson, Stuart H.M. Butchart, Kit M.
Kovacs, Brett R. Scheffers, et al. 2015. "Assessing Species Vulnerability to Climate Change."
Nature Climate Change 5 (3): 215–24. https://doi.org/10.1038/nclimate2448.
Ponti, Raquel, Angel Arcones, Xavier Ferrer, and David R. Vieites. 2020. "Seasonal Climatic Niches
Diverge in Migratory Birds." *Ibis* 162 (2): 318–30.
Qian, Hong. 2010. "Environment–Richness Relationships for Mammals, Birds, Reptiles, and Amphibians
at Global and Regional Scales." *Ecological Research* 25 (3): 629–37.
R Core Team. 2021. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R
Foundation for Statistical Computing. https://www.R-project.org/.
Rahmstorf, S., and D. Coumou. 2011. "Increase of Extreme Events in a Warming World." *Proceedings of*
the National Academy of Sciences 108 (44): 17905–9. https://doi.org/10.1073/pnas.1101766108.
Román-Palacios, Cristian, and John J. Wiens. 2020. "Recent Responses to Climate Change Reveal the
Drivers of Species Extinction and Survival." *Proceedings of the National Academy of Sciences*
117 (8): 4211–17. https://doi.org/10.1073/pnas.1913007117.
Rushing, Clark S., J. Andrew Royle, David J. Ziolkowski, and Keith L. Pardieck. 2020. "Migratory
Behavior and Winter Geography Drive Differential Range Shifts of Eastern Birds in Response to
Recent Climate Change." *Proceedings of the National Academy of Sciences*, May, 202000299.
https://doi.org/10.1073/pnas.2000299117.
Shew, Justin J., Clayton K. Nielsen, and Donald W. Sparling. 2019. "Finer-Scale Habitat Predicts Nest
Survival in Grassland Birds More than Management and Landscape: A Multi-Scale Perspective."
Journal of Applied Ecology 56 (4): 929–45.
Shipley, J. Ryan, Cornelia W. Twining, Conor C. Taff, Maren N. Vitousek, Andrea Flack, and David W.
Winkler. 2020. "Birds Advancing Lay Dates with Warming Springs Face Greater Risk of Chick
Mortality." *Proceedings of the National Academy of Sciences* 117 (41): 25590–94.
Somveille, Marius, Ana SL Rodrigues, and Andrea Manica. 2015. "Why Do Birds Migrate? A
Macroecological Perspective." *Global Ecology and Biogeography* 24 (6): 664–74.
———. 2018. "Energy Efficiency Drives the Global Seasonal Distribution of Birds." *Nature Ecology &*
Evolution 2 (6): 962–69.

Tingley, M. W., W. B. Monahan, S. R. Beissinger, and C. Moritz. 2009. "Birds Track Their Grinnellian
Niche through a Century of Climate Change." *Proceedings of the National Academy of Sciences*
106 (Supplement_2): 19637–43. https://doi.org/10.1073/pnas.0901562106.
Ummenhofer, Caroline C., and Gerald A. Meehl. 2017. "Extreme Weather and Climate Events with
Ecological Relevance: A Review." *Philosophical Transactions of the Royal Society B: Biological*
Sciences 372 (1723): 20160135. https://doi.org/10.1098/rstb.2016.0135.
Watanabe, Yuuki Y. 2016. "Flight Mode Affects Allometry of Migration Range in Birds." *Ecology*
Letters 19 (8): 907–14.
Wickham, Hadley. 2011. "Ggplot2." *Wiley Interdisciplinary Reviews: Computational Statistics* 3 (2):
180–85.
Wickham, Hadley, Mara Averick, Jennifer Bryan, Winston Chang, Lucy D'Agostino McGowan, Romain
François, Garrett Grolemund, Alex Hayes, Lionel Henry, and Jim Hester. 2019. "Welcome to the
Tidyverse." *Journal of Open Source Software* 4 (43): 1686.
Wickham, Hadley, and Maintainer Hadley Wickham. 2020. "Package 'Httr.'"
Wilman, Hamish, Jonathan Belmaker, Jennifer Simpson, Carolina de la Rosa, Marcelo M. Rivadeneira,
and Walter Jetz. 2014. "EltonTraits 1.0: Species-Level Foraging Attributes of the World's Birds
and Mammals: *Ecological Archives* E095-178." *Ecology* 95 (7): 2027–2027.
https://doi.org/10.1890/13-1917.1.
Winger, Benjamin M., Giorgia G. Auteri, Teresa M. Pegan, and Brian C. Weeks. 2019. "A Long Winter
for the Red Queen: Rethinking the Evolution of Seasonal Migration." *Biological Reviews* 94 (3):
737–52.
Winkler, David W, Miles K Luo, and Eldar Rakhimberdiev. 2013. "Temperature Effects on Food Supply
and Chick Mortality in Tree Swallows (Tachycineta Bicolor)," 12.
Zink, Robert M. 2011. "The Evolution of Avian Migration." *Biological Journal of the Linnean Society*
104 (2): 237–50.
Zurell, Damaris, Laure Gallien, Catherine H. Graham, and Niklaus E. Zimmermann. 2018. "Do Long-
Distance Migratory Birds Track Their Niche through Seasons?" *Journal of Biogeography* 45 (7):
1459–68. https://doi.org/10.1111/jbi.13351.
697

Supporting Information

Title: Diverse strategies for tracking seasonal environmental niches at hemispheric scale

Jeremy Cohen*[1,2] & Walter Jetz[1,2]

[1]Department of Ecology and Evolutionary Biology, Yale University, 165 Prospect Street, New Haven, CT, 06520 USA
[2]Center for Biodiversity and Global Change, Yale University, 165 Prospect Street, New Haven, CT, 06520 USA

*Correspondence to: jeremy3cohen@gmail.com

Figure S1. Seasonal and locational niche centroid similarity are inversely related. Niche-tracking species maximize niche similarity between breeding and nonbreeding seasons (defined as seasonal similarity, x-axis) and minimize the similarity between the nonbreeding season niche and the stationary winter niche (defined as locational similarity, y-axis). The reverse is true for non-tracker species. Tropical residents are an exception to this rule; these species have overlapping niche space for all three components, thus maximizing both seasonal and locational similarity. Dashed lines represent means and gray shading represents 95% confidence interval for trend line.

Figure S2: Interspecific variation in locational niche tracking of niche position and breadth.
Locational niche similarity represents the similarity of the overwintering niche to the winter conditions at
breeding locations. Points represent species-level variation in locational similarity of niche position and
breadth between the breeding and overwintering seasons for 619 species of North American birds.
Marginal density plots illustrate a positive relationship between tracking locational niche position and
breadth. Darker points reflect greater values of seasonal positional niche similarity and are primarily
distributed among points with lower values of locational positional similarity.

Appendix 1. Seasonal niche dissimilarity for 619 species of North American birds (sorted taxonomically). Dissimilarity is partitioned into Mahalanobis distance and determinant ratio for both seasonal (breeding/overwintering comparison) and locational (overwintering/winter at breeding sites comparison) dissimilarity. Position and breadth similarity (visualized and used in analyses) was generated from these values by ln-transforming and sign-flipping.

(appendix_1.csv)

Appendix 2. Updates to avian phylogeny. We updated the avian phylogeny of Jetz et al (2012) to the 2021 Clements taxonomy by harmonizing species names, as follows.

(appendix_2.csv)

Table S1. Model tables (pgls and ANOVA) summarizing functional trait drivers of seasonal niche position and breadth similarity across species in multivariate models.

Mahalonobis distance

PGLS table	Coefficient	SE	t-value	p-value
Intercept	-1.745	0.270	-6.472	<0.001
Migration distance	0.428	0.041	10.359	<0.001
Body mass (log)	-0.798	0.097	-8.233	<0.001
Diet (herbivore)	-0.647	0.178	-3.626	<0.001
Diet (insectivore)	-0.048	0.177	-0.275	0.784
Diet (omnivore)	-0.394	0.139	-2.836	0.005
Habitat (generalist)	0.001	0.174	0.005	0.996
Habitat (open)	-0.381	0.110	-3.470	0.001
Habitat (water)	0.364	0.172	2.110	0.035

Mahalonobis distance

ANOVA table	DF	F-value	p-value
Intercept	1	121.713	<0.001
Migration distance	1	197.317	<0.001
Body mass (log)	1	93.094	<0.001
Diet	3	5.702	0.001
Habitat preference	3	9.675	<0.001

Determinant ratio

PGLS table	Coefficient	SE	t-value	p-value
Intercept	1.360	0.262	5.183	<0.001
Migration distance	0.250	0.040	6.212	<0.001
Body mass (log)	0.525	0.094	5.565	<0.001
Diet (herbivore)	-0.053	0.174	-0.305	0.760
Diet (insectivore)	0.206	0.172	1.196	0.232
Diet (omnivore)	0.102	0.135	0.754	0.451
Habitat (generalist)	-0.081	0.169	-0.481	0.631
Habitat (open)	-0.400	0.107	-3.738	<0.001
Habitat (water)	0.319	0.168	1.905	0.057

Determinant ratio

ANOVA table	DF	F-value	p-value
Intercept	1	185.261	<0.001
Migration distance	1	37.702	<0.001
Body mass (log)	1	51.669	<0.001
Diet	3	2.201	0.087

| Habitat preference | 3 | 10.145 | <0.001 |

Table S2. Model tables (pgls and ANOVA) summarizing functional trait drivers of locational niche position similarity across species in multivariate models.

Mahalonobis distance

PGLS table	Coefficient	SE	t-value	p-value
Intercept	0.182	0.282	0.645	0.519
Migration distance	-0.331	0.043	-7.660	<0.001
Body mass (log)	0.952	0.101	9.389	<0.001
Diet (herbivore)	-0.215	0.187	-1.155	0.249
Diet (insectivore)	-0.176	0.185	-0.956	0.340
Diet (omnivore)	-0.092	0.145	-0.637	0.524
Habitat (generalist)	0.822	0.182	4.519	<0.001
Habitat (open)	0.415	0.115	3.609	<0.001
Habitat (water)	0.515	0.180	2.861	0.004

Mahalonobis distance

ANOVA table	DF	F-value	p-value
Intercept	1	11.162	0.001
Migration distance	1	111.730	<0.001
Body mass (log)	1	111.154	<0.001
Diet	3	0.703	0.551
Habitat preference	3	7.757	<0.001

Table S3. Model tables (pgls) summarizing functional trait drivers of seasonal niche position and breadth similarity across species in univariate models.

Mahalonobis distance PGLS table	Coefficient	SE	t-value	p-value
Intercept	-3.060	0.214	-14.291	0.000
Migration distance	0.539	0.042	12.747	0.000

Mahalonobis distance PGLS table	Coefficient	SE	t-value	p-value
Intercept	-1.039	0.219	-4.744	0.000
Body mass (log)	-1.057	0.097	-10.883	0.000

Mahalonobis distance PGLS table	Coefficient	SE	t-value	p-value
(Intercept)	-1.254	0.257	-4.876	0.000
Diet (herbivore)	-1.287	0.202	-6.380	0.000
Diet (insectivore)	0.085	0.208	0.410	0.682
Diet (omnivore)	-0.681	0.155	-4.409	0.000

Mahalonobis distance PGLS table	Coefficient	SE	t-value	p-value
Intercept	-1.864	0.260	-7.168	0.000
Habitat (generalist)	-0.356	0.211	-1.684	0.093
Habitat (open)	-0.375	0.134	-2.796	0.005
Habitat (water)	0.121	0.208	0.578	0.563

Determinant ratio PGLS table	Coefficient	SE	t-value	p-value
Intercept	1.891	0.201	9.393	<0.001
Migration distance	0.229	0.040	5.770	<0.001

Determinant ratio PGLS table	Coefficient	SE	t-value	p-value
Intercept	1.938	0.201	9.659	<0.001
Body mass (log)	0.480	0.089	5.397	<0.001

Determinant ratio PGLS table	Coefficient	SE	t-value	p-value
(Intercept)	2.092	0.227	9.213	<0.001
Diet (herbivore)	0.071	0.178	0.401	0.688
Diet (insectivore)	0.354	0.183	1.934	0.054
Diet (omnivore)	0.435	0.136	3.190	0.001

Determinant ratio PGLS table	Coefficient	SE	t-value	p-value
Intercept	2.304	0.218	10.572	<0.001
Habitat (generalist)	-0.161	0.177	-0.909	0.364
Habitat (open)	-0.437	0.113	-3.881	<0.001
Habitat (water)	0.499	0.175	2.858	0.004

Table S4. Model tables (pgls) summarizing functional trait drivers of locational niche position similarity across species in univariate models.

Mahalonobis distance PGLS table	Coefficient	SE	t-value	p-value
Intercept	1.485	0.224	6.643	<0.001
Migration distance	-0.424	0.044	-9.610	<0.001

Mahalonobis distance PGLS table	Coefficient	SE	t-value	p-value
Intercept	-0.357	0.215	-1.658	0.098
Body mass (log)	1.119	0.095	11.721	<0.001

Mahalonobis distance PGLS table	Coefficient	SE	t-value	p-value
(Intercept)	0.331	0.265	1.252	0.211
Diet (herbivore)	0.521	0.208	2.510	0.012
Diet (insectivore)	-0.255	0.214	-1.194	0.233
Diet (omnivore)	0.351	0.159	2.207	0.028

Mahalonobis distance PGLS table	Coefficient	SE	t-value	p-value
Intercept	0.091	0.256	0.354	0.723
Habitat (generalist)	1.068	0.208	5.124	<0.001
Habitat (open)	0.408	0.132	3.085	0.002
Habitat (water)	0.774	0.205	3.770	<0.001

Table S5. Metrics describing phylogenetic signal in seasonal niche tracking. Asterisks denote significance level (*** < 0.001 < ** < 0.01 < * < 0.05).

	Blomberg's K		Lambda	
	Seasonal dissimilarity	**Locational dissimilarity**	**Seasonal dissimilarity**	**Locational dissimilarity**
Mahalonobis distance	0.083***	0.115***	0.664***	0.663***
Determinant Ratio	0.057*		0.278***	