The roles of isolation and interspecific interaction in generating the functional diversity of an insular mammal radiation

Abstract

Communities that occupy similar environments but vary in the richness of closely related species can illuminate how functional variation and species richness interact to fill ecological space in the absence of abiotic filtering, though this has yet to be explored on an oceanic island where the processes of community assembly may differ from continental settings. In discrete montane communities on the island of Sulawesi, local murine rodent (rats and mice) richness ranges from 7 to 23 species. We measured 17 morphological, ecological, and isotopic traits, both individually and grouped into 5 multivariate traits in 40 species, to test for the expansion or packing of functional space among nine murine communities. We employed a novel probabilistic approach for integrating intraspecific and community-level trait variance into functional richness. Trait-specific and phylogenetic diversity patterns indicate dynamic community assembly due to variable niche expansion and packing on multiple niche axes. Locomotion and covarying traits such as tail length emerged as a fundamental axis of ecological variation, expanding functional space and enabling the niche packing of other traits such as diet and body size. Though trait divergence often explains functional diversity in island communities, we found that phylogenetic diversity facilitates functional space expansion in some conserved traits such as cranial shape, while more labile traits are overdispersed both within and between island clades, suggesting a role of niche complementarity. Our results evoke interspecific interactions, differences in trait lability, and the independent evolutionary trajectories of each of Sulawesi's 6 murine clades as central to generating the exceptional functional diversity and species richness in this exceptional, insular radiation.

27   Bahasa Indonesia Abstract translation to be provided in subsequent version.

Introduction

Whether at a local, continental, or global scale, species richness is not evenly distributed across the landscape. This unevenness emerges from both environmental and resource heterogeneity among communities, and from the interactions among the species within local communities. Local species richness is often positively correlated with the complexity of habitat structure and diversity of available resources. The observation that different localities with similar habitat structure, resource availability, and historical access often contain ecologically similar communities led to the prediction that species only co-occur if they partition niche space along some axis, otherwise one will be excluded through competition (Hutchinson 1957, MacArthur & Levins 1967, May & MacArthur 1972, Brown & Lieberman 1973, Brown 1975, Pianka 1974, M'Closkey 1978). Competitive exclusion can be mitigated if two co-occurring species use a narrower breadth of resources, producing a more densely packed community niche space ("niche packing"), of if they exploit habitats or resources that are unused or non-existent in low-resource or species-poor communities, leading to a larger community niche space ("niche expansion"; MacArthur 1965, 1970, Pigot et al. 2016, Oliveira et al. 2020). The foundational work on competition's role in community assembly relied on empirical data from continental communities with equal biogeographic accessibility but variation in primary productivity or habitat complexity (Brown & Lieberman 1973, Brown 1975, M'Closkey 1978, Karr & James 1975, Pianka 1974, Weiher & Keddy 1995), or from archipelagos where island size or geographic complexity determines resource availability and habitat area, both of which influence overall species richness (Wilson 1961, Diamond 1975, Lister 1976, Gillespie 2004, Losos 2009, Losos & Ricklefs 2009). Less explored are insular areas of similar habitat structure and resource

availability, but with discrete communities that vary in species richness ("species richness

anomalies", Swenson et al. 2016 pg. E83). Yet, these anomalies offer powerful systems for

interrogating the role of competition in the distribution of functional diversity within and among

communities because the effects of abiotic processes such as habitat filtering are minimized

(Swenson et al 2016, Li et al. 2017). Most studies of species richness anomalies have examined

plants in continental settings (Latham & Ricklefs 1993, Swenson et al. 2016, Xu et al. 2019)

where environmental and historical processes influence the regional species pool. Few, if any,

studies have tested the role of competition in the assembly of discrete communities nested within

an insular setting where species pools are formed through long-distance colonization and in-situ

diversification.

Oceanic islands are often hotspots of diversification and endemism, and their unique

biogeographic and historical conditions can decouple functional and phylogenetic diversity

patterns, which helps illuminate the ecological and evolutionary processes that structure local

communities (Losos & Ricklefs 2009). Lineages that arrive and diversify in isolated locations

regularly undergo ecological shifts that result in behaviors or phenotypes that are uncommon or

non-existent in source communities (Millen 2006, Pinto et al. 2008, Esselstyn et al. 2012, 2015,

2021, Stroud & Losos 2016). The presence of high functional and ecological disparity within an

endemic radiation may affect community structure in multiple ways. First, there is some

evidence that community assembly occurs differently on continents and islands. Using an

assortment of phenotypic and behavioral proxies for resource use across a variety of spatial

scales, niche packing is frequently invoked as the primary process of assembly in species-rich

animal communities on continents (Brown 1975, Karr & James 1975, M'Closkey 1978,

MacArthur & MacArthur 1961, Pianka 1974, Pigot et al. 2016, Van de Perre et al. 2020).

However, niche expansion has found some support among isolated island communities of many

closely related species (Lister 1976). Second, multiple lineages evolving in sympatry in isolated

settings can reduce the degree of phylogenetic niche conservatism, and instead promote niche

divergence (Losos et al. 2003). Decreased niche conservatism affects community assembly by

limiting the impact of phylogenetic diversity on community niche breadth. Importantly, by

quantifying trait differences among closely related species, decreased niche conservatism can

illuminate which traits are most evolutionary labile and/or important for resource partitioning

(Losos et al. 2003, Hiller et al. 2019, Dorey et al. 2020, Stroud 2021).

The murine rodent fauna (rats and mice in the subfamily Murinae) of Sulawesi,

Indonesia, a mountainous, tropical, oceanic island at the center of the Wallacean biodiversity

hotspot (Figure 1), is an intriguing system for testing patterns of community niche occupancy.

First, the murine diversity of Sulawesi is exceptional, with at least 48 endemic species stemming

from seven colonist ancestors that arrived from circa 6 Ma to < 1 Ma (Figure S1, Rowe et al.

2019, Handika et al. 2021). While some colonists spawned small radiations of species, others are

evidenced by only a single living species (Table 1, Figure S1). Each clade is widespread on

Sulawesi and most contribute to species richness of all local murine communities on the island.

Second, Sulawesi contains some of the most unusual rodent forms found anywhere (Figure 1b,

Esselstyn et al. 2012, 2015, Rowe et al. 2014) living in sympatry with more typical "rat-like"

morphologies and ecologies. Sulawesi murines have an array of cranial shapes that reflect their

dietary preferences (Figure 1b; Esselstyn et al. 2012, 2015, Martinez et al. 2018), consume a

broad range of foods such as fruit, seeds, fungi, leaves, roots, and earthworms (Musser 2014,

Rowe et al. 2016a), have body sizes ranging from 10 to 500g, and occupy a variety of locomotor

modes (arboreal; scansorial; terrestrial; amphibious, Nations et al. 2021). Third, due to the

topographic complexity of Sulawesi and the positive correlation between elevation and small

mammal diversity in this region (Heaney 2001, Esselstyn et al. 2021), the murine communities

on the islands are partitioned into discrete, montane assemblages (Figure 1). The local montane

murine communities, defined as the species living on a mountain from the upper-lowland forest

to the peak, range from 7 to 23 species, with the upper limit being perhaps the most diverse local

community of closely related terrestrial mammals on Earth. Lastly, the variation in local

community richness does not appear to be determined by environmental or habitat differences.

Maximum elevation, which may correlate with the area occupied by different habitats, appears to

play only a minor role in species richness (Figure 1a), and net primary productivity is nearly

constant across the island's montane regions (Imhoff et al. 2004), suggesting that environmental

filtering, often a central process of community assembly (Webb et al. 2002, Cavender-Bares et

al. 2004, Li et al. 2017), does not greatly affect the functional diversity of Sulawesi's murine

communities. Together, these properties generate a fascinating hierarchical organization: a

species pool of small mammals with disparate ecologies and morphologies living in discrete

montane communities that, due to their habitat similarity and striking disparity in richness,

represent species-richness anomalies, all of which lie within a remote, oceanic island. Such a

system presents an intriguing natural experiment for testing alternative hypotheses of how

communities assemble to occupy ecological niche space.

Figure 1: Small mammal surveys of nine mountains a) on the oceanic island of Sulawesi revealed varying murine
rodent species richness across the island. Each mountain on the map is labeled with the maximum elevation and the
number of murine rodent species present. All but Latimojong were surveyed within 600m of the summit. b) The
diversity of Sulawesi murines is exceptional and includes unique forms that live alongside species with more
"typical" ecologies and morphologies, as demonstrated from surfaces of cranial μCT scans of *i – Rattus hoffmanni,* a
"typical" murine morphology and ecology, *ii* – the shrew rat *Echiothrix leucura* with its extremely elongate rostrum
and soft invertebrate diet, *iii* – the arboreal *Haeromys minahassae*, with a short rostrum and very small size. c)
Photographs: *i – Rattus hoffmanni*, *ii –Echiothrix leucura*, and *iii – Haeromys minahassae*.

Here, we test whether increased species richness in environmentally similar communities

of montane rodents leads to the expansion or packing of ecological niche space. To infer the

ecological niche breadth of each montane assemblage, we quantify diet, trophic dimension, and

microhabitat using 12 individual functional and ecological traits and five multivariate trait

complexes to estimate both the volume and density of community functional space, which we

define as the sum of the n-dimensional functional volumes of the species therein. Importantly,

individual traits or groups of traits related to the same niche axis can reveal distinct, interacting

processes (Spasojevic & Suding 2012, Pigot et al. 2016, Kohli et al. 2021). Additionally, we

examine the role of phylogenetic niche conservatism and trait lability in the assembly of these

nine montane communities by testing for the influence of phylogenetic diversity and inter-clade

variation on functional space occupancy.

Clade	Crown Age	Species Sampled
Echiothrix	5.11 (4.49-5.69)	10 (10)
Maxomys	3.72 (3.21-4.29)	5 (5)
Bunomys	3.45 (3.09-3.81)	15 (15)
Margaretamys	2.92 (2.29-3.51)	2 (4)
Rattus*	1.57 (1.22-1.95)	6 (6)
Haeromys[†]	Indeterminate	1 (1)

Table 1: Ages of six clades descended from Sulawesi colonists. Median age in millions of years is reported with
95% credible intervals in parentheses. Species sampled reports the number of species in this study with overall clade
richness in parentheses and demonstrates near complete sampling. The two unsampled *Margaretamys* species were
not detected in the localities in this study. All ages taken from Rowe et al. 2019. *There were likely two
colonizations by the ancestors of native *Rattus* spp., the second of which occurred 1.16-0.6 Ma (Rowe et al. 2019),
however, all *Rattus* in this study form a clade relative to other Sulawesi murines. Two human commensal *Rattus*
spp. found on Sulawesi were excluded. [†]The age of arrival of the Haeromys clade is unknown as other species of
Haeromys from Borneo have yet to be included in phylogenetic analyses.

151   **Materials and Methods**

The distribution of species- We compiled occurrence records from nine small mammal

inventories of mountain regions on Sulawesi, Indonesia, including one well-documented

mountain surveyed from 1973 to 1976 by Guy Musser and colleagues (Mt. Nokilalaki; Musser,

2014) and eight mountains surveyed between 2011 and 2016 (Ambang, Bawakaraeng,

Buliohuto, Dako, Gandang Dewata, Katopasa, Latimojong, and Torompupu; Figure 1a).

All surveys began in lower primary forest near the line of anthropogenic forest clearing (1100m

to 1500m) and extended to upper-montane forests. All surveys extended to within 600m

elevation of the summit except for Latimojong (highest survey site at 2535m, summit at 3400m).

Historical surveys by Musser lasted several months and were conducted over four years,

employing a mix of snap traps and live traps. Modern surveys (2011-16) lasted an average of 17

days (11-25) and employed similar collection methods, including a mix of snap traps, live traps,

and 20-30L pitfall buckets. All the murine rodent species known from the sampled localities

(Musser 2014, Wilson et al. 2019) were collected during these modern surveys, indicating a

thorough sampling effort. Five new taxa that were discovered during these expeditions have been

described (Musser 2014, Esselstyn et al. 2012, Rowe et al. 2014, Esselstyn et al. 2015, Rowe et

al. 2016b) and several new locality records resulted (Achmadi, et al., 2014; Handika, et al.,

2021). Specimens from all surveys were deposited in the Museum Zoologicum Bogoriense

(MZB), Bogor, Indonesia; the American Museum of Natural History (AMNH), New York, USA;

Museums Victoria (MV), Melbourne, Australia; the Museum of Vertebrate Zoology (MVZ),

Berkeley, USA; the Field Museum of Natural History (FMNH), Chicago, USA; and the

Louisiana State University Museum of Natural Science (LSUMZ), Baton Rouge, USA.

Functional trait data collection and processing- We compiled or generated functional trait

values for 11 continuous traits and one discrete trait for all available Sulawesi murines from each

of the nine communities. Detailed information on traits, sample sizes, data processing, and

multivariate trait composition are available in Table S1. All data, models, and output files are

available in Dryad Repository (to be added prior to publication).

Morphological data collection- We assembled external measurement data from 630 specimens,

including head-body length (mm), tail length (mm), hind-foot length (mm), ear length (mm), and

mass (g), from previously published sources (Wilson et al. 2019, Nations et al. 2021) and online

museum databases. Measured specimens were from the nine surveyed mountains and other

localities on Sulawesi. To obtain ecologically relevant features of external measurements and

mitigate the influence of size in some of our analyses, we calculated three commonly used ratios:

Relative tail length (tail length / (head-body length + tail length)), relative hind-foot length (hind-

foot length / head-body length), and relative ear length (ear length / head-body length) (Nations

et al. 2021, Table S1).

The shape of a rodent's skull and lower jaw provides a wealth of indirect ecological

information on foraging, feeding, and sensory processing, and is often used as a proxy for

fundamental dietary niche (Samuels 2009). We generated μCT scans of the cranium of 64

specimens from 38 species and the mandible of 61 specimens from 36 species (Table S1). Scans

were generated from specimens collected in the nine surveys, as well as from previously

collected museum materials. Stacks of 2D Tiff files were imported to MorphoDig, where 3D

landmarks were placed on cropped volume renderings (Lebrun, 2018). We placed 67 cranial

landmarks (Figure S2) on the left side of the skull, unless damage caused us to use the right side,

in which case we reversed the rendering on the Z-axis. In separate renderings, we placed 20

landmarks on the left dentary of the mandible (Figure S2). Landmarks were exported from

MorphoDig as .stv files and imported into the R package geomorph v.4.0.3 for processing

(Adams et al. 2021, Baken et al. 2021). Missing landmarks (11 of 4288 cranial and three of 1220

mandible landmarks) were imputed, a generalized Procrustes analysis (GPA) superimposition

was performed, and shape coordinates were subjected to a principal components analysis. We

retained the centroid size (an estimate of total size) and the scores from the first 36 principal

components of the cranium and 20 axes of the mandible, each representing >95% of the shape

variation of the element.

Stable isotope data collection- Approximately 1-2 grams of hairs were plucked from the rump of

286 dry museum specimens collected on six focal surveys. Isotopic values can vary regionally

(Fry 2006); therefore, we collected hair samples from multiple individuals of each species from

each locality (mean = 4.6 specimens/species/locality, range = 1-14). Nitrogen stable isotope

values (δ^{15}N) act as a proxy for consumer trophic position as they generally increase by 3–5%

per trophic level (DeNiro & Epstein 1981). Carbon isotope values (δ^{13}C) generally exhibit little

to no change with trophic position and are commonly used as proxies of consumer's basal carbon

resource use (e.g. use of differing primary production energy pathways; DeNiro & Epstein

1978). Combined, these two metrics are commonly used to quantify the "isotopic niche" of

consumers, which can act as a useful proxy of species realized dietary niche (Newsome et al.

2007, Ben-David & Flaherty 2012). Stable isotope values are reported in delta notation in per mil

units. Samples were processed at the Stable Isotope Ecology Laboratory, Department of

Oceanography & Coastal Sciences, Louisiana State University. Details on sample cleaning,

processing, and analysis are in the Supporting Methods.

Locomotor mode data collection- Nations et al. (2019, 2021) classified Sulawesi murines into

one of four discrete locomotor modes: Arboreal, (climbing is integral to survival); General

(navigates a variety of substrates and habitat strata); Terrestrial (on the ground surface); and

Amphibious, (dependent on aquatic habitats for foraging). Nations et al. (2021) showed that

locomotor transitions in the evolutionary history of murines occurred in a set order (Arboreal ↔

General ↔ Terrestrial ↔ Amphibious) and therefore we treat these modes as ordinal categories.

Combined traits – We combined our 11 continuous traits into five multivariate traits that

represent distinct niche dimensions: (1) head shape (cranium shape PC 1-36 and mandible shape

PC 1-20); (2) isotopic niche space (δ^{15}N and δ^{13}C stable isotope values); (3) body proportions

(head-body length, relative tail length, relative hind-foot length, and relative ear length); (4) body

size (log(mass) and head-body length); and (5) total morphological shape (all nine morphological

traits). We did this by combining the predicted trait values for each community (see below).

Estimating species' trait values- The sample size of morphological and isotopic data varied

between species and mountain communities, and in some cases was limited to one individual,

such as with the amphibious *Waiomys mammasae*, a species known from a single specimen

(Rowe et al. 2014). To mitigate uneven sampling, we estimated a probability distribution of

species trait values using partial-pooling in a multilevel Bayesian model. Unlike complete

pooling (one global mean estimated for all combined samples) or no-pooling (one mean

estimated per species, independent of all others), partial-pooling estimates a mean for each

species as well as the variance among species, which serves as an adaptive prior that is common

to all the species' means (McElreath 2020). The Bayesian partial-pooling modeling allowed us to

incorporate intraspecific variation in trait value predictions while avoiding point estimates such

as averages, which discard valuable information. This approach prevents unbalanced estimates

by using the trait-variance probability estimates from well-sampled species to inform variance

estimates of species with fewer samples (Gelman & Hill 2006, McElreath 2020). All analyses

were conducted in the probabilistic programming language Stan (Carpenter et al. 2017) within

the R library brms v. 2.17.0 (Bürkner 2018). Subsequent data processing and figuring relied on

the R libraries tidyverse v. 1.3.1 (Wickham et al. 2020), furrr v. 0.3.0 (Vaughan 2021), and

tidybayes v. 3.0.2 (Kay 2020). All data, scripts, and output files are available in Zenodo

Repository (to be added prior to publication) and on GitHub (to be added).

For each continuous trait (Table S1), we used the trait value as the response variable, and

used species as a group-level predictor. All traits were scaled to unity prior to analyses. We used

the student-*t* distribution to describe the response variable to minimize the influence of rare,

extreme observations (a.k.a. 'robust regression', Kruschke 2013, McElreath 2020). Each model

included four chains with 4000 iterations of warm-up and 1000 sampling iterations. Posterior

predictions from the four chains were combined, resulting in 4000 samples per trait per species.

To mitigate the potential geographic signal in stable isotope values (Fry 2006), and facilitate

randomization (see below), the posterior estimates of the δ^{15}N and δ^{13}C stable isotope values

were estimated with models that included an additional 'community' group-level effect, but

otherwise were identical to the models described above (Supporting Methods). These stable

isotope models generated posterior values of species' isotopic measurements conditioned on the

locality, enabling comparisons among communities and randomized sampling for null models

(Table S1, Figure S3). Full details of the model, prior, and chain-estimation are described in the

Supporting Methods.

Estimating functional space volume – We defined functional space volume as the total volume of

n-dimensional trait space occupied by species in a community, and for consistency and clarity

we use the term 'functional space volume' for all individual and multivariate traits, regardless of

dimension. We used the sum of the variance of each trait value to estimate the volume of

community functional space, which is less sensitive to outliers and outperforms other metrics

such as ellipse volumes, convex hull volumes, and hypervolumes, especially for functional

spaces with many axes (Li et al. 2017, Guillerme et al. 2020). We estimated a posterior

distribution of the variance of each trait for each community by first grouping the species' trait

value estimates by community. Then, for each of the 4000 posterior draws (one draw ranging

between 7 and 23 values, depending on the richness of the community), we estimated the

variance of the trait, resulting in a distribution of 4000 variance values for each trait for each

community. We estimated the combined multivariate trait space variance by summing the

variances of each trait, then dividing by the number of traits. We estimated the 89% probability

values for the variance of each individual and combined trait space. We estimated the size of

locomotor-mode space by calculating the variance of each community.

Estimating functional space density- We estimated the density of species within functional trait

space with the mean nearest neighbor (NN) metric (Guillerme et al. 2020). Our methods follow

the estimates of functional space volume above. First, we grouped species' trait values by

community, then estimated the individual trait density for each posterior draw using the NN

metric in the disparity() function of the R library dispRity (Guillerme 2018), resulting in a

distribution of 4000 NN values for each trait and combined trait space. To estimate locomotor

mode density, we dummy coded locomotor mode into four binary columns (one per mode) and

performed a redundancy analysis (Legendre & Legendre 2012) with the rda() function in the

vegan package (Oksanen et al. 2019). We then extracted the PC scores for each species, grouped

the species by community, and estimated the NN value for each community as with the

continuous traits above, in this case generating only a single NN value per community rather

than a distribution.

Null models of functional space volume and density- We used null models to determine the

difference between the functional space of our nine communities and randomly assembled

communities. For each community we created 1000 randomized null communities of $n_{community}$

species for each of the nine localities using the independent swap algorithm (Gotelli 2001). We

then calculated both volume and density as above for each individual (n=12) and combined

(n=5) trait for each of the 1000 randomized community samples. Standardized effect size (SES)

was calculated as:

$$SES = \frac{Trait_{observed} - Mean(Trait_{Random})}{SD(Trait_{Random})},$$

where $Trait_{observed}$ is the vector of 4000 samples of the density or volume of the given trait,

$Mean(Trait_{Random})$ and $SD(Trait_{Random})$ are the mean and standard deviation of the density

or volume values of the given trait from the 1000 random species assemblies. Positive SES

values indicate greater than random functional space volume or lower than random functional

space density (overdispersion), meaning that species in that community are occupying functional

space outside the range of other communities and/or their niches are farther apart in functional

space (niche expansion). Negative SES values indicate lower than random functional space

volume and greater than random functional space density (underdispersion or clustering),

meaning that species in that community are occupying less ecological space than other

communities and/or their niches are closer together in functional space (niche packing; Oliveria

et al. 2020).

Effect of species richness on community functional space volume and density- If niche expansion

is the primary mode of resource partitioning in Sulawesi murine communities, then we expected

that functional space volume will increase with increasing species richness, and that functional

space density will remain stable as richness increases. Whereas if niche packing is occurring, we

expected that functional space density will increase with species richness, and that there will be

no effect of richness on functional space volume. To quantify the effect size of richness

($\beta_{richness}$) on functional space volume and density, we used Bayesian linear regression models

that include the measurement error of the predictor variable to estimate the effect size of species

richness on volume or density. We used species richness as the response variable, and the mean

and standard error of the trait estimates for each community as the predictor. Measurement error

models contain vastly more information per observation than a single point observation and

provide robust estimates despite the small number of sampled communities (Bürkner 2018,

McElreath 2020). We also calculated the Bayesian R^2 value for each regression (Gelman et al.

2019). The models, priors, and chain estimations are detailed in the Supporting Methods.

Phylogenetic diversity as a path to trait disparity- The ecological space occupied by a

community may depend on which lineages or clades are present (Webb 2000). Trait disparity

appears to vary among clades of Sulawesi murines, and if so, then the volume and densities of

functional spaces may be more influenced by phylogenetic diversity than ecological factors. To

determine how phylogenetic diversity mediates functional space occupancy in the nine murine

communities, we estimated the phylogenetic diversity (PD) of each community using Faith's

metric of branch length (Faith 1996). We removed all but the Sulawesi species (n = 35) from a

time-calibrated phylogenetic hypothesis of Murinae (Nations et al. 2021). For the analyses

described above, we had trait data for four species that are not included in this phylogeny:

Maxomys wattsi, Rattus bontanus, *Rattus mollicomulus*, and *Rattus marmosurus*. We manually

added these four species into the tree using the R package phytools (Revell 2012). Details are

found in the Supplemental Methods. We estimated Faith's metric of PD and the SES value of PD

using the R package picante (Kembell et al. 2010). We used the independent swap algorithm to

sample 1000 random communities and used linear modeling in brms to estimate the effect size of

species richness on SES PD. *Haeromys minahassae* is the only representative of its genus on

Sulawesi, and, due to its large phylogenetic distance from other Sulawesi murines, may have an

oversized impact on estimates of community PD. Therefore, we repeated the estimates of SES

PD and effect size above without *Haeromys minahassae*. To quantify the trait disparity within

and among Sulawesi murine clades, we grouped the posterior distributions of the species'

predicted trait values by clade, then estimated the variance of each of the 12 univariate traits. We

plotted the trait value variance for each community along with the species' predicted trait values

(Figure S4)

Figure 2: The effect of species richness on community functional space volume (a, b) and density (c, d). X-axes show the $\beta_{richness}$ estimates (the regression slope) for each trait space (y-axis) on the left and the Bayesian R^2 for each $\beta_{richness}$ estimate on the right. Colored point intervals show 89% probability of $\beta_{richness}$ estimates (effect size), with color varying by $\beta_{richness}$ value. Black point intervals show 50% posterior estimates of Bayesian R^2 for each trait space. Four of the five multivariate traits (**a**) and seven of the 12 individual traits (**b**), show an increase in functional space volume (trait variance) with greater richness (i.e., positive $\beta_{richness}$). Phylogenetic diversity also increases with species richness (**b**). All multivariate traits (**c**) and individual traits (**d**) show a stable or, surprisingly, decreasing functional space density (increased NN distance) with greater species richness.

Results

Models of trait values- All parameters in the Bayesian multilevel model estimates of trait-value

distributions and measurement-error estimates of the species richness effect on trait values had

ESS > 1000 and a Gelman-Rubin diagnostic $\hat{R} \leq 1.01$, indicating convergence. Raw volume and

density values differed from the volume and density of SES. To avoid the confounding influence

of species richness in volume and density estimates we report and discuss only the SES values

(Swenson 2014).

Effects of Species Richness on Functional Space Volume and Density- Functional space volume

was positively correlated with species richness in seven of the 12 traits (locomotor mode, tail

length, head-body length, cranium shape, mass, δ^{15}N, and hind foot length), as shown by the

positive $\beta_{richness}$ slope values from the Bayesian linear models (Figures 3, Table S3) and the

greater variance with higher community richness (Figure 3). The remaining five traits (ear

length, cranium size, dentary size, dentary shape, and δ^{13}C) did not substantially increase in

volume with increased species richness. Among the multivariate traits, the SES values of total

morphological shape increased substantially with species richness while head shape and isotopic

niche space show a moderate probability of increasing with species richness (Figures 3, 4, Table

S2, S3).

391   Linear regression showed that species richness had little to no effect on functional space

density (NN SES values) for 11 of the 12 individual traits and three of five multivariate traits

(Figures 3, S4, Table S2), consistent with niche expansion. The exceptions were a positive

$\beta_{richness}$ slope (i.e., decreased density with richness) for locomotor mode, total morphological

shape, and external proportions, suggesting an extreme overdispersion of morphology and

locomotor diversity in the richest communities.

Phylogenetic Diversity and Functional space- Phylogenetic diversity (PD) increased with species

richness (Figures 3b, 4d, Table S2). Including *Haeromys minahassae* in the estimates of PD

changed the PD SES values of the individual communities but had minimal impact on linear

regressions (Table S4). Results including *H. minahassae* had a slope ($\beta_{richness}$) of 0.143 (0.03,

0.252) while excluding *H. minahassae* generated a slope of 0.169 (89% C.I. of 0.041, 0.295), and

we therefore present the results that include *H. minahassae*. Katopasa, a community of 12

murines on Sulawesi's Eastern Peninsula, has the lowest PD SES value followed by

Bawakaraeng on the Southwestern Peninsula (seven species). Gandang Dewata, the richest

community (23 species), has the highest PD SES value (Figure 3a, Table S2). As suspected,

functional space volumes vary between clades, though not consistently (Figure S5). For example,

the Echiothrix clade has the highest variance in cranial shape, dentary shape, and dentary size but

low trait variance for hind-foot length and $\delta^{15}N$. In contrast, the Maxomys clade has low variance

for all trait values except tail length and $\delta^{15}N$ (Figure S5). This results in different densities of

functional space occupation among traits, where some trait volumes, such as cranial space, are

strongly influenced by phylogeny, while others, such as isotopic niche space, have high and low

values distributed among clades (Figure 4).

Figure 3: Estimated functional volume SES plotted against species richness: **a**) Multivariate trait volumes, **b**) Skull traits, **c**) External measurements, and **d**) Isotopic measures, locomotor mode, and phylogenetic diversity. Points represent the mean SES values and error bars indicate the 89% credible intervals. X-axis tick marks show species richness for each community. Values equal to zero are consistent with null expectations, positive values indicate overdispersion, and negative values show underdispersion (trait clustering). Dotted lines depict the 89% interval of the null distribution. The plot of trait densities is shown in the supporting information (Figure S4).

Figure 4: The mode of functional space filling varies among traits. The bivariate plots on the left depict the 2D functional spaces. Each black shape is the mean value of a particular species in each clade (shown in legend). The convex hull colors match the color of the community functional volume SES distributions in the right-hand columns. The thick convex hull lines are mean values, and 500 random samples from the posterior of each trait are shown in thin hull lines. The number of species in each community and each clade used in this study is shown next to the SES distributions in parentheses. All values were scaled to z-scores prior to analyses. **a**) Cranial morphospace values are mostly clustered tightly near the mean of each axis, apart from Echiothrix species and the single *Haeromys* species. Intraspecific variance is also relatively low on these axes. **b**) Intraspecific variance is high in isotopic niche space. High and low δ¹⁵N values are distributed among clades, but that is not the case for δ¹³C values. **c**) Large values of hind foot length belong to members of two clades, while large and small tail length values are dispersed among all six clades, reducing the influence of phylogenetic diversity on tail length disparity.

Discussion

Variation in species richness among communities of closely related species that inhabit

similar environments provides a unique window to explore how interspecific competition may

affect community functional richness absent confounding factors like environmental variation

(Roughgarden 1976, Swenson et al. 2016). Unfortunately, said communities are rare in nature

(Swenson et al. 2016, Van de Perre et al. 2020). Here we used a large dataset of ecological and

morphological traits along with a novel probabilistic approach to estimate the changes in

community niche occupancy across a richness gradient on an oceanic island. The functional

space of most traits increased with greater species richness among the nine murine communities

studied while there was no change in functional space density, consistent with limiting similarity.

Locomotor mode disparity has the strongest positive relationship with species richness, but the

functional volumes of skull and body shape (indicators of diet, locomotion, and microhabitat use

in murines and vertebrates in general; Peters 1986, Losos 2009, Martinez et al. 2018, Nations et

al. 2021), also strongly increase with species richness (Figure 2). Measurements of overall body

size, cranial size, dentary shape and size, ear length, and $\delta^{13}C$ values all demonstrate either a

weak signal of niche packing, or no signal across the species richness gradient. While the volume

of many functional spaces increases with richness, functional space density shows little

correlation with species richness for most traits, and it surprisingly decreases with increased

richness for external morphology and locomotor mode. These results, along with a general

underdispersion of trait values in low-richness communities and overdispersion in high-richness

communities (Table S2, Figures 4 & S4), suggest that species in rich communities mitigate

competitive interactions by occupying underused niche space. We also found that increasing

phylogenetic diversity is a means of increasing functional space occupation. Combined, these

results point to species interactions as a mechanism for both phenotypic and phylogenetic

overdispersion (Webb et al. 2002, Cavender-Bares et al. 2004, Li et al. 2017) and suggest that,

given adequate resources, structural complexity, and evolutionary time, species can adapt to

unique regions of ecospace, often far from the average trait value, which minimizes niche

overlap and cultivates exceptional richness.

Whether in a single desert valley (Brown 1975), or across continental (Maestri &

Patterson 2016, Kohli et al. 2022) and global latitudinal gradients (Karr & James 1975, Pellissier

et al. 2018), structurally complex habitats are thought to foster higher species diversity. High

plant diversity creates a more complex, vertically structured habitat matrix for other plant and

animal species to occupy and has long been tied to higher animal richness (Hutchinson 1959,

MacArthur & MacArthur 1961, Scheffers et al. 2013, Oliveira & Scheffers 2019). Our estimates

of locomotor-mode occupancy clearly demonstrate that vertical habitat partitioning is critical to

maintaining high species richness in Sulawesi murines (Figure 2, Table S2). Strikingly, the

density of locomotor trait space decreases along the richness gradient, indicating a very high

level of trait overdispersion (Figures 3c & S4). Arboreal, Terrestrial, and Amphibious locomotor

modes each provide access to different microhabitats that contain similar resources. Among the

23 species found on Mt. Gandang Dewata, the amphibious *Waiomys mamasae*, the terrestrial

Paucidentomys vermidax, and the arboreal *Sommeromys macrorhinos*, all consume invertebrates

and have similar δ^{15}N values, yet they are unlikely to compete for resources due to their distinct

microhabitat use, a pattern observed in other insular communities of closely related vertebrates

(e.g., Jamaican *Anolis*; Shoener 1974). It's worth noting that when the lone amphibious species

Waiomys mamasae, only known from Gandang Dewata, is removed from the data, the locomotor

variance of Gandang Dewata remains the highest among the communities. Morphological

measurements are often used to infer locomotor mode in a variety of eco-evolutionary contexts

(Ricklefs & Travis 1980, Samuels and Van Valkenburgh 2009, Pianka et al. 2017, Verde-

Arregoitia et al. 2019), and our estimates of tail-length variance, a trait correlated with

locomotion in murines (Nations et al. 2021), increases with richness at nearly the same rate as

locomotor mode (Figure 2). Importantly, we demonstrate that locomotor mode rankings

themselves, especially if shown to be structured in an ordinal manner (Nations et al. 2021), can

be used directly to estimate community locomotor variance, and provide a simple way to test

hypotheses of niche structure and variance across wide taxonomic levels and geographic scales

where continuous trait data are lacking.

Combining many traits into one multivariate measure of functional diversity is a common

approach in evolutionary ecology. Ordination techniques were especially promoted to overcome

pitfalls from early community ecology studies that used few, largely subjective measures of

resource use (Ricklefs & Travis 1980). However, merging traits into multivariate axes masks

trait-specific processes related to functional space (Spasojevic & Suding 2012, Astor et al. 2014).

Indeed, our trait volume and density estimates reveal distinct patterns between individual and

combined traits. For example, isotopic niche space, a combined signal of δ^{15}N and δ^{13}C values

commonly used in terrestrial ecological studies, exhibits an equivocal signal (Figure 2b), but

individual isotopic values reveal that the packing signal originates from the static δ^{13}C value

along the species richness gradient. The δ^{15}N value, a signal of trophic level, expands along the

richness gradient, a result that would be overlooked in multivariate analyses. In contrast to δ^{15}N,

dentary shape exhibits an equivocal signal, despite its presumed relationship to dietary niche

(Maestri et al. 2016, Kohli et al. 2019). These opposing patterns are only illuminated by

analyzing individual traits (Spasojevic & Suding 2012). If we performed this study using only

cranial size and dentary shape, which are thought to capture the important axes of size and diet in

murine rodents (Rowsey et al. 2019, 2020) and mammals in general (Prevosti et al. 2012,

Grossnickle 2020), our results would suggest that functional volume does not increase with

species richness.

The disparity of traits within a clade mediates the impact of phylogenetic diversity on

community functional space volume. If trait values are phylogenetically clustered, then

increasing phylogenetic diversity is necessary for community niche expansion to occur. But if

trait values are phylogenetically overdispersed, niche packing, expansion, or, as we found, both

could result from increased phylogenetic diversity. Phylogenetic niche conservatism is of

particular interest in communities that are assembled through a mix of colonization and in situ

speciation, such as Sulawesi murines, Caribbean anoles, or Hawaiian spiders (Gillespie 2004,

Losos 2009, Rowe et al. 2019), as the distribution of traits among clades likely depends on their

biogeographic history. Niche divergence has been hypothesized to overcome niche conservatism

in communities with an extended history of coevolution, such as those on old, oceanic islands,

likely diminishing the signal of phylogenetic trait clustering (Losos et al. 2003). We find that

niche divergence and conservatism may occur simultaneously on different functional traits. For

example, the elongate, highly distinct skulls of some species in the Echiothrix clade, the

descendants of the first murine colonists on Sulawesi, set them apart from other clades in skull

shape and, despite their relatively low abundance, these species overcontribute to community

cranial and dentary shape volumes (Figures 3b, 4b, S5). Unlike cranial shape however, the

Echiothrix clade occupies a very constrained portion of δ^{15}N trophic space. Additionally, trophic

level estimates from δ^{15}N values are notably dispersed among clades (Figures 4c, S5), and high

phylogenetic diversity is not necessary for high isotopic niche space estimates. Are there innate

differences in these traits that could lead to opposing patterns of niche conservatism?

The evolutionary lability of ecologically important traits determines the rate of

convergence and divergence possible within a given time frame and can directly influence

dispersion of trait values among species (Cavender-Bares et al. 2004). The traits with the highest

within-clade variance — $\delta^{15}N$, tail length, body size — are all thought to be evolutionarily labile.

Changes in tail length, body size, and intestinal tract morphology can occur on brief evolutionary

time scales (Powell & King 1997, Naya et al. 2008, Kingsley et al. 2017, 2021), whereas

morphological changes in cranial shape, such as substantial rostral elongation and the reduction

of molar grinding area, may take much longer. Indeed, insular species are known to have rapidly

expanded breadths of diet and labile morphological traits following colonization (Stuart et al.

2014) and subsequent speciation (Wilson 1959, 1961, Lister 1976, Millien 2006, Rowe et al.

2016a). But the extreme cranial morphologies of some Sulawesi murines, particularly those of

the oldest radiation on the island (Echiothrix clade), are likely the result of a long process of in-

situ evolution (Rowe et al. 2019). In other words, on shorter time scales, similarities in some

slowly evolving traits may lead to higher divergences in more labile traits, as evidenced in the

Maxomys and Bunomys clades (Figure S5). The opposite pattern may also occur, but only

following sufficient evolutionary time. We propose that the evolutionary lability of traits is a

determinant of trait value dispersion (Webb et al. 2002, Cavender-Bares et al. 2004), which

directly relates to our inferences of functional space occupancy in Sulawesi murine communities.

The theory of niche complementarity suggests that a pair of coexisting species that are similar in

one trait should diverge in another trait (Schoener 1974). Yet, in an isolated setting, the lability

of the traits in question necessarily determines the degree of complementarity possible within a

given time frame. Linking trait lability with niche complementarity in an island system has

important implications for the generation of functional and taxonomic diversity and may further

illuminate the process of niche filling in insular, adaptive radiations.

Conclusion

Here we provide evidence that limiting similarity in functional traits reflecting

locomotion and microhabitat use plays an important role in the assembly of discrete, montane

small mammal communities on an oceanic island. Our results contrast with recent studies that

recovered niche packing as key to increased species richness in continental tropical vertebrate

communities (Pigot et al. 2016, Peixoto et al. 201, Pellissier et al. 2018, Van de Perre 2020,

Dehling et al. 2022, Hughes et al. 2022). Furthermore, our results counter the predictions that

niche packing is expected to occur if resources remain constant among communities

(Roughgarden 1976). Yet, Roughgarden (1976) posited that the predicted relationship between

species richness and resources may be different on remote islands, as the distance from source

populations may affect both community richness and the functional trait values of the species

present. The organization of distinct montane communities on an oceanic island may be just such

an example. The regional murine species pool of Sulawesi is itself structured by idiosyncratic

immigration, diversification, and (presumably) extinction dynamics, and the resultant species'

functional traits. The species in these communities that occupy the edges of some functional

spaces, such as cranial shape (Figure 4), represent morphologies and ecologies found only on

Sulawesi or other large, oceanic islands. And the "imperfect isolation" (sensu Samonds et al

2013) of Sulawesi allows for other regional murine taxa, often represented by more "average"

murine phenotypes, to periodically colonize the island, adding more species into the center of

functional space (Figure 4). The presence of such disparate phenotypes and ecologies on

Sulawesi produces a larger functional space reserve than is available in most, if any, continental

rodent systems, nurturing both niche expansion and high species richness. Therefore, the

complex topography, isolation, abundant resources, and sequential colonization of Sulawesi

might lead to species assembly processes that are typical of other large, oceanic islands, but are

atypical of continental systems, "sky islands", geologically younger islands, or regions with less

abundant resources (deserts, high latitude habitats), but the presence of this general pattern

remains untested.

590   **Literature Cited**

Achmadi, A. S., K. C. Rowe, and J. A. Esselstyn. 2014. New records of two rarely encountered,

endemic rats (Rodentia: Muridae: Murinae) from Gunung Gandangdewata, West

Sulawesi province. Treubia 41:51-60.

Adams, D. C., M. Collyer, and A. Kaliontzopoulou. 2021. Geomorph: Software for geometric

morphometric analyses. R package version 4.0.3.

Astor, T., J. Strengbom, M. P. Berg, L. Lenoir, B. Marteinsdóttir, and J. Bengtsson. 2014.

Underdispersion and overdispersion of traits in terrestrial snail communities on islands.

Ecology and Evolution 4:2090-2102.

Baken, E. K., M. L. Collyer, A. Kaliontzopoulou, and D. C. Adams. 2021. Geomorph v4.0 and

gmShiny: enhanced analytics and a new graphical interface for a comprehensive

morphometric experience. Methods in Ecology and Evolution. 12:2355-2363.

Baltensperger, A. P., F. Huettmann, J. C. Hagelin, and J. M. Welker. 2015. Quantifying trophic

niche spaces of small mammals using stable isotopes (δ^{15}N and δ^{13}C) at two scales across

Alaska. Canadian Journal of Zoology 93:579-588.

Ben-David, M., and E. A. Flaherty. 2012. Stable isotopes in mammalian research: a beginner's

guide. Journal of Mammalogy 93:312-328.

Brown, J. H., and G. A. Lieberman. 1973. Resource utilization and coexistence of seed□eating

desert rodents in sand dune habitats. Ecology 54:788-797.

Brown, J. H. 1975. Geographical Ecology of Desert Rodents. Pages 315-341 *in* M. L. Cody and

J. M. Diamond, editors. Ecology and Evolution of Communities. The Belknap Press,

Cambridge.

Bürkner, P.-C. 2018. Advanced Bayesian Multilevel Modeling with the R Package brms. The R

Journal 10:395–411.

Carpenter, B., A. Gelman, M. D. Hoffman, D. Lee, B. Goodrich, M. Betancourt, M. Brubaker, J.

Guo, P. Li, and A. Riddell. 2017. Stan: A Probabilistic Programming Language. Journal

of Statistical Software 76.

Cavender-Bares, J., D. D. Ackerly, D. A. Baum, and F. A. Bazzaz. 2004. Phylogenetic

Overdispersion in Floridian Oak Communities. The American Naturalist 163:823-843.

Dehling, D. M., G. V. Dalla Riva, M. C. Hutchinson, and D. B. Stouffer. 2022. Niche packing

and local coexistence in a megadiverse guild of frugivorous birds are mediated by fruit

dependence and shifts in interaction frequencies. The American Naturalist 199:855–868.

DeNiro, M. J., and S. Epstein. 1978. Influence of diet on the distribution of carbon isotopes in
animals. Geochimica et Cosmochimica Acta 42:495-506.

DeNiro, M. J., and S. Epstein. 1981. Influence of diet on the distribution of nitrogen isotopes in
animals. Journal of Archeological Science 45:341-351.

Des Roches, S., D. M. Post, N. E. Turley, J. K. Bailey, A. P. Hendry, M. T. Kinnison, J. A.
Schweitzer, and E. P. Palkovacs. 2018. The ecological importance of intraspecific
variation. Nature Ecology & Evolution 2:57-64.

Diamond, J. M. 1975. Assembly of Species Communities. Pages 342-444 *in* M. L. Cody and J.
M. Diamond, editors. Ecology and Evolution of Communities. The Belknap Press,
Cambridge.

Dorey, J. B., S. V. Groom, E. H. Freedman, C. S. Matthews, O. K. Davies, E. J. Deans, C.
Rebola, M. I. Stevens, M. S. Lee, and M. P. Schwarz. 2020. Radiation of tropical island
bees and the role of phylogenetic niche conservatism as an important driver of
biodiversity. Proceedings of the Royal Society B 287:20200045.

Esselstyn, J. A., A. S. Achmadi, and K. C. Rowe. 2012. Evolutionary novelty in a rat with no
molars. Biology Letters 8:990-993.

Esselstyn, J. A., A. S. Achmadi, H. Handika, and K. C. Rowe. 2015. A hog-nosed shrew rat
(Rodentia: Muridae) from Sulawesi Island, Indonesia. Journal of Mammalogy 96:895-
907.

Esselstyn, J. A., A. S. Achmadi, H. Handika, M. T. Swanson, T. C. Giarla, and K. C. Rowe.
2021. Fourteen New, Endemic Species of Shrew (Genus *Crocidura)* from Sulawesi
Reveal a Spectacular Island Radiation. Bulletin of the American Museum of Natural
History 454:1-108.

Faith, D. P. 1996. Conservation priorities and phylogenetic pattern. Conservation Biology

10:1286-1289.

Fry, B. 2006. Stable isotope ecology. Springer, New York.

Galetti, M., R. R. Rodarte, C. L. Neves, M. Moreira, and R. Costa-Pereira. 2016. Trophic Niche

Differentiation in Rodents and Marsupials Revealed by Stable Isotopes. PLOS One

11:e0152494.

Gelman A, Hill J. 2006. Data Analysis Using Regression and Multilevel/Hierarchical Models.

Cambridge: Cambridge University Press.

Gelman, A., B. Goodrich, J. Gabry, and A. Vehtari. 2019. R-squared for Bayesian regression

models. The American Statistician 73:307-309.

Gillespie, R. 2004. Community assembly through adaptive radiation in Hawaiian spiders.

Science 303:356-359.

Gotelli, N. J. 2001. Research frontiers in null model analysis. Global Ecology and Biogeography

10:337-343.

Grossnickle, D. M. 2020. Feeding ecology has a stronger evolutionary influence on functional

morphology than on body mass in mammals. Evolution 74:610-628.

Guillerme T. 2018. dispRity: A modular R package for measuring disparity. Methods in Ecology

and Evolution 9:1755-1763.

Guillerme, T., M. N. Puttick, A. E. Marcy, and V. Weisbecker. 2020. Shifting spaces: Which

disparity or dissimilarity measurement best summarize occupancy in multidimensional

spaces? Ecology and Evolution 10:7261-7275.

Handika, H., A. S. Achmadi, J. A. Esselstyn, and K. C. Rowe. 2021. Molecular and

morphological systematics of the Bunomys division (Rodentia: Muridae), an endemic

radiation on Sulawesi. Zoologica Scripta 50:141-154.

Heaney, L. R. 2001. Small mammal diversity along elevational gradients in the Philippines: an

assessment of patterns and hypotheses. Global Ecology and Biogeography 10:15–39.

Hiller, A. E., M. S. Koo, K. R. Goodman, K. L. Shaw, P. M. O'Grady, and R. G. Gillespie. 2019.

Niche conservatism predominates in adaptive radiation: comparing the diversification of

Hawaiian arthropods using ecological niche modelling. Biological Journal of the Linnean

Society 127:479-492.

Hughes, E. C., D. P. Edwards, J. A. Bright, E. J. Capp, C. R. Cooney, Z. K. Varley, and G. H.

Thomas. 2022. Global biogeographic patterns of avian morphological diversity. Ecology

Letters 25:598-610.

Hutchinson, G. E. 1957. Concluding remarks. Cold Spring Harbor Symposia on Quantitative

Biology 22:415-427.

Hutchinson, G. E. 1959. Homage to Santa Rosalia or Why Are There So Many Kinds of

Animals? The American Naturalist 93:145-159.

Imhoff, M. L., L. Bounoua, T. Ricketts, C. Loucks, R. Harriss, and W. T. Lawrence. 2004.

Global patterns in human consumption of net primary production. Nature 429:870-873.

Karr, J. R., and F. C. James. 1975. Ecomorphological Configurations and Convergent Evolution.

Pages 258-291 *in* M. L. Cody and J. M. Diamond, editors. Ecology and Evolution of

Communities. The Belknap Press, Cambridge.

Kay, M. 2020. Tidybayes: Tidy data and geoms for Bayesian models. R package version 2.1.1.

Kembel, S. W., P. D. Cowan, W. K. Helmus, W. K. Cornwell, H. Morlon, D. D. Ackerly, S. P.

Blomberg, and C. O. Webb. 2010. Picante: R tools for integrating phylogenies and

ecology. Bioinformatics 26:1463-1464.

Kingsley, E. P., K. M. Kozak, S. P. Pfeifer, D. S. Yang, and H. E. Hoekstra. 2017. The ultimate

and proximate mechanisms driving the evolution of long tails in forest deer mice.

Evolution 71:261-273.

Kingsley, E. P., E. R. Hager, J.-M. Lassance, K. M. Turner, O. S. Harringmeyer, C. Kirby, B. I.

Neugeboren, and H. E. Hoekstra. 2024. Adaptive tail-length evolution in deer mice is

associated with differential Hoxd13 expression in early development. Nature Ecology &

Evolution:1–15.

Kohli, B. A., and M. A. Jarzyna. 2021. Pitfalls of ignoring trait resolution when drawing

conclusions about ecological processes. Global Ecology and Biogeography 30:1139–

1152.

Kohli, B. A., M. J. Miyajima, and M. A. Jarzyna. 2022. Elevational diversity patterns of rodents

differ between wet and arid mountains. Global Ecology and Biogeography 31:1726-1740.

Kohli, B. A., and R. J. Rowe. 2019. Beyond guilds: the promise of continuous traits for

mammalian functional diversity. Journal of Mammalogy 100:285-298.

Kohli, B. A., R. D. Stevens, E. A. Rickart, and R. J. Rowe. 2021. Mammals on mountainsides

revisited: Trait☐based tests of assembly reveal the importance of abiotic filters. Journal

of Biogeography 48:1606-1621.

Kruschke, J. K. 2013. Bayesian estimation supersedes the t test. Journal of Experimental

Psychology: General 142:573.

Latham, R. E., and R. E. Ricklefs. 1993. Global Patterns of Tree Species Richness in Moist

Forests: Energy-Diversity Theory Does Not Account for Variation in Species Richness.

Oikos 67:325–333.

Lebrun, R. 2018. MorphoDig, an open-source 3D freeware dedicated to biology. IPC5, Paris,

France.

Legendre, P., and L. Legendre. 2012. Numerical ecology. Elsevier, Amsterdam.

Li, Y., B. Shipley, J. N. Price, V. de L. Dantas, R. Tamme, M. Westoby, A. Siefert, B. S.

Schamp, M. J. Spasojevic, V. Jung, D. C. Laughlin, S. J. Richardson, Y. L. Bagousse-

Pinguet, C. Schöb, A. Gazol, H. C. Prentice, N. Gross, J. Overton, M. V. Cianciaruso, F.

Louault, C. Kamiyama, T. Nakashizuka, K. Hikosaka, T. Sasaki, M. Katabuchi, C.

Frenette Dussault, S. Gaucherand, N. Chen, M. Vandewalle, and M. A. Batalha. 2018.

Habitat filtering determines the functional niche occupancy of plant communities

worldwide. Journal of Ecology 106:1001–1009.

Lister, B. C. 1976. The nature of niche expansion in West Indian Anolis lizards I: ecological

consequences of reduced competition. Evolution 30:659-676.

Losos, J. B. 2009. Lizards in an evolutionary tree: ecology and adaptive radiation of anoles.

University of California Press, Berkeley.

Losos, J. B., M. Leal, R. E. Glor, and K. D. Queiroz. 2003. Niche lability in the evolution of a

Caribbean lizard community. Nature 424:542–545.

Losos, J. B., and R. E. Ricklefs. 2009. Adaptation and diversification on islands. Nature

457:830-836.

M'Closkey, R. T. 1978. Niche separation and assembly in four species of Sonoran Desert

rodents. The American Naturalist 112:683-694.

MacArthur, R. H. 1965. Patterns of species diversity. Biological Reviews 40:510-533.

MacArthur, R. H. 1970. Species packing and competitive equilibrium for many species.

Theoretical Population Biology 1:1-11.

MacArthur, R., and R. Levins. 1967. The limiting similarity, convergence, and divergence of

coexisting species. The American Naturalist 101:377-385.

MacArthur, R. H., and J. W. MacArthur. 1961. On bird species diversity. Ecology 42:594-598.

Maestri, R., B. Patterson, R. Fornel, L. Monteiro, and T. De Freitas. 2016. Diet, bite force and

skull morphology in the generalist rodent morphotype. Journal of Evolutionary Biology

29:2191-2204.

Maestri, R., and B. D. Patterson. 2016. Patterns of species richness and turnover for the South

American rodent fauna. PLoS One 11:e0151895.

Martinez, Q., R. Lebrun, A. S. Achmadi, J. A. Esselstyn, A. R. Evans, L. R. Heaney, R. P.

Miguez, K. C. Rowe, and P.-H. Fabre. 2018. Convergent evolution of an extreme dietary

specialization, the olfactory system of worm-eating rodents. Scientific Reports 8:17806.

May, R. M., and R. H. Mac Arthur. 1972. Niche overlap as a function of environmental

variability. Proceedings of the National Academy of Sciences 69:1109-1113.

McElreath R. 2020 Statistical Rethinking v2. Taylor & Francis Group, Boca Raton.

Millien, V. 2006. Morphological Evolution Is Accelerated among Island Mammals. PLOS

Biology 4:e321.

Musser, G. G. 2014. A systematic review of Sulawesi Bunomys (Muridae, Murinae) with the

description of two new species. Bulletin of the American Museum of Natural History

392.

Nations, J. A., L. R. Heaney, T. C. Demos, A. S. Achmadi, K. C. Rowe, and J. A. Esselstyn.

2019. A simple skeletal measurement effectively predicts climbing behavior in a diverse

clade of small mammals. Biological Journal of the Linnean Society 128:323–336.

Nations, J. A., G. G. Mount, S. M. Morere, A. S. Achmadi, K. C. Rowe, and J. A. Esselstyn.

2021. Locomotory mode transitions alter phenotypic evolution and lineage diversification

in an ecologically rich clade of mammals. Evolution 75:376-393.

Naya, D. E., F. Bozinovic, and W. H. Karasov. 2008. Latitudinal trends in digestive flexibility:

testing the climatic variability hypothesis with data on the intestinal length of rodents.

The American Naturalist 172:122-134.

Newsome, S. D., C. Martinez del Rio, S. Bearhop, and D. L. Phillips. 2007. A niche for isotopic

ecology. Frontiers in Ecology and the Environment 5:429-436.

Oksanen, J, F. Guillaume Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R.

Minchin, R. B. O'Hara, G. L. Simpson, P. Solymos, M. Henry, H. Stevens, E. Szoecs and

H. Wagner (2019). Vegan: Community Ecology Package. R package version 2.5-6.

Oliveira, B. F., and B. R. Scheffers. 2019. Vertical stratification influences global patterns of

biodiversity. Ecography 42:249-249.

Oliveira, B. F., J. M. Flenniken, R. P. Guralnick, S. E. Williams, and B. R. Scheffers. 2020.

Historical environmental stability drives discordant niche filling dynamics across

phylogenetic scales. Journal of Biogeography 47:807-816.

Peixoto, F. P., P. H. P. Braga, and P. Mendes. 2018. A synthesis of ecological and evolutionary

determinants of bat diversity across spatial scales. BMC ecology 18:1–14.

Pellissier, V., J. Y. Barnagaud, W. D. Kissling, Ç. Şekercioğlu, and J. C. Svenning. 2018. Niche

packing and expansion account for species richness–productivity relationships in global

bird assemblages. Global Ecology and Biogeography 27:604-615.

Peters, R. H. 1986. The Ecological Implications of Body Size. Cambridge University Press,

Cambridge.

Pianka, E. R. 1974. Niche overlap and diffuse competition. Proceedings of the National

Academy of Sciences 71:2141-2145.

Pianka, E. R., L. J. Vitt, N. Pelegrin, D. B. Fitzgerald, and K. O. Winemiller. 2017. Toward a

Periodic Table of Niches, or Exploring the Lizard Niche Hypervolume. The American

Naturalist 190:601-616.

Pigot, A. L., C. H. Trisos, and J. A. Tobias. 2016. Functional traits reveal the expansion and

packing of ecological niche space underlying an elevational diversity gradient in

passerine birds. Proceedings of the Royal Society B: Biological Sciences 283:20152013.

Pinto, G., D. L. Mahler, L. J. Harmon, and J. B. Losos. 2008. Testing the island effect in

adaptive radiation: rates and patterns of morphological diversification in Caribbean and

mainland Anolis lizards. Proceedings of the Royal Society B: Biological Sciences

275:2749–2757.

Powell, R. A., and C. M. King. 1997. Variation in body size, sexual dimorphism and age-specific

survival in stoats, Mustela erminea (Mammalia: Carnivora), with fluctuating food

supplies. Biological Journal of the Linnean Society 62:165-194.

Prevosti, F. J., G. F. Turazzini, M. D. Ercoli, and E. Hingst-Zaher. 2012. Mandible shape in

marsupial and placental carnivorous mammals: a morphological comparative study using

geometric morphometrics. Zoological Journal of the Linnean Society 164:836-855.

Revell, L. J. 2012. Phytools: an R package for phylogenetic comparative biology (and other

things). Methods in Ecology and Evolution 3:217-223.

Ricklefs, R. E., and J. Travis. 1980. A Morphological Approach to the Study of Avian

Community Organization. The Auk 97:321-338.

Rowe, K. C., A. S. Achmadi, and J. A. Esselstyn. 2014. Convergent evolution of aquatic

foraging in a new genus and species (Rodentia: Muridae) from Sulawesi Island,

Indonesia. Zootaxa 3815:541-564.

Rowe K.C., A.S. Achmadi, and J.A. Esselstyn. 2016a. Repeated evolution of carnivory among

Indo-Australian rodents. Evolution 70:653-665.

Rowe, K. C., A. S. Achmadi, and J. A. Esselstyn. 2016b. A new genus and species of

omnivorous rodent (Muridae: Murinae) from Sulawesi, nested within a clade of endemic

carnivores. Journal of Mammalogy 97:978-991.

Rowe, K. C., A. S. Achmadi, P. H. Fabre, J. J. Schenk, S. J. Steppan, and J. A. Esselstyn. 2019.

Oceanic islands of Wallacea as a source for dispersal and diversification of murine

rodents. Journal of Biogeography 46:2752-2768.

Rowsey, D. M., L. R. Heaney, and S. A. Jansa. 2019. Tempo and mode of mandibular shape and

size evolution reveal mixed support for incumbency effects in two clades of

island☐endemic rodents (Muridae: Murinae). Evolution 73:1411-1427.

Rowsey, D. M., R. M. Keenan, and S. A. Jansa. 2020. Dietary morphology of two island-

endemic murid rodent clades is consistent with persistent, incumbent-imposed

competitive interactions. Proceedings of the Royal Society B 287:20192746.

Roughgarden, J. 1976. Resource partitioning among competing species–a coevolutionary

approach. Theoretical Population Biology 9:388-424.

Samonds, K. E., L. R. Godfrey, J. R. Ali, S. M. Goodman, M. Vences, M. R. Sutherland, M. T.

824   Irwin, and D. W. Krause. 2013. Imperfect Isolation: Factors and Filters Shaping

825   Madagascar's Extant Vertebrate Fauna. PLOS ONE 8:e62086.

826   Samuels, J. X. 2009. Cranial morphology and dietary habits of rodents. Zoological Journal of the

827   Linnean Society 156:864–888.

828   Samuels, J. X., and B. Van Valkenburgh. 2008. Skeletal indicators of locomotor adaptations in

829   living and extinct rodents. Journal of Morphology 269:1387-1411.

830   Scheffers, B. R., B. L. Phillips, W. F. Laurance, N. S. Sodhi, A. Diesmos, and S. E. Williams.

831   2013. Increasing arboreality with altitude: a novel biogeographic dimension. Proceedings

832   of the Royal Society B 280:20131581.

833   Schoener, T. W. 1974. Resource Partitioning in Ecological Communities: Research on how

834   similar species divide resources helps reveal the natural regulation of species diversity.

835   Science 185:27-39.

836   Sikes, R. S., and Animal Care and Use Committee of the American Society of Mammalogists.

837   2016. 2016 Guidelines of the American Society of Mammalogists for the use of wild

838   mammals in research and education. Journal of Mammalogy 97:663–688.

839   Spasojevic, M. J., and K. N. Suding. 2012. Inferring community assembly mechanisms from

840   functional diversity patterns: the importance of multiple assembly processes. Journal of

841   Ecology 100:652-661.

842   Stroud, J. T., and J. B. Losos. 2016. Ecological Opportunity and Adaptive Radiation. Annual

843   Review of Ecology, Evolution, and Systematics 47:507-532.

Stroud, J. T. 2021. Island species experience higher niche expansion and lower niche

conservatism during invasion. Proceedings of the National Academy of Sciences 118:

e2018949118.

Stuart, Y. E., T. S. Campbell, P. a. Hohenlohe, R. G. Reynolds, L. J. Revell, and J. B. Losos.

2014. Rapid evolution of a native species following invasion by a congener. Science

346:463-466.

Swenson, N. G. 2014. Functional and Phylogenetic Ecology in R. Springer, New York.

Swenson, N. G., and M. D. Weiser. 2014. On the packing and filling of functional space in

eastern North American tree assemblages. Ecography 37:1056–1062.

Swenson, N. G., M. D. Weiser, L. Mao, S. Normand, M. Á. Rodríguez, L. Lin, M. Cao, and J.-C.

Svenning. 2016. Constancy in functional space across a species richness anomaly. The

American Naturalist 187:E83-E92.

Van de Perre, F., M. R. Willig, S. J. Presley, I. J.-C. Mukinzi, M. S. Gambalemoke, H. Leirs, and

E. Verheyen. 2020. Functional volumes, niche packing and species richness:

biogeographic legacies in the Congo Basin. Royal Society Open Science 7:191582.

Vaughan, D, and M. Dancho 2021. furrr: Apply Mapping Functions in Parallel using Futures. R

package version 0.2.2.

Verde Arregoitia, L. D., D. O. Fisher, and M. Schweizer. 2017. Morphology captures diet and

locomotor types in rodents. Royal Society Open Science 4:160957.

Violle, C., B. J. Enquist, B. J. McGill, L. Jiang, C. H. Albert, C. Hulshof, V. Jung, and J.

Messier. 2012. The return of the variance: intraspecific variability in community ecology.

Trends in Ecology & Evolution 27:244-252.

Webb, C. O. 2000. Exploring the phylogenetic structure of ecological communities: an example

867   for rain forest trees. The American Naturalist 156:145-155.

868   Webb, C. O., D. D. Ackerly, M. A. McPeek, and M. J. Donoghue. 2002. Phylogenies and

869   community ecology. Annual Review of Ecology and Systematics 33:475-505.

870   Weiher, E., and P. A. Keddy. 1995. Assembly rules, null models, and trait dispersion: new

871   questions from old patterns. Oikos:159-164.

872   Wickham, H., M. Averick, J. Bryan, W. Chang, L. D. A. McGowan, R. François, G. Grolemund,

873   A. Hayes, L. Henry, and J. Hester. 2019. Welcome to the Tidyverse. Journal of Open

874   Source Software 4:1686.

875   Wilson, D. E., R. A. Mittermeier, and T. E. Lacher. 2019. Handbook of the Mammals of the

876   World. Lynx Editions, Barcelona.

877   Wilson, E. O. 1959. Adaptive shift and dispersal in a tropical ant fauna. Evolution 13:122-144.

878   Wilson, E. O. 1961. The Nature of the Taxon Cycle in the Melanesian Ant Fauna. The American

879   Naturalist 95:169-193.

880   Woodward, G., and A. G. Hildrew. 2002. Body size determinants of niche overlap and

881   intraguild predation within a complex food web. Journal of Animal Ecology 71:1063-

882   1074.

883   Xu, X., D. Dimitrov, N. Shrestha, C. Rahbek, and Z. Wang. 2019. A consistent species richness–

884   climate relationship for oaks across the Northern Hemisphere. Global Ecology and

885   Biogeography 28:1051–1066.

References Only Cited in the Online Supplement

Achmadi, A. S. 2010. Taxonomic Status of Spiny Rats (Maxomys Jentink, Rodentia) from Indonesia and Malaysia Based on Morphological Study. Treubia 37:49-82.

Fry, B. 2006. Stable isotope ecology. Springer, New York.

Hückstädt, L., J. Burns, P. Koch, B. McDonald, D. E. Crocker, and D. Costa. 2012. Diet of a specialist in a changing environment: the crabeater seal along the western Antarctic Peninsula. Marine Ecology Progress Series 455:287-301.

Kruschke, J. K. 2013. Bayesian estimation supersedes the t test. Journal of Experimental Psychology: General 142:573.

McElreath R. 2020 Statistical Rethinking v2. Taylor & Francis Group. Boca Raton.

Musser, G. G., and M. D. Carleton. 2005. Superfamily Muroidea. Pages 894–1531 in D. E. Wilson, Reeder D.M., editor. Mammal species of the world: a taxonomic and geographic reference. Johns Hopkins Univ. Press, Baltimore.

911