1 # The roles of isolation and interspecific interaction in generating the
2 # functional diversity of an insular mammal radiation
3
4

5 Jonathan A. Nations*[1,2], Brooks A. Kohli[3], Heru Handika[2], Anang S. Achmadi[4], Michael J.
6 Polito[5,6], Kevin C. Rowe[7], Jacob A. Esselstyn[2]
7
8
[1]Florida Museum of Natural History, University of Florida, Gainesville, Florida, 32611, United
States
[2]Museum of Natural Science and Department of Biological Sciences, Louisiana State University,
Baton Rouge, Louisiana, 70803, United States
[3]Biological Sciences, Ohio University, Athens, Ohio, 45701, United States
[4]Museum Zoologicum Bogoriense, Research Center for Ecology and Ethnobiology, National
Research and Innovation Agency (BRIN), Cibinong, Jawa Barat 16911, Indonesia
[5]Department of Oceanography and Coastal Sciences, Louisiana State University, Baton Rouge,
Louisiana, 70803, United States
[6]Department of Ocean Sciences, University of California, Santa Cruz, California, 95064, United
States
[7]Sciences Department, Museums Victoria, Melbourne, Victoria 3001, Australia
21
22 *Correspondence to be sent to: Florida Museum of Natural History, University of Florida,
23 Gainesville, Florida, 32611, United States; E-mail: jnations@floridamuseum.ufl.edu

Abstract

Communities that occupy similar environments but vary in the richness of closely related species

can illuminate how functional variation and species richness interact to fill ecological space in

the absence of abiotic filtering, though this has yet to be explored on an oceanic island where the

processes of community assembly may differ from continental settings. In discrete montane

communities on the island of Sulawesi, local murine rodent (rats and mice) richness ranges from

7 to 23 species. We measured 17 morphological, ecological, and isotopic traits, both individually

and grouped into 5 multivariate traits in 40 species, to test for the expansion or packing of

functional space among nine murine communities. We employed a novel probabilistic approach

for integrating intraspecific and community-level trait variance into functional richness. Trait-

specific and phylogenetic diversity patterns indicate dynamic community assembly due to

variable niche expansion and packing on multiple niche axes. Locomotion and covarying traits

such as tail length emerged as a fundamental axis of ecological variation, expanding functional

space and enabling the niche packing of other traits such as diet and body size. Though trait

divergence often explains functional diversity in island communities, we found that phylogenetic

diversity facilitates functional space expansion in some conserved traits such as cranial shape,

while more labile traits are overdispersed both within and between island clades, suggesting a

role of niche complementarity. Our results evoke interspecific interactions, differences in trait

lability, and the independent evolutionary trajectories of each of Sulawesi's 6 murine clades as

central to generating the exceptional functional diversity and species richness in this exceptional,

insular radiation.

Keywords: Community Niche Space, Functional Morphology, Stable Isotopes, Murinae,

Bayesian, Sulawesi, Indonesia

    Introduction

Whether at a local, continental, or global scale, species richness is not evenly distributed across

the landscape. This unevenness emerges from both environmental and resource heterogeneity

among communities, and from the interactions among the species within local communities.

Local species richness is often positively correlated with the complexity of habitat structure and

diversity of available resources (Tews et al. 2004). The observation that different localities with

similar habitat structure, resource availability, and historical access often contain ecologically

similar communities led to the prediction that species only co-occur if they partition niche space

along some axis (herein we consider the niche to be the size and shape of multivariate ecological

space that a species utilizes), otherwise one will be excluded through competition (Hutchinson

1957, MacArthur & Levins 1967, May & MacArthur 1972, Brown & Lieberman 1973, Brown

1975, Pianka 1974, M'Closkey 1978). Competitive exclusion can be mitigated if two co-

occurring species use a narrower breadth of resources, producing a more densely packed

community niche space ("niche packing"), or if they exploit habitats or resources that are unused

or non-existent in low-resource or species-poor communities, leading to a larger community

niche space ("niche expansion"; MacArthur 1965, 1970, Pigot et al. 2016, Oliveira et al. 2020).

The foundational work on competition's role in community assembly relied on empirical data

from continental communities with equal biogeographic accessibility but variation in primary

productivity or habitat complexity (Brown & Lieberman 1973, Brown 1975, M'Closkey 1978,

Karr & James 1975, Pianka 1974, Weiher & Keddy 1995), or from archipelagos where island

size or geographic complexity determines resource availability and habitat area, both of which

influence overall species richness (Wilson 1961, Diamond 1975, Lister 1976, Gillespie 2004,

Losos 2009, Losos & Ricklefs 2009). Less explored are insular areas of similar habitat structure

and resource availability, but with discrete communities that vary in species richness ("species

richness anomalies", Swenson et al. 2016 pg. E83). Yet, these anomalies offer powerful systems

for interrogating the role of competition in the distribution of functional diversity within and

among communities because the effects of abiotic processes such as habitat filtering are

minimized (Swenson et al 2016, Li et al. 2017). Most studies of species richness anomalies have

examined plants in continental settings (Latham & Ricklefs 1993, Swenson et al. 2016, Xu et al.

2019) where environmental and historical processes influence the regional species pool. Few, if

any, studies have tested the role of competition in the assembly of discrete communities nested

within an insular setting where species pools are formed through long-distance colonization and

in-situ diversification.

Oceanic islands are often hotspots of diversification and endemism, and their unique

biogeographic and historical conditions can help illuminate the ecological and evolutionary

processes that structure local communities (Losos & Ricklefs 2009). Lineages that arrive and

diversify in isolated locations regularly undergo ecological shifts that result in behaviors or

phenotypes that are uncommon or non-existent in source communities (Carlquist 1966, Millen

2006, Pinto et al. 2008, Esselstyn et al. 2012, 2015, 2021, Stroud & Losos 2016). The presence

of high functional and ecological disparity within an endemic radiation may affect community

structure in multiple ways. First, there is some evidence that community assembly occurs

differently on continents and islands. Using an assortment of phenotypic and behavioral proxies

for resource use across a variety of spatial scales, niche packing is frequently invoked as the

primary process of assembly in species-rich animal communities on continents (Brown 1975,

Karr & James 1975, M'Closkey 1978, MacArthur & MacArthur 1961, Pianka 1974, Pigot et al.

2016, Van de Perre et al. 2020). However, niche expansion has found some support among

isolated island communities of many closely related species (Lister 1976). Second, multiple

lineages evolving in sympatry within an insular setting often promotes niche divergence through

adaptive diversification, reducing phylogenetic niche conservatism (Losos et al. 2003). As a

result, the niche breadth of a community within an insular setting, despite the lower phylogenetic

diversity, may equal that of a similar continental community. Importantly, quantifying trait

differences following adaptive diversification among closely related species can illuminate

which traits are most evolutionary labile and/or important for resource partitioning (Losos et al.

2003, Hiller et al. 2019, Dorey et al. 2020, Stroud 2021).

The murine rodent fauna (rats and mice in the subfamily Murinae) of Sulawesi,

Indonesia, a mountainous, wet tropical, oceanic island at the center of the Wallacean biodiversity

hotspot (Figure 1), is an intriguing system for testing patterns of community niche occupancy.

First, the murine diversity of Sulawesi is exceptional, with at least 48 endemic species stemming

from seven colonist ancestors that arrived from circa 6 Ma to < 1 Ma (Figure S1, Rowe et al.

2019, Handika et al. 2021). While some colonists spawned small radiations of species, others are

evidenced by only a single living species (Table 1, Figure S1). Each clade is widespread on

Sulawesi and most contribute to species richness of all local murine communities on the island.

Second, Sulawesi contains some of the most unusual rodent forms found anywhere (Figure 1b,

Esselstyn et al. 2012, 2015, Rowe et al. 2014) living in sympatry with more typical "rat-like"

morphologies and ecologies. Sulawesi murines have an array of cranial shapes that reflect their

dietary preferences (Figure 1b; Esselstyn et al. 2012, 2015, Martinez et al. 2018), consume a

broad range of foods such as fruit, seeds, fungi, leaves, roots, and earthworms (Musser 2014,

Rowe et al. 2016a), have body sizes ranging from 10 to 500g, and occupy a variety of locomotor

modes (arboreal; scansorial; terrestrial; amphibious, Nations et al. 2021). Third, due to the

topographic complexity of Sulawesi and the positive correlation between elevation and small

mammal diversity in this region (Heaney 2001, Esselstyn et al. 2021), the murine communities

on the islands are partitioned into discrete, montane assemblages (Figure 1). The local montane

murine communities, defined as the species living on a mountain from the upper-lowland forest

to the peak, range from 7 to 23 species, with the upper limit being, to our knowledge, the most

diverse local community of closely related terrestrial mammals on Earth. Lastly, the variation in

local community richness does not appear to be determined by environmental or habitat

differences among mountains but is likely the result of the colonization process outward from the

central core to the peninsulas during the island's formation (Hall 2013, Nugraha & Hall 2018,

Handika et al. 2021). Maximum elevation, which may correlate with the area occupied by

different habitats, appears to play only a minor role in species richness (Figure 1a), and net

primary productivity is nearly constant across the island's montane regions (Imhoff et al. 2004),

suggesting that environmental filtering, often a central process of community assembly (Webb et

al. 2002, Cavender-Bares et al. 2004, Li et al. 2017), does not greatly affect the functional

diversity of Sulawesi's murine communities. Together, these properties generate a fascinating

hierarchical organization: a species pool of small mammals with disparate ecologies and

morphologies living in discrete montane communities that, due to their habitat similarity and

striking disparity in richness, represent species-richness anomalies, all of which lie within a

remote, oceanic island. Such a system presents a compelling natural experiment for testing

alternative hypotheses of how communities assemble to occupy ecological niche space

Figure 1: Small mammal surveys of nine mountains a) on the oceanic island of Sulawesi revealed varying murine
rodent species richness across the island. Each mountain on the map is labeled with the maximum elevation and the
number of murine rodent species present. All but Latimojong were surveyed within 600m of the summit. b) The
diversity of Sulawesi murines is exceptional and includes unique forms that live alongside species with more
"typical" ecologies and morphologies, as demonstrated from surfaces of cranial μCT scans of *i – Rattus hoffmanni,* a
"typical" murine morphology and ecology, *ii* – the shrew rat *Echiothrix leucura* with its extremely elongate rostrum
and soft invertebrate diet, *iii* – the arboreal *Haeromys minahassae*, with a short rostrum and very small size. c)
Photographs: *i – Rattus hoffmanni*, *ii –Echiothrix leucura*, and *iii – Haeromys minahassae*.

Here, we test whether increased species richness in environmentally similar communities

of montane rodents leads to the expansion or packing of ecological niche space. Observational

data, including behavior, life history, and movement patterns, are scarce for the nocturnal,

secretive Sulawesi murines. Much of what we know about these species comes from museum

specimens and their associated metadata (e.g., locality and habitat details, forest strata

preferences, and morphological measurements). Therefore, to infer the ecological niche breadth

of each montane assemblage, we quantify diet, trophic dimension, and microhabitat using 12

individual functional and ecological traits and five multivariate trait complexes to estimate both

the volume and density of community functional space, which we define as the sum of the n-

dimensional functional spaces of the species therein. Importantly, individual traits or groups of

traits related to the same niche axis can reveal distinct, interacting processes (Spasojevic &

Suding 2012, Pigot et al. 2016, Kohli et al. 2021). Additionally, we examine the role of

phylogenetic niche conservatism and trait lability in the assembly of these nine montane

communities by testing for the influence of phylogenetic diversity and inter-clade variation on

functional space occupancy.

188   **Materials and Methods**

The distribution of species- We compiled occurrence records from nine small mammal

inventories of mountain regions on Sulawesi, Indonesia, including one well-documented

mountain surveyed from 1973 to 1976 by Guy Musser and colleagues (Mt. Nokilalaki; Musser,

2014) and eight mountains surveyed between 2011 and 2016 (Ambang, Bawakaraeng,

Buliohuto, Dako, Gandang Dewata, Katopasa, Latimojong, and Torompupu; Figure 1a).

All surveys began in lower primary forest near the line of anthropogenic forest clearing (1100m

to 1500m) and extended to upper-montane forests. All surveys extended to within 600m

elevation of the summit except for Latimojong (highest survey site at 2535m, summit at 3400m).

Trapping records show that there are no Sulawesi murines restricted to elevations above 2500m,

or to habitats within 600m of the summit (Musser 2014). Historical surveys by Musser lasted

several months and were conducted over four years, employing a mix of snap traps and live

traps. Modern surveys (2011-16) lasted an average of 17 days (11-25) and employed similar

collection methods, including a mix of snap traps, live traps, and 20-30L pitfall buckets. All the

murine rodent species known from the sampled localities (Musser 2014, Wilson et al. 2019) were

collected during these modern surveys, indicating a thorough sampling effort. Five new taxa that

were discovered during these expeditions have been described (Musser 2014, Esselstyn et al.

2012, Rowe et al. 2014, Esselstyn et al. 2015, Rowe et al. 2016b) and several new locality

records resulted (Achmadi, et al., 2014; Handika, et al., 2021). Specimens from all surveys were

deposited in the Museum Zoologicum Bogoriense (MZB), Bogor, Indonesia; the American

Museum of Natural History (AMNH), New York, USA; Museums Victoria (MV), Melbourne,

Australia; the Museum of Vertebrate Zoology (MVZ), Berkeley, USA; the Field Museum of

Natural History (FMNH), Chicago, USA; and the Louisiana State University Museum of Natural

Science (LSUMZ), Baton Rouge, USA.

Clade	Crown Age	Species Sampled
Echiothrix	5.11 (4.49-5.69)	10 (10)
Maxomys	3.72 (3.21-4.29)	5 (5)
Bunomys	3.45 (3.09-3.81)	15 (15)
Margaretamys	2.92 (2.29-3.51)	2 (4)
Rattus*	1.57 (1.22-1.95)	6 (6)
Haeromys[†]	Indeterminate	1 (1)

Table 1: Ages of six clades descended from Sulawesi colonists. Median age in millions of years is reported with
95% credible intervals in parentheses. Species sampled reports the number of species in this study with overall clade
richness in parentheses and demonstrates near complete sampling. The two unsampled *Margaretamys* species were
not detected in the localities in this study. All ages taken from Rowe et al. 2019. *There were likely two
colonizations by the ancestors of native *Rattus* spp., the second of which occurred 1.16-0.6 Ma (Rowe et al. 2019),
however, all *Rattus* in this study form a clade relative to other Sulawesi murines. Two human commensal *Rattus*
spp. found on Sulawesi were excluded. [†]The age of arrival of the Haeromys clade is unknown as other species of
Haeromys from Borneo have yet to be included in phylogenetic analyses.

Functional trait data collection and processing- We compiled or generated functional trait

values for 11 continuous traits and one discrete trait for all available Sulawesi murines from each

of the nine communities. Individual Sulawesi murine species exhibit little intraspecific

morphological variation among localities, far less than the morphological differences among

species (Musser 2014), and we therefore combined measurements of individual species from

multiple localities and estimated trait distributions using probabilistic methods to overcome the

limited availability of some traits. Detailed information on traits, sample sizes, data processing,

and multivariate trait composition are available in Table S1. All data, models, and output files

are available in Dryad Repository (to be added prior to publication).

Morphological data collection- We assembled external measurement data from 630 specimens,

including head-body length (mm), tail length (mm), hind-foot length (mm), ear length (mm), and

mass (g), from previously published sources (Wilson et al. 2019, Nations et al. 2021) and online

museum databases. Measured specimens were from the nine surveyed mountains and other

localities on Sulawesi. To obtain ecologically relevant features of external measurements and

mitigate the influence of size in some of our analyses, we calculated three commonly used ratios:

Relative tail length (tail length / (head-body length + tail length)), relative hind-foot length (hind-

foot length / head-body length), and relative ear length (ear length / head-body length) (Nations

et al. 2021, Table S1).

The shape of a rodent's skull and lower jaw provides a wealth of indirect ecological

information on foraging, feeding, and sensory processing, and is often used as a proxy for

fundamental dietary niche (Samuels 2009). We generated μCT scans of the cranium of 64

specimens from 38 species and the mandible of 61 specimens from 36 species (Table S1). Scans

were generated from specimens collected in the nine surveys, as well as from previously

collected museum materials. Stacks of 2D Tiff files were imported to MorphoDig, where 3D

landmarks were placed on cropped volume renderings (Lebrun, 2018). We placed 67 cranial

landmarks (Figure S2) on the left side of the skull, unless damage caused us to use the right side,

in which case we reversed the rendering on the Z-axis. In separate renderings, we placed 20

landmarks on the left dentary of the mandible (Figure S2). Landmarks were exported from

MorphoDig as .stv files and imported into the R package geomorph v.4.0.3 for processing

(Adams et al. 2021, Baken et al. 2021). Missing landmarks (11 of 4288 cranial and three of 1220

mandible landmarks) were imputed, a generalized Procrustes analysis (GPA) superimposition

was performed, and shape coordinates were subjected to a principal components analysis. We

retained the centroid size (an estimate of total size) and the scores from the first 36 principal

components of the cranium and 20 axes of the mandible, each representing >95% of the shape

variation of the element.

Stable isotope data collection- Approximately 1-2 grams of hairs were plucked from the rump of

286 dry museum specimens collected on six focal surveys (Ambang, Bawakaraeng, Buliohuto,

Dako, Gandang Dewata, and Latimojong). Isotopic values can vary regionally (Fry 2006);

therefore, we collected hair samples from multiple individuals of each species from each locality

(mean = 4.6 specimens/species/locality, range = 1-14). Nitrogen stable isotope values (δ^{15}N) act

as a proxy for consumer trophic position as they generally increase by 3–5% per trophic level

(DeNiro & Epstein 1981). Carbon isotope values (δ^{13}C) generally exhibit little to no change with

trophic position and are commonly used as proxies of consumer's basal carbon resource use (e.g.

use of differing primary production energy pathways; DeNiro & Epstein 1978). Combined, these

two metrics are commonly used to quantify the "isotopic niche" of consumers, which can act as a

useful proxy of species realized dietary niche (Newsome et al. 2007, Ben-David & Flaherty

2012). Stable isotope values are reported in delta notation in per mil units. Samples were

processed at the Stable Isotope Ecology Laboratory, Department of Oceanography & Coastal

Sciences, Louisiana State University. Details on sample cleaning, processing, and analysis are in

the Supporting Methods.

Locomotor mode data collection- We used the locomotor classification scheme from Nations et

al. (2019, 2021) to group each Sulawesi murine into one of four discrete locomotor modes:

Arboreal, (climbing is integral to survival); General (navigates a variety of substrates and habitat

strata); Terrestrial (on the ground surface); and Amphibious, (dependent on aquatic habitats for

foraging).

Combined traits – We combined our 11 continuous traits into five multivariate traits that

represent distinct niche dimensions: (1) head shape (cranium shape PC 1-36 and mandible shape

PC 1-20); (2) isotopic niche space (δ^{15}N and δ^{13}C stable isotope values); (3) body proportions

(head-body length, relative tail length, relative hind-foot length, and relative ear length); (4) body

size (log(mass) and head-body length); and (5) total morphological shape (all nine morphological

traits). We did this by combining the predicted trait values for each community (see below).

Estimating species' trait values- The sample size of morphological and isotopic data varied

between species and mountain communities, and in some cases was limited to one individual,

such as with the amphibious *Waiomys mammasae*, a species known from a single specimen

(Rowe et al. 2014). To mitigate uneven sampling, we estimated a probability distribution of

species trait values using partial-pooling in a multilevel Bayesian model. Unlike complete

pooling (one global mean estimated for all combined samples) or no-pooling (one mean

estimated per species, independent of all others), partial-pooling estimates a mean for each

species as well as the variance among species, which serves as an adaptive prior that is common

to all the species' means (McElreath 2020). The Bayesian partial-pooling modeling allowed us to

incorporate intraspecific variation in trait value predictions while avoiding point estimates such

as averages, which discard valuable information. This approach prevents unbalanced estimates

by using the trait-variance probability estimates from well-sampled species to inform variance

estimates of species with fewer samples (Gelman & Hill 2006, McElreath 2020). All analyses

were conducted in the probabilistic programming language Stan (Carpenter et al. 2017) within

the R library brms v. 2.17.0 (Bürkner 2018). Subsequent data processing and figuring relied on

the R libraries tidyverse v. 1.3.1 (Wickham et al. 2020), furrr v. 0.3.0 (Vaughan 2021), and

tidybayes v. 3.0.2 (Kay 2020). All data, scripts, and output files are available in Zenodo

Repository (to be added prior to publication) and on GitHub (to be added).

For each continuous trait (Table S1), we used the trait value as the response variable, and

used species as a group-level predictor. All traits were scaled to unity prior to analyses. We used

the student-*t* distribution to describe the response variable to minimize the influence of rare,

extreme observations (a.k.a. 'robust regression', Kruschke 2013, McElreath 2020). Each model

included four chains with 4000 iterations of warm-up and 1000 sampling iterations. Posterior

predictions from the four chains were combined, resulting in 4000 samples per trait per species.

To mitigate the potential geographic signal in stable isotope values (Fry 2006), and facilitate

randomization (see below), the posterior estimates of the δ^{15}N and δ^{13}C stable isotope values

were estimated with models that included an additional 'community' group-level effect, but

otherwise were identical to the models described above (Supporting Methods). These stable

isotope models generated posterior values of species' isotopic measurements conditioned on the

locality, enabling comparisons among communities and randomized sampling for null models

(Table S1, Figure S3). Full details of the model, prior, and chain-estimation are described in the

Supporting Methods.

Estimating functional space volume – We defined functional space volume as the total volume of

n-dimensional trait space occupied by species in a community, and for consistency and clarity

we use the term 'functional space volume' for all individual and multivariate traits, regardless of

dimension. We used the sum of the variance of each trait value to estimate the volume of

community functional space for each community, which is less sensitive to outliers and

outperforms other metrics such as ellipse volumes, convex hull volumes, and hypervolumes,

especially for functional spaces with many axes (Li et al. 2017, Guillerme et al. 2020). We

estimated a posterior distribution of the variance of each trait for each community by first

grouping the species' trait value estimates by community. Then, for each of the 4000 posterior

draws (one draw ranging between 7 and 23 values, depending on the richness of the community),

we estimated the variance of the trait, resulting in a distribution of 4000 variance values for each

trait for each community. We estimated the combined multivariate trait space variance by

summing the variances of each trait, then dividing by the number of traits. We estimated the 89%

probability values for the variance of each individual and combined trait space. To estimate

locomotor mode variance, we dummy coded locomotor mode into four binary columns (one per

mode) and performed a redundancy analysis (Legendre & Legendre 2012) with the rda()

function in the vegan package (Oksanen et al. 2019). We then extracted the PC scores for each

species and calculated the sum of variances of the three PC axes for each community.

Estimating functional space density- We estimated the density of species within functional trait

space with the mean nearest neighbor (NN) metric (Guillerme et al. 2020). Our methods follow

the estimates of functional space volume above. First, we grouped species' trait values by

community, then estimated the individual trait density for each posterior draw using the NN

metric in the disparity() function of the R library dispRity (Guillerme 2018), resulting in a

distribution of 4000 NN values for each trait and combined trait space. To estimate locomotor

mode density, performed a redundancy analysis on the dummy-coded locomotor data as above.

We then extracted the PC scores for each species, grouped the species by community, and

estimated the NN value for each community as with the continuous traits above, in this case

generating only a single NN value per community rather than a distribution.

Null models of functional space volume and density- We used null models to determine the

difference between the functional space of our nine communities and randomly assembled

communities. For each community we created 1000 randomized null communities of $n_{community}$

species for each of the nine localities using the independent swap algorithm (Gotelli 2001). We

then calculated both volume and density as above for each individual (n=12) and combined

(n=5) trait for each of the 1000 randomized community samples. Standardized effect size (SES)

was calculated as:

$$SES = \frac{Trait_{observed} - Mean(Trait_{Random})}{SD(Trait_{Random})},$$

where $Trait_{observed}$ is the vector of 4000 samples of the density or volume of the given trait,

$Mean(Trait_{Random})$ and $SD(Trait_{Random})$ are the mean and standard deviation of the density

or volume values of the given trait from the 1000 random species assemblies. Positive SES

values indicate greater than random functional space volume or lower than random functional

space density (overdispersion), meaning that species in that community are occupying functional

space outside the range of other communities and/or their niches are farther apart in functional

space (niche expansion). Negative SES values indicate lower than random functional space

volume and greater than random functional space density (underdispersion or clustering),

meaning that species in that community are occupying less ecological space than other

communities and/or their niches are closer together in functional space (niche packing; Oliveria

et al. 2020).

Effect of species richness on community functional space volume and density- If niche expansion

is the primary mode of resource partitioning in Sulawesi murine communities, then we expected

that functional space volume will increase with increasing species richness, and that functional

space density will remain stable as richness increases. Whereas if niche packing is occurring, we

expected that functional space density will increase with species richness, and that there will be

no effect of richness on functional space volume. To quantify the effect size of richness

($\beta_{richness}$) on functional space volume and density, we used Bayesian linear regression models

that include the measurement error of the predictor variable to estimate the effect size of species

richness on volume or density. We used species richness as the predictor variable, and the mean

and standard error of the trait estimates for each community as the response. Measurement error

models contain vastly more information per observation than a single point observation and

provide robust estimates despite the small number of sampled communities (Bürkner 2018,

McElreath 2020). We also calculated the Bayesian R^2 value for each regression (Gelman et al.

2019). The models, priors, and chain estimations are detailed in the Supporting Methods.

Phylogenetic diversity as a path to trait disparity- The ecological space occupied by a

community may depend on which lineages or clades are present (Webb 2000). Trait disparity

appears to vary among clades of Sulawesi murines, and if so, then the volume and densities of

functional spaces may be more influenced by phylogenetic diversity than ecological factors. To

determine how phylogenetic diversity mediates functional space occupancy in the nine murine

communities, we estimated the phylogenetic diversity (PD) of each community using Faith's

metric of branch length (Faith 1996). We removed all but the Sulawesi species (n = 35) from a

time-calibrated phylogenetic hypothesis of Murinae (Nations et al. 2021). For the analyses

described above, we had trait data for four species that are not included in this phylogeny:

Maxomys wattsi, Rattus bontanus, *Rattus mollicomulus*, and *Rattus marmosurus*. We manually

added these four species into the tree using the R package phytools (Revell 2012). Details are

found in the Supplemental Methods. We used the R package picante (Kembell et al. 2010) to

estimated Faith's metric of PD, and to sample 1000 random communities in order to calculate the

SES value of PD (SES PD). We used linear modeling in brms to estimate the effect size of

species richness on SES PD. *Haeromys minahassae* is the only representative of its genus on

Sulawesi, and, due to its large phylogenetic distance from other Sulawesi murines, may have an

oversized impact on estimates of community PD. Therefore, we repeated the estimates of SES

PD and effect size above without *Haeromys minahassae*. To quantify the trait disparity within

and among Sulawesi murine clades, we grouped the posterior distributions of the species'

predicted trait values by clade, then estimated the variance of each of the 12 univariate traits. We

plotted the trait value variance for each community along with the species' predicted trait values

(Figure S4)

Figure 2: The effect of species richness on community functional space volume (a, b) and density (c, d). X-axes
show the $\beta_{richness}$ estimates (the regression slope) for each trait space (y-axis) on the left and the Bayesian R^2 for
each $\beta_{richness}$ estimate on the right. Colored point intervals show 89% probability of $\beta_{richness}$ estimates (effect
size), with color varying by $\beta_{richness}$ value. Black point intervals show 50% posterior estimates of Bayesian R^2 for
each trait space. Density was estimated using the mean nearest neighbor (NN), and a high NN distance indicates low
density. Four of the five multivariate traits (**a**) and seven of the 12 individual traits (**b**), show an increase in
functional space volume (trait variance) with greater richness (i.e., positive $\beta_{richness}$). Phylogenetic diversity also
increases with species richness (**b**). All multivariate traits (**c**) and individual traits (**d**) show a stable or, surprisingly,
decreasing functional space density (increased NN distance) with greater species richness.
423

424   **Results**

Models of trait values- All parameters in the Bayesian multilevel model estimates of trait-value

distributions and measurement-error estimates of the species richness effect on trait values had

ESS > 1000 and a Gelman-Rubin diagnostic $\hat{R} \leq 1.01$, indicating convergence. Raw volume and

density values differed from the volume and density of SES. To avoid the confounding influence

of species richness in volume and density estimates we report and discuss only the SES values

(Swenson 2014).

Effects of Species Richness on Functional Space Volume and Density- Functional space volume

was positively correlated with species richness in seven of the 12 traits (locomotor mode, tail

length, head-body length, cranium shape, mass, δ^{15}N, and hind foot length), as shown by the

positive $\beta_{richness}$ slope values from the Bayesian linear models (Figures 3, Table S3) and the

greater variance with higher community richness (Figure 3). The remaining five traits (ear

length, cranium size, dentary size, dentary shape, and δ^{13}C) did not substantially increase in

volume with increased species richness. Among the multivariate traits, the SES values of total

morphological shape increased substantially with species richness while head shape and isotopic

niche space show a moderate probability of increasing with species richness (Figures 3, 4, Table

S2, S3).

Linear regression showed that species richness had little to no effect on functional space

density (NN SES values) for 11 of the 12 individual traits and three of five multivariate traits

(Figures 3, S4, Table S2), consistent with niche expansion. The exceptions were a positive

$\beta_{richness}$ slope (i.e., decreased density with richness) for locomotor mode, total morphological

shape, and external proportions, suggesting an extreme overdispersion of morphology and

locomotor diversity in the richest communities.

Figure 3: Estimated functional volume SES plotted against species richness: **a**) Multivariate trait volumes, **b**) Skull traits, **c**) External measurements, and **d**) Isotopic measures, locomotor mode, and phylogenetic diversity. Points represent the mean SES values and error bars indicate the 89% credible intervals. X-axis tick marks show species richness for each community. Values equal to zero are consistent with null expectations, positive values indicate overdispersion, and negative values show underdispersion (trait clustering). Dotted lines depict the 89% interval of the null distribution. The plot of trait densities is shown in the supporting information (Figure S4).

Phylogenetic Diversity and Functional space- Phylogenetic diversity (PD) increased with species

richness (Figures 3b, 4d, Table S2). Including *Haeromys minahassae* in the estimates of PD

changed the PD SES values of the individual communities but had minimal impact on linear

regressions (Table S4). Results including *H. minahassae* had a slope ($\beta_{richness}$) of 0.143 (0.03,

0.252) while excluding *H. minahassae* generated a slope of 0.169 (89% C.I. of 0.041, 0.295), and

we therefore present the results that include *H. minahassae*. Katopasa, a community of 12

murines on Sulawesi's Eastern Peninsula, has the lowest PD SES value followed by

Bawakaraeng on the Southwestern Peninsula (seven species). Gandang Dewata, the richest

community (23 species), has the highest PD SES value (Figure 3a, Table S2). As suspected,

functional space volumes vary between clades, though not consistently (Figure S5). For example,

the Echiothrix clade has the highest variance in cranial shape, dentary shape, and dentary size but

low trait variance for hind-foot length and $\delta^{15}N$. In contrast, the Maxomys clade has low variance

for all trait values except tail length and $\delta^{15}N$ (Figure S5). This results in different densities of

functional space occupation among traits, where some trait volumes, such as cranial space, are

strongly influenced by phylogeny, while others, such as isotopic niche space, have high and low

values distributed among clades (Figure 4).

Figure 4: The mode of functional space filling varies among traits. The bivariate plots on the left depict the 2D functional spaces. Each black shape is the mean value of a particular species in each clade (shown in legend). The convex hull colors match the color of the community functional volume SES distributions in the right-hand columns. The thick convex hull lines are mean values, and 500 random samples from the posterior of each trait are shown in thin hull lines. The number of species in each community and each clade used in this study is shown next to the SES distributions in parentheses. All values were scaled to z-scores prior to analyses. **a)** Cranial morphospace values are mostly clustered tightly near the mean of each axis, apart from Echiothrix species and the single *Haeromys* species. Intraspecific variance is also relatively low on these axes. **b)** Intraspecific variance is high in isotopic niche space. High and low $\delta^{15}N$ values are distributed among clades, but that is not the case for $\delta^{13}C$ values. **c)** Large values of hind foot length belong to members of two clades, while large and small tail length values are dispersed among all six clades, reducing the influence of phylogenetic diversity on tail length disparity.

Discussion

Variation in species richness among communities of closely related species that inhabit similar environments provides a unique window to explore how interspecific competition may affect community functional richness in the absence of confounding factors like environmental

variation (Roughgarden 1976, Swenson et al. 2016). Unfortunately, these species richness

anomalies are uncommon (Swenson et al. 2016, Van de Perre et al. 2020). Here we used a large

dataset of ecological and morphological traits to estimate the changes in community niche

occupancy across a richness gradient on an oceanic island. Though the trait estimates of some

species necessarily stem from a small number of specimens, and therefore may be subject to

error, our novel probabilistic approach incorporates measurement error and species-level

variation into the posterior distribution and propagates this uncertainty through the estimation of

community niche space. The functional space of most traits increased with greater species

richness among the nine murine communities studied while there was no change in functional

space density, consistent with limiting similarity. Locomotor mode disparity has the strongest

positive relationship with species richness, but the functional volumes of skull and body shape

(indicators of diet, locomotion, and microhabitat use in murines and vertebrates in general;

Peters 1986, Losos 2009, Martinez et al. 2018, Nations et al. 2021), also strongly increase with

species richness (Figure 2). Measurements of overall body size, cranial size, dentary shape and

size, ear length, and $\delta^{13}C$ values all demonstrate either a weak signal of niche packing, or no

signal across the species richness gradient. While the volume of many functional spaces

increases with richness, functional space density shows little correlation with species richness for

most traits, and it surprisingly decreases with increased richness for external morphology and

locomotor mode. These results, along with a general underdispersion of trait values in low-

richness communities and overdispersion in high-richness communities (Table S2, Figures 4 &

S4), suggest that species in rich communities mitigate competitive interactions by occupying

underused niche space. We also found that increasing phylogenetic diversity is a means of

increasing functional space occupation. Combined, these results point to species interactions as a

mechanism for both phenotypic and phylogenetic overdispersion (Webb et al. 2002, Cavender-

Bares et al. 2004, Li et al. 2017) and suggest that, given adequate resources, structural

complexity, and evolutionary time, lineages can evolve to occupy unique regions of ecospace,

often far from the average trait value, which minimizes niche overlap and cultivates exceptional

richness.

Whether in a single desert valley (Brown 1975), or across continental (Maestri &

Patterson 2016, Kohli et al. 2022) and global latitudinal gradients (Karr & James 1975, Pellissier

et al. 2018), structurally complex habitats are thought to foster higher species diversity. High

plant diversity creates a more complex, vertically structured habitat matrix for other plant and

animal species to occupy and has long been tied to higher animal richness (Hutchinson 1959,

MacArthur & MacArthur 1961, Scheffers et al. 2013, Oliveira & Scheffers 2019). Our estimates

of locomotor-mode occupancy clearly demonstrate that vertical habitat partitioning is critical to

maintaining high species richness in Sulawesi murines (Figure 2, Table S2). Strikingly, the

density of locomotor trait space decreases along the richness gradient, indicating a very high

level of trait overdispersion (Figures 3c & S4). Arboreal, Terrestrial, and Amphibious locomotor

modes each provide access to different microhabitats that contain similar resources. Among the

23 species found on Mt. Gandang Dewata, the amphibious *Waiomys mamasae*, the terrestrial

Paucidentomys vermidax, and the arboreal *Sommeromys macrorhinos*, all consume invertebrates

and have similar δ^{15}N values, yet they are unlikely to compete for resources due to their distinct

microhabitat use, a pattern observed in other insular communities of closely related vertebrates

(e.g., Jamaican *Anolis*; Shoener 1974). It's worth noting that when the lone amphibious species

Waiomys mamasae, only known from Gandang Dewata, is removed from the data, the locomotor

variance of Gandang Dewata remains the highest among the communities. Morphological

measurements are often used to infer locomotor mode in a variety of eco-evolutionary contexts

(Ricklefs & Travis 1980, Samuels and Van Valkenburgh 2009, Pianka et al. 2017, Verde-

Arregoitia et al. 2019), and our estimates of tail-length variance, a trait correlated with

locomotion in murines (Nations et al. 2021), increases with richness at nearly the same rate as

locomotor mode (Figure 2). Importantly, our results suggest that locomotor mode rankings may

be an effective way to estimate community locomotor variance where continuous trait data are

lacking.

Combining many traits into one multivariate measure of functional diversity is a common

approach in evolutionary ecology. Ordination techniques were especially promoted to overcome

pitfalls from early community ecology studies that used few, largely subjective measures of

resource use (Ricklefs & Travis 1980). However, merging traits into multivariate axes masks

trait-specific processes related to functional space (Spasojevic & Suding 2012, Astor et al. 2014).

Indeed, our trait volume and density estimates reveal distinct patterns between individual and

combined traits. For example, isotopic niche space, a combined signal of δ^{15}N and δ^{13}C values

commonly used in terrestrial ecological studies, exhibits an equivocal signal (Figure 2b), but

individual isotopic values reveal that the packing signal originates from the static δ^{13}C value

along the species richness gradient. The δ^{15}N value, a signal of trophic level, expands along the

richness gradient, a result that would be overlooked in multivariate analyses. In contrast to δ^{15}N,

dentary shape exhibits an equivocal signal, despite its presumed relationship to dietary niche

(Maestri et al. 2016, Kohli et al. 2019). These opposing patterns are only illuminated by

analyzing individual traits (Spasojevic & Suding 2012). If we performed this study using only

cranial size and dentary shape, which are thought to capture the important axes of size and diet in

murine rodents (Rowsey et al. 2019, 2020) and mammals in general (Prevosti et al. 2012,

Grossnickle 2020), our results would suggest that functional volume does not increase with

species richness.

The disparity of traits within a clade mediates the impact of phylogenetic diversity on

community functional space volume. If trait values are phylogenetically clustered, then

increasing phylogenetic diversity is necessary for community niche expansion to occur. But if

trait values are phylogenetically overdispersed, niche packing, expansion, or, as we found, both

could result from increased phylogenetic diversity. The distribution of traits among clades is of

particular interest in communities that are assembled through a mix of colonization and in situ

speciation, such as Sulawesi murines, Caribbean anoles, or Hawaiian spiders (Gillespie 2004,

Losos 2009, Rowe et al. 2019). Niche divergence has been hypothesized to overcome niche

conservatism in communities with an extended history of coevolution, such as those on oceanic

islands, likely diminishing the signal of phylogenetic trait clustering (Losos et al. 2003). We find

that niche divergence and conservatism may occur simultaneously on different functional traits.

For example, the elongate, highly distinct skulls of some species in the Echiothrix clade, the

descendants of the first murine colonists on Sulawesi, set them apart from other clades in skull

shape and, despite their relatively low abundance, these species overcontribute to community

cranial and dentary shape volumes (Figures 3b, 4b, S5). Unlike cranial shape however, the

Echiothrix clade occupies a very constrained portion of δ^{15}N trophic space. Additionally, trophic

level estimates from δ^{15}N values are notably dispersed among clades (Figures 4c, S5), and high

phylogenetic diversity is not necessary for high isotopic niche space estimates. Are there innate

differences in these traits that could lead to opposing patterns of niche conservatism?

The evolutionary lability of ecologically important traits determines the rate of

convergence and divergence possible within a given time frame and can directly influence

dispersion of trait values among species (Cavender-Bares et al. 2004). The traits with the highest

within-clade variance — $\delta^{15}N$, tail length, body size — are all thought to be evolutionarily labile.

Changes in tail length, body size, and intestinal tract morphology can occur on brief evolutionary

time scales (Powell & King 1997, Naya et al. 2008, Kingsley et al. 2017, 2021), whereas

morphological changes in cranial shape, such as substantial rostral elongation and the reduction

of molar grinding area, may take much longer. Indeed, insular species are known to have rapidly

expanded breadths of diet and labile morphological traits following colonization (Stuart et al.

2014) and subsequent speciation (Wilson 1959, 1961, Lister 1976, Millien 2006, Rowe et al.

2016a). But the extreme cranial morphologies of some Sulawesi murines, particularly those of

the oldest radiation on the island (Echiothrix clade), are likely the result of a long process of in-

situ evolution (Rowe et al. 2019). In other words, on shorter time scales, similarities in some

slowly evolving traits may lead to higher divergences in more labile traits, as evidenced in the

Maxomys and Bunomys clades (Figure S5). The opposite pattern may also occur, but only

following sufficient evolutionary time. We propose that the evolutionary lability of traits is a

determinant of trait value dispersion (Webb et al. 2002, Cavender-Bares et al. 2004), which

directly relates to our inferences of functional space occupancy in Sulawesi murine communities.

The theory of niche complementarity suggests that a pair of coexisting species that are similar in

one trait should diverge in another trait (Schoener 1974). Yet, in an isolated setting, the lability

of the traits in question necessarily determines the degree of complementarity possible within a

given time frame. Linking trait lability with niche complementarity in an island system has

important implications for the generation of functional and taxonomic diversity and may further

illuminate the process of niche filling in insular, adaptive radiations.

616   **Conclusion**

618   Here we provide evidence that limiting similarity in functional traits reflecting

locomotion and microhabitat use plays an important role in the assembly of discrete, montane

small mammal communities on an oceanic island. Our results contrast with recent studies that

recovered niche packing as key to increased species richness in continental tropical vertebrate

communities (Pigot et al. 2016, Peixoto et al. 201, Pellissier et al. 2018, Van de Perre 2020,

Dehling et al. 2022, Hughes et al. 2022). Furthermore, our results counter the predictions that

niche packing is expected to occur if resources remain constant among communities

(Roughgarden 1976). Yet, Roughgarden (1976) posited that the predicted relationship between

species richness and resources may be different on remote islands, as the distance from source

populations may affect both community richness and the functional trait values of the species

present. The organization of distinct montane communities on an oceanic island may be just such

an example. The regional murine species pool of Sulawesi is itself structured by idiosyncratic

immigration, diversification, and (presumably) extinction dynamics, and the resultant species'

functional traits. The species in these communities that occupy the edges of some functional

spaces, such as cranial shape (Figure 4), represent morphologies and ecologies found only on

Sulawesi or other large, oceanic islands. And the "imperfect isolation" (sensu Samonds et al

2013) of Sulawesi allows for other regional murine taxa, often represented by more "average"

murine phenotypes, to periodically colonize the island, adding more species into the center of

functional space (Figure 4). The presence of such disparate phenotypes and ecologies on

Sulawesi (Esselstyn et al. 2012, 2015, Rowe et al. 2014) produces a larger functional space

reserve than is available in most, if any, continental rodent systems, nurturing both niche

expansion and high species richness. Therefore, the complex topography, isolation, abundant

resources, and sequential colonization of Sulawesi might lead to species assembly processes that

are typical of other large, oceanic islands, but are atypical of continental systems, "sky islands",

geologically younger islands, or regions with less abundant resources (deserts, high latitude

habitats), but the presence of this general pattern remains untested.

Literature Cited

Achmadi, A. S., K. C. Rowe, and J. A. Esselstyn. 2014. New records of two rarely encountered,

endemic rats (Rodentia: Muridae: Murinae) from Gunung Gandangdewata, West

Sulawesi province. Treubia 41:51-60.

Adams, D. C., M. Collyer, and A. Kaliontzopoulou. 2021. Geomorph: Software for geometric

morphometric analyses. R package version 4.0.3.

Astor, T., J. Strengbom, M. P. Berg, L. Lenoir, B. Marteinsdóttir, and J. Bengtsson. 2014.

Underdispersion and overdispersion of traits in terrestrial snail communities on islands.

Ecology and Evolution 4:2090-2102.

Baken, E. K., M. L. Collyer, A. Kaliontzopoulou, and D. C. Adams. 2021. Geomorph v4.0 and

gmShiny: enhanced analytics and a new graphical interface for a comprehensive

morphometric experience. Methods in Ecology and Evolution. 12:2355-2363.

Baltensperger, A. P., F. Huettmann, J. C. Hagelin, and J. M. Welker. 2015. Quantifying trophic

niche spaces of small mammals using stable isotopes (δ^{15}N and δ^{13}C) at two scales across

Alaska. Canadian Journal of Zoology 93:579-588.

Ben-David, M., and E. A. Flaherty. 2012. Stable isotopes in mammalian research: a beginner's

guide. Journal of Mammalogy 93:312-328.

Brown, J. H., and G. A. Lieberman. 1973. Resource utilization and coexistence of seed⬚eating

desert rodents in sand dune habitats. Ecology 54:788-797.

Brown, J. H. 1975. Geographical Ecology of Desert Rodents. Pages 315-341 *in* M. L. Cody and

J. M. Diamond, editors. Ecology and Evolution of Communities. The Belknap Press,

Cambridge.

Bürkner, P.-C. 2018. Advanced Bayesian Multilevel Modeling with the R Package brms. The R

Journal 10:395–411.

Carlquist, S. 1966. The Biota of Long-Distance Dispersal. I. Principles of Dispersal and

Evolution. The Quarterly Review of Biology 41: 247–270.

Carpenter, B., A. Gelman, M. D. Hoffman, D. Lee, B. Goodrich, M. Betancourt, M. Brubaker, J.

Guo, P. Li, and A. Riddell. 2017. Stan: A Probabilistic Programming Language. Journal

of Statistical Software 76.

Cavender-Bares, J., D. D. Ackerly, D. A. Baum, and F. A. Bazzaz. 2004. Phylogenetic

Overdispersion in Floridian Oak Communities. The American Naturalist 163:823-843.

Dehling, D. M., G. V. Dalla Riva, M. C. Hutchinson, and D. B. Stouffer. 2022. Niche packing

and local coexistence in a megadiverse guild of frugivorous birds are mediated by fruit

dependence and shifts in interaction frequencies. The American Naturalist 199:855–868.

DeNiro, M. J., and S. Epstein. 1978. Influence of diet on the distribution of carbon isotopes in

animals. Geochimica et Cosmochimica Acta 42:495-506.

DeNiro, M. J., and S. Epstein. 1981. Influence of diet on the distribution of nitrogen isotopes in

animals. Journal of Archeological Science 45:341-351.

Des Roches, S., D. M. Post, N. E. Turley, J. K. Bailey, A. P. Hendry, M. T. Kinnison, J. A.

Schweitzer, and E. P. Palkovacs. 2018. The ecological importance of intraspecific

variation. Nature Ecology & Evolution 2:57-64.

Diamond, J. M. 1975. Assembly of Species Communities. Pages 342-444 *in* M. L. Cody and J.

M. Diamond, editors. Ecology and Evolution of Communities. The Belknap Press,

Cambridge.

Dorey, J. B., S. V. Groom, E. H. Freedman, C. S. Matthews, O. K. Davies, E. J. Deans, C.

Rebola, M. I. Stevens, M. S. Lee, and M. P. Schwarz. 2020. Radiation of tropical island

bees and the role of phylogenetic niche conservatism as an important driver of

biodiversity. Proceedings of the Royal Society B 287:20200045.

Esselstyn, J. A., A. S. Achmadi, and K. C. Rowe. 2012. Evolutionary novelty in a rat with no

molars. Biology Letters 8:990-993.

Esselstyn, J. A., A. S. Achmadi, H. Handika, and K. C. Rowe. 2015. A hog-nosed shrew rat

(Rodentia: Muridae) from Sulawesi Island, Indonesia. Journal of Mammalogy 96:895-

907.

Esselstyn, J. A., A. S. Achmadi, H. Handika, M. T. Swanson, T. C. Giarla, and K. C. Rowe.

2021. Fourteen New, Endemic Species of Shrew (Genus *Crocidura)* from Sulawesi

Reveal a Spectacular Island Radiation. Bulletin of the American Museum of Natural

History 454:1-108.

Faith, D. P. 1996. Conservation priorities and phylogenetic pattern. Conservation Biology

10:1286-1289.

Fry, B. 2006. Stable isotope ecology. Springer, New York.

Galetti, M., R. R. Rodarte, C. L. Neves, M. Moreira, and R. Costa-Pereira. 2016. Trophic Niche

Differentiation in Rodents and Marsupials Revealed by Stable Isotopes. PLOS One

11:e0152494.

Gelman A, Hill J. 2006. Data Analysis Using Regression and Multilevel/Hierarchical Models.

Cambridge: Cambridge University Press.

Gelman, A., B. Goodrich, J. Gabry, and A. Vehtari. 2019. R-squared for Bayesian regression

models. The American Statistician 73:307-309.

Gillespie, R. 2004. Community assembly through adaptive radiation in Hawaiian spiders.

Science 303:356-359.

Gotelli, N. J. 2001. Research frontiers in null model analysis. Global Ecology and Biogeography

10:337-343.

Grossnickle, D. M. 2020. Feeding ecology has a stronger evolutionary influence on functional

morphology than on body mass in mammals. Evolution 74:610-628.

Guillerme T. 2018. dispRity: A modular R package for measuring disparity. Methods in Ecology
and Evolution 9:1755-1763.

Guillerme, T., M. N. Puttick, A. E. Marcy, and V. Weisbecker. 2020. Shifting spaces: Which
disparity or dissimilarity measurement best summarize occupancy in multidimensional
spaces? Ecology and Evolution 10:7261-7275.

Hall, R. 2013. The palaeogeography of Sundaland and Wallacea since the Late Jurassic. Journal
of Limnology 72(s2):1-17.

Handika, H., A. S. Achmadi, J. A. Esselstyn, and K. C. Rowe. 2021. Molecular and
morphological systematics of the Bunomys division (Rodentia: Muridae), an endemic
radiation on Sulawesi. Zoologica Scripta 50:141-154.

Heaney, L. R. 2001. Small mammal diversity along elevational gradients in the Philippines: an
assessment of patterns and hypotheses. Global Ecology and Biogeography 10:15–39.

Hiller, A. E., M. S. Koo, K. R. Goodman, K. L. Shaw, P. M. O'Grady, and R. G. Gillespie. 2019.
Niche conservatism predominates in adaptive radiation: comparing the diversification of
Hawaiian arthropods using ecological niche modelling. Biological Journal of the Linnean
Society 127:479-492.

Hughes, E. C., D. P. Edwards, J. A. Bright, E. J. Capp, C. R. Cooney, Z. K. Varley, and G. H.
Thomas. 2022. Global biogeographic patterns of avian morphological diversity. Ecology
Letters 25:598-610.

Hutchinson, G. E. 1957. Concluding remarks. Cold Spring Harbor Symposia on Quantitative
Biology 22:415-427.

Hutchinson, G. E. 1959. Homage to Santa Rosalia or Why Are There So Many Kinds of
Animals? The American Naturalist 93:145-159.

Imhoff, M. L., L. Bounoua, T. Ricketts, C. Loucks, R. Harriss, and W. T. Lawrence. 2004.

Global patterns in human consumption of net primary production. Nature 429:870-873.

Karr, J. R., and F. C. James. 1975. Ecomorphological Configurations and Convergent Evolution.

Pages 258-291 *in* M. L. Cody and J. M. Diamond, editors. Ecology and Evolution of

Communities. The Belknap Press, Cambridge.

Kay, M. 2020. Tidybayes: Tidy data and geoms for Bayesian models. R package version 2.1.1.

Kembel, S. W., P. D. Cowan, W. K. Helmus, W. K. Cornwell, H. Morlon, D. D. Ackerly, S. P.

Blomberg, and C. O. Webb. 2010. Picante: R tools for integrating phylogenies and

ecology. Bioinformatics 26:1463-1464.

Kingsley, E. P., K. M. Kozak, S. P. Pfeifer, D. S. Yang, and H. E. Hoekstra. 2017. The ultimate

and proximate mechanisms driving the evolution of long tails in forest deer mice.

Evolution 71:261-273.

Kingsley, E. P., E. R. Hager, J.-M. Lassance, K. M. Turner, O. S. Harringmeyer, C. Kirby, B. I.

Neugeboren, and H. E. Hoekstra. 2024. Adaptive tail-length evolution in deer mice is

associated with differential Hoxd13 expression in early development. Nature Ecology &

Evolution:1–15.

Kohli, B. A., and M. A. Jarzyna. 2021. Pitfalls of ignoring trait resolution when drawing

conclusions about ecological processes. Global Ecology and Biogeography 30:1139–

1152.

Kohli, B. A., M. J. Miyajima, and M. A. Jarzyna. 2022. Elevational diversity patterns of rodents

differ between wet and arid mountains. Global Ecology and Biogeography 31:1726-1740.

Kohli, B. A., and R. J. Rowe. 2019. Beyond guilds: the promise of continuous traits for

mammalian functional diversity. Journal of Mammalogy 100:285-298.

Kohli, B. A., R. D. Stevens, E. A. Rickart, and R. J. Rowe. 2021. Mammals on mountainsides

revisited: Trait☐based tests of assembly reveal the importance of abiotic filters. Journal

of Biogeography 48:1606-1621.

Kruschke, J. K. 2013. Bayesian estimation supersedes the t test. Journal of Experimental

Psychology: General 142:573.

Latham, R. E., and R. E. Ricklefs. 1993. Global Patterns of Tree Species Richness in Moist

Forests: Energy-Diversity Theory Does Not Account for Variation in Species Richness.

Oikos 67:325–333.

Lebrun, R. 2018. MorphoDig, an open-source 3D freeware dedicated to biology. IPC5, Paris,

France.

Legendre, P., and L. Legendre. 2012. Numerical ecology. Elsevier, Amsterdam.

Li, Y., B. Shipley, J. N. Price, V. de L. Dantas, R. Tamme, M. Westoby, A. Siefert, B. S.

Schamp, M. J. Spasojevic, V. Jung, D. C. Laughlin, S. J. Richardson, Y. L. Bagousse-

Pinguet, C. Schöb, A. Gazol, H. C. Prentice, N. Gross, J. Overton, M. V. Cianciaruso, F.

Louault, C. Kamiyama, T. Nakashizuka, K. Hikosaka, T. Sasaki, M. Katabuchi, C.

Frenette Dussault, S. Gaucherand, N. Chen, M. Vandewalle, and M. A. Batalha. 2018.

Habitat filtering determines the functional niche occupancy of plant communities

worldwide. Journal of Ecology 106:1001–1009.

Lister, B. C. 1976. The nature of niche expansion in West Indian Anolis lizards I: ecological

consequences of reduced competition. Evolution 30:659-676.

Losos, J. B. 2009. Lizards in an evolutionary tree: ecology and adaptive radiation of anoles.

University of California Press, Berkeley.

Losos, J. B., M. Leal, R. E. Glor, and K. D. Queiroz. 2003. Niche lability in the evolution of a

Caribbean lizard community. Nature 424:542–545.

Losos, J. B., and R. E. Ricklefs. 2009. Adaptation and diversification on islands. Nature

457:830-836.

M'Closkey, R. T. 1978. Niche separation and assembly in four species of Sonoran Desert

rodents. The American Naturalist 112:683-694.

MacArthur, R. H. 1965. Patterns of species diversity. Biological Reviews 40:510-533.

MacArthur, R. H. 1970. Species packing and competitive equilibrium for many species.

Theoretical Population Biology 1:1-11.

MacArthur, R., and R. Levins. 1967. The limiting similarity, convergence, and divergence of

coexisting species. The American Naturalist 101:377-385.

MacArthur, R. H., and J. W. MacArthur. 1961. On bird species diversity. Ecology 42:594-598.

Maestri, R., B. Patterson, R. Fornel, L. Monteiro, and T. De Freitas. 2016. Diet, bite force and

skull morphology in the generalist rodent morphotype. Journal of Evolutionary Biology

29:2191-2204.

Maestri, R., and B. D. Patterson. 2016. Patterns of species richness and turnover for the South

American rodent fauna. PLoS One 11:e0151895.

Martinez, Q., R. Lebrun, A. S. Achmadi, J. A. Esselstyn, A. R. Evans, L. R. Heaney, R. P.

Miguez, K. C. Rowe, and P.-H. Fabre. 2018. Convergent evolution of an extreme dietary

specialization, the olfactory system of worm-eating rodents. Scientific Reports 8:17806.

May, R. M., and R. H. Mac Arthur. 1972. Niche overlap as a function of environmental

variability. Proceedings of the National Academy of Sciences 69:1109-1113.

McElreath R. 2020 Statistical Rethinking v2. Taylor & Francis Group, Boca Raton.

Millien, V. 2006. Morphological Evolution Is Accelerated among Island Mammals. PLOS

Biology 4:e321.

Musser, G. G. 2014. A systematic review of Sulawesi Bunomys (Muridae, Murinae) with the

description of two new species. Bulletin of the American Museum of Natural History

392.

Nations, J. A., L. R. Heaney, T. C. Demos, A. S. Achmadi, K. C. Rowe, and J. A. Esselstyn.

2019. A simple skeletal measurement effectively predicts climbing behavior in a diverse

clade of small mammals. Biological Journal of the Linnean Society 128:323–336.

Nations, J. A., G. G. Mount, S. M. Morere, A. S. Achmadi, K. C. Rowe, and J. A. Esselstyn.

2021. Locomotory mode transitions alter phenotypic evolution and lineage diversification

in an ecologically rich clade of mammals. Evolution 75:376-393.

Naya, D. E., F. Bozinovic, and W. H. Karasov. 2008. Latitudinal trends in digestive flexibility:

testing the climatic variability hypothesis with data on the intestinal length of rodents.

The American Naturalist 172:122-134.

Newsome, S. D., C. Martinez del Rio, S. Bearhop, and D. L. Phillips. 2007. A niche for isotopic

ecology. Frontiers in Ecology and the Environment 5:429-436.

Nugraha, A. M. S. and Hall, R. 2018. Late cenozoic palaeogeography of Sulawesi, Indonesia.

Palaeogeography, Palaeoclimatology, Palaeoecology 490: 191–209.

Oksanen, J, F. Guillaume Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R.

Minchin, R. B. O'Hara, G. L. Simpson, P. Solymos, M. Henry, H. Stevens, E. Szoecs and

H. Wagner (2019). Vegan: Community Ecology Package. R package version 2.5-6.

Oliveira, B. F., and B. R. Scheffers. 2019. Vertical stratification influences global patterns of

biodiversity. Ecography 42:249-249.

Oliveira, B. F., J. M. Flenniken, R. P. Guralnick, S. E. Williams, and B. R. Scheffers. 2020.

Historical environmental stability drives discordant niche filling dynamics across

phylogenetic scales. Journal of Biogeography 47:807-816.

Peixoto, F. P., P. H. P. Braga, and P. Mendes. 2018. A synthesis of ecological and evolutionary

determinants of bat diversity across spatial scales. BMC ecology 18:1–14.

Pellissier, V., J. Y. Barnagaud, W. D. Kissling, Ç. Şekercioğlu, and J. C. Svenning. 2018. Niche

packing and expansion account for species richness–productivity relationships in global

bird assemblages. Global Ecology and Biogeography 27:604-615.

Peters, R. H. 1986. The Ecological Implications of Body Size. Cambridge University Press,

Cambridge.

Pianka, E. R. 1974. Niche overlap and diffuse competition. Proceedings of the National

Academy of Sciences 71:2141-2145.

Pianka, E. R., L. J. Vitt, N. Pelegrin, D. B. Fitzgerald, and K. O. Winemiller. 2017. Toward a

Periodic Table of Niches, or Exploring the Lizard Niche Hypervolume. The American

Naturalist 190:601-616.

Pigot, A. L., C. H. Trisos, and J. A. Tobias. 2016. Functional traits reveal the expansion and

packing of ecological niche space underlying an elevational diversity gradient in

passerine birds. Proceedings of the Royal Society B: Biological Sciences 283:20152013.

Pinto, G., D. L. Mahler, L. J. Harmon, and J. B. Losos. 2008. Testing the island effect in

adaptive radiation: rates and patterns of morphological diversification in Caribbean and

mainland Anolis lizards. Proceedings of the Royal Society B: Biological Sciences

275:2749–2757.

Powell, R. A., and C. M. King. 1997. Variation in body size, sexual dimorphism and age-specific

survival in stoats, Mustela erminea (Mammalia: Carnivora), with fluctuating food

supplies. Biological Journal of the Linnean Society 62:165-194.

Prevosti, F. J., G. F. Turazzini, M. D. Ercoli, and E. Hingst-Zaher. 2012. Mandible shape in

marsupial and placental carnivorous mammals: a morphological comparative study using

geometric morphometrics. Zoological Journal of the Linnean Society 164:836-855.

Revell, L. J. 2012. Phytools: an R package for phylogenetic comparative biology (and other

things). Methods in Ecology and Evolution 3:217-223.

Ricklefs, R. E., and J. Travis. 1980. A Morphological Approach to the Study of Avian

Community Organization. The Auk 97:321-338.

Rowe, K. C., A. S. Achmadi, and J. A. Esselstyn. 2014. Convergent evolution of aquatic

foraging in a new genus and species (Rodentia: Muridae) from Sulawesi Island,

Indonesia. Zootaxa 3815:541-564.

Rowe K.C., A.S. Achmadi, and J.A. Esselstyn. 2016a. Repeated evolution of carnivory among

Indo-Australian rodents. Evolution 70:653-665.

Rowe, K. C., A. S. Achmadi, and J. A. Esselstyn. 2016b. A new genus and species of

omnivorous rodent (Muridae: Murinae) from Sulawesi, nested within a clade of endemic

carnivores. Journal of Mammalogy 97:978-991.

Rowe, K. C., A. S. Achmadi, P. H. Fabre, J. J. Schenk, S. J. Steppan, and J. A. Esselstyn. 2019.

Oceanic islands of Wallacea as a source for dispersal and diversification of murine

rodents. Journal of Biogeography 46:2752-2768.

Rowsey, D. M., L. R. Heaney, and S. A. Jansa. 2019. Tempo and mode of mandibular shape and

size evolution reveal mixed support for incumbency effects in two clades of

island□endemic rodents (Muridae: Murinae). Evolution 73:1411-1427.

Rowsey, D. M., R. M. Keenan, and S. A. Jansa. 2020. Dietary morphology of two island-

endemic murid rodent clades is consistent with persistent, incumbent-imposed

competitive interactions. Proceedings of the Royal Society B 287:20192746.

Roughgarden, J. 1976. Resource partitioning among competing species–a coevolutionary

approach. Theoretical Population Biology 9:388-424.

Samonds, K. E., L. R. Godfrey, J. R. Ali, S. M. Goodman, M. Vences, M. R. Sutherland, M. T.

Irwin, and D. W. Krause. 2013. Imperfect Isolation: Factors and Filters Shaping

Madagascar's Extant Vertebrate Fauna. PLOS ONE 8:e62086.

Samuels, J. X. 2009. Cranial morphology and dietary habits of rodents. Zoological Journal of the

Linnean Society 156:864–888.

Samuels, J. X., and B. Van Valkenburgh. 2008. Skeletal indicators of locomotor adaptations in

living and extinct rodents. Journal of Morphology 269:1387-1411.

Scheffers, B. R., B. L. Phillips, W. F. Laurance, N. S. Sodhi, A. Diesmos, and S. E. Williams.

2013. Increasing arboreality with altitude: a novel biogeographic dimension. Proceedings

of the Royal Society B 280:20131581.

Schoener, T. W. 1974. Resource Partitioning in Ecological Communities: Research on how

similar species divide resources helps reveal the natural regulation of species diversity.

Science 185:27-39.

Sikes, R. S., and Animal Care and Use Committee of the American Society of Mammalogists.

2016. 2016 Guidelines of the American Society of Mammalogists for the use of wild

mammals in research and education. Journal of Mammalogy 97:663–688.

Spasojevic, M. J., and K. N. Suding. 2012. Inferring community assembly mechanisms from

functional diversity patterns: the importance of multiple assembly processes. Journal of

Ecology 100:652-661.

Stroud, J. T., and J. B. Losos. 2016. Ecological Opportunity and Adaptive Radiation. Annual

Review of Ecology, Evolution, and Systematics 47:507-532.

Stroud, J. T. 2021. Island species experience higher niche expansion and lower niche

conservatism during invasion. Proceedings of the National Academy of Sciences 118:

e2018949118.

Stuart, Y. E., T. S. Campbell, P. a. Hohenlohe, R. G. Reynolds, L. J. Revell, and J. B. Losos.

2014. Rapid evolution of a native species following invasion by a congener. Science

346:463-466.

Swenson, N. G. 2014. Functional and Phylogenetic Ecology in R. Springer, New York.

Swenson, N. G., and M. D. Weiser. 2014. On the packing and filling of functional space in

eastern North American tree assemblages. Ecography 37:1056–1062.

Swenson, N. G., M. D. Weiser, L. Mao, S. Normand, M. Á. Rodríguez, L. Lin, M. Cao, and J.-C.

Svenning. 2016. Constancy in functional space across a species richness anomaly. The

American Naturalist 187:E83-E92.

Tews, J., Brose, U., Grimm, V., Tielbörger, K., Wichmann, M. C., Schwager, M. and Jeltsch, F.

2004. Animal species diversity driven by habitat heterogeneity/diversity: the importance

of keystone structures. Journal of Biogeography 31: 79–92.

Van de Perre, F., M. R. Willig, S. J. Presley, I. J.-C. Mukinzi, M. S. Gambalemoke, H. Leirs, and

E. Verheyen. 2020. Functional volumes, niche packing and species richness:

biogeographic legacies in the Congo Basin. Royal Society Open Science 7:191582.

Vaughan, D, and M. Dancho 2021. furrr: Apply Mapping Functions in Parallel using Futures. R

package version 0.2.2.

Verde Arregoitia, L. D., D. O. Fisher, and M. Schweizer. 2017. Morphology captures diet and

locomotor types in rodents. Royal Society Open Science 4:160957.

Violle, C., B. J. Enquist, B. J. McGill, L. Jiang, C. H. Albert, C. Hulshof, V. Jung, and J.

Messier. 2012. The return of the variance: intraspecific variability in community ecology.

Trends in Ecology & Evolution 27:244-252.

Webb, C. O. 2000. Exploring the phylogenetic structure of ecological communities: an example

for rain forest trees. The American Naturalist 156:145-155.

Webb, C. O., D. D. Ackerly, M. A. McPeek, and M. J. Donoghue. 2002. Phylogenies and

community ecology. Annual Review of Ecology and Systematics 33:475-505.

Weiher, E., and P. A. Keddy. 1995. Assembly rules, null models, and trait dispersion: new

questions from old patterns. Oikos:159-164.

Wickham, H., M. Averick, J. Bryan, W. Chang, L. D. A. McGowan, R. François, G. Grolemund,

A. Hayes, L. Henry, and J. Hester. 2019. Welcome to the Tidyverse. Journal of Open

Source Software 4:1686.

Wilson, D. E., R. A. Mittermeier, and T. E. Lacher. 2019. Handbook of the Mammals of the

World. Lynx Editions, Barcelona.

Wilson, E. O. 1959. Adaptive shift and dispersal in a tropical ant fauna. Evolution 13:122-144.

Wilson, E. O. 1961. The Nature of the Taxon Cycle in the Melanesian Ant Fauna. The American

947   Naturalist 95:169-193.

948   Woodward, G., and A. G. Hildrew. 2002. Body□size determinants of niche overlap and

949   intraguild predation within a complex food web. Journal of Animal Ecology 71:1063-

950   1074.

951   Xu, X., D. Dimitrov, N. Shrestha, C. Rahbek, and Z. Wang. 2019. A consistent species richness–

952   climate relationship for oaks across the Northern Hemisphere. Global Ecology and

953   Biogeography 28:1051–1066.

965   **References Only Cited in the Online Supplement**

966   Achmadi, A. S. 2010. Taxonomic Status of Spiny Rats (Maxomys Jentink, Rodentia) from

967   Indonesia and Malaysia Based on Morphological Study. Treubia 37:49-82.

Fry, B. 2006. Stable isotope ecology. Springer, New York.

Hückstädt, L., J. Burns, P. Koch, B. McDonald, D. E. Crocker, and D. Costa. 2012. Diet of a

specialist in a changing environment: the crabeater seal along the western Antarctic

Peninsula. Marine Ecology Progress Series 455:287-301.

Kruschke, J. K. 2013. Bayesian estimation supersedes the t test. Journal of Experimental

Psychology: General 142:573.

McElreath R. 2020 Statistical Rethinking v2. Taylor & Francis Group. Boca Raton.

Musser, G. G., and M. D. Carleton. 2005. Superfamily Muroidea. Pages 894–1531 in D. E.

Wilson, Reeder D.M., editor. Mammal species of the world: a taxonomic and geographic

reference. Johns Hopkins Univ. Press, Baltimore.

Figure 1: Small mammal surveys of nine mountains a) on the oceanic island of Sulawesi revealed varying murine
rodent species richness across the island. Each mountain on the map is labeled with the maximum elevation and the
number of murine rodent species present. All but Latimojong were surveyed within 600m of the summit. b) The
diversity of Sulawesi murines is exceptional and includes unique forms that live alongside species with more
"typical" ecologies and morphologies, as demonstrated from surfaces of cranial μCT scans of *i – Rattus hoffmanni,* a

"typical" murine morphology and ecology, *ii* – the shrew rat *Echiothrix leucura* with its extremely elongate rostrum
and soft invertebrate diet, *iii* – the arboreal *Haeromys minahassae*, with a short rostrum and very small size. c)
Photographs: *i* – *Rattus hoffmanni*, *ii* –*Echiothrix leucura*, and *iii* – *Haeromys minahassae*.
989
990
991

Clade	Crown Age	Species Sampled
Echiothrix	5.11 (4.49-5.69)	10 (10)
Maxomys	3.72 (3.21-4.29)	5 (5)
Bunomys	3.45 (3.09-3.81)	15 (15)
Margaretamys	2.92 (2.29-3.51)	2 (4)
Rattus*	1.57 (1.22-1.95)	6 (6)
Haeromys[†]	Indeterminate	1 (1)

Table 1: Ages of six clades descended from Sulawesi colonists. Median age in millions of years is reported with
95% credible intervals in parentheses. Species sampled reports the number of species in this study with overall clade
richness in parentheses and demonstrates near complete sampling. The two unsampled *Margaretamys* species were
not detected in the localities in this study. All ages taken from Rowe et al. 2019. *There were likely two
colonizations by the ancestors of native *Rattus* spp., the second of which occurred 1.16-0.6 Ma (Rowe et al. 2019),
however, all *Rattus* in this study form a clade relative to other Sulawesi murines. Two human commensal *Rattus*
spp. found on Sulawesi were excluded. [†]The age of arrival of the Haeromys clade is unknown as other species of
Haeromys from Borneo have yet to be included in phylogenetic analyses.
1000

1001

Figure 2: The effect of species richness on community functional space volume (a, b) and density (c, d). X-axes
show the $\beta_{richness}$ estimates (the regression slope) for each trait space (y-axis) on the left and the Bayesian R^2 for

each $\beta_{richness}$ estimate on the right. Colored point intervals show 89% probability of $\beta_{richness}$ estimates (effect
size), with color varying by $\beta_{richness}$ value. Black point intervals show 50% posterior estimates of Bayesian R^2 for
each trait space. Density was estimated using the mean nearest neighbor (NN), and a high NN distance indicates low
density. Four of the five multivariate traits (**a**) and seven of the 12 individual traits (**b**), show an increase in
functional space volume (trait variance) with greater richness (i.e., positive $\beta_{richness}$). Phylogenetic diversity also
increases with species richness (**b**). All multivariate traits (**c**) and individual traits (**d**) show a stable or, surprisingly,
decreasing functional space density (increased NN distance) with greater species richness.

1024

Figure 3: Estimated functional volume SES plotted against species richness: **a**) Multivariate trait volumes, **b**) Skull traits, **c**) External measurements, and **d**) Isotopic measures, locomotor mode, and phylogenetic diversity. Points represent the mean SES values and error bars indicate the 89% credible intervals. X-axis tick marks show species richness for each community. Values equal to zero are consistent with null expectations, positive values indicate overdispersion, and negative values show underdispersion (trait clustering). Dotted lines depict the 89% interval of the null distribution. The plot of trait densities is shown in the supporting information (Figure S4).

1035

1036

Figure 4: The mode of functional space filling varies among traits. The bivariate plots on the left depict the 2D
functional spaces. Each black shape is the mean value of a particular species in each clade (shown in legend). The
convex hull colors match the color of the community functional volume SES distributions in the right-hand columns.
The thick convex hull lines are mean values, and 500 random samples from the posterior of each trait are shown in
thin hull lines. The number of species in each community and each clade used in this study is shown next to the SES
distributions in parentheses. All values were scaled to z-scores prior to analyses. **a**) Cranial morphospace values are
mostly clustered tightly near the mean of each axis, apart from Echiothrix species and the single *Haeromys* species.
Intraspecific variance is also relatively low on these axes. **b**) Intraspecific variance is high in isotopic niche space.
High and low δ^{15}N values are distributed among clades, but that is not the case for δ^{13}C values. **c**) Large values of
hind foot length belong to members of two clades, while large and small tail length values are dispersed among all
six clades, reducing the influence of phylogenetic diversity on tail length disparity.

1048
1049