MAIT cells and MR1 play an immunosuppressive role in glioblastoma through the induction of neutrophils and MDSCs

Taijun Hana and Masaki Terabe

Basic Immunology Section, Neuro-Oncology Branch, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA

Running title:

Immunosuppressive role of MAIT cell and MR1 in GBM

Corresponding:

Masaki Terabe, Ph.D, Basic Immunology Section, Neuro-Oncology Branch, National Cancer Institute, National Institutes of Health, Building 37, Room 1016, 37 Convent Drive, Bethesda, MD 20892, USA

Tel: 240-760-6731

E-mail: terabe@mail.nih.gov

Funding: This work was supported by the intramural program of the National Institutes of Health, National Cancer Institute, Center for Cancer Research (ZIA BC 011877).

Conflict of Interest: The authors have no conflicts of interest to declare.

24   **Authorship:** Conception and design: T.H, M.T. Data analysis: T.H. Manuscript writing: T.H.,

25   M.T. Final approval: All authors approved.

Abstract

Background: Major histocompatibility complex class I-related protein (MR1) is an antigen-presenting molecule for Mucosal-Associated Invariant T (MAIT) cells. Although the expression level of the *MR1* gene was reported to correlate with worse survival in glioma patients, a detailed mechanism of this association has not been elucidated. This study aimed to understand the mechanism by which *MR1* levels affect glioblastoma (GBM) patient survival and any potential involvement of MAIT cells in this mechanism.

Methods: Multi-layered gene expression analysis, including direct sequence alignment and search, were conducted on The Cancer Genome Atlas (TCGA)-GBM data. The immunological impact of MAIT cells in GBM was evaluated using multiple analytical methods, including statistical correlation analysis and Gene Set Enrichment Analysis. Pan-cancer TCGA data was utilized to evaluate the role of MAIT cells in other cancers.

Results: Intact *MR1* was expressed in a broad spectrum of cell types in GBM, and its expression level was inversely correlated with the survival of GBM patients. Significant positive correlations were found between the expression of *MR1*, the amount of MAIT cell infiltration, and the activity of tumor-associated neutrophils (TANs)/myeloid-derived suppressor cells (MDSCs) in GBM. Simultaneously, the activity of TANs/MDSCs was significantly inversely correlated with the survival of GBM patients. Similar associations between MAIT/MR1 and TANs/MDSCs were observed in four additional cancer types.

Conclusions: Our observations shed light on the possible immunosuppressive role of MAIT cells and MR1 in GBM. It was indicated that *MR1* up-regulation leads to activation of MAIT cells, which in turn induce the activation and infiltration of TANs/MDSCs. Our results suggest the existence of a novel immunosuppressive MAIT/MR1-TAN/MDSC pathway whose activation

contributes to the poorer prognosis of GBM patients and which necessitates novel strategies to

counteract this immunosuppressive role of MAIT cells.

Keywords: Glioblastoma, MAIT cell, MR1, neutrophil, MDSC, immunosuppression

Key points:

・ The expression level of *MR1* inversely correlates with the survival of GBM patients.

・ *MR1* expression levels and MAIT cell infiltration correlated.

・ MAIT cell and MR1 play immunosuppressive roles in GBM through neutrophil and MDSC.

Importance of the study

MR1-restricted MAIT cells are unconventional T cells that have the potential to regulate tumor

immunity. However, their immunobiology in GBM has not been understood. *MR1* expression

levels were reported to correlate with worse survival in glioma patients, while the involvement of

MAIT cells in the correlation was not clear. Using the TCGA-GBM data, we observed a fraction

of GBM patients have MAIT cells in the tumor and significant correlations between the expression

level of MR1, MAIT cell infiltration, and activities of TANs/MDSCs. Genes involved in multiple

TAN/MDSC-mediated immunosuppressive pathways were upregulated in MAIT-positive patients.

Importantly, both *MR1* expression levels and TAN/MDSC activation negatively correlated with

patient survival. The results suggest the existence of a novel immunosuppressive MAIT/MR1-

TAN/MDSC pathway whose activation contributes to the poorer prognosis of GBM patients, and

which provides novel strategies to counteract immune suppression in GBM patients.

Introduction

Glioblastoma (GBM) is the most devastating and common primary brain tumor in adults. Despite a multi-modal standard of care, which includes surgical resection followed by radichemotherapy, five-year survival remains around 10%[1], underscoring the need for better treatment. Immunotherapies, such as immune checkpoint blockers (ICB), have shown promise in treatments of other cancer types, but to date have not shown clinical efficacy in the treatment of GBM[2]. This lack of efficacy has been attributed to a highly immunosuppressive microenvironment, which includes a high content of immunosuppressive myeloid cells[3]. However, factors that activate or induce those immunosuppressive myeloid cells are still under investigation.

Numerous studies are underway to elucidate mechanisms of immunosuppression in GBM. Recently, it was reported that a transcriptomic expression level of major histocompatibility complex class I-related protein (MR1) significantly correlated to the poor survival of glioma patients[4]. However, the underlying mechanism by which the *MR1* expression level correlates with poor survival was not fully determined. MR1 is a non-classical major histocompatibility complex (MHC) class I protein and is the antigen-presenting molecule for Mucosal-Associated Invariant T (MAIT) cells. The only known function of MR1 is antigen presentation to MAIT cells. MR1 has been found in almost all mammals, and its genome sequence is highly conserved in many species, suggesting the importance of MR1 across the mammalian kingdom. The repertoire of antigens presented by MR1 is, however not well understood. It has been shown that riboflavin-derivatives, bacterial metabolites, are a major type of antigen presented to MAIT cells. However, it was recently reported that MR1 might load undefined cancer-derived antigens and present them to MR1-restricted T cells[5]. MAIT cells express an invariant T cell receptor (TCR) alpha chain, which utilizes Vα7.2Jα33 in humans[6]. Currently, several additional TCR gene segment combinations,

such as Vα7.2Jα12 and Vα7.2Jα20, have been reported for the invariant TCRα chain. Although

MAIT cells were discovered as a T cell population enriched in gut lamina propria, they are

prevalent in non-mucosal tissues in humans.

MAIT cells can secrete multiple cytokines upon their activation. They can secrete Th1-type

cytokines such as IFN-γ and TNF and Th17-type cytokines such as IL-17, IL-22, and IL-21. Their

cytokine profile can be modified depending on the environment they are in[7]. Although the

functions of each cytokine cannot be simply stated because of their complexity, the Th1-type

cytokines tend to produce immune responses needed for killing intracellular pathogens and cancer

cells. Th17-type cytokines induce immune responses against fungus infection. Among them, IL-

17 is known to promote the local attraction and activation of specific myeloid cells, such as

neutrophils, through the induction of chemokines[8,9].

The role of MAIT cells in cancer is uncertain, and there is no consensus on whether activated

MAIT cells are more likely to promote or inhibit cancer progression[7,10]. Since MAIT cells can also

secrete perforin and granzyme B, they can potentially lyse cancer cells. Some reports suggest that

MAIT cells can kill cancer cells including glioma-derived cells under certain *in vitro* conditions,

but their ability to target human cancers *in vivo* has not been confirmed[11,12]. On the other hand,

Yan, et al., using *in vivo* mouse tumor models, demonstrated that MAIT cells promote tumor

initiation, growth, and metastasis[13]. Some clinical studies reported an inverse correlation between

the number of tumor-infiltrating MAIT cells and survival in multiple types of cancer[14,15]. The

discrepancy in observed MAIT cell functions *in vitro* and *in vivo* or clinical conditions suggests

that the immunological functions of MAIT cells can be modified by their surrounding environment.

Interaction between GBM and MAIT cells has not been well studied. Peterfalvi, et al., using RT-

PCR, suggested the infiltration of MAIT cells in clinical GBM samples, but their clinical impact

or function in GBM remained to be elucidated[16].

We explored the roles of MR1 and MAIT cells in GBM by using large patient cohort data sets.

We found a significant correlation between *MR1* expression and GBM patients' poor survival and

significant correlations among *MR1*, MAIT cells, and the activity of tumor-associated neutrophils

(TANs)/myeloid-derived suppressor cells (MDSCs). Based on the findings, we propose a

hypothesis that *MR1* up-regulation with infiltration of MAIT cells in GBM activates

TANs/MDSCs that express multiple immunosuppressive mechanisms. The MAIT/MR1-

TAN/MDSC pathway can be a mechanism that contributes to the lack of effective response to

immunotherapy and provides the poor prognosis of GBM.

Results

***MR1* expression levels correlated with worse overall survival of GBM patients**

The Cancer Genome Atlas (TCGA)-GBM expression data showed that *MR1* expression levels

varied among the patient-derived GBM samples (Figure_1a). We then examined the correlation

between *MR1* expression levels and GBM patients' overall survival. We used data from TCGA

and the Chinese Glioma Genome Atlas (CGGA)[17] and observed a significant negative correlation

between *MR1* expression levels and GBM survival (Figure_1b), with higher expressions of *MR1*

being associated with worse survival, in agreement with the previously reported prognostic value

of *MR1* expression in GBM[4]. We also investigated whether the observed association could be

attributed to functional mutations in *MR1* including the recently-reported loss of antigen

presentation function of the substitution (*MR1*-R31H) in a human patient[18], but found a very low

frequency (0.06%) of genomic mutations in a range of glioma data sets including 3391 glioma

Figure_1: The *MR1* gene status and its prognostic correlation in GBM
(a) The RNA-expression data of TCGA-GBM 168 core samples were analyzed to see the expression levels of *MR1*. (b) The correlation between *MR1* expression and GBM patient survival. Clinical data available TCGA-GBM 153 patient data and Chinese Glioma Genome Atlas (CGGA) 85 primary GBM patient data were analyzed. Cut off FPKM: TCGA-MR1 = 3.67, CGGA-MR1 = 9.59, Log-rank test (c) The mutational schema of *MR1* locus in 3391 samples of glioma. Only 2 missense mutations were detected. (d) H3K27ac ChIP viewer image in *MR1* locus of GBM and high-grade glioma samples, GBM2w (IDHwt GBM, GSM1824806), AA15m (IDHmut-R132H, Grade3 Astrocytoma, GSM1824808). (e) Single-cell RNA-seq data of 3589 cells from GBM tissues of four patients were analyzed with a tSNE plot. All colored dots represent cells expressing *MR1*.

samples, and none of which had the reported loss of function mutation *MR1*-R31H (Figure_1c).

Additionally, we examined the epigenetic transcription status of *MR1* using available H3K27ac

ChIP data[19] of GBM and grade 3 astrocytoma, and found that the *MR1* promotor was active across

these samples (Figure_1d). Given the bulk tissue nature of TCGA samples, we also considered

whether the *MR1* expression in a subset of the samples could be due to the larger contribution of

micro-environment components in those samples. We utilized single-cell RNA-sequencing (RNA-

seq) data[20] and found that *MR1* was widely expressed in both glioma cells and tumor-infiltrating

leukocytes (TILs) in GBM(Figure_1e), and that *MR1* expression was relatively upregulated in the

tumor core compared to the tumor periphery.

All these observations indicated that intact *MR1* is widely transcribed in GBM tissue and

suggested that it is involved in the tumor pathogenesis of GBM.

The level of *MR1* expression correlates with the activity of TANs and MDSCs

To elucidate a potential mechanism to explain the correlation between high *MR1* expression

and poor prognosis, we extracted the top 500 significantly co-expressing genes (Supplementary

table_1) of *MR1* by using TCGA-GBM gene expression data. We further analyzed those 500 genes

for correlations with clinical outcomes[21]. Among the top 500 genes co-expressed with *MR1,* 158

genes significantly correlated with disease prognosis (Supplementary table_2) (Figure_2a). The

pathway analysis of the 158 genes showed strong enrichment ($q < 2.0 \times 10e\text{-}10$) of the neutrophil

degranulation pathway (R-HSA-6798695) (Figure_2b). Since neutrophil degranulation (ND)

results from neutrophil activation, this data indicated that intra-tumoral *MR1* up-regulation

correlates with neutrophil activation in GBM tissue, which could impact the clinical prognosis.

Thirty genes (Supplementary table_3) among the 158 genes were classified into the ND pathway,

and we call these 30 genes ND30. It is known that the biological characteristics of tumor infiltrated

neutrophils and blood circulating neutrophils are significantly different because the tumor

microenvironment modifies their activities[22]. Thus, tumor infiltrated neutrophils are typically

called tumor-associated neutrophils (TANs). Since we focused on tumor-derived transcriptomic

data, all neutrophils in this study will be called TANs. Recently, it was reported that ND pathway

genes were also up-regulated in activated myeloid-derived suppressor cells (MDSCs)[23]. Thus, the

observed up-regulation of the ND pathway can be an indication of MDSC activation as well.

Additionally, it is also known that TANs and MDSCs share many morphological, functional, and

transcriptomic characteristics and are difficult to distinguish from each other[24,25]. Therefore, the

activities of TANs and MDSCs were evaluated simultaneously in the current study.

Figure_2: Correlations of *MR1* expression and the activity of TANs/MDSCs
(a) A schematic illustration of the extracted top 500 significantly co-expressing genes with *MR1* and the prognosis related 158 genes among them. The horizontal axis is the Spearman's *r* values for each gene's correlation level with *MR1*, and the vertical axis is the *q*-value (Benjamini-Hochberg adjusted *p*-value) for each gene. (b) The enriched pathways of the 158 genes. The bar graph shows the enriched *q*-values for each pathway. (c) Correlation scatter plots between *MR1* gene expression, NDsig-index, and MDSCsig-index among the 168 TCGA-GBM samples.

To examine the correlation between the activities of TANs and MDSCs in GBM, we calculated

the ND signature gene index (NDsig-index, as the average fragments per kilobase of transcript per

million mapped fragments, FPKM, of the 479 genes of ND pathway signature, R-HSA-6798695,

Supplementary table_3) and MDSC-signature gene[26] index (MDSCsig-index, as the average

FPKM of the 80 MDSC-signature genes, Supplementary table_3) for each GBM sample.

Interestingly, those two scores among samples were highly correlated (Spearman's *r* = 0.889, *p* <

0.0001), and significant correlations between *MR1* and MDSCsig-index and *MR1* and NDsig-

index were also confirmed (Figure_2c). These data suggest that the *MR1* and the activity of

TANs/MDSCs in GBM samples are highly correlated, and the up-regulation of *MR1* possibly

induces the activation of TANs/MDSCs.

MAIT cell infiltrated GBM samples had a significantly higher level of *MR1* expression

Since MR1 is an antigen-presenting molecule for MAIT cells, presumably, there is no direct

interaction between MR1 and TANs/MDSCs. Thus, we hypothesized that MAIT cells are involved

in the MR1-TAN/MDSC pathway that negatively impacts GBM prognosis. One previous study

reported on the MAIT cell infiltration into GBM, but the results have not been verified by

multifaceted analysis[16]. To detect MAIT cell infiltration into GBM, we took advantage of the

expression of a unique invariant T cell receptor alpha chain, which utilizes a Vα7.2 (TRAV1-2)

gene segment combined with either a Jα33 (TRAJ33), a Jα12 (TRAJ12), or a Jα20 (TRAJ20) gene

segment, by MAIT cells. The raw-sequence FASTQ files of TCGA-GBM samples were

comprehensively searched to find the characteristic sequences for the gene segments of TRAV1-

2, TRAJ33, TRAJ12, and TRAJ20. The samples containing the reads of TRAV1-2 and one of

TRAJ33, TRAJ12, or TRAJ20 simultaneously were considered positive for MAIT cell infiltration.

Fourteen out of the 168 (8.3%) TCGA-GBM samples were identified as "MAIT-positive" samples

(Figure_3a, Supplementary table_4). We then evaluated the transcriptomic expression of the

MAIT cell surface receptors. Since CCR5, CCR6, CXCR6, CD161, IL-12R, and IL-18R are

known to be expressed in MAIT cells, we examined the expressions of these genes in comparison

with MAIT-positive/negative samples. All 6 genes had significant up-regulation in MAIT-positive

samples (Figure_3b). The gene expression of *IL17A*(IL-17), *TNF*, and *IFNG*(IFN-γ), which are

Figure_3: Multi-faceted evaluation of the MAIT infiltrated samples in TCGA-GBM
(a) The levels of MAIT-cell infiltration in TCGA-GBM samples. The 8.3% (14/168) of samples were detected as MAIT cell infiltration positive samples. The infiltration level was calculated with the formula, [The detected read count of the MAIT-target read of the sample/ The total RNA read count of the sample]. (b)(c) Violin plots showing the gene expression level of MAIT cell surface receptors/secreting cytokines. (d) The information on MAIT cell infiltration was integrated with other clinical/molecular indicators. Molecular indicators' information was obtained from a previously reported database[27], and the neural samples were integrated into the proneural samples. (e) The ratio-bar chart shows the distribution of GBM subtypes among MAIT-positive samples and the total TCGA-GBM samples. (f) A violin plot showing the *MR1* gene expression level in the MAIT positive/negative groups.

typical cytokines secreted by MAIT cells, was also examined. Significant up-regulation of *IL17A*

and *TNF* in MAIT-positive samples was observed. However, no significant difference in *IFNG*

expression was detected (Figure_3c). These data support our conclusion that samples containing

transcripts of the invariant TCRα gene segment were infiltrated by MAIT cells. The cytokine gene

expression data also raised the possibility that MAIT cells in GBM tissue may have a defect in

producing IFN-γ.

The attributes of the MAIT-positive/negative samples were clarified as follows. Although there

was no isocitrate dehydrogenase (IDH) mutant case in MAIT-positive samples, there was no

statistically significant deviation in the distribution of the IDH mutant cases among the MAIT-

positive/negative samples (Binomial test, $p = 1$). Other indicators also did not differentiate the

MAIT-positive sample, and the MAIT-positive samples were statistically evenly distributed in

multiple GBM subtypes (Figure_3d, e). These findings indicate that MAIT cell infiltration is not

significantly associated with conventional attributes. We then examined the *MR1* expression status

of MAIT-positive samples and found that MAIT-positive samples had a significantly higher

expression level of *MR1* than MAIT-negative samples (Figure_3f). This important finding

indicates that the *MR1* up-regulation correlates with the increased MAIT cell infiltration into GBM

tissue.

Multiple pathways of TANs/MDSCs contributed to the immunosuppression in MAIT-

positive samples

To further understand the transcriptomic differences between MAIT positive and negative

samples, we conducted a comprehensive Gene Set Enrichment Analysis (GSEA) using Gene

Ontology Biological Process (GOBP) signatures. Interestingly, the most significantly enriched

process among all GOBP-processes, including non-immunological processes in MAIT-positive

samples, was an immunological process, the negative regulation of immune response

(GO:0050777) (Figure_4a). Since the negative regulation of immune response is an integrated

pathway including many immunosuppressive processes, this significant enrichment suggests that

multiple immunosuppressive mechanisms are more active in MAIT-positive samples than in

MAIT-negative samples. As described above, we observed that the *MR1* expression level

correlates with TAN/MDSC activation levels (Figure_2c). Thus, we extracted the neutrophil-

related pathways from all the pathways enriched in MAIT-positive samples. Many neutrophil

migration/chemotaxis and neutrophil activation-related pathways were significantly enriched in

MAIT-positive samples (Figure_4b). Since GOBP does not have appropriate MDSC-related

pathways, we could not examine the MDSC.

We next examined the TAN/MDSC activities in MAIT-positive/negative samples. The

expression levels of ND30 genes in MAIT-positive samples were clearly up-regulated compared

with in MAIT-negative samples (Figure_4c). We also calculated an "ND30-index", the mean

FPKM of the ND30 genes for each GBM sample, and the ND30-index showed a significantly

higher score in the MAIT-positive samples (Figure_4d). GSEA, using the signature genes of ND

pathway and the gene set of MDSC signature, showed significant enrichment of those signatures

in MAIT positive samples (Figure_4e). These data suggested that TANs/MDSCs are highly

activated in MAIT-positive samples compared with MAIT-negative samples.

TANs/MDSCs are known to suppress cancer immunity through various mechanisms[28-30]. For

instance, the up-regulation of arginase 1 (ARG1) expression depletes the local arginine[31], and the

low arginine level impairs T cells by down-regulating CD3 ζ chains, which are necessary for a

TCR signalling[32]. One of the other immunosuppressive mechanisms of TANs/MDSCs is through

the induction of Foxp3$^+$CD4$^+$ regulatory T cells (Treg). Tregs are attracted by CCL17 and CCL22

and induced by TGF-β[33]. CCL17 and TGF-β can be secreted by TANs[34,35] and MDSCs[36]. Although

TANs/MDSCs do not secrete CCL22 directly, they can contribute to its local up-regulation

through M2 macrophage polarization[37,38]. TANs/MDSCs also express PD-L1[39], and the up-

regulation of indoleamine 2,3-dioxygenase (IDO) depletes the local tryptophan and produces

kynurenine that inhibits T cell proliferation. The IL-10 and TGF-β produced by TANs/MDSCs

suppress T cell proliferation/activity as well[40-42]. We examined the gene expressions of those

molecules and found that all the genes were significantly up-regulated in MAIT-positive samples

compared with MAIT-negative samples (Figure_4f). We observed the up-regulation of *ARG1* and

the arginine reporter gene, *ASS1* (up-regulated upon the local arginine depletion), in MAIT-

positive samples (Figure_4g). Notably, *CCL17* in the MAIT-positive samples had an

approximately 10 times higher expression level on average compared with the MAIT-negative

samples. Increased Treg infiltration in MAIT-positive samples was indicated by a significant

increase of *FOXP3,* a typical Treg marker gene, in MAIT-positive samples (Figure_4h). CCR4, a

chemokine receptor expressed on Treg, which contributes to Treg-chemotaxis by binding to

CCL17, was also highly up-regulated in MAIT-positive samples (Figure_4h). Increased Treg

infiltration in MAIT positive samples was further confirmed using previously reported 22 Treg

signature genes[43] in GSEA（Supplementary table_3）(Figure_4i). These data suggested that the

multiple immunosuppressive pathways mediated by TANs/MDSCs were active in the MAIT-

positive samples. We then evaluated the impact of TANs/MDSCs on the GBM clinical course

(Figure_4j). Significant negative correlations of TAN/MDSC activity with the overall survival

indicated that their activities have an unfavorable impact on the GBM malignancy.

Altogether, the data strongly suggested that MAIT cells activated by *MR1* up-regulation induce

the activation and infiltration of TANs/MDSCs, which exert multiple immunosuppressive

mechanisms in GBM.

Figure_4: Evaluations of the activity of TANs/MDSCs in MAIT-positive/negative samples
(a) The enrichment plot of GSEA-GOBP's negative regulation of immune response in MAIT-positive/negative samples. (b) The enrichment results of neutrophil-related pathways in GSEA-GOBP in MAIT-positive samples. All pathways shown were significantly enriched ($q < 0.05$). (c) Heatmap of the gene expression level of ND30 genes for each sample. The vertical column represents one GBM sample. MAIT-positive samples were gathered on the right. (d) A Violin plot showing the ND30-index level. (e) GSEA enrichment plots of neutrophil degranulation pathway and MDSC signature. (f) Scatter chart showing the differential expression levels of the immunosuppressive pathway genes related to TANs/MDSCs among MAIT-positive/negative samples. The horizontal axis shows the fold change (log2 converted) as follows. [Mean FPKM of MAIT-positive samples / Mean FPKM of MAIT-negative samples] The vertical axis shows the p-value (-log2 converted) of the unpaired t-test that compares the expression levels of MAIT-positive/negative samples for the gene. (g) Violin plots showing the *ASS1* or *ARG1* gene expression level. (h) Violin plots showing the expression level of *FOXP3* and *CCR4*. (i) GSEA enrichment plot, using our Treg-signature gene set. (j) The prognostic correlation of NDsig-index

and MDSCsig-index with TCGA-GBM clinical data. Cut off: NDsig-index = 41.89, MDSCsig-index = 27.48, Log-rank test

Pan-cancer analysis showed correlations between the activity of TANs/MDSCs and the

***MR1*'s prognostic correlation**

The results obtained by GBM gene expression data sets strongly suggested the existence of a

novel MAIT/MR1-TAN/MDSC pathway that potentially suppresses tumor immune responses. We

next asked if the correlations found in GBM can also be found in other types of cancer. In the

current study, we could not access to enough FASTQ raw data files to examine the MAIT cell

infiltration in different cancer types. Thus, we focused on investigating the relationship between

MR1 expression and the prognosis of 14 major cancer types（breast cancer, cervical cancer,

colorectal cancer, head and neck cancer, liver cancer, lung cancer, melanoma, ovarian cancer,

pancreatic cancer, prostate cancer, renal cancer, stomach cancer, testis cancer, thyroid cancer）

using TCGA transcriptomic data for each cancer types. In 5 cancer types（colorectal cancer, head

and neck cancer, pancreatic cancer, stomach cancer, and testis cancer), high expression of *MR1*

was correlated with poor survival, whereas in 2 cancer types（lung cancer, renal cancer）the

relationship was reversed with expression levels of *MR1* positively correlated with survival

(Figure_5a). The remaining 7 cancer types had no significant difference in correlation with

prognosis. We call these 5 cancer types that *MR1* correlates with worse survival "MR1-

unfavorable cancers" and call the 2 cancer types that *MR1* correlates with better survival "MR1-

favorable cancers". These results suggested that the intra-tumoral biological pathways associated

with MR1 may be different among cancer types. As we evaluated previously, we examined the

transcriptomic expression of the MAIT cell surface receptor genes in the MR1-high/low groups of

these MR1-unfavorable/favorable cancers. In all seven cancer types, most of the MAIT-surface

genes were significantly up-regulated in the MR1-high group (Figure_S1). These findings suggest

that the increased level of *MR1* expression increased the amount of infiltrating MAIT cells.

297   We then asked if TANs/MDSCs are involved in the *MR1*-involving mechanism that

significantly correlates with the disease prognosis. We performed GSEA using the customized

signature gene set of NDsig genes and MDSCsig genes (Supplementary table_3) and found that 4

cancer types (colorectal cancer, pancreatic cancer, stomach cancer, and testis cancer) in MR1-

unfavorable cancers had significant (FDR-*q* < 0.25) or strong trend in up-regulation of both ND

and MDSC signatures in the MR1-high group (Figure_5b). On the other hand, the signatures were

not correlated with *MR1* expression levels in all MR1-favorable cancers (renal, and lung cancer)

and one MR1-unfavorable cancer (head and neck) (detailed results are in Supplementary table_5).

We confirmed this observation by using other signatures of GSEA, which was performed using

the GOBP signature, and all neutrophil-related pathways that were significantly enriched in each

MR1 high group were extracted. The 4 cancer types, which had significant up-regulation of

TAN/MDSC activities along with *MR1* up-regulation, showed significant up-regulation of

multiple neutrophil-related pathways, such as positive regulation of neutrophil migration,

neutrophil extravasation, neutrophil activation involved in immune response, regulation of

neutrophil chemotaxis, neutrophil-mediated immunity, in MR1-high group (Figure_5c). In

contrast, head and neck, renal, and lung cancers indicated no significant up-regulation of

neutrophil-related pathways in the MR1-high or low group. These results well supported the

hypothesis that the MAIT/MR1-TAN/MDSC pathway is active in MR1-unfavorable cancers. It is

important to note that MAIT/MR1 did not correlate with TAN/MDSC activities in MR1-favorable

cancer types.

Figure_5: Correlations between *MR1*, prognosis, and TANs/MDSCs in the pan-cancer analysis

(a) The prognostic correlation of *MR1* in multiple cancer types. All red lines represent the MR1-high group, and blue lines represent the MR1-low group. (b) Enrichment plots of the GSEA using neutrophil degranulation pathway and our MDSCsig gene set for multiple cancer types. MR1-high/low groups were compared. (c) The enrichment results of neutrophil related pathways in GSEA-GOBP in the MR1-high group. All pathways shown were significantly enriched (*q* < 0.05).

Discussion

In this study, we proposed a novel MAIT/MR1-TAN/MDSC pathway as a potential mechanism for a negative impact of *MR1* expression on GBM prognosis. To our knowledge, this is the first study revealing the function of MAIT cells in glioblastoma. The transcriptional data of TCGA-GBM revealed that high expression of *MR1* was significantly correlated with GBM's poor survival. This observation was consistent with a previous report and was also confirmed by the CGGA database[4]. Comprehensive gene expression data analysis and the direct search of the raw sequence data of TCGA-GBM showed that 8.3% of the GBM samples had MAIT cell infiltration, and the intra-tumoral *MR1* expression significantly correlated with the infiltration level of MAIT cells. The GSEA, comparing the MAIT-positive/negative samples, showed that tumor immunity in MAIT-positive samples was significantly negatively regulated, supporting an immunosuppressive role of MAIT cells. Further analysis showed that the MAIT cell infiltration was significantly correlated with the activity of the immunosuppressive cell population, TANs/MDSCs. Simultaneously, TAN/MDSC activity was significantly correlated to the poor survival of GBM. These data suggested that *MR1* up-regulation activates MAIT cells that negatively impact the GBM prognosis through the activation of TANs/MDSCs. The pan-cancer analysis also strongly suggested the activity of the MAIT/MR1-TAN/MDSC pathway in 4 additional cancer types.

There are ongoing debates about whether the MAIT cell is an ally or an enemy of cancer immunity[10]. A few reports demonstrated that MAIT cells can kill cancer cells including glioma-derived cells under limited experimental conditions[11,12]. However, this finding does not necessarily mean that they can kill cancer cells *in vivo*. No studies based on clinical data or *in vivo* experiments provided evidence of the tumor-suppressive ability of MAIT cells. On the contrary, their tumor-

promoting ability has been suggested by multiple reports based on clinical data and *in vivo*

models[13-15]. Duan, et al. showed the possibility that tumor-infiltrating MAIT cells are

reprogrammed in the tumor milieu to produce IL-8 and suppress the ability to produce IFN-γ and

IL-17[14]. Another study reported that tumor-infiltrating MAIT cells enhanced the production of IL-

17 and TNF-α while an IFN-γ production level was not changed[13]. In the current study, the

expression levels of *IL17A* and *TNF* were significantly higher in MAIT-positive samples while

there was no statistical difference in *IFNG* expression between MAIT-positive and negative

samples. Although details in altered functions of MAIT cells observed are different among studies,

it seems that a cytokine profile of MAIT cells in cancer-bearing individuals is skewed away from

type I immune response. The alteration of the IFN-γ producing function of MAIT cells seems to

be like what has been reported on another unconventional T cell, natural killer T (NKT) cells.

Dhodapkar M, et al. reported that the IFN-γ secreting ability of NKT cells circulating in

progressive cancer patients was suppressed[44].

Importantly, in renal and lung cancer, in which *MR1* expression levels correlated with better

prognosis, there was no correlation between *MR1* expression levels and the signatures of TANs or

MDSCs. This was in striking contrast to five cancer types including GBM, in which the disease

prognosis had a negative correlation with *MR1* expression levels. These data suggest that MAIT

cells have multiple faces in the context of the regulation of tumor immunity. Only when the

interplay between MAIT cells and TANs/MDSCs occurs do MAIT cells have a negative impact

on disease outcomes. There may be multiple factors that contribute to changing the roles of MAIT

cells among different cancer types. One factor may be the tumor microenvironment as each cancer

type has a different tumor microenvironment (TME) associated with MR1 and MAIT cells. Since

the immunological phenotype of MAIT cells can change depending on the environment[13,14], the

inter-cancer-type difference in TME possibly explains the different observations about

MAIT/MR1-TAN/MDSC pathway in multiple cancer types. MR1-favorable cancers, such as lung

cancer and renal cancer, may have a TME that does not guide MAIT cells towards the phenotype

that can affect TANs/MDSCs through the secretion of IL-17. TME may include antigens available

for MR1 to present to MAIT cells. Although it has been shown that vitamin B metabolites derived

from commensal bacteria are recognized by MAIT cells, there are yet to be characterized cancer-

derived antigens[5].

The level of TAN infiltration has been reported to correlate with the grades of glioma, and

some tumor-promoting mechanisms by TANs in GBM have been suggested[45-47]. However, the

mechanisms attracting TANs to the GBM tissue, and their activation mechanism are not well

understood. Our study observed a significant correlation between the amount of MAIT cell

infiltration and the activation level of TANs/MDSCs. MAIT cells rapidly secrete a large amount

of cytokines including IL-17[48], which is important in this context as it is well established that IL-

17 can promote the local attraction and activation of TANs through the induction of chemokines[8,9].

In fact, our data showed the significant up-regulation of *IL17A* in MAIT-positive samples, and not

only the activation of TANs/MDSCs but also multiple neutrophil attraction pathways were up-

regulated in MAIT-positive samples. Since neutrophils can exert strong cytotoxicity, they may

play anti-tumor roles. However, the biological function of neutrophils can be modified depending

on the tissue environment, especially by the local TGF-β status. Under a TGF-β depleted

environment, neutrophils tend to acquire the anti-tumor phenotype, called N1; whereas, under the

TGF-β sufficient environment, they tend to acquire the pro-tumor phenotype, called N2[29]. The

immunosuppressive and tumor-promoting role of N2-TAN has been reported in multiple cancer

types[29,30]. Since the expression level of *TGFB1* in MAIT-positive samples was significantly up-

regulated (Figure 4f), the TME of MAIT-positive patients is the preferred environment for TANs

to acquire the pro-tumor phenotype, N2. Thus, the TANs in MAIT-positive GBM samples are

considered to play a relatively immunosuppressive role and can be a cause of GBM's poor

prognosis.

MDSCs suppress cancer immunity[28] and in GBM, their immunosuppressive role and the

correlation between their infiltration and poor survival was reported[49]. He D, et al. showed that

MDSCs have phenotypic plasticity in their immunosuppressive function using *in vivo* cancer

model and IL-17 was required for the development of MDSCs in tumor-bearing mice and for

exerting MDSC's pro-tumor functions[50]. Guan X, et al. also observed a significant correlation

between the tumor IL-17 level and the MDSC infiltration level into renal cancer[51]. These reports

support our hypothesis that the IL-17 secreted from MAIT cells can attract MDSCs and promote

their immunosuppressive function in GBM.

We identified the samples with MAIT cell infiltration in TCGA-GBM by directly searching

TCR transcripts in the FASTQ files of their bulk RNA-seq. Only one previous report suggested

the presence of MAIT cells in GBM by detecting TCR transcripts by PCR with a limited number

of samples[16]. The present study is the first to use next-generation sequencer data for the purpose.

As a result, 8.3% of the samples were determined to have MAIT cell infiltration. Strictly, the

interpretation of this result must consider the limitation of bulk RNA-seq's resolution. Generally,

more than one hundred million cells can be included in a tumor sample submitted for bulk RNA-

seq[52], while the average total read (total of paired-end reads) per sample of TCGA-GBM was

approximately five hundred million reads. Hence on average, less than 5 reads for each cell may

be included in the bulk RNA-seq data, and only highly contained transcripts are detected. The

MAIT-positive samples we found should be considered a particularly high amount of MAIT-

infiltrated samples, and therefore it is likely that the method we used underestimated the number

of samples containing MAIT cells.

Figure_6: The MAIT/MR1-TAN/MDSC pathway
The activation of MAIT cell, along with the *MR1* up-regulation, activates and chemo-attracts granulocytes, such as TANs/MDSCs, through IL-17 and TNF production. The activated TANs/MDSCs induce Tregs via the production of TGF-β, CCL17, and CCL22, and Tregs suppress the T cell-mediated immunity. Simultaneously, TANs/MDSCs suppress T cells via multiple pathways. The suppression of T cell immunity provides a poor prognosis for GBM.

This study analyzed the clinical molecular data related to GBM, MR1, and MAIT cells based

on significant correlations in the context of disease outcomes. Hence, it is impossible to

definitively prove the direction of the causal relationship. For example, the correlation between

MAIT cells and TANs/MDSCs can also be interpreted as that TANs/MDSCs activated MAIT cells.

This point is one limitation of the current study. Although we proposed the mechanism that MAIT

cells can activate TANs/MDSCs via IL-17, there is the potential of some reverse pathways.

Another limitation of our study is that although most analyses were performed using transcriptomic

data for each cancer type, the transcriptomic levels do not always agree with the protein levels and

their functional status. MR1 is known to be expressed on the cell surface with the presence of

antigens. However, it does not function as antigen-presenting molecule when it is not loaded with

antigens even with their transcriptomic expression[53]. Therefore, ideally, comprehensive proteomic

data of a large GBM dataset should be analyzed for the purpose. However, such comprehensive

proteomics data for GBM is not currently available, thus we have leveraged the available data for

the analysis.

In conclusion, we propose a novel MAIT/MR1-TAN/MDSC pathway that can contribute to

the poor prognosis of GBM (Figure_6). MAIT cells can be a missing link between GBM and

TAN/MDSC's activation and infiltration, which was previously reported. Our study focused on *in*

silico analysis using cancer big data. Further research using experimental methods is needed to test

the hypothesis. However, our observations shed light on the possible immunosuppressive role of

MR1 and MAIT cells in GBM.

Material and methods

The definition of glioblastoma (GBM) in this study

The WHO's central nervous tumor classification was revised in 2021, and IDH mutant grade 4

astrocytoma was excluded from the previously defined "glioblastoma"[54]. However, after

examining various data, we determined that it was not essential to divide the grade 4 gliomas into

multiple subtypes along with our study of MR1 and MAIT cells. We considered that the role of

MR1 and MAIT cells in glioma could be a consistent phenomenon in high-grade gliomas.

Therefore, our definition of GBM in this study is in line with the previous (WHO-2016) definition

of WHO. The disease concept is along with the TCGA-GBM dataset, in which GBMs are all the

grade 4 gliomas of IDH mutant and wild type.

Identification of MR1 mutational frequency, the epigenetic transcriptional status, the

transcriptional distribution in GBM tissue

The mutational frequency was confirmed using cBioportal[55]. A total of 3391 multi-grade glioma

samples were examined. The cBioportal's datasets of Glioblastoma (TCGA, Cell 2013), Low-

Grade Gliomas (UCSF, Science 2014), Merged Cohort of LGG and GBM (TCGA, Cell 2016),

Anaplastic Oligodendroglioma and Anaplastic Oligoastrocytoma (MSKCC, Neuro Oncol 2017),

Glioma (MSK, Nature 2019), Glioma (MSKCC, Clin Cancer Res 2019), Brain Lower Grade

Glioma (TCGA, PanCancer Atlas) were used. The H3K27ac ChIP data of GBM2w (IDHwt GBM,

GSM1824806), AA15m (IDHmut-R132H, Grade3 Astrocytoma, GSM1824808) were used to

examine the epigenetic transcriptional status of *MR1* in glioma. IgV_2.11.9 was used to visualize

the Bed.file data. Single-cell RNA-seq analyzing web tool, GBM-seq[20], was used to examine the

MR1's transcriptional distribution in GBM tissue. The generated tSNE plot of *MR1* expression

analysis was used.

The TCGA gene expression analysis and the prognostic examination

The TCGA-GBM core 168 samples'(Supplementary table_4) gene expression data was

downloaded from Genomic Data Commons Data Portal（https://portal.gdc.cancer.gov/）. All the

gene expression analyses were conducted using this set of data. Prognostic examinations with

specific gene expression in this study were done using the Log-rank test, and all cut-off thresholds

were decided using TCGA survival data analyzer[21,56]. TCGA prognostic analyses were conducted

using the TCGA-GBM 153 samples (Supplementary table_6) with available clinical data[21].

Chinese Glioma Genome Atlas (CGGA) data analysis was performed using the CGGA data

portal[17]. The dataset of "mRNAseq_325" was used with the order of *MR1* & survival analysis

option. The result of WHO grade IV survival (primary glioma) was used. When comparing specific

gene expressions between two groups, a two-tailed unpaired Student's t-test was used to analyze,

and $p < 0.05$ was considered statistically significant. When fold changes in gene expression were

calculated, the mean FPKM of both groups was used for the calculation. For pan-cancer analysis,

the TCGA expression and clinical data for all cancer types were obtained from The Human Protein

Atlas[21]. All analytical methods used are the same as GBM. All gene expression data analyzed in

this study was performed using an FPKM unit.

Extracting the MR1 co-expressing genes and gene set pathway analysis

The *MR1* co-expressing genes were extracted with cBioportal using the RNA Seq V2 RSEM data

set of Glioblastoma Multiforme (TCGA, PanCancer Atlas, Cell 2018). "Co-expression" analysis

mode with "MR1" was conducted, and the top 500 co-expression genes were extracted. All the co-

expression or correlation analyses of this study were performed using Spearman's correlation test

using GraphPad Prism8(GraphPad Software, La Jolla, California, USA), and $p < 0.05$ was used as

the significant threshold. To extract the GBM prognosis-related genes from the top 500 *MR1* co-

expression genes, the GBM prognostic gene database[21] was utilized. The 158 genes significantly

correlating the poor survival of GBM ($p < 0.05$) were extracted (Supplementary table_2). Gene set

pathway analyses of this study were conducted with g:Profiler[57].

Analyses using the signature of Neutrophil degranulation, MDSC, and Treg

The gene set of Neutrophil degranulation (REACTOME: R-HSA-6798695, 479 genes) was used

to generate the NDsig-index for each TCGA sample. All FPKM values of the 479 genes were

averaged for each sample, and the value was defined as NDsig-index for each sample. The MDSC

characteristic signature gene set was obtained from a previous study[26]. Since the gene set was

originally composed of 105 murine genes, we extracted 80 genes from them that exist in humans,

and the 80 genes were defined as MDSC signature genes (Supplementary table_3). The MDSCsig-

index was calculated similarly to the NDsig-index using the 80 MDSC signature genes. The

signature gene set for Treg was obtained from a previous study[43]. The 22 signature genes

differentially expressed in Tregs were used for our analysis (Supplementary table_3).

Detecting the MAIT-positive samples in TCGA-GBM

All FASTQ files of 168 TCGA-GBM core samples were downloaded from the Genomic Data

Commons Data Portal Legacy Archive（https://portal.gdc.cancer.gov/legacy-archive/）. Access

to TCGA-controlled data and studies using them were approved by the National Cancer Institute

(#28406). Reads having the characteristic T cell receptor sequences (Supplementary table_7) of

MAIT cells were searched with the FASTQ files. It was confirmed that all the searched sequences

were unique in the whole genome, and no other loci have the same sequence using the GGGenome-

ultrafast sequence search (https://gggenome.dbcls.jp/en/). The samples containing the reads of

TRAV1-2 and TRAJ33 or TRAJ12 or TRAJ20 simultaneously were classified as the samples with

the MAIT cell infiltration and were defined as MAIT-positive. The MAIT-infiltration level of each

sample was calculated with the formula, [The detected read count of the TRAV1-2 read of the

sample/ The total RNA read count of the sample]. All these analyses were conducted on the NIH-

high performance computer, the Biowulf cluster system.

Gene expression heatmap and Gene Set Enrichment Analysis (GSEA)

The FPKM values of ND30 genes for each sample were extracted to draw the heatmap of

Figure_4c. Intergenic normalization was performed with GraphPad Prism8(GraphPad Software,

La Jolla, California, USA), and each cell was colored along with the expression levels. GSEAs of

this study were conducted using the expression data of core 168 TCGA-GBM samples and the

pan-cancer samples. GSEA_4.2.3 was used to run the 1000 permutations analyses using the

signatures   of   c2.cp.reactome.v7.5.1.symbols.gmt,   c5.go.bp.v7.5.1.symbols.gmt,   and

c2.cp.kegg.v7.5.1.symbols.gmt. GSEA with the signature of ND, MDSC, and Treg was conducted

using our customized "gmx" signature files made from their characteristic signatures.

Acknowledgments

We thank Dr. Mark R. Gilbert and Dr. Orieta Celiku for the critical reading of the manuscript and

the NIH HPC group for providing the Biowulf cluster system. This work was supported by the

intramural program of the National Institutes of Health, National Cancer Institute, Center for

Cancer Research (ZIA BC 011877).

References

1. Stupp R, Taillibert S, Kanner A, et al. Effect of Tumor-Treating Fields Plus Maintenance Temozolomide vs Maintenance Temozolomide Alone on Survival in Patients With Glioblastoma: A Randomized Clinical Trial. *Jama.* 2017; 318(23):2306-2316.

2. Omuro A, Vlahovic G, Lim M, et al. Nivolumab with or without ipilimumab in patients with recurrent glioblastoma: results from exploratory phase I cohorts of CheckMate 143. *Neuro-oncology.* 2018; 20(5):674-686.

3. Klemm F, Maas RR, Bowman RL, et al. Interrogation of the Microenvironmental Landscape in Brain Tumors Reveals Disease-Specific Alterations of Immune Cells. *Cell.* 2020; 181(7):1643-1660 e1617.

4. Kubica P, Lara-Velazquez M, Bam M, et al. MR1 overexpression correlates with poor clinical prognosis in glioma patients. *Neuro-Oncology Advances.* 2021.

5. Crowther MD, Dolton G, Legut M, et al. Genome-wide CRISPR-Cas9 screening reveals ubiquitous T cell cancer targeting via the monomorphic MHC class I-related protein MR1. *Nature immunology.* 2020; 21(2):178-185.

6. Treiner E, Duban L, Bahram S, et al. Selection of evolutionarily conserved mucosal-associated invariant T cells by MR1. *Nature.* 2003; 422(6928):164-169.

7. Cogswell DT, Gapin L, Tobin HM, McCarter MD, Tobin RP. MAIT Cells: Partners or Enemies in Cancer Immunotherapy? *Cancers.* 2021; 13(7).

8. Parajuli P, Mittal S. Role of IL-17 in Glioma Progression. *J Spine Neurosurg.* 2013; Suppl 1.

9. Wu L, Awaji M, Saxena S, Varney ML, Sharma B, Singh RK. IL-17-CXC Chemokine Receptor 2 Axis Facilitates Breast Cancer Progression by Up-Regulating Neutrophil Recruitment. *The American journal of pathology.* 2020; 190(1):222-233.

10. Haeryfar SMM, Shaler CR, Rudak PT. Mucosa-associated invariant T cells in malignancies: a faithful friend or formidable foe? *Cancer Immunol Immunother.* 2018; 67(12):1885-1896.

11. Gherardin NA, Loh L, Admojo L, et al. Enumeration, functional responses and cytotoxic capacity of MAIT cells in newly diagnosed and relapsed multiple myeloma. *Scientific reports.* 2018; 8(1):4159.

12. Priya R, Brutkiewicz RR. MR1 Tetramer-Based Artificial APCs Expand MAIT Cells from Human Peripheral Blood That Effectively Kill Glioblastoma Cells. *Immunohorizons.* 2021; 5(6):500-511.

13. Yan J, Allen S, McDonald E, et al. MAIT Cells Promote Tumor Initiation, Growth, and Metastases via Tumor MR1. *Cancer discovery.* 2020; 10(1):124-141.

14. Duan M, Goswami S, Shi JY, et al. Activated and Exhausted MAIT Cells Foster Disease Progression and Indicate Poor Outcome in Hepatocellular Carcinoma. *Clinical cancer research : an official journal of the American Association for Cancer Research.* 2019; 25(11):3304-3316.

15. Zabijak L, Attencourt C, Guignant C, et al. Increased tumor infiltration by mucosal-associated invariant T cells correlates with poor survival in colorectal cancer patients. *Cancer Immunol Immunother.* 2015; 64(12):1601-1608.

16. Peterfalvi A, Gomori E, Magyarlaki T, et al. Invariant Valpha7.2-Jalpha33 TCR is expressed in human kidney and brain tumors indicating infiltration by mucosal-associated invariant T (MAIT) cells. *Int Immunol.* 2008; 20(12):1517-1525.

17. Zhao Z, Zhang KN, Wang Q, et al. Chinese Glioma Genome Atlas (CGGA): A
Comprehensive Resource with Functional Genomic Data from Chinese Glioma Patients.
Genomics Proteomics Bioinformatics. 2021; 19(1):1-12.
18. Howson LJ, Awad W, von Borstel A, et al. Absence of mucosal-associated invariant T
cells in a person with a homozygous point mutation in MR1. *Sci Immunol.* 2020; 5(49).
19. Flavahan WA, Drier Y, Liau BB, et al. Insulator dysfunction and oncogene activation in
IDH mutant gliomas. *Nature.* 2016; 529(7584):110-114.
20. Darmanis S, Sloan SA, Croote D, et al. Single-Cell RNA-Seq Analysis of Infiltrating
Neoplastic Cells at the Migrating Front of Human Glioblastoma. *Cell reports.* 2017;
21(5):1399-1410.
21. Uhlen M, Zhang C, Lee S, et al. A pathology atlas of the human cancer transcriptome.
Science (New York, N.Y.). 2017; 357(6352).
22. Mollinedo F. Neutrophil Degranulation, Plasticity, and Cancer Metastasis. *Trends
Immunol.* 2019; 40(3):228-242.
23. Dean MJ, Ochoa JB, Sanchez-Pino MD, et al. Severe COVID-19 Is Characterized by an
Impaired Type I Interferon Response and Elevated Levels of Arginase Producing
Granulocytic Myeloid Derived Suppressor Cells. *Frontiers in immunology.* 2021;
12:695972.
24. Zhou J, Nefedova Y, Lei A, Gabrilovich D. Neutrophils and PMN-MDSC: Their
biological role and interaction with stromal cells. *Semin Immunol.* 2018; 35:19-28.
25. Aarts CEM, Kuijpers TW. Neutrophils as myeloid-derived suppressor cells. *Eur J Clin
Invest.* 2018; 48 Suppl 2:e12989.
26. Alshetaiwi H, Pervolarakis N, McIntyre LL, et al. Defining the emergence of myeloid-
derived suppressor cells in breast cancer using single-cell transcriptomics. *Sci Immunol.*
2020; 5(44).
27. Brennan CW, Verhaak RG, McKenna A, et al. The somatic genomic landscape of
glioblastoma. *Cell.* 2013; 155(2):462-477.
28. Yang Y, Li C, Liu T, Dai X, Bazhin AV. Myeloid-Derived Suppressor Cells in Tumors:
From Mechanisms to Antigen Specificity and Microenvironmental Regulation. *Frontiers
in immunology.* 2020; 11:1371.
29. Masucci MT, Minopoli M, Carriero MV. Tumor Associated Neutrophils. Their Role in
Tumorigenesis, Metastasis, Prognosis and Therapy. *Frontiers in oncology.* 2019; 9:1146.
30. Michaeli J, Shaul ME, Mishalian I, et al. Tumor-associated neutrophils induce apoptosis
of non-activated CD8 T-cells in a TNFα and NO-dependent mechanism, promoting a
tumor-supportive environment. *Oncoimmunology.* 2017; 6(11):e1356965.
31. Grzywa TM, Sosnowska A, Matryba P, et al. Myeloid Cell-Derived Arginase in Cancer
Immune Response. *Frontiers in immunology.* 2020; 11:938.
32. Rodriguez PC, Zea AH, DeSalvo J, et al. L-arginine consumption by macrophages
modulates the expression of CD3 zeta chain in T lymphocytes. *Journal of immunology
(Baltimore, Md. : 1950).* 2003; 171(3):1232-1239.
33. Shen Y, Wei Y, Wang Z, et al. TGF-β regulates hepatocellular carcinoma progression by
inducing Treg cell polarization. *Cell Physiol Biochem.* 2015; 35(4):1623-1632.
34. Chu HW, Trudeau JB, Balzar S, Wenzel SE. Peripheral blood and airway tissue
expression of transforming growth factor beta by neutrophils in asthmatic subjects and
normal control subjects. *J Allergy Clin Immunol.* 2000; 106(6):1115-1123.

616	**35.**	Zhou SL, Zhou ZJ, Hu ZQ, et al. Tumor-Associated Neutrophils Recruit Macrophages
617		and T-Regulatory Cells to Promote Progression of Hepatocellular Carcinoma and
618		Resistance to Sorafenib. *Gastroenterology.* 2016; 150(7):1646-1658.e1617.
619	**36.**	Veglia F, Sanseviero E, Gabrilovich DI. Myeloid-derived suppressor cells in the era of
620		increasing myeloid cell diversity. *Nat Rev Immunol.* 2021; 21(8):485-498.
621	**37.**	Sinha P, Clements VK, Bunt SK, Albelda SM, Ostrand-Rosenberg S. Cross-talk between
622		myeloid-derived suppressor cells and macrophages subverts tumor immunity toward a
623		type 2 response. *Journal of immunology (Baltimore, Md. : 1950).* 2007; 179(2):977-983.
624	**38.**	Wei C, Yang C, Wang S, et al. M2 macrophages confer resistance to 5-fluorouracil in
625		colorectal cancer through the activation of CCL22/PI3K/AKT signaling. *OncoTargets*
626		*and therapy.* 2019; 12:3051-3063.
627	**39.**	Ballbach M, Dannert A, Singh A, et al. Expression of checkpoint molecules on myeloid-
628		derived suppressor cells. *Immunol Lett.* 2017; 192:1-6.
629	**40.**	Li F, Zhao Y, Wei L, Li S, Liu J. Tumor-infiltrating Treg, MDSC, and IDO expression
630		associated with outcomes of neoadjuvant chemotherapy of breast cancer. *Cancer biology*
631		*& therapy.* 2018; 19(8):695-705.
632	**41.**	Taga K, Tosato G. IL-10 inhibits human T cell proliferation and IL-2 production. *Journal*
633		*of immunology (Baltimore, Md. : 1950).* 1992; 148(4):1143-1148.
634	**42.**	Dahmani A, Delisle JS. TGF-β in T Cell Biology: Implications for Cancer
635		Immunotherapy. *Cancers.* 2018; 10(6).
636	**43.**	Pesenacker AM, Wang AY, Singh A, et al. A Regulatory T-Cell Gene Signature Is a
637		Specific and Sensitive Biomarker to Identify Children With New-Onset Type 1 Diabetes.
638		*Diabetes.* 2016; 65(4):1031-1039.
639	**44.**	Dhodapkar MV, Geller MD, Chang DH, et al. A reversible defect in natural killer T cell
640		function characterizes the progression of premalignant to malignant multiple myeloma.
641		*The Journal of experimental medicine.* 2003; 197(12):1667-1676.
642	**45.**	Liang J, Piao Y, Holmes L, et al. Neutrophils promote the malignant glioma phenotype
643		through S100A4. *Clinical cancer research : an official journal of the American*
644		*Association for Cancer Research.* 2014; 20(1):187-198.
645	**46.**	Sippel TR, White J, Nag K, et al. Neutrophil degranulation and immunosuppression in
646		patients with GBM: restoration of cellular immune function by targeting arginase I.
647		*Clinical cancer research : an official journal of the American Association for Cancer*
648		*Research.* 2011; 17(22):6992-7002.
649	**47.**	Fossati G, Ricevuti G, Edwards SW, Walker C, Dalton A, Rossi ML. Neutrophil
650		infiltration into human gliomas. *Acta neuropathologica.* 1999; 98(4):349-354.
651	**48.**	Dusseaux M, Martin E, Serriari N, et al. Human MAIT cells are xenobiotic-resistant,
652		tissue-targeted, CD161hi IL-17-secreting T cells. *Blood.* 2011; 117(4):1250-1259.
653	**49.**	De Leo A, Ugolini A, Veglia F. Myeloid Cells in Glioblastoma Microenvironment. *Cells.*
654		2020; 10(1).
655	**50.**	He D, Li H, Yusuf N, et al. IL-17 promotes tumor development through the induction of
656		tumor promoting microenvironments at tumor sites and myeloid-derived suppressor cells.
657		*Journal of immunology (Baltimore, Md. : 1950).* 2010; 184(5):2281-2288.
658	**51.**	Guan X, Liu Z, Zhang J, Jin X. Myeloid-derived suppressor cell accumulation in renal
659		cell carcinoma is correlated with CCL2, IL-17 and IL-18 expression in blood and tumors.
660		*Adv Clin Exp Med.* 2018; 27(7):947-953.

52. Del Monte U. Does the cell number 10(9) still really fit one gram of tumor tissue? *Cell*
cycle (Georgetown, Tex.). 2009; 8(3):505-506.
53. McWilliam HEG, Villadangos JA. How MR1 Presents a Pathogen Metabolic Signature
to Mucosal-Associated Invariant T (MAIT) Cells. *Trends Immunol.* 2017; 38(9):679-689.
54. Louis DN, Perry A, Wesseling P, et al. The 2021 WHO Classification of Tumors of the
Central Nervous System: a summary. *Neuro-oncology.* 2021; 23(8):1231-1251.
55. Cerami E, Gao J, Dogrusoz U, et al. The cBio cancer genomics portal: an open platform
for exploring multidimensional cancer genomics data. *Cancer discovery.* 2012; 2(5):401-
404.
56. Budczies J, Klauschen F, Sinn BV, et al. Cutoff Finder: a comprehensive and
straightforward Web application enabling rapid biomarker cutoff optimization. *PloS one.*
2012; 7(12):e51862.
57. Raudvere U, Kolberg L, Kuzmin I, et al. g:Profiler: a web server for functional
enrichment analysis and conversions of gene lists (2019 update). *Nucleic acids research.*
2019; 47(W1):W191-w198.

 Figure legends

Figure_1

The *MR1* gene status and its prognostic correlation in GBM

(a) The RNA-expression data of TCGA-GBM 168 core samples were analyzed to see the

expression levels of *MR1*. (b) The correlation between *MR1* expression and GBM survival.

Clinical data available TCGA-GBM 153 patients' data and Chinese Glioma Genome Atlas

(CGGA) 85 primary GBM patients' data were analyzed. Cut off FPKM: TCGA-MR1 = 3.67,

CGGA-MR1 = 9.59, Log-rank test (c) The mutational schema of *MR1* locus in 3391 samples of

glioma. Only 2 missense mutations were detected. (d) H3K27ac ChIP viewer image in *MR1* locus

of GBM and high-grade glioma samples, GBM2w (IDHwt GBM, GSM1824806), AA15m

(IDHmut-R132H, Grade3 Astrocytoma, GSM1824808). (e) Single-cell RNA-seq data of 3589

cells from GBM tissues of four patients were analyzed with a tSNE plot. All colored dots represent

cells expressing *MR1*.

Figure_2

Correlations of *MR1* expression and the activity of TANs/MDSCs

(a) A schematic illustration of the extracted top 500 significantly co-expressing genes with *MR1*

and the prognosis related 158 genes among them. The horizontal axis is the Spearman's *r* values

for each gene's correlation level with *MR1*, and the vertical axis is the *q*-value (Benjamini-

Hochberg adjusted *p*-value) for each gene. (b) The enriched pathways of the 158 genes. The bar

chart shows the enriched *q*-values for each pathway. (c) Correlation scatter charts between *MR1*

gene expression, NDsig-index, and MDSCsig-index among the 168 TCGA-GBM samples.

Figure_3

Multi-faceted evaluation of the MAIT infiltrated samples in TCGA-GBM

(a) The levels of MAIT-cell infiltration in TCGA-GBM samples. The 8.3% (14/168) of samples

were detected as MAIT cell infiltration positive samples. The infiltration level was calculated with

the formula, [The detected read count of the MAIT-target read of the sample/ The total RNA read

count of the sample]. (b)(c) Violin plots showing the gene expression level of MAIT cell surface

receptors/secreting cytokines. (d) The information on MAIT cell infiltration was integrated with

other clinical/molecular indicators. Molecular indicators' information was obtained from a

previously reported database[27], and the neural samples were integrated into the proneural samples.

(e) The ratio-bar chart shows the distribution of GBM subtypes among MAIT-positive samples

and the total TCGA-GBM samples. (f) A violin plot showing the *MR1* gene expression level in

the MAIT positive/negative groups.

Figure_4

Evaluations of the activity of TANs/MDSCs in MAIT-positive/negative samples

(a) The enrichment plot of GSEA-GOBP's negative regulation of immune response in MAIT-

positive/negative samples. (b) The enrichment results of neutrophil-related pathways in GSEA-

GOBP in MAIT-positive samples. All pathways shown were significantly enriched ($q < 0.05$). (c)

Heatmap of the gene expression level of ND30 genes for each sample. The vertical column

represents one GBM sample. MAIT-positive samples were gathered on the right. (d) A Violin plot

showing the ND30-index level. (e) GSEA enrichment plots of neutrophil degranulation pathway

and MDSC signature. (f) Scatter chart showing the differential expression levels of the

immunosuppressive pathway genes related to TANs/MDSCs among MAIT-positive/negative

samples. The horizontal axis shows the fold change (log2 converted) as follows. [Mean FPKM of

MAIT-positive samples / Mean FPKM of MAIT-negative samples] The vertical axis shows the *p*-

value (-log2 converted) of the unpaired t-test that compares the expression levels of MAIT-

positive/negative samples for the gene. (g) Violin plots showing the *ASS1* or *ARG1* gene

expression level. (h) Violin plots showing the expression level of *FOXP3* and *CCR4*. (i) GSEA

enrichment plot, using our Treg-signature gene set. (j) The prognostic correlation of NDsig-index

and MDSCsig-index with TCGA-GBM clinical data. Cut off: NDsig-index = 41.89, MDSCsig-

index = 27.48, Log-rank test

Figure_5

Correlations between *MR1*, prognosis, and TANs/MDSCs in the pan-cancer analysis

(a) The prognostic correlation of *MR1* in multiple cancer types. All red lines represent the MR1-

high group, and blue lines represent the MR1-low group. (b) Enrichment plots of the GSEA using

neutrophil degranulation pathway and our MDSCsig gene set for multiple cancer types. MR1-

high/low groups were compared. (c) The enrichment results of neutrophil related pathways in

GSEA-GOBP in the MR1-high group. All pathways shown were significantly enriched ($q < 0.05$).

Figure_6

The MAIT/MR1-TAN/MDSC pathway

The activation of MAIT cell, along with the *MR1* up-regulation, activates and chemo-attracts

granulocytes, such as TANs/MDSCs, through IL-17 and TNF production. The activated

TANs/MDSCs induce Tregs via the production of TGF-β, CCL17, and CCL22, and Tregs suppress

the T cell-mediated immunity. Simultaneously, TANs/MDSCs suppress T cells via multiple

pathways. The suppression of T cell immunity provides a poor prognosis for GBM.

Supplementary figure

Figure_S1: The differential expressions of MAIT cell surface receptor genes in MR1-high/low group with pan-cancer analysis

Scatter charts show the differential expression levels of the MAIT cell surface receptor genes in the MR1-high/low group. The horizontal axis shows the fold change (log2 converted) as follows. [Mean FPKM of MR1-high group / Mean FPKM of MR1-low group] The vertical axis shows the significance (unpaired t-test).