The snapdragon genomes reveal the evolutionary dynamics of the S locus supergene

Sihui Zhu[†,1,2,3], **Yu'e Zhang**[†,4], **Lucy Copsy**[5], **Qianqian Han**[3,4], **Dongfeng Zheng**[1,2,3], **Enrico Coen**[5], **Yongbiao Xue**[*,1,2,3,4]

[1] National Genomics Data Center & CAS Key Laboratory of Genome Sciences and Information, Beijing Institute of Genomics, Chinese Academy of Sciences, Beijing 100101, China

[2] China National Center for Bioinformation, Beijing 100101, China

[3] University of Chinese Academy of Sciences, Beijing 100049, China

[4] State Key Laboratory of Plant Cell and Chromosome Engineering, Institute of Genetics and Developmental Biology, and the Innovation Academy of Seed Design, Chinese Academy of Sciences, Beijing 100101, China

[5] John Innes Centre, Norwich NR47UH, UK.

† These authors contributed equally to this work.

* **Correspondence**: Yongbiao Xue (ybxue@genetics.ac.cn)

1 **Competing Interest Statement:** The authors declare no competing

2 interest.

3 **Keywords:** snapdragon, *S*-locus, supergene, evolutionary genomics, *SLFs*

1 Abstract

The multi-allelic S-locus, containing a pistil S-$RNase$ and dozens of S-locus F-

box (SLF), underlies genetic control of self-incompatibility (SI) in *Antirrhinum*

hispanicum. The genus *Antirrhinum*, harboring such a SI system has been

used as a model to study self-incompatibility extensively. However, there have

been limited studies on the genomic organization of the S-locus supergene

due to a lack of high-quality genomic data. Here, we present the

chromosome-level reference and haplotype-resolved genome assemblies of a

self-incompatible *Antirrhinum hispanicum* line, AhS_7S_8. Alongside with the

draft genome of *Misopates orontium*, comparative genomics reveals that

A.hispanicum diverged from its self-compatible cousin 12.3 million years ago

(Mya). Expanded gene families enriched in distinct functional terms implied

different evolutionary trajectories of outcrossing and selfing species. For the

first time, two complete *A.hispanicum* S-haplotypes spanning ~1.2Mb and

containing 32 $SLFs$ were reconstructed, while most of the $SLFs$ derived from

retroelement-mediated proximal or tandem duplication approximately 122 Mya.

Moreover, we detected a candidate *cis*-transcription factor associated with

regulating $SLFs$ expression, and two miRNAs may control the expression of

this transcription factor. Inter-specific S-locus and intra-specific S-haplotype

comparisons revealed the dynamic nature and polymorphism of the S-locus

supergene mediated by continuous gene duplication, segmental translocation

or loss, and TE-involved transposition events. Our data provides an excellent

resource for future research on the evolutionary studies on *S-RNase*-based

self-incompatibility system.

1 Introduction

Self-incompatibility (SI) is a molecular recognition system that prevents inbreeding and promotes outcrossing in hermaphroditic flowering plants, thereby maintaining genetic diversity and helping angiosperms expand into a wide range of habitats. The molecular mechanisms of SI in eudicots have been extensively studied for decades. In Solanaceae, Rosaceae, Rubiaceae, Rutaceae and Plantaginaceae, SI is controlled by a single supergene called the *S*-locus, with a variety of haplotypes[1]. The *S*-locus generates the pistil *S* determinant (*S*-ribonuclease with cytotoxicity) and the pollen *S* determinant (dozens of pollen-specific *S*-locus F-box genes called *SLFs*) that mediate self/non-self recognition. A unique *S*-haplotype consists of specific *SLFs* and a specific *S-RNase*. Successful fertilization only occurs between individuals harboring different *S*-haplotypes, and the S-RNase can be degraded by non-self SLF products collectively[2]. Reconstructing the evolutionary history of a supergene is necessary to understand how such a functional module originate and spread in diverse taxons[3, 4]. Supergenes controlling complex phenotypes, which include Batesian mimicry in butterfly[5] and alternative social organization in fire ant colonies[6], have long inspired both empirical and theoretical studies on them. However, in plant SI species, most previous reports focused on the discussions of self-incompatible genes themselves, few systematic analyses of the origin and evolution of the *S*-locus at the whole genome level, and the

impact of gene duplication on the shaping process of the functional locus was not considered, presenting a critical knowledge gap.

The breakdown of SI results in a mating system change from outcrossing to self-fertilization in angiosperms. SI has been lost frequently during angiosperm diversification[7], and the transition to self-compatibility (SC) are usually accompanied by accumulation of loss-of-mutations of the *S*-locus[8]. There has been intense interest in the role of parallel molecular changes underlying the evolutionary shifts to selfing associated with loss of SI[9]. Loss-of-function or loss-of-expression of *S-RNase* can confer self-compatibility (SC) in some Solanaceae species[10, 11], which was also observed in snapdragon[12,13]. Therefore, it has long been served as an ideal system to explore the impact of mating system shifts and dynamics of adaptation. With the advent of the post genome era, accurate and contiguous genome assembly, the critical basis for understanding genomic diversity and evolutionary trajectories, is available now. In this study, we assembled a chromosome-level reference genome with a total length of ~480Mb of self-incompatible *Antirrhinum hispanicum* (AhS_7S_8, heterozygous genotype at *S-locus*, hereafter referred to as 'SI-Ah'). Also, high-quality draft genomes of self-incompatible *Antirrhinum linkianum* and self-compatible *Misopates orontium* were present here. These sequence resources could provide us an opportunity to perform comparative genomic analyses to capture consequences of the self-incompatibility breakdown and obtain insights into the evolutionary genomics basis of the loci controlling self-

incompatibility. We also took the advantage of HiFi data and Hi-C reads to assemble two haploid genomes of SI-Ah, each contains an intact *S*-haplotype. In this study, the complete structure and sequence of the *Antirrhinum S*-locus allowed us to inspect how and when the *SLF* genes proliferate, as well as variations between different *S*-haplotypes. The genomic data and analysis reported in this work are of great value for understanding the genetic bases of adaptive characteristic of species, and provide insights into the evolution of the *S*-locus in snapdragon.

Results

Genome assembly and evaluation

We combined different sequencing strategies to derive *A.hispanicum* reference genome assembly: Illumina pair-ended reads, PacBio single-molecule reads, BioNano optical maps, and Hi-C sequencing reads (Fig. S1 and Table S1). The self-incompatible line *A.hispanicum* of AhS_7S_8 was estimated to be with genome size of ~473.4Mb and a modest heterozygosity of 0.59% (Fig. S2) based on the 21-mer spectrum. Firstly, 36.0Gb of continue long reads (CLR) and 19.1Gb of circular consensus reads (CCS) were obtained on PacBio platform for primary assembly using different assemblers. Then, a total of 203.3Gb BioNano data and 69.9Gb Hi-C paired-end reads were produced to scaffold the contigs into 8 pseudo-chromosomes. In the end, we obtained three sets of chromosome-level genome assemblies of AhS_7S_8, a

consensus assembly with mosaic sequence structure (Fig. 1) and two

haplotype-resolved assemblies each contains a copy of the *S*-locus (S_7

haplome with S_7-haplotype and S_8 haplome with S_8-haplotype). Yet long

stretch of gaps exists in S_8-haplotype of S_8 haplome and cannot be filled up

directly with the sequencing data of AhS_7S_8 due to heterochromatic

characteristics of the *S*-locus[14]. Therefore, to overcome these difficulties in

assembly of the *S*-locus, we applied a genetic approach using a cross

between AhS_7S_8 and self-compatible relative *A.majus* (S_CS_C)[15], generating

two F1 (AhS_7S_C and AhS_8S_C). Since the F1 individuals have haploid

chromosome sets from AhS_7S_8, their sequencing data were used for error

correction and gap filling to improve the haplome assemblies. Comparing

A.hispanicum phased genome sequences with the consensus assembly

revealed generally good alignment except some gaps and inversions in few

chromosomes, which might be caused by the heterozygous region collapse in

Canu assembly (Fig. S4). To facilitate comparative analyses, we also

sequenced and assembled the genomes of self-incompatible *Antirrhinum*

linkianum and self-compatible *Misopates orontium.* The heterozygosity and

genome size of *A.linkianum* is very close to that of *A.hispanicum*, while

M.orontium exhibits heterozygosity rate of 0.21% and approximately 24.1%

reduction in genome size (Table S1).

The BUSCO assessment results indicated that 95.0-98.7% of conserved

genes were completely recovered in these assemblies (Fig. S6 and Table S6).

The S_7 and S_8 haplomes were combined and evaluated under the protein mode, and 98.6% (2,090) of BUSCO genes were complete, while 92.2% (1,955) being duplicated. A total of 30,956 *Antirrhinum* EST records downloaded from NCBI (2021-04-23) were mapped to the genome assemblies, with nearly 90% of which can be mapped (Table S5). PE reads from the DNA and RNA libraries from this study were mapped to the genome assemblies respectively. The mapping rate ranged from 70% to 94% (properly mapped or uniquely mapped), as listed in Table S5. In addition, LTR Assembly Index(LAI)[16] based on LTR-RTs was calculated to evaluate genome assembly continuity. The LAI values of the two haplomes exceed 20, which demonstrates a 'golden' standard level, and the draft genome of other two species also reached 'reference' grade quality (19.3 and 21.9 for *A.linkianum* and *M.orontium*, respectively). Cumulatively, the results above suggested the reliability of all the nuclear genome assemblies. In addition, circular chloroplast and mitochondrial genomes were also retrieved from contigs assembly, with length of 144.8kb and 527.4kb respectively (Fig. S5).

Particularly, the *S*-locus of self-incompatible snapdragon which spans more than 1.2Mb was completely assembled with a continuous BioNano molecular cmap as strong evidence (Fig. S7). The *A.hispanicum* genome assemblies are, to the best of our knowledge, one of the most contiguous and complete *de novo* plant genomes covering the complex *S*-locus supergene to date.

Genome annotation

Repetitive elements accounted for approximately 43% of *Antirrhinum* genome (Table S4), and long terminal repeats (LTRs) being the largest member of transposons families, covering ~23% of the nucleus genome (Fig. 1A), of which, the Ty3/gypsy (12.0%) and Ty1/copia (11.6%) are the most abundant transposable elements (TEs). DNA transposons comprises ~12% of the *Antirrhinum* genomes (Table S4). According to theoretical modeling, if the effects of TE insertion are mostly recessive or co-dominant, increased homozygosity would result in more efficient elimination[17], which therefore should be excepted have lesser TE in selfing species. As we observed, the total amount of TE in self-compatible *M.orontium* (38.0%) is apparently lower than self-incompatible species (Table S4). The insertion time of intact LRT-RTs in *Antirrhinum* based on the divergence of two flanking terminal repeat sequences revealed a recent burst of LTR-RTs at 0.3 Mya, and indicated that most old LTR-RTs have been lost (Fig. S9). The younger insertions have not accumulated enough substitution in two flanking repeats, making this type of genome more difficult to assemble correctly without long-range information. Gene models of all genome assemblies were predicted followed the same analysis pipeline as described in method. For consensus assembly of *A.hispanicum*, 42,667 protein-coding genes were predicted from the soft-masked genome, 89.3% of which could be annotated in at least one functional database, and other assemblies gave close genome annotation results (Table

S7). The average transcript and CDS length of protein-coding genes were 3,023 bp and 1,146bp, respectively. The average exon and intron lengths were 232 bp and 479 bp, with 4.8 exons per gene on average. The length distribution of CDSs, exons, and introns of snapdragon show similar patterns to other plant species (Fig. S10), which confirmed the credibility of gene prediction results. The GC content is distributed unevenly in most pseudo-chromosomes and tends to be higher in gene-poor regions. The density of protein-coding genes on the pseudo-chromosomes was highest in distal regions and declined in centromeric regions (Fig. 1), suggesting that gene distribution is not random. As expected, the pattern of gene presence is opposite to that of LTR-RT elements and methylation level. A total of 2,112 transcription factors belong to 58 families were annotated, including bHLH, ERF, MYB, C2H2, NAC, and B3 families with largest number of members (Fig. S11). Analysis of small-RNAseq libraries from four tissues identified 108 known miRNA genes of 42 families of *A.hispanicum* and 105 miRNA genes of 42 families (details listed in Table S9). In addition, 719 transfer RNA genes (tRNAs), 311 small nuclear RNA genes (snRNAs), and 791 ribosomal RNA (rRNAs) genes for *A.hispanicum* were predicted (Table S8).

Taken together, three parts of snapdragon genome (nuclear, chloroplast, mitochondria) are of high completeness and correctness, making them a valuable resource for comparative and evolutionary studies.

Comparative genomic analysis

We employed the consensus-assembly to represent *A.hispanicum* in comparative genomic analysis to conduct phylogenomic analysis. Proteomes from *A.hispanicum, A.linkianum*, and *M.orontium*, together with 15 other species (Table S10) were clustered into 36,295 orthogroups (Table S11) and used for phylogenetic relationship construction subsequently. In Lamiids, the family Plantaginaceae, to which *Antirrhinum* and *Misopates* belongs, form a monophyletic group that diversified recently. The species tree indicated that the divergence of Plantaginaceae from Lamiales order common ancestor occurred ~55.8 Mya during Paleocene-Eocene transition (Fig. 2A). After that, *M.orontium* diverged from *Antirrhinum* genus and lost self-incompatibility assuming self-compatibility is ancestral state in Plantaginaceae[7]. *A.hispanicum* and *A.linkianum* split at around 3.5 Mya [95% CI=1.6-5.3 Mya], which is collided with emergence of modern Mediterranean climate and rapid speciation of *Antirrhinum* lineage, and in agreement with previously reported results using various datasets and methodologies[18].

To identify WGD events in *Antirrhinum*, we first compared the SI-Ah genome with the reconstructed ancient eudicot (AEK)[19] genome, and *Vitis vinifera* genome which have underwent none large-scale genome duplication after whole-genome triplication (WGT) of AEK[20]. Synteny analysis of the SI-Ah with AEK and *Vitis vinifera* genomes showed synteny depths ratio of 1:6 and 1:2 respectively (Fig. S12), and large-scale segmental duplications can be

observed from the self-alignment of SI-Ah (Fig. 1). The Ks distribution of

paranomes displayed two peaks (Fig. 2B), indicating *Antirrhinum* genus have

at least experienced another one WGD after the gamma-WGT.

To approximately date the WGD in snapdragon, the distributions of

synonymous substitutions per synonymous site (Ks) among paralogous genes

within a genome were examined. The distributions plots of Ks for

A.hispanicum, *C.rattanii*[21] and *A.linkianum* paranomes showed a distinct peak

at 2.21 echoing the well-reported gamma-WGT occurred in the common

ancestors of all eudicots approximately 117 Mya[22](Fig. 2B and S14), as well

as an additional peak at ~0.61 corresponding to a later duplication event

shared among Plantaginaceae members. Therefore, the younger WGD was

dated to occur at ~32.3 Mya, posterior to divergence of *Antirrhinum* and other

Lamiales plants, indicating a Plantaginaceae-specific WGD event. The

Antirrhinum lineage neutral mutation rate was inferred using the formula

$\mu=D/2T$, where D is the evolutionary distance of two *Antirrhinum* species

(peak value of Ks log-transformed distribution = 0.04, Fig. 2C), and T is the

divergence time of the two species (3.5Mya). The neutral mutation rate for

Antirrhinum genus was estimated to be 5.7e-9 (95% CI: 3.8e-9 - 1.3e-8)

substitution per site per year, which is very close to the result estimated by

using genotyping-by-sequencing dataset[18].

Loss of self-incompatibility occurred many times during angiosperms evolution

history[7, 8]. Although two species have split up for millions of years, synteny

block analysis revealed strong collinearity between *A.hispanicum* and *M.orontium* (Fig. S13). And to investigate the evolutionary trajectory of SI- and SC- snapdragons, gene family analysis was conducted based on orthologous relationship of 18 plant species (Fig. 2A). A total of 445 and 294 gene families were expanded and contracted respectively in *Antirrhinum* lineage. The results also revealed 704 (3,230 genes) expanded and 1,692 (2,956 genes) contracted gene families in *A.hispanicum*, 1,097 (4,787 genes) expanded and 1,392 (4,015 genes) contracted gene families in *A.linkianum* respectively. (Fig. 2A). Moreover, 702 gene families (4,305 genes) were found in expansion, while 1209 families (1,771 genes) were found in contraction in the self-compatible *M.orontium*. These rapidly evolving gene families might give us some clues about consequences of losing self-incompatibility in *M.orontium* lineage.

The GO enrichment analysis for 445 expanded gene families of self-incompatible *Antirrhinum* branch showed that the significantly enriched GO terms mostly involved 'binding', 'enzyme activity', and 'response' (Fig. 2D). While in *M.orontium*, enriched GO terms of expanded families mostly related to photosynthesis (Fig. 2E), and KEGG enrichment analysis revealed that the expanded genes were mainly implicated in phenylpropanoid, flavonoid, sesquiterpenoid and triterpenoid biosynthesis (Fig. S16), which may contribute to the survival of self-compatible species facing environmental changes on the way to an evolutionary 'dead-end'[23]. For the complete list of

GO enrichment analysis results, see Table S12 and S13.

Position, structure, and flanking genes of the *S*-locus supergene

Identifying sequence details of the *S*-locus supergene would be the initial step towards understanding its evolution in snapdragon. The *Antirrhinum S*-locus was found to locate in peri-telomeric region on the short arm of chromosome 8[14], as opposed to sub-centromeric area in *Solanaceous* species. A total of 275 genes encoding F-box protein were annotated in the entire snapdragon genome (Fig. 1). A cluster of 32 *SLF* (*S*-locus F-box) genes, which match to the structural characteristics of the candidate *S*-locus, located in chromosome 8 (Fig. 1). Gapless *S*-locus of 1.25Mb in *A.hispanicum* and pseudo *S*-locus of 804kb in *A.majus*[15] are both reconstructed with BioNano long-range molecules as support evidence (Fig. S7 and S8), whereas the *S*-loci of *A.linkianum* and *M.orontium* were identified by synteny alignment with *A.hispanicum*. Except *SLFs*, the *S*-locus also contains genes encoding G-beta repeat, matE, zinc finger, cold acclimation, auxin responsive proteins, and dozens of genes with unknown function or TE-related function (listed in Table S16). Additionally, the *S*-locus was surrounded by two copies of organ specific protein at left side (OSP1: *AnH01G33547.01*, OSP2: *AnH01G33548.01*) and five copies of phytocyanin genes at right side (here named PC1: *AnH01G33649.01*, PC2: *AnH01G33650.01*, PC3: *AnH01G33651.01*, PC4:

AnH01G33652.01, PC5: *AnH01G33653.01*) (Fig. 3E). The contiguity and completeness of the genome assembly, together with solid annotation, give rise to the possibility of proposing an origin model of the *S*-locus supergene. There is no doubt that gene duplication plays a key role in shaping process of such a complex supergene. It has been proved that tandem or proximal duplicates evolved toward functional roles in plant self-defense and adaptation to dramatic changing environments[24]. To inspect how multiple *SLFs* were generated and proliferated, phylogenetic tree of 275 *SLFs* was built (Fig. 3A). So, if all the *SLFs* originated via WGDs or large segmental duplication, their paralogs should colocalize elsewhere in the genome and have similar divergence with the *S*-locus genes. On the contrary, if *SLFs* were the results of stepwise local duplications, they would exhibit less differences and there should be more distant paralogs scattered elsewhere in the genome. The *SLFs* phylogenetic tree proves that the situation is the latter one. The phylogenetic tree along with expression heatmap of 275 F-box genes indicated *SLFs* not only share closer relationship, but also display similar pollen-specific expression pattern which distinguished from other F-box genes (Fig. 3A). This phenomenon has not changed much in cultivar *A.majus* after its domestication (Fig. S21). Furthermore, based on classification of gene duplication type, the two OSPs derived from the gamma-WGD followed by a recent tandem duplication (Table S20), while the PCs arise from tandem duplication asynchronously assuming constant substitution rate, both can be

supported by gene phylogenetic tree topology of their paralogs (Fig. S22,

S23). As for *SLFs*, 24 and 6 of them are proximal duplication and tandem

duplication genes respectively, and other 2 *SLFs* are dispersed duplication

derived (Table S16). Besides, 27 of 32 *SLFs* are intron-less, suggesting *SLFs*

proliferated most likely through retroelement-mediated way. *Ks* distribution of

pairwise *SLFs* and non-*S*-locus F-box genes were compared, and the

synonymous substitutions rates of *SLF* paralogs are much lower than non-*S*-

locus F-box paralogs (Fig. 3B). With the estimated *Antirrhinum* mutation rate,

the peak value of *SLFs* paralogs *Ks* distribution (ranging from 1.35 to 1.40)

corresponding 122 Mya[25] (Fig. 3B), indicated that the *S*-locus structure is a

very conserved and long lived supergene survived from a long-term balancing

selection[26, 27]. This observation concludes that the *S*-allelic polymorphisms

should be inherited from common ancestors and the differentiations began

before the currently extant species formation. Dissection of *SLFs* and flanking

genes indicate that the *S*-locus is still experiencing gene gaining in

Antirrhinum (Table S20).

Using MEME suit analysis, we identified a significant motif

(GATCCTAXAATATCTC) located upstream of most *SLFs*, and the motif

annotation implied *SLFs* are likely to be co-regulated by a MYB-related family

transcription factor (Fig. 3C). By examining expression of all the MYB-related

family members in the genome, *AnH01G01437.01, AnH01G01435.01,*

AnH01G03933.01, AnH01G28556.01, AnH01G01110.01, AnH01G16778.01,

AnH01G14798.01, AnH01G03969.01, AnH01G42539.01, AnH01G33501.01, AnH01G41740.01, AnH01G31594.01, and *AnH01G17038.01* were expressed in pollen but barely in pistil tissue (Fig. 3D). Among them, *AnH01G33501.01* is located about 1.2Mb upstream of the *S*-locus, making it a potential regulator involved in activating expression of the *SLFs*. Moreover, by examining miRNA target prediction results, two candidate miRNAs (aof-miR396b and zma-miR408b-5p) were revealed to bind with different regions of *AnH01G33501.01*, and their expression patterns exhibited negative correlation with the TF (Fig. S25). The miR396 family member, aof-miR396b, was believed to influence GRF transcription factors expression[28], while over-expression of zma-miR408b in maize were reported to respond to abiotic stress to maintain leaf growth and improve biomass[29]. The results implied that those two miRNAs may also play an important role in self-incompatibility.

From the intraspecies similarity matrix heatmap of *SLFs* of SI-Ah, we can also observe the divergence among paralogs are much higher than interspecific orthologous pairs (Fig. S24). The pairwise sequence identity of 32 *SLF* paralogs ranged from 33.1 to 77.3%. And the closer the physical distance between two *SLFs*, the higher the sequence similarity and identity (Fig. 4A). In addition, the values of the non-synonymous/synonymous substitutions rate ratio (Ka/Ks) of nearly all *SLFs* from different allele below one is a signature of purifying selection to maintain necessary protein functions to detoxify S-RNase collectively (Fig. 4B).

Evolution history of the *Antirrhinum S*-locus

Comparison of microsynteny revealed the presence of the *S*-locus remnant in other *S-RNase*-based self-incompatible species, including *Chiococca alba*[30] (Rubiaceae), *Citrus clementina*[31] (Rutaceae), *Physalis pubescens*[32] (Solanaceae), *Prunus davidiana*[33] (Rosaceae), but not in the syntenic regions of species outside Eudicotyledoneae, such as *Oryza sativa* (Poaceae). The distribution pattern also suggested that the *S*-locus emerged early right after the divergence of Eudicotyledoneae from with other Mesangiosperms. In *P.davidiana*, core part of the whole *S*-locus was translocated to a different chromosome, and only four *F-boxes* are still there occupying a drastically contracted region (Fig. 5A). Among the four genes, two of them have 4 and 3 orthologous *SLFs* in SI-Ah implying that the multiple copies formed after species divergence. The same contraction was observed in *Chiococca* except with a few more F-box genes. Whereas in citrus genome, none of the homologous *SLFs* were found. However, in *Physalis pubescens*, a Solanaceae member phylogenetically closer with *Antirrhinum*, almost all *SLFs* were lined up in same order with *Antirrhinum* and well preserved. Therefore, it would be reasonable to infer that the protype of extant *S*-locus with only several *SLFs* and *S-RNase* appeared in the common ancestor of eudicots[25, 34, 35]. Then, accompanied by speciation, *SLFs* genes underwent loss and gain at different rate in different lineages (Fig. 5F). If the new *SLFN* copy can recognize more non-self S-RNase[12], then it is advantageous for carrying this

new duplication and spreading through the population by genetic drift[36]. Gene acquisition was most likely through tandem or proximal duplications from ancestor copies, and the losses mainly involved large fragment deletion.

The continuity of genome assemblies in this study also provides an opportunity to make comparison of S-haplotypes from same or very close species. For the sake of simplicity, we use 'S_7-locus' and 'S_8-locus' to refer the S-locus from the S_7- and S_8- haplotypes containing S-RNase and SLFs respectively, and their SLF genes as 'S_7-SLF' and 'S_8-SLF', while Al-S-locus refers the S-locus of self-incompatible $A.linkianum$. Am-S_C-locus and Mo-S_C-locus which contain no S-RNase denotes ψS-locus of self-compatible $A.majus$ and $M.orontium$. There are 32 $SLFs$ annotated in the region of S_7-locus (S_7-SLF1 ~ S_7-SLF32) and S_8-locus (S_8-SLF1 ~ S_8-SLF32) respectively. Their position in chromosome and sequence information were listed in Table S16. We compared different S-haplotypes to identify polymorphism of the supergene.

Synteny plot showed structure and gene content of the flanking ends of different S-haplotypes were well conserved. The S_7-locus has high collinearity with Am-S_C-locus, S_8 locus and Al-S-locus except a small inversion at the start of alignment (Fig. 5B-E). From the comparison between Mo-S_C-locus and S_7-locus, a large deletion with a length of about 556kb could be observed, which involved S-RNase and more than ten $SLFs$ as well as dozens of functional genes in S_7-haplotype. These genes include ATPase, Cytokinin, DUF4218,

other ones mostly are uncharacterized proteins. This structural variation can be responsible for the loss of self-incompatibility in *M.orontium*. Whereas between the S_7 and S_8 locus, a small inversion could be observed, which may involve with recombination suppression, and serval neighbor genes of S_7-*RNase* have no corresponding orthologs in S_8 haplotype (Fig. 5E), suggesting a divergent evolution history in different regions of the *S*-locus. The relative position of *S-RNase* in Al-*S*-locus being different from *A.hispanicum* (Fig. 5D), indicated that the changing physical position *S-RNase* does not impact its function and the dynamic change of the polymorphic *S*-locus in *Antirrhinum* has been very active. And we did observe a nearby CACTA transposable element in the vicinity of *A.linkianum S-RNase*, *Tam*, which may lead to the gene transposition.

Together, these results revealed the dynamic nature of the *S*-locus supergene during evolution and speciation in Eudicotyledoneae, which involves continuous gene duplication, segmental translocation or loss, and TE-mediated transposition.

Allelic expression and DNA methylation profiling

Since haplotype-resolved assembly is available and gene order is highly consistent between two haplomes (Fig. S27-34), we can determine two alleles of a physical locus and investigate allelic gene expression directly. Base on

synteny and homology relationship, 79,398 genes (90.6% of all predicted genes) were found to have homologs on the counterpart haplotype (see "Method"). Most allelic genes displayed low level of sequence dissimilarity (Fig. 6A). To understand the expression profile of allelic genes, RNA-seq datasets were analyzed using allele-aware quantification tools with combined coding sequences of two haplomes as reference. In the survey across four different tissues, most expressed loci did not exhibit large variance, only 2,504 gene pairs (3.2% of total genes, see Fig. S36) displayed significant expression imbalance between two alleles (|fold-change| ≥ 2, p ≤ 0.01), suggesting that most alleles were unbiasedly expressed in the SI-Ah genome. We also analyzed four WGBS libraries to profile methylation state of two haplomes. About 51.6-59.5 million PE reads were generated for each sample (Table S2). About 92.4-92.9% of total reads can be properly mapped to the haplomes. Reads for each sample were mapped to the snapdragon chloroplast genome too, the conversion rate higher than 99% confirming reliability of the experiments. Methycytosines were identified in CpG, CHH, and CHG contexts. A large fraction of cytosines in CpG (~80.6%) and CHG (~55%) contexts were methylated, but a smaller fraction of cytosines in CHH (~8%) context were methylated. No significant methylation level variations were observed between two haplomes (Fig. S27-34), similar to that reported in potato[37].

After examining *SLFs* and *S-RNase* expression of two haplotypes, we found

that S_7-$RNase$ and S_8-$RNase$ expressed indiscriminately and fairly high in AhS_7S_8 pistil (Table S18), which is consistent with the situation in crab cactus[38], and they are nearly undetectable in leaf and petal. Those two genes only expressed in pistils of AhS_7S_C and AhS_8S_C respectively as expected, yet at higher level than that of AhS_7S_8 pistil. Since the gene sequences did not change after one crossing experiment, it is very likely that RNase expression was changed through other mechanisms. As for the expression profiles of $SLFs$ in pollen tissue of three different genotypes, S_7-$SLFs$ and S_8-$SLFs$ did show an overall decline in two offspring samples but tend to be reduced generally. Those genes could be divided into two groups. Group1 consists of 4 genes, were highly expressed in pollen of AhS_7S_8 with obvious tissue-specificities, but barely expressed in two hybrid progenies, and these genes displayed allelic bias. Another 28 $SLFs$ in cluster 2 maintain expression patterns in three genotypes, but the expression level in two hybrid progenies are slightly lower than that of AhS_7S_8 (Fig. 6B). We assumed that group2 are necessary for degrading toxic S-RNase.

Epigenetic modifications are believed to be crucial in controlling gene expression[39]. To determine whether methylation level is responsible for the difference between two groups, we compared the DNA methylation in the pollen from AhS_7S_8. We calculated the average DNA methylation rates for the $SLFs$ gene regions and found that DNA methylation in gene body are higher than flanking regions and the highest methylation rate of mCs was in the CpG

context, followed by CHG and CHH (Fig. 6C). The average methylation rates of two groups in the CpG, CHG, and CHH are comparable in gene body and flanking regions. Thus, the similar levels and patterns of methylation of two groups of *SLFs* may not significantly explain the variance in expression. Therefore, the difference may be caused by other kinds of factors. Thus, our data show that some genes are grouped into one cluster can be regulated differentially, irrespective of their relative position within the locus.

Discussion

Assembling genomes of self-incompatible plants has always been a challenge due to inevitable heterozygosity and large portions of repetitive sequences. In current study, we utilized multiple sequencing technologies and the genetic strategy to obtain chromosome-level genome of self-incompatible *A.hispanicum*. Benefiting from the robust assembly, we can determine the genomic landscape of the first self-incompatible snapdragon. Along with high quality genomes of other two Plantaginaceae species, *A.linkianum* (2n=16) and *M.orontium* (2n=16), these assemblies would be valuable resources and enabled us to understand the genome evolution of *Antirrhinum* lineage, gaining knowledge of the proximate and ultimate consequences of losing self-incompatibility[8]. Apart from the shared gamma WGT in all eudicots,

A.hispanicum underwent a Plantaginaceae-specific WGD event dated back to the Eocene, which may have been involved in large scale chromosomal rearrangement and TEs proliferation. The phylogenomic analysis clearly reveals that *A.hispancium* split from its relative *A.linkianum* during the Pliocene Epoch, coinciding with establishment of modern Mediterranean climate in the Mediterranean basin[40]. Mating system shift is ubiquitous throughout angiosperms, the evolutionary consequences of transition have attracted extensive attention for a long time[41, 42]. The comparative genomic analysis indicated that after losing self-incompatibility, though the genomic architecture of SC- *M.orontium* has not changed much, the gene contents have undergone certain fluctuations. To increase environmental tolerance, the self-compatible organism recruits alternative biological metabolic pathways to improve survival rate, and produced more plant compounds to cope with various external stimuli, since it is no longer possible for self-compatible species through recombination to generate new variants to adapt to the volatile environments.

Genetic architecture and evolutionary history of the supergenes have long fascinated biologists. Owning to the long-range sequencing information, the complete *S*-locus consisting of dozens of genes was successfully reconstructed, and our results indicated that *SLF* members within this gene cluster share common *cis*-regulatory elements, ensuring a spatially coordinated expression of protein products to degrade *S-RNase*. Additionally,

the physical position and expression pattern of *AnH01G33501* suggested that it may be a candidate transcription factor regulating downstream *SLFs* expression. Moreover, two miRNAs were considered to act as regulators to control expression of this transcription factor[43]. Based on calculation of Ks of *SLFs*, we found that the ancestral *SLF* had an ancient origin. During the long evolutionary process, different ancestral *SLFs* faced different fate (deletion, expansion, contraction, pseudogenization) in different lineages. Frequent and rapid acquisition of new *SLFs* were adapted to the allelic diversity of *S-RNase*[12] and contributed to the complexity of the extant *Antirrhinum* *S*-locus supergene. Duplication type analysis of all the extant *SLFs* revealed that most of them originated earlier than two WGDs in Plantaginaceae through tandem or proximal duplication, yet the flanking genes of the *S*-locus are still expanding gradually. Thus, the snapdragon *S*-locus supergene has charm of both old and young.

Structural variations such as translocations and inversion are considered drivers of genome evolution and speciation[44, 45]. These rearrangements have likely contributed to the divergence of these species because they can lead to reproductive isolation if they are large enough to interfere with chromosome pairing and recombination during meiosis[46]. By comparing sequence of two *S*-haplotypes, we speculated such an inversion at this locus may be involved in recombination suppression, thus maintain the integrity and specificity of every single haplotype. Inter-specific and intra-specific *S*-haplotype comparison

revealed continuous dynamic changes exist in the snapdragon *S*-locus supergene, and the hyper-divergence may reveal the cause of reproductive isolation and speciation. These findings further enriched our understanding of the *S*-locus from a genome-wide perspective. Ultimately, our study provides unprecedented insights into the genome dynamics of this ornamental flower and offers abundant genomic resource for studies on *Antirrhinum*. A combination of advanced sequencing technologies, comparative genomics, and multi-omics datasets will help further decipher the genetic mechanisms of the *S*-locus evolution, thereby expediting the processes of horticulture and genetic research in the future.

Materials and methods

Sample collection and library construction

All plant samples were grown and collected from green house. For *de novo* assemblies, fresh young leaf tissues of 3 *Antirrhinum hispanicum* lines (AhS_7S_8, AhS_7Sc, and AhS_8Sc), *Antirrhinum linkianum,* and *Misopates orontium* were collected for DNA extraction using cetyltrimethylammonium bromide (CTAB) method. PCR-free paired-end DNA libraries of ~450bp insert size for each individual were prepared and then sequenced on an Illumina HiSeq4000 platform, following the manufacturer's instructions. For PacBio Sequel II sequencing, high molecular weight (HMW) genomic DNA was extracted to construct SMRTbell libraries with insert size of 20kb, using

SMRTbell Express Template Prep Kit 2.0 (PacBio, #100-938-900) and sequenced on PacBio Sequel II system. Another library of 15kb size was sequenced to generate high-qualitied consensus reads (HiFi reads) for AhS_7S_8. For BioNano optical maps sequencing, HMW genomic DNA of AhS_7S_8 were extracted using BioNanoPrepTM Plant DNA Isolation Kit (BioNano Genomics), following BioNano Prep Plant Tissue DNA Isolation Base Protocol (part # 30068). Briefly, fresh leaves were collected and fixed with formaldehyde, then followed by DNA purification and extraction. Prepared nuclear DNA was labeled using BspQ1 as restriction enzyme with BioNano Prep Labeling Kit (Berry Genomics, Beijing). The fluorescently labelled DNA was stained at room temperature and then loaded onto a Saphyr chips to scan on the BioNano Genomics Saphyr System by the sequencing provider Berry Genomics Corporation (Beijing, China).

To obtain a chromosome-level genome, Hi-C library was constructed for AhS_7S_8. Young leaves were fixed with formaldehyde and lysed, then cross-linked DNA samples were digested with restriction enzyme DpnII overnight. After repairing sticky-ends and reversing the cross-linking, DNA fragments were purified, sheared, enriched and sequenced on Illumina NovaSeq6000 under paired-end mode. Four representative tissues, including leaf, petal, pollen, and stamen, of three species, plus pollen and pistil of the lines AhS_7S_C and AhS_8S_C were collected in the green house for RNA-seq. Two biological replicates were prepared for pistil and pollen. Strand-specific RNA-seq

libraries were prepared following the manufacturer's recommended standard

protocol, and then sequenced on Illumina platform under paired-end mode. At

least 20 million paired-end reads were generated for each library. Small RNAs

were separated from total RNA described above using PAGE gel, then

followed adapter ligation, reverse transcription, PCR product purification, and

library quality control. Samples were sequenced using Nextseq500 (50 bp

single-end) to yield 20 million reads per sample. Same tissues were also used

for whole-genome bisulfite sequencing. Genomic DNA was extracted and

fragmented into ~450bp, then treated with bisulfite using the EZ DNA

Methylation-Gold Kit according to manufacturer's instructions. The libraries

were sequenced on a HiSeq4000 system and at least 50 million PE-reads for

each library were generated.

Genome assembly and evaluation

To estimate the genome size independent of assembly, we characterized the

basic genome feature using high-quality Illumina short reads. Firstly, PE reads

were trimmed using Trim_galore (v0.6.1)[47] with parameters "-q25 --stringency

3" to remove low-quality bases and adapters. Based on the 21-mer spectrum

derived from clean data, heterozygosity rate and haploid genome length of

three species were evaluated using software GenomeScope2[48] .

For line AhS_7S_8, a consensus assembly consisting of collapsed sequences,

and haplotype-resolved assemblies consisting of scaffolds from phased allele,

were generated. Firstly, we removed reads of poor quality (RQ < 0.8) and

short length (< 1kb) in PacBio subreads from the 20kb library before assembly.

The contig-level assembly was performed on filtered subreads using Canu

package (v1.9)[49] with parameters "genomeSize=550m corMinCoverage=2

corOutCoverage=200 'batOptions = -dg 3 -db 3 -dr 1 -ca 500 -cp 50'". After

that, alternative haplotig sequences were removed using purge_dups[50], only

primary contigs were kept for further scaffolding. The draft genomes of

A.linkianum and *M.orontium* were assembled using the same procedure. On

the other hand, to obtain haplotype-resolved genome assemblies of AhS_7S_8,

~40X PacBio HiFi reads along with Hi-C data were utilized to separate

parental allele using FALCON-phase[51]. The resulting two phased contig-level

haplotigs were used for downstream analysis independently. The contig

assemblies were scaffolded into pseudo-chromosomes following the same

pipeline described below (Fig. S1).

Low-quality BioNano optical molecular maps with length shorter than 150kb or

label density less than 9 per 100kb were removed. After that, the BioNano

genome maps combining with contigs were fed into BioNano Solve hybrid

scaffolding pipeline v3.6 (BioNano genomics) to produce hybrid scaffolds

under non-haplotype aware mode. When there were conflicts between the

sequence maps and optical molecular maps, both were cut at the conflict site

and assembled with parameters "-B2 -N2". Followed by BioNano scaffolding,

Hi-C data was utilized to cluster contigs into 8 groups subsequently. Clean Hi-

C reads were then mapped to the BioNano-derived scaffolds using BWA[52].

The chromosome-level assembly was furtherly generated using the 3D-DNA

analysis pipeline to correct mis-join, order, orientation, and translocation from

the assembly[53]. Manual review and refinement of the candidate assembly was

performed in Juicerbox Assembly Tools (v1.11.08)[54] for interactive correction.

Then the long pseudo-chromosomes were regenerated using script "run-asm-

pipeline-post-review.sh -s finalize --sort-output --bulid-gapped-map" in 3D-

DNA package with reviewed assembly file as input. Three chromosome-level

assemblies of AhS_7S_8 were constructed, and we named them as consensus-

assembly SI-Ah (the mosaic one), S_7 haplome and S_8 haplome encompassing

S_7 haplotype and S_8 haplotype, respectively). The sequences of two

haplomes were named as Chr*.1 and Chr*.2 to distinguish homologous

chromosomes. After manual curation, we used ultra-high-depth PacBio

dataset of two offsprings and PBJelly[55] to fill gaps in haplome assemblies,

followed by two rounds of polish with gcpp program

(https://github.com/PacificBiosciences/gcpp) in pbbioconda package.

The organellar genome of *Antirrhinum* were identified using BLASTn against

the *A.hispanicum* contigs with plant plastid sequences as reference

(https://www.ncbi.nlm.nih.gov/genome/organelle/). The organellar genome

sequences were submitted to CHLOROBOX[56] website to annotate

automatically and visualize genome maps.

The continuity and completeness of genome assemblies were benchmarked

using long terminal repeat (LTR) Assembly Index (LAI)[16] and BUSCO

embryophyta_odb10[57] core genesets, which include 2121 single-copy

orthologous genes. Also, the PacBio reads and Illumina PE reads were

mapped to respective assemblies using minimap2 and BWA[52], to evaluate

correctness. MUMmer package[58] was used to inspect the difference between

pair-wise genome assemblies. SNV (SNPs and InDels) were called using

show-snps with parameters '-rITHC', then parsed and collapsed adjacent

mutation/variation with our custom python script. While structural variation

(SVs) including duplication, deletion, inversion, and translocation between two

phased assemblies were called based on reciprocal alignment from nucmer

and show-coords.

Annotation of repetitive DNA elements

To identify repetitive elements within each genome sequences, firstly we used

RepeatModeler (open-1.0.11) to build a *de novo* repetitive elements library

from the assembled genome sequences independently. GenomeTools suite[59]

was used to annotate LTR-RTs with protein HMMs from the Pfam database. In

addition, LTR_retriever[60] was used to utilize output of GenomeTools and

LTR_FINDER[61] to generate whole-genome LTR annotation. Parameters of

LTR_FINDER and LTRharvest were set requiring minimum and maximum

LTR length of 100 bp and 7 kb, respectively.

Each of these libraries were classified with RepeatClassifier, followed by

combining and removing redundancy using USERACH (https://www.drive5.com/usearch/). Then the unclassified sequences library was analyzed using BLASTX to search against transposase and plant protein databases to remove putative protein-coding genes. We submitted unknown repetitive sequences to CENSOR (https://www.girinst.org/censor/index.php) to get further classified and annotated. *De-novo* searches for miniatures inverted repeat transposable element (MITEs) used MITE_Hunter[62] software with parameters "-w 2500 -n 5 -L 80 -I 80 -m 2 -S 12345678 -P 1". Finally, all genome assemblies were soft-masked in company with the repeat library using RepeatMasker (open-4-0-9)[63] with parameters "-div 22 -cutoff 220 - xsmall".

Gene prediction and functional annotation

The gene prediction was performed using BRAKER2[64] annotation pipeline with parameters '--etpmode --softmasking', which integrates RNA-seq datasets and evolutionary related protein sequences, as well as *ab initio* prediction results. Clean RNA-seq reads were aligned to the repeat-mask genomes using STAR[65], then converted and merged to BAM format. The hint files were incorporated with *ab initio* prediction tool AUGUSTUS[66] and GeneMark-ETP[67] to train gene models. The predicted genes with protein length < 120 or TPM < 0.5 in any RNA-seq sample were removed. The *SLF* gene models included in the *S*-locus were examined in Genome browser and

manually curated.

The tRNA genes were annotated by homologous searching against the whole genome using tRNAscan-SE(v2.0)[68], rRNA and snRNA genes were identified using cmscan program in INFERNAL(v1.1.2)[69] package to search from the Rfam database. Four small-RNA sequencing datasets were collapsed, combining with known miRNA sequence download from mirBase(v22)[70] as input of miRDP2(v1.1.4)[71] to identify miRNA genes across the genome. Next, miRNA mature sequences were extracted to predict potential target gene using miRANDA (http://www.microrna.org) by searching against all predicted full-length cDNA sequences.

To perform functional analysis of the predicted gene models, protein sequences were search against the InterPro consortium databases including PfamA, PROSITE, TIGRFAM, SMART, SuperFamily, and PRINTS as well as Gene Ontology database and pathways (KEGG, Reactome) databases using InterProScan (v5.51-85.0)[72]. The protein sequences were also submitted to plantTFDB online server[73] to identify transcription factors and assign TFs families. In addition, predicted genes were also annotated biological function using diamond[74] to search against NCBI non-redundant protein and UniProtKB (downloaded on 21st Mar 2021, UniProt Consortium)[75] database with an e-value of 1e-5. Genes were annotated to the eggnog database using eggNOG-mapper[76].

Whole-genome duplication and intergenomic analysis

Syntenic blocks (at least 5 genes per block) of *A.hispanicum* were identified using MCscan (Python version)[77] with default parameters. Intra-genome syntenic relationships were visualized using Circos[78] in Fig1. We also compared *A.hispanicum* genome with several plant genomes, including *S.miltiorrhiza*, *S.lycopersicum*, *A.coerulea*, and *V.vinifera*. Dotplots for genome pairwise synteny was generated using the command 'python -m jcvi.graphics.dotplot --cmap=Spectral --diverge=Spectral'.

For the *Ks* plots, we used wgd package[79] with inflation factor of 1.5 and ML estimation times of 3 for reciprocal best hits search and MCL clustering. The synonymous substitution rate (*Ks*) for paralogs and orthologs were calculated using codeml program of PAML package[80]. We plotted output data from result files using custom python scripts with Gaussian mixture model function to fit the distribution (1-5 components) and determined peak *Ks* values. Estimated mean peak values were used for dating WGD events.

Comparative genomics and divergence time estimation

Orthologous genes relationships were built based on the predicted proteomes deprived from consensus assembled *A.hispanicum* and other angiosperm species listed in (Table S11) using OrthoFinder2[81]. Only longest protein sequences were selected to represent each gene model. Rooted species tree inferred by OrthoFinder2 using STRIDE[82] and STAG[83] algorithm was used as

a starting species tree for downstream analysis.

The species divergence time was estimated using MCMCtree in PAML[80] package with branch lengths estimated by BASEML, with *Amborella* as outgroup. The Markov chain Monte Carlo (MCMC) process consists of 500,000 burn-in iterations and 400,000 sampling iterations. The same parameters were executed twice to confirm the results were robust. Species divergence time for *Amborella trichopoda- Arabidopsis thaliana* (173-199 Mya), *Vitis vinifera-Petunia axillaris* (111-131 Mya) and *Solanum lycopersicum–Solanum tuberosum* (5.23-9.40 Mya) which were obtained from TimeTree database[84] were used to calibrate the estimation model, and constrained the root age <200 Mya.

To determine the gene family expansion and contraction, orthologous genes count table and phylogenetic species tree topology inferred by OrthoFinder2 were taken into latest CAFE5[85], which employed a random birth and death model to determine expansion and contraction in gene families of given node. Corresponding p value was provided for each node and branch of phylogeny tree and cutoff 0.05 was used to identify gene families undergo expanded or contraction significantly at a specific lineage or species. KEGG and GO enrichment analysis of expanded gene family members were performed with Fisher's exact test and adjusted by the Benjamini-Hochberge method to identify significantly enriched terms.

Gene expression analysis

Apart from aiding gene model prediction, RNA-seq datasets were also used for transcripts quantification. For expression analysis, we used STAR[65] to map clean RNA-seq reads to reference genomes with parameters '--quantMode TranscriptomeSAM --outSAMstrandField intronMotif --alignIntronMax 6000 --alignIntronMin 50'. The transcripts per million (TPM) values were obtained using expectation maximization tool rsem[86]. The reproducibility of RNA-seq samples was assessed using spearman correlation coefficient. Samples from same tissue display strong correlation with a Pearson's correlation coefficient of $r > 0.85$ (Fig. S29), indicating good reproducibility. Owing to haplotype-resolved assembly and the gene structure annotation of *A.hispanicum* genome, allelic genes from a same locus can be identified using a synteny-based strategy along with identity-based method. Reciprocal best hit between two haplotypes were identified using MCSCAN[77] at first, and genes not in synteny block were searched against coding sequence counterparts to fill up the allelic relationship table. We used MUSCLE[87] to align coding sequences of two allelic genes, and then calculated Levenshtein edit distance to measure allelic divergence. The divergence rate was defined as the number of edit distance divided by the total length of aligned bases.

Allelic transcripts quantification was conducted using cleaned RNA-seq datasets and allele-aware tool kallisto v.0.46.0[88]. We applied this software to obtain the expression levels read counts and TPM of genes of both

1 haplotypes. Differential expression analysis of allelic genes was performed

2 using R package edgeR[89]. Cutoff criteria of allele imbalanced expressed

3 genes were set as adjusted P value < 0.01, false discovery rate < 0.01 and

4 $|\log_2(FC)|>1$.

Phylogenetic analyses of genes

F-box genes in whole genome were identified by searching Interproscan

annotation results. Sequences annotated with both PF00646 (F-box domain)

and PF01344 (Kelch motif) or PF08387 (FBD domain) or PF08268 (F-box

associated domain) were considered as F-box genes. And the F-box genes

located in the S-locus region of snapdragon were considered as potential

SLFs. Organ specific protein and paralogs were selected using PF10950 as

keyword, while PF02298 for plastocyanin gene. Protein sequence alignments

were constructed using MUSCLE[87] and manually checked. The maximum

likelihood phylogenetic gene trees were constructed using raxml-ng[90] with 100

replicates of bootstrap and parameter '-m MF'. And duplicate gene were

classified into different categories using DupGen_finder[24] with parameters '-d

15'.

Cis-regulatory element analysis of predicted *SLFs* promoters

20 *Cis*-regulatory elements are specific DNA sequences that are located

upstream of gene coding part and involved in regulation of genes expression by binding with transcription factors (TFs). We explored the upstream 2000bp sequences of 32 *SLFs* of *A.hispanicum* to discover TF binding sites using MEME[91], and Tomtom was used for comparison against JASPAR database[92] of the discovered motif.

Whole-genome bisulfite sequencing analysis

The raw WGBS reads were processed to remove adapter sequences and low-quality bases using Trim_galore with default parameters. The cleaned reads were then mapped to the two haplomes using abismal command from MethPipe[93] package with parameters '-m 0.02'. All reads were mapped to the chloroplast genome (normally unmethylated) of snapdragon to estimate bisulfite conversion rate. The non-conversion ratio of chloroplast genome Cs to Ts was considered ad a measure of error rate.

Each cytosine of sequencing depth $\geqslant 5$ were seen as true *methylcytosines* sites. Methylation level at every single *methylcytosine* site was estimated as a probability based on the ratio of methylated to the total reads mapped to that loci. Methylation level in genes and 2kb flanking regions was determined using custom Python scripts. Gene body refers to the genomic region from start to stop codon coordinates in the gff file. Each gene and its flanking regions were partitioned into ten bins of equal size and average methylation level in each bin was calculated by dividing the reads indicating methylation

by total reads observed in the respective bin.

Data availability

All raw genome sequencing datasets, assembled genome sequences, and predicted gene models have been deposited at the GSA database[94] in the National Genomics Data Center, Beijing Institute of Genomics (BIG), Chinese Academy of Sciences and China National Center for Bioinformation, under accession numbers PRJCA006918 that are publicly accessible at http://ngdc.cncb.ac.cn/gsa. RNA-seq datasets have been deposited under accession id CRA005238.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (32030007) and the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB27010302).

Conflict of interests

The authors declare no competing interests.

Author contributions

Y.X. designed and supervised the research; Y.E.Z. and Q.Q.H performed the experiments and prepared plant samples for sequencing; L.C. provided plant

seeds; S.Z. analyzed the data; D.Z. contributed data visualization; S.Z., Y.X. and E.C. wrote the manuscript. All authors discussed the results and commented on the final manuscript.

References

1. Takayama, S. & Isogai, A. Self-incompatibility in plants. *Annu. Rev. Plant Biol.* **56**, 467–489 (2005).

2. Kubo, K. *et al.* Collaborative Non-Self Recognition System in S-RNase–Based Self-Incompatibility. *Science* **330**, 796–799 (2010).

3. Potente, G. *et al.* Comparative Genomics Elucidates the Origin of a Supergene Controlling Floral Heteromorphism. *Mol. Biol. Evol.* **39**, msac035 (2022).

4. Hager, E. R. *et al.* A chromosomal inversion contributes to divergence in multiple traits between deer mouse ecotypes. *Science* **377**, 399–405 (2022).

5. Joron, M. *et al.* Chromosomal rearrangements maintain a polymorphic supergene controlling butterfly mimicry. *Nature* **477**, 203–206 (2011).

6. Wang, J. *et al.* A Y-like social chromosome causes alternative colony organization in fire ants. *Nature* **493**, 664–668 (2013).

7. Zhao, H. *et al.* Origin, loss, and regain of self-incompatibility in angiosperms. *Plant Cell* **34**, 579–596 (2022).

8. Igic, B., Lande, R. & Kohn, J. R. Loss of Self‑Incompatibility and Its Evolutionary Consequences. *Int. J. Plant Sci.* **169**, 93–104 (2008).

9. Shimizu, K. K. & Tsuchimatsu, T. Evolution of Selfing: Recurrent Patterns in Molecular Adaptation. *Annu. Rev. Ecol. Evol. Syst.* **46**, 593–622 (2015).

10. Golz, J. F., Clarke, A. E., Newbigin, E. & Anderson, M. A relic S-RNase is expressed in the styles of self-compatible Nicotiana sylvestris. *Plant J.* **16**, 591–599 (1998).

11. Markova, D. N. *et al.* Evolutionary history of two pollen self-incompatibility factors reveals alternate routes to self-compatibility within Solanum. *Am. J. Bot.* **104**, 1904–1919 (2017).

12. Xue, Y., Carpenter, R., Dickinson, H. G. & Coen, E. S. Origin of allelic diversity in antirrhinum S locus RNases. *Plant Cell* **8**, 805–814 (1996).

13. Qiao, H. *et al.* The F-box protein AhSLF-S2 controls the pollen function of S-RNase-based self-incompatibility. *Plant Cell* **16**, 2307–2322 (2004).

14. Yang, Q., Zhang, D., Li, Q., Cheng, Z. & Xue, Y. Heterochromatic and genetic features are consistent with recombination suppression of the self-incompatibility locus in Antirrhinum. *Plant J.* **51**, 140–151 (2007).

15. Li, M. *et al.* Genome structure and evolution of Antirrhinum majus L. *Nat. Plants* **5**, 174–183 (2019).

16. Ou, S., Chen, J. & Jiang, N. Assessing genome assembly quality using the LTR Assembly Index (LAI). *Nucleic Acids Res.* **46**, e126 (2018).

17. Wright, S. I. & Schoen, D. J. Transposon dynamics and the breeding system. *Genetica* **107**, 139 (1999).

18. Otero, A., Fernández-Mazuecos, M. & Vargas, P. Evolution in the Model Genus Antirrhinum Based on Phylogenomics of Topotypic Material. *Front. Plant Sci.* **12**, 631178 (2021).

19. Murat, F., Armero, A., Pont, C., Klopp, C. & Salse, J. Reconstructing the genome of the most recent common ancestor of flowering plants. *Nat. Genet.* **49**, 490–496 (2017).

20. Jaillon, O. *et al.* The grapevine genome sequence suggests ancestral hexaploidization in major angiosperm phyla. *Nature* **449**, 463–468 (2007).

21. Frazee, L. J. *et al.* New genomic resources and comparative analyses reveal differences in floral gene expression in selfing and outcrossing Collinsia sister species. *G3 (Bethesda).* **11**, jkab177 (2021).

22. Jiao, Y. *et al.* A genome triplication associated with early diversification of the core eudicots. *Genome Biol.* **13**, R3 (2012).

23. Igic, B. & Busch, J. W. Is self-fertilization an evolutionary dead end? *New Phytol.* **198**, 386–397 (2013).

24. Qiao, X. *et al.* Gene duplication and evolution in recurring polyploidization-diploidization cycles in plants. *Genome Biol.* **20**, 1–23 (2019).

25. Igic, B. & Kohn, J. R. Evolutionary relationships among self-

incompatibility RNases. *Proc Natl Acad Sci U S A.* **98**, 13167–13171

(2001).

26. Takahata, N. A simple genealogical structure of strongly balanced allelic

lines and trans-species evolution of polymorphism. *Proc. Natl. Acad. Sci.*

87, 2419–2423 (1990).

27. Vekemans, X. & Slatkin, M. Gene and allelic genealogies at a

gametophytic self-incompatibility locus. *Genetics* **137**, 1157–1165

(1994).

28. Debernardi, J. M., Rodriguez, R. E., Mecchia, M. A. & Palatnik, J. F.

Functional specialization of the plant miR396 regulatory network

through distinct microRNA-target interactions. *PLoS Genet.* **8**,

e1002419 (2012).

29. Aydinoglu, F. Elucidating the regulatory roles of microRNAs in maize

(Zea mays L.) leaf growth response to chilling stress. *Planta* **251**, 38

(2020).

30. He, Z. *et al.* Evolution of coastal forests based on a full set of mangrove

genomes. *Nat. Ecol. Evol.* **6**, 738–749 (2022).

31. Wu, G. A. *et al.* Sequencing of diverse mandarin, pummelo and orange

genomes reveals complex history of admixture during citrus

domestication. *Nat. Biotechnol.* **32**, 656–662 (2014).

32. Lu, J. *et al.* The Physalis floridana genome provides insights into the

biochemical and morphological evolution of Physalis fruits. *Hortic. Res.*

8, 244 (2021).

33. Cao, K. *et al.* Chromosome-level genome assemblies of four wild peach species provide insights into genome evolution and genetic basis of stress resistance. *BMC Biol.* **20**, 139 (2022).

34. Steinbachs, J. E. & Holsinger, K. E. S-RNase-mediated gametophytic self-incompatibility is ancestral in eudicots. *Mol. Biol. Evol.* **19**, 825–829 (2002).

35. Kubo, K. I. *et al.* Gene duplication and genetic exchange drive the evolution of S-RNase-based self-incompatibility in Petunia. *Nat. Plants* **1**, (2015).

36. Fujii, S., Kubo, K. I. & Takayama, S. Non-self- and self-recognition models in plant self-incompatibility. *Nat. Plants* **2**, 16130 (2016).

37. Zhou, Q. *et al.* Haplotype-resolved genome analyses of a heterozygous diploid potato. *Nat. Genet.* **52**, 1018–1023 (2020).

38. Ramanauskas, K. & Igić, B. RNase-based self-incompatibility in cacti. *New Phytol.* **231**, 2039–2049 (2021).

39. Gibney, E. R. & Nolan, C. M. Epigenetics and gene expression. *Heredity (Edinb).* **105**, 4–13 (2010).

40. Vargas, P., Carrió, E., Guzmán, B., Amat, E. & Güemes, J. A geographical pattern of Antirrhinum (Scrophulariaceae) speciation since the Pliocene based on plastid and nuclear DNA polymorphisms. *J. Biogeogr.* **36**, 1297–1312 (2009).

41. Goodwillie, C. *et al.* Correlated evolution of mating system and floral display traits in flowering plants and its implications for the distribution of mating system variation. *New Phytol.* **185**, 311–321 (2010).

42. Willi, Y. & Määttänen, K. Evolutionary dynamics of mating system shifts in Arabidopsis lyrata. *J. Evol. Biol.* **23**, 2123–2131 (2010).

43. Finnegan, E. J., Liang, D. & Wang, M.-B. Self-incompatibility: Smi silences through a novel sRNA pathway. *Trends Plant Sci.* **16**, 238–241 (2011).

44. Dodsworth, S., Leitch, A. R. & Leitch, I. J. Genome size diversity in angiosperms and its influence on gene space. *Curr. Opin. Genet. Dev.* **35**, 73–78 (2015).

45. Bowers, J. E., Chapman, B. A., Rong, J. & Paterson, A. H. Unravelling angiosperm genome evolution by phylogenetic analysis of chromosomal duplication events. *Nature* **422**, 433–438 (2003).

46. Lamichhaney, S. *et al.* Structural genomic changes underlie alternative reproductive strategies in the ruff (Philomachus pugnax). *Nat. Genet.* **48**, 84–88 (2015).

47. Krueger, F. Trim Galore!: A wrapper tool around Cutadapt and FastQC to consistently apply quality and adapter trimming to FastQ files. *Babraham Inst.* (2015).

48. Ranallo-Benavidez, T. R., Jaron, K. S. & Schatz, M. C. GenomeScope 2.0 and Smudgeplot for reference-free profiling of polyploid genomes.

Nat. Commun. **11**, 1432 (2020).

49. Koren, S. *et al.* Canu: Scalable and accurate long-read assembly via adaptive κ-mer weighting and repeat separation. *Genome Res.* **27**, 722–736 (2017).

50. Guan, D. *et al.* Identifying and removing haplotypic duplication in primary genome assemblies. *Bioinformatics* **36**, 2896–2898 (2020).

51. Kronenberg, Z. N. *et al.* Extended haplotype-phasing of long-read de novo genome assemblies using Hi-C. *Nat. Commun.* **12**, 1935 (2021).

52. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).

53. Olga, D. *et al.* De novo assembly of the Aedes aegypti genome using Hi-C yields chromosome-length scaffolds. *Science* **356**, 92–95 (2017).

54. Durand, N. C. *et al.* Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments. *Cell Syst.* **3**, 95–98 (2016).

55. English, A. C. *et al.* Mind the Gap: Upgrading Genomes with Pacific Biosciences RS Long-Read Sequencing Technology. *PLoS One* **7**, e47768 (2012).

56. Tillich, M. *et al.* GeSeq - Versatile and accurate annotation of organelle genomes. *Nucleic Acids Res.* **45**, W6–W11 (2017).

57. Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M. BUSCO: Assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**, 3210–3212

(2015).

58. Marçais, G. *et al.* MUMmer4: A fast and versatile genome alignment system. *PLoS Comput. Biol.* **14**, 1–14 (2018).

59. Gremme, G., Steinbiss, S. & Kurtz, S. Genome tools: A comprehensive software library for efficient processing of structured genome annotations. *IEEE/ACM Trans. Comput. Biol. Bioinforma.* **10**, 645–656 (2013).

60. Ou, S. & Jiang, N. LTR_retriever: A highly accurate and sensitive program for identification of long terminal repeat retrotransposons. *Plant Physiol.* **176**, 1410–1422 (2018).

61. Xu, Z. & Wang, H. LTR-FINDER: An efficient tool for the prediction of full-length LTR retrotransposons. *Nucleic Acids Res.* **35**, W265–W268 (2007).

62. Han, Y. & Wessler, S. R. MITE-Hunter: A program for discovering miniature inverted-repeat transposable elements from genomic sequences. *Nucleic Acids Res.* **38**, 1–8 (2010).

63. Smit, A., Hubley, R. & Green, P. RepeatMasker Open-4.0. *http://www.repeatmasker.org* (2013).

64. Brůna, T., Hoff, K. J., Lomsadze, A., Stanke, M. & Borodovsky, M. BRAKER2: automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein database. *NAR genomics Bioinforma.* **3**, lqaa108 (2021).

65. Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

66. Stanke, M. *et al.* AUGUSTUS: ab initio prediction of alternative transcripts. *Nucleic Acids Res.* **34**, W435–W439 (2006).

67. Brůna, T., Lomsadze, A. & Borodovsky, M. GeneMark-EP+: eukaryotic gene prediction with self-training in the space of genes and proteins. *NAR Genomics Bioinforma.* **2**, 1–14 (2020).

68. Lowe, T. M. & Eddy, S. R. tRNAscan-SE: A Program for Improved Detection of Transfer RNA Genes in Genomic Sequence. *Nucleic Acids Res.* **25**, 955–964 (1997).

69. Nawrocki, E. P. & Eddy, S. R. Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics* **29**, 2933–2935 (2013).

70. Kozomara, A., Birgaoanu, M. & Griffiths-Jones, S. MiRBase: From microRNA sequences to function. *Nucleic Acids Res.* **47**, D155–D162 (2019).

71. Kuang, Z., Wang, Y., Li, L. & Yang, X. MiRDeep-P2: Accurate and fast analysis of the microRNA transcriptome in plants. *Bioinformatics* **35**, 2521–2522 (2019).

72. Quevillon, E. *et al.* InterProScan: Protein domains identifier. *Nucleic Acids Res.* **33**, W116–W120 (2005).

73. Jin, J. *et al.* PlantTFDB 4.0: Toward a central hub for transcription factors and regulatory interactions in plants. *Nucleic Acids Res.* **45**,

D1040–D1045 (2017).

74. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND. *Nat. Methods* **12**, 59–60 (2014).

75. Bateman, A. *et al.* UniProt: The universal protein knowledgebase in 2021. *Nucleic Acids Res.* **49**, D480–D489 (2021).

76. Huerta-Cepas, J. *et al.* EggNOG 5.0: A hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Res.* **47**, D309–D314 (2019).

77. Tang, H. *et al.* Synteny and collinearity in plant genomes. *Science* 320 486–488 (2008).

78. Krzywinski, M. *et al.* Circos: An information aesthetic for comparative genomics. *Genome Res.* **19**, 1639–1645 (2009).

79. Zwaenepoel, A. & Van De Peer, Y. Wgd-simple command line tools for the analysis of ancient whole-genome duplications. *Bioinformatics* **35**, 2153–2155 (2019).

80. Yang, Z. PAML 4: Phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24**, 1586–1591 (2007).

81. Emms, D. M. & Kelly, S. OrthoFinder: Phylogenetic orthology inference for comparative genomics. *Genome Biol.* **20**, 238 (2019).

82. Emms, D. M. & Kelly, S. STRIDE: Species tree root inference from gene duplication events. *Mol. Biol. Evol.* **34**, 3267–3278 (2017).

83. Emms, D. M. STAG: Species Tree Inference from All Genes. *bioRxiv*

(2018) doi:10.1101/267914.

84. Kumar, S., Stecher, G., Suleski, M. & Hedges, S. B. TimeTree: A Resource for Timelines, Timetrees, and Divergence Times. *Mol. Biol. Evol.* **34**, 1812–1819 (2017).

85. Mendes, F. K., Vanderpool, D., Fulton, B. & Hahn, M. W. CAFE 5 models variation in evolutionary rates among gene families. *Bioinformatics* **36**, 5516–5518 (2020).

86. Li, B. & Dewey, C. N. RSEM: Accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

87. Edgar, R. C. MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* **32**, 1792–1797 (2004).

88. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

89. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: A Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

90. Kozlov, A. M., Darriba, D., Flouri, T., Morel, B. & Stamatakis, A. RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* **35**, 4453–4455 (2019).

91. Bailey, T. L. *et al.* MEME Suite: Tools for motif discovery and searching.

Nucleic Acids Res. **37**, 202–208 (2009).

92. Fornes, O. *et al.* JASPAR 2020: Update of the open-Access database of transcription factor binding profiles. *Nucleic Acids Res.* **48**, D87–D92 (2020).

93. Song, Q. *et al.* A reference methylome database and analysis pipeline to facilitate integrative and comparative epigenomics. *PLoS One* **8**, e81148 (2013).

94. Database Resources of the National Genomics Data Center, China National Center for Bioinformation in 2022. *Nucleic Acids Res.* **50**, D27–D38 (2022).

Fig.1 | Genomic features of *A.hispanicum*. The outermost layer is circular

ideogram of the eight pseudomolecules. Track **b-j** corresponds to the

distribution of the GCcontent (red denote higher than average whole-genome

level, and green vice versa), Ty3/Gypsy, Ty1/Copia retrotransposons (higher

density shown in darker green), gene density, methylation level, expression

level of four tissues (petal, pollen, pistil, and leaf) respectively, all calculated in

250kb non-overlap window. Track **k** corresponds to A/B compartments

inferred by using Hi-C data. Track **l** corresponds to SNP/InDel density

distribution. Track **m** corresponds to location of miRNA genes. Track **n** corresponds to location of all F-box genes, blue and red represent forward and reverse strand respectively. Track **o** marker "T" corresponds to location of ribonuclease T2 genes. Track **p** corresponds to inter-chromosome syntenic blocks. The grey sector marked *S*-locus in Chr8.

Fig. 2 | Comparative genomics analysis and lineage expanded gene

families of *Antirrhinum*. A Phylogeny position of the snapdragon and gene

family evolution. The left panel shows phylogenetic tree of 18 species. The

black star denotes the Plantaginaceae-specific WGD. The pie diagram on each branch of the tree represents the proportion of gene families undergoing expansion (red) or contraction (blue) events. The right panel shows the distribution of single copy, two copy, multiple copy, unique paralogs, and other orthologs in plant species. **B** Paranome *Ks* distribution of three Plantaginaceae members and *Salvia miltiorrhiza*. **C** *Ks* distribution of one-to-one orthologs identified between *A.hispanicum* and other species. The density curves were fitted using Gaussian mixture model, and the peak values marked with dotted dash lines. **D-E** Gene Ontology enrichment analysis of expanded family members in *Antirrhinum* lineage and *M.orontium* branch. The color of circle represents the scaled odds ratio. The size of circle represents the gene number of GO terms.

Fig.3 | Identification of a candidate regulator of *SLFs*. A Phylogenetic tree

reconstructed based on the amino acid sequences of all F-box genes from

A.hispanicum genome. The maximal likelihood phylogenetic tree was calcu-

lated using RAxML-NG with 1000 bootstrap. The color of gene ID denoted the

chromosome the gene within as indicated in the legend. Heatmap color repre-

sent Z-scores derived from RNA-seq expression data for each gene. *SLF*

genes are marked with black stars. **B** The red line is synonymous substitution

rate distribution of 32 *SLFs*, while the green one denotes other non-*SLF* F-box

genes in the genome. **C** The bottom panel represents motif found by scanning

upstream 1000bp of 32 *SLFs* using MEME, and the top panel is a Tomtom

alignment for annotation the putative transcription factor binding site. **D** The

cluster heatmap of expression matrix of all MYB-related transcription factor

family members in SI-Ah genome. The columns represent 4 tissues. The leg-

end shows the range of log2(TPM+0.5), and TPM is mean value of two biolog-

ical replicates. Genes marked in black box may be involved in controlling the

expression of *SLFs*, since they are expressed in pollen but not in pistil. **E**

Schematic structure and regulatory relationship of snapdragon *S*-locus.

Fig.4 | The correlation of position and similarity of *SLFs*. A Heatmap

showing sequence identity (lower left triangle) and sequence similarity (upper

right triangle) between *SLFs* in SI-Ah. The genes in x-axis and y-axis are

ordered by physical position on the chromosome. **B** Pairwise Ka/Ks analysis

for all *SLF*s and ribonuclease gene in the *S*-locus between SI-Ah and *A.majus,*

as well as *A.linkianum*. Genes are ordered by genomic coordinates, and label

of ribonuclease gene is highlighted in red.

Fig. 5 | Evolution of the *Antirrhinum S*-locus supergene. A Microsynteny

of the S-locus between *A.hispanicum* with other four species (*Chiococca alba*,

Citrus clementina, *Physalis pubescens*, and *Prunus davidiana*). The red arrow

denotes the *S-RNase* of *A.hispanicum*. **B-D** The micro collinearity patterns of

S-haplotype in *A.majus*, *M.orontinum*, and *A.linkianum*, with S_7-haplotype

from *A.hispanicum*. **E** The microsynteny of S_7-haplotype vs. S_8-haplotype. The

green curve lines and the red lines denote *SLFs* and *RNase* gene respectively,

other genes are colored grey. The Tam downstream 11kb of *S-RNase* is

marked by a blue arrow, and the TE is on the same strand with *S-RNase*. **F**

Possible formation process of the extant *Antirrhinum* S-locus. The color of

SLFs represents newly evolved ones to deal with the evolution of the *S-*

RNase with new specificity.

Fig.6 | Allelic expression and methylation profile of *SLFs* for *AhS₇S₈* of

***A.hispanicum*. A** The distribution of allelic divergence of genes. The

divergence was measured using Levenshtein edit distance. **B** Allelic

expression heatmap of *SLFs* in three genotypes. Every two columns

represent allelic gene pairs of two *S*-haplotypes (HF: pollen, HZ: pistil). *SLF*

starts with 'p' indicated pseudo- F-box gene lack of F-box domain or F-box

protein interaction domain, but they are the best hit of the counterpart F-box

allele. **C** Methylation level within group1 (higher expressed) and group2 (lower

expressed) *SLFs* and their 2kb flanking (upstream and downstream) regions

in CpG, CHG, and CHH context. Each region was divided into 10 bins of equal

size and the methylation level in each bin is displayed in the line graph.