Th1 cells alter the inflammatory signature of IL-6 by channeling STAT transcription factors to *Alu*-like retroelements

David Millrine[1,2,3], Ana Cardus Figueras[1,2], Javier Uceda Fernandez[1,2], Robert Andrews[1,2], Barbara Szomolay[1,2], Benjamin C Cossins[1,2], Christopher M. Rice[4], Jasmine Li[5], Victoria J Tyrrell[1,2], Louise McLeod[6,7], Peter Holmans[8], Valerie B O'Donnell[1,2], Philip R Taylor[1,2,9], Stephen J. Turner[5], Brendan J. Jenkins[6,7], Gareth W Jones[4], Nicholas Topley[1,2], Nigel M Williams[8], & Simon A Jones[1,2*]

Affiliations–

[1]Division of Infection & Immunity, School of Medicine, Cardiff University, Cardiff, Wales, UK

[2]Systems Immunity University Research Institute, Cardiff University, Cardiff, Wales, UK

[3]Medical Research Council Protein Phosphorylation & Ubiquitylation Unit (MRC-PPU), School of Life Sciences, University of Dundee, Dundee UK

[4]School of Cellular & Molecular Medicine, University of Bristol, Bristol, UK

[5]Department of Microbiology, Biomedicine Discovery Institute, Monash University, Clayton, Victoria 3800, Australia

[6]Centre for Innate Immunity & Infectious Diseases, Hudson Institute of Medical Research, Clayton, Victoria 3168, Australia

[7]Department of Molecular translational Science, Faculty of Medicine, Nursing & Health Sciences, Monash University, Clayton, Victoria 3800, Australia.

[8]Division of Psychological Medicine & Clinical Neuroscience, School of Medicine, Cardiff University, Cardiff, Wales, UK.

[9]UK Dementia Research Institute at Cardiff, Cardiff University, Cardiff, Wales, UK

*Corresponding author–

Professor Simon A. Jones, Division of Infection & Immunity, School of Medicine, Cardiff University, Heath Park, Cardiff, CF14 4XN, Wales, UK.

Tel: +44 29 2068 7325; Fax: +44 29 2068 7303; E-mail: JonesSA@cf.ac.uk

Abstract –

Cytokines that signal *via* STAT1 and STAT3 transcription factors instruct decisions affecting tissue homeostasis, anti-microbial host defense, and inflammation-induced tissue injury. To understand the coordination of these activities, we applied RNA-seq, ChIP-seq, and ATAC-seq to identify the transcriptional output of STAT1 and STAT3 in peritoneal tissues during acute resolving inflammation and inflammation primed to drive fibrosis. Bioinformatics focussed on the transcriptional signature of IL-6 in both settings and examined how pro-fibrotic IFNγ-secreting CD4[+] T-cells altered the interpretation of STAT1 and STAT3 cytokine cues. In resolving inflammation, STAT1 and STAT3 cooperated to drive stromal gene expression affecting anti-microbial immunity and tissue homeostasis. The introduction of IFNγ-secreting CD4[+] T-cells altered this transcriptional program and channeled STAT1 and STAT3 to a previously latent GAS motif in *Alu*-like elements. STAT1 and STAT3 binding to this conserved sequence revealed evidence of reciprocal cross-regulation and gene signatures relevant to pathophysiology. Thus, effector T-cells re-tune the transcriptional output of IL-6 by shaping a regulatory interplay between STAT1 and STAT3 in inflammation.

Key words –

Cytokines, Inflammation, Fibrosis, Transcription Factors, Retroelements, Retrotransposons

Introduction –

Cytokines are essential for development, tissue homeostasis, and the regulation of inflammation (Dinarello, 2007). The intracellular signaling pathways controlling these activities are intrinsically linked, and aberrant host defense compromises tissue integrity and physiological function. Patients treated with anti-cytokine therapies often provide evidence of this relationship (Choy *et al*, 2013). For example, interleukin (IL)-6 inhibition is less effective in diseases where IL-6 maintains tissue homeostasis and barrier immunity (Andrews *et al*, 2018; Harbour *et al*, 2020; Hunter & Jones, 2015; Jones & Jenkins, 2018; Jones *et al*, 2011; Navarini *et al*, 2011; Schett *et al*, 2013; Taniguchi *et al*, 2015). How IL-6 operating through a single receptor signaling cassette coordinates the maintenance of tissue physiology and the transition to pathophysiology is unknown.

Interleukin-6 regulates cellular responses *via* receptor activation of Janus kinases (Jak) and members of the Signal Transducer and Activator of Transcription (STAT) family (Hunter & Jones, 2015; Jones & Jenkins, 2018). While IL-6 employs other signaling intermediates, IL-6 instructs cell decisions primarily through STAT1 and STAT3 transcription factors (Costa-Pereira *et al*, 2002; Hirahara *et al*, 2015; Jones *et al*, 2013; Peters *et al*, 2015; Twohig, 2019; Villarino *et al*, 2017; Wilmes *et al*, 2021). These proteins share a complex regulatory interplay, and gene ablation studies show that STAT1 and STAT3 often counteract each other or engage shared enhancers (Avalle *et al*, 2012; Hunter & Jones, 2015; Martinez-Fabregas *et al*, 2020; Vahedi *et al*, 2015). These interactions re-tune the interpretation of cytokine cues, instructing alternate patterns of gene regulation (Fielding *et al*, 2014; Grohmann *et al*, 2018; Hong *et al*, 2002; Jones *et al*, 2015; Jones *et al.*, 2013; Lin & Leonard, 2019; Martinez-Fabregas *et al.*, 2020; Twohig, 2019; Wiede *et al*, 2019).

Resident tissue cells respond to immune challenge by steering decision-making processes affecting disease outcome (Dinarello, 2007; Gomes & Teichmann, 2020; Jones, 2005; Krausgruber *et al*, 2020). These activities rely on cytokine networks that promote communication between stromal tissue and infiltrating leukocytes (Jones, 2005). In bacterial peritonitis, IL-6 controls anti-microbial immunity and the resolution of inflammation by steering the transition from innate to adaptive immunity (Catar *et al*, 2017; Cho *et al*, 2014; Hurst *et al*, 2001; Jones *et al*, 2010; McLoughlin *et al*, 2004; McLoughlin *et al*, 2005). This process requires IL-6R shedding from infiltrating neutrophils, which promotes IL-6 trans-signaling and STAT3-driven outcomes in resident stromal cells (Fielding *et al*, 2008; Hurst *et al.*,

2001; Jones *et al*, 1999; McLoughlin *et al.*, 2004; McLoughlin *et al*, 2003). Repeated immune activation alters this response, with IL-6 supporting an expansion of pro-fibrotic IFNγ-secreting Th1 cells, which initiate tissue injury through activation of stromal STAT1 signaling (Fielding *et al.*, 2014). To understand how the effector properties of Th1 cells impact the contribution of STAT1 and STAT3 in acute resolving peritonitis, we have applied next-generation sequencing methods to examine the stromal response to inflammation. Our analysis shows that Th1 cells alter the transcriptional output of IL-6 by re-directing STAT transcription factors to a GAS-like motif in *Alu*-like retroelements. These results offer an explanation of how effector lymphocytes shape the stromal response to inflammation by modifying the interpretation of cytokine cues.

Results –

***Effector Th1 cells shape stromal responses to peritonitis*–** To understand how pro-fibrotic Th1 cells shape the stromal cell response to acute peritonitis, we challenged (i.p.) mice with a cell-free supernatant from *Staphylococcus epidermidis* (SES) (Fielding *et al.*, 2014; Hurst *et al.*, 2001; Jenkins *et al*, 2005). Mice received SES alone or co-administered with CD4$^+$ T-cells expanded *ex vivo* into Th1 cells (normalized to 10^6 IFNγ-secreting T-cells) (Fielding *et al.*, 2014). To control for transcriptional changes induced by Th1 cells (SES+Th1), a separate group of mice received SES and an equivalent number of naïve CD4$^+$ T-cells (**Fig-1**). At 3 and 6 hours post-stimulation, the peritoneum was harvested and prepared for RNA-seq. K-means clustering was restricted to transcripts differentially regulated (Log2FC>1.75, P$_{adj}$<0.05) under each condition (**Fig-1A**). Applying statistical tools (silhouette, gap statistics, and elbow), we identified four clusters of gene regulation in mice receiving SES alone or SES+Th1 (**Fig-1B**). We identified 821 differentially regulated transcripts whose expression altered in (at least) one time point. SES regulated a total of 225 genes, with the activities of Th1 cells affecting another 673 genes. These included genes affecting vascularization, epithelial morphogenesis, and hyperplasia (e.g., *Angpt4*, *Egr3*, *Igfbp2*, *Ntf3*, *Tbxa2r*). Others displayed involvement in innate sensing pathways (e.g., *Ifitm1*, *Marco*, *Myd88*) and leukocyte infiltration (e.g., *Ccl3*, *Ccl5*, *Ccl2*, *Cxcl1*, *Cxcl10*, *Icam1*, *Vcam1*) (**Fig-1C**). The presence of Th1 cells altered gene expression in each cluster, with molecular pathway analysis highlighting an enrichment of transcripts attributed to STAT1 and interferon signaling (**Fig-2A** & **2B**; Supplemental Figure-1). To substantiate these findings, we used the Upstream Regulator tool in Ingenuity Pathway Analysis to predict transcriptional mechanisms accounting for changes in gene expression in each dataset. Consistent with the SES activation of TLR2 (Colmont *et al*, 2011), this alogothmn identified genes controlled by the NF-κB pathway (**Fig-2C**). It also identified genes linked with STAT1, STAT3, and several IRF transcription factors (**Fig-2C**). Transcripts affiliated with STAT1 and STAT3 showed considerable enrichment in SES+Th1 treated mice, suggesting a potential link between IL-6 and IFNγ signaling in steering SES-induced outcomes (Fielding *et al.*, 2014; Hurst *et al.*, 2001; McLoughlin *et al.*, 2003; Robson *et al*, 2001).

***Th1 cells modify the properties of IL-6*–** To explore the link between IL-6 and IFNγ in SES-induced peritonitis, we used immunodetection methods to quantify changes in their production as a reponse

to SES challenge (**Supplemental Figure-2A**). Peritoneal IFNγ remained below the limit of detection in mice treated with SES alone. However, IFNγ levels were substantially enhanced by the introduction of Th1 cells (**Supplemental Figure-2A**). The presence of Th1 cells also increased IL-6 bioavailability by promoting increases in IL-6 and sIL-6R (**Supplemental Figure-2A**). To establish if Th1 cells alter the transcriptional output of IL-6, we performed RNA-seq on peritoneal tissues from SES challenged wild type (*wt*) and *Il6*[-/-] mice. K-means clustering identified 241 significantly altered transcripts (Log2FC>1.75, P_{adj}<0.05) impacted by the loss of IL-6 (**Fig-3A** & **3B**, **Supplemental Figure-2B-D**). These included gene required for tissue homeostasis (e.g., *Atoh1*, *Cldn5*, *Fgfbp1*, *Npnt*, *Oxtr*, *Pdx1*, *Pgr*, *Stab2*), host defense (e.g., *Trim52*, *C7*), leukocyte recruitment (e.g., *Ccl8*, *Ccl17*, *Ccl22*, *Ccl24*), and adhesion (e.g., *Selp*, *Adgrb2*) (**Fig-3C**). To identify the stromal cells responding to IL-6 in SES-induced inflammation, we compared our results against a single cell RNA-seq dataset (GEO GSM4053741) from mouse omental CD45[-]CD41[-]Ter119[-]CD31[-]PDPN[+/-] cells (Jackson-Jones *et al*, 2020). This analysis defined roles for peritoneal fibroblasts (including *Ccl11*[+] *Pdgfra*[+] and *Matn2*[+] *Pdgfra*[+] subsets) and mesothelial cells (including *Ifit*[+] and *Cxcl13*[+] mesothelial cells) in shaping IL-6 responses (**Supplemental Figure-2E**).

Our initial bioinformatic predictions identified for IL-6 in governing innate immune outcomes. To verify this connection, we developed a flow cytometric method to compare the effector properties of circulating and infiltrating Ly-6B[hi] Ly-6G[hi] neutrophils from *wt* and *Il6*[-/-] mice. Circulating neutrophils were loaded *ex vivo* with the peroxidase substrate 3'(p-aminophenyl) fluorescein (APF) and exposed to opsonized *Staphylococcus epidermidis* labeled with DDAO-far-red. These reporter dyes were used to track neutrophil respiratory burst and phagocytosis capabilities. Circulating Ly-6B[hi] Ly-6G[hi] neutrophils from *wt* and *Il6*[-/-] mice showed no differences in effector functions (**Supplemental Figure-2F**). However, infiltrating neutrophils from *Il6*[-/-] mice treated (i.p.) with 5x10[8] cfu fluorescently labeled *Staphylococcus epidermidis* displayed impaired neutrophil function (**Fig-3D**, **Supplemental Figure-2G**). We further confirmed these findings by visualizing fluorescent bacteria in neutrophils using imaging flow cytometry (**Supplemental Figure-2H**). Since IL-6 requires sIL-6R to regulate stromal responses within the peritoneal cavity, we conducted an identical experiment in *wt* mice treated with the IL-6 trans-signaling antagonist soluble gp130 (sgp130). Treatment with sgp130 significantly reduced the effector properties of infiltrating neutrophils in infected mice (**Fig-3D**). This defect was reversed by reconstituting IL-6 signaling (via the administration [i.p.] of a chimeric IL-6-

sIL-6R fusion protein) in *Il6*[-/-] mice (**Fig-3D**, Supplemental Figure-2H). Thus, IL-6 governs neutrophil responses to local infection.

Th1 cells alter STAT transcription factor activity– STAT1 and STAT3 transcription factors become rapidly activated following SES challenge, with maximal activation coinciding with the 3 and 6 hours chosen for our RNA-seq analysis (Fielding *et al.*, 2014; Fielding *et al.*, 2008; McLoughlin *et al.*, 2005) (Supplemental Figure-3A). STAT1 activities often shape the transcriptional output of IL-6 and STAT3 (Costa-Pereira *et al.*, 2002; Twohig, 2019; Vahedi *et al*, 2012; Villarino *et al.*, 2017). To test if this relationship is seen in response to SES, we extracted peritoneal tissues from *gp130*[Y757F:Y757F] mice challenged with SES. These animals possess a single tyrosine-to-phenylalanine substitution in the cytoplasmic domain of gp130 that prevents the negative regulation of STAT1 and STAT3 following cytokine activation (Jenkins *et al.*, 2005; McLoughlin *et al.*, 2005). Immunoblot for tyrosine-phosphorylated STAT1 (pY-STAT1) and STAT3 (pY-STAT3) showed that SES triggers a prolonged STAT transcription factor activation in these mice (Supplemental Figure-3A). Here, partial *Stat3* ablation in *gp130*[Y757F:Y757F] mice (*gp130*[Y757F:Y757F]:*Stat3*[+/-]) extended the duration of pY-STAT1 activity in response to SES challenge (Supplemental Figure-3A). Thus STAT1 and STAT3 activities are interlinked in SES inflammation and may explain how Th1 cells impact the transcriptional output of IL-6. We, therefore, applied chromatin immunoprecipitation-sequencing (ChIP-seq) to investigate how STAT1 and STAT3 transcription factors engage the genome following SES challenge (**Fig-4**, Supplemental Figure-3B). Our analysis identified sequencing peaks displaying a four-fold enrichment above input (P<0.0001; false discovery rate 0.05). These mapped to transcriptional start sites (TSS), exons, introns, and intergenic regions (Supplemental Figure-3C). Motif enrichment analysis (MEME-ChIP) confirmed the specificity of these interactions and identified consensus DNA binding motifs for STAT1 or STAT3 beneath the mapped sequencing peaks (Supplemental Figure-3B).

The presence of Th1 cells noticeably altered the genomic localization of STAT transcription factors following SES stimulation (**Fig-4A**). Following treatment with SES alone, STAT1 and STAT3 worked in close partnership and often bound loci in nearby proximity. Th1 cells increased the number of sequencing peaks identified by ChIP-seq for STAT1 and STAT3 (Supplemental Figure-3C). This increase in STAT binding was consistent with the enhanced gene regulation associated with Th1 cell involvement (**Fig-1** & **Fig-3**). Comparing STAT transcription factor binding in SES+Th1 datasets from

wt and *Il6*[-/-] mice, we also identified genomic loci displaying evidence of STAT1 and STAT3 cross-regulation (**Fig-5**). Here, STAT transcription factors often bound similar genomic coordinates at downstream sequences distal of TSS. In DNA samples from *wt* mice, these sites were STAT3 occupied. However, in *Il6* deficiency, these same sites showed increases in STAT1 binding. Promoters displaying this form of cross-regulation included genes involved in cytoskeletal organization (e.g., *Mtus2*, *Actb*, *Rhpn2, Wasf1*), metabolism (e.g., *Angptl4*, *Mtor*, *Neu2*, *Pgm1*), and tissue remodeling (e.g., *Timp1*, *Col2a1*, *Vegf*) (**Fig-5A**). This switch from STAT3 to STAT1 coincided with transcriptional changes marked by the induction or suppression of gene expression under *Il6* deficiency (**Fig-5B**). These included the suppression of genes affecting cellular differentiation (e.g., *Dnaic1*, *Eif2s3y*) and increases in genes linked with matrix protein biosynthesis (e.g., *Acan*, *Npnt*, *Col2a1*). Thus, STAT1 and STAT3 coordinate differential impacts on the control of tissue homeostasis.

To substantiate the significance of genomic interactions, we used Assay for Transposase-Accessible Chromatin-sequencing (ATAC-seq) to confirm links between STAT transcription factor binding and chromatin accessibility in SES+Th1 treated mice (**Fig-6**). The heatmap profiles show chromatin accessibility across the entire genome for *wt* and *Il6*[-/-] mice (**Fig-6A**). Adopting a differential binding analysis (*Diffbind*) of ATAC-seq datasets, we identified open chromatin regions linked with IL-6 bioactivity. These sites showed enrichment at TSS (**Fig-6B**). Access to these sites was partially restricted by the absence of *Il6*, suggesting that IL-6 promotes chromatin remodeling in inflammation (**Fig-6C**). Motif enrichment analysis of the DNA sequences aligned to these peaks identified consensus sites consistent with the computational predictions shown in Figure 2C. These included STAT and IRF transcription factors and others (e.g., KLF15, NRF1, HOXOA13, and several zinc-binding factors) contributing to tissue homeostasis and epigenetic control (**Fig-6D**).

***STAT transcription factors engage Alu-like retroelements**–* The proximity of STAT1 and STAT3 binding to consensus sequences for other transcription factors suggested links to regulatory regions such as super-enhancers. We, therefore, mapped the genomic localization of P300 in peritoneal tissue extracts from mice challenged with SES+Th1 cells (**Fig-7A**). This histone acetyltransferase controls chromatin remodeling and often localizes active or poised enhancers, where P300 functions as a scaffolding factor and co-regulator of transcription factor activity (Hutchins *et al*, 2013; Villarino *et al.*, 2017). ChIP-seq analysis for P300 identified sequencing peaks sharing STAT transcription factor

binding (Supplemental Figure-3B). However, P300 mapping represented a small proportion (<11%) of the total sequencing peaks identified by STAT1 and STAT3 ChIP-seq (**Fig-7A**). Although 60% of the P300 loci showed a switch in STAT transcription factor binding under *Il6* deficiency, the transition from STAT3 to STAT1 was more prominent at loci lacking P300 (**Fig-7A**). Promoters displaying this motif included genes for *Stat6*, *Adamts1*, *Socs1*, and several interferon regulated genes (*Irf1*, *Irf9*, *Mx2*, *Il15*, *Ifit1*). These loci displayed STAT3 binding in samples from *wt* mice and STAT1 binding in datasets from *Il6⁻/⁻* mice (**Fig-7A**). A limited number of genomic loci showed binding for both STAT1 and STAT3 (**Fig-7A**).

Combining *wt* and *Il6⁻/⁻* mice datasets, motif analysis identified a centrally enriched sequence (CCTGTAATCCCAGC) with 90-95% identity to annotated GAS elements (MA0137.3, MA0144.2; Supplemental Figure-3B) in our STAT1 and STAT3 ChIP-seq data (**Fig-7B**). Given the conserved nature of this sequence, we used coordinate mapping to define genomic regions with proximity to the CCTGTAATCCCAGC motif in the murine genome. Here, sequence analysis showed the CCTGTAATCCCAGC motif to reside in short interspersed nuclear elements (SINE) classified by the RepeatMasker bioinformatic tool (**Fig-8**) (Ferrari *et al*, 2020). Analysis located STAT1 and STAT3 binding to a subset of SINE resembling B1 Alu elements (**Fig-8A**, Supplemental Figure-4). These elements display conserved architectures that include the CCTGTAATCCCAGC sequence (termed the GAS-Alu motif), residing close to a Pol-II A-box and flanked by consensus sites for T-bet and Runx3 (**Fig-8B**). Genomic DNA from SES-challenged mice showed no significant interaction of STAT transcription factors with the GAS-Alu motif. Thus, Th1 cells modify Jak-STAT cytokine signaling by re-directing STAT factors to genomic *Alu*-like retroelements.

The GAS-like motif identifies immune pathways linked to human physiology– Based on the sequence homology between murine and human Alu elements, we tested whether the GAS-Alu motif correlated with single nucleotide polymorphisms (SNP) in human disease (**Fig-9A**). From publicly available genome-wide association studies (GWAS), we identified GAS-Alu motifs in enhancer sequences designated by the FANTOM5 consortium. Our analysis revealed 8423 sequences, mapping to 2334 genes (Entrez) (Supplemental Figure-5A). Genes affiliated with the GAS-Alu motif were involved in various processes, including thrombopoietin (e.g., *PRKCB*) and VEGF (e.g., *PDGFC, ACTG2*) and integrin (e.g., *RAPGEF1*) signaling. Others regulate leukocyte signaling (e.g., *VAV1, CACNG3,*

PPP3CB) and migration (*ARHGAP8*, *ACTG2*), and tissue turnover (e.g., *MYO10*, *ARHGEF19*). These included several mapped by our ChIP-seq of murine STAT1 and STAT3. For example, *ARHGEF19* (*Arhgef19*), *COL5A* (*Col5a*), *MYO10* (*Myo10*), *PPARG* (*Pparg*), *PRKCB* (*Prkcb*), and *RABGEF1* (*Rabgef1*).

Next, we downloaded GWAS summary statistics (n=2505) hosted by major repositories (NHGRI-EBI, CTGLab, NCBI) and tested for FANTOM5 genesets and gene signatures of IL-6 and IFNγ activity (available from msigdb; Broad Institute) using MAGMA. These were compared to a series of randomized and shuffled gene lists. Genes aligned to the HLA locus were excluded from our datasets to control against the high degree of linkage disequilibrium at these loci. MAGMA output files were compared against enhancers mapped by the FANTOM5 project and identified 257 GWAS displaying enrichment of the GAS-Alu motif (**Supplemental Figure-5B**). Pairwise comparison (Pearson) and hierarchical clustering of these GWAS datasets identified three clusters based on gene identity (**Fig-9A**). Genes in Cluster-2 contained an enrichment of IL-6 and IFNγ gene signatures and included genes involved in immune regulation and pathophysiology (**Fig-9B** & **9C**). These include traits assigned as polygenic and monogenic disease variants (Vuckovic *et al*, 2020). Thus, the GAS-Alu motif classifies examples of Jak-STAT cytokine signaling in immune pathology.

Discussion –

Cytokines regulate transcriptional processes that maintain tissue homeostasis, protective immunity, and outcomes affecting inflammation-induced tissue injury (Borthwick *et al*, 2013; Duffield *et al*, 2013). In bacterial peritonitis, IL-6 and IFNγ are critical determinants of these outcomes (Catar *et al.*, 2017; Fielding *et al.*, 2014; Hurst *et al.*, 2001; Jenkins *et al.*, 2005; Jones *et al.*, 2010; Krausgruber *et al.*, 2020; McLoughlin *et al.*, 2005; McLoughlin *et al.*, 2003; Modur *et al*, 1997; Navarini *et al.*, 2011; Robson *et al.*, 2001; Schett *et al.*, 2013; Xing *et al*, 1998; Yang *et al*, 2020). Here, IL-6 compromises tissue repair by supporting the expansion of pro-fibrotic IFNγ-secreting CD4$^+$ T-cells (Fielding *et al.*, 2014). While these cytokines rely on STAT1 and STAT3 signaling (Fielding *et al.*, 2014; Fielding *et al.*, 2008; Jenkins *et al.*, 2005; McLoughlin *et al.*, 2005), it is unclear how these transcription factors coordinate anti-microbial host immunity, tissue scarring, and fibrosis. To understand their involvement in these processes, we tracked the stromal activity of STAT1 and STAT3 transcription factors in peritoneal tissues following an inflammatory challenge. Our analysis shows that IFNγ-secreting CD4$^+$ T-cells alter the transcriptional output of IL-6 by channeling STAT transcription factors to previously unoccupied genomic loci.

Jak-STAT signaling is complex and includes an interplay between individual STAT transcription factors (Costa-Pereira *et al.*, 2002; Ernst *et al*, 2008; Hirahara *et al.*, 2015; Jones *et al.*, 2013; Olbrich & Freeman, 2018; Qing & Stark, 2004; Villarino *et al.*, 2017; Yang *et al.*, 2020; Yang *et al*, 2011). For example, patients with *STAT1* gain-of-function or *STAT3* loss-of-function mutations often display similar clinical features. These include increased susceptibility to infections at barrier surfaces, eczema-type rashes, and bowel perforations (Abusleme *et al*, 2018; Goel *et al*, 2018; Liu *et al*, 2011; Luis & Calva-Mercado, 2018; Milner *et al*, 2015; Olbrich & Freeman, 2018; Toubiana *et al*, 2016). These clinical phenotypes are also present in patients lacking *IL6R* or mice lacking *Il6* and often reflect the role of STAT3 in innate immunity (Holland *et al*, 2007; Kano *et al*, 2003; Kopf *et al*, 1994; Matsukawa *et al*, 2003; Ramsay *et al*, 1994; Spencer *et al*, 2019; Welte *et al*, 2003). Consistent with these findings, our analysis shows that IL-6 controls stromal activities that enhance the phagocytic properties of infiltrating neutrophils. The inclusion of Th1 cells enhances this anti-microbial response, contributing to an increase in various interferon-responsive genes supporting host immunity. During acute resolving inflammation, a limited number of IFNγ-secreting CD4$^+$ T-cells are present in the

peritoneal cavity (Fielding *et al.*, 2014; Jones *et al.*, 2010). This number significantly increases as a response to repeated peritonitis, and their retention leads to compromised tissue homeostasis and fibrosis following activation (Fielding *et al.*, 2014; Jones *et al.*, 2010). Our analysis reveals that this change in inflammatory status impacts various cellular processes that include an alteration in IL-6 bioactivity. These data point towards a transcriptional interplay between IL-6 and IFNγ that impacts how STAT transcription factors shape inflammatory outcomes.

Studies of cancer cells and *ex vivo* stimulated T-cells detail how cross-regulation between STAT1 and STAT3 modulates target gene expression (Costa-Pereira *et al.*, 2002; Hirahara *et al.*, 2015; Hong *et al.*, 2002; Jones *et al.*, 2013; Twohig, 2019). Our studies suggest that IFNγ secretion by Th1 cells may facilitate such interactions in stromal tissues following inflammatory activation. In this context, Th1 cells directly impacted the transcriptional output of IL-6 and were responsible for directing STAT1 and STAT3 to previously latent enhancers. Genes affiliated with these loci contribute to tissue remodelling, fibrosis, solute transport, membrane permeability, and hypoxia. Thus, our data points towards an agonist-specific repertoire of latent enhancers employed to sense and interpret changes in the tissue microenvironment (Ostuni *et al*, 2013). For example, transcription factors linked with myeloid cell development (e.g., PU.1) often instruct the binding of NF-κB, AP-1, and interferon response factors to the genome (Natoli *et al*, 2011). Our results are, therefore, consistent with theories that genes linked by roles in related biological functions commonly share similar mechanisms of transcriptional control (Tong *et al*, 2016). In this regard, a close inspection of the DNA sequences enriched for STAT transcription factor binding identified a conserved motif in *Alu*-like retroelements (Chen & Yang, 2017; Vassetzky & Kramerov, 2013). Retroelements are endogenous components of eukaryotic genomes. They support non-allelic recombination, polyadenylation, alternative splicing, and the transcription of gene-rich regions (Chen & Yang, 2017; Tajnik *et al*, 2015). Significantly, retroelements possess consensus binding motifs for various transcription factors and often display evidence of DNA methylation, suggesting an involvement in gene regulation (Polak & Domany, 2006; Xie *et al*, 2009). Functional genomic studies in cell lines of stromal or immune cell origin demonstrating the binding of basic leucine zipper transcription factors, the aryl hydrocarbon receptor and other transcriptional regulators to retroelements (Lu *et al*, 2020; Roman *et al*, 2008; Sun *et al*, 2018). Our analysis showed that IL-6 signaling, in association with IFNγ-secreting Th1 cells,

promotes STAT3 binding to *Alu* sequences. STAT3 binding was, however, lost in *Il6* deficiency. Instead, these same sites showed STAT1 binding. Thus, *Alu*-like retroelements may represent sentinels of transcriptional cross-regulation in stromal tissues.

What is the significance of STAT transcription factor binding to *Alu* sequences? Are these interactions relevant to human disease or treatment responses to tocilizumab, tofacitinib, and others? Addressing these questions is challenging due to the complexities of mapping repetitive DNA elements. Our data imply a link between *Alu*-like sequences and tissue pathology. Moreover, GWAS commonly identify human *Alu* polymorphisms linked with interferonopathies or diseases characterized by alterations in STAT1 activity (Hung *et al*, 2015; Payer *et al*, 2017; Volkman & Stetson, 2014; Zeng *et al*, 2014). Our analysis of human GWAS datasets revealed several targets also identified in mice treated with SES and Th1 cells. These data support a role for epigenetic modifiers that regulate the accessibility of transcription factors to specific enhancers under certain inflammatory settings or disease processes. Future evaluation of these events will open new opportunities to understand how cytokine cues are interpreted or fine-tuned to direct physiology or pathophysiology.

Acknowledgements –

This manuscript is dedicated to Dr Javier Uceda Fernandez, a dearly loved friend and colleague who was tragically taken from us on the 29th of August 2018. Kidney Research UK (Reference RP-024-20160304 awarded to SAJ, DF, NT, GWJ), Versus Arthritis (Reference 20770, 19796, 20305 awarded to SAJ, VOD, NMW & GWJ), and the National Health and Medical Research Council of Australia (to BJJ) provided grant support for this project. JUF was recipient of a la Caixa PhD Studentship administered through the British Council, and BCC is supported by a PhD studentship from the Systems Immunity University Research Institute at Cardiff. JL is recipient of a Rutherford Fellowship Grant. PRT is recipient of a Wellcome Trust Investigator Award (Reference 107964/Z/15/Z) and receives funding through the UK Dementia Research Institute. Bioinformatic analysis was developed with support from the Systems Immunity University Research Institute in Cardiff.

Materials and methods–

Animals– All procedures were performed under UK Home Office project license P05D6A456. Inbred wild type C57BL/6 male mice were purchased from Charles River UK. IL-6-deficient (*Il6*$^{-/-}$) (Kopf *et al.*, 1994) were bred under approved UK Home Office guidelines in Cardiff University. The *gp130*$^{Y757F:Y757F}$ and *gp130*$^{Y757F:Y757F}$:*Stat3*$^{+/-}$ mice have been previously described (Jenkins *et al.*, 2005). Experiments were approved by the Animal Ethics Committee and included genetically matched *gp130*$^{+/+}$ littermate controls. All experiments were performed using age-matched 8-12 weeks old mice.

SES-Induced Peritoneal Inflammation– A lyophilized cell-free supernatant prepared from *Staphylococcus epidermidis* (SES), whose activity had been standardized using an *in vitro* cell-based CXCL8 bioassay, was used to induce acute peritoneal inflammation (Hurst *et al.*, 2001). Mice were administered (i.p.) with a defined dose of SES (500µl). Soluble gp130Fc was added (i.p.) to WT mice as indicated. At 3 and 6 hours post-inflammatory challenge, mice were sacrificed and the peritoneal cavity lavaged with 2ml ice cold PBS. The peritoneal membrane was harvested at the same time points.

Transfer of Th1 Cells– To replicate events promoting peritoneal fibrosis some mice receiving SES were also simultaneously administered with either naïve CD4$^+$ T-cells or CD4$^+$ T-cells conditioned *ex vivo* towards a Th1 phenotype. Briefly, splenic naïve CD4$^+$ T-cells (CD4$^+$CD25$^-$CD44^{lo}CD62L^{hi}) were flow sorted and place in coated plates with anti-CD3e (145-2C11) and 5µg/ml soluble anti-CD28 (37.51) antibodies. Cells were cultured for 4 days in the presence of 20ng/ml murine recombinant IL-12 (R&D systems; 419-ML). The proportion of IFNγ-secreting CD4$^+$ T-cells (Th1 cells) was determined by intracellular flow cytometry using antibodies against to CD4 (RM4-5), IFNγ (XMG1.2), IL-17 (TC11-18H10.1), IL-4 (11B11) and IL-13 (eBio13A). Based on this analysis, T-cells were washed in ice cold PBS, re-suspended in a 500µl aliquot of PBS reconstituted SES, and administered via the intraperitoneal route. Here, a cell concentration of 5-10 x10^5 CD4$^+$ T-cells was administered to reflect the proportion of Th1 cells recruited to the peritoneal cavity under acute SES challenge. Where indicated, control mice were administered the same number of sorted naïve (Th0) CD4$^+$ T-cells. Changes in the inflammatory infiltrate were analyzed by direct counting (Coulter Z2, Beckman

Coulter), differential cell counting and flow cytometry approaches using antibodies against defined leukocyte subsets. The peritoneal membrane was harvested at 3 and 6h after the injections.

***Fluorescent labelling of bacteria*–** An inoculum of *Staphylococcus epidermidis* ATCC 12228 (5×10^8 cfu/mouse) was prepared from log phase cultures under sterile conditions. The suspension was centrifuged, and bacteria labelled for 20 min at 37°C in pre-warmed PBS containing Cell Trace Far Red (CT-FR) (Life Technologies) (1 µM or 8 µM for *ex vivo* and *in vivo* experiments, respectively). For *ex vivo* experiments, bacteria were serum-opsonized while, for *in vivo* experiments were centrifuged and washed 3 times in PBS, before resuspending in sterile PBS.

***Ex vivo neutrophil effector function assay*–** Whole blood was collected by cardiac puncture into tubes containing 5mM EDTA. Samples were diluted 1:10 and washed 3 times in ice cold PBS. Cells were resuspended in serum-free RPMI 1640 containing 5 µM of 3'-(p-aminophenyl) fluorescein (APF) and incubated for 30 min at 37°C. APF-loaded cells were split into 100 µl aliquots and cultured at 37°C with an equal volume of pre-warmed, opsonized Cell Trace Far Red (CT-FR; 1 µM) labelled *S. epidermidis*. Cells were incubated for set time intervals (0-30 min) and transferred to an iced water bath prior to preparation for flow cytometry using a Beckman Coulter Cyan-ADP flow cytometer.

***In vivo neutrophil effector function assay*–** Mice were administered (i.p.) with Cell Trace Far Red (CT-FR; 8 µM) labelled *S. epidermidis*. An independent group of *Il6*[-/-] mice received a dose of CT-FR labelled *S. epidermidis* together with the IL-6-sIL-6R chimeric fusion protein HDS (50 ng/ml). Six hours after bacterial challenge, the peritoneal cavity was lavaged with 2 ml of RPMI 1640 containing 5 µM of 3'-(p-aminophenyl) fluorescein (APF). Neutrophil phagocytosis and respiratory burst activity were examined by flow cytometry using a Beckman Coulter Cyan-ADP flow cytometer and analyzed using *Summit* (software v4.3, Beckman-Coulter) or FlowJo 10 (TreeStar).

***Imaging flow cytometry analysis*–** Lavaged neutrophils were immune-stained for Ly6G (1A8), cells were resuspended in 100 µl of sterile PBS, and events (>8000 events per sample) were acquired at a low imaging rate and 60X amplification using the Amnis ImageStream®X Mark II Imaging Flow Cytometer (Amnis). Neutrophils were gated according to Ly6G staining. Phagocytic activity was expressed as a Phagocytic Index, which reflects the number of bacteria ingested by an individual Ly6G[+] neutrophil during the incubation period. To evaluate the phagocytic index of neutrophils undergoing phagocytosis, the Application "Spot Counting" in ImageStream software IDEAS (Amnis)

was used. Spots were counted based on the distribution of CT-FR bacterial staining. The efficiency of phagocytic uptake was quantified by examining cells displaying either a low (1-2 counts) or a high number (3 counts or more) of ingested bacteria.

Immunoblotting of peritoneal tissues – Protein was extracted from frozen peritoneal biopsies using ice-cold lysis buffer. Samples were pre-cleared of cellular debris before separation by SDS-PAGE and immunoblotting with specific antibodies against STAT1, STAT3 and tyrosine phosphorylated forms of STAT1 (pY-STAT1) and STAT3 (pY-STAT3) (Jenkins *et al.,* 2005). Immuno-labelled proteins were detected using either the enhanced chemiluminescence (ECL) detection system (Amersham Biosciences) or Odyssey Infrared Imaging System (LI-COR, Lincoln, New England) with the appropriate secondary antibodies as per the manufacturer's instructions.

RNA-seq – Peritoneal membrane sections (80mg tissue) were dissociated in 1ml buffer RLT (QIAGEN) supplemented with β-mercaptoethanol (1:100 v:v) using a handheld electric homogeniser (Benchmark Scientific). Lysate was diluted 1:3 in distilled water and digested in 0.2mg/ml proteinase-K (Invitrogen; 25530049) for 10 minutes at 55°C. Lysate was cleared and RNA precipitated in 70% ethanol. Total RNA was extracted using the RNeasy Mini kit (QIAGEN) following the manufacturer's instructions. 2-4mg of mRNA was processed to generate the libraries. Cytoplasmic, mitochondrial, and ribosomal RNA was depleted using the Ribominus transcriptome isolation kit (Ambion; K155001). Libraries were prepared using the RNA-seq kit v2 (Life technologies; 4475936) and sequencing on an ion torrent (Thermo Fisher).

Chromatin Immunoprecipitation (ChIP)-seq – Excised peritoneal membranes were immediately frozen in liquid nitrogen and stored at -80°C until use. Tissues were diced and ground to a fine powder with intermittent addition of liquid nitrogen. Genomic DNA was extracted, crosslinked, and fragmented by sonication prior to treatment with 2µg/ml of anti-STAT1 (sc-592, Santa Cruz Biotechnology), anti-STAT3 (sc-482, Santa Cruz Biotechnology), anti-P300 (05-257, Millipore), or isotype control antibodies. Immunoprecipitation was conducted overnight at 4°C under continuous gentle agitation. Antigen-antibody complexes were captured using Protein-A/G magnetic beads, washed and DNA fragments liberated following treatment with proteinase-K and extraction with phenol-chloroform. Biological repeats from 3 independent tissue extracts were pooled and concentrated before the library preparation and next generation sequencing. ChIP libraries were

prepared according to the manufacturer's instructions (Illumina Truseq DNA ChIP kit; RS-122-2001). Size selection (200-400bp) was determined using a Blue Pippin (Sage Science) system employing 2% agarose cartridges (Sage Science; BDF2003). Appropriate library size distribution was confirmed using an Agilent 2100 bioanalyser and final concentrations determined using Qubit (Invitrogen). Libraries were sequenced on an Illumina HiSeq4000.

***ATAC-seq*–** Excised peritoneal membranes were immediately frozen in liquid nitrogen and stored at -80°C until use. Tissues were diced and ground to a fine powder with intermittent addition of liquid nitrogen. Omni-ATAC seq was performed as described previously (Corces *et al*, 2017). Briefly, 100,000 nuclei per sample were isolated using an iodixanol gradient and ATAC-seq was subsequently performed according to the original protocol (Buenrostro *et al*, 2013) using Nextera DNA Sample preparation kit (Illumina, FC-121-1030). After library amplification, DNA was isolated using Qiagen MiniElute kit and size selection was performed using Blue Pippin (Sage Science) system employing 2% agarose cartridges (Sage Science; BDF2003). Libraries were sequenced on an Illumina HiSeq4000.

***Quantification and Statistical Analysis*–** No randomization and no blinding were used for the animal experiments. Whenever possible, the investigator was partially blinded for assessing the outcome. All data were analyzed using Prism 8 (GraphPad Prism, La Jolla, CA). Information on the types of Statistical methods used, the sample size and number of repetitions are listed in the Legends.

***RNA-seq data analysis*–** Raw fastq were mapped using Torrent SuiteTM to the *mm10* reference genome and counts were assigned to transcripts using featureCounts (Liao *et al*, 2014) with the GRCm38.84 Ensembl gene build GTF. Differential gene expression analyses used the DESeq2 package (Love *et al*, 2014). Genes were discarded from the analysis differential expression failed to be significant (significance: adj.pval < 0.05, Benjamini-Hochberg correction for multiple testing). Differentially regulated genes were uploaded into Ingenuity Pathway Analysis (IPA) (QIAGEN) for functional analysis.

***ChIP-seq data analysis*–** Between 40-70M reads were obtained for each sample. These were trimmed with Trim Galore (https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) and assessed for quality using FastQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Reads were mapped to the mouse mm10 reference genome using bwa (Li & Durbin, 2009). Peaks were called using macs2 (Zhang *et al*, 2008), using the BAMPE model (adj.pval 0.05).

***ATAC-seq analysis*– **P**aired-end reads were processed with Trim Galore (https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) and assessed for quality using FastQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) prior to mapping to the mouse *mm10* reference genome (Li & Durbin, 2009). Peaks were called using macs2 and the BAMPE model (adj.pval 0.05) (Zhang *et al.*, 2008). Differential open-region analysis used Diffbind in Bioconductor

(http://bioconductor.org/packages/release/bioc/vignettes/DiffBind/inst/doc/DiffBind.pdf).

Histograms depicting distance from TSS were generated using the ChIPseeker R-package and show the peak count fraction.

***FANTOM5 enhancer analysis*– Sequences annotated as enhancers by the FANTOM5 consortium were downloaded from Slidebase (250bp pad) (http://slidebase.binf.ku.dk). Motif occurrences (n=8423, p<0.00001) in these sequences were identified using the FIMO algorithm (Meme ChIP-suite; http://meme-suite.org/tools/fimo). Genes in proximity to the identified sequences (defined as 2kb upstream or downstream) were mapped to the ensembl GRCh38 (hg38) build and visualized using the ClueGO cytoscape pluggin). Genome-wide motif occurrences were visualized with RIdeogram R-package (https://cran.r-project.org/web/packages/RIdeogram/vignettes/RIdeogram.html).

***Motif identification*– Sequences under ChIP peaks (q<0.05) were obtained using the bedtools getfasta command against the ensemble *mm10* genome build. Sequence fasta files were uploaded to Meme ChIP (http://meme-suite.org/tools/meme-chip) and searched against the murine HOCOMOCO (v11 CORE) and Eukaryote DNA databases. The *de novo* GAS-like motif was enriched by both Centrimo and Meme algorithms.

***Spaced motif analysis*– Analysis was conducted with Spamo (Meme-ChIP-suite) using default parameters. Secondary motifs occur within 150bp of the user-provided primary motif (GAS-like). All secondary motifs are referenced in the HOCOMOCO (v11 CORE) database. Input sequences, including a 250bp pad, were derived from the ChIP-seq data or enhancer annotations in FANTOM5.

***Multiple Sequence Alignment*– Consensus sequences (Hidden Markov Models) corresponding to murine and human Alu-family transposable elements were downloaded from the Dfam database (https://dfam.org/) - Mouse; B1_mm, B1_mus1, PB1D11, B1_Mur2, B1_Mur4, PB1, B1_Mur1, B1_Mur3, PB1D10, B1_Mus2, B1_F1, PB1D7, PB1D9, B1F2, B1F, B2_Mm1a, B2_Mm1t, B2_Mm2, B3,

B3A, B4, B4A, ID_B1, Human; AluY, AluSc, AluJB, AluJo, AluJr, AluJr4, AluSc5, AluSc8, AluSg, AluSg4, AluSg7, AluSp, AluSq, AluSq10, AluSq2, AluSq4, AluSx, AluSx1, AluSx3, AluSx4, AluSz, AluSz6, AluYa5, AluYa8, AluYb8, AluYb9, AluYc, AluYc3, AluYd8, AluYh9, AluYk11, AluYk12, AluYk4, AluYg6, AluYk3, AluYm1, AluYk2, AluYe6, AluYi6, AluYe5, AluYi6_4d, AluYf1, AluYh3, AluYj4, AluYh7, FAM, FLAMA, FLAMC, FRAM. Sequences were aligned using MUSCLE (European Bioinformatics Institute) using the default ClustalW output and visualized in Jalview.

Gene Set Enrichment Analysis (GSEA)– A ranked gene list was prepared for each dataset using the differential gene expression analysis. Using the GSEA pre-ranked function, enrichment profiles were generated against the biological processes (C-5; 7573 gene sets) gene ontology database. For visualization, GSEA output files were loaded into Cytoscape using the enrichment-map plugin. For small gene sets, ontology enrichments were performed using either the mSigdb overlap tool (broad institute) or the Metascape online tool. Network aesthetics (e.g., color, spacing) were modified in adobe illustrator. Network statistical thresholds are below p<0.01, q<0.01.

Repeatmasker overlaps– Alu and L1 sequence coordinates were derived from the repeatmasker definitions downloaded from the UCSC table browser (*mm10*). Repeats overlapping STAT binding were identified using the bedtools intersect algorithm. The absolute number of Alu and L1 sequences overlapping each ChIP dataset was calculated in R studio and plotted using the circlize package.

Visualization and annotation– Heatmap visualization in Morpheus (broad institute) and Pheatmap (R package) using colours defined in the viridis and R Color Brewer packages. Figures were prepared using the ggplot2 R package (R package), graphpad prism8, and arranged in Adobe Illustrator. Data filtering for complex heatmaps was achieved using Dplyr (R package). IL-6 regulated genes were defined as those meeting the statistical threshold (*Il6^-/-* vs wt padj<0.05 with a log2 fold change greater than 1.75 or less than -1.75) in at least one experimental condition. STAT regulated genes were similarly described as those with ChIP signal under the statistical threshold (q<0.05) in at least one experimental condition.

Generation of gene sets for MAGMA analysis– Sequences annotated as enhancers by the FANTOM5 consortium were downloaded from Slidebase (http://slidebase.binf.ku.dk/) including a 200bp pad. To identify motif occurrences, these sequences were entered to the FIMO web server (Meme ChIP suite) together with the MEME-formatted motif extracted from our ChIP-sequencing analyses.

Identified gene sets were derived by mapping coordinates to the HG19 reference genome (genes within 2kb) using PAVIS (https://manticore.niehs.nih.gov/pavis2/). In parallel, FANTOM5 sequences containing Alu sequences were defined by intersecting (bedtools) FANTOM5 coordinates with the Repeatmasker database (UCSC table browser). Coordinates were mapped to genes using PAVIS. Randomised control gene sets were derived as follows. (1) FIMO output coordinates were 'shuffled' across FANTOM sequences or genome wide (Hg19 reference genome) using the Bedtools shuffle feature. (2) Random bed files (8000x 500bp sequences) were generated using the Bedtools Random against either the FANTOM5 enhancer sequences or genome wide (hg19). Three gene sets were generated for each control. Hallmark gene sets were downloaded from msigdb (broad Institute). Genes mapping to the MHC locus were downloaded from the UCSC table browser. This gene list was used to filter MHC-locus genes from all gene sets prior to MAGMA analysis (Dplyr R-package).

MAGMA– GWAS summary statistics (Hg19) were downloaded using FTP links supplied by the GWAS atlas (https://atlas.ctglab.nl/). After re-formatting for compatibility with MAGMA, summary statistics were mapped to genes using the build-37 gene locations file (NCBI37.3.gene.loc). Gene results files were next generated for each summary statistic. The GWAS atlas was used to refence N (number of study participants) for each study. Finally, gene set analysis was performed against gene sets generated as described above. Results files were read into R and relevant columns extracted using R base functions. For genes, p-values were extracted from MAGMA output files ending genes.sets.out. For gene sets, p-values were extracted from output files ending.gsa.out. P-values were merged into a single dataframe for correlation analysis (R-base function) or filtering (Dplyr R-package). Heatmaps were generated using the Pretty heatmap R package (Pheatmap).

Data and Code Availability– RNA-seq, ChIP-seq and ATAC-seq datasets reported in this paper have been deposited in ArrayExpress under accession code EMBL-EBIE-MTAB-10087. The authors declare that all relevant data supporting the findings of this study are available on request. R scripts for performing the main steps of analysis are available from the Lead contact on reasonable request.

References–

Abusleme L, Diaz PI, Freeman AF, Greenwell-Wild T, Brenchley L, Desai JV, Ng WI, Holland SM, Lionakis MS, Segre JA *et al* (2018) Human defects in STAT3 promote oral mucosal fungal and bacterial dysbiosis. *JCI Insight* 3

Andrews C, McLean MH, Durum SK (2018) Cytokine Tuning of Intestinal Epithelial Function. *Front Immunol* 9: 1270

Avalle L, Pensa S, Regis G, Novelli F, Poli V (2012) STAT1 and STAT3 in tumorigenesis: A matter of balance. *JAKSTAT* 1: 65-72

Borthwick LA, Wynn TA, Fisher AJ (2013) Cytokine mediated tissue fibrosis. *Biochim Biophys Acta* 1832: 1049-1060

Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ (2013) Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nat Methods* 10: 1213-1218

Catar R, Witowski J, Zhu N, Lucht C, Derrac Soria A, Uceda Fernandez J, Chen L, Jones SA, Fielding CA, Rudolf A *et al* (2017) IL-6 Trans-Signaling Links Inflammation with Angiogenesis in the Peritoneal Membrane. *J Am Soc Nephrol* 28: 1188-1199

Chen LL, Yang L (2017) ALUternative Regulation for Gene Expression. *Trends Cell Biol* 27: 480-490

Cho Y, Johnson DW, Vesey DA, Hawley CM, Pascoe EM, Clarke M, Topley N, bal ANZTI (2014) Dialysate interleukin-6 predicts increasing peritoneal solute transport rate in incident peritoneal dialysis patients. *BMC Nephrol* 15: 8

Choy EH, Kavanaugh AF, Jones SA (2013) The problem of choice: current biologic agents and future prospects in RA. *Nat Rev Rheumatol* 9: 154-163

Colmont CS, Raby AC, Dioszeghy V, Lebouder E, Foster TL, Jones SA, Labeta MO, Fielding CA, Topley N (2011) Human peritoneal mesothelial cells respond to bacterial ligands through a specific subset of Toll-like receptors. *Nephrol Dial Transplant* 26: 4079-4090

Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA, Vesuna S, Satpathy AT, Rubin AJ, Montine KS, Wu B *et al* (2017) An improved ATAC-seq protocol reduces background and enables interrogation of frozen tissues. *Nat Methods* 14: 959-962

Costa-Pereira AP, Tininini S, Strobl B, Alonzi T, Schlaak JF, Is'harc H, Gesualdo I, Newman SJ, Kerr IM, Poli V (2002) Mutational switch of an IL-6 response to an interferon-gamma-like response. *Proc Natl Acad Sci U S A* 99: 8043-8047

Dinarello C (2007) Historical insights into cytokines. *Eur J Immunol* 37: S34-45

Duffield JS, Lupher M, Thannickal VJ, Wynn TA (2013) Host responses in tissue repair and fibrosis. *Annu Rev Pathol* 8: 241-276

Ernst M, Najdovska M, Grail D, Lundgren-May T, Buchert M, Tye H, Matthews VB, Armes J, Bhathal PS, Hughes NR *et al* (2008) STAT3 and STAT1 mediate IL-11-dependent and inflammation-associated gastric tumorigenesis in gp130 receptor mutant mice. *The Journal of clinical investigation* 118: 1727-1738

Ferrari R, de Llobet Cucalon LI, Di Vona C, Le Dilly F, Vidal E, Lioutas A, Oliete JQ, Jochem L, Cutts E, Dieci G *et al* (2020) TFIIIC Binding to Alu Elements Controls Gene Expression via Chromatin Looping and Histone Acetylation. *Mol Cell* 77: 475-487.e411

Fielding CA, Jones GW, McLoughlin RM, McLeod L, Hammond VJ, Uceda J, Williams AS, Lambie M, Foster TL, Liao CT *et al* (2014) Interleukin-6 signaling drives fibrosis in unresolved inflammation. *Immunity* 40: 40-50

Fielding CA, McLoughlin RM, McLeod L, Colmont CS, Najdovska M, Grail D, Ernst M, Jones SA, Topley N, Jenkins BJ (2008) IL-6 regulates neutrophil trafficking during acute inflammation via STAT3. *J Immunol* 181: 2189-2195

Goel S, Sahu S, Minz RW, Singh S, Suri D, Oh YM, Rawat A, Sehgal S, Saikia B (2018) STAT3-Mediated Transcriptional Regulation of Osteopontin in STAT3 Loss-of-Function Related Hyper IgE Syndrome. *Front Immunol* 9: 1080

Gomes T, Teichmann SA (2020) An antiviral response beyond immune cells. *Nature* 583: 206-207

Grohmann M, Wiede F, Dodd GT, Gurzov EN, Ooi GJ, Butt T, Rasmiena AA, Kaur S, Gulati T, Goh PK *et al* (2018) Obesity Drives STAT-1-Dependent NASH and STAT-3-Dependent HCC. *Cell* 175: 1289-1306 e1220

Harbour SN, DiToro DF, Witte SJ, Zindl CL, Gao M, Schoeb TR, Jones GW, Jones SA, Hatton RD, Weaver CT (2020) TH17 cells require ongoing classic IL-6 receptor signaling to retain transcriptional and functional identity. *Sci Immunol* 5

Hirahara K, Onodera A, Villarino AV, Bonelli M, Sciume G, Laurence A, Sun HW, Brooks SR, Vahedi G, Shih HY *et al* (2015) Asymmetric Action of STAT Transcription Factors Drives Transcriptional Outputs and Cytokine Specificity. *Immunity* 42: 877-889

Holland SM, DeLeo FR, Elloumi HZ, Hsu AP, Uzel G, Brodsky N, Freeman AF, Demidowich A, Davis J, Turner ML *et al* (2007) STAT3 mutations in the hyper-IgE syndrome. *N Engl J Med* 357: 1608-1619

Hong F, Jaruga B, Kim WH, Radaeva S, El-Assal ON, Tian Z, Nguyen VA, Gao B (2002) Opposing roles of STAT1 and STAT3 in T cell-mediated hepatitis: regulation by SOCS. *The Journal of clinical investigation* 110: 1503-1513

Hung T, Pratt GA, Sundararaman B, Townsend MJ, Chaivorapol C, Bhangale T, Graham RR, Ortmann W, Criswell LA, Yeo GW *et al* (2015) The Ro60 autoantigen binds endogenous retroelements and regulates inflammatory gene expression. *Science* 350: 455-459

Hunter CA, Jones SA (2015) IL-6 as a keystone cytokine in health and disease. *Nat Immunol* 16: 448-457

Hurst SM, Wilkinson TS, McLoughlin RM, Jones S, Horiuchi S, Yamamoto N, Rose-John S, Fuller GM, Topley N, Jones SA (2001) Il-6 and its soluble receptor orchestrate a temporal switch in the pattern of leukocyte recruitment seen during acute inflammation. *Immunity* 14: 705-714

Hutchins AP, Diez D, Miranda-Saavedra D (2013) Genomic and computational approaches to dissect the mechanisms of STAT3's universal and cell type-specific functions. *JAKSTAT* 2: e25097

Jackson-Jones LH, Smith P, Portman JR, Magalhaes MS, Mylonas KJ, Vermeren MM, Nixon M, Henderson BEP, Dobie R, Vermeren S *et al* (2020) Stromal Cells Covering Omental Fat-Associated Lymphoid Clusters Trigger Formation of Neutrophil Aggregates to Capture Peritoneal Contaminants. *Immunity* 52: 700-715 e706

Jenkins BJ, Grail D, Nheu T, Najdovska M, Wang B, Waring P, Inglese M, McLoughlin RM, Jones SA, Topley N *et al* (2005) Hyperactivation of Stat3 in gp130 mutant mice promotes gastric hyperproliferation and desensitizes TGF-beta signaling. *Nature medicine* 11: 845-852

Jones GW, Bombardieri M, Greenhill CJ, McLeod L, Nerviani A, Rocher-Ros V, Cardus A, Williams AS, Pitzalis C, Jenkins BJ *et al* (2015) Interleukin-27 inhibits ectopic lymphoid-like structure development in early inflammatory arthritis. *J Exp Med* 212: 1793-1802

Jones GW, Greenhill CJ, Williams JO, Nowell MA, Williams AS, Jenkins BJ, Jones SA (2013) Exacerbated inflammatory arthritis in response to hyperactive gp130 signalling is independent of IL-17A. *Ann Rheum Dis* 72: 1738-1742

Jones GW, McLoughlin RM, Hammond VJ, Parker CR, Williams JD, Malhotra R, Scheller J, Williams AS, Rose-John S, Topley N *et al* (2010) Loss of CD4+ T cell IL-6R expression during inflammation underlines a role for IL-6 trans signaling in the local maintenance of Th17 cells. *J Immunol* 184: 2130-2139

Jones SA (2005) Directing transition from innate to acquired immunity: defining a role for IL-6. *J Immunol* 175: 3463-3468

Jones SA, Jenkins BJ (2018) Recent insights into targeting the IL-6 cytokine family in inflammatory diseases and cancer. *Nat Rev Immunol* 18: 773-789

Jones SA, Novick D, Horiuchi S, Yamamoto N, Szalai AJ, Fuller GM (1999) C-reactive protein: a physiological activator of interleukin 6 receptor shedding. *J Exp Med* 189: 599-604

Jones SA, Scheller J, Rose-John S (2011) Therapeutic strategies for the clinical blockade of IL-6/gp130 signaling. *The Journal of clinical investigation* 121: 3375-3383

Kano A, Wolfgang MJ, Gao Q, Jacoby J, Chai GX, Hansen W, Iwamoto Y, Pober JS, Flavell RA, Fu XY (2003) Endothelial cells require STAT3 for protection against endotoxin-induced inflammation. *J Exp Med* 198: 1517-1525

Kopf M, Baumann H, Freer G, Freudenberg M, Lamers M, Kishimoto T, Zinkernagel R, Bluethmann H, Kohler G (1994) Impaired immune and acute-phase responses in interleukin-6-deficient mice. *Nature* 368: 339-342

Krausgruber T, Fortelny N, Fife-Gernedl V, Senekowitsch M, Schuster LC, Lercher A, Nemc A, Schmidl C, Rendeiro AF, Bergthaler A *et al* (2020) Structural cells are key regulators of organ-specific immune responses. *Nature* 583: 296-302

Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25: 1754-1760

Liao Y, Smyth GK, Shi W (2014) featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 30: 923-930

Lin JX, Leonard WJ (2019) Fine-Tuning Cytokine Signals. *Annu Rev Immunol* 37: 295-324

Liu L, Okada S, Kong XF, Kreins AY, Cypowyj S, Abhyankar A, Toubiana J, Itan Y, Audry M, Nitschke P *et al* (2011) Gain-of-function human STAT1 mutations impair IL-17 immunity and underlie chronic mucocutaneous candidiasis. *J Exp Med* 208: 1635-1648

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15: 550

Lu JY, Shao W, Chang L, Yin Y, Li T, Zhang H, Hong Y, Percharde M, Guo L, Wu Z *et al* (2020) Genomic Repeats Categorize Genes with Distinct Functions for Orchestrated Regulation. *Cell Rep* 30: 3296-3311 e3295

Luis BAL, Calva-Mercado JJ (2018) Recurrent Spontaneous Intestinal Perforation due to STAT1 GOF Mutation. *Am J Gastroenterol* 113: 1057-1058

Martinez-Fabregas J, Wang L, Pohler E, Cozzani A, Wilmes S, Kazemian M, Mitra S, Moraga I (2020) CDK8 Fine-Tunes IL-6 Transcriptional Activities by Limiting STAT3 Resident Time at the Gene Loci. *Cell Rep* 33: 108545

Matsukawa A, Takeda K, Kudo S, Maeda T, Kagayama M, Akira S (2003) Aberrant inflammation and lethality to septic peritonitis in mice lacking STAT3 in macrophages and neutrophils. *J Immunol* 171: 6198-6205

McLoughlin RM, Hurst SM, Nowell MA, Harris DA, Horiuchi S, Morgan LW, Wilkinson TS, Yamamoto N, Topley N, Jones SA (2004) Differential regulation of neutrophil-activating chemokines by IL-6 and its soluble receptor isoforms. *J Immunol* 172: 5676-5683

McLoughlin RM, Jenkins BJ, Grail D, Williams AS, Fielding CA, Parker CR, Ernst M, Topley N, Jones SA (2005) IL-6 trans-signaling via STAT3 directs T cell infiltration in acute inflammation. *Proc Natl Acad Sci U S A* 102: 9589-9594

McLoughlin RM, Witowski J, Robson RL, Wilkinson TS, Hurst SM, Williams AS, Williams JD, Rose-John S, Jones SA, Topley N (2003) Interplay between IFN-gamma and IL-6 signaling governs neutrophil trafficking and apoptosis during acute inflammation. *The Journal of clinical investigation* 112: 598-607

Milner JD, Vogel TP, Forbes L, Ma CA, Stray-Pedersen A, Niemela JE, Lyons JJ, Engelhardt KR, Zhang Y, Topcagic N *et al* (2015) Early-onset lymphoproliferation and autoimmunity caused by germline STAT3 gain-of-function mutations. *Blood* 125: 591-599

Modur V, Li Y, Zimmerman GA, Prescott SM, McIntyre TM (1997) Retrograde inflammatory signaling from neutrophils to endothelial cells by soluble interleukin-6 receptor alpha. *The Journal of clinical investigation* 100: 2752-2756

Natoli G, Ghisletti S, Barozzi I (2011) The genomic landscapes of inflammation. *Genes Dev* 25: 101-106

Navarini AA, French LE, Hofbauer GF (2011) Interrupting IL-6-receptor signaling improves atopic dermatitis but associates with bacterial superinfection. *The Journal of allergy and clinical immunology* 128: 1128-1130

Olbrich P, Freeman AF (2018) STAT1 and STAT3 mutations: important lessons for clinical immunologists. *Expert Rev Clin Immunol* 14: 1029-1041

Ostuni R, Piccolo V, Barozzi I, Polletti S, Termanini A, Bonifacio S, Curina A, Prosperini E, Ghisletti S, Natoli G (2013) Latent enhancers activated by stimulation in differentiated cells. *Cell* 152: 157-171

Payer LM, Steranka JP, Yang WR, Kryatova M, Medabalimi S, Ardeljan D, Liu C, Boeke JD, Avramopoulos D, Burns KH (2017) Structural variants caused by Alu insertions are associated with risks for many human diseases. *Proc Natl Acad Sci U S A* 114: E3984-E3992

Peters A, Fowler KD, Chalmin F, Merkler D, Kuchroo VK, Pot C (2015) IL-27 Induces Th17 Differentiation in the Absence of STAT1 Signaling. *J Immunol* 195: 4144-4153

Polak P, Domany E (2006) Alu elements contain many binding sites for transcription factors and may play a role in regulation of developmental processes. *BMC Genomics* 7: 133

Qing Y, Stark GR (2004) Alternative activation of STAT1 and STAT3 in response to interferon-gamma. *J Biol Chem* 279: 41679-41685

Ramsay AJ, Husband AJ, Ramshaw IA, Bao S, Matthaei KI, Koehler G, Kopf M (1994) The role of interleukin-6 in mucosal IgA antibody responses in vivo. *Science* 264: 561-563

Robson RL, McLoughlin RM, Witowski J, Loetscher P, Wilkinson TS, Jones SA, Topley N (2001) Differential regulation of chemokine production in human peritoneal mesothelial cells: IFN-gamma controls neutrophil migration across the mesothelium in vitro and in vivo. *J Immunol* 167: 1028-1038

Roman AC, Benitez DA, Carvajal-Gonzalez JM, Fernandez-Salguero PM (2008) Genome-wide B1 retrotransposon binds the transcription factors dioxin receptor and Slug and regulates gene expression in vivo. *Proc Natl Acad Sci U S A* 105: 1632-1637

Schett G, Elewaut D, McInnes IB, Dayer JM, Neurath MF (2013) How cytokine networks fuel inflammation: Toward a cytokine-based disease taxonomy. *Nature medicine* 19: 822-824

Spencer S, Kostel Bal S, Egner W, Lango Allen H, Raza SI, Ma CA, Gurel M, Zhang Y, Sun G, Sabroe RA *et al* (2019) Loss of the interleukin-6 receptor causes immunodeficiency, atopy, and abnormal inflammatory responses. *J Exp Med*

Sun X, Wang X, Tang Z, Grivainis M, Kahler D, Yun C, Mita P, Fenyo D, Boeke JD (2018) Transcription factor profiling reveals molecular choreography and key regulators of human retrotransposon expression. *Proc Natl Acad Sci U S A* 115: E5526-E5535

Tajnik M, Vigilante A, Braun S, Hanel H, Luscombe NM, Ule J, Zarnack K, Konig J (2015) Intergenic Alu exonisation facilitates the evolution of tissue-specific transcript ends. *Nucleic Acids Res* 43: 10492-10505

Taniguchi K, Wu LW, Grivennikov SI, de Jong PR, Lian I, Yu FX, Wang K, Ho SB, Boland BS, Chang JT *et al* (2015) A gp130-Src-YAP module links inflammation to epithelial regeneration. *Nature* 519: 57-62

Tong AJ, Liu X, Thomas BJ, Lissner MM, Baker MR, Senagolage MD, Allred AL, Barish GD, Smale ST (2016) A Stringent Systems Approach Uncovers Gene-Specific Mechanisms Regulating Inflammation. *Cell* 165: 165-179

Toubiana J, Okada S, Hiller J, Oleastro M, Lagos Gomez M, Aldave Becerra JC, Ouachee-Chardin M, Fouyssac F, Girisha KM, Etzioni A *et al* (2016) Heterozygous STAT1 gain-of-function mutations underlie an unexpectedly broad clinical phenotype. *Blood* 127: 3154-3164

Twohig J, Cardus Figueras, A., Andrews, R., Wiede, F., Cossins, BC., Derrac Soria, A., Lewis, M., Townsend, MJ., Millrine, D., Li, J., Hill, D., Uceda Fernandez, J., Liu, X., Szomolay, B., Pepper, CJ., Taylor, PR., Pitzalis, C., Tiganis, T. Williams, NM., Jones, GW. & Jones, SA. (2019) Naïve T-cell activation re-tunes STAT1 signaling to deliver unique cytokine responses in memory CD4 T-cells. *Nature Immunology* 20: 458-470

Vahedi G, Kanno Y, Furumoto Y, Jiang K, Parker SC, Erdos MR, Davis SR, Roychoudhuri R, Restifo NP, Gadina M *et al* (2015) Super-enhancers delineate disease-associated regulatory nodes in T cells. *Nature* 520: 558-562

Vahedi G, Takahashi H, Nakayamada S, Sun HW, Sartorelli V, Kanno Y, O'Shea JJ (2012) STATs shape the active enhancer landscape of T cell populations. *Cell* 151: 981-993

Vassetzky NS, Kramerov DA (2013) SINEBase: a database and tool for SINE analysis. *Nucleic Acids Res* 41: D83-89

Villarino AV, Kanno Y, O'Shea JJ (2017) Mechanisms and consequences of Jak-STAT signaling in the immune system. *Nat Immunol* 18: 374-384

Volkman HE, Stetson DB (2014) The enemy within: endogenous retroelements and autoimmune disease. *Nat Immunol* 15: 415-422

Vuckovic D, Bao EL, Akbari P, Lareau CA, Mousas A, Jiang T, Chen MH, Raffield LM, Tardaguila M, Huffman JE *et al* (2020) The Polygenic and Monogenic Basis of Blood Traits and Diseases. *Cell* 182: 1214-1231 e1211

Welte T, Zhang SS, Wang T, Zhang Z, Hesslein DG, Yin Z, Kano A, Iwamoto Y, Li E, Craft JE *et al* (2003) STAT3 deletion during hematopoiesis causes Crohn's disease-like pathogenesis and lethality: a critical role of STAT3 in innate immunity. *Proc Natl Acad Sci U S A* 100: 1879-1884

Wiede F, Brodnicki TC, Goh PK, Leong YA, Jones GW, Yu D, Baxter AG, Jones SA, Kay TWH, Tiganis T (2019) T-Cell-Specific PTPN2 Deficiency in NOD Mice Accelerates the Development of Type 1 Diabetes and Autoimmune Comorbidities. *Diabetes* 68: 1251-1266

Wilmes S, Jeffrey PA, Martinez-Fabregas J, Hafer M, Fyfe PK, Pohler E, Gaggero S, Lopez-Garcia M, Lythe G, Taylor C *et al* (2021) Competitive binding of STATs to receptor phospho-Tyr motifs accounts for altered cytokine responses. *Elife* 10

Xie H, Wang M, Bonaldo Mde F, Smith C, Rajaram V, Goldman S, Tomita T, Soares MB (2009) High-throughput sequence-based epigenomic analysis of Alu repeats in human cerebellum. *Nucleic Acids Res* 37: 4331-4340

Xing Z, Gauldie J, Cox G, Baumann H, Jordana M, Lei XF, Achong MK (1998) IL-6 is an antiinflammatory cytokine required for controlling local or systemic acute inflammatory responses. *The Journal of clinical investigation* 101: 311-320

Yang X, Yan H, Jiang N, Yu Z, Yuan J, Ni Z, Fang W (2020) IL-6 trans-signaling drives a STAT3-dependent pathway that leads to structural alterations of the peritoneal membrane. *Am J Physiol Renal Physiol* 318: F338-F353

Yang XP, Ghoreschi K, Steward-Tharp SM, Rodriguez-Canales J, Zhu J, Grainger JR, Hirahara K, Sun HW, Wei L, Vahedi G *et al* (2011) Opposing regulation of the locus encoding IL-17 through direct, reciprocal actions of STAT3 and STAT5. *Nat Immunol* 12: 247-254

Zeng M, Hu Z, Shi X, Li X, Zhan X, Li XD, Wang J, Choi JH, Wang KW, Purrington T *et al* (2014) MAVS, cGAS, and endogenous retroviruses in T-independent B cell responses. *Science* 346: 1486-1492

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M, Li W *et al* (2008) Model-based analysis of ChIP-Seq (MACS). *Genome Biol* 9: R137

Figures & Legends

Figure-1

Th1 cells augment the stromal response to acute inflammation. (A) RNA-seq was performed on stromal tissues extracted from SES challenged mice. Volcano plots show differential expression analysis (LIMMA) and statistical thresholding ($p_{adj}<0.05$, $Log_2FC>1.75$) of datasets at 3 and 6 hours post treatment. Mice received (i.p.) SES alone (SES) or SES in combination with Th1 cells (SES+Th1). Each test condition was compared to untreated mice (SES), or mice receiving SES and naïve CD4[+] T-cells. **(B)** K-means clustering (n=4) of data shown in Panel-A (heatmap). UMAP visualizations show the distribution of gene clusters and their link to SES (blue) or SES+Th1 (red) datasets. **(C)** Gene Ontology (msigdb) shows the top-5 biological processes for each cluster (left). Each cluster descriptor is coded to match UMAP clustering colors in Panel-B. Representative examples of genes in each cluster are presented as heatmaps (right).

Figure-2

Regulatory signatures of cytokine signalling in acute inflammation. (A) Ingenuity Pathway Analysis of biological processes involved in SES-induced inflammation. **(B)** Gene set enrichment analysis shows the impact of Th1 cells on Jak-STAT cytokine signalling and interferon regulated outcomes. A summary of the complete GSEA analysis is provided in *Supplemental Figure-1*. **(C)** Ingenuity Pathway Analysis of upstream regulators. Predicted p-value (Fisher's exact test) identifies gene signatures characteristic of specific transcription factors linked to TLR (NF-κB, Rel-A and shown), interferon (Yellow; IRF family members are shown) and Jak-STAT cytokine signaling (Red; STAT1 and STAT3 are displayed).

Figure-3

Th1 cells shape the transcriptional output of IL-6. (A) Heatmap of IL-6 regulated transcripts identified by differential expression analysis (LIMMA) of *wt* versus *Il6*[-/-] mice. Data is arranged by K-means clustering (n=3) of significantly regulated transcripts (Cluster-1; red, Cluster-2; grey, Cluster-3; blue). **(B)** UMAP visualization of all IL-6 regulated transcripts ($p_{adj}<0.05$, $Log_2FC>1.75$). **(C)** Alignment of representative transcripts from each cluster against the top-20 biological processes identified by gene ontology enrichment analysis of 241 IL-6 regulated transcripts (msigdb). **(D)** Flow cytometric analysis of infiltrating neutrophil effector function (see *Supplemental Figure-2*). Peritonitis was induced in WT mice by administration of fluorescently labelled Staphylococcus epidermidis ($5x10^8$ CFU) in the presence of sgp130Fc (250ng/mouse). Changes in phagocytosis and respiratory burst are shown (mean ± SEM, n=4; **p<0.01, *p<0.05).

Figure-4

ChIP-seq analysis of STAT1 and STAT3 involvement in SES-induced inflammation. Genomic DNA from the peritoneal membrane of mice challenged with SES alone or SES+Th1 was extracted at 3 hours. Peak calling and downstream processing are described in *Materials and Methods*. Volcano plots summarize ChIP-seq profiling. Each dot represents a peak (Grey = q>0.05 and/or $\log_2FC>1.75$). Peaks below the significance (q<0.05) and above the $\log_2FC$ (>1.75) cutoff values are highlighted in blue (*wt*) and red (*Il6$^{-/-}$*).

Figure-5

STAT1 and STAT3 interplay shapes gene regulation. (A) Volcano plot of RNA-seq data showing differentially regulated gene expression (*Il6$^{-/-}$* vs wt; $p_{adj}<0.05$, $\log_2FC>1.75$) in SES+Th1 treated *Il6$^{-/-}$* mice (3-hour post administration). Differential gene regulation is shown for representative genes displaying reciprocal STAT1 and STAT3 binding in ChIP-seq datasets from *wt* and *Il6$^{-/-}$* mice (summarized in inset). **(B)** Euclidean clustering of the 33 genes depicted in Panel-A.

Figure-6

Mapping of chromatin accessibility by ATAC-seq. (A) Heatmap visualization of ATAC-seq profiling of peritoneal extracts from mice treated with SES and Th1 cells (Top). The peak count frequency of sequence reads associated with transcription start sites (TSS) is shown for *wt* and *Il6$^{-/-}$* mice (Bottom). **(B)** Histogram shows chromatin accessibility at TSS linked with IL-6 regulated genes (Top). Graph shows the genomic distribution of IL-6 regulated loci (Bottom). **(C)** Fold change (*Il6$^{-/-}$* vs *wt*) in ATAC-seq reads at indicated genomic features. **(D)** Motif enrichment analysis (MEME-ChIP) of genomic regions identified in differential binding analysis of ATAC-seq datasets. Annotations identify putative transcription factor motifs associated with sequencing peaks.

Figure-7

ChIP-seq provides evidence of STAT1-STAT3 cross-regulation. (A) Circos plots show the relative binding of STAT transcription factors to putative enhancers defined by either P300 binding (either *wt* or *Il6$^{-/-}$* mice) or sites bearing homology to a *de novo* GAS-like motif identified by STAT1 and STAT3 ChIP-seq (Top). Binding to these sites was calculated using the Bedtools intersect algorithm. **(B)** Motif enrichment analysis of STAT3 ChIP-seq dataset from wt mice (SES+Th1). Pairwise comparison using the Pearson method was generated using Motif Alignment and Search Tool (MAST; Meme ChIP suite). Annotation shows the source algorithm of each motif (Centrimo (green), Meme (orange), and DREME (grey); Meme ChIP suite). A cluster is highlighted (hashed line) that maps to a sequence displaying homology with a GAS-like motif.

Figure-8

Identification of a GAS-like motif in Alu elements. (A) Multiple sequence alignment (Muscle; EBI) of human and mouse retroelement sequences downloaded from the dfam database (n=66 sequences). The Alu-GAS motif (CCTGTAATCCCAGC) identified by STAT1 and STAT3 ChIP-seq is located. Conserved regions are shown in blue, and a summary of the sequence identity is shown in orange (0-100%). **(B)** An annotated summary retroelement sequence is shown (top) locating representative secondary motifs (+/-150bp) relative to the Alu-GAS motif. Spaced motif analysis (Spamo; Meme ChIP suite) of these secondary sites are shown. **(C)** Quantitation of retroelement binding in ChIP-seq datasets based on repeatmasker annotations.

Figure 9

Association of GAS-Alu motif with human physiology. (A) MAGMA enrichment analysis of genes identified through enhancer profiling (FANTOM5 consortium) of genes linked to the GAS-Alu motif (see *Supplemental Figure 5*). Analysis identified significant enrichment for associations in 257 GWAS. For each GWAS, a vector was formed of the -log p-values of the motif-linked genes, and Pearson correlations calculated for each pair of p-value vectors. UMAP distribution identifies GWAS datasets sharing common traits. Those displaying links to immune cell function are shown (comprising Cluster-2). **(B)** Top gene-wide p-values identified in Cluster-2 phenotypes. Horizontal color bar identifies GWAS phenotypes linked to immune cell regulation (grey; n=68); immunopathology (red; n=13); and other (pink; n=3). **(C)** Gene Ontology enrichment analysis of the genes identified in Panel-B (n=53).

Figure-1

Figure-2

A

B

C

Figure-4

A

Figure-5

Figure-6

Figure-7

A

wt   GAS-like   STAT1   P300   STAT3

Il6⁻/⁻   GAS-like   STAT1   P300   STAT3

B

STAT3 ChIP/SES+Th1

MIN   Pearson   MAX

GAS-like

■ Centrimo
■ DREME
■ MEME

Figure-8

A

CCTGTAATCCCAGC

Homo Sapiens

7SLRNA

Mus Musculus

%Identity

B

RUNX PolIII A-box STAT1/STAT3 TBX21

GGCCGGG**CGGCGG**G**TGGC**TCACG**CCTGTAATCCCAGC**ACTTTG**GGAGGCCGAGAG**GCGGGCGGATCACTTGAG
GTCAGGAGTTCGAGACCTAGTGCGCTATGCCGATCGGGTGTCCGCACTAAGCCTGGCATCAACATGGTGAA
ACCCCGTCTCTACTAAAAATACAAAAAATTAGCCGGGCGTGGTGGCGGGGC***CCTGTAGTCCCAGC***TACTCGG
AGAGGCTGAGGCAGGAGAATGGCGTGAACCCCGGGAGGCGGAGCTTGCAGTGATGATCAGTGCCGAGATCGC
GCCTGTGAATAGCCACTGCAGTCCGCACTCCAGCCTGGGCGACAAGAGCGAGACTCCGTTCTC**AAAAAAAAA**
AAAAAAAAAAAAAAAAAAAAAA   **IRF**

STAT3 ChIP
- IRF3 e-value=4.52E61
- TBX21 e-value=1.31E74
- RUNX3 e-value=1.01E107

STAT1 ChIP
- IRF3 e-value=5.31E121
- TBX21 e-value=1.36E218
- RUNX3 e-value=5.68E294

Fantom5 enhancer
- TBX21 e-value=1.02E21
- RUNX3 e-value=4.35E46

Homo sapiens

C

Retroelement binding (absolute)

0   3E3   6E3   9E3

SES
- STAT1
- STAT3
- STAT1
- STAT3

SES+Th1
- STAT1
- STAT3
- STAT1
- STAT3

wt
Il6⁻/⁻

L1   B1