Chronic Jetlag Accelerates Pancreatic Neoplasia in Conditional *Kras*-Mutant Mice

Authors: Patrick B Schwartz, MD[1], Morgan T Walcheck, PhD[1], Manabu Nukaya, PhD[1], Derek M

Pavelec, PhD[2], Kristina A Matkowskyj[3,4,5], and Sean M Ronnekleiv-Kelly, MD[1,5]

Affiliations:

[1]Department of Surgery, Division of Surgical Oncology, University of Wisconsin School of Medicine and

Public Health, Madison, WI

[2]Biotechnology Center, University of Wisconsin, Madison, WI

[3]Department of Pathology and Laboratory Medicine, University of Wisconsin School of Medicine and

Public Health, Madison, Wisconsin.

[4]William S Middleton Memorial Veterans Hospital, Madison, Wisconsin.

[5]University of Wisconsin Carbone Cancer Center, University of Wisconsin School of Medicine and

Public Health, Madison, WI

Corresponding Author: Sean M Ronnekleiv-Kelly, MD (ronnekleiv-kelly@surgery.wisc.edu)

Address: 600 Highland Ave. Bx 7375, K3/705 CSC, Madison, WI, 53792

Phone: (608) 262-2025 **Fax:** (608) 252-0913

Running Title: Chronic Jetlag Accelerates Pancreatic Neoplasia

21 **Disclosure:** The authors declare no conflict of interest.

Abstract:

Misalignment of the circadian clock compared to environmental cues causes circadian desynchrony, which is pervasive in humans. Clock misalignment can lead to various pathologies including obesity and diabetes, both of which are associated with pancreatic ductal adenocarcinoma - a devastating cancer with an 80% five-year mortality rate. Although circadian desynchrony is associated with an increased risk of several solid-organ cancers, the correlation between clock misalignment and pancreas cancer is unclear. Using a chronic jetlag model, we investigated the impact of clock misalignment on pancreas cancer initiation in mice harboring a pancreas-specific activated *Kras* mutation. We found that chronic jetlag accelerated the development of pancreatic cancer precursor lesions, with a concomitant increase in precursor lesion grade. Cell-autonomous knock-out of the clock in pancreatic epithelial cells of *Kras*-mutant mice demonstrated no acceleration of precursor lesion formation, indicating non-cell-autonomous clock dysfunction was responsible for the expedited tumor development. Therefore, we applied single-cell RNA sequencing over time and identified fibroblasts as the cell population manifesting the greatest clock-dependent changes, with enrichment of specific cancer-associated fibroblast pathways due to circadian misalignment. Collectively, these results suggest fibroblasts as the putative target of chronic jetlag-induced accelerated pancreas cancer initiation.

Keywords: Circadian, Pancreas, Jetlag, Pancreatic Ductal Adenocarcinoma, Misalignment

Introduction:

Circadian clock proteins coordinate daily biological rhythms with an oscillation of 24 hours (McIntosh et al. 2010). Control from the central clock in the hypothalamus assimilates environmental cues (e.g. day/night cycles) to modulate peripheral organ rhythmic expression of core circadian genes and behavioral patterns (e.g. sleep/wake cycle) (McIntosh et al. 2010). In addition to central control, nearly all cells have an autonomous circadian clock based on transcriptional and translational feedback mechanisms of core clock genes (CCGs) (Lee et al. 2001; Preitner et al. 2002; Sato et al. 2004; Takahashi 2017). Collectively, the CCGs of the central and peripheral clocks orchestrate the transcription of circadian output genes, also known as clock-controlled genes (Takahashi 2017; Cox and Takahashi 2019). The central and peripheral clocks in mammals (and other organisms) have evolved to anticipate environmental changes that occur over the 24-hr day/night cycle, and in doing so, appropriately sequence complex internal processes to align with external cues (Takahashi 2017). For instance, in humans, energy intake, physical activity, and cognitive activities are anticipated to occur during the day while rest and sleep are expected to occur at night (Pittendrigh 1993; Wright et al. 2013). Consequently, this network of circadian control coordinates the optimal timing of critical cell and organ processes such as metabolism, cell division, apoptosis, and immune function (Matsuo et al. 2003; Bass and Takahashi 2010; Lee and Sancar 2011; Scheiermann et al. 2013).

Notably, the internal clock can be adjusted, or re-entrained, based on changes to environmental zeitgebers or 'time-givers', such as through alteration of the day/night cycle (Wright et al. 2013). However, persistent misalignment between environmental cues compared to the internal clocks can result in circadian desynchrony with associated aberrant gene signaling (Kettner et al. 2016; West et al. 2017). Unfortunately, this type of clock dysfunction (misalignment or circadian desynchrony) is pervasive in humans and this condition affects a substantial proportion of the human population, particularly shift workers, whose clock timing can be altered dramatically compared to environmental zeitgebers by work schedule requirements (Bass and Lazar 2016). Further, this environmental or behavior-induced

desynchrony can have profound implications for human health including the promotion of various

pathologies such as the metabolic syndromes of obesity and diabetes (Arble et al. 2009; Roenneberg et al.

2012; Perelis et al. 2016). There is also a strong correlation between circadian clock dysfunction and the

risk of developing several different solid organ cancers, including lung cancer, liver cancer, and colon

cancer (Schernhammer et al. 2003; Papagiannakopoulos et al. 2016; Kettner et al. 2016). Meanwhile, a

history of shiftwork has been shown to increase the risk of pancreatic ductal adenocarcinoma (PDAC) in

men over 2-fold (Parent et al. 2012); concordantly, obesity and diabetes are both risk factors for the

development of PDAC (Schernhammer et al. 2003; Marcheva et al. 2010; Kettner et al. 2016). Despite

this converging evidence and the substantive data linking the circadian clock to pancreas-specific disease

processes (Marcheva et al. 2010; Perelis et al. 2015), the correlation between circadian desynchrony and

the risk of developing PDAC remains unclear.

We sought to establish the connection between clock misalignment – a process that is highly prevalent in

humans – and PDAC because this would have important implications for understanding cancer

pathogenesis and risk mitigation in this deadly disease. Risk factor reduction leading to cancer prevention

is particularly relevant in this highly lethal malignancy because few people survive after cancer diagnosis:

of the 60,430 people estimated to be diagnosed in 2021, approximately 48,220 (~80%) will die of their

disease (Siegel et al. 2021). As a foundation for uncovering the link between circadian clock dysfunction

and the development of PDAC, we investigated whether circadian misalignment impacted tumor

initiation within the pancreas. We used *LSL-Kras$^{G12D/+}$; Pdx1-Cre* mice (KC) that harbor a *Kras* mutation

in pancreatic acinar and ductal cells (Hingorani et al. 2003). KC mice develop the full spectrum of

neoplastic changes including acinar-to-ductal metaplasia (earliest lesion), pancreatic intra-epithelial

neoplasia 1 (PanIN-1), higher grade PanIN lesions (PanIN-2, -3), and ultimately PDAC (Hingorani et al.

2003). We implemented a well-characterized model of chronic jetlag (CJ) that is known to cause

environmental-induced misalignment and phase-shift of the clock (Filipski et al. 2004;

Papagiannakopoulos et al. 2016; Schwartz et al. 2021) and is meant to mimic the environmental and

behavioral misalignment experienced by humans (Vetter 2020). The abrupt change in timing of the

light/dark cycle with CJ causes a period of desynchrony, with differences in the rate of re-entrainment by

internal clocks or behavior (Yamazaki et al. 2000; West and Bechtold 2015). Previously, we have found

that behavioral re-entrainment occurs much more rapidly than re-entrainment of the pancreatic clock

(Schwartz et al. 2021). Moreover, this type of protocol (CJ) has been shown to promote the growth of

several other cancers, including osteosarcoma, and hepatocellular carcinoma in mice (Filipski et al. 2004;

Filipski et al. 2009; Kettner et al. 2016).

In the current study, we hypothesized that circadian clock desynchrony through CJ would accelerate the

initiation of PanIN-1 and progression to higher-grade PanINs or PDAC. We found that the pancreata of

KC mice subjected to CJ developed increased fibroinflammatory infiltrate and neoplastic lesion formation

(i.e. PanINs) compared to KC mice under standard conditions. To understand whether cell-autonomous

clock dysfunction was responsible for the accelerated PanIN development, we generated KC mice with an

abolished clock (*Bmal1* knock-out) in the pancreatic ductal and acinar cells (cellular origin of PDAC).

These mice did not replicate the phenotype observed in the CJ group of KC mice, indicating a non-cell-

autonomous role of clock dysfunction that is facilitating the PanIN progression. To investigate further, we

used single-cell RNA sequencing (scRNA-seq) and identified cancer-associated fibroblasts (CAFs) as the

putative target of the CJ-induced phenotype. This work serves as a foundation for understanding the link

between circadian misalignment through CJ and the acceleration of PDAC development.

Materials and Methods:

Mouse Care and Husbandry

All animal studies were conducted according to an approved protocol (M005959) by the University of

Wisconsin School of Medicine and Public Health (UW SMPH) Institutional Animal Care and Use

Committee (IACUC). Male and female mice were housed in an Assessment and Accreditation of

Laboratory Animal Care (AALAC) accredited selective pathogen-free facility (UW Medical Sciences

Center) on corncob bedding with chow diet (Mouse diet 9F 5020; PMI Nutrition International) and water

ad libitum.

Conditional *LSL-Kras*$^{G12D/+}$ (K) mice (B6.129S4-Kras tm4Tyj/J (#008179)) were bred with mice

expressing Cre-Recombinase driven by the *Pdx1* promoter (C) (B6.FVB-Tg(Pdx1-cre)6Tuv/J) (#014647))

to generate *LSL-Kras*$^{G12D/+}$, *Pdx1-Cre* (KC) mice (Hingorani et al. 2003). The KC mouse has been well

characterized as a pancreas pre-cancer model (Hingorani et al. 2003). To evaluate the effect of loss of

clock function on KC mouse pathology the conditional *Bmal1*$^{fx/fx}$ mouse, kindly gifted by Professor

Christopher Bradfield (University of Wisconsin-Madison, Madison, WI), was crossed with the KC mouse

to generate the *Bmal1*$^{fx/fx}$, *LSL-Kras*$^{G12D/+}$; *Pdx1-Cre* (BKC) mouse model (Liu et al. 2014; Johnson et al.

2014). To identify cell types targeted by *Pdx1-Cre* recombinase in the KC mouse model, the KC mouse

was additionally crossed with the *LSL-tdTomato* reporter (Ai14) mouse (B6.Cg-

Gt(ROSA)26Sortm14(CAG-tdTomato)Hze/J (#007914)) to produce the KCT mouse (*LSL-R26*$^{CAG-tdTomato}$;

LSL-Kras$^{G12D/+}$; *Pdx1-Cre*). The presence of each allele, including activated *Kras*-mutation, was

confirmed with genotyping PCR according to Jackson Laboratory's (Bar Harbor, ME) protocols as

previously described (Liu et al. 2014; Walcheck et al. 2021). The K, C, and T mice were purchased from

Jackson Laboratory.

Chronic Jetlag Protocol

Starting at 4-6 weeks of age, KC mice were subjected to either a normal circadian (standard lighting) 12-

hour/12-hour light: dark (LD) cycle condition, or a chronic jetlag protocol (CJ) known to induce circadian

desynchrony and mimic altered environmental exposure experienced by humans (Filipski et al. 2004;

Papagiannakopoulos et al. 2016; Vetter 2020; Schwartz et al. 2021). CJ consisted of a 12-hour/12-hour

LD cycle, phase-shifted forward 4 hours every 2-3 days. Mice were kept under their respective conditions

until the study endpoint at 5 and 9 months of age.

Histologic Analysis

Mouse pancreatic tissue was fixed in 10% neutral buffered formalin for 48 hours and then stored in 70%

ethanol until further processing. Tissue was processed, paraffin embedded, sectioned onto slides, and

stained with hematoxylin and eosin (H&E) by the University of Wisconsin Carbone Cancer Center

(UWCCC) Experimental Animal Pathology Laboratory (EAPL) Core. Each pancreas was sectioned at

200 μm steps for at least 6 slides. All slides processed were assessed for the presence of pancreas cancer

precursor lesions (pancreatic intraepithelial neoplasia or PanIN) along with grading on a scale of 1-3 (1 =

lowest grade, 2 = intermediate grade, and 3 = high grade) and/or pancreatic ductal adenocarcinoma

(PDAC). This was performed by a blinded, board-certified, gastrointestinal pathologist (KAM) at 5

months (n = 36 normal circadian [24 male and 12 female] vs n = 26 CJ [14 male and 12 female]) and 9

months age (n = 33 normal circadian [19 male and 14 female] vs n = 39 CJ [23 male and 16 female]).

Additional BKC mice (n = 24 [14 male and 10 female]) were assessed at 5 months. Categorical

comparisons between groups were made with the Chi-squared or Fisher's Exact test as appropriate. KC

mice develop fibroinflammatory infiltrate associated with adjacent PanINs and are collectively part of the

pancreatic changes with neoplasm formation (Hingorani et al. 2003). Quantification of the total percent

pancreas involvement by the fibroinflammatory infiltrate and PanINs (referred to collectively as

FI/PanIN) was made by 3 separate reviewers, and differences were tested using Student's t-test between

groups. For evaluation, two slides were selected from each pancreas comprising the largest area of

available tissue. Values are presented as the mean with associated standard deviation (SD). Inter-rater

reliability statistics were computed and the intraclass correlation coefficient is reported with associated

95% confidence intervals.

Immunohistochemistry

KCT mouse pancreas samples were formalin-fixed, paraffin embedded, and sectioned by the EAPL Core

facility. Immunohistochemistry (IHC) for red fluorescence protein (tdTomato) was also performed by

EAPL. Sections were deparaffinized in xylenes and hydrated through graded alcohols to distilled water.

Antigens were retrieved using citrate buffer pH 6.0 (10 mM Citric Acid and 0.05% tween 20).

Endogenous peroxidase was blocked with 0.3% H_2O_2 and blocking of non-specific binding was

performed using 10% goat serum. Sections were incubated with rabbit anti-RFP antibody (600-401-379,

Rockland Inc, Pottstown, PA) (1:1600) followed by ImmPRESS goat anti-rabbit IgG (MP-7451, Vector

Laboratories, Burlingame, CA). Detection was performed using the DAB substrate kit (8059S, Cell

Signaling Technology, Danvers, MA). Samples were counterstained using Mayer's hematoxylin

(MHS32, Millipore-Sigma, St. Louis, MO).

scRNAseq Sample Preparation, Library Construction, and Sequencing

Five- and 9-month-old normal circadian and CJ KC mice were sacrificed with cervical dislocation and the

pancreas was rapidly dissected (< 60 seconds) and placed in cold PBS (Thermo Fisher Scientific,

Waltham, MA). One male and female pancreas was pooled from each condition and time point after

demonstration of no discernable sex-specific differences (Takele Assefa et al. 2020). Pancreas samples

were minced until the tissue was < 1 mm in diameter, and then centrifuged at 900 RPM for 5 minutes to

remove fat contamination. Samples were then incubated with 5 mL of Collagenase P (Sigma Aldrich, St.

Louis, MO) in HBSS (Thermo Fisher Scientific) at 37° C for 15 minutes while being shaken at 100 RPM.

Additionally, samples were manually agitated every 2-3 minutes. The digestion reaction was quenched

with 10 mL R10 media. Samples were then centrifuged at 900 RPM and washed twice with 10 mL of

R10. Samples were then strained in a gradient fashion using 500 μm, 100 μm, and 40 μm strainers.

Samples were then live/dead sorted into single-cell populations and were transferred to the University of

Wisconsin Gene Expression Center (GEC) for Library Preparation.

In brief, libraries were constructed according to the Chromium NextGEM Single Cell 3' Reagent Kit v3.1

User Guide, Rev.D (10x Genomics, Pleasanton, CA). Single-cell suspension cell concentration and

viability were quantified on the Countess II (Thermo Fisher Scientific) using 0.4% Trypan Blue

(Invitrogen, Carlsbad, CA). Single-cell droplet encapsulation was performed using the Single Cell G Chip

and the 10x Genomics Chromium Controller. Following the completion of the Chromium run, the Gel

Beads in Emulsion (GEMs) were transferred to emulsion-safe strip tubes for GEM-RT using an

Eppendorf MasterCycler Pro thermocycler (Eppendorf, Hamburg, Germany). Following reverse

transcription (RT), GEMs were broken, and the pooled single-cell cDNA was collected in Recovery

Agent, purified using DynaBeads MyOne Silane beads (Thermo Fisher Scientific), and amplified with the

appropriate number of PCR cycles based on the number of cells targeted for recovery. Post-cDNA

amplified product was purified using SPRIselect (Beckman Coulter, Brea, CA) and quantified on a

Bioanalyzer 2100 (Agilent, Santa Clara, CA) using the High Sensitivity DNA kit. Adapters were then

added to the libraries after fragmentation, end repair, A-tailing, and double-sided size selection using

SPRIselect. Following adapter ligation, libraries were cleaned up using SPRIselect and sample-specific

indexes (Chromium i7 Multiplex Kit, 10x Genomics) were added by sample index PCR using the

appropriate number of PCR cycles based on the cDNA yield calculated by the Bioanalyzer quantification.

After sample index PCR, samples were double-size selected using SPRIselect and quantified by the Qubit

High Sensitivity DNA Kit (Thermo Fisher Scientific). Libraries were sequenced on a NovaSeq6000

(Illumina, San Diego, CA).

scRNAseq Data Analysis

Single-cell RNAseq data were analyzed by the UW Bioinformatics Resource Center. Data were

demultiplexed using the Cell Ranger Single Cell Software Suite, mkfastq command wrapped around

Illumina's bcl2fastq. The MiSeq balancing run was quality-controlled using calculations based on UMI-

tools (Smith et al. 2017). Sample libraries were balanced for the number of estimated reads per cell and

run on an Illumina NovaSeq system. Cell Ranger software was then used to perform demultiplexing,

alignment, filtering, barcode counting, UMI counting, and gene expression estimation for each sample

according to the 10x Genomics documentation (https://support.10xgenomics.com/single-cell-gene-

expression/software/pipelines/latest/what-is-cell-ranger). The gene expression estimates from each sample

were then aggregated using Cellranger (cellranger aggr) to compare experimental groups with normalized

sequencing-depth and expression data.

Gene expression data was then processed using the Seurat version 4 pipeline for data integration (Hao et

al. 2021). Data was loaded into Seurat. Low quality (<200 features or >20% mitochondrial gene content)

and doublets were removed. Cells expressing more than 1% hemoglobin gene content were filtered out.

The normal circadian and CJ KC data were merged and integrated using the SCTransform v2 wrapper

with 3000 unique features (Choudhary and Satija 2022). SCTransform allows for normalization and better

batch correction based on a regularized negative binomial regression. The regression additionally allowed

for correction of the cell cycle effect and mitochondrial gene content (Nestorowa et al. 2016). A principal

component analysis (PCA) was performed with the number of dimensions (principal components)

selected based on an elbow plot. Data were visualized using the Uniform Manifold Approximation and

Projection for Dimension Reduction (UMAP) (McInnes et al. 2020). Cell identities were called based on

the expression of known markers for each cell type and scored using the ScType package to cross-validate

identities following the default settings for SCT-transformed data in the provided vignette (Ianevski et al.

2022). Differential expression between conditions and time points was performed between clusters or

conditions in Seurat using the MAST test based on a log fold change threshold of 0.25 and a false

discovery rate (FDR) corrected p value of $p < 0.05$. The MAST test within Seurat is based on the MAST

package, which is specifically designed to handle differential expression in scRNA-seq data and utilizes a

hurdle model (Finak et al. 2015). Gene Set Enrichment Analysis (GSEA) was conducted with the fgsea

package leveraged against the oncogenic signature gene set (c6.all.v7.5.1.symbols) and biologic process

subset of gene ontology (GO BP) (c5.go.v7.5.1.symbols) (Sergushichev 2016). Pseudotime trajectory

analysis was performed with the Monocle 3 package (Trapnell et al. 2014). Genes correlated with

pseudotime progression were computed using the Moran's I test and those genes with an FDR $p < 0.05$

were considered significant. All scRNAseq data is publicly available through gene expression omnibus

(GEO) (Accession number: GSE209629). All analyses in this manuscript were conducted in R version

4.2.0 (Vienna, Austria) or GraphPad Prism version 8.3.1 (Dotmatics, Boston, MA).

Results:

To test the hypothesis that circadian misalignment by chronic jetlag (CJ) accelerates tumor initiation (i.e.

PanIN or pre-cancerous lesion formation) and development of PDAC, we elected to utilize the *LSL-*

Kras^G12D/+, *Pdx1-Cre* (KC) mouse model – a well-characterized PDAC development model – and

assessed the burden of neoplastic lesions (i.e. PanIN) and PDAC (Hingorani et al. 2003). We have

previously shown that the CJ protocol results in circadian misalignment with phase-shift of core clock

gene (CCG) expression and alteration of diurnally expressed genes in the pancreas compared to mice

under normal circadian (standard lighting) conditions (Schwartz et al. 2021). Furthermore, we have

demonstrated that after normalizing the time of lights on/lights off (i.e. light/dark cycle) in CJ mice,

complete re-entrainment of the pancreatic clock does not occur out to day 10 post-normalization; this is in

contrast to behavioral re-entrainment which occurred by 3 days (Schwartz et al. 2021). Therefore, the CJ

protocol is a robust protocol to evaluate the impact of circadian misalignment on the tumor initiation and

progression of pancreatic pre-cancerous lesions.

We first evaluated KC mice at age 5 months, because existing literature demonstrates that 5-month KC

mice develop neoplastic changes consisting of early PanINs (i.e. PanIN-1) but most of the pancreas

remains normal (Hingorani et al. 2003). Therefore, evaluating the extent of pancreas involvement by

PanIN at this time point (not incidence) should allow for the detection of any acceleration in tumor

initiation due to circadian misalignment (CJ). All sections were scored for their degree of pancreatic

involvement by neoplastic lesions (e.g. PanINs) and the associated fibroinflammatory infiltrate, hereafter

referred to as FI/PanIN. This was calculated by determining the percent of the pancreas that was involved

with FI/PanIN by 3 independent reviewers, and there was an excellent intraclass correlation coefficient of

0.896 (95% CI 0.868-0.919; $p < 0.01$). We found that KC mice that were exposed to CJ (n = 26)

displayed an approximate 2-fold mean increase (± standard deviation [SD]) in FI/PanIN involvement of

the pancreas at 5 months compared to KC mice under standard lighting conditions (n = 36) (22.47% ±

19.08 vs 12.73% ± 12.08; $p = 0.027$, **Figure 1**). There were no sex-specific differences at 5 months for

normal circadian (males 13.14% ± 12.68 vs. females 11.93% ± 11.27; $p = 0.77$) or CJ KC mice (males

17.27% ± 12.94 vs. females 28.53% ± 23.55; $p = 0.16$).

Due to the expected progression of FI/PanIN at 9 months, the extent becomes less relevant (Hingorani et

al. 2003). Instead, by 9 months, the incidence of higher grade PanINs (i.e. PanIN-2 and PanIN-3) and

development of PDAC better reflects the progression of pre-cancerous lesions into cancer. This is because

i) not all PanIN-1s progress to higher-grade lesions, ii) PanIN-2,-3, and PDAC represent tumor

progression, and iii) KC mice are expected to develop a significant incidence of PanIN-2, -3, and PDAC

with increasing age (Hingorani et al. 2003). In contrast, higher-grade lesions are rarely observed by

approximately 5 months of age (Hingorani et al. 2003). Concordantly, we found that KC mice under

normal circadian conditions exhibited almost exclusively PanIN-1, with only 5.6% (n = 2/36)

demonstrating a PanIN-2 lesion at age 5 months; no 5-month KC mice under normal conditions

developed PanIN-3 or PDAC (**Figure 2A/C**). Interestingly, although not significant, 7.7% (n = 2/26) of

KC mice subjected to CJ developed PanIN-2, and one mouse developed dedifferentiated PDAC which

had replaced the entirety of the pancreatic parenchyma (**Figure 2A/C**).

As expected, the incidence of advanced lesions increased by 9 months of age in KC mice under normal

circadian conditions to 3% (n = 1/33) harboring PanIN-2, 6.1% (n = 2/33) developing PanIN-3, and

15.2% (n = 5/33) of mice progressing to PDAC (**Figure 2B/C**). Notably, although the incidence of

PanIN-3 (10.3% [n = 4/39]) and PDAC (10.3% [n = 4/39]) was similar between CJ and normal circadian

KC mice at 9 months, there was a significant increase in PanIN-2 formation in KC mice due to CJ (23.1%

[n = 9/39] vs 3% [n = 1/33], $p = 0.02$) (**Figure 2B-E**). There were no sex-specific differences in PanIN-2

incidence for normal circadian (male 0/19 (0%) vs. female 1/14 (7.1%); $p = 0.42$) and CJ mice (male 4/23

(17.4%) vs. female 5/16 (31.2%); $p = 0.44$). Collectively, this data demonstrates that circadian

desynchrony promoted by CJ in KC mice caused accelerated development of FI/PanIN at 5 months

leading to an increased incidence of higher grade PanINs (i.e. PanIN-2) by 9 months, and these findings

were independent of sex. This strongly supports the assertion that circadian misalignment (desynchrony)

facilitates tumor initiation and progression of the neoplastic PanIN lesions.

Pancreatic Clock Gene Deletion Abrogated Tumor Initiation

Different cell populations within the pancreas can lead to tumor development. For instance, islet cells can

transform into neuroendocrine tumors (Zhang et al. 2013; Li and Xie 2022), whereas acinar-to-ductal

metaplasia is the initiating process in PanIN formation (and consequently PDAC) (Chuvin et al. 2017).

Therefore, we hypothesized that misalignment of the clock within acinar cells was responsible for the

accelerated PanIN phenotype seen in CJ KC mice. In turn, we employed a cell-autonomous *Bmal1* knock-

out model to test whether circadian disruption in acinar cells facilitates tumor initiation and progression of

the pancreas cancer precursor lesions. In support of this hypothesis, previous studies in hepatocellular

carcinoma (HCC) have revealed a concordant phenotype between CJ and cell-autonomous clock

disruption (*Bmal1* knock-out) in mice predisposed to develop HCC (Kettner et al. 2016). We crossed the

KC mouse with a mouse harboring a conditional deletion of the core clock gene *Bmal1* to produce

Bmal1$^{fx/fx}$; LSL-Kras$^{G12D/+}$; Pdx1-Cre (BKC) mice, which possess *Bmal1* deletion in cells concomitantly

expressing *Pdx1* and the *Kras^{G12D}* mutation (Johnson et al. 2014). BMAL1 is a central transcriptional

regulator of the clock and is therefore essential for clock functionality (Bunger et al. 2000). We evaluated

the degree of FI/PanIN in BKC mice under standard lighting conditions at 5 months and, contrary to our

hypothesis, we found that cell-autonomous clock disruption caused resistance to FI/PanIN development

(**Figure 3A-D**). Compared to normal circadian KC mice, BKC mice had scant development of FI/PanIN

(12.73% ± 12.08 (KC) vs 1.59% ± 0.65 (BKC); $p < 0.01$), with essentially histologically normal pancreas,

and no mice exhibited a higher-grade lesion than PanIN-1 (**Figure 3A-D**). Additionally, this suppressed

PanIN development in BKC mice was not sex-dependent (male 1.64% ± 0.74 vs. female 1.52% ± 0.53; p

=0.65).

The observed results with cell-autonomous clock disruption were discordant with our expectations and

demonstrated that *Bmal1* knock-out in the targeted cells did not recapitulate the accelerated PanIN

phenotype seen with KC mice under (global) CJ conditions. *Pdx1* is expressed early during pancreatic

development and later to commit pancreatic progenitor cells to islets (Offield et al. 1996; Hingorani et al.

2003). To understand which cell populations in BKC mice exhibited loss of BMAL1 (location of *Pdx1*

expression), we crossed the KC mouse with the Ai14 'marker' mouse that conditionally expresses

tdTomato in the presence of *Pdx1*-driven Cre-recombinase (**Figure 3E-F**). This revealed tdTomato

expression (*Kras* mutation and *Bmal1* deletion) in acinar cells, ductal cells, PanINs, and islet cells.

Concomitantly, expression was absent in the fibroinflammatory infiltrate (fibroblasts, collagen, and

immune cells) and blood vessels surrounding the PanINs. Thus, in mice predisposed to pancreas

neoplasm formation, circadian disruption in acinar cells produced a different phenotype compared to CJ,

suggesting a different cell population may be responsible for the observed accelerated PanIN formation

and progression in CJ KC mice.

Single-cell RNA Sequencing of KC Mice Recapitulates the Highly Heterogeneous Tumor

Microenvironment

Prior work in the liver revealed that cell-autonomous knock-out of the clock in liver hepatocytes

(Albumin-Cre*; Bmal1*$^{\text{fx/fx}}$) drove a nearly identical phenotype compared to CJ mice (non-alcoholic fatty

liver disease progressing to HCC) (Kettner et al. 2016). However, we found that in the pancreas of KC

mice this was not evident. We expected that this was a consequence of non-cell autonomous changes to

the circadian clock (i.e. not acinar cells or PanINs) – whether by loss of clock function or by

misalignment – are responsible for accelerating the neoplastic progression. To examine further, we tested

the hypothesis that misalignment of the clock in an alternative cell population caused the accelerated

PanIN formation and progression; this was done by evaluating the expression of individual cells in the

tumor microenvironment over time. We performed single-cell RNA sequencing (scRNAseq) from pooled

pancreatic samples (1 male/female) from both normal circadian and CJ KC mice at 5 and 9 months

(**Figure 4A/B**). Notably, male and female mice were pooled because we consistently demonstrated no

sex-specific effect within either the normal circadian or the CJ group of KC mice at 5 months or 9

months. After filtering and dimensionality reduction, a total of 14 different heterogeneous cell types were

annotated using the ScType package comprising 42,211 cells (**Figure 4A-B & Supplemental Data 1-2**)

(Ianevski et al. 2022). These included acinar cells, several classes of immune cells, endothelial cells, and

fibroblast populations.

We first began by subsetting the pancreatic epithelial cells, which should encompass both normal acinar

and developing PanINs, and used pseudotime trajectory analysis to track cell-state changes that occurred

from 5 to 9 months. The expectation was that this population would gain signatures of neoplastic

transformation over time, given the loss of normal acinar cell density and increase in PanINs from 5 to 9

months. Pseudotime was set to zero at the point with the highest fraction of cells from the 5-month mice

and allowed to progress (**Figure 4C**). Moran's I autocorrelation analysis was then performed to obtain

genes that varied across the trajectory (**Supplemental Data 3**). Gene set enrichment analysis (GSEA) of

the genes that correlated with pseudotime progression revealed an association with multiple *Kras*-driven

cancer cell line signatures – corresponding precisely to the development of *Kras*-driven pancreatic

neoplasia over time (**Figure 4D**). Therefore, we identified a collection of pancreatic epithelial cells that

appeared to represent the transition from acinar cells to PanIN, coinciding with the increased PanIN

formation seen in histopathology from 5 to 9 months. These findings established a foundation for the

subsequent analysis of the clock to understand the effects of circadian misalignment (CJ) in the acinar cell

and PanIN population, which would then help to support or refute our new hypothesis of a non-cell-

autonomous cell population driving the accelerated PanIN phenotype.

Chronic Jetlag Promotes Differential Clock Gene Expression in Fibroblasts

We performed differential gene expression comparing the collective 5- and 9-month normal circadian and

CJ mouse cells, and assessed for differences in the CCG and clock-associated genes (including *Tef*, *Hlf*,

Bhlhe40, *Bhlhe41*, *Nfil3*, *Rorc*, *Dbp*, and *Ciart*) (**Supplemental Data 4**). We previously showed that CJ-

induced clock misalignment in the pancreas caused differential expression of the core clock genes

(Schwartz et al. 2021), and thus we expected populations of cells manifesting misalignment of the clock

between normal circadian and CJ conditions would similarly demonstrate CCG differential expression. In

support of our new hypothesis, the acinar cell/PanIN clusters that we identified and confirmed from the

pseudotime analysis did not exhibit differential expression of any of the CCGs or the clock-associated

genes between clusters derived from normal circadian or CJ KC mice. We then evaluated the other

clusters, and we found that endothelial cells, macrophages, memory CD4 T cells, and monocytes each had

a single clock-associated gene differentially expressed (**Figure 5A**). Interestingly, fibroblasts were the

only cell type that exhibited differential expression of CCGs and clock-associated genes as a result of

circadian misalignment, including *Per2*, *Per3*, *Nr1d2*, *Hlf*, *Tef*, and *Dbp* (**Figure 5A-B**). Fibroblasts are

an integral part of the PDAC microenvironment and are known to exhibit circadian rhythmicity when

compared to naïve fibroblasts (Parascandolo et al. 2020). The lack of other cell types exhibiting changes

in clock-specific gene expression suggests fibroblasts may be the target of the CJ-induced circadian

misalignment and accelerated neoplastic phenotype. Concordantly, fibroblasts were among the population

of cells that were negative for Pdx1 expression (in the KCT 'marker' mouse), corresponding to an intact

clock function in the BKC mice.

Fibroblasts Exhibit Time-Dependent Effects Secondary to Chronic Jetlag

To further evaluate the fibroblasts within our dataset, we isolated three different subclusters (**Figure 6A**).

Using existing markers generated for cancer-associated fibroblasts (CAFs), including inflammatory CAFs

(iCAFs), myofibroblast CAFs (mCAFs), and antigen-presenting CAFs (apCAFs), we assigned identities

with ScType (**Supplementary Data 5-6**) (Elyada et al. 2019). Of the three clusters, we identified all three

as iCAFs – with the expression of known markers, such as *Cle3b*, *Il6*, *Has1*, and *Col14a1* (**Figure 6B**).

Fibroblast cluster 3 exhibited some characteristics consistent with apCAFs, including expression of

multiple major histocompatibility complex (MHC) II components *H2-Aa*, *H2-Eb1*, and *H2-ab1* and *Cd74*,

the invariant chain of MHC II (Figure 6B). We then performed differential gene expression and GSEA

comparing the iCAFs between normal circadian and CJ groups at 5 and 9 months (**Figure 7A-B** &

Supplemental Data 7). We found that multiple metabolic pathways in fibroblasts were enriched in the CJ

group at 5 months (**Figure 7A**), and our 9-month comparison revealed enrichment of several apoptosis-

related pathways, including 'NEGATIVE_REGULATION_OF_CELL_DEATH' and

'REGULATION_OF_APOPTOTIC_SIGNALING_PATHWAY' (**Figure 7B**). To evaluate if CJ was

simply increasing fibrosis (i.e. collagen deposition) to facilitate these changes, we compared the collagen

content from 5- and 9-month normal circadian and CJ KC mice (**Supplemental Data 8**). We found no

differences in the collagen content at either time point. We, therefore, identified that the iCAFs in the

developing tumor microenvironment of KC mice had early enrichment of metabolic pathways in response

to CJ, with late enrichment of apoptosis-related pathways. Enrichment of these pathways in fibroblasts as

a consequence of CJ corresponds with the finding that fibroblasts manifest differential expression of core

clock genes (i.e. affected by misalignment of the clock). When combined with the observation that acinar

cells/PanINs demonstrated no differential gene expression of CCGs, and abolishing the clock in acinar

cells did not replicate the CJ-induced phenotype, these results suggest CJ causes clock desynchrony in the

fibroblast population to alter signaling and accelerate the formation and progression of PanINs.

Discussion:

There has been a substantial volume of literature examining circadian desynchrony, which encompasses

the misalignment between environmental or behavioral cycles and the endogenous clock (Vetter 2020).

Persistent misalignment between environmental cues (light/dark cycle) and behavioral cues (feeding)

compared to the internal clocks can result in aberrant gene signaling (Kettner et al. 2016; Sulli et al. 2018)

with consequent disease pathology (Scheer et al. 2009; Lee et al. 2013; Morris et al. 2015). For instance,

in healthy human adults subjected to a forced circadian desynchrony protocol that replicated shift work

(12-hour reversal of light/dark cycle for 3 days), there was a misalignment of the circadian clock relative

to the timing of the light/dark cycle causing glucose intolerance as well as reduced insulin sensitivity –

findings indicative of impaired glucose metabolism that occurs with the onset of type II diabetes (Morris

et al. 2015). This coincides with epidemiological findings of an increased risk of type II diabetes with

shift work (Pan et al. 2011) and is congruent with the increased risk of obesity and metabolic syndrome in

individuals performing rotating shift work schedules (Lin et al. 2009; Antunes et al. 2010). Unfortunately,

the metabolic reprogramming and chronic inflammation that ensues following these diagnoses can drive

the formation of several cancers such as hepatocellular carcinoma (Campbell et al. 2016), which

commences a path of 'neoplastic change' consisting of non-alcoholic fatty liver disease, non-alcoholic

steatohepatitis, and liver cancer (Simon et al. 2021). The same process of neoplastic transformation in the

liver can occur due to circadian misalignment: C57Bl/6J mice subjected to chronic jetlag conditions

experienced an acceleration of non-alcoholic fatty liver disease and an increased incidence of

hepatocellular carcinoma compared to mice under standard lighting conditions (Kettner et al. 2016).

Notably, the evidence correlating circadian desynchrony and liver cancer is mirrored in a similar

metabolic organ – the pancreas. There is a strong correlation between obesity and diabetes, and risk of

developing pancreatic ductal adenocarcinoma (PDAC). A potential link between circadian misalignment

(associated with obesity and diabetes) and PDAC can be inferred from this correlation, and this putative

link is strengthened by epidemiologic data reporting an increased risk of PDAC in shift workers (Parent et

al. 2012). Therefore, based on this cumulative evidence, we sought to test the hypothesis that circadian

clock desynchrony through chronic jetlag (CJ) would accelerate the initiation and progression of PanINs

in mice predisposed to develop pre-cancerous lesions (PanIN-1,-2,-3) and PDAC.

Our work revealed that circadian misalignment through CJ increased tumor initiation and progression of

PanINs, which are pancreas cancer pre-cursor lesions that can ultimately develop into PDAC. Our

findings of accelerated neoplasm progression in the current study are alarming considering the highly

lethal nature of PDAC (Siegel et al. 2021), and the prevalence of circadian desynchrony in society

(Antunes et al. 2010). This type of clock dysfunction affects a significant percentage of the population,

such as shift workers under imposed non-daytime schedules, who experience chronic misalignment

between their endogenous clock and environmental cues (Bass and Lazar 2016). A less severe but highly

prevalent form of misalignment may also be present through extended exposure to light at night, extended

duration of dietary intake, or significant alteration in sleep patterns (social jetlag) (Roenneberg et al.

2012; Chang et al. 2015; Bass and Lazar 2016; Fishbein et al. 2021). To study the potential implications

of these types of clock desynchrony in the development of PDAC, we utilized a mouse model in which

Kras-mutation in pancreatic acinar cells promotes acinar-to-ductal metaplasia, followed by PanIN

formation, and ultimately PDAC. Consequently, a pertinent question for the present work is the

applicability of the model to humans. Interestingly, in humans, PanINs are frequently found in the

pancreas and their prevalence increases with age (Matsuda et al. 2017). They are a result of acinar

metaplastic transformation to a ductal cell phenotype, typically in response to injury or inflammation (De

La O et al. 2008; Reichert and Rustgi 2011; Chuvin et al. 2017). In an autopsy study of 173 cases with no

evidence of PDAC (mean age 80.5 years), PanIN-1 was present in 77% of pancreas specimens, PanIN-2

was present in 28%, and 4% of the pancreas specimens harbored PanIN-3 (Matsuda et al. 2017).

Consistent with our KC model, PanIN-1s were always present in the pancreas when PanIN-2 or -3 were

identified. These autopsy findings were also confirmed in a separate study (Longnecker and Suriawinata

2022). Thus, our model is concordant with the neoplastic process that occurs in humans and relevant for

studying PanIN progression. Our findings of CJ-induced acceleration of advanced PanIN formation could

be applicable to many individuals who unknowingly harbor PanINs and experience consistent circadian

misalignment.

PanINs are microscopic and early PanINs are unable to be detected with the current imaging modalities

(Kanda et al. 2012). Therefore, human studies of circadian misalignment driving PanIN progression have

not been feasible. Yet, the *in vivo* findings in our study warrant consideration of future investigations

designed to evaluate risk in humans and possibly risk mitigation. Given the widespread presence of

circadian misalignment, and the prevalence of PanINs, a suitable population of individuals to target for

risk mitigation of circadian desynchrony may prove tenable. For example, there is convincing evidence

that patients who have a family history of PDAC demonstrate a greater propensity for PanIN formation,

are more likely to harbor advanced PanINs, and, therefore, are at higher risk for developing PDAC. In a

study by Shi and colleagues evaluating patients who underwent pancreatic resection for PDAC, those

with a family history of PDAC (without a germline mutation) demonstrated a rate of 1.51 PanIN

lesions/cm^2 in the pancreas (i.e. non-cancer pancreas) compared to 0.55 PanIN lesion/cm^2 in patients with

sporadic PDAC (Shi et al. 2009). The majority of the lesions were PanIN-1, with fewer PanIN-2 and

PanIN-3 lesions identified (Shi et al. 2009); again mirroring the findings in our KC model where there is

multifocal lesion formation, and the majority of the lesions at 9 months were PanIN-1 with fewer PanIN-2

and PanIN-3. Moreover, in the study by Shi *et al* (Shi et al. 2009), PanIN lesions were found in the

pancreas remote from invasive cancer, which supports the findings from autopsy studies (Matsuda et al.

2017; Longnecker and Suriawinata 2022) that not all PanINs progress to cancer. Similar to the etiology of

PanINs in our model, greater than 95% of human PanINs harbor a *Kras* mutation (including 92% of

PanIN-1) (Kanda et al. 2012), and it is generally accepted that the fraction of *Kras*-mutant cells in PanINs

increases with PanIN grade, indicating clonal expansion of the *Kras*-mutant cells that have been

'transformed' (Kanda et al. 2012). However, the drivers of this transformation from PanIN-1 to higher-

grade PanINs or PDAC are poorly understood. Consequently, our study forms a strong basis for further

investigation to understand how circadian misalignment promotes the progression of PanIN lesions. This

should lead to prevention strategies (e.g. chemoprevention or behavioral modification) that can then be

applied to high-risk patients to prevent the progression of PanINs to cancer (Miller et al. 2016).

Incidentally, these high-risk patients are already being captured in high-risk clinics, where individuals

with a strong family history or known pre-cursor lesions in the pancreas are being followed to identify

cancer transformation at an early stage (Shin et al. 2015; Miller et al. 2016; Dbouk et al. 2022). This

would be an ideal group to target for risk mitigation or chemoprevention.

To better understand the basis for our observations of accelerated neoplasm formation with circadian

misalignment (CJ), we implemented a conditional mouse model to achieve cell-autonomous core clock

gene knock-out. We chose a conditional knock-out as opposed to a global knock-out because of the

known difficulty in disentangling the effects in a global knock-out model (e.g. *Bmal1*$^{-/-}$). For example, the

opposing effects in glucose metabolism and insulin sensitivity with knock-out of the clock in the pancreas

and liver can mask glucose intolerance phenotypes that are otherwise readily apparent with islet-cell-

specific clock disruption (Lamia et al. 2008; Marcheva et al. 2010; Perelis et al. 2016). Furthermore,

chronic jetlag exposure is a 'global' effect and we aimed to home in on the cell population responsible for

the observed phenotype. In this way, the use of core clock gene knock-out models has been indispensable

in ascertaining clock-dependent mechanisms for various pathologies. Oftentimes, the phenotypes among

circadian disrupted groups are concordant, whether by circadian desynchrony or through genetic

alteration. For instance, Lee and colleagues found that the cell-autonomous deletion of *Bmal1* in β-cells of

the pancreas resulted in the loss of glucose-stimulated insulin secretion, and the blunted insulin secretion

with peripheral insulin resistance was replicated by environmentally-induced circadian misalignment (i.e.

6-hr light phase advancement protocol) (Lee et al. 2013). In a separate study, mice predisposed to

colorectal cancer development were subjected to circadian misalignment via constant lighting conditions

and showed a 2-fold increase in tumor initiation compared to those housed under diurnal conditions

(Stokes et al. 2021). In this study, Stokes *et al*. showed that global and epithelial loss of *Bmal1* similarly

induced tumor initiation, suggesting that *Bmal1* expression in the colon epithelial cells (cell of origin for

colorectal cancer) is similarly 'targeted' by perturbations in the clock (altered lighting) and by clock

disruption (*Bmal1* deletion). Comparable results were seen by Kettner *et al*. when evaluating rates of non-

alcoholic fatty liver disease and hepatocellular carcinoma between cell-autonomous *Bmal1* knock-out

mice and mice subjected to circadian desynchrony (Kettner et al. 2016). Thus, we expected that cell-

autonomous clock disruption in KC mice (pancreatic endocrine and exocrine *Bmal1* knock-out) would

demonstrate a PanIN phenotype concordant with the circadian misaligned KC mice. However, when we

crossed our KC mice with the condition *Bmal1^{fx/fx}* mouse, we found a discordant phenotype in pancreatic

tumor initiation compared to circadian misalignment. Although the results were unanticipated, they did

provide insight and enabled us to conclude that disruption of the clock in acinar cells does not facilitate

PanIN formation or progression. This led us to investigate separate putatively involved cell populations.

To better understand individual cell population involvement, and to build a foundation for discerning how

pancreatic neoplasia more rapidly progressed in CJ conditions, we performed scRNAseq at both 5 and 9

months. We found that the pancreatic epithelial cells, including the PanINs and acinar cells, showed a

time-dependent cancer signature on pseudotime analysis consistent with the observed histopathologic

changes. Confirming the time-dependent cancer signature enabled us to examine these cells primarily for

changes to the clock, to determine if the scRNAseq data from circadian misaligned mice complemented

the *Bmal1* cell-autonomous knock-out data. Subsequently, we assessed 13 other cell types for evidence of

differential gene expression between normal circadian and CJ, and we found that fibroblasts –not the

pancreatic epithelial cells – were the only cell type with multiple CCG and clock-associated gene

changes. This finding supports the notion that CJ augments fibroblasts in the developing tumor

microenvironment and not necessarily the epithelial/PanINs themselves. Notably, this also reinforces the

differences identified in the cell-autonomous *Bmal1* knock-out model versus the circadian desynchrony

model – misalignment may be targeting the fibroblast population to accelerate neoplasm formation and

progression. Meanwhile, in the *LSL-Bmal1$^{fx/fx}$; Pdx1-Cre; LSL-Kras$^{G12D/+}$* mice, *Bmal1* (and thus the

clock) is intact in the mesenchymal cells: IHC in KC mice conditionally expressing tdTomato

demonstrated that there was *Pdx1*-Cre-directed expression in the acinar, ductal, islet, and PanINs, but not

the mesenchymal lineage cells, including the fibroblasts and endothelial cells. We used additional

expression markers to subset the fibroblasts and we identified all 3 clusters as inflammatory CAFs

(iCAFs) based on established markers (Elyada et al. 2019). iCAFs secrete IL-6, CXCL9, and TGFβ and

can be both immunosuppressive as well as immune-promoting (Sahai et al. 2020). Interestingly, we did

not observe any differences in the amount of pancreatic collagen content at 5 or 9 months between normal

circadian and CJ KC mice, suggesting a role for paracrine signaling in mediating the effects of CJ by

fibroblasts rather than extracellular matrix production (i.e. desmoplasia). While this requires additional

consequential investigation and will be the focus of future work, it is important to note that other

investigators have uncovered the substantive involvement of the fibroblast clock in cancer. In a pan-

cancer analysis of The Cancer Genome Atlas tumor samples, Wu and colleagues employed a new

computational approach (LTM) to explore clock-controlled pathways in cancer (Wu et al. 2022). The

authors found that the variation in clock strength across 9 separate cancers was directly related to the

proportion of fibroblasts in the samples and concluded that this was consequent to fibroblasts possessing a

more robust circadian clock than the adjacent cancer cells within the tumor microenvironment. Moreover,

fibroblasts are known to have an essential role in fostering cancer progression in the pancreas cancer

microenvironment (Hanahan and Coussens 2012; Helms et al. 2020). As fibroblast density increases with

the advancement of PanINs (i.e. PanIN-1 to PanIN-2/3) towards invasive cancer, the cancer-associated

fibroblasts (CAFs) display a 'wound-healing' type response that supports neoplastic proliferation through

direct and indirect metabolic contributions (Helms et al. 2020). CAFs display differences in rhythmic

gene expression compared to naïve fibroblasts (Parascandolo et al. 2020), and the wound healing type

response in fibroblasts has been previously shown to be strongly clock dependent (Hoyle et al. 2017),

with disruption of the clock profoundly affecting this pathway. Concordantly, after identifying and

characterizing the CAFs in our mouse pancreas samples, we performed GSEA at 5 and 9 months. This

demonstrated that at 5 months CJ promoted metabolic gene enrichment in the CAFs, and by 9 months,

there was enrichment of multiple cell death and apoptosis-related pathways. It is therefore conceivable

that the fibroblasts – which harbor a strong clock and coevolve with PanINs – underlie the circadian

desynchrony-induced acceleration of PanIN formation and progression. To investigate further whether

circadian disruption in the fibroblasts may recapitulate the circadian misalignment (CJ) phenotype, in

future work we aim to generate *Col1a1-Cre; Bmal1$^{fx/fx}$* in FLP recombinase *Kras*-mutant mice (*Pdx1-*

FlpO; *FSF-Kras$^{G12D/+}$*) (Wu et al. 2017). This would be effective in abolishing the fibroblast clock in the

developing tumor microenvironment because all the identified fibroblast populations expressed high

levels of *Col1a1* (scRNAseq).

A limitation to consider from the scRNAseq data was that we did not identify islet cell clusters and were

therefore unable to discern whether differential expression of CCGs was present in this population of

cells. However, Pdx1 is expressed in endocrine cells in the pancreas, and therefore the BKC mice (which

did not display an accelerated PanIN phenotype) would have a disrupted clock in this cell population

(Offield et al. 1996; Hingorani et al. 2003). While clock disruption via *Bmal1* knock-out in islet cells

produces metabolic changes that could ostensibly alter the formation or growth trajectory of tumors

(PanINs), the prevailing phenotype with BMAL1 deficiency in islet cells – and also with circadian

misalignment – is glucose intolerance, hypoinsulinemia, and hyperglycemia (i.e. characteristics of

diabetes) (Marcheva et al. 2010; Lee et al. 2013; Perelis et al. 2016). Given that diabetes is a risk factor

for PDAC development (Abbruzzese et al. 2018), and the lack of phenotype (expedited PanIN formation)

in the BKC mice, it seems unlikely that circadian misalignment in the islets would have contributed to

accelerated PanIN formation and progression. Regardless, we were not able to evaluate this cell

population in the scRNAseq analysis, and so this remains a limitation. To investigate further, one could

consider beta-cell specific *Bmal1* knock-out (*Rip-Cre; Bmal1$^{fx/fx}$*) in FLP recombinase *Kras*-mutant mice

(*Pdx1-FlpO*; *FSF-Kras$^{G12D/+}$*) to determine if the accelerated PanIN development emerges (Wu et al.

2017). Another consideration is the intriguing observation that loss of pancreatic *Bmal1* (acinar cells,

ductal cells, islets, PanINs) resulted in near abolishment of PanIN formation, and not just an absence of

accelerated PanIN formation. This finding speaks to the complex interplay of the circadian clock amongst

cell populations in the tumor microenvironment. In future work, we aim to employ various genetically

engineered mouse models to disentangle these findings, which should have clinical relevance for the

multitude of individuals that harbor PanINs while concomitantly experiencing chronic circadian

misalignment.

Although previous evidence suggests a role for the circadian clock in PDAC growth and spread, this is the

first study to show perturbations of the clock increases tumor initiation in PDAC (Jiang et al. 2016; Jiang

et al. 2018). Our study provides a foundation for future work to expand our understanding of how a

dyssynchronous clock orchestrates cancer development in the pancreas.

Figure Legends:

Figure 1: *Pancreatic fibroinflammation/PanIN development is increased by chronic jetlag at 5 months.*

A. Boxplot comparing the mean (with associated 25th and 75th quantiles) percent pancreatic involvement of FI/PanIN at 5 months for normal circadian KC (n= 36) and CJ KC mice (n = 26). **B.** Representative H&E images of normal circadian (upper) and chronic jetlag (lower) mice at 40x (left) and 100x (right). Scale bars are shown for reference. [Annotation: PanIN (P), acinar (A) cells, islet (I) cells, ductal (D) cells, vascular (V) cells (i.e. endothelial cells), and fibroinflammatory (F) infiltrate]

Figure 2: *Chronic jetlag accelerates the grade of PanIN lesions.* Bar charts indicating the incidence of PanIN-1-3 and PDAC in normal circadian (black) and CJ (grey) conditions at **A.** 5 months (normal circadian n = 36 vs. CJ n = 26) and **B.** 9 months (normal circadian n = 33 vs. CJ n = 39). **C.** Table showing the Fischer's Exact Test comparisons between 5- and 9-month normal circadian and CJ mice. **D.** Representative 200x H&E image from a CJ mouse demonstrating PanIN-1 (arrow) and PanIN-2 lesion (arrowhead). **E.** Representative 200x H&E image from a CJ mouse demonstrating a PanIN-3 lesion (arrowhead) and PDAC off the same gland (dashed circle). Scale bars are shown for reference.

Figure 3: *Pancreatic cell-autonomous deletion of Bmal1 does not accelerate pancreatic fibroinflammatory infiltrate/PanIN development at 5 months*. **A.** Boxplot comparing the mean (with associated 25th and 75th quantiles) percent pancreatic involvement of FI/PanIN at 5 months for normal circadian KC (n= 36) and BKC mice (n = 24). Representative H&E images demonstrating the FI/PanIN from a BKC mouse at 40x (**B**), 100x (**C**), and 200x (**D**) To evaluate the cell types affected in BKC, KC mice were crossed with Ai14 reporter mice to create KCT mice. H&E stained sections (**E**) and immunohistochemistry for tdTomato (**F**) were compared to evaluate cell-specific Pdx1-driven Cre expression in KC mice. Scale bars are shown for reference. [annotation: PanIN (P), acinar (A) cells, islet (I) cells, ductal (D) cells, vascular (V) cells (i.e. endothelial cells), and fibroinflammatory (F) infiltrate]

Figure 4: *The single-cell landscape of KC mice recapitulates the developing pancreatic cancer tumor microenvironment.* **A.** UMAP demonstrating the 14 diverse cell types and 42,211 cells in the scRNAseq dataset. **B.** UMAPs split by time (5 and 9 months) and condition (normal circadian and CJ) **C.**

Pseudotime trajectory analysis was performed for the subset of pancreatic acinar cells with 0 pseudotime

set to the point with the highest fraction of cells at 5 months. **D.** Gene Set Enrichment Analysis (GSEA)

was performed for the oncogenic signature gene set associated with pseudotime progression and ordered

by the log adjusted p value.

Figure 5: *Fibroblasts were preferentially affected by chronic jetlag.* **A.** Table of core clock genes (CCGs)

and clock-associated genes differentially expressed when comparing normal circadian and CJ conditions

of each cell type. **B.** Representative UMAP dimensionality plots demonstrating cellular expression (blue

color) of select CCG and clock-associated genes in the subset of fibroblasts.

Figure 6: *Fibroblasts subclusters in the developing pancreatic tumor microenvironment were identified*

as inflammatory cancer-associated fibroblasts. **A.** A UMAP demonstrating the three fibroblast

subclusters identified on dimensionality reduction. **B.** Violin plots with select markers for Pan-,

Inflammatory-, Myofibroblastic-, and Antigen Presenting-Cancer Associated Fibroblasts (CAFs).

Figure 7: *Chronic jetlag influences fibroblast expression patterns across time.* Figures with The Gene Set

Enrichment Analysis (GSEA) demonstrating the gene ontology biologic process (GO BP) pathways

enriched in CJ compared to normal circadian fibroblasts at **A.** five and **B.** nine months. The top 10

pathways at each time point are ordered by their log-adjusted p value.

<u>Supplementary Data:</u>

<u>Supplemental Data 1</u>: Marker database for single-cell RNA sequencing cell-type identification

<u>Supplemental Data 2</u>: ScType Scores for overall single-cell RNA sequencing dataset

<u>Supplemental Data 3</u>: Moran's I pseudotime analysis for the pancreatic acinar cells

<u>Supplemental Data 4</u>: Differential gene expression between normal circadian and chronic jetlag

conditions for all cell types

<u>Supplemental Data 5</u>: Marker database for fibroblast subtype identification

<u>Supplemental Data 6</u>: ScType Scores for fibroblast subtype identification

<u>Supplemental Data 7</u>: Differential gene expression between normal circadian and chronic jetlag

conditions for fibroblasts at 5 and 9 months

<u>Supplemental Data 8</u>: Quantification of collagen content in 5-month normal circadian and chronic jetlag

KC mice

Acknowledgments:

We would like to take the opportunity to acknowledge the University of Wisconsin Carbone Cancer

Center Support Grant P30 CA014520, which provides funding for the Experimental Animal Pathology

Lab Core.

We would also to acknowledge the Gene Expression Center and the Bioinformatics Resource Center

(BRC) at the University of Wisconsin, Madison for their contributions to this work.

Finally, we would like to thank the Michael W. Oglesby Foundation for their funding support of our work

in circadian disruption and pancreas pathology.

Grants:

The research reported in this publication was supported by the Department of Defense Peer Reviewed

Cancer Research Program Award CA190176 (Grants.gov ID GRANT12935023) and the University of

Wisconsin Institute for Clinical and Translational Research KL2 Award (UL1TR002373 and

KL2TR002374) [SRK]. Funding support for this work was also provided by the National Institutes of

Health under Award Number T32 ES007015 [PBS]. The content is solely the responsibility of the authors

and does not necessarily represent the official views of the National Institutes of Health.

Data Statement:

Single-cell RNA sequencing data is made publicly available through gene expression omnibus (GEO)

(Accession number: GSE209629). All other data that support the findings of this study are available from

the corresponding author, SRK, upon reasonable request.

<u>References:</u>

Abbruzzese JL, Andersen DK, Borrebaeck CAK, Chari ST, Costello E, Cruz-Monserrate Z, Eibl G,

Engleman EG, Fisher WE, Habtezion A, et al. 2018. The Interface of Pancreatic Cancer With Diabetes,

Obesity, and Inflammation: Research Gaps and Opportunities: Summary of a National Institute of

Diabetes and Digestive and Kidney Diseases Workshop. Pancreas. 47(5):516–525.

https://doi.org/10.1097/MPA.0000000000001037

Antunes LC, Levandovski R, Dantas G, Caumo W, Hidalgo MP. 2010. Obesity and shift work:

chronobiological aspects. Nutr Res Rev. 23(1):155–168. https://doi.org/10.1017/S0954422410000016

Arble DM, Bass J, Laposky AD, Vitaterna MH, Turek FW. 2009. Circadian timing of food intake

contributes to weight gain. Obesity (Silver Spring). 17(11):2100–2102.

https://doi.org/10.1038/oby.2009.264

Bass J, Lazar MA. 2016. Circadian time signatures of fitness and disease. Science. 354(6315):994–999.

https://doi.org/10.1126/science.aah4965

Bass J, Takahashi JS. 2010. Circadian integration of metabolism and energetics. Science.

330(6009):1349–1354. https://doi.org/10.1126/science.1195027

Bunger MK, Wilsbacher LD, Moran SM, Clendenin C, Radcliffe LA, Hogenesch JB, Simon MC,

Takahashi JS, Bradfield CA. 2000. Mop3 is an essential component of the master circadian pacemaker in

mammals. Cell. 103(7):1009–1017. https://doi.org/10.1016/s0092-8674(00)00205-1

Campbell PT, Newton CC, Freedman ND, Koshiol J, Alavanja MC, Beane Freeman LE, Buring JE, Chan

AT, Chong DQ, Datta M, et al. 2016. Body Mass Index, Waist Circumference, Diabetes, and Risk of

Liver Cancer for U.S. Adults. Cancer Res. 76(20):6076–6083. https://doi.org/10.1158/0008-5472.CAN-

16-0787

Chang A-M, Aeschbach D, Duffy JF, Czeisler CA. 2015. Evening use of light-emitting eReaders

negatively affects sleep, circadian timing, and next-morning alertness. Proc Natl Acad Sci U S A.

112(4):1232–1237. https://doi.org/10.1073/pnas.1418490112

Choudhary S, Satija R. 2022. Comparison and evaluation of statistical error models for scRNA-seq.

Genome Biology. 23(1):27. https://doi.org/10.1186/s13059-021-02584-9

Chuvin N, Vincent DF, Pommier RM, Alcaraz LB, Gout J, Caligaris C, Yacoub K, Cardot V, Roger E,

Kaniewski B, et al. 2017. Acinar-to-Ductal Metaplasia Induced by Transforming Growth Factor Beta

Facilitates KRASG12D-driven Pancreatic Tumorigenesis. Cell Mol Gastroenterol Hepatol. 4(2):263–282.

https://doi.org/10.1016/j.jcmgh.2017.05.005

Cox KH, Takahashi JS. 2019. Circadian clock genes and the transcriptional architecture of the clock

mechanism. J Mol Endocrinol. 63(4):R93–R102. https://doi.org/10.1530/JME-19-0153

Dbouk M, Katona BW, Brand RE, Chak A, Syngal S, Farrell JJ, Kastrinos F, Stoffel EM, Blackford AL,

Rustgi AK, et al. 2022. The Multicenter Cancer of Pancreas Screening Study: Impact on Stage and

Survival. J Clin Oncol. 40(28):3257–3266. https://doi.org/10.1200/JCO.22.00298

De La O J-P, Emerson LL, Goodman JL, Froebe SC, Illum BE, Curtis AB, Murtaugh LC. 2008. Notch

and Kras reprogram pancreatic acinar cells to ductal intraepithelial neoplasia. Proc Natl Acad Sci U S A.

105(48):18907–18912. https://doi.org/10.1073/pnas.0810111105

Elyada E, Bolisetty M, Laise P, Flynn WF, Courtois ET, Burkhart RA, Teinor JA, Belleau P, Biffi G,

Lucito MS, et al. 2019. Cross-Species Single-Cell Analysis of Pancreatic Ductal Adenocarcinoma

Reveals Antigen-Presenting Cancer-Associated Fibroblasts. Cancer Discov. 9(8):1102–1123.

https://doi.org/10.1158/2159-8290.CD-19-0094

Filipski E, Delaunay F, King VM, Wu M-W, Claustrat B, Gréchez-Cassiau A, Guettier C, Hastings MH,

Francis L. 2004. Effects of chronic jet lag on tumor progression in mice. Cancer Res. 64(21):7879–7885.

https://doi.org/10.1158/0008-5472.CAN-04-0674

Filipski E, Subramanian P, Carrière J, Guettier C, Barbason H, Lévi F. 2009. Circadian disruption

accelerates liver carcinogenesis in mice. Mutat Res. 680(1–2):95–105.

https://doi.org/10.1016/j.mrgentox.2009.10.002

Finak G, McDavid A, Yajima M, Deng J, Gersuk V, Shalek AK, Slichter CK, Miller HW, McElrath MJ,

Prlic M, et al. 2015. MAST: a flexible statistical framework for assessing transcriptional changes and

characterizing heterogeneity in single-cell RNA sequencing data. Genome Biol [Internet]. [accessed 2021

Feb 4] 16. https://doi.org/10.1186/s13059-015-0844-5

Fishbein AB, Knutson KL, Zee PC. 2021. Circadian disruption and human health. J Clin Invest.

131(19):e148286. https://doi.org/10.1172/JCI148286

Hanahan D, Coussens LM. 2012. Accessories to the crime: functions of cells recruited to the tumor

microenvironment. Cancer Cell. 21(3):309–322. https://doi.org/10.1016/j.ccr.2012.02.022

Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, Lee MJ, Wilk AJ, Darby C, Zager

M, et al. 2021. Integrated analysis of multimodal single-cell data. Cell. 184(13):3573-3587.e29.

https://doi.org/10.1016/j.cell.2021.04.048

Helms E, Onate MK, Sherman MH. 2020. Fibroblast Heterogeneity in the Pancreatic Tumor

Microenvironment. Cancer Discov. 10(5):648–656. https://doi.org/10.1158/2159-8290.CD-19-1353

Hingorani SR, Petricoin EF, Maitra A, Rajapakse V, King C, Jacobetz MA, Ross S, Conrads TP,

Veenstra TD, Hitt BA, et al. 2003. Preinvasive and invasive ductal pancreatic cancer and its early

detection in the mouse. Cancer Cell. 4(6):437–450. https://doi.org/10.1016/s1535-6108(03)00309-x

Hoyle NP, Seinkmane E, Putker M, Feeney KA, Krogager TP, Chesham JE, Bray LK, Thomas JM, Dunn

K, Blaikley J, O'Neill JS. 2017. Circadian actin dynamics drive rhythmic fibroblast mobilization during

wound healing. Sci Transl Med. 9(415):eaal2774. https://doi.org/10.1126/scitranslmed.aal2774

Ianevski A, Giri AK, Aittokallio T. 2022. Fully-automated and ultra-fast cell-type identification using

specific marker combinations from single-cell transcriptomic data. Nat Commun. 13(1):1246.

https://doi.org/10.1038/s41467-022-28803-w

Jiang W, Zhao S, Jiang X, Zhang E, Hu G, Hu B, Zheng P, Xiao J, Lu Z, Lu Y, et al. 2016. The circadian

clock gene Bmal1 acts as a potential anti-oncogene in pancreatic cancer by activating the p53 tumor

suppressor pathway. Cancer Lett. 371(2):314–325. https://doi.org/10.1016/j.canlet.2015.12.002

Jiang W, Zhao S, Shen J, Guo L, Sun Y, Zhu Y, Ma Z, Zhang X, Hu Y, Xiao W, et al. 2018. The MiR-

135b-BMAL1-YY1 loop disturbs pancreatic clockwork to promote tumourigenesis and chemoresistance.

Cell Death Dis. 9(2):149. https://doi.org/10.1038/s41419-017-0233-y

Johnson BP, Walisser JA, Liu Y, Shen AL, McDearmon EL, Moran SM, McIntosh BE, Vollrath AL,

Schook AC, Takahashi JS, Bradfield CA. 2014. Hepatocyte circadian clock controls acetaminophen

bioactivation through NADPH-cytochrome P450 oxidoreductase. Proc Natl Acad Sci U S A.

111(52):18757–18762. https://doi.org/10.1073/pnas.1421708111

Kanda M, Matthaei H, Wu J, Hong S-M, Yu J, Borges M, Hruban RH, Maitra A, Kinzler K, Vogelstein

B, Goggins M. 2012. Presence of somatic mutations in most early-stage pancreatic intraepithelial

neoplasia. Gastroenterology. 142(4):730-733.e9. https://doi.org/10.1053/j.gastro.2011.12.042

Kettner NM, Voicu H, Finegold MJ, Coarfa C, Sreekumar A, Putluri N, Katchy CA, Lee C, Moore DD,

Fu L. 2016. Circadian Homeostasis of Liver Metabolism Suppresses Hepatocarcinogenesis. Cancer Cell.

30(6):909–924. https://doi.org/10.1016/j.ccell.2016.10.007

Lamia KA, Storch K-F, Weitz CJ. 2008. Physiological significance of a peripheral tissue circadian clock.

Proc Natl Acad Sci U S A. 105(39):15172–15177. https://doi.org/10.1073/pnas.0806717105

Lee C, Etchegaray JP, Cagampang FR, Loudon AS, Reppert SM. 2001. Posttranslational mechanisms

regulate the mammalian circadian clock. Cell. 107(7):855–867. https://doi.org/10.1016/s0092-

8674(01)00610-9

Lee J, Moulik M, Fang Z, Saha P, Zou F, Xu Y, Nelson DL, Ma K, Moore DD, Yechoor VK. 2013.

Bmal1 and β-cell clock are required for adaptation to circadian disruption, and their loss of function leads

to oxidative stress-induced β-cell failure in mice. Mol Cell Biol. 33(11):2327–2338.

https://doi.org/10.1128/MCB.01421-12

Lee JH, Sancar A. 2011. Regulation of apoptosis by the circadian clock through NF-kappaB signaling.

Proc Natl Acad Sci U S A. 108(29):12036–12041. https://doi.org/10.1073/pnas.1108125108

Li S, Xie K. 2022. Ductal metaplasia in pancreas. Biochim Biophys Acta Rev Cancer. 1877(2):188698.

https://doi.org/10.1016/j.bbcan.2022.188698

Lin Y-C, Hsiao T-J, Chen P-C. 2009. Persistent rotating shift-work exposure accelerates development of

metabolic syndrome among middle-aged female employees: a five-year follow-up. Chronobiol Int.

26(4):740–755. https://doi.org/10.1080/07420520902929029

Liu Y, Johnson BP, Shen AL, Wallisser JA, Krentz KJ, Moran SM, Sullivan R, Glover E, Parlow AF,

Drinkwater NR, et al. 2014. Loss of BMAL1 in ovarian steroidogenic cells results in implantation failure

in female mice. PNAS. 111(39):14295–14300. https://doi.org/10.1073/pnas.1209249111

Longnecker DS, Suriawinata AA. 2022. Incidence of Pancreatic Intraepithelial Neoplasia in an Autopsy

Series. Pancreas. 51(4):305–309. https://doi.org/10.1097/MPA.0000000000002027

Marcheva B, Ramsey KM, Buhr ED, Kobayashi Y, Su H, Ko CH, Ivanova G, Omura C, Mo S, Vitaterna

MH, et al. 2010. Disruption of the clock components CLOCK and BMAL1 leads to hypoinsulinaemia and

diabetes. Nature. 466(7306):627–631. https://doi.org/10.1038/nature09253

Matsuda Y, Furukawa T, Yachida S, Nishimura M, Seki A, Nonaka K, Aida J, Takubo K, Ishiwata T,

Kimura W, et al. 2017. The Prevalence and Clinicopathological Characteristics of High-Grade Pancreatic

Intraepithelial Neoplasia: Autopsy Study Evaluating the Entire Pancreatic Parenchyma. Pancreas.

46(5):658–664. https://doi.org/10.1097/MPA.0000000000000786

Matsuo T, Yamaguchi S, Mitsui S, Emi A, Shimoda F, Okamura H. 2003. Control mechanism of the

circadian clock for timing of cell division in vivo. Science. 302(5643):255–259.

https://doi.org/10.1126/science.1086271

McInnes L, Healy J, Melville J. 2020. UMAP: Uniform Manifold Approximation and Projection for

Dimension Reduction. arXiv:180203426 [cs, stat] [Internet]. [accessed 2021 Feb 4].

http://arxiv.org/abs/1802.03426

McIntosh BE, Hogenesch JB, Bradfield CA. 2010. Mammalian Per-Arnt-Sim proteins in environmental

adaptation. Annu Rev Physiol. 72:625–645. https://doi.org/10.1146/annurev-physiol-021909-135922

Miller MS, Allen P, Brentnall TA, Goggins M, Hruban RH, Petersen GM, Rao CV, Whitcomb DC, Brand

RE, Chari ST, et al. 2016. Pancreatic Cancer Chemoprevention Translational Workshop: Meeting Report.

Pancreas. 45(8):1080–1091. https://doi.org/10.1097/MPA.0000000000000705

Morris CJ, Yang JN, Garcia JI, Myers S, Bozzi I, Wang W, Buxton OM, Shea SA, Scheer FAJL. 2015.

Endogenous circadian system and circadian misalignment impact glucose tolerance via separate

mechanisms in humans. Proc Natl Acad Sci U S A. 112(17):E2225-2234.

https://doi.org/10.1073/pnas.1418955112

Nestorowa S, Hamey FK, Pijuan Sala B, Diamanti E, Shepherd M, Laurenti E, Wilson NK, Kent DG,

Göttgens B. 2016. A single-cell resolution map of mouse hematopoietic stem and progenitor cell

differentiation. Blood. 128(8):e20–e31. https://doi.org/10.1182/blood-2016-05-716480

Offield MF, Jetton TL, Labosky PA, Ray M, Stein RW, Magnuson MA, Hogan BL, Wright CV. 1996.

PDX-1 is required for pancreatic outgrowth and differentiation of the rostral duodenum. Development.

122(3):983–995. https://doi.org/10.1242/dev.122.3.983

Pan A, Schernhammer ES, Sun Q, Hu FB. 2011. Rotating night shift work and risk of type 2 diabetes:

two prospective cohort studies in women. PLoS Med. 8(12):e1001141.

https://doi.org/10.1371/journal.pmed.1001141

Papagiannakopoulos T, Bauer MR, Davidson SM, Heimann M, Subbaraj L, Bhutkar A, Bartlebaugh J,

Vander Heiden MG, Jacks T. 2016. Circadian Rhythm Disruption Promotes Lung Tumorigenesis. Cell

Metab. 24(2):324–331. https://doi.org/10.1016/j.cmet.2016.07.001

Parascandolo A, Bonavita R, Astaburuaga R, Sciuto A, Reggio S, Barra E, Corcione F, Salvatore M,

Mazzoccoli G, Relógio A, Laukkanen MO. 2020. Effect of naive and cancer-educated fibroblasts on

colon cancer cell circadian growth rhythm. Cell Death Dis. 11(4):289. https://doi.org/10.1038/s41419-

020-2468-2

Parent M-É, El-Zein M, Rousseau M-C, Pintos J, Siemiatycki J. 2012. Night work and the risk of cancer

among men. Am J Epidemiol. 176(9):751–759. https://doi.org/10.1093/aje/kws318

Perelis M, Marcheva B, Ramsey KM, Schipma MJ, Hutchison AL, Taguchi A, Peek CB, Hong H, Huang

W, Omura C, et al. 2015. Pancreatic β cell enhancers regulate rhythmic transcription of genes controlling

insulin secretion. Science. 350(6261):aac4250. https://doi.org/10.1126/science.aac4250

Perelis M, Ramsey KM, Marcheva B, Bass J. 2016. Circadian Transcription from Beta Cell Function to

Diabetes Pathophysiology. J Biol Rhythms. 31(4):323–336. https://doi.org/10.1177/0748730416656949

Pittendrigh CS. 1993. Temporal organization: reflections of a Darwinian clock-watcher. Annu Rev

Physiol. 55:16–54. https://doi.org/10.1146/annurev.ph.55.030193.000313

Preitner N, Damiola F, Lopez-Molina L, Zakany J, Duboule D, Albrecht U, Schibler U. 2002. The orphan

nuclear receptor REV-ERBalpha controls circadian transcription within the positive limb of the

mammalian circadian oscillator. Cell. 110(2):251–260. https://doi.org/10.1016/s0092-8674(02)00825-5

Reichert M, Rustgi AK. 2011. Pancreatic ductal cells in development, regeneration, and neoplasia. J Clin

Invest. 121(12):4572–4578. https://doi.org/10.1172/JCI57131

Roenneberg T, Allebrandt KV, Merrow M, Vetter C. 2012. Social jetlag and obesity. Curr Biol.

22(10):939–943. https://doi.org/10.1016/j.cub.2012.03.038

Sahai E, Astsaturov I, Cukierman E, DeNardo DG, Egeblad M, Evans RM, Fearon D, Greten FR,

Hingorani SR, Hunter T, et al. 2020. A framework for advancing our understanding of cancer-associated

fibroblasts. Nat Rev Cancer. 20(3):174–186. https://doi.org/10.1038/s41568-019-0238-1

Sato TK, Panda S, Miraglia LJ, Reyes TM, Rudic RD, McNamara P, Naik KA, FitzGerald GA, Kay SA,

Hogenesch JB. 2004. A functional genomics strategy reveals Rora as a component of the mammalian

circadian clock. Neuron. 43(4):527–537. https://doi.org/10.1016/j.neuron.2004.07.018

Scheer FAJL, Hilton MF, Mantzoros CS, Shea SA. 2009. Adverse metabolic and cardiovascular

consequences of circadian misalignment. Proc Natl Acad Sci U S A. 106(11):4453–4458.

https://doi.org/10.1073/pnas.0808180106

Scheiermann C, Kunisaki Y, Frenette PS. 2013. Circadian control of the immune system. Nat Rev

Immunol. 13(3):190–198. https://doi.org/10.1038/nri3386

Schernhammer ES, Laden F, Speizer FE, Willett WC, Hunter DJ, Kawachi I, Fuchs CS, Colditz GA.

2003. Night-shift work and risk of colorectal cancer in the nurses' health study. J Natl Cancer Inst.

95(11):825–828. https://doi.org/10.1093/jnci/95.11.825

Schwartz PB, Walcheck MT, Berres ME, Nakaya M, Wu G, Carrillo ND, Matkowskyj KA, Ronnekleiv-

Kelly SM. 2021. Chronic Jetlag-Induced Alterations in Pancreatic Diurnal Gene Expression. Physiol

Genomics. https://doi.org/10.1152/physiolgenomics.00022.2021

Sergushichev AA. 2016. An algorithm for fast preranked gene set enrichment analysis using cumulative

statistic calculation. bioRxiv.:060012. https://doi.org/10.1101/060012

Shi C, Klein AP, Goggins M, Maitra A, Canto M, Ali S, Schulick R, Palmisano E, Hruban RH. 2009.

Increased Prevalence of Precursor Lesions in Familial Pancreatic Cancer Patients. Clin Cancer Res.

15(24):7737–7743. https://doi.org/10.1158/1078-0432.CCR-09-0004

Shin EJ, Topazian M, Goggins MG, Syngal S, Saltzman JR, Lee JH, Farrell JJ, Canto MI. 2015. Linear-

array EUS improves detection of pancreatic lesions in high-risk individuals: a randomized tandem study.

Gastrointest Endosc. 82(5):812–818. https://doi.org/10.1016/j.gie.2015.02.028

Siegel RL, Miller KD, Fuchs HE, Jemal A. 2021. Cancer Statistics, 2021. CA Cancer J Clin. 71(1):7–33.

https://doi.org/10.3322/caac.21654

Simon TG, Roelstraete B, Sharma R, Khalili H, Hagström H, Ludvigsson JF. 2021. Cancer Risk in

Patients With Biopsy-Confirmed Nonalcoholic Fatty Liver Disease: A Population-Based Cohort Study.

Hepatology. 74(5):2410–2423. https://doi.org/10.1002/hep.31845

Smith T, Heger A, Sudbery I. 2017. UMI-tools: modeling sequencing errors in Unique Molecular

Identifiers to improve quantification accuracy. Genome Res. 27(3):491–499.

https://doi.org/10.1101/gr.209601.116

Stokes K, Nunes M, Trombley C, Flôres DEFL, Wu G, Taleb Z, Alkhateeb A, Banskota S, Harris C,

Love OP, et al. 2021. The Circadian Clock Gene, Bmal1, Regulates Intestinal Stem Cell Signaling and

Represses Tumor Initiation. Cell Mol Gastroenterol Hepatol. 12(5):1847-1872.e0.

https://doi.org/10.1016/j.jcmgh.2021.08.001

Sulli G, Manoogian ENC, Taub PR, Panda S. 2018. Training the Circadian Clock, Clocking the Drugs,

and Drugging the Clock to Prevent, Manage, and Treat Chronic Diseases. Trends Pharmacol Sci.

39(9):812–827. https://doi.org/10.1016/j.tips.2018.07.003

Takahashi JS. 2017. Transcriptional architecture of the mammalian circadian clock. Nat Rev Genet.

18(3):164–179. https://doi.org/10.1038/nrg.2016.150

Takele Assefa A, Vandesompele J, Thas O. 2020. On the utility of RNA sample pooling to optimize cost

and statistical power in RNA sequencing experiments. BMC Genomics. 21(1):312.

https://doi.org/10.1186/s12864-020-6721-y

Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, Lennon NJ, Livak KJ, Mikkelsen TS,

Rinn JL. 2014. The dynamics and regulators of cell fate decisions are revealed by pseudotemporal

ordering of single cells. Nat Biotechnol. 32(4):381–386. https://doi.org/10.1038/nbt.2859

Vetter C. 2020. Circadian disruption: What do we actually mean? Eur J Neurosci. 51(1):531–550.

https://doi.org/10.1111/ejn.14255

Walcheck MT, Matkowskyj KA, Turco A, Blaine-Sauer S, Nukaya M, Noel J, Ronnekleiv OK,

Ronnekleiv-Kelly SM. 2021. Sex-dependent development of Kras-induced anal squamous cell carcinoma

in mice. PLoS One. 16(11):e0259245. https://doi.org/10.1371/journal.pone.0259245

West AC, Bechtold DA. 2015. The cost of circadian desynchrony: Evidence, insights and open questions.

Bioessays. 37(7):777–788. https://doi.org/10.1002/bies.201400173

West AC, Smith L, Ray DW, Loudon ASI, Brown TM, Bechtold DA. 2017. Misalignment with the

external light environment drives metabolic and cardiac dysfunction. Nat Commun. 8(1):417.

https://doi.org/10.1038/s41467-017-00462-2

Wright KP, McHill AW, Birks BR, Griffin BR, Rusterholz T, Chinoy ED. 2013. Entrainment of the

human circadian clock to the natural light-dark cycle. Curr Biol. 23(16):1554–1558.

https://doi.org/10.1016/j.cub.2013.06.039

Wu G, Ruben MD, Francey LJ, Lee YY, Anafi RC, Hogenesch JB. 2022. An in silico genome-wide

screen for circadian clock strength in human samples [Internet]. [accessed 2022 Sep

8]:2022.05.10.491250. https://doi.org/10.1101/2022.05.10.491250

Wu J, Liu X, Nayak SG, Pitarresi JR, Cuitiño MC, Yu L, Hildreth BE, Thies KA, Schilling DJ, Fernandez

SA, et al. 2017. Generation of a pancreatic cancer model using a Pdx1-Flp recombinase knock-in allele.

PLoS One. 12(9):e0184984. https://doi.org/10.1371/journal.pone.0184984

Yamazaki S, Numano R, Abe M, Hida A, Takahashi R, Ueda M, Block GD, Sakaki Y, Menaker M, Tei

H. 2000. Resetting central and peripheral circadian oscillators in transgenic rats. Science. 288(5466):682–

685. https://doi.org/10.1126/science.288.5466.682

894 Zhang J, Francois R, Iyer R, Seshadri M, Zajac-Kaye M, Hochwald SN. 2013. Current understanding of

895 the molecular biology of pancreatic neuroendocrine tumors. J Natl Cancer Inst. 105(14):1005–1017.

896 https://doi.org/10.1093/jnci/djt135

A. 5 months

B. 9 months

C.

	5 months			9 months		
	Normal Circadian	Chronic Jetlag	*p value*	Normal Circadian	Chronic Jetlag	*p value*
PanIN1	100%	96.20%	0.43	100%	100%	1.00
PanIN2	5.7%	7.70%	1.00	3.0%	23.70%	0.02
PanIN3	0%	0%	1.00	6.10%	10.30%	0.68
PDAC	0%	3.80%	0.43	15.20%	10.30%	0.72

D.

E.

A.

Cell Type	Gene Name	Log2 FC	Adj p value
Endothelial Cell	*Tef*	0.326049301	0.001081832
Fibroblast	*Dbp*	0.606422244	7.90E-208
Fibroblast	*Per3*	0.574324977	1.74E-209
Fibroblast	*Tef*	0.395985989	2.46E-123
Fibroblast	*Nr1d2*	0.334812842	3.82E-79
Fibroblast	*Hlf*	0.333177448	5.96E-101
Fibroblast	*Per2*	0.270340607	1.22E-130
Macrophage	*Dbp*	0.425914793	4.73E-111
Memory CD4 T Cell	*Bhlhe40*	-0.480979686	4.00E-22
Non-Classical Monocyte	*Dbp*	0.341766669	1.68E-18

B.