**Chronic Jetlag Accelerates Pancreatic Neoplasia in Conditional *Kras*-Mutant Mice**

**Authors:** Patrick B Schwartz, MD[1], Morgan T Walcheck, PhD[1], Manabu Nukaya, PhD[1], Derek M

Pavelec, PhD[2], Kristina A Matkowskyj[3,4,5], and Sean M Ronnekleiv-Kelly, MD[1,5]

**Affiliations:**

[1]Department of Surgery, Division of Surgical Oncology, University of Wisconsin School of Medicine and

Public Health, Madison, WI

[2]Biotechnology Center, University of Wisconsin, Madison, WI

[3]Department of Pathology and Laboratory Medicine, University of Wisconsin School of Medicine and

Public Health, Madison, Wisconsin.

[4]William S Middleton Memorial Veterans Hospital, Madison, Wisconsin.

[5]University of Wisconsin Carbone Cancer Center, University of Wisconsin School of Medicine and

Public Health, Madison, WI

**Corresponding Author:** Sean M Ronnekleiv-Kelly, MD (ronnekleiv-kelly@surgery.wisc.edu)

**Address:**  600 Highland Ave. Bx 7375, K3/705 CSC, Madison, WI, 53792

**Phone:** (608) 262-2025 **Fax:** (608) 252-0913

**Running Title:** Chronic Jetlag Accelerates Pancreatic Neoplasia

**Disclosure:** The authors declare no conflict of interest.

**Abstract:**

Pancreatic ductal adenocarcinoma (PDAC) is a devastating cancer with an 80% five-year mortality rate –

reflecting an inadequate understanding of pertinent drivers of cancer development. One potential driver of

PDAC development is circadian disruption, known to control a myriad of pancreatic processes. Clock

dysfunction is associated with multiple risk factors for PDAC, including obesity, diabetes, and heightened

inflammation, and has been shown to accelerate the development of other solid-organ cancers. Using a

well-characterized model of chronic jetlag (CJ) to mimic irregular sleep patterns, we subjected mice

harboring a pancreas-specific activated *Kras* mutation (KC mice) to either a normal lighting pattern or CJ

and observed the resultant histopathologic changes. We found that CJ accelerated the development of

fibroinflammatory infiltrate/pancreatic intraepithelial neoplasia and increased the grade of the pancreatic

lesions. Applying single-cell RNA sequencing over multiple time points, we further identified fibroblasts

in the developing pancreatic tumor microenvironment as a putative target of CJ-induced changes. Our

results suggest a role for clock dysfunction in PDAC initiation which may hold relevance for preventative

efforts.

43    **Keywords:** Circadian, Pancreas, Jetlag, Pancreatic Ductal Adenocarcinoma, Mouse

## Introduction:

Pancreatic ductal adenocarcinoma (PDAC) is a highly lethal cancer due to its early spread and the limited effectiveness of current therapies. Of the 60,430 people estimated to be diagnosed in 2021, approximately 48,220 (~80%) will die of their disease.[1] This high mortality rate reflects, in part, an inadequate understanding of essential factors driving PDAC development. Insight into these factors could lead to new therapeutic or preventive strategies. One factor that has a strong association with cancer development is the disruption of circadian rhythm caused by irregular sleep patterns.[2]

Circadian clock proteins control daily biological rhythms with an oscillation of 24 hours.[3] Control from the central clock in the hypothalamus assimilates cues from day/night cycles (e.g. sleep patterns) and food intake to modulate peripheral organ rhythmic expression of core circadian genes.[3] In addition to central control, nearly all cells have an autonomous circadian clock based on transcriptional feedback mechanisms of core clock genes (CCGs). Collectively, the CCGs tightly regulate clock-controlled genes (e.g. downstream genes controlled by the clock) that modulate a myriad of cellular processes. Circadian disruption is a condition that affects a substantial proportion of the U.S. population, particularly shift workers and military personnel, whose sleep patterns can be altered dramatically by work schedule requirements.[4] Dysregulation of normal circadian rhythms can result in non-rhythmic and aberrant gene signaling, which can substantially alter organ function and promote various pathologies such as cancer development.[5] Studies have shown a link between circadian disruption and the risk of developing colorectal, lung, and liver cancers, among others.[2,5,6] Furthermore, disruption of this highly regulated process often results in diabetes mellitus, obesity, and increased inflammation- common risk factors for PDAC development.[5,7,8]

We sought to evaluate the effects of a chronic phase shift protocol, known to mimic sleep deprivation (also known as chronic jetlag [CJ]), on PDAC development.[6,9] To accomplish this, we conducted our experiments using the well-established $Kras^{G12D/+}$; $Pdx1$-Cre (KC) model of PDAC development.[10] After phase shift protocol, the pancreata of CJ KC mice underwent histopathologic evaluation revealing

increased fibroinflammatory/pre-neoplastic lesions (i.e. pancreatic intraepithelial neoplasia or PanINs)

during PDAC development, resulting in increased PanIN grade. Follow-up single-cell RNA sequencing

(scRNA-seq) revealed a tumor microenvironment consistent with our histologic analysis. We further

identified cancer-associated fibroblasts (CAFs) as the putative target of the effects of CJ. This work

serves as a foundation for the understanding of CJ-induced acceleration of PDAC development.

**Materials and Methodology:**

*Mouse Care and Husbandry*

All animal studies were conducted according to an approved protocol (M005959) by the University of

Wisconsin School of Medicine and Public Health (UW SMPH) Institutional Animal Care and Use

Committee (IACUC). Male and female mice were housed in an Assessment and Accreditation of

Laboratory Animal Care (AALAC) accredited selective pathogen-free facility (UW Medical Sciences

Center) on corncob bedding with chow diet (Mouse diet 9F 5020; PMI Nutrition International) and water

*ad libitum*.

Conditional $Kras^{G12D/+}$ (K) mice (B6.129S4-Kras tm4Tyj/J (#008179)) bred with mice expressing Cre-

Recombinase driven by the $Pdx1$ promoter (C) (B6.FVB-Tg(Pdx1-cre)6Tuv/J) (#014647)) to generate

$Kras^{G12D/+}$, $Pdx1$-Cre (KC) mice.[10] The KC mouse has been well characterized previously.[10] To evaluate

the effect of loss of clock function on KC mouse pathology the conditional $Arntl^{fx/fx}$ mouse, kindly gifted

by Professor Christopher Bradfield, was crossed with the KC mouse to generate the $Arntl^{fx/fx}$, $Kras^{G12D/+}$;

$Pdx1$-Cre (AKC) mouse model.[11,12] To identify cell types targeted by $Pdx1$-Cre recombinase in the KC

mouse model, the KC mouse was additionally crossed with the conditional Ai14 mouse expressing

tdTomato (B6.Cg-Gt(ROSA)26Sortm14(CAG-tdTomato)Hze/J (#007914)) to produce the KCT mouse.

The presence of each allele was confirmed with qRT-PCR through Transnetyx (Transnetyx Inc, Cordova,

TN). The K, C, and T mice were purchased from Jackson Laboratory (Bar Harbor, ME).

*Chronic Jetlag Protocol*

Starting at 4-6 weeks of age, KC mice were subjected to either a normal circadian (NC) 12-hour light:

dark (LD) cycle condition, or a chronic jetlag protocol (CJ) known to mimic altered sleep-wake cycles.[6,9]

CJ consisted of a 12-hour LD cycle, phase-shifted forward 4 hours every 2-3 days. Mice were kept under

their respective conditions until the study endpoint at 5 and 9 months of age. Kaplan Meier survival

analysis between NC and CJ KC mice was made by log-rank testing.

*Histologic Analysis*

Mouse pancreatic tissue was fixed in 10% neutral buffered formalin for 48 hours and then stored in 70%

ethanol until slide preparation. Slides were processed, embedded, sectioned, and stained with hematoxylin

and eosin (H&E) by the University of Wisconsin Carbone Cancer Center (UWCCC) Experimental

Animal Pathology Laboratory (EAPL) Core. All slides were assessed for the presence of precursor lesions

(PanIN) along with grading on a scale of 1-3 and/or pancreatic ductal adenocarcinoma (PDAC) by a

blinded, board-certified, gastrointestinal pathologist (KAM). Categorical comparisons between groups

were made with the Chi-squared or Fisher's Exact test as appropriate. Quantification of the total percent

pancreas involvement by the fibroinflammatory infiltrate, atrophy, and/or neoplastic lesions (i.e. PanINs)

during the development of PDAC was made by 3 separate reviewers, and differences were tested using

Student's t-test between groups. Inter-rater reliability statistics were computed and the intraclass

correlation coefficient is reported with associated 95% confidence intervals.

*Immunohistochemistry*

KCT mouse pancreas samples were formalin-fixed, embedded, and sectioned as above by the EAPL Core

facility. Immunohistochemistry (IHC) for red fluorescence protein (tdTomato) was also performed by

EAPL. Sections were deparaffinized in xylenes and hydrated through graded alcohols to distilled water.

Antigens were retrieved using citrate buffer pH 6.0 (10 mM Citric Acid and 0.05% tween 20).

Endogenous peroxidase was blocked with 0.3% $H_2O_2$ in PBS for 10 minutes at room temperature and

blocking of non-specific binding was performed using 10% goat serum. Sections were incubated with

rabbit anti-RFP antibody (600-401-379, Rockland Inc, Pottstown, PA) (1:1600) overnight at 4ºC. After

rinsing, sections were incubated with ImmPRESS goat anti-rabbit IgG (MP-7451, Vector Laboratories,

Burlingame, CA) for 30 minutes at room temperature. Detection was performed using the DAB substrate

kit (8059S, Cell Signaling Technology, Danvers, MA). Samples were counterstained using Mayer's

hematoxylin (MHS32, Millipore-Sigma, St. Louis, MO) for one minute, followed by rinsing and

coverslipping of slides.

*scRNAseq Sample Preparation, Library Construction, and Sequencing*

Five- and 9-month-old NC and CJ KC mice were sacrificed with cervical dislocation and the pancreas

was rapidly dissected ($< 60$ seconds) and placed in cold PBS (Thermo Fisher Scientific, Waltham, MA).

One male and female pancreas was pooled from each condition and time point. Pancreas samples were

minced until the tissue was $< 1$ mm in diameter, and then centrifuged at 900 RPM for 5 minutes to

remove fat contamination. Samples were then incubated with 5 mL of Collagenase P (Sigma Aldrich, St.

Louis, MO) in HBSS (Thermo Fisher Scientific) at $37°$ C for 15 minutes while being shaken at 100 RPM.

Additionally, samples were manually agitated every 2-3 minutes. The digestion reaction was quenched

with 10 mL R10 media. Samples were then centrifuged at 900 RPM and washed twice with 10 mL of

R10. Samples were then strained in a gradient fashion using 500 μm, 100 μm, and 40 μm strainers.

Samples were then live/dead sorted into single-cell populations and were transferred to the University of

Wisconsin Gene Expression Center (GEC) for Library Preparation.

In brief, libraries were constructed according to the Chromium NextGEM Single Cell 3' Reagent Kit v3.1

User Guide, Rev.D (10x Genomics, Pleasanton, CA). Single-cell suspension cell concentration and

viability were quantified on the Countess II (Thermo Fisher Scientific) using 0.4% Trypan Blue

(Invitrogen, Carlsbad, CA). Single-cell droplet encapsulation was performed using the Single Cell G Chip

and the 10x Genomics Chromium Controller. Following the completion of the Chromium run the GEMs

were transferred to emulsion-safe strip tubes for GEM-RT using an Eppendorf MasterCycler Pro

thermocycler (Eppendorf, Hamburg, Germany). Following RT, GEMs were broken, and the pooled

single-cell cDNA was collected in Recovery Agent, purified using DynaBeads MyOne Silane beads

(Thermo Fisher Scientific), and amplified with the appropriate number of PCR cycles based on the

number of cells targeted for recovery. Post-cDNA amplified product was purified using SPRIselect

(Beckman Coulter, Brea, CA) and quantified on a Bioanalyzer 2100 (Agilent, Santa Clara, CA) using the

High Sensitivity DNA kit. Adapters were then added to the libraries after fragmentation, end repair, A-

tailing, and double-sided size selection using SPRIselect. Following adapter ligation, libraries were

cleaned up using SPRIselect and sample-specific indexes (Chromium i7 Multiplex Kit, 10x Genomics)

were added by sample index PCR using the appropriate number of PCR cycles based on the cDNA yield

calculated by the Bioanalyzer quantification. After sample index PCR, samples were double-size selected

using SPRIselect and quantified by the Qubit High Sensitivity DNA Kit (Thermo Fisher Scientific).

Libraries were sequenced on a NovaSeq6000 (Illumina, San Diego, CA).

*scRNAseq Data Analysis*

Single-cell RNAseq data were analyzed by the UW Bioinformatics Resource Center. Data were

demultiplexed using the Cell Ranger Single Cell Software Suite, mkfastq command wrapped around

Illumina's bcl2fastq. The MiSeq balancing run was quality-controlled using calculations based on UMI-

tools.[13] Sample libraries were balanced for the number of estimated reads per cell and run on an Illumina

NovaSeq system. Cell Ranger software was then to perform demultiplexing, alignment, filtering, barcode

counting, UMI counting, and gene expression estimation for each sample according to the 10x Genomics

documentation (https://support.10xgenomics.com/single-cell-gene-

expression/software/pipelines/latest/what-is-cell-ranger). The gene expression estimates from each

sample were then aggregated using Cellranger (cellranger aggr) to compare experimental groups with

normalized sequencing-depth and expression data.

Gene expression data was then processed using the Seurat version 4 pipeline for data integration.[14] Data

was loaded into Seurat. Low quality (<200 features or >20% mitochondrial gene content) and doublets

were removed. Cells expressing more than 1% hemoglobin gene content were filtered out. The NC and CJ

KC data were merged and integrated using the SCTransform v2 wrapper with 3000 unique features.[15]

SCTransform allows for normalization and better batch correction based on a regularized negative

binomial regression. The regression additionally allowed for correction of the cell cycle effect and

mitochondrial gene content.[16] A principal component analysis (PCA) was performed with the number of

dimensions (principal components) selected based on an elbow plot. Data were visualized using the

Uniform Manifold Approximation and Projection for Dimension Reduction (UMAP).[17] Cell identities

were called based on the expression of known markers for each cell type and scored using the ScType

package to cross-validate identities following the default settings for SCT transformed data in the

provided vignette.[18] Differential expression between conditions and time points was performed between

clusters or conditions in Seurat using the MAST test based on a log fold change threshold of 0.25 and a

false discovery rate (FDR) corrected $p$ value of $p < 0.05$. The MAST test within Seurat is based on the

MAST package, which is specifically designed to handle differential expression in scRNA-seq data and

utilizes a hurdle model.[19] Gene Set Enrichment Analysis (GSEA) was conducted with the fgsea package

leveraged against oncogenic signature gene set (c6.all.v7.5.1.symbols) and biologic process subset of

gene ontology (GO BP) (c5.go.v7.5.1.symbols).[20] Pseudotime trajectory analysis was performed with the

Monocle 3 package.[21] Genes correlated with pseudotime progression were computed using the Moran's I

test and those genes with an FDR $p < 0.05$ were considered significant. \All analyses in this manuscript

were conducted in R version 4.2.0 (Vienna, Austria) or GraphPad Prism version 8.3.1 (Dotmatics,

Boston, MA).

**Results:**

*Chronic jetlag promotes pancreatic tumor initiation*

To better understand the effect of chronic jetlag (CJ) on pancreatic cancer development, we elected to

utilize the well-characterized $Kras^{G12D/+}$, *Pdx1*-Cre (KC) mouse model and assessed the burden of

neoplastic lesions (i.e. PanIN and PDAC). All slides were scored for their degree of involvement by 3

independent reviewers with an excellent intraclass correlation coefficient of 0.925 (95% CI 0.911-0.938;

$p < 0.01$). We found that there was a 39.52% increase (12.73% vs. 52.25%) in pancreatic involvement by

neoplastic lesions (e.g. PanINs) and the associated fibroinflammatory infiltrate (hereafter referred to as

FI/PanIN) from five to nine months in normal circadian (NC) KC mice (**Figure 1A**). Interestingly, KC

mice that were exposed to CJ were found to have an approximately 2-fold increase in FI/PanIN at 5

months of development (12.73% vs. 22.47%; **Figure 1A**) – which equilibrated by 9 months (52.25% vs.

43.38%; **Figure 1B**).

After demonstrating CJ increased the degree of FI/PanIN at five months, we evaluated the grade of

neoplasia at both five and nine months. Each pancreas was evaluated for PanIN1-3 and PDAC. All NC

and CJ KC mice at five and nine months exhibited PanIN1 except a single five-month CJ KC mouse that

had evidence of dedifferentiated PDAC which had replaced the entirety of the pancreatic parenchyma

(**Figure 2A**). When evaluating higher grade lesions, few mice developed PanIN2-3 or PDAC at 5 months

(**Figure 2A**). However, at 9 months, KC mice subjected to CJ had an increased incidence of PanIN2

(**Figure 2B**). Therefore, CJ promoted a greater degree of FI/PanIN initiation leading to increased PanIN

grade by nine months.

*Pancreatic clock gene deletion abrogated tumor initiation*

After establishing the effect of CJ on pancreatic tumor development, we next assessed whether circadian

disruption within the developing tumor cells similarly demonstrated an accelerated phenotype. We

crossed the KC mouse with a conditional deletion of the core clock gene *Arntl* to produce *Arntl*^fx/fx;

*Kras*^G12D/+; *Pdx1*-Cre (AKC) mice, which possess *Arntl* deletion in cells expressing the *Kras*^G12D

mutation.[11] *Arntl* is a central transcriptional regulator of the clock and essential for clock functionality.[22]

We evaluated the degree of FI/PanIN in NC AKC mice at 5 months and, surprisingly, we found that the

lesions had nearly disappeared. Compared to KC mice, AKC mice had an average of 1.59% of pancreatic

involvement (vs. 12.73%; $p < 0.01$), and no mice exhibited a higher-grade lesion than a PanIN1 (**Figure

3A**). To understand which cell populations may cause resistance to pancreatic neoplasm formation with

loss of *Arntl*, we crossed the KC mouse with the conditional Ai14 mouse expressing tdTomato (**Figure

**3B**). This revealed tdTomato expression (and thus *Kras*-mutation and *Arntl* deletion) in acinar cells,

ductal cells, PanINs and islet cells, with an absence of expression in the fibroinflammatory infiltrate

(chronic pancreatitis/atrophy) surrounding the PanINs. Thus, although CJ enhances pancreatic tumor

initiation in our mouse model, the loss of *Arntl* expression in pancreatic cells revealed that an intact

circadian clock is likely needed to promote the developing pancreatic FI/PanIN.

*Single-cell RNA sequencing of KC mice recapitulates the highly heterogeneous tumor microenvironment*

To better understand how CJ was modulating the pancreatic epithelial cells within the tumor

microenvironment to progress toward neoplasia, we evaluated the developing tumor microenvironment

over time. To accomplish this, we performed single-cell RNA sequencing (scRNAseq) from pooled

pancreatic samples (1 male/female) from both NC and CJ KC mice at 5 and 9 months (**Figure 4B**). After

filtering and dimensionality reduction, a total of 14 different heterogeneous cell types were annotated

using the ScType package comprising 42,211 cells (**Figure 4A & Supplemental Data 1-2**).[18]

We started by subsetting the pancreatic epithelial cells, which should encompass both normal acinar and

developing PanINs, and used pseudotime trajectory analysis to track cell-state changes that occurred from

5 to 9 months. Pseudotime was set to zero at the point with the highest fraction of cells from the 5-month

mice and allowed to progress (**Figure 4C**). A Moran's I autocorrelation analysis was then performed to

obtain genes that varied across the trajectory (**Supplemental Data 3**). Gene set enrichment analysis

(GSEA) of the genes correlated with pseudotime progression was associated with multiple *Kras*-driven

cancer cell line signatures – corresponding to the development of pancreatic neoplasia over time (**Figure**

**4D**). Therefore, we identified a collection of pancreatic epithelial cells that appeared to represent the

developing pancreatic cancer microenvironment, which provided a foundation for the subsequent analysis

to understand the effects of CJ.

*Chronic jetlag promotes differential clock gene expression in fibroblasts*

We performed DGE across the collective 5- and 9-month mice for differences in core clock gene (CCG)

and clock-associated gene expression between the NC and CJ conditions (**Supplemental Data 4**). We

hypothesized that clusters with significantly altered expression would be important for downstream

investigation. Perhaps most intriguing, the acinar cell/PanIN clusters that we identified and confirmed

from the pseudotime analysis did not exhibit DGE. We found that endothelial cells, macrophages,

memory CD4 T cells, and monocytes each had a single clock-associated gene differentially expressed

(**Figure 5A**). Interestingly, fibroblasts were the only cell type that exhibited DGE of CCGs and clock-

associated genes, including *Per2*, *Per3*, *Nr1d2*, *Hlf*, *Tef*, and *Dbp* (**Figure 5A-B**). Fibroblasts are an

integral part of the PDAC microenvironment and are known to exhibit circadian rhythmicity when

compared to naïve fibroblasts.[23] The lack of other cell types exhibiting changes in clock-specific gene

expression suggests fibroblasts may be the target of the CJ-induced accelerated neoplastic phenotype.

*Fibroblasts exhibit time-dependent effects secondary to chronic jetlag*

To further evaluate the fibroblasts within our dataset, we isolated three different subclusters (**Figure 6A**).

Using existing markers generated for cancer-associated fibroblasts (CAFs), including inflammatory CAFs

(iCAFs), myofibroblast CAFs (mCAFs), and antigen-presenting CAFs (apCAFs), we assigned identities

with ScType (**Supplementary Data 5-6**).[24] Of the three clusters, we identified all three as iCAFs – with

the expression of known markers, such as *Cle3b*, *Il6*, *Has1*, and *Col14a1* (**Figure 6B**). Notably, fibroblast

Cluster 3 exhibited some characteristics consistent with apCAFs, including expression of multiple MHC

II components *H2-Aa*, *H2-Eb1*, and *H2-ab1* and *Cd74*, the invariant chain of MHC II (Figure 6B). We

then performed DGE and GSEA between NC and CJ iCAFs at 5 and 9 months (**Figure 7A-B** &

**Supplemental Data 7**). Consistent with our prior work on the effects of CJ on mouse pancreas, we found

that multiple metabolic pathways in fibroblasts were enriched at 5 months (**Figure 7A**).[9,25] Notably, our

9-month comparison revealed enrichment of several apoptosis-related pathways, including

'NEGATIVE_REGULATION_OF_CELL_DEATH' and

'REGULATION_OF_APOPTOTIC_SIGNALING_PATHWAY' (**Figure 7B**). To evaluate if CJ was

simply increasing fibrosis (i.e. collagen deposition) to facilitate these changes, we compared the collagen

content from five- and nine-month NC and CJ KC mice (**Supplemental Data 8**). We found no

differences in the collagen content at either timepoint. We, therefore, identified that the iCAFs in the

developing tumor microenvironment of KC mice had early enrichment of metabolic pathways in response

to CJ with late enrichment of apoptosis-related pathways – suggesting CJ augments fibroblast signaling to

promote tumor formation.

**Discussion:**

Pancreatic ductal adenocarcinoma (PDAC) is an aggressive malignancy that has a high rate of mortality.[1]

This high rate of mortality is, at least in part, due to an incomplete understanding of the pathogenesis of

PDAC. One pathogenic factor being explored in cancers, including PDAC, is the circadian clock.[5,6,26,27]

To further evaluate the role of clock dysfunction in PDAC development, we subjected mice to a well-

characterized chronic jetlag (CJ) protocol.[9,25] Interestingly, in KC mice prone to PDAC formation, we

found that CJ promoted an increase in FI/PanIN development leading to more advanced lesions.[10] We

then used a series of scRNAseq experiments to identify fibroblasts as the susceptible cell type to CJ

within the developing PDAC tumor microenvironment. Further analysis revealed an enrichment of

apoptosis-related pathways, including the negative regulation of cellular death. Future experiments will

explore the role of clock dysfunction in promoting alterations in fibroblast signaling and how these

alterations promote the acceleration of PanIN formation.

To evaluate the development of PDAC, we selected a well-characterized model of PDAC development,

the KC mouse model.[10] Hingorani *et al*. characterized their mice at 2.25, 4.5, and 9 months. They noted

that few animals had dysplastic ducts before 5 months of age and higher-grade lesions were rare before 7-

10 months. Overall, the group noted few mice developed and succumbed to PDAC (6.89%). Therefore,

we selected 5 months as a timepoint where we could potentially note changes in the early initiation of

neoplasia during PDAC development, and 9 months to evaluate for the development of higher-grade

lesions. Similar to the original description, in mice housed under normal lighting conditions (NC) we

observed that between 5 and 9 months there was an increase in the FI/PanIN lesions. We further identified

an increase in the grade of PanIN lesions over time. When we subjected our KC mice to chronic jetlag

(CJ), we noted that there was an increase in the amount of FI/PanIN formation (22.47% vs. 12.73%)

compared to KC NC mice at 5 months. By 9 months, this appeared to correspond to an increased number

of PanIN2 lesions (23.7% vs. 5.1%). Overall, the rate of PDAC was low in either group at nine months

(NC = 15.4% vs. CJ = 10.5%). To better understand whether the observed increase in PanIN2 lesions

would translate to increased PDAC, a longer follow-up interval or a more aggressive PDAC mouse model

outside the current experimental scope (such as the KPC mouse harboring an additional mutation in *Tp53*)

would be required.[28] Thus, we identified CJ as a driver of PDAC initiation leading to an increase in PanIN

grade. Although previous evidence suggests a role for the clock in PDAC progression, this is the first

study to show perturbations in the clock increase tumor initiation in PDAC development.[29,30]

Recently, studies have shown similar effects on initiation in other cancer models. In mice predisposed to

colorectal cancer development, mice subjected to constant lighting showed a 2-fold increase in tumor

initiation compared to those housed under diurnal conditions.[31] In this study, Stokes *et al*. also showed

that global and epithelial loss of *Arntl* recapitulated the increases in tumor initiation, suggesting that *Arntl*

is a tumor suppressor in colorectal cancer development and that perturbations in the clock (altered

lighting) caused a similar phenotype to clock disruption (*Arntl* deletion). When we crossed our KC mice

with the condition $Arntl^{fx/fx}$ mouse, we found a discordant phenotype in pancreatic tumor initiation

compared to colorectal cancer. Specifically, there was a near abolishment of all FI/PanIN lesions at 5

months (1.59% AKC vs. 12.53% NC KC vs 22.47% CJ KC) – suggesting that an intact *Arntl* is required

for the development of PDAC. Previously, in our examination of the effects of CJ on the wildtype

pancreatic transcriptome, we found that diurnal rhythmicity was maintained, but widespread changes in

metabolic gene expression compared to wildtype NC mice were observed.[9] We also demonstrated that

CJ-induced misalignment of LD cues in wildtype mice was associated with changes in the pancreatic

lipidome characteristic of pathogenic states, such as diabetes mellitus – a known risk factor for PDAC.[25]

Other investigations have shown that circadian misalignment leads to higher rates of death, changes in

cardiac conduction physiology, and hepatocellular carcinoma.[32–35] Thus, although CJ does not cause clock

disruption (clock rhythmicity is maintained), misalignment can generate significant pathology.

Concordantly, in our model, it is more likely that circadian misalignment – where the internal clock is

dyssynchronous with external zeitgebers – rather than arrhythmicity, is the mechanism by which CJ

exerts its effects.[32] The findings in our study highlight a key distinction between behavioral models that

cause a phase shift in the clock and genetically engineered mouse models causing clock disruption: there

may not be an overlap in phenotype owing to differing effects on the clock, or the population of cells

impacted by behavioral versus genetic manipulation may be different. For instance, the pancreatic

epithelial cells were targeted by the conditional deletion of *Arntl* whereas CJ would ostensibly have a

more global impact. Future work will delve into whether conditional clock disruption in the apparent CJ

targeted cell population (i.e. fibroblasts) demonstrates a similar or dissimilar phenotype of accelerated

pancreatic FI/PanIN development.

To better understand how pancreatic neoplasia progressed in the setting of both NC and CJ conditions,

scRNAseq was performed at both 5 and 9 months. We found that the pancreatic epithelial cells, including

the PanINs and acinar cells, showed a time-dependent cancer signature on pseudotime analysis consistent

with the observed histopathologic changes. Subsequently, when we examined the 14 different cell types

for evidence of DGE between NC and CJ, we found that fibroblasts – and not the pancreatic epithelial

cells – were the only cell type with multiple CCG and clock-associated gene changes. This finding further

supports the notion that CJ augments aspects of the developing tumor microenvironment but not

necessarily the epithelial/PanINs themselves. Notably, IHC with KC mice conditionally expressing

tdTomato demonstrated that there was *Pdx1*-Cre-directed expression in the acinar, ductal, islet, and

PanINs, but not the mesenchymal lineage cells, including the fibroblasts and endothelial cells. Others

have also found that clock dysfunction in the supportive stroma of the tumor microenvironment may play

a critical role.[36] Alexander *et al*. noted that myeloid-specific deletion of *Arntl* led to increased growth of

melanoma tumors in mice. Furthermore, they found a similar result when they co-injected melanoma

along with exogenous macrophages harboring a deletion of *Arntl* compared to wildtype macrophages.

Collectively, these data indicated that the supportive components of the tumor microenvironment, such as

fibroblasts, may coordinate the observed increases in pancreatic neoplasia secondary to CJ.

When fibroblasts were subsetted in our scRNAseq experiment, we further identified all 3 clusters as

inflammatory CAFs (iCAFs) based on known markers.[24] CAFs are known to display differences in

rhythmic gene expression compared to naïve fibroblasts.[23] iCAFs secrete IL-6, CXCL9, and TGFβ and

can be both immunosuppressive as well as immune promoting.[37] Accordingly, we observed that CJ was

associated with enrichment in genes associated with leukocyte motility. Interestingly, we did not observe

any differences in the amount of collagen content at 5 or 9 months between NC and CJ KC mice,

suggesting a role for paracrine signaling in mediating the effects of CJ by fibroblasts rather than

extracellular matrix production (i.e. desmoplasia). After identifying and characterizing the CAFs, we

performed GSEA at 5 and 9 months, which demonstrated that at 5 months CJ promoted metabolic gene

enrichment – consistent with our prior observations of CJ on wildtype pancreas.[9] However, by 9 months,

there was enrichment of multiple cell death and apoptosis-related pathways. Correspondingly, CAFs

generate pro-tumor signals to support the developing cancer cells and have been shown to promote

decreased cancer cell death and resistance to apoptosis when co-cultured.[38]

Collectively, these experiments shed light on a novel role for the clock in mediating PDAC initiation

during development. Future work will be required to elucidate the mechanistic underpinnings which

mediate the effects of CJ on PDAC development. Identification of this connection could eventually lead

to preventative strategies for patients who are at high-risk for PDAC development, such as those with

strong family history or individuals with pancreatic pre-cancerous lesions.

 **Figure Legends:**

**Figure 1:** *Pancreatic fibroinflammation/PanIN development is increased by chronic jetlag.* **A.** At 5

months, CJ KC mice (n = 26) and NC KC mice (n= 36) were evaluated for pancreatic

fibroinflammation/PanIN (FI/PanIN )development. CJ was found to increase the degree of pancreatic

FI/PanIN development compared to NC mice (22.47% vs. 12.73%; *p* = 0.027). **B.** At 9 months the degree

of pancreatic FI/PanIN development was not different between CJ mice (n = 39) and NC (n = 33)

(43.38% vs. 52.25%; *p* = 0.06).

**Figure 2:** *Chronic jetlag accelerates the grade of PanIN lesions.* KC mice subjected to either NC (black)

or CJ (grey) conditions were systematically evaluated for PanINs1-3 and PDAC at 5 (n = 36 vs. n = 26)

and 9 months of age (n = 33 vs. n = 39). **A.** At 5 months, few mice exhibited lesions above PanIN1 or

PDAC **B.** At 9 months, KC mice subjected to CJ exhibited significantly higher rates of PanIN2 compared

to NC KC mice (23.7% vs. 5.1%; *p* =0.02). **C.** Kaplan Meier log-rank survival was not different between

NC (n = 37) and CJ (n = 43) KC mice (*p* = 0.31) during the 9-month study period. **D.** The categorical

comparisons between 5- and 9-month KC mice with Fisher's exact testing.

**Figure 3:** *Pancreatic cell deletion of Arntl abrogates pancreatic fibroinflammatory infiltrate/PanIN*

*development*. **A.** Five-month NC KC (n = 36) and NC AKC mice (n = 24) were evaluated for pancreatic

fibroinflammation/PanIN (FI/PanIN). Compared to KC mice, AKC mice were found to have their

pancreatic FI/PanIN abrogated (12.73% vs. 1.59%; *p* = 3.2e-06). **B.** To evaluate the cell types affected by

AKO, KC mice were crossed with Ai14 mice to create KCT mice. H&E stained sections (left) and

immunohistochemistry for tdTomato (right) were compared to evaluate cell-specific *Pdx-1* cre expression

in KC mice, which was found to be limited to pancreatic epithelial cells, including PanINs (P), acinar (A),

islet (I), and ductal (D) cells. Expression was not seen in vascular (V) cells (i.e. endothelial cells) or the

fibroinflammatory (F) infiltrate, including fibroblasts and immune cells.

**Figure 4:** *The single-cell landscape of KC mice recapitulates the developing pancreatic cancer tumor*

*microenvironment.* **A.** A total of 14 diverse cell types were identified in the collective single-cell RNA

sequencing (scRNAseq) dataset comprising 42,211 cells. **B.** Both 5- and 9-month time points were

assayed to correspond to histopathologic analyses. **C.** Pseudotime trajectory analysis was performed for

the subset of pancreatic acinar cells with 0 pseudotime set to the point with the highest fraction of cells at

5 months. **D.** Gene Set Enrichment Analysis (GSEA) was performed for the oncogenic signature gene set

associated with pseudotime progression and ordered by the log adjusted p value.

**Figure 5:** *Fibroblasts were preferentially affected by chronic jetlag.* **A.** Table of core clock genes (CCGs)

and clock-associated genes differentially expressed when comparing NC and CJ conditions of each cell

type. **B.** Representative UMAP dimensionality plots demonstrating cellular expression (blue color) of

select CCG and clock-associated genes in the subset of fibroblasts.

**Figure 6:** *Fibroblasts subclusters in the developing pancreatic tumor microenvironment were identified*

*as inflammatory cancer-associated fibroblasts.* **A.** A UMAP demonstrating the three fibroblast

subclusters identified on dimensionality reduction. **B.** Violin plots with select markers for Pan-,

Inflammatory-, Myofibroblastic-, and Antigen Presenting-Cancer Associated Fibroblasts (CAFs). Clusters

1-3 all express markers consistent with Pan-CAFs and Inflammatory CAFs. Cluster 3 also expresses

markers consistent with antigen-presenting CAFs.

**Figure 7:** *Chronic jetlag influences fibroblast expression patterns across time.* Figures with The Gene Set

Enrichment Analysis (GSEA) demonstrating the gene ontology biologic process (GO BP) pathways

enriched in fibroblasts from the NC and CJ conditions at **A.** five and **B.** nine months. The top 10

pathways at each timepoint are ordered by their log-adjusted p value.

**Supplementary Data:**

<u>Supplemental Data 1</u>: Marker database for single-cell RNA sequencing cell-type identification

Supplemental Data 2: ScType Scores for overall single-cell RNA sequencing dataset

Supplemental Data 3: Moran's I pseudotime analysis for the pancreatic acinar cells

Supplemental Data 4: Differential gene expression between normal circadian and chronic jetlag

conditions for all cell types

Supplemental Data 5: Marker database for fibroblast subtype identification

Supplemental Data 6: ScType Scores for fibroblast subtype identification

Supplemental Data 7: Differential gene expression between normal circadian and chronic jetlag

conditions for fibroblasts at 5 and 9 months

Supplemental Data 8: Quantification of collagen content in 5-month normal circadian and chronic jetlag

KC mice

**Acknowledgments:**

We would like to take the opportunity to acknowledge the University of Wisconsin Carbone Cancer

Center Support Grant P30 CA014520, which provides funding for the Experimental Animal Pathology

Lab Core.

We would also to acknowledge the Gene Expression Center and the Bioinformatics Resource Center

(BRC) at the University of Wisconsin, Madison for their contributions to this work.

Finally, we would like to thank the Michael W. Oglesby Foundation for their funding support of our work

in circadian disruption and pancreas pathology.

**Grants:**

The research reported in this publication was supported by the Department of Defense Peer Reviewed Cancer Research Program Award CA190176 (Grants.gov ID GRANT12935023) and the University of Wisconsin Institute for Clinical and Translational Research KL2 Award (UL1TR002373 and KL2TR002374) [SRK]. Funding support for this work was also provided by the National Institutes of Health under Award Number T32 ES007015 [PBS]. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

**References:**

1. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer Statistics, 2021. *CA Cancer J Clin*. 2021;71(1):7-33. doi:10.3322/caac.21654

2. Schernhammer ES, Laden F, Speizer FE, et al. Night-shift work and risk of colorectal cancer in the nurses' health study. *J Natl Cancer Inst*. 2003;95(11):825-828. doi:10.1093/jnci/95.11.825

3. McIntosh BE, Hogenesch JB, Bradfield CA. Mammalian Per-Arnt-Sim proteins in environmental adaptation. *Annu Rev Physiol*. 2010;72:625-645. doi:10.1146/annurev-physiol-021909-135922

4. Bass J, Lazar MA. Circadian time signatures of fitness and disease. *Science*. 2016;354(6315):994-999. doi:10.1126/science.aah4965

5. Kettner NM, Voicu H, Finegold MJ, et al. Circadian Homeostasis of Liver Metabolism Suppresses Hepatocarcinogenesis. *Cancer Cell*. 2016;30(6):909-924. doi:10.1016/j.ccell.2016.10.007

6. Papagiannakopoulos T, Bauer MR, Davidson SM, et al. Circadian Rhythm Disruption Promotes Lung Tumorigenesis. *Cell Metab*. 2016;24(2):324-331. doi:10.1016/j.cmet.2016.07.001

7. Marcheva B, Ramsey KM, Buhr ED, et al. Disruption of the clock components CLOCK and BMAL1 leads to hypoinsulinaemia and diabetes. *Nature*. 2010;466(7306):627-631. doi:10.1038/nature09253

8.   Roenneberg T, Allebrandt KV, Merrow M, Vetter C. Social jetlag and obesity. *Curr Biol*.

2012;22(10):939-943. doi:10.1016/j.cub.2012.03.038

9.   Schwartz PB, Walcheck MT, Berres ME, et al. Chronic Jetlag-Induced Alterations in Pancreatic

Diurnal Gene Expression. *Physiol Genomics*. Published online May 31, 2021.

doi:10.1152/physiolgenomics.00022.2021

10. Hingorani SR, Petricoin EF, Maitra A, et al. Preinvasive and invasive ductal pancreatic cancer and its

early detection in the mouse. *Cancer Cell*. 2003;4(6):437-450. doi:10.1016/s1535-6108(03)00309-x

11. Johnson BP, Walisser JA, Liu Y, et al. Hepatocyte circadian clock controls acetaminophen

bioactivation through NADPH-cytochrome P450 oxidoreductase. *Proc Natl Acad Sci U S A*.

2014;111(52):18757-18762. doi:10.1073/pnas.1421708111

12. Liu Y, Johnson BP, Shen AL, et al. Loss of BMAL1 in ovarian steroidogenic cells results in

implantation failure in female mice. *PNAS*. 2014;111(39):14295-14300.

doi:10.1073/pnas.1209249111

13. Smith T, Heger A, Sudbery I. UMI-tools: modeling sequencing errors in Unique Molecular

Identifiers to improve quantification accuracy. *Genome Res*. 2017;27(3):491-499.

doi:10.1101/gr.209601.116

14. Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal single-cell data. *Cell*.

2021;184(13):3573-3587.e29. doi:10.1016/j.cell.2021.04.048

15. Choudhary S, Satija R. Comparison and evaluation of statistical error models for scRNA-seq.

*Genome Biology*. 2022;23(1):27. doi:10.1186/s13059-021-02584-9

16. Nestorowa S, Hamey FK, Pijuan Sala B, et al. A single-cell resolution map of mouse hematopoietic

stem and progenitor cell differentiation. *Blood*. 2016;128(8):e20-e31. doi:10.1182/blood-2016-05-

716480

17. McInnes L, Healy J, Melville J. UMAP: Uniform Manifold Approximation and Projection for

Dimension Reduction. *arXiv:180203426 [cs, stat]*. Published online September 17, 2020. Accessed

February 4, 2021. http://arxiv.org/abs/1802.03426

18. Ianevski A, Giri AK, Aittokallio T. Fully-automated and ultra-fast cell-type identification using

specific marker combinations from single-cell transcriptomic data. *Nat Commun*. 2022;13(1):1246.

doi:10.1038/s41467-022-28803-w

19. Finak G, McDavid A, Yajima M, et al. MAST: a flexible statistical framework for assessing

transcriptional changes and characterizing heterogeneity in single-cell RNA sequencing data.

*Genome Biol*. 2015;16. doi:10.1186/s13059-015-0844-5

20. Sergushichev AA. An algorithm for fast preranked gene set enrichment analysis using cumulative

statistic calculation. *bioRxiv*. Published online June 20, 2016:060012. doi:10.1101/060012

21. Trapnell C, Cacchiarelli D, Grimsby J, et al. The dynamics and regulators of cell fate decisions are

revealed by pseudotemporal ordering of single cells. *Nat Biotechnol*. 2014;32(4):381-386.

doi:10.1038/nbt.2859

22. Bunger MK, Wilsbacher LD, Moran SM, et al. Mop3 is an essential component of the master

circadian pacemaker in mammals. *Cell*. 2000;103(7):1009-1017. doi:10.1016/s0092-8674(00)00205-

1

23. Parascandolo A, Bonavita R, Astaburuaga R, et al. Effect of naive and cancer-educated fibroblasts on

colon cancer cell circadian growth rhythm. *Cell Death Dis*. 2020;11(4):289. doi:10.1038/s41419-020-

2468-2

24. Elyada E, Bolisetty M, Laise P, et al. Cross-Species Single-Cell Analysis of Pancreatic Ductal

Adenocarcinoma Reveals Antigen-Presenting Cancer-Associated Fibroblasts. *Cancer Discov*.

2019;9(8):1102-1123. doi:10.1158/2159-8290.CD-19-0094

25. Schwartz PB, Barrett-Wilt GA, Ronnekleiv-Kelly SM. Chronic Jetlag Alters the Landscape of the

Pancreatic Lipidome. *Pancreas*. 2022;51(1):80-89. doi:10.1097/MPA.0000000000001962

26. Relles D, Sendecki J, Chipitsyna G, Hyslop T, Yeo CJ, Arafat HA. Circadian gene expression and

clinicopathologic correlates in pancreatic cancer. *J Gastrointest Surg*. 2013;17(3):443-450.

doi:10.1007/s11605-012-2112-2

27. Li W, Liu L, Liu D, et al. Decreased circadian component Bmal1 predicts tumor progression and

poor prognosis in human pancreatic ductal adenocarcinoma. *Biochem Biophys Res Commun*.

2016;472(1):156-162. doi:10.1016/j.bbrc.2016.02.087

28. Hingorani SR, Wang L, Multani AS, et al. Trp53R172H and KrasG12D cooperate to promote

chromosomal instability and widely metastatic pancreatic ductal adenocarcinoma in mice. *Cancer

Cell*. 2005;7(5):469-483. doi:10.1016/j.ccr.2005.04.023

29. Jiang W, Zhao S, Shen J, et al. The MiR-135b-BMAL1-YY1 loop disturbs pancreatic clockwork to

promote tumourigenesis and chemoresistance. *Cell Death Dis*. 2018;9(2):149. doi:10.1038/s41419-

017-0233-y

30. Jiang W, Zhao S, Jiang X, et al. The circadian clock gene Bmal1 acts as a potential anti-oncogene in
pancreatic cancer by activating the p53 tumor suppressor pathway. *Cancer Lett*. 2016;371(2):314-
325. doi:10.1016/j.canlet.2015.12.002

31. Stokes K, Nunes M, Trombley C, et al. The Circadian Clock Gene, Bmal1, Regulates Intestinal Stem
Cell Signaling and Represses Tumor Initiation. *Cell Mol Gastroenterol Hepatol*. 2021;12(5):1847-
1872.e0. doi:10.1016/j.jcmgh.2021.08.001

32. West AC, Smith L, Ray DW, Loudon ASI, Brown TM, Bechtold DA. Misalignment with the external
light environment drives metabolic and cardiac dysfunction. *Nat Commun*. 2017;8(1):417.
doi:10.1038/s41467-017-00462-2

33. Acosta-Rodríguez V, Rijo-Ferreira F, Izumo M, et al. Circadian alignment of early onset caloric
restriction promotes longevity in male C57BL/6J mice. *Science*. 2022;376(6598):1192-1202.
doi:10.1126/science.abk0297

34. Inokawa H, Umemura Y, Shimba A, et al. Chronic circadian misalignment accelerates immune
senescence and abbreviates lifespan in mice. *Sci Rep*. 2020;10. doi:10.1038/s41598-020-59541-y

35. VoPham T, Weaver MD, Vetter C, et al. Circadian Misalignment and Hepatocellular Carcinoma
Incidence in the United States. *Cancer Epidemiol Biomarkers Prev*. 2018;27(7):719-727.
doi:10.1158/1055-9965.EPI-17-1052

36. Alexander RK, Liou YH, Knudsen NH, et al. Bmal1 integrates mitochondrial metabolism and
macrophage activation. *Elife*. 2020;9:e54090. doi:10.7554/eLife.54090

37. Sahai E, Astsaturov I, Cukierman E, et al. A framework for advancing our understanding of cancer-
associated fibroblasts. *Nat Rev Cancer*. 2020;20(3):174-186. doi:10.1038/s41568-019-0238-1

38. Olumi AF, Grossfeld GD, Hayward SW, Carroll PR, Tlsty TD, Cunha GR. Carcinoma-associated

Fibroblasts Direct Tumor Progression of Initiated Human Prostatic Epithelium1. *Cancer Research*.

1999;59(19):5002-5011.

**A.**

**B.**

**A.**

| Cell Type | Gene Name | Log2 FC | Adj p value |
|---|---|---|---|
| Endothelial Cell | *Tef* | 0.326049301 | 0.001081832 |
| Fibroblast | *Dbp* | 0.606422244 | 7.90E-208 |
| Fibroblast | *Per3* | 0.574324977 | 1.74E-209 |
| Fibroblast | *Tef* | 0.395985989 | 2.46E-123 |
| Fibroblast | *Nr1d2* | 0.334812842 | 3.82E-79 |
| Fibroblast | *Hlf* | 0.333177448 | 5.96E-101 |
| Fibroblast | *Per2* | 0.270340607 | 1.22E-130 |
| Macrophage | *Dbp* | 0.425914793 | 4.73E-111 |
| Memory CD4 T Cell | *Bhlhe40* | -0.480979686 | 4.00E-22 |
| Non-Classical Monocyte | *Dbp* | 0.341766669 | 1.68E-18 |

**B.**