Multi-environmental population phenotyping suggests the higher risk of wheat

***Rht-B1b* and *Rht-D1b* cultivars in global warming scenarios**

Zihui Liu[1,3], Zunkai Hu[1,3], Zhiwei Zhu[1], Junmei Cao[2], Jialiang Zhang[1], Xiangyu Ma[1],

Xinzhong Zhang[2], Xiaoming Wang[1], Wanquan Ji[1], Shengbao Xu[1*]

[1]State Key Laboratory of Crop Stress Biology for Arid Areas, College of Agronomy,

Northwest A&F University, Yangling, Shaanxi, China.

[2]Research Institute of Grain Crops, Xinjiang Academy of Agricultural Sciences,

Urumqi, Xinjiang Uygur Autonomous Region, China.

[3]These authors contributed equally: Zihui Liu, Zunkai Hu.

* Correspondence: xushb@nwsuaf.edu.cn

ABSTRACT

After six decades practice, the semi-dwarf alleles *Rht-B1b* and *Rht-D1b* (also called

Rht-1 and *Rht-2*) had been applied into around 70% current wheat cultivars, laid the

foundation for the worldwide wheat production supply, then the agronomic traits

controlled by the two alleles still keep unclear except dwarfing wheat. Here 13

agronomic traits were investigated in 400 wheat accessions with seven environments,

uncovered the genetic effects of *Rht-B1b* and *Rht-D1b* on wheat structure and yield

traits in different genetic backgrounds and environments, and the distinct genetic

effects between *Rht-B1b* and *Rht-D1b*, suggesting that the introduction of green

revolution alleles profoundly shaped the agronomy traits of modern wheat cultivars.

The late-sowing assays and regression analysis based on the phenotypic and their

meteorological data showed the accessions containing *Rht-B1b* and *Rht-D1b* are more

sensitive to the temperature increase, and the *Rht-D1b* may lost additional 8% yield

compared the cultivars without the green revolution alleles if the average temperature

increases 1 °C. These results suggest the application of green revolution are facing

more challenges to maintain the futural wheat production supply in global warming

scenarios.

The introduction of the semi-dwarfing trait into wheat cultivars during the 1960s

and 1970s, confers great lodging resistance, allowing high fertilizer input, thereby

significantly enhance yield of crop, was a defining as the "Green Revolution" [1,2]. The

Green Revolution alleles in wheat include the *Rht-B1b* and *Rht-D1b* (also named as

Rht-1 and *Rht-2*), which introduce premature stop codons in the N-terminal coding

region of DELLA proteins to alter the GA signaling [3,4]. Approximately 70% of

cultivars in a worldwide winter wheat panel that were released in the 21st century

carry one of these two "Green Revolution" alleles [5].

After sixty years application, people found the green revolution alleles not only

function in reducing the plant height, but also involved in a broad of agronomic traits

[6,7]. However, limited by few genetic backgrounds and environmental conditions, the

agronomic traits controlled by green revolution alleles showed inconsistent

conclusions in previous studies [8-10]. In addition, two green alleles, the *Rht-B1b* and

Rht-D1b, showed minor difference in controlling wheat traits [8,11,12], which may

related their adoption to the distinct wheat region in USA [13], Europe [14,15] and China

[16,17], then the underlying difference still keeps unknown. These studies suggest the

agronomic phenotypes controlled by *Rht-B1b* and *Rht-D1b* are far from fully

dissected, which obstacles the study on underlying regulatory mechanism and its

precisely application in wheat breeding and production.

In addition, current practices had demonstrated that the wheat cultivars

containing *Rht-B1b* and *Rht-D1b* are more vulnerable to drought stress [18,19], indicating

a potential weakness of *Rht-B1b* and *Rht-D1b* in adapt to environmental stress, which

may influence the application of green revolution alleles in futural global warming

scenario. As the most adopted wheat alleles, their adaptation to high temperature has

to be evaluated to ensure wheat production security for the future.

Here, 13 agronomic traits of *Rht-B1b* and *Rht-D1b* were investigated in seven

environments across three years on 400 wheat accessions, demonstrating the global

genetic effects of *Rht-B1b* and *Rht-D1b* on wheat traits and its environmental stability,

providing the reliable phenotypes and its occurrence probability in field for the

precisely application in current and futural wheat breeding. The heat sensitivity of

green revolution alleles suggests current wheat breeding and production have to

adjusted to face the global warm scenarios.

The bread wheat population for agronomic phenotyping

A total 400 wheat accessions were collected from all over the world, and were

genotyped by RNA-seq [20] and confirmed by the KASP assay, including 105 *Rht-B1b*

(*B1b)* accessions, 98 *Rht-D1b* (*D1b*) accessions, and 197 accessions without *B1b* and

D1b (refer to *NO* thereafter) (Figure 1A). Thirteen agronomic traits were investigated

in seven environments across three years (Table S1). The correlations analysis

suggests the plant height in this population is significantly correlated with peduncle

length, total tiller number, kernel number per spike and thousand kernel weight

(Figure 1B), indicating that these traits may be affected by the introduction of green

revolution alleles.

The genetic effects of *Rht-B1b* and *Rht-D1b* on wheat plant structure and yield

The BLUE (best linear unbiased estimator) value of agronomic traits in seven

environments was used to evaluate the difference between each genotype, and 1000

permutation tests were performed on the 50% random sampling combination in each

group to evaluate genomic background independence, termed as genomic probability

(GP). We found that all the plant structure related traits are significantly different

between *NO* and dwarf group (the *B1b* and *D1b*), and also significantly different

between *B1b* and *D1b* with high GP (> 0.75 in Figure 2). Notably, the *Rht-D1b*

generally displayed a stronger effect in altering structure traits, consistent with the

previous observation that the dwarf effect from *D1b* is significantly and stably higher

than that in *B1b* [5].

However, the general significant difference can't be detected on the spike and

kernel related traits between the dwarf and *NO* group (Figure 2). Interestingly, the

D1b confers higher kernel length, kernel width and final kernel weight, result in 2.1

microgram increase for each kernel (Figure 2), equal to 6% yield increase compared

to the *NO*. In contrast, the *B1b* showed higher kernel number per spike, which result

in 4.2 kernels increase for one spike, contribute around 8% yield increase compared to

that of *NO*. The significantly difference of kernel number and kernel weight are also

observed between the cultivar group of *B1b* and *D1b* with medium GP (0.5-0.8),

highlighting that the distinct positive effect of *Rht-B1b* and *Rht-D1b* on wheat yield,

providing new insights for higher yield conferred by green revolution alleles [1,18].

Environmental variations of Green Revolution genes

To evaluate the environmental stability of the effect of green revolution alleles,

the environmental probability (EP), which present the occurrence probability with

significant difference in investigated environments, was created to describe the

environmental stability (Figure 2, Table S1). We found the effect of green revolution

alleles were stable in plant structure traits with high EP, especially *D1b*, which

showed the significant difference compared to *NO* group in more than five

environments out of all seven investigated environments, indicating that the effects of

green alleles on wheat structure traits is relative stable in different environments.

For spike related trait, only the kernel number per spike controlled by *B1b*

showed high environmental stability (>70%), while the effect of *D1b* on spike length,

thousand kernel weight and kernel width showed medium environmental stability (≥

43.9%) (Figure 2). These results suggested the roles of green revolution alleles in

kernel traits are sensitive to the environments.

Functional verification of *Rht-B1b* and *Rht-D1b*

To evaluate the pleiotropism of *B1b* and *D1b*, the GWAS with resample model

inclusion probability were performed on of the wheat traits, showing six and seven

traits were significantly associated with *B1b* and *D1b* respectively in this nature

population (Table 1, Table S2), supporting the pleiotropism of *Rht-B1b* and *Rht-D1b*

in regulating the wheat traits (Table 1). Considering different environmental response

of wheat accessions is also derived from the genomic difference, we combined the EP

and the GP, and we found the green alleles could be identified in 91% wheat traits in

GWAS if the combined probability is over 0.7 in given trait (Table 1).

The pleiotropism of green alleles was further supported by three F_2 populations

constructed by the accessions with *B1b* and *D1b* respectively (Figure 3), and support

the function of *B1b* in reducing SL (Figure 3) although the SL displayed a low

combined probability (Table 1). Moreover, F_2 population showed that the *B1b* also has

a role in enhancing TKW, but its effect is much lower than that of *D1b* (Figure 3,

Table 1).

These investigations suggest the introduction of *Rht-B1b* and *Rht-D1b*

substantially altered multitude traits in current wheat cultivars by their genetic effect.

However, the alterations conferred by green revolution alleles occur in field with

different probability for the different genomic background.

Table 1 Verification of *Rht-B1b* and *Rht-D1b* for all plant structure and yield related traits.

Traits	*Rht-B1b*					*Rht-D1b*				
	EP (%)	GP	CP (%)	PE (%)	GWASP (%)	EP (%)	GP	CP (%)	PE (%)	GWASP (%)
PH	100	1	100	-22.6*	100	100	1	100	-27.7*	100
PL	100	1	100	-19.7*	85.7	100	1	100	-27.9*	85.7
TTN	85.7	0.77	66.0	-8.65*	0	100	0.99	99.0	-19*	14.3
PTN	85.7	0.75	64.3	-8.09*	0	100	0.98	98.0	-18.3*	42.9
FLAN	71.4	0.90	64.3	-10.9*	0	100	0.98	98.0	-19.8*	85.7
SL	42.9	0.93	39.9	-0.42	0	85.7	0.92	78.8	-5.1*	0
TSN	57.1	0.57	32.5	+0.37	0	42.9	0.93	39.9	+0.69	0
FSN	57.1	0.85	48.5	+1.9*	0	71.4	0.95	67.8	+1.53*	0
SS	14.3	0.60	8.60	-1.59	0	57.1	0.61	34.8	-6.35*	0
KNS	71.4	0.99	70.7	+8.95*	100	57.1	0.75	42.8	+3.47	0
TKW	57.1	0.83	47.4	+2.59	71.4	85.7	0.91	78.0	+6.32*	57.1
KL	14.3	0.69	9.90	+0.16	57.1	71.4	0.85	60.7	+2.2*	0
KW	28.6	0.60	17.2	+1.04	71.4	85.7	0.89	76.3	+4.5*	28.6

EP, Environmental probability, presents the ratio of the number of environments
showing significant difference / all investigated environments (seven environments);

GP, Genetic probability, presents the ratio of comparison showing significantly
difference / all comparison (the 1000 permutation);

CP, Combined probability, presents the probability of EP * GP;

PE, Phenotype effect, presenting the effect of dwarf allele on the given traits;

GWASP, GWAS probability, presents the number of environments detecting dwarf
allele / all environments in GWAS analysis.

Spatiotemporal application of the Green Revolution genes in China

In our wheat population, including 87 landraces (LA) and 255 historical/modern

cultivars (MC) (Table S1). We found the application frequency of *B1b* and *D1b*

dramatically increase in MC (Figure 4A). The *D1b* was preferentially adopted by the

Chinese cultivars in the Yellow and Huai wheat zone (50.4%), while the *Rht-B1b* was

more adopted by Chinese cultivars in Yangtze River winter wheat zone and the

introduced cultivars from abroad (Table S3), consistent to that there is a distinct

environmental adoption of two green revolution genes [15,21,22]. The distinct

preferentially adoption of *B1b* and *D1b* displayed a clearer trend in the CMC released

after 2000. The total application frequency of *B1b* and *D1b* increased from 45.9% to

78.1% in CMC, which is released before and after 2000 (Figure 4B), highlighting the

green revolution alleles still are the major dwarf alleles applied in current wheat

varieties.

The heat adaptation potential of *Rht-B1b* and *Rht-D1b*

To evaluate the thermotolerance of green revolution alleles for the global

warming challenges, late-sowing field assays [23,24] were performed in three years,

which confers higher temperatures for different developmental stages of wheat

(Figure 5A, Table S5).

We found that almost investigated wheat traits significantly changed by the late-

sowing (Figure 5B, Table S6). For plant structure, the plant height was decreased

about 30%, and the tiller number were decreased more than 50% in late-sowing

condition. However, for yield traits, only the thousand kernel weight were decreased

more than 15%, indicating that the vegetative growth of wheat was more sensitive to

the late sowing (Figure 5B, Table S6). Notably, the agronomic traits of green

revolution alleles accessions showed more severe influences by the late-sowing

(Figure 5B), indicating the accessions with green revolution alleles may be more

sensitive to the temperature increase than accessions in *NO* group, especially the

accessions containing *Rht-D1b*.

The effect of green alleles in response to increasing temperature

To further evaluate the effects of high temperature on *Rht-B1b* and *Rht-D1b*, the

meteorological data, including the air temperature, sunlight hours, relative humidity

and solar radiation in seven environments was collected (Table S7, S8), then integrate

the phenotypic data from the seven environments for a multivariate regression

analysis [25,26], and we found the temperature in different developmental stage

contributes the substantial major effect to all traits of wheat population (Table S9).

Next, the average temperatures at different developmental stage were selected for

the regression analysis (Table S10). With the regression equations, we predict the

traits alterations when the temperature increases 1 °C in whole growth season,

showing the tilling number, flag leaf angle, grain weight and plant height are the most

temperature sensitive traits (Figure 6A), and the cultivars containing $B1b$ and $D1b$

were more sensitive to the increasing temperature compared to NO accessions,

generally consistent to the estimation by the late sowing (Figure 5B).

The yield per plant (YPP) will reduce around 20% if the temperature increases

1 °C (Figure 6B). Compared with NO cultivars, the YPP of cultivars containing $B1b$

and $D1b$ showed an additional 3.7% and 8.7% penalty, respectively, if the temperature

increases 1 °C (Figure 6B). Meanwhile, the late sowing in three years contribute an

average of 3.16 °C temperature increase, contribute 66.4 - 72.3% penalty in YPP

(Figure 6C), and an additional 3.2% and 5.9 % YPP penalty were observed in the

cultivars containing $B1b$ and $D1b$ respectively, which showed the similar decreased

trend with the predicted result (6.3% and 9.9% for $B1b$ and $D1b$ respectively) by the

regression analysis (Figure 6B), supporting that the cultivars containing green

revolution alleles are vulnerable to temperature increase. Next, with another

regression was performed on the YPP variations and the temperature variation based

on the same dataset, showing an additional 4.8% and 7.5% yield penalty in cultivars

containing $B1b$ and $D1b$ (Figure 6D), which are similar to the results from the

regression on the YPP and average temperature (Figure 6B). Thus, the late sowing

assay and regression analysis strongly suggest that the cultivars containing green

revolution alleles are more sensitive to the temperature increase, will cause a more

severe wheat yield penalty in the global warming scenarios.

Discussion

Wheat as one of the most important crop, provides 20% food for human being [27]. With the introduction of the dwarf genes by the Green Revolution, the yield of wheat was increased year by year [28]. Up to now, about 70% of cultivars in the world carry one of the green revolution genes [5], which laid the foundation for wheat production supply for the increasing world population. Therefore, any agronomic traits and/or stress-resistance related to the green revolution alleles may influence the global wheat production and food supply security.

In this study, we clarified multitude agronomic traits controlled by the green alleles, which closely depend on the genomic background, and their phenotypes occur in filed with different probability. This result updates our understanding about crop phenotypes in field, they are not a fixed, but display a dynamic output depend on different genomic background. The understanding and global traits may provide helpful reference for precisely applying green revolution alleles in current and futural wheat breeding and production.

Global warming has resulted in large variations in wheat yield, with expected losses reaching up to 6% for each 1 °C rise in temperature [29] and warmer regions are likely to suffer more yield loss with increasing temperature than cooler regions [30]. Compare with the preindustrial era, the global average surface temperature was about 1 °C higher [31], if worldwide carbon emissions continue at the current rate, global warming is likely to exceed 1.5 °C between 2030 and 2050, and even more than 3–5 °C at the end of the 21st century [32]. The situation is urgent to ensure the food supply with the global warming.

Generally, the higher temperature boosts the wheat development in booting, heading, anthesis, and maturity [33,34], and displaying a comprehensive alteration in wheat traits, consequent to a yield penalty (Figure 5 and Figure 6). For human, the yield is the most important target for sustainability. The yield of wheat is genetically determined by three individual components: spike number (the productive tiller number) per unit area, grain number per spike and kernel weight [35]. Although the

grain number per spike is significantly inhibited by the heat [36,37], the tiller number and

kernel weight is more sensitive to the temperature increase (Figure 5 and 6), response

to the major yield loss. Previous studies reported the productive tillers is significantly

inhibited by the early high temperature by accelerate its development [33,38]. Our data

further showed that the tiller number reduction by heat contributes a major yield

penalty, highlight the importance in maintaining the tiller number for securing wheat

yield. In addition, the plant height is also highly sensitive to the temperature increase

(Figure 5B and Figure 6A), further supporting that the biomass plays critical

foundation for the grain yield and stress resistance [18,39]. The higher biomass would be

reconsidered in the wheat breeding program. Therefore, adjusting sowing time for

bigger biomass and finding the heat-resistance QTL would be urgent mission for the

global warming challenge [39] to maintain the wheat supply. Notably, the KNS is not

sensitive to the temperature increase, implying the big spike with higher KNS may

have the higher heat tolerance to avoid the yield penalty in the global warming

scenarios, which may also be considered in current and futural wheat breeding

program.

Our results showed the cultivars containing *Rht-B1b* and *Rht-D1b* have an

increased KNS and kernel weight (Figure 2 and figure 6A), then still showed a

significant lower YPP (Figure 6B), which would result from the significant reduced

tillers number (Figure 6A). However, in wheat production, the cultivars containing

green revolution alleles substantially increased the wheat yield per unit [1,18], implying

a higher planting density were applied in cultivars containing green alleles. In this

study, we also identified a significant decrease in flag leaf angle conferred by green

revolution alleles, which would confer a more compact wheat plant structure, to allow

a higher planting density and contribute to the significant yield increase per unit area.

The higher planting density provides a plausible explanation for the yield increase of

cultivars containing green revolution [40-42]. However, these phenotypes controlled by

green revolution alleles are sensitive to the temperature increase (Figure 5). The tiller

number is the most heat sensitive traits for wheat yield components, showing a

dramatic decrease with the temperature increase. To maintain the yield, the reduced

tiller number require higher planting density for compensation. However, the flag leaf

angle is also significantly increased with the temperature increase (Figure 5), which

would prevent the higher planting density practices. Thus, the dilemma between tiller

number and flag leaf angle in high temperature has to be resolved in futural wheat

breeding. For the cultivars containing green alleles, especially containing *Rht-D1b*,

the dilemma more prominent because both tiller number and flag leaf angle getting

worse with the temperature increase, compared to the cultivars without green

revolution alleles, which may further limit the production of cultivars with green

alleles in the global warming scenarios.

Climate change may outpace current wheat breeding yield improvements [30] , and

more recently released winter wheat varieties were less able to resist high-temperature

extremes than older varieties [43], we speculate that the increased frequency of the green

revolution alleles in current wheat cultivars (Figure 4B) may partly response to the

decreased heat resistance of current cultivars. To ensure the food security, we may

consider more dwarf alleles for futural substitution, or improve the thermotolerant

defect of green revolution alleles in time.

Methods

Plant materials and growing conditions

400 accessions of bread wheat were collected in this study, including modern

cultivars and landraces from worldwide wheat accessions (Table S1). All materials

were planted in seven environments (E1--E7) at Yangling (34°28′N, 108°07′E,

altitude 517 m) and Chongzhou (30°63′N, 103°67′E, altitude 1300 m) of China in

2018-2021 (Table S11). Field experiments were arranged in a randomized complete

block design with each variety planted in three replicates. Each accession was sown

with 1 m in length and a row spacing of 20 cm, with 10 seeds per row. Field

management was consistent with local practices with watering for wheat production.

Three F_2 populations (Zhongluo08-3 × Ningmai12, Yannong22 × Qingfeng1 and

MY72 × Changwei18) were planted at the Jize county of Hebei province (36°90′N,

114°80′E, altitude 42 m) in 2021-2022 growth season. Field management was

consistent with local practices for wheat production.

Agronomic traits measurements

The traits of plant structure and yield were measured at physiological maturity.

Three plant individuals selected from each row were used to investigate plant height

(PH), peduncle length (PL), total tiller number (TTN), productive tiller number (PTN),

flag leaf angle (FLAN), spike length (SL), total spikelet number (TSN), fertile

spikelet number (FSN) and kernel number per spike (KNS) according to established

protocols [8,44-46]. Days to heading was assessed as the interval between the date of

seeding emergence and the date at which 50% of spikes per row emerged from the

flag leaf, days to flowering was assessed as the interval between the date of seeding

emergence and the date at which 50% of spikes per row have extruded at least one

anther [47]. Further, we calculated spikelet spacing (SS, SL/TSN) and yield per plant

(YPP, PTN×KNS×TKW). In addition, thousand kernel weight (TKW), kernel length

(KL) and kernel width (KW) were recorded using the rapid SC-G grain appearance

quality image analysis system developed by Hangzhou WSeen Detection Technology

Co, Ltd, China [45].

DNA extraction and genotyping

Genomic DNA was extracted from wheat flag leaves of heading stage plants

following the modified Cetyl Trimethylammonium Bromide (CTAB) method for

genotyping [48]. Kompetitive allele-specific PCR (KASP) markers were developed by

the mutation site of *Rht-B1b* and *Rht-D1b* to identify all the accessions [49]. All primers

were listed in Supplementary Table S12. The KASP assay was performed as following:

assays were set up as 5 μL reactions containing 2.2 μL template DNA dried to assay

plate, 2.5 μL of V4.0 2x Kaspar mix (LGC Group, Teddington, UK), 2.444 μL ddH$_2$O

and 0.056 μL primer mix; and the PCR cycling included hotstart at 94 °C for 15 min,

followed by 10 touchdown cycles (94 °C for 20 s; touchdown 61°C, -0.6 °C per cycle,

1 min), then followed by 35 cycles of amplification (94 °C for 20 s; 55 °C for 1 min).

Genome-wide association study

The GWAS were performed on the SNPs derived population RNA-seq [20] with

resample model inclusion probability [50]. In details, 60% of each group were randomly

selected without replacement and forward regression was performed, and repeated

100 times. The SNPs that were selected in the regression model in five or more

subsamples were considered significant (RMIP $\geq$ 0.05) in this study. GWAS was

implemented with the GAPIT version 3 in R (v3.6.1), using the FarmCPU [51] on each

bootstrapped dataset. A Bonferroni-corrected threshold probability based on

individual tests was calculated to correct for multiple comparisons using 1/N, where

N is the effective SNP number calculated with the Genetic type 1 Error Calculator

(v0.2) [52]. The significant *P*-value thresholds for plant height and peduncle length were

8.58×10^{-4} in this population, but a more severe value $P < 1 \times 10^{-4}$ was selected to

declare marker-trait associations (MTAs) in this study. To identify the traits that *Rht-*

B1b or *Rht-D1b* are involved in regulating, those detected in at least two

environments were considered to be stable identification.

Statistical analysis

The best linear unbiased estimate (BLUE) values for phenotypic under seven

environments were obtained using a mixed linear model by genotype as a fixed effect

and replicates as random effects. Analysis of variance (ANOVA) was performed to

test the homogeneity of genotype and phenotypic based on linear mixed model using

the Python SciPy module. Differences between phenotypic of different genotypes

were tested by *t-test* [53]. If there is homogeneity of variance between the two sets of

data, the default argument to the function *stats.ttest_ind* is used; otherwise, the

parameter *equal_var* is set to *False*. All analyses were conducted using *lme4* (Bates

et al., 2015) and *lsmeains* packages [54] in R3.6.2 [55].

Permutation test

To reduce the interferes by co-selected SNPs and/or genes, the permutation test [56]

was performed on each genotype pairs to evaluate the effect of different genomic

background. The wheat population was divided into three groups by genotypes (*NO*,

B1b, *D1b*), 50% of the accessions were randomly selected to test the significance

between different groups. 1000 permutations were performed to examine the

occurrence probability with significant differences between two groups.

Multiple linear regression analysis of climate factors

Weather data were acquired from the China Meteorological Data Service Center

(https://data.cma.cn). We collected 5 climate variables, including the maximum and

minimum temperatures, sunlight hours, relative humidity and the solar radiation in

seven environments. According to the heading and flowering time of each accession

and 5 climate variables, the average temperature, the average sunlight hours, the

average relative humidity and the average solar radiation at seeding stage, flowering

stage and filling stage for each accession were collected according their

developmental process (Table S7). Multiple regression analysis was used to examine

the effect of meteorological factors on phenotypes [25]. The equation was:

$Y=a+bX_1+cX_2+dX_3+eX_4+fX_5+gX_6+hX_7+iX_8+jX_9+kX_{10}+lX_{11}+mX_{12}$

Where Y represents the phenotypic data. X_1, X_2, X_3, X_4, X_5, X_6, X_7, X_8, X_9,

X_{10}, X_{11}, X_{12}, indicate the average temperature, the average sunlight hours, the

average relative humidity and the average solar radiation at seeding stage, flowering

stage and filling stage, respectively. b, c, d, e, f, g, h, i, j, k, l, m are the regression

coefficients, and a is a constant. The Ordinary least square (*OLS*) was used for

equation fitting, $P < 0.05$.

The *step* function of R3.6.2 was used for step-based regression, and the

eliminated variables were selected according to Akaike Information Criterion (AIC).

The meteorological factors with the smallest values from AIC were always removed

until the equation was constructed successfully. The calculation formula of AIC:

$AIC=2k-2\ln(L)$. L is the maximum likelihood function of the regression model. K is

the number of parameters. The AIC of the model obtained by eliminating different

variables were compared, the eliminated variables were selected in turn, and the final

retained variables were ranked in importance from small to large according to AIC.

The *vif* function of R3.6.2 was used to test whether the model has multicollinearity.

The Variance Inflation Factor (VIF) was used to measure the severity of

multicollinearity in multiple linear regression model, VIF<10, with no

multicollinearity; $10 \leq$ VIF<100, strong multicollinearity; VIF $\geq$ 100, severe

multicollinearity. The error rate of multiple linear regression model was calculated by

sigma function in R3.6.2.

Multiple linear regression analysis of average temperature

Multiple regression analysis was used to examine the effect of average

temperature at different stage on phenotypes. The equation was:

$$Y=m_1+m_2Xa+m_3Xb+m_4Xc$$

Where Y represents the phenotypic data. Xa, Xb, Xc represents the average

temperature at seeding stage, flowering stage and filling stage, respectively. m_2, m_3,

m_4 are the regression coefficients, and m_1 is a constant. The traits were predicted by

1 °C increase in the average temperature of all developmental stage.

The regression between YPP variations and the temperature variation. The

variation of YPP and temperature can be obtained by subtracting BLUE in each

environment. Based on the regression equation, the YPP was predicted by 1 °C

increase in the average temperature of all developmental stage.

Data availability

All data, models, or code generated or used during the study are available from

the corresponding author by request. The phenotypes and SNP data are shown in

https://iwheat.net/resource/. Meanwhile this data will keeps update every year, and we

welcome more team adopt this population, phenotypes and meteorologic data, and

may collect more information when we get more responses from users.

References

1.   Hedden, P. The genes of the Green Revolution. *Trends Genet* **19**, 5-9 (2003).

2.   Peng, J. *et al.* 'Green revolution' genes encode mutant gibberellin response modulators. *Nature*
400, 256-61 (1999).

3.   Wu, J. *et al.* Dominant and pleiotropic effects of a GAI gene in wheat results from a lack of
interaction between DELLA and GID1. *Plant Physiol* **157**, 2120-30 (2011).

4.   Liu, Q., Wu, K., Harberd, N.P. & Fu, X. Green Revolution DELLAs: From translational
reinitiation to future sustainable agriculture. *Mol Plant* **14**, 547-549 (2021).

5.   Wurschum, T., Langer, S.M., Longin, C.F.H., Tucker, M.R. & Leiser, W.L. A modern Green
Revolution gene for reduced height in wheat. *Plant J* **92**, 892-903 (2017).

6.   Lanning, S.P., Martin, J.M., Stougaard, R.N., Guillen-Portal, F.R. & Talbert, L.E. Evaluation
of Near-Isogenic Lines for Three Height-Reducing Genes in Hard Red Spring Wheat. *Crop*
Science **52**, 1145-1152 (2012).

7.   Sherman, J.D. *et al.* Genetics of End-Use Quality Differences between a Modern and
Historical Spring Wheat. *Crop Science* **54**, 1972-1980 (2014).

8.   Jobson, E.M., Johnston, R.E., Oiestad, A.J., Martin, J.M. & Giroux, M.J. The Impact of the
Wheat Rht-B1b Semi-Dwarfing Allele on Photosynthesis and Seed Development Under Field
Conditions. *Front Plant Sci* **10**, 51 (2019).

9.   Würschum, T., Langer, S.M. & Longin, C.F.H. Genetic control of plant height in European
winter wheat cultivars. *Theoretical and Applied Genetics* **128**, 865-874 (2015).

10. Youssefian, S., Kirby, E. & Gale, M. Pleiotropic effects of the GA-insensitive Rht dwarfing
genes in wheat. 1. Effects on development of the ear, stem and leaves. *Field Crops Research*
28, 179-190 (1992).

11. Bush, M.G. & Evans, L.T. Growth and development in tall and dwarf isogenic lines of spring
wheat. *Field Crops Research* **18**, 243-270 (1988).

12. Velu, G., Singh, R.P., Huerta, J. & Guzmán, C. Genetic impact of Rht dwarfing genes on grain
micronutrients concentration in wheat. *Field Crops Res* **214**, 373-377 (2017).

13. Guedira, M. *et al.* Distribution of Rht Genes in Modern and Historic Winter Wheat Cultivars
from the Eastern and Central USA. *Crop Science* **50**, 1811-1822 (2010).

14. Wurschum, T. *et al.* Exploiting the Rht portfolio for hybrid wheat breeding. *Theor Appl Genet*
131, 1433-1442 (2018).

15. Wurschum, T., Langer, S.M. & Longin, C.F. Genetic control of plant height in European
winter wheat cultivars. *Theor Appl Genet* **128**, 865-74 (2015).

16. Gao, Z., Shi, Z., Zhang, A. & Guo, J. Distribution of genes associated with yield potential and
water-saving in Chinese Zone II wheat detected by developed functional markers. *J Genet* **94**,
35-42 (2015).

17. Zhang, X., Yang, S., Zhou, Y., He, Z. & Xia, X. Distribution of the *Rht-B1b*, *Rht-D1b* and
Rht8 reduced height genes in autumn-sown Chinese wheats detected by molecular markers.
Euphytica **152**, 109-116 (2006).

18. Jatayev, S. *et al.* Green revolution 'stumbles' in a dry environment: Dwarf wheat with Rht
genes fails to produce higher grain yield than taller plants under drought. *Plant Cell Environ*
43, 2355-2364 (2020).

19. Monneveux, P., Jing, R.L. & Misra, S.C. Phenotyping for drought adaptation in wheat using
physiological traits. *Frontiers in Physiology* **3**, 429 (2012).

20. Wang, X. *et al.* Population Transcriptome and Phenotype Reveal that the *Rht-D1b* Contribute

a Larger Seedling Roots to Modern Wheat Cultivars. *bioRxiv*, 2022.06.02.494553 (2022).

21.   Grogan, S.M. *et al.* Allelic Variation in Developmental Genes and Effects on Winter Wheat
436   Heading Date in the U.S. Great Plains. *PLoS One* **11**, e0152852 (2016).

437   22.   Zhao, J. *et al.* Global status of 47 major wheat loci controlling yield, quality, adaptation and
438   stress resistance selected over the last century. *BMC Plant Biol* **19**, 5 (2019).

439   23.   Chauhan, S., Srivalli, S., Nautiyal, A.R. & Khanna-Chopra, R. Wheat cultivars differing in
440   heat tolerance show a differential response to monocarpic senescence under high-temperature
441   stress and the involvement of serine proteases. *Photosynthetica* **47**, 536-547 (2009).

442   24.   Tiwari, C. *et al.* Exploring the possibility of obtaining terminal heat tolerance in a doubled
443   haploid population of spring wheat (*Triticum aestivum* L.) in the eastern Gangetic plains of
444   India. *Field Crops Research* **135**, 1-9 (2012).

445   25.   Tao, Z., Wang, H., Dai, J., Alatalo, J. & Ge, Q. Modeling spatiotemporal variations in leaf
446   coloring date of three tree species across China. *Agricultural and Forest Meteorology* **249**,
447   310-318 (2018).

448   26.   Chmielewski, F.M. & Potts, J.M. The relationship between crop yields from an experiment in
449   southern England and long-term climate variations. *Agricultural & Forest Meteorology* **73**,
450   43-66 (1995).

451   27.   Shewry, P.R. & Hey, S.J. The contribution of wheat to human diet and health. *Food Energy
452   Secur* **4**, 178-202 (2015).

453   28.   Gao, F. *et al.* Genetic Progress in Grain Yield and Physiological Traits in Chinese Wheat
454   Cultivars of Southern Yellow and Huai Valley since 1950. *Crop Science* **57**, 760-773 (2017).

455   29.   Liu, B. *et al.* Similar estimates of temperature impacts on global wheat yield by three
456   independent methods. *Nature Climate Change* **6**, 1130-1136 (2016).

457   30.   Zhang, T. *et al.* Climate change may outpace current wheat breeding yield improvements in
458   North America. *Nat Commun* **13**, 1-9 (2022).

459   31.   Zhao, C. *et al.* Temperature increase reduces global yields of major crops in four independent
460   estimates. *Proc Natl Acad Sci U S A* **114**, 9326-9331 (2017).

461   32.   Arias, P. *et al.* Climate Change 2021: The Physical Science Basis. Contribution of Working
462   Group14 I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change;
463   Technical Summary. (2021).

464   33.   Rahman, M.A., Chikushi, J., Yoshida, S. & Karim, A. Growth and yield components of wheat
465   genotypes exposed to high temperature stress under control environment. *Bangladesh Journal
466   of Agricultural Research* **34**, 360-372 (2009).

467   34.   Lian, X. *et al.* Summer soil drying exacerbated by earlier spring greening of northern
468   vegetation. *Sci Adv* **6**, eaax0255 (2020).

469   35.   Xiao, J. *et al.* Wheat genomic study for genetic improvement of traits in China. *Sci China Life
470   Sci* **65**, 1718-1775 (2022).

471   36.   Shpiler, L. & Blum, A. Differential reaction of wheat cultivars to hot environments. *Euphytica*
472   **35**, 483-492 (1986).

473   37.   Dias, A.S. & Lidon, F.C. Evaluation of Grain Filling Rate and Duration in Bread and Durum
474   Wheat, under Heat Stress after Anthesis. *Journal of Agronomy and Crop Science* **195**, 137-147
475   (2009).

476   38.   McMaster, G.S. Phenology, development, and growth of the wheat (*Triticum aestivum* L.)
477   shoot apex: a review. *Advances in agronomy* (1997).

39. Langridge, P. & Reynolds, M. Breeding for drought and heat tolerance in wheat. *Theor Appl*
Genet **134**, 1753-1769 (2021).
40. Evenson, R.E. & Gollin, D. Assessing the impact of the Green Revolution, 1960 to 2000.
science **300**, 758-762 (2003).
41. Naruoka, Y. *et al.* Identification of quantitative trait loci for productive tiller number and its
relationship to agronomic traits in spring wheat. *Theor Appl Genet* **123**, 1043-53 (2011).
42. Sinclair, T.R. & Sheehy, J.E. Erect leaves and photosynthesis in rice. *Science* **283**, 1455-1455
(1999).
43. Tack, J., Barkley, A. & Nalley, L.L. Effect of warming temperatures on US wheat yields. *Proc*
Natl Acad Sci U S A **112**, 6931-6 (2015).
44. Fan, X. *et al.* QTLs for flag leaf size and their influence on yield-related traits in wheat
(*Triticum aestivum* L.). *Molecular Breeding* **35**, 24 (2015).
45. Guan, P. *et al.* Use of near-isogenic lines to precisely map and validate a major QTL for grain
weight on chromosome 4AL in bread wheat (*Triticum aestivum* L.). *Theor Appl Genet* **132**,
2367-2379 (2019).
46. Sun, C. *et al.* Genome-wide association study for 13 agronomic traits reveals distribution of
superior alleles in bread wheat from the Yellow and Huai Valley of China. *Plant Biotechnol J*
15, 953-969 (2017).
47. Li, L. *et al.* Genetic dissection of drought and heat-responsive agronomic traits in wheat. *Plant*
Cell Environ **42**, 2540-2553 (2019).
48. Wehner, G.G., Balko, C.C., Enders, M.M., Humbeck, K.K. & Ordon, F.F. Identification of
genomic regions involved in tolerance to drought stress and drought stress induced leaf
senescence in juvenile barley. *BMC Plant Biol* **15**, 1-15 (2015).
49. Ma, J. *et al.* Identification and validation of a major and stably expressed QTL for spikelet
number per spike in bread wheat. *Theor Appl Genet* **132**, 3155-3167 (2019).
50. Hirsch, C.N. *et al.* Insights into the effects of long-term artificial selection on seed size in
maize. *Genetics* **198**, 409-21 (2014).
51. Liu, X., Huang, M., Fan, B., Buckler, E.S. & Zhang, Z. Iterative Usage of Fixed and Random
Effect Models for Powerful and Efficient Genome-Wide Association Studies. *PLoS Genet* **12**,
e1005767 (2016).
52. Wu, J. *et al.* A large-scale genomic association analysis identifies the candidate causal genes
conferring stripe rust resistance under multiple field environments. *Plant Biotechnol J* **19**,
177-191 (2021).
53. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python.
Nature methods **17**, 261-272 (2020).
54. Lenth, R.V. Least-Squares Means: The R Package lsmeans. *Journal of Statistical Software* **69**,
1-33 (2016).
55. Bates, D.M., Maechler, M., Bolker, B.M. & Walker, S.C. Package lme4: linear mixed-effects
models using Eigen and S4. *Journal of statistical software* **67**(2014).
56. Zhou, Y. *et al.* Triticum population sequencing provides insights into wheat adaptation. *Nat*
Genet **52**, 1412-1422 (2020).
519

Acknowledgements

This work was supported by "Integration of Two Chains" Key Research and
Development Projects of Shaanxi Province "Wheat Seed Industry Innovation Project";
grants from the National Natural Science Foundation of China [31571756 and
31501380] and Key R&D Program of Yangling Seed Industry Innovation Center
(Ylzy-xm-01). We thank Guoyue Chen and Zhien Pu for wheat field investigation in
Sichuan Province, and thank Yufeng Zhang for wheat hybrid group field investigation
in Hebei Province, and thank Bing Liu for the valuable comments to this study.

Author contributions

S.X. designed the concept and experiments. Z.L., J.Z. and X.M. performed field
investigation. J.C and X.Z performed the related germplasm identification. Z.L., Z.H.,
Z.Z., X.W. W.J., and S.X. analyzed the data and wrote the manuscript. All the authors
were involved in the revision of the manuscript and approved the final manuscript.

Competing interests

The authors declare no competing interests.
535

Figures

Figure. 1 The wheat population and agronomic traits. (A) The number of wheat

accessions with different genotypes in the population. *NO*, wheat accessions without

Rht-B1b and *Rht-D1b*; *B1b*, wheat accessions containing *Rht-B1b*; *D1b*, wheat

accessions containing *Rht-D1b*. The number indicates the accessions number in each

genotype. (B) The correlations analysis between different agronomic traits by the best

linear unbiased estimate (BLUE) values. PH, plant height; PL, peduncle length; TTN,

total tiller number; PTN, productive tiller number; FLAN, flag leaf angle; SL, spike

length; TSN, total spikelet number; FSN, fertile spikelet number; SS, spikelet spacing;

KL, kernel length; KW, kernel width; KNS, kernel number per spike; TKW, thousand

kernel weight. Red lines mean positive correlation and gray lines mean negative

correlation. The correlations were analyzed by the "corrplot" in the R package using

the BLUE values in Table S1.

Figure. 2 The function of the *Rht-B1b* and *Rht-D1b* on wheat traits. The *Rht-B1b* and

Rht-D1b were compared to that of *NO* accessions, the letters at the top of violin plots

indicate the significant difference between each group (two-sided Dunn's Kruskal–

Wallis test, $P < 0.01$). The permutation tests (1000) were performed on the 50%

random samples combination in each group, the numbers represent the ratio of

comparison showing significantly difference / all comparison (1000). The percentages

in parentheses indicate environmental probability, the number of environments that

have significant difference effect / the total number of environments (7). The data

come from the BLUE values (n=400) of seven environments, as shown in Table S1.

Figure 3. Validation on the genetic role of the *Rht-B1b* and *Rht-D1b* in three F2
populations. The parents are Zhongluo08-3 (*Rht-D1b*); Ningmai12 (*Rht-B1b*),
Yannong22 (*Rht-B1b*); Qingfeng1 (*Rht-D1b*); MY72 (*Rht-B1b*) and Changwei18
(*Rht-D1b*), respectively. Different letters indicate statistical significance, determined
using two-sided Dunn's Kruskal–Wallis test ($P < 0.05$). *B1aD1a* refers *Rht-B1a* &
Rht-D1a; *B1bD1a* refers *Rht-B1b* & *Rht-D1a*; *B1aD1b* refers *Rht-B1a* & *Rht-D1b*;
B1bD1b refers *Rht-B1b* & *Rht-D1b*.

Figure. 4 The spatiotemporal distribution of the *Rht-B1b* and *Rht-D1b*. (A) The distribution of the *Rht-B1b* and *Rht-D1b*. LA, landraces; Y&H, Yellow and Huai winter wheat zone; YTS, Yangtze River winter wheat zone; OTW, other wheat zone in China; IC, Introduced worldwide cultivars to China. The numbers in the bar indicate the number of wheat accessions, respectively. (B) The application frequency of the *Rht-B1b* and *Rht-D1b* before and after 2000. B-2000, the cultivars was released before 2000; P-2000, the cultivars was released post 2000.

Figure. 5 The phenotypic variations with late-sowing. (A) The temperature variations with late-sowing. N, normal-sowing; L, late-sowing. The sowing date and temperature was shown in Table S4 and S5. (B) The phenotypic alteration with late-sowing. The data is the phenotypic data in late-sowing / phenotypic data in normal-sowing. Different letters on the bars indicate the significant differences ($P < 0.05$, two-sided Dunn's Kruskal–Wallis test), as shown in Table S6. The gray dash lines indicate the value 1.

Figure. 6 The phenotypes of the *Rht-B1b* and *Rht-D1b* in response to temperature increase. (A) The alterations of wheat traits when the temperature increased 1°C. Normal, the normal temperature of wheat growth; +1°C indicates the average temperature increase 1°C of all developmental stage. The number indicates the phenotype calculated from the regression equation with normal temperatures; the percentages indicate the phenotypic changed ratio of corresponding genotype wheat accessions when the average temperature increased 1°C. (B) The predicted yield per plant (YPP) with temperature increase by regression analysis. The YPP was estimated by productive tiller number × kernel number per spike × kernel weight. 3.16°C indicates the average temperature increase of late-sowing compared to normal-sowing. (C) The YPP with normal-sowing and late-sowing. The number indicates the YPP of normal-sowing, the percentages indicate the YPP changed ratio in late sowing compared with normal sowing. NS, normal-sowing; LS, late-sowing. (D) The variations of YPP according to the temperature variations. The percentages indicate the YPP changed ratio of corresponding genotype wheat accessions when the average temperature increased 1°C and 3.16°C.

Additional information

Supplemental Table 1. The composition and phenotypes of 400 wheat accessions in seven environments.

Supplemental Table 2. GWAS of 13 wheat traits with RMIP (≥5).

Supplemental Table 3. The distributions of wheat accessions in different agroecological regions.

Supplemental Table 4. The duration days of all accessions at different stage.

Supplemental Table 5. The average temperature of different developmental stage in normal-sowing and late-sowing.

Supplemental Table 6. The alterations of wheat traits with late-sowing.

Supplemental Table 7. The meteorological data for all wheat accessions.

Supplemental Table 8. The heading and flowering time of all wheat accessions.

Supplemental Table 9. The multiple linear regression equations with multitude meteorological factors.

Supplemental Table 10. The multiple linear regression equations with average temperatures.

Supplemental Table 11. The filed investigation in seven environments.

Supplemental Table 12. The KASP primer sequence.