Multi-environmental population phenotyping suggest the higher risks of wheat

Rht-B1b **and** ***Rht-D1b*** **cultivars in global warming scenarios**

Zihui Liu[1,3], Zunkai Hu[1,3], Xiangjun Lai[1], Junmei Cao[2], Jialiang Zhang[1], Xiangyu Ma[1],

Xinzhong Zhang[2], Xiaoming Wang[1], Wanquan Ji[1], Shengbao Xu[1*]

[1]State Key Laboratory of Crop Stress Biology for Arid Areas, College of Agronomy,

Northwest A&F University, Yangling, Shaanxi, China.

[2]Research Institute of Grain Crops, Xinjiang Academy of Agricultural Sciences,

Urumqi, Xinjiang Uygur Autonomous Region, China.

[3]These authors contributed equally: Zihui Liu, Zunkai Hu.

* Correspondence: xushb@nwsuaf.edu.cn

Short Title: The filed phenotyping of *Rht-B1b* and *Rht-D1b*

One-sentence Summary: The *Rht-B1b* and *Rht-D1b* controls broadly wheat

agronomic traits, which are vulnerable to temperature increase.

The authors responsible for distribution of materials integral to the findings presented

in this article in accordance with the policy described in the Instructions for Authors

(www.plantcell.org) is: Shengbao Xu (xushb@nwsuaf.edu.cn).

The E-mail address of authors:

Zihui Liu, lzh@nwsuaf.edu.cn

Zunkai Hu, huzunkai@nwafu.edu.cn

Xiangjun Lai, lxj2820@nwafu.edu.cn

Junmei Cao, caojm0508@126.com

Jialiang Zhang, nxyzjl@nwafu.edu.cn

Xiangyu Ma, maxiangyu@nwafu.edu.cn

Xinzhong Zhang, zxinzhong1031@163.com

Xiaoming Wang, wangxm@nwsuaf.edu.cn

Wanquan Ji, jiwanquan2003@126.com

Shengbao Xu, xushb@nwsuaf.edu.cn

ABSTRACT

After six decades practice, the semi-dwarf alleles *Rht-B1b* and *Rht-D1b*, (also called *Rht-1* and *Rht-2*) had been applied into more than 70% current wheat cultivars, laid the foundation for the worldwide wheat production supply, then the phenotypes of the two alleles still keeps unclear. Here 400 wheat accessions, including 106 *Rht-B1b* accessions, 99 *Rht-D1b* accessions, were systematic investigated on 16 agronomic traits in seven environments, uncover the effects of two well-known alleles on wheat structure and yield traits in different genetic backgrounds and ecosystems, providing important reference for further application in different genetic and environmental conditions. The regression analysis based on the phenotypic and their meteorological data showed the temperature increase at seedling stage severely reduce the tiller number and result in huge yield penalty per plant. The accession containing *Rht-D1b* may lost about 23.5% yield if the temperature increases 1□ in average. This population phenotyping provides the global phenotypes to dissect the molecular mechanism of *Rht-B1b*, *Rht-D1b* and GA signaling in wheat development and environmental adaptation. This study recommends a new approach to get the global phenotypes conferred by a specific allele with natural environmental conditions, which benefits the data from field to lab.

INTRODUCTION

The introduction of the semi-dwarfing trait into wheat cultivars during the 1960s and 1970s, confers great lodging resistance, allowing high fertilizer input, thereby significantly enhance yield of wheat, was a defining as the "Green Revolution" (Peng et al., 1999; Hedden, 2003). Approximately 70% of cultivars in a worldwide winter wheat panel that were released in the 21st century carry one of these two "Green Revolution" alleles (Wurschum et al., 2017).

There are two Rht (Reduced Height) alleles, *Rht-B1b* and *Rht-D1b* (also named as *Rht1* and *Rht2*), which introduce premature stop codons in the N-terminal coding region of DELLA proteins to alter the GA signaling (Wu et al., 2011; Liu et al., 2021). Accumulating data showed the Gibberellins interacts with multiple phytohormones, such as auxin (Löfke et al., 2013), strigolactones (Marzec, 2017; Liao et al., 2019), brassinosteroids (Ross and Quittenden, 2016) and ethylene (Qin et al., 2022), to plays broadly role in regulating plant development and environmental adaptation (Claeys et al., 2014). However, the GA synthesis, transportation and regulatory role is spatiotemporally dependent (Topham et al., 2017; Binenbaum et al., 2018; Van De Velde et al., 2021; Qin et al., 2022), challenges the functional study of *Rht-B1b* and *Rht-D1b*. Recent study suggests the *Rht-B1b* and *Rht-D1b* produce N-terminal truncated proteins through translational re-initiation, which may confer new function beyond of GA signaling (Van De Velde et al., 2021). All these prevent our understanding on the regulatory mechanism of the green alleles in wheat.

On the other side, after sixty years application the green alleles were found implication in a broad agronomic traits in addition to reducing the plant height (Lanning et al., 2012; Sherman et al., 2014). However, limited by few genetic backgrounds and environmental conditions in previous studies, the agronomic traits controlled by green alleles always showed inconsistent conclusion (Jobson et al., 2019). In addition, two green alleles, the *Rht-B1b* and *Rht-D1b*, showed minor difference in controlling wheat traits (Bush and Evans, 1988; Velu et al., 2017; Jobson et al., 2019), which may related their adoption to the distinct wheat region in USA

(Guedira et al., 2010), Europe (Wurschum et al., 2015; Wurschum et al., 2018) and

China (Zhang et al., 2006; Gao et al., 2015), then the underlying mechanism still

keeps unknown. These observations suggested the agronomic phenotypes of *Rht-B1b*

and *Rht-D1b* are far from fully dissected, which prevent the study on underlying

regulatory mechanism and its precisely application in wheat breeding and production.

Current practices had demonstrated that the wheat cultivars containing *Rht-B1b*

and *Rht-D1b* are more vulnerable to drought stress (Monneveux et al., 2012; Jatayev

et al., 2020), indicating a potential weakness of *Rht-B1b* and *Rht-D1b* in adapt to

environmental stress, which may influence the green alleles application in futural

global warming scenario. As the most adopted wheat alleles, their adaptation to

temperature increase has to be evaluated to ensure wheat production security for the

future.

Here, 16 agronomic traits of *Rht-B1b* and *Rht-D1b* were investigated in seven

environments across three years on 400 wheat accessions, demonstrating the global

effects of *Rht-B1b* and *Rht-D1b* on wheat traits and its environmental stability,

providing the global phenotypes for studying the molecular mechanism of *Rht-B1b*

and *Rht-D1b* in wheat development and environmental adaptation, and the potential

risk for futural wheat breeding and production.

RESULTS

The bread wheat population for agronomic phenotyping

A total 400 wheat accessions were collected from all over the world, and were

genotyped by KASP assay, including 105 *Rht-B1b* (*B1b*) accessions, 98 *Rht-D1b*

(*D1b*) accessions, and 197 accessions without the two alleles (refer to *NO* thereafter)

(Figure 1A). Sixteen agronomic traits on plant structure and yield components were

investigated in seven environments across three years (Table S1). The correlations

analysis suggests the plant height in this population is significantly correlated with

peduncle length, total tiller number, flag leaf length, flag leaf width, kernel number

per spike and thousand kernel weight (Figure 1B), indicating that these traits may be

affected by the dwarf gene introduction.

108   The effects of *Rht-B1b* and *Rht-D1b* introduction on wheat plant structure and

109   yield

To acquire the phenotypes of *Rht-B1b* and *Rht-D1b*, the BLUE (best linear

unbiased estimator) value of the agronomic traits in seven environments were

compared to that of *NO* accessions. Meanwhile, 1000 permutation tests were

performed on the 50% random samples combination in each group to evaluate genetic

background dependence. We found that all the plant structure related traits are

significantly altered by the green alleles (Figure 2A) with low genetic background

dependences (> 800 times significantly difference out of 1000 times test). Meanwhile,

all the traits also showed significantly difference and low genetic background

dependence between the *Rht-B1b* and *Rht-D1b* accessions, except the trait flag leaf

area, which showed significantly difference between *Rht-B1b* and *Rht-D1b* but this

difference is medium dependent on the genetic background (540 significant difference

in 1000 permutation test). Moreover, the *Rht-D1b* generally displayed a stronger

effect in altering structure traits, consistent with the previous observation that the

dwarf effect from *Rht-D1b* is significantly and stably higher than that in *Rht-B1b*

(Wurschum et al., 2017).

However, the general significant difference can't be detected on the yield related

traits between the dwarf and *NO* group (Figure 2B). The *Rht-D1b* has a role in

reducing the spike length and the spikelet spacing but still maintain the similar

spikelet number, suggesting a compact spike structure in the wheat cultivars

containing *Rht-D1b*. Meanwhile, the *Rht-D1b* confers higher kernel length, kernel

width and final kernel weight, result in 2.1 microgram increase for each kernel

(Figure. 2B), equal to 6% yield increase. In contrast, the *Rht-B1b* showed higher

fertile spikelet number and kernel number per spike, which result in 4.2 kernels

increase per spike, contribute 8% yield increase compared to that of *NO*. The

significantly difference of kernel number and kernel weight are also observed between

the group of *Rht-B1b* and *Rht-D1b* with medium genetic background dependence,

highlighting that the distinct positive effect of *Rht-B1b* and *Rht-D1b* on the wheat

yield, providing new insights for higher yield conferred by green alleles (Hedden,

2003; Jatayev et al., 2020).

Environmental variations of Green Revolution genes

To evaluate the environmental stability of the green alleles, the difference of

traits was further compared in different environment (Figure 2, Table. S1). We found

the effect of green alleles were stable on plant structure traits in all environments,

especially *Rht-D1b*, which showed the significant difference in five environments at

least out of seven environments compared to *NO* group, indicating that the effects of

green alleles on wheat structure traits is stable in different environments.

For yield related trait, only the kernel number per spike controlled by *Rht-B1b*

showed high environmental stability (>70%), while the effect of *Rht-D1b* on spike

length, thousand kernel weight and kernel width showed medium environmental

stability (≥43.9%) (Figure 2). These results suggested the effects of green alleles on

yield traits tend to be affected by the environments.

Spatiotemporal application of the Green Revolution genes in China

In our wheat population, including 87 landraces (LA) and 255 historical/modern

cultivars (MC) (Table. S1), which contains 30 Chinese landraces and 162 Chinese

modern cultivars (CMC) (after 1970s) (Figure 3A). In China, which have no accession in

LA contains the Green Revolution genes, while more than 67% accessions in CMC

contains one of Green Revolution alleles. The higher frequency of *Rht-D1b* (45.1%)

was observed in Chinese cultivars, and enriched in the Yellow and Huai wheat zone

(83.6%) out of 10 Chinese agroecological wheat zones (Table S2). Which further

supporting that a distinct environmental application of two Green Revolution genes

(Wurschum et al., 2015; Grogan et al., 2016; Zhao et al., 2019), and highlighting that

the importance of *Rht-D1b* in wheat production and breeding in the biggest wheat

region of China.

The application frequency of *Rht-B1b* or *Rht-D1b* increased from 57.3% to 77.8%

in MC which is released before and after 2000 (Figure 3B), highlighting the green

alleles are the major dwarf gene applied in current wheat varieties.

The heat adaptation potential of *Rht-B1b* and *Rht-D1b*

To evaluate the thermotolerance of green alleles for facing the Global warming

challenges, Late-sowing field assays (Chauhan et al., 2009; Tiwari et al., 2012) were

performed in three years, which conferred higher temperatures for each growth stage

of wheat (Figure 4A). Almost investigated traits significantly changed with the late-

sowing (Figure 4B, Table S5), indicating that wheat traits are susceptible to late-

sowing.

For plant structure, the plant height and flag leaf area were decreased about 30%,

and the tiller number were decreased more than 50% in late-sowing condition,

however, for yield traits, only the thousand kernel weight were decreased more than

15%, indicating that the vegetative growth stage of wheat was more sensitive to the

environmental alterations (Figure 4B, Table S5).

Notably, the agronomic traits of green alleles accessions showed more severe

influences by the late-sowing, indicating the green alleles are more sensitive to the

temperature increase than *NO* (Figure 4B), especially the accessions with *Rht-D1b*.

The effect of green alleles in response to increasing temperature

To further confirm the effects of high temperature on green alleles, we collected

all traits data of 400 accessions in seven environments and the available

meteorological data in seven environments for multivariate regression analysis

(Chmielewski and Potts, 1995; Tao et al., 2018). We found relative humidity and solar

radiation is not significantly contribute influence to wheat traits variation, while the

temperature in different developmental stage contributes the substantial effect to all

traits of wheat population (Table S8).

The temperature showed the necessary variations on the different accessions at

different developmental stage in seven environments (Figure 5A). The regression

analysis shows that the average temperature at different stage significantly affected

wheat traits (Table S9). Based on the regression equations, we predict the traits

alterations when the average temperature increase 1 and 2 □ (Figure 5B and C). We

found the tilling number, flag leaf angle, grain weight and plant height are the most

temperature sensitive traits, generally consistent to the estimation by the late sowing

(Figure 4B). Both *Rht-B1b* and *Rht-D1b* were more severe influenced by the

increasing temperature, compared to *NO* accessions.

This regression analysis showed the temperature increase in seedling stage is the

most severe disruption for the wheat growth, except the flag leaf angle is mostly

influenced by the temperature in filling stage (Table S8). The yield per plant reduces

around 20% and 38% if the average temperature increases 1 and 2 □ (Figure 5D). The

yield penalty of cultivars containing green alleles is about 25% higher compared to

that in *NO* cultivars. Further calculation demonstrates that the temperature increases at

seedling stage cause the most penalty for the yield per plant, which directly inhibits

the tiller number (Figure 5E). This regression analysis strongly suggest that green

alleles is more sensitive to the temperature increase, will cause a severe yield penalty

in the global warming scenarios.

DISCUSSION

Decades application had proved the capability of green alleles in controlling plant

structure and increasing wheat yield (Kertesz et al., 1991; Jin et al., 2015; Jobson et

al., 2019). Our results further showed the *Rht-B1b* and *Rht-D1b* reduced tillers, flag

leaf length, and flag leaf angle, resulted in more compact wheat plant structure, which

allow a higher planting density and contribute to the significant yield increase per unit

area, as demonstrated in rice and maize (Sinclair and Sheehy, 1999; Evenson and

Gollin, 2003). Meanwhile, these two alleles caused the increase of flag leaf width,

which made the leaf shape more reasonable and increased the transverse light

receiving area for enhancing photosynthetic efficiency (Gale et al., 1974; Morgan et

al., 1990), these findings may provide references for wheat structure improvement.

Notably the more erect flag leaf was observed in accessions containing green alleles,

which is unknown traits related to the GA signaling, need further clarified. It would

play valuable role in improving modern wheat cultivars.

In rice, the GA negatively regulated tiller number through SLR1-MOC1 mechanism

(Xu et al., 2021), showing the interaction mechanism between GA and strigolactones

(Marzec, 2017). However, we found the green alleles reduce the tiller number in

wheat, indicating a different mechanism in wheat to control the tiller number through

the DELLA. The reduced tiller may confer a benefit for the dry-land wheat yield, but

limited the yield potential under well water conditions (Houshmandfar et al., 2020).

We speculate that the less tiller number would be another reason to require a higher

plant density for the wheat cultivars containing green alleles.

Although many studies have reported that the semi-dwarfing genes affect wheat yield

related traits such in number, grain number per spike and thousand kernel weight

(Pinthus and Levy, 1983; Gale and Youssefian, 1985; Richards, 1992; Zhang et al.,

2013), but no convinced conclusions have been reached so far. In this study, we found

the *Rht-B1b* contribute more the kernel number per spike while *Rht-D1b* confers

higher kernel weight in different genetic background and environments (Figure 2),

highlighting the distinct way to increase wheat yield, this finding is great significance

for wheat production and breeding.

Our data suggested that the phenotypes controlled by green alleles is easily influenced

by the temperature increase, consistent to GA signaling participate in environmental

perception (Mo et al., 2018; Bao et al., 2020; Tang et al., 2021; Xu et al., 2021). The

increased temperature confers broad negative effect on the wheat structure and yield

traits (Figure 5). We found the tiller number is the most heat sensitive traits for wheat

yield. The reduced tiller number would require higher planting density for

maintaining wheat production. However, the flag leaf angle is also significantly

increased by the temperature increase (Figure 5), which would prevent the higher

planting density practice. Thus the dilemma between tiller number and flag leaf angle

has to resolved in futural wheat breeding. For the cultivars containing green alleles,

especially containing *Rht-D1b*, temperature increase would cause more yield loss to

250 risk futural food supply.

252 **MATERIALS AND METHODS**

253 **Plant materials and growing conditions**

254 400 accessions of bread wheat were collected in this study, including modern cultivars

255 and landraces from worldwide regions (Table S1). All materials were planted in seven

256 environments (E1--E7) at Yangling (34°28′N, 108°07′E, altitude 517 m) and

257 Chongzhou (30°63′N, 103°67′E, altitude 1300 m) of China in 2018-2021 (Table S10).

258 The experiments were arranged in a randomized complete block design with each

259 variety planted in three replicates. Each accession was sown with 1 m in length and a

260 row spacing of 20 cm, with 10 seeds per row. Field management was consistent with

261 local practices for wheat production.

262 **Agronomic traits measurements**

263 The traits of plant structure and yield were measured at physiological maturity. Three

264 healthy individuals selected from each row were used to investigate plant height (PH),

265 peduncle length (PL), total tiller number (TTN), productive tiller number (PTN), flag

266 leaf length (FLL), flag leaf width (FLW), flag leaf angle (FLAN), spike length (SL),

267 total spikelet number (TSN), fertile spikelet number (FSN) and kernel number per

268 spike (KNS) according to established protocols (Fan et al., 2015; Sun et al., 2017;

269 Guan et al., 2019; Jobson et al., 2019). Days to heading was assessed as the interval

270 between the date of seeding emergence and the date at which 50% of spikes per row

271 emerged from the flag leaf, days to flowering was assessed as the interval between the

272 date of seeding emergence and the date at which 50% of spikes per row have extruded

273 at least one anther (Kiseleva et al., 2016; Li et al., 2019). Further, we calculated flag

274 leaf area (FLA, FLL × FLW × 0.83), spikelet spacing (SS, SL / TSN) and yield per

275 plant (YPP, PTN×KNS×TKW). In addition, thousand kernel weight (TKW), kernel

276 length (KL) and kernel width (KW) were recorded using the rapid SC-G grain

277 appearance quality image analysis system developed by Hangzhou WSeen Detection

Technology Co, Ltd, China (Guan et al., 2019).

DNA extraction and genotyping

Genomic DNA was extracted from wheat flag leaves of heading stage plants

following the modified Cetyl Trimethylammonium Bromide (CTAB) method for

genotyping (Wehner et al., 2015). Kompetitive allele-specific PCR (KASP) markers

were developed by the mutation site of *Rht-B1b* and *Rht-D1b* to identify all the

accessions (Ma et al., 2019). All primers were listed in Supplementary Table S11. The

KASP assay was performed as following: assays were set up as 5 μL reactions

containing 2.2 μL template DNA dried to assay plate, 2.5 μL of V4.0 2x Kaspar mix

(LGC Group, Teddington, UK), 2.444 μL ddH$_2$O and 0.056 μL primer mix; and the

PCR cycling included hotstart at 94 °C for 15 min, followed by 10 touchdown cycles

(94 °C for 20 s; touchdown 61°C, -0.6 °C per cycle, 1 min), then followed by 35

cycles of amplification (94 °C for 20 s; 55 °C for 1 min).

Statistical analysis

The best linear unbiased estimate (BLUE) values for phenotypic under seven

environments were obtained using a mixed linear model by genotype as a fixed effect

and replicates as random effects. Analysis of variance (ANOVA) was performed to

test the homogeneity of genotype and phenotypic based on linear mixed model using

the Python SciPy module. Differences between phenotypic of different genotypes

were tested by *t-test* (Virtanen et al., 2020). If there is homogeneity of variance

between the two sets of data, the default argument to the function *stats.ttest_ind* is

used; otherwise, the parameter *equal_var* is set to *False*. All analyses were conducted

using *lme4* (Bates et al., 2015) and *lsmeains* packages (Lenth, 2016) in R3.6.2 (Bates

et al., 2014).

Permutation test

To avoid the co-selected SNPs and/or genes interferes, the permutation test was

performed on each genotype pairs to evaluate the effect of different genomic

background. The wheat population was divided into three groups by genotypes (*NO*,

B1b, *D1b*), 50% of the accessions were randomly selected to test the significance

between different groups. 1000 permutations were performed to examine how often

significant differences between groups occurred in by chance alone. Significant

differences between groups were considered to be reliable if they occurred in a

frequency more than 800; not vice versa (Wang et al., 2018; Xu et al., 2020).

Multivariate regression analysis

Multiple linear regression analysis of climate factors

Weather data were taken from the China Meteorological Data Service Center

(https://data.cma.cn). We collected 6 climate variables, that is, the maximum and

minimum temperatures, precipitation, sunlight hours and relative humidity and the

solar radiation in seven environments. According to the heading and flowering time of

each accession and 6 climate variables, the average maximum temperature, the

average minimum temperature, the mean precipitation, the sunlight hours, the average

relative humidity and the solar radiation at seeding stage, flowering stage and filling

stage for every accessions were collected (Table S6). Multiple regression analysis was

used to examine the effect of meteorological factors on phenotypes (Chmielewski and

Potts, 1995; Tao et al., 2018). The equation was:

$Y=a+bX_1+cX_2+dX_3+eX_4+fX_5+gX_6+hX_7+iX_8+jX_9+kX_{10}+lX_{11}+mX_{12}+nX_{13}+$

$oX_{14}+pX_{15}+qX_{16}+rX_{17}+sX_{18}$

where Y represents the phenotypic data. $X_1, X_2, X_3, X_4, X_5, X_6, X_7, X_8, X_9, X_{10}, X_{11}, X_{12},$

$X_{13}, X_{14}, X_{15}, X_{16}, X_{17}, X_{18}$ indicate the average maximum temperature, the average

minimum temperature, the mean precipitation, the sunlight hours, the average relative

humidity and the solar radiation at seeding stage, flowering stage and filling stage,

respectively. b, c, d, e, f, g, h, i, j, k, l, m, n, o, p, q, r, s are the regression coefficients,

and a is a constant.

Multiple linear regression analysis of average temperature at different stage

Multiple regression analysis was used to examine the effect of average temperature at

different stage on phenotypes. The equation was:

$$Y=m_1+m_2Xa+m_3Xb+m_4Xc$$

where Y represents the phenotypic data. Xa, Xb, Xc represents the average temperature

at seeding stage, flowering stage and filling stage, respectively. m_2, m_3, m_4 are the

regression coefficients, and m_1 is a constant.

We used *fit()* function to construct the general equation for regression analysis, and

then *drop1()* function was used to conduct stepwise regression for the constructed

equation. Meteorological factors with the smallest values from Akaike Information

Criterion (AIC) were always removed until the equation was constructed successfully.

The criteria of the constructed equation were as follows: the significance of T-test for

the regression coefficient of independent variables was $Pt < 0.05$; the significance of

F-test for the whole equation was $Pf < 0.05$, and the R-squared > 0.8.

Supplemental Data

Supplemental Table 1. The information and phenotypes of 400 wheat accessions in

seven environments.

Supplemental Table 2. The distributions of wheat accessions in different

agroecological regions.

Supplemental Table 3. The durations days for all wheat accessions at different stage.

Supplemental Table 4. The average temperature of wheat different sowing date at

different growth stages.

Supplemental Table 5. The alterations of traits with late-sowing.

Supplemental Table 6. The meteorological data in different periods and years for all

wheat accessions.

Supplemental Table 7. The heading and flowering time for all wheat accessions.

Supplemental Table 8. The multiple linear regression equations with multiple

meteorological factors.

Supplemental Table 9. The multiple linear regression equations with temperatures.

Supplemental Table 10. The sowing information of seven environments.

Supplemental Table 11. The KASP primer sequence.

ACKNOWLEDGEMENTS

This work was supported by Key Research and Development Projects of Shaanxi

Province (Collection, Exploration, Identification and Utilization of Elite Wheat

Germplasm) and grants from the National Natural Science Foundation of China

[31571756 and 31501380]. We thank Guoyue Chen and Zhien Pu for wheat field

investigation in Sichuan Province, and thank Yufeng Zhang for wheat hybrid group

field investigation in Hebei Province.

AUTHOR CONTRIBUTIONS

S.X. designed the concept and experiments. Z.L., X.L., J.Z. and X.M. performed field

investigation. J.C and X.Z performed the phenotypic investigation of transgenic lines

in field. Z.L., Z.H., W.J., X.L. and S.X. analyzed the data and wrote the manuscript.

All the authors were involved in the revision of the manuscript and approved the final

manuscript.

Data availability statement

All data, models, or code generated or used during the study are available from the

corresponding author by request. The phenotypes and SNP data are shown in

https://iwheat.net/resource/.

REFERENCES

Bao, S., Hua, C., Shen, L., and Yu, H. (2020). New insights into gibberellin signaling in

regulating flowering in Arabidopsis. J Integr Plant Biol **62**, 118-131.

Bates, D.M., Maechler, M., Bolker, B.M., and Walker, S.C. (2014). Package lme4: linear

mixed-effects models using Eigen and S4. Journal of statistical software **67**.

Binenbaum, J., Weinstain, R., and Shani, E. (2018). Gibberellin Localization and Transport in

Plants. Trends Plant Sci **23**, 410-421.

Bush, M.G., and Evans, L.T. (1988). Growth and development in tall and dwarf isogenic lines

of spring wheat. Field Crops Research **18**, 243-270.

Chauhan, S., Srivalli, S., Nautiyal, A.R., and Khanna-Chopra, R. (2009). Wheat cultivars

differing in heat tolerance show a differential response to monocarpic senescence

under   high-temperature   stress   and   the   involvement   of   serine   proteases.

Photosynthetica **47**, 536-547.

Chmielewski, F.M., and Potts, J.M. (1995). The relationship between crop yields from an

experiment   in   southern   England   and   long-term   climate   variations.   Agricultural   &

Forest Meteorology **73**, 43-66.

Claeys, H., De Bodt, S., and Inzé, D. (2014). Gibberellins and DELLAs: central nodes in

growth regulatory networks. Trends Plant Sci **19**, 231-239.

Evenson, R.E., and Gollin, D. (2003). Assessing the impact of the green revolution, 1960 to

2000. Science **300**, 758-762.

Fan, X., Cui, F., Zhao, C., Zhang, W., Yang, L., Zhao, X., Han, J., Su, Q., Ji, J., Zhao, Z.,

Tong, Y., and Li, J. (2015). QTLs for flag leaf size and their influence on yield-related

traits in wheat (Triticum aestivum L.). Molecular Breeding **35**, 24.

Gale, M.D., and Youssefian, S. (1985). Chapter 1–Dwarfing genes in wheat. Progress in Plant

Breeding–1, 1-35.

Gale, M.D., Edrich, J., and Lupton, F.G.H. (1974). Photosynthetic rates and the effects of

applied gibberellin in some dwarf, semi-dwarf and tall wheat varieties (Triticum

aestivum). The Journal of Agricultural Science **83**, 43-46.

Gao, Z., Shi, Z., Zhang, A., and Guo, J. (2015). Distribution of genes associated with yield

potential and water-saving in Chinese Zone II wheat detected by developed functional

markers. Journal of genetics **94**, 35-42.

Grogan, S.M., Brown-Guedira, G., Haley, S.D., McMaster, G.S., Reid, S.D., Smith, J., and

Byrne, P.F. (2016). Allelic Variation in Developmental Genes and Effects on Winter

Wheat Heading Date in the U.S. Great Plains. PLoS One **11**, e0152852.

Guan, P., Di, N., Mu, Q., Shen, X., Wang, Y., Wang, X., Yu, K., Song, W., Chen, Y., Xin, M.,

Hu, Z., Guo, W., Yao, Y., Ni, Z., Sun, Q., and Peng, H. (2019). Use of near-isogenic

lines to precisely map and validate a major QTL for grain weight on chromosome 4AL

in bread wheat (Triticum aestivum L.). Theor Appl Genet **132**, 2367-2379.

Guedira, M., Brown-Guedira, G., Van Sanford, D., Sneller, C., Souza, E., and Marshall, D.

(2010). Distribution of Rht Genes in Modern and Historic Winter Wheat Cultivars from

the Eastern and Central USA. Crop Science - CROP SCI **50**.

Hedden, P. (2003). The genes of the Green Revolution. Trends in genetics : TIG **19**, 5-9.

Houshmandfar, A., Ota, N., O'Leary, G.J., Zheng, B., Chen, Y., Tausz-Posch, S., Fitzgerald,

G.J., Richards, R., Rebetzke, G.J., and Tausz, M. (2020). A reduced-tillering trait

shows small but important yield gains in dryland wheat production. Glob Chang Biol

26, 4056-4067.

Jatayev, S., Sukhikh, I., Vavilova, V., Smolenskaya, S.E., Goncharov, N.P., Kurishbayev, A.,

Zotova, L., Absattarova, A., Serikbay, D., Hu, Y.G., Borisjuk, N., Gupta, N.K., Jacobs,

B., de Groot, S., Koekemoer, F., Alharthi, B., Lethola, K., Cu, D.T., Schramm, C.,

Anderson, P., Jenkins, C.L.D., Soole, K.L., Shavrukov, Y., and Langridge, P. (2020).

Green revolution 'stumbles' in a dry environment: Dwarf wheat with Rht genes fails to

produce higher grain yield than taller plants under drought. Plant Cell Environ **43,**

2355-2364.

Jin, K., Shen, J., Ashton, R.W., White, R.P., Dodd, I.C., Phillips, A.L., Parry, M.A., and

Whalley, W.R. (2015). The effect of impedance to root growth on plant architecture in

wheat. Plant Soil **392,** 323-332.

Jobson, E.M., Johnston, R.E., Oiestad, A.J., Martin, J.M., and Giroux, M.J. (2019). The

Impact of the Wheat Rht-B1b Semi-Dwarfing Allele on Photosynthesis and Seed

Development Under Field Conditions. Front Plant Sci **10,** 51.

Kertesz, Z., Flintham, J.E., and Gale, M.D. (1991). EFFECTS OF Rht DWARFING GENES

ON WHEAT GRAIN YIELD AND ITS COMPONENTS UNDER EASTERN

EUROPEAN CONDITIONS. Cereal Research Communications **19,** 297-304.

Kiseleva, A.A., Shcherban, A.B., Leonova, I.N., Frenkel, Z., and Salina, E.A. (2016).

Identification of new heading date determinants in wheat 5B chromosome. BMC Plant

Biol **16 Suppl 1,** 8.

Lanning, S.P., Martin, J.M., Stougaard, R.N., Guillen-Portal, F.R., and Talbert, L.E. (2012).

Evaluation of Near-Isogenic Lines for Three Height-Reducing Genes in Hard Red

Spring Wheat. Crop Science **52,** 1145.

Lenth, R.V. (2016). Least-Squares Means: The R Package lsmeans. Journal of Statistical

451    Software **69**, 1 - 33.

452 **Li, L., Mao, X., Wang, J., Chang, X., Reynolds, M., and Jing, R.** (2019). Genetic dissection of

453    drought and heat-responsive agronomic traits in wheat. Plant Cell Environ **42**, 2540-

454    2553.

455 **Liao, Z., Yu, H., Duan, J., Yuan, K., Yu, C., Meng, X., Kou, L., Chen, M., Jing, Y., Liu, G.,**

456    **Smith, S.M., and Li, J.** (2019). SLR1 inhibits MOC1 degradation to coordinate tiller

457    number and plant height in rice. Nat Commun **10**, 2738.

458 **Liu, Q., Wu, K., Harberd, N.P., and Fu, X.** (2021). Green Revolution DELLAs: From

459    translational reinitiation to future sustainable agriculture. Mol Plant **14**, 547-549.

460 **Löfke, C., Zwiewka, M., Heilmann, I., Van Montagu, M.C.E., Teichmann, T., and Friml, J.**

461    (2013). Asymmetric gibberellin signaling regulates vacuolar trafficking of PIN auxin

462    transporters during root gravitropism. Proc Natl Acad Sci U S A **110**, 3627-3632.

463 **Ma, J., Ding, P., Liu, J., Li, T., Zou, Y., Habib, A., Mu, Y., Tang, H., Jiang, Q., Liu, Y., Chen,**

464    **G., Wang, J., Deng, M., Qi, P., Li, W., Pu, Z., Zheng, Y., Wei, Y., and Lan, X.** (2019).

465    Identification and validation of a major and stably expressed QTL for spikelet number

466    per spike in bread wheat. Theor Appl Genet **132**, 3155-3167.

467 **Marzec, M.** (2017). Strigolactones and Gibberellins: A New Couple in the Phytohormone

468    World? Trends Plant Sci **22**, 813-815.

469 **Mo, Y., Vanzetti, L.S., Hale, I., Spagnolo, E.J., Guidobaldi, F., Al-Oboudi, J., Odle, N., Pearce,**

470    **S., Helguera, M., and Dubcovsky, J.** (2018). Identification and characterization of

471    Rht25, a locus on chromosome arm 6AS affecting wheat plant height, heading time,

472    and spike development. Theor Appl Genet **131**, 2021-2035.

Monneveux, P., Jing, R.L., and Misra, S.C. (2012). Phenotyping for drought adaptation in

wheat using physiological traits. Front. Physiol. **3**, 12.

Morgan, J.A., Lecain, D.R., and Wells, R. (1990). Semidwarfing Genes Concentrate

Photosynthetic Machinery and Affect Leaf Gas Exchange of Wheat. Crop Science **30**,

602-608.

Peng, J., Richards, D.E., Hartley, N.M., Murphy, G.P., Devos, K.M., Flintham, J.E., Beales, J.,

Fish, L.J., Worland, A.J., Pelica, F., Sudhakar, D., Christou, P., Snape, J.W., Gale,

M.D., and Harberd, N.P. (1999). 'Green revolution' genes encode mutant gibberellin

response modulators. Nature **400**, 256-261.

Pinthus, M.J., and Levy, A.A. (1983). The relationship between the Rht 1 and Rht 2 dwarfing

genes and grain weight in Triticum aestivum L. spring wheat. Theoretical & Applied

Genetics **66**, 153-157.

Qin, H., Pandey, B.K., Li, Y., Huang, G., Wang, J., Quan, R., Zhou, J., Zhou, Y., Miao, Y.,

Zhang, D., Bennett, M.J., and Huang, R. (2022). Orchestration of ethylene and

gibberellin signals determines primary root elongation in rice. Plant Cell **34**, 1273-

1288.

Richards, R. (1992). The effect of dwarfing genes in spring wheat in dry environments. I.

Agronomic characteristics. Australian Journal of Agricultural Research **43**, 517-527.

Ross, J.J., and Quittenden, L.J. (2016). Interactions between Brassinosteroids and

Gibberellins: Synthesis or Signaling? Plant Cell **28**, 829-832.

Sherman, J.D., Nash, D., Lanning, S.P., Martin, J.M., Blake, N.K., Morris, C.F., and Talbert,

L.E. (2014). Genetics of End-Use Quality Differences between a Modern and

Historical Spring Wheat. Crop Science **54**, 1972-1980.

Sinclair, T.R., and Sheehy, J.E. (1999). Erect Leaves and Photosynthesis in Rice. Science

283, 1455-1455.

Sun, C., Zhang, F., Yan, X., Zhang, X., Dong, Z., Cui, D., and Chen, F. (2017). Genome-wide

association study for 13 agronomic traits reveals distribution of superior alleles in

bread wheat from the Yellow and Huai Valley of China. Plant Biotechnol J **15**, 953-

969.

Tang, T., Botwright Acuna, T., Spielmeyer, W., and Richards, R.A. (2021). Effect of

gibberellin-sensitive Rht18 and gibberellin-insensitive Rht-D1b dwarfing genes on

vegetative and reproductive growth in bread wheat. J Exp Bot **72**, 445-458.

Tao, Z., Wang, H., Dai, J., Alatalo, J., and Ge, Q. (2018). Modeling spatiotemporal variations

in leaf coloring date of three tree species across China. Agricultural and Forest

Meteorology **249**, 310-318.

Tiwari, C., Wallwork, H., Dhari, R., Arun, B., Mishra, V.K., and Joshi, A.K. (2012). Exploring

the possibility of obtaining terminal heat tolerance in a doubled haploid population of

spring wheat (Triticum aestivum L.) in the eastern Gangetic plains of India. Field

Crops Research **135**.

Topham, A.T., Taylor, R.E., Yan, D., Nambara, E., Johnston, I.G., and Bassel, G.W. (2017).

Temperature variability is integrated by a spatially embedded decision-making center

to break dormancy in Arabidopsis seeds. Proceedings of the National Academy of

Sciences of the United States of America **114**, 6629-6634.

Van De Velde, K., Thomas, S.G., Heyse, F., Kaspar, R., Van Der Straeten, D., and Rohde, A.

(2021). N-terminal truncated RHT-1 proteins generated by translational reinitiation

cause semi-dwarfing of wheat Green Revolution alleles. Mol Plant **14**, 679-687.

Velu, G., Singh, R.P., Huerta, J., and Guzmán, C. (2017). Genetic impact of Rht dwarfing

genes on grain micronutrients concentration in wheat. Field Crops Res **214**, 373-377.

Wang, X., Chen, Q., Wu, Y., Lemmon, Z.H., Xu, G., Huang, C., Liang, Y., Xu, D., Li, D.,

Doebley, J.F., and Tian, F. (2018). Genome-wide Analysis of Transcriptional

Variability in a Large Maize-Teosinte Population. Mol Plant **11**, 443-459.

Wehner, G.G., Balko, C.C., Enders, M.M., Humbeck, K.K., and Ordon, F.F. (2015).

Identification of genomic regions involved in tolerance to drought stress and drought

stress induced leaf senescence in juvenile barley. BMC Plant Biol **15**, 125.

Wu, J., Kong, X., Wan, J., Liu, X., Zhang, X., Guo, X., Zhou, R., Zhao, G., Jing, R., Fu, X.,

and Jia, J. (2011). Dominant and pleiotropic effects of a GAI gene in wheat results

from a lack of interaction between DELLA and GID1. Plant Physiol **157**, 2120-2130.

Wurschum, T., Langer, S.M., and Longin, C.F. (2015). Genetic control of plant height in

European winter wheat cultivars. Theor Appl Genet **128**, 865-874.

Wurschum, T., Langer, S.M., Longin, C.F.H., Tucker, M.R., and Leiser, W.L. (2017). A

modern Green Revolution gene for reduced height in wheat. Plant J **92**, 892-903.

Wurschum, T., Liu, G., Boeven, P.H.G., Longin, C.F.H., Mirdita, V., Kazman, E., Zhao, Y.,

and Reif, J.C. (2018). Exploiting the Rht portfolio for hybrid wheat breeding. TAG.

Theoretical and applied genetics. Theoretische und angewandte Genetik **131**, 1433-

1442.

Xu, P., Chen, H., Li, T., Xu, F., Mao, Z., Cao, X., Miao, L., Du, S., Hua, J., Zhao, J., Guo, T.,

Kou, S., Wang, W., and Yang, H.Q. (2021). Blue light-dependent interactions of

CRY1 with GID1 and DELLA proteins regulate gibberellin signaling and

photomorphogenesis in Arabidopsis. Plant Cell **33**, 2375-2394.

Xu, W., Li, K., Li, S., Hou, Q., Zhang, Y., Liu, K., and Sun, Q. (2020). The R-Loop Atlas of

Arabidopsis Development and Responses to Environmental Stimuli. Plant Cell **32**,

888-903.

Zhang, J., Dell, B., Biddulph, B., Drake-Brockman, F., Walker, E., Khan, N., Wong, D.,

Hayden, M., and Appels, R. (2013). Wild-type alleles of Rht-B1 and Rht-D1 as

independent determinants of thousand-grain weight and kernel number per spike in

wheat. Molecular Breeding **32**, 771-783.

Zhang, X., Yang, S., Zhou, Y., He, Z., and Xia, X. (2006). Distribution of the Rht-B1b, Rht-

D1b and Rht8 reduced height genes in autumn-sown Chinese wheats detected by

molecular markers. Euphytica **152**, 109-116.

Zhao, J., Wang, Z., Liu, H., Zhao, J., Li, T., Hou, J., Zhang, X., and Hao, C. (2019). Global

status of 47 major wheat loci controlling yield, quality, adaptation and stress

resistance selected over the last century. BMC Plant Biol **19**, 5.

Figure. 1 The number of accessions with different genotypes in population and the correlation analysis between different agronomic traits.

(A) The number of wheat accessions with different genotypes in the population by KASP assay. *NO*, wheat accessions without *Rht-B1b* and *Rht-D1b*; *B1b*, wheat accessions containing *Rht-B1b*; *D1b*, wheat accessions containing *Rht-D1b*. The number indicates wheat accessions number in each genotype.

(B) The correlations analysis between different agronomic traits by the best linear unbiased estimate (BLUE) values. PH, plant height; PL, peduncle length; TTN, total tiller number; PTN, productive tiller number; FLL, flag leaf length; FLW, flag leaf width; FLA, flag leaf area; FLAN, flag leaf angle; SL, spike length; TSN, total spikelet number; FSN, fertile spikelet number; SS, spikelet spacing; KL, kernel length; KW, kernel width; KNS, kernel number per spike; TKW, thousand kernel weight. Red lines mean positive correlation and gray lines mean negative correlation. The correlations were analyzed by the "corrplot" in the R package using the BLUE values in Table S1.

Figure. 2 The function of the *Rht-B1b* and *Rht-D1b* on wheat traits.

(A) The function of the *Rht-B1b* and *Rht-D1b* on wheat structure. The letters at the top of violin plots indicate the significant difference between each group (two-sided Dunn's Kruskal–Wallis test, $p < 0.01$). The numbers represent the ratio of comparison showing significantly difference / all comparison, the 1000 permutation tests. Red number marks the ratio less 800. The percentages in parentheses indicate environmental stability, the number of environments that have the same effect divided by the total number of environments. The data come from the BLUE values (n=400) of seven conditions, as shown in Table S1.

(B) The function of the *Rht-B1b* and *Rht-D1b* on wheat yield.

Figure. 3 The spatiotemporal distribution of the *Rht-B1b* and *Rht-D1b* in China.

(A) The introduction and distribution of the *Rht-B1b* and *Rht-D1b* in Chinese landraces and Chinese modern cultivars. Y&H, Yellow and Huai wheat zone (blue); OT, Other agroecological wheat zone (red). The numbers at the top of the bars means the total number of wheat accessions. The numbers in the blue and red bar indicate the number of wheat accessions in different agroecological wheat zone, respectively.

(B) The application time of the *Rht-B1b* and *Rht-D1b* in Chinese modern cultivars (MC). B-2000, the cultivars was released from 1970's to 2000; P-2000, post 2000.

Figure. 4 The phenotypic variations with late-sowing.

(A) The temperature variations with late-sowing. N, normal sowing time conditions; L, late sowing time conditions. The sowing date and temperature was shown in Table S3 and S4.

(B) The phenotypic alteration with late sowing. The data is the phenotypic data in late-sowing/ phenotypic data in normal-sowing. Different letters on the bars indicate the significant differences ($p < 0.05$, two-sided Dunn's Kruskal–Wallis test), as shown in Table S5. The gray dash lines indicate the value 1.

Figure. 5 The phenotypes of the *Rht-B1b* and *Rht-D1b* in response to temperature increase.

(A) The temperature variations in seven different environments. The environments information, temperature and the developmental stage of each accession as shown in Table S6, S7 and S11.

(B) The structure traits alterations with average temperature increased 0.5 and 1℃. The phenotypic value of Normal, +0.5℃ and +1℃, was calculated by regression equation with average temperatures in observed temperature, the average temperature increased +0.5℃, +1℃ respectively. The percentages the bars indicate the phenotypic changed ratio of corresponding genotype wheat accessions when the average temperature increased 0.5 and 1℃.

(C) The alterations of wheat yield traits when the average temperature increased 0.5 and 1℃. The phenotypic value of Normal, +0.5℃ and +1℃, was calculated by regression equation with average temperatures in observed temperature, the average temperature increased +0.5℃, +1℃ respectively.

(D) The alterations of wheat yield per plant (YPP) with temperature increase. The YPP was estimated by productive tiller number * kernel number per spike * kernel weight in (C).

(E) The alterations of wheat yield per plant with temperature increased 0.5 and 1℃ at different developmental stage. The YPP was estimated by productive tiller number * kernel number per spike * kernel weight, which was calculated by the regression equation with the average temperature at sowing-heading, heading-flowering and grain-filling stages of wheat, respectively.