1 **Conservation networks do not match ecological requirements of amphibians**
2

3 **Authors**: Matutini Florence, Jacques Baudry, Marie-Josée Fortin, Guillaume Pain,
4 Joséphine Pithon
5
6

MATUTINI Florence – florence.matutini@gmail.com - BAGAP, INRAE, Institut Agro,
ESA, 49000, Angers, France ; Office français de la biodiversité, 5 Imp. de Saint-Thibault,
78610 Auffargis
BAUDRY Jacques –jmhbaudry2@gmail.com - BAGAP, INRAE, Institut Agro, ESA,
35042, Rennes, France
FORTIN Marie-Josée - mariejosee.fortin@utoronto.ca - Dept of Ecology and
Evolutionary Biology, Univ. of Toronto, Toronto, ON, Canada M5S 3B2
PAIN Guillaume – g.pain@groupe-esa.com - BAGAP, INRAE, Institut Agro, ESA,
49000, Angers, France
PITHON Joséphine - j.pithon@groupe-esa.com - BAGAP, INRAE, Institut Agro, ESA,
49000, Angers, France
18

19 **Corresponding author**:
20 Matutini Florence, florence.matutini@gmail.com - +33 603962008 - Office français de la
21 biodiversité, 5 Imp. de Saint-Thibault, 78610 Auffargis

22
23
24
25

Abstract

1. Amphibians are among the most threatened taxa as they are highly sensitive to habitat degradation and fragmentation. They are considered as model species to evaluate habitats quality in agricultural landscapes. In France, all amphibian species have a protected status requiring recovery plans for their conservation. Conservation networks combining protected areas and green infrastructure can help the maintenance of their habitats and while favouring their movement in fragmented landscape such as farmlands. Yet, assessing the effectiveness of conservation networks is challenging.

2. Here, we compared ecological requirements of amphibian species with existing conservation network coverage in a human-dominated region of western France. First, we mapped suitable habitat distributions for nine species of amphibian with varying ecological requirements and mobility. Second, we used stacking species distribution modelling (SSDM) to produce multi-species habitat suitability maps. Then, we defined species and multi-species core habitats and performed a connectivity analysis using Circuitscape theory to identify spatial continuity in suitable habitats at regional scale. Finally, we compared different suitability maps with existing conservation networks to assess conservation coverage and efficiency.

3. We highlighted a mismatch between the most suitable amphibian habitats at regional scale and the conservation network, both for common species and for species of high conservation concern. We also found two bottlenecks between areas of suitable habitat which might be crucial for population movements provoked by global change, especially for species associated with hedgerow mosaic landscapes. These bottlenecks were not covered by any form of protection and are located in an intensive farmland context.

4. *Synthesis and applications* - We advocate the need to better integrate agricultural landscape mosaics into protected areas as well as to protect the associated extensive agricultural practices suitable for biodiversity, including extensive livestock farming. We also emphasize the importance of interacting layers of green infrastructure for amphibian conservation and the need for it to be effectively considered in land-use planning policies. Further investigations are needed to strengthen and restore habitat continuities in area with regional bottlenecks marked by rare and degraded semi-natural elements.

Keywords

1 Introduction

The biodiversity decline is still accelerating across the world affecting both scarce and common species (Dirzo et al., 2014; Gaston & Fuller, 2008) especially in agroecosystems (Kleijn et al., 2011; Seibold et al., 2019). To palliate this decline, the establishment of protected areas (PAs) is a relevant conservation tool in a human dominated world (Rodrigues & Cazalis, 2020). However, PAs alone are not sufficient to protect biodiversity and they should be part of an ensemble of management and planning strategies aiming to create functional conservation networks. Green infrastructure (GI) has been proposed as a way to address this issue (Chatzimentor et al., 2020; Salomaa et al., 2017). Such initiatives aim to maintain a coherent and functional network of suitable habitats and to improve biodiversity integration into land-use planning. GI design is based on the *ecological network* concept (Opdam et al., 2006) emphasizing complementarity of core habitat patches and ecological corridors at different spatial and temporal scales (Opdam et al., 2006; Salomaa et al., 2017).

Evaluating the efficiency of conservation networks (here PAs and GI) is a real challenge (e.g., Rodrigues & Cazalis, 2020). In France, GI has mostly been defined based on spatial analysis of land-cover data without considering species data. It is based on a simplistic classification of ecosystems reduced to broad categories of land cover data (i.e. different layers for forest or wetlands separately) with a poor consideration of habitat complementation. In contrast, PAs have usually been defined according to the needs of focal endangered species and/or habitats, often charismatic species such as birds and endangered plants, poorly representative of species-habitat relations (Rodrigues et al., 2004). Gap analysis and conservation prioritization are common approaches for a first-step assessment of the effectiveness of a conservation network and to inform decisions for conservation measures (Jennings 2000; Moilanen et al. 2005). To conduct a gap analysis, outputs of spatial modelling are particularly useful (Ahmadi et al., 2020). Species distribution models (SDMs) are commonly used to make spatially explicit predictions of the suitable environment of a given species from biological data (Guisan & Zimmermann, 2000) and multi-specific approaches have recently been developed (Scherrer et al., 2018; Thuiller et al., 2015). In addition, recent methods integrate both graph theory and SDMs for ecological network modelling (Duflot et al., 2018; Godet & Clauzel, 2020). These multi-species and integrative approaches could be relevant for gap analyses and spatial prioritization but remain underused (Foltête et al., 2020).

In western Europe, traditional hedgerow landscapes are of high conservation concern for numerous taxa such as plants, amphibians, birds, bats and arthropods (Baudry et al., 2000; Boissinot et al., 2019). These rural landscapes are associated with dense hedgerow networks interconnected with a mosaic of small interlocked patches of pastures, cultivated fields and bushes with high pond density (Burel & Baudry, 1995). They have been greatly impacted by agricultural intensification and have declined from 40 % to 80 % in Europe (Bazin & Schmutz, 1994). Amphibians are good candidates to assess the ecological quality of human-dominated landscape mosaics and to question the efficiency of existing conservation networks. Indeed, They are among the most threatened taxa with rapid and widespread population declines in both natural and modified habitats (Stuart et al., 2004). They have a bi-phasic life cycle with different ecological requirements inducing seasonal

migrations between habitat patches (Sinsch, 2014). Their poor mobility and their impermeable skin increases their sensitivity to anthropogenic disturbances, pollution and ecological fragmentation induced by urbanization and intensive agriculture (Cushman, 2006; Hamer & McDonnell, 2008; Stuart et al., 2004). Hence, some authors consider amphibians as good ecological indicators of general environmental health (Collins and Storfer 2003; Díaz-García et al. 2017 but see Sewell & Griffiths, 2009). In addition, amphibians have been the target of several citizen science programs in different countries around the world (De Solla et al., 2005; Petrovan & Schmidt, 2016) leading to a large amount of available data over wide areas which could be relevant for conservation tools assessment (Snäll et al., 2016).

In a human-dominated region of western France, we compared the predicted habitat requirements of amphibian species with existing conservation network coverage (of both protected areas and green infrastructure). Specifically, we: (1) produced single and multi-species habitat suitability maps for amphibians with differing ecological requirements and mobility (2) compared these habitat suitability maps with existing conservation networks (PAs and GI), and (3) identified gaps in conservation coverage which could be a priority for new potential conservation areas.

2   Material and Methods

2.1   Regional context and studied species

Ecological context

Our study was performed in a region of western France (Pays-de-la-Loire) covering 32,082 km². Intensification of agricultural practices have greatly transformed the landscape over the last century from a complex matrix of small fields, meadows and woody elements to more homogeneous landscapes with larger monocultural fields. With 21 species of amphibian (out of 43 species recorded at national scale), of which 12 are priority species for conservation, the region has considerable responsibility for the preservation of amphibian species and their habitats, including traditional hedgerow landscapes and wetlands.

Studied species

Nine amphibian species in the Pays-de-la-Loire region were studied (see Table 1 for species list). Three species have high conservation value (*Pelodytes punctatus*, *Triturus marmoratus* and *Rana temporaria* i.e., regional priority index > 1; Table 1). Three form a "FOR" group of forest species or species associated with dense hedgerow mosaic landscape (*T. marmoratus, R. temporaria* and *Salamandra salamandra*) and five form a "GEN" group of more generalist species (*Lissotriton helveticus*, *Bufo spinosus, Hyla arborea, Rana dalmatina* and *Triturus cristatus).* *P. punctatus* is a pioneer species mainly associated with disturbed open environments. Finally, the "ALL" group contained all nine studied species. Species were selected according to data availability and previous studies (Matutini, Baudry, Fortin, et al., 2021; Matutini, Baudry, Pain, et al., 2021).

Biodiversity conservation policies

In France, there are different conservation tools including protected areas, green
infrastructures, and inventoried sites with recognised biodiversity value but that are not
protected (Categories "PA","GI" and "INV" respectively in Table 2). We defined five
groups of conservation tool based on IUCN categories and protection levels (Table 2 and
Fig. 1). For two types of PAs (SCEN and ENS in PA-Group 2), site boundaries were not
available. As a result, only a point location per site was considered in the analysis (see
Appendix 2). If one conservation area was overlapped by another, we classed the area into
the group with the strongest protection level and therefore all PAs were excluded from GI.
French GI is divided into different land cover or landscape classes (Table 2). Each GI layer
includes two types of elements: habitat nodes, considered as "biodiversity reservoirs", and
"corridors" ensuring potential continuity between nodes based on geographical criteria.
163
2.2 **Single and multi-species habitat suitability maps**
We used 100 m-resolution habitat suitability maps resulting from a previous study of the
same nine species in Pays-de-la-Loire (Matutini, Baudry, Fortin, et al., 2021). This used
heterogeneous data from citizen sciences completed by professional surveys to perform
and evaluate (with independent dataset) presence-only habitat suitability modelling with
functional friction-based and multi-scale predictors (see Matutini, Baudry, Fortin, et al.,
2021 and Matutini, Baudry, Pain, et al., 2021 for full details and methods).
We produced multi-species suitability maps by compiling individual species maps
(Stack-SDM, e.g. Ferrier & Guisan, 2006) using a simple probabilistic stacking method
(pSSDM, see Zurell et al., 2019). We summed continuous individual maps without binary
"presence-absence" conversion (Calabrese et al., 2014; Scherrer et al., 2020) for three focal
species groups: (1) forest species (FOR), (2) generalist species (GEN), and (3) all species
(ALL). According to the ecological context of our region for a best integration of mosaic
effect for landscape evaluation, we summed species maps at 500 m-resolution after 100 m-
pixels aggregation using maximum suitability values (see Appendix 1 for additional
information and selection tests). Hence, we compiled single species standard deviation
maps to obtain a general representation of levels of confidence in our mapped distributions.
To assess the accuracy of the three pSSDMs, we compared multi-species suitability index
(i.e., the sum of single species suitability values from 0 to 1) with species richness from
independent pond monitoring (Matutini, Baudry, Pain, et al., 2021). We calculated a
community-AUC (Scherrer et al., 2020) and R^2 value for each species group with 500
iterations.
186
187 2.3 **Reclassification and core habitat node definition**
188 For each species and stacked species group, we reclassified the final suitability map into
189 five categories of potential habitat: not suitable (below the ten-percentile value P_{10th}
190 method in Huang et al 2020), Less suitable (P_{10th}-55%), moderately suitable (55%-65%),
191 suitable (65%-80%), and highly suitable (>80%).
192
193 2.4 **Network patterns**
194 To map potential connections between each highly suitable patch (i.e., patch with a
195 suitability index up to 80%) at regional scale, we used circuit theory with Circuitscape

software (McRae et al., 2016). Circuit theory considered each cell as an electric node connected to neighbouring cells by resistors and defined by a landscape resistance (or conductance) value (McRae et al., 2008). We used highly suitable core habitats as source patches, and a friction map as a measure of conductance (i.e., conductance of each raster 100 m cell for movement). This method provides an exploration of potential links between nodes according to landscape permeability combining habitat suitability and connectivity modelling (Ahmadi et al., 2020; Koen et al., 2014).

Habitat suitability could however be a poor proxy of permeability as species may move in unsuitable habitats (Keeley et al., 2017). Hence, the friction map was computed using different friction cost values derived from the multi-species habitat map and landcover data (Keeley et al., 2016, 2017). Therefore, we considered moderately to highly suitable habitat classes (i.e., from 65% to 100%) as permeable features except for pixels crossed by a linear barrier. In less suitable habitats (i.e., under the 65% threshold), we calculated a friction cost based on a 100m-grid analysis performed over 10 m-resolution landscape layers followed by a classification procedure (see Appendix 3). 100m-resolution were selected to reduce computational power needed for Circuitscape analysis and to consider landscape complementation of fine-scale landscape elements (e.g., ditches, hedgerows and field margins, Pope et al., 2000; Mazerolle, 2005). The cost values of impermeable barriers (i.e., high-density urban area, highspeed train line, highway and dual carriageways) were considered as infinite. Permeable features facilitating movement across barriers (e.g., viaduct, wildlife pass and others, see Table 2) were digitized from aerial photographs. To reduce border effects, we allocated resistance values to a 5km-buffer around the studied region which were randomly selected if land-cover data were not available.

2.5 **Gap analysis and prioritizing for conservation**

We based the gap analysis framework on a method used by Ahmadi et al. (2020) that compared reclassified suitability maps with areas covered by PA or GI. We calculated "total conservation coverage" (i.e., the proportion of suitable habitat overlapping a conservation area), "conservation efficiency" (i.e., the proportion of conserved areas overlapping suitable habitats) and plotted histograms showing the total distribution of suitability values covered by different types of conservation area. To complete the analysis, we overlaid the existing conservation network with the final regional maps obtained using Circuitscape for each species group to discuss network configuration in relation to potential regional continuities. All spatial analysis were performed using R environment v. 3.5.3 (R Development Core Team, 2019).

3 **Results**
3.1 **Multi-species maps and core habitat definition**
For each group, the community-AUC varied from 0.76 (forest species) to 0.87 (generalist species) and R^2 from 0.23 (forest species) to 0.31 (nine studied species). The suitable maps are shown in Fig. 1 while associated confidence maps are available Appendix 4. For the nine studied species, highly suitable habitats covered 12% of the region and suitable habitats covered 49% of the region (Fig. 1). Highly suitable habitat for the forest species,

240   covered 11% and suitable habitat covered 17% of the region. Larger amounts of highly
241   suitable and suitable habitats were available to generalist species, covering 33% and 40%
242   of the region respectively.
243

244   3.2   **Protected areas (PAs)**
245   The distribution of protected areas along the habitat suitability gradients modelled for each
246   species or species group (FOR, GEN, ALL) is shown Fig. 2.
247

Conservation coverage
PAs covered a relatively low proportion of highly suitable habitat: 15%, 19% and 8% for
all species, forest species and generalist species respectively (Table 4). The most strictly
protected areas protection (PA group 1) covered even lower proportions of suitable habitat:
1% of highly suitable habitat and 2% of suitable habitat for each species group. Similarly,
for two of the three species of high conservation concern, PA covered only a relatively low
17% or 26% of highly suitable habitat, for *T. marmoratus* and *R. temporaria* respectively.
Only for *P. punctatus* was conservation coverage high: 90% of highly suitable habitat was
included in PA.
257

Conservation efficiency
Among PA-Group 1, no sites with the highest level of protection (names "RNN", see Table
2) contained highly suitable area for any species group or individual forest species except
P. punctatus (12% of RNN areas contain high suitable habitat) (Table 4). Other PA-
Group 1 protection levels were covered by 7 to 24% of multi-species core area and by 3 to
24% of species-specific highly suitable habitat, with a higher proportion for *P. punctatus*.
Regional Park (PA-Group 2) and Natura 2000 network (PA-Group 3) included less than
16% multi-species core area and lower proportions for the two-forest species individually.
For SCEN and ENS (in PA-Group 2) 26%, 12% and 8% of these sites were located in
highly suitable habitats for generalists, forest species and all nine species respectively.
Overall, PAs contain a higher proportion of highly suitable habitat for *P. punctatus* than for
other species.
270

3.3   **Inventoried sites (INV)**
Inventoried sites with known ecological value but no legal protection (INV-Group 5)
covered 13% of the region. 20% to 34% of their areas were covered by multi-species
highly suitable habitat (24 to 57% for suitable area). 11%, 21% and 1% are covered by
highly suitable habitat of *T. marmoratus*, *R. temporaria* and *P. Punctatus* respectively
(Table 4). INV-Group 5 covered the most suitable and highly suitable habitat of all studied
groups and species except for *P. punctatus* with a poor overlapping between this species
issues and areas of INV-Group 5 (i.e., 2%).
279

3.4   **Green infrastructure (GI)**
GI included and covered a larger area of highly suitable and suitable habitat for amphibian
species than PA particularly hedgerow networks BOCAGE followed by WOOD and WET
(see Table 2). Globally, GI covered around 50% of potential habitat for amphibian species.

Among the five GI's classes (Table 4), GI relating to traditional hedgerow landscapes
(BOCAGE) were the most widespread and covered from 16% to 33% of highly suitable
habitat of species or species groups (especially *T. marmoratus* with 33% and excluding *P.*
punctatus with only 2%). In addition, the GI BOCAGE contained 42% highly suitable
habitat for generalist species and between 12-21% for other species except *P. punctatus*.
GI relating to woodlands (WOOD) and wetlands (WET) covered more than 10% of highly
suitable habitat except for *T. marmoratus* and *P. punctatus*. GI relating to coastland
(COAST) and natural open grasslands (OPEN) covered small areas compared with other
GI and also a low area of suitable habitat for studied species.
293
Potential multi-habitat continuities patterns
The distribution patterns associated with highly suitable and suitable habitats are shown
Fig. 1 and supplemented in Fig. 3 by a more detailed analysis of the spatial organisation
and potential links between habitat patches. The central and the southern parts of the
region have a few patches of highly suitable habitat, especially for forest species.
Generalist species have a more homogeneous distribution of suitable habitats spreads over
the region. Suitable and highly suitable habitat distribution patterns associated with forest
species are mostly common with other species, with the exception of large suitable forest
patches especially at the north of the region. Indeed, 92% and 81% of highly and suitable
habitat of FOR species (i.e., suitability index > 65%) are include in GEN and ALL species'
high and suitable habitat respectively.
Regarding potential multi-specific regional continuities, two major sectors form
"bottlenecks" to north-south (sector B, Fig. 3) or east-west continuity axis and potential
movements (sector A, Fig. 3). This is particularly marked for forest species. These two
sectors contain few protected areas and show a cover mismatch with GI (Fig. 3).
309

4   Discussion

We highlighted a mismatch between the most suitable amphibian habitats at regional scale
and the existing conservation network, in particular protected areas (in coverage and
efficiency). Indeed, habitat patches with a high suitability index associated with common
or slightly rarer species were poorly covered by PA except for one species of high
conservation concern: *P. punctatus*. Yet, most suitable areas were better covered by GI
(especially GI layers associated with wood, hedgerow landscapes and wetlands) and by
INV sites (inventoried sites with ecological value but no regulatory protection). Difference
in coverage and efficiency obtain for PAs and INV highlight that some sites with
amphibian conservation issues are known but not protected. In addition, we identified two
narrow "bottlenecks" in suitable habitat continuities at regional scale which are only very
sparsely covered by the conservation network (PAs and GI).
322

4.1   General limits and interpretation framework

In addition to the limits related to the presence-only SDM detailed in Matutini, Baudry,
Fortin, et al. (2021) recently developed stacking-SDM methods require further
development to improve their evaluation. SDM are useful tools to predict the distribution
of habitat suitability but rarely predict the biology of populations (Lee◻Yaw et al., 2022).

The multi-species networks identified should be considered simply as maps of highly
suitable habitat for biodiversity (habitat perspective) - measured with highly sensitive and
biphasic amphibian species – rather than as a real multilayer network of amphibian
communities (species perspective). In addition, resistance values based on literature and
expert opinions or derived from SDM values are still debated (see Godet & Clauzel, 2020
for methods comparison and discussion) and information about species movement and
dispersion processes are still very scarce in the literature (Pittman et al. 2014). A more
rigorous approach would be to integrate independent genetic data to assess our final maps
(in addition to the external presence-absence dataset use to evaluate our models),
particularly in the two bottleneck sectors. However, these data were not available. This
might be worthwhile pursuing for post-validation and further investigations.
Our results must also be interpreted keeping in mind the spatial and temporal scales
involved. Amphibians have poor movement capacities, rarely exceeding a few kilometres
during dispersal movements (Sinsch, 2014). In this work, continuities were identified from
habitat distributions at regional scale and species level reflecting long-term ecological
processes and population movements across decades. This study takes a first step towards
assessing potential barriers to species distribution shifts in response to global change, but
does not directly include landscape dynamics and climate change scenarios. Rather it
provides elements at regional scale to guide regional policies and further studies to
improve multi-scale integration and efficiency of measures for biodiversity conservation.
348
4.2 **A mismatch for the conservation of amphibians**
Two main reasons may explain the mismatch between location of PAs and amphibian
habitat requirements. Firstly, PAs are generally defined according to the needs of rare
and/or emblematic species or habitats). In the study region, most PA focus on birds or
plants species, large open wetlands near the western coast or the Loire River valley but
only a few target species of amphibian and these tend to be rare species (e.g., *Pelobates*
cultripes, *Bombina variegata*). Most local, rare species of high conservation concern are
pioneer species and/or associated with loose or sandy soils with poor vegetation that are
found in particular on the coast, along the Loire River and in some quarries. Their ecology
contrasts with the other studied species, except for *P. punctatus* which shares many of the
same ecological requirements, explaining the better conservation coverage and efficiency
of PA for this species.
Secondly, PA definition in the region presents a "location bias". Indeed, PAs are
defined primarily in areas of low economic interest, less used for human activities
(Rodrigues et al., 2004). This bias is common worldwide and widely described in the
literature (Ahmadi et al., 2020; Joppa & Pfaff, 2009). It can be especially problematic for
species living in human-dominated landscapes, like most amphibian species. For example,
Rodrigues et al. (2004) demonstrated that amphibians contain a large number of "gap
species" worldwide, i.e., they are the biological group least covered by the global PA
network. In addition, our results show a potential "network bias" similar to Ahmadi et al.
(2020), i.e., the extension of PA is established without sufficiently considering their
position within the network. Indeed, the current PA sites (and the sites intended for the

future extension of PAs – see Appendix 4) are not located in the two areas with highest potential defined by our primary analysis of the regional continuity of suitable habitats.

4.3 **Green infrastructure (GI) as a complementary tool for biodiversity conservation**

Our results show GI offers better coverage of highly suitable habitats than PAs for amphibian species groups, but this remains partial, in particular in the two identified "bottleneck" sectors. Indeed, agricultural areas are poorly covered by PAs, generally due to a conflict of economic interest (Rodrigues et al. 2004) while GI provides complementary coverage in areas dominated by human activities. Agricultural landscapes with high conservation value for biodiversity such as hedgerow mosaic landscapes depend on the maintenance of traditional farming practices that may even be inhibited by PAs conservation (Schmitz et al., 2017)

In addition, we underline the importance of GI layers associated with wetlands, woodlands and traditional hedgerow mosaic landscapes for amphibians. Consequently, our results highlight the importance of considering several GI layers with potential interactions. GI tends to compartmentalize conservation issues by splitting them into distinct layers of habitat types considered as independent "subnetworks", where the "habitat" concept is limited to landcover categories (see Pilosof et al., 2017 for general perspective on multi-layer nature of ecological networks). Species survival is however conditioned by different ecological processes which generally go beyond a single type of land cover class. Some bat species, for example, can roost in buildings or old trees in summer, caves in winter, forage in open areas and wetlands and use hedgerow networks to transit or migrate. Species other than amphibians may also have very contrasting requirements during their life cycle (e.g., many holometabolous insects and odonates) with, for example, distinct aquatic and terrestrial phases.

Lastly, we expected a higher overlap of the GI layer called Bocage (traditional hedgerow mosaics) based on knowledge of amphibian ecology (Boissinot et al., 2019). This GI layer was mapped using combined land use data (i.e., hedges, ponds and meadows only) at 1km-resolution with a poor consideration of the interactions between these elements (i.e., simple metrics as density or area related to hedges, ponds and meadows were combined independently). Of course, other elements of the landscape are important for amphibians, such as small streams and wooded patches, as well as more complex processes related to landscape complementation and structural heterogeneity (Boissinot et al., 2019). Amphibian habitat network studies mainly focus on pond networks but natural or semi-natural forest patches, especially mature forest (e.g. >70 years) and even small woodlots (Boissinot et al. 2015), are essential for amphibian population persistence, in particular in agricultural regions (Hartel et al., 2010; Collins & Fahrig 2017).

Therefore, current methods for defining green infrastructure, based on landcover data, should be complemented by integrating biological data. An integrative approach using habitat suitability mapping combined with connectivity modelling, using biological observations, seems relevant for integrating species-habitat relations into ecological network definition at different spatial scales (Clauzel & Godet 2020).

4.4 **Implications for conservation**

Protected areas seem insufficient and poorly suited to maintaining amphibians regionally. Green infrastructure (GI) has better efficiency and coverage but is not regulatory (unprotected) and covers only 16 to 33% of highly suitable habitat. An important weakness of GI is its exclusive reliance on land use data without including biological data and landscape complementation as well as the multi-layered nature of ecological networks. Ecological niche modelling tools such as SDM, coupled with ecological connectivity models, are of particular interest for improving the ecological realism of identified networks and supporting related decisions. To make this possible, strengthening the interactions between researchers and stakeholders is essential, especially because these modelling tools need to be used and interpreted with care.

The second major result concerns bottlenecks. We identified "bottleneck" sectors not covered by any conservation tools such as PA or GI despite high potential conservation value. These areas might be especially important for amphibian populations in a context of global change requiring distribution shifts to maintain suitable conditions for population viability. Indeed, according to Préau et al. (2019), important distribution shifts at regional scale are expected for certain amphibian species based on global change predictions. These two "bottlenecks" are located in a context of open fields and intensive farming which are considered by nature protection organisations to have limited potential for biodiversity. In these areas, some semi-natural elements such as small woods associated with wetlands may nonetheless play a key role in maintaining functionality and connectivity of the network at regional scale and should be strengthened. In these two areas, habitat continuities are maintained by riparian woods (sometime very thin and degraded) and associated river landscapes with a higher proportion of pastures and hedgerows, as well as small wood patches close to wetlands including ponds. As a result, these two areas need further investigation and could be targeted by conservation and restoration measures. These results also underline the importance of multi-scale and multi-level analysis for habitat and connectivity studies (e.g., McGarigal et al. 2016).

Finally, the different maps obtained from this study provide support for prioritizing certain regional policies with different objectives: (1) strengthening PA network and GI, in particular in the 2 bottleneck sectors. (Appendix 5 provides more specific information to prioritize new protected area from a regional preselected set of sites according to our results); (2) to identify potential conflict points with transport network (i.e., roads and railways); (3) spatially prioritize actions supporting agroecological transition, particularly in the two bottleneck areas considered as "degraded" for biodiversity. Conservation actions for amphibian conservation often focus on ponds and pond networks. The complementarity of ponds (or wetlands) and forest patches should be considered to define conservation actions (Cushman, 2006; Denoël & Lehmann, 2006; Pope et al., 2000) as well as certain associated traditional farming practices in western Europe (Boissinot et al., 2019; Hartel et al., 2010). In our region, amphibians have interactions with traditional agricultural practices and with related landscape dynamics. The success of conservation measures depends on our ability to integrate biodiversity issues into human-dominated agricultural landscapes shaped by farming practices.

Acknowledgements

This work would not have been possible without the support of the Pays de la Loire Herpetological Group, the CPIE Regional Union and the French BirdLife partner (LPO). We are especially grateful to Morgane Sineau and Benoit Marchadour who coordinate regional naturalists' databases. We also acknowledge the many naturalists involved, for access to data and for additional fieldwork, especially Dorian Angot, Baptiste Gaboriau and Martin Bonhomme. We thank Jean Secondi and Aurélien Besnard for providing helpful discussion. Our work was supported by funding from Ecole Supérieure d'Agricultures d'Angers, Angers Loire Metropole, The French Society for Ecology and Evolution (SFE[2]) and "Humanité et Biodiversité".

Authors' contributions

Florence Matutini: Conceptualization (Equal), Data curation (Lead), Formal analysis (Lead), Funding acquisition (Supporting), Methodology (Lead), Investigation (Lead), Visualization (Lead), Writing-original draft (Lead), Writing-review & editing (Lead); **Marie-Josée Fortin:** Conceptualization (Equal), Methodology (Supporting), Supervision (Supporting), Writing-original draft (Supporting), Writing-review & editing (Supporting); **Jacques Baudry, Guillaume Pain & Josephine Pithon:** Conceptualization (Equal), Methodology (Supporting), Supervision (Equal), Writing-original draft (Supporting), Writing-review & editing (Supporting)

References

Ahmadi, M., Farhadinia, M. S., Cushman, S. A., Hemami, M. R., Nezami Balouchi, B., Jowkar, H., & Macdonald, D. W. (2020). Species and space: a combined gap analysis to guide management planning of conservation areas. Landscape Ecology, 35(7), 1505–1517. https://doi.org/10.1007/s10980-020-01033-5

Baudry, J., Bunce, R. G. H., & Burel, F. (2000). Hedgerow diversity : An international perspective on their origin , function and management. March. https://doi.org/10.1006/jema.2000.0358

Bazin, P., & Schmutz, T. (1994). La mise en place de nos bocages en europe et leur déclin. Revue Forestière Française, S, 115. https://doi.org/10.4267/2042/26606

Boissinot, A., Besnard, A., & Lourdais, O. (2019). Amphibian diversity in farmlands: Combined influences of breeding-site and landscape attributes in western France. Agriculture, Ecosystems & Environment, 269(September 2018), 51–61. https://doi.org/10.1016/j.agee.2018.09.016

Boissinot, A., Grillet, P., Besnard, A., & Lourdais, O. (2015). Small woods positively influence the occurrence and abundance of the common frog (Rana temporaria) in a traditional farming landscape. Amphibia Reptilia, 36(4), 417–424. https://doi.org/10.1163/15685381-00003013

Burel, F., & Baudry, J. (1995). Social, aesthetic and ecological aspects of hedgerows in rural landscapes as a framework for greenways. Landscape and Urban Planning, 33(1–3), 327–340. https://doi.org/10.1016/0169-2046(94)02026-C

Calabrese, J. M., Certain, G., Kraan, C., & Dormann, C. F. (2014). Stacking species distribution models and adjusting bias by linking them to macroecological models. Global Ecology and Biogeography, 23(1), 99–112. https://doi.org/10.1111/geb.12102

Chatzimentor, A., Apostolopoulou, E., & Mazaris, A. D. (2020). A review of green infrastructure research in Europe: Challenges and opportunities. Landscape and Urban Planning, 198(March), 103775. https://doi.org/10.1016/j.landurbplan.2020.103775

Clauzel, C., & Godet, C. (2020). Combining spatial modeling tools and biological data for improved multispecies assessment in restoration areas. Biological Conservation, 250(August), 108713. https://doi.org/10.1016/j.biocon.2020.108713

Collins, J. P., & Storfer, A. (2003). Global amphibian declines: Sorting the hypotheses. Diversity and Distributions, 9(2), 89–98. https://doi.org/10.1046/j.1472-4642.2003.00012.x

Collins, S. J., & Fahrig, L. (2017). Responses of anurans to composition and configuration of agricultural landscapes. Agriculture, Ecosystems and Environment, 239, 399–409. https://doi.org/10.1016/j.agee.2016.12.038

Cushman, S. A. (2006). Effects of habitat loss and fragmentation on amphibians: A review and prospectus. Biological Conservation, 128(2), 231–240. https://doi.org/10.1016/j.biocon.2005.09.031

De Solla, S. R., Shirose, L. J., Fernie, K. J., Barrett, G. C., Brousseau, C. S., & Bishop, C. a. (2005). Effect of sampling effort and species detectability on volunteer based anuran monitoring programs. Biological Conservation, 121, 585–594. https://doi.org/10.1016/j.biocon.2004.06.018

DeMaynadier, P. G., & Hunter, M. L. (2008). Effects of Silvicultural Edges on the Distribution and Abundance of Amphibians in Maine. Conservation Biology, 12(2), 340–352. https://doi.org/10.1111/j.1523-1739.1998.96412.x

Denoël, M., & Lehmann, A. (2006). Multi-scale effect of landscape processes and habitat quality on newt abundance: Implications for conservation. Biological Conservation, 130(4), 495–504. https://doi.org/10.1016/j.biocon.2006.01.009

Díaz-García, J. M., Pineda, E., López-Barrera, F., & Moreno, C. E. (2017). Amphibian species and functional diversity as indicators of restoration success in tropical montane forest. Biodiversity and Conservation, 26(11), 2569–2589. https://doi.org/10.1007/s10531-017-1372-2

Dirzo, R., Young, H. S., Galetti, M., Ceballos, G., Isaac, N. J. B., & Collen, B. (2014). Defaunation in the Anthropocene. Science, 345(6195), 401–406. https://doi.org/10.1126/science.1251817

Duflot, R., Avon, C., Roche, P., & Bergès, L. (2018). Combining habitat suitability models and spatial graphs for more effective landscape conservation planning: An applied methodological framework and a species case study. Journal for Nature Conservation, 46(May 2017), 38–47. https://doi.org/10.1016/j.jnc.2018.08.005

Ferrier, S., & Guisan, A. (2006). Spatial modelling of biodiversity at the community level. Journal of Applied Ecology, 43(3), 393–404. https://doi.org/10.1111/j.1365-2664.2006.01149.x

Foltête, J. C., Savary, P., Clauzel, C., Bourgeois, M., Girardet, X., Saharoui, Y., Vuidel, G., & Garnier, S. (2020). Coupling landscape graph modeling and biological data: a review. Landscape Ecology, 7. https://doi.org/10.1007/s10980-020-00998-7

Gaston, K. J., & Fuller, R. A. (2008). Commonness, population depletion and conservation biology. Trends in Ecology and Evolution, 23(1), 14–19. https://doi.org/10.1016/j.tree.2007.11.001

Godet, C., & Clauzel, C. (2020). Comparison of landscape graph modelling methods for analysing pond network connectivity. Landscape Ecology, 9. https://doi.org/10.1007/s10980-020-01164-9

Guisan, A., & Zimmermann, N. E. (2000). Predictive habitat distribution models in ecology. Ecological Modelling, 135(2–3), 147–186. https://doi.org/10.1016/S0304-3800(00)00354-9

Hamer, A. J., & McDonnell, M. J. (2008). Amphibian ecology and conservation in the urbanising world: A review. Biological Conservation, 141(10), 2432–2449. https://doi.org/10.1016/j.biocon.2008.07.020

Hartel, T., Schweiger, O., Öllerer, K., Cogălniceanu, D., & Arntzen, J. W. (2010). Amphibian distribution in a traditionally managed rural landscape of Eastern Europe: Probing the effect of landscape composition. Biological Conservation, 143(5), 1118–1124. https://doi.org/10.1016/j.biocon.2010.02.006

Jennings, M. D. (2000). Gap analysis: Concepts, methods, and recent results. Landscape Ecology, 15(1), 5–20. https://doi.org/10.1023/A:1008184408300

Joppa, L. N., & Pfaff, A. (2009). High and far: Biases in the location of protected areas. PLoS ONE, 4(12), 1–6. https://doi.org/10.1371/journal.pone.0008273

Keeley, A. T. H., Beier, P., & Gagnon, J. W. (2016). Estimating landscape resistance from habitat suitability: effects of data source and nonlinearities. Landscape Ecology. https://doi.org/10.1007/s10980-016-0387-5

562 Keeley, A. T. H., Beier, P., Keeley, B. W., & Fagan, M. E. (2017). Habitat suitability is a poor
563   proxy for landscape connectivity during dispersal and mating movements. Landscape and Urban
564   Planning, 161, 90–102. https://doi.org/10.1016/j.landurbplan.2017.01.007

565 Kleijn, D., Rundlöf, M., Scheper, J., Smith, H. G., & Tscharntke, T. (2011). Does conservation on
566   farmland contribute to halting the biodiversity decline? Trends in Ecology and Evolution, 26(9),
567   474–481. https://doi.org/10.1016/j.tree.2011.05.009

568 Koen, E. L., Bowman, J., Sadowski, C., & Walpole, A. A. (2014). Landscape connectivity for
569   wildlife: Development and validation of multispecies linkage maps. Methods in Ecology and
570   Evolution, 5(7), 626–633. https://doi.org/10.1111/2041-210X.12197

571 Lee☐Yaw, J. A., McCune, J. L., Pironon, S., & Sheth, S. N. (2022). Species distribution models
572   rarely predict the biology of real populations. Ecography, 2022(6), 1–16.
573   https://doi.org/10.1111/ecog.05877

574 Matutini, F., Baudry, J., Fortin, M. J., Pain, G., & Pithon, J. (2021). Integrating landscape
575   resistance and multi-scale predictor of habitat selection for amphibian distribution modelling at
576   large scale. Landscape Ecology, 36(12), 3557–3573. https://doi.org/10.1007/s10980-021-01327-
577   2

578 Matutini, F., Baudry, J., Pain, G., Sineau, M., & Pithon, J. (2021). How citizen science could
579   improve species distribution models and their independent assessment. Ecology and Evolution,
580   November 2020, ece3.7210. https://doi.org/10.1002/ece3.7210

581 Mazerolle, M. J. (2005). Drainage ditches facilitate frog movements in a hostile landscape.
582   Landscape Ecology, 20(5), 579–590. https://doi.org/10.1007/s10980-004-3977-6

583 Mcrae, B. H., Dickson, B. G., Keitt, T. H., Shah, V. B., Mcrae, B. H., Dickson, B. G., Keitt, T. H.,
584   & Shah, V. B. (2008). Using Circuit Theory to Model Connectivity in Ecology , Evolution , and
585   Conservation. Ecology, 89(10), 2712–2724.

586 Mcrae, B., Shah, V., & Edelman, A. (2016). Circuitscape: modeling landscape connectivity to
587   promote conservation and human health. The Nature Conservancy, 1–14.

588 Moilanen, A., Franco, A. M. A., Early, R. I., Fox, R., Wintle, B., & Thomas, C. D. (2005).
589   Prioritizing multiple-use landscapes for conservation: Methods for large multi-species planning
590   problems. Proceedings of the Royal Society B: Biological Sciences, 272(1575), 1885–1891.
591   https://doi.org/10.1098/rspb.2005.3164

592 Opdam, P., Steingröver, E., & Rooij, S. Van. (2006). Ecological networks: A spatial concept for
593   multi-actor planning of sustainable landscapes. Landscape and Urban Planning, 75(3–4), 322–
594   332. https://doi.org/10.1016/j.landurbplan.2005.02.015

595 Petrovan, S. O., & Schmidt, B. R. (2016). Volunteer conservation action data reveals large-scale
596   and long-term negative population trends of a widespread amphibian, the common toad (Bufo
597   bufo). PLoS ONE, 11(10), 1–12. https://doi.org/10.1371/journal.pone.0161943

598 Pilosof, S., Porter, M. A., Pascual, M., & Kéfi, S. (2017). The multilayer nature of ecological
599   networks. In Nature Ecology and Evolution (Vol. 1, Issue 4). Nature Publishing Group.
600   https://doi.org/10.1038/s41559-017-0101

601 Pope, S. E., Fahrig, L., & Merriam, H. G. (2000). Landscape complementation and metapopulation
602   effects on leopard frog populations. Ecology, 81(9), 2498–2508. https://doi.org/10.1890/0012-
603   9658(2000)081[2498:LCAMEO]2.0.CO;2

Rodrigues, A. S. L., & Cazalis, V. (2020). The multifaceted challenge of evaluating protected area
effectiveness. Nature Communications, 1–4. https://doi.org/10.1038/s41467-020-18989-2

Rodrigues, A. S. L., J, A., MI, B., L, B., M, B., RM, C., LD, F., GA, D. F., KJ, G., M, H., & JS., L.
(2004). Effectiveness of the global protected area network in representing species diversity.
Nature, 640–643. https://doi.org/10.1038/nature02459.1.

Salomaa, A., Paloniemi, R., Kotiaho, J. S., Kettunen, M., Apostolopoulou, E., & Cent, J. (2017).
Can green infrastructure help to conserve biodiversity? Environment and Planning C:
Government and Policy, 35(2), 265–288. https://doi.org/10.1177/0263774X16649363

Scherrer, D., D'Amen, M., Fernandes, R. F., Mateo, R. G., & Guisan, A. (2018). How to best
threshold and validate stacked species assemblages? Community optimisation might hold the
answer. Methods in Ecology and Evolution, 9(10), 2155–2166. https://doi.org/10.1111/2041-
210X.13041

Scherrer, D., Mod, H. K., & Guisan, A. (2020). How to evaluate community predictions without
thresholding? Methods in Ecology and Evolution, 11(1), 51–63. https://doi.org/10.1111/2041-
210X.13312

Schmitz, M. F., Herrero-Jáuregui, C., Arnaiz-Schmitz, C., Sánchez, I. A., Rescia, A. J., & Pineda,
F. D. (2017). Evaluating the Role of a Protected Area on Hedgerow Conservation: The Case of
a Spanish Cultural Landscape. Land Degradation and Development, 28(3), 833–842.
https://doi.org/10.1002/ldr.2659

Seibold, S., Gossner, M. M., Simons, N. K., Blüthgen, N., Müller, J., Ambarlı, D., Ammer, C.,
Bauhus, J., Fischer, M., Habel, J. C., Linsenmair, K. E., Nauss, T., Penone, C., Prati, D., Schall,
P., Schulze, E. D., Vogt, J., Wöllauer, S., & Weisser, W. W. (2019). Arthropod decline in
grasslands and forests is associated with landscape-level drivers. Nature, 574(7780), 671–674.
https://doi.org/10.1038/s41586-019-1684-3

Sewell, D., & Griffiths, R. A. (2009). Can a single amphibian species be a good biodiversity
indicator? Diversity, 1(2), 102–117. https://doi.org/10.3390/d1020102

Sinsch, U. (2014). Movement ecology of amphibians: from individual migratory behaviour to
spatially structured populations in heterogeneous landscapes 1 , 2. Canadian Journal of Zoology,
92(6), 491–502. https://doi.org/10.1139/cjz-2013-0028

Snäll, T., Lehtomäki, J., Arponen, A., Elith, J., & Moilanen, A. (2016). Green Infrastructure Design
Based on Spatial Conservation Prioritization and Modeling of Biodiversity Features and
Ecosystem Services. Environmental Management, 57(2), 251–256.
https://doi.org/10.1007/s00267-015-0613-y

Stuart, S. N., Chanson, J. S., Cox, N. A., Young, B. E., Rodrigues, A. S. L., Fischman, D. L., &
Waller, R. W. (2004). Status and trends of amphibian declines and extinctions worldwide.
Science, 306(5702), 1783–1786. https://doi.org/10.1126/science.1103538

Thuiller, W., Pollock, L. J., Gueguen, M., & Münkemüller, T. (2015). From species distributions to
meta-communities. 18(12), 1321–1328. https://doi.org/10.1111/ele.12526.

Fig. 1. Identification of highly suitable and suitable multi-specific patches of habitat using stacked habitat suitability map for three species groups.

Legend

- ■ Group 1 - high conservation level (UICN class from I to IV)
- ■ Group 2 (UICN class V)
- ■ Group 3 (Natura 2000)
- □ All region area (no protected area)

Fig. 2. Distribution of pixels along the regional habitat suitability gradient covered or not by protected area. Suitability index (from the habitat suitability map) is calculated for each species and varies from 0 to 1. Suitability index of the three groups (ALL, FOR and GEN) is the sum of the individual species index related to each group (pSSDM). The frequency corresponds to a number of pixels covered by protected areas from the Group 1, 2, 3 in relation to the whole region area.

Fig. 3. Large scale permeability between high suitability areas and associated continuities patterns confronted with regional green infrastructure. A and B show potential conflict area and issues for continuity conservation at regional scale with a mismatch with the green infrastructure network.

Table 1. Description of the nine studied species: conservation status and number of presence-only data available for analysis. RL FR: France red list; RL Region: regional red list of Pays-de-la-Loire; Priority: regional priority level for species conservation from the regional red list defined by Marchadour et al. (2009); Hab. Dir. N2000: habitats directive classification (Natura 2000); Pop trends: regional population trends from Marchadour et al. (2009); Nb presence data: Number of presence-only data were calculated after a 100 m-resolution rasterization and correspond to the number of pixels with at least one observation of the species.

	Species	RL FR (2008)	RL Region (2009)	Priority (region)	Hab. Dir. N2000	Pop trends (1998-2008)	Nb presence data (Atlas)
Species used for SSDM	*Pelodytes punctatus*	LC	LC	2		↘	1714
	Triturus marmoratus	LC	LC	3	IV	↘	906
	Rana temporaria	LC	VU	2	V	↘	848
	Lissotriton helveticus	LC	LC	1		?	4580
	Bufo spinosus	LC	LC	0		?	6071
	Hyla arborea	LC	LC	0	IV	↘	4555
	Rana dalmatina	LC	LC	0	IV	?	6118
	Triturus cristatus	LC	LC	0	II & IV	↘	1040
	Salamandra salamandra	LC	LC	0		?	3514

Table 2. Classification of main conservation areas according to protection levels used in Pays-de-la-Loire (France). PA: protected area; GI: Green infrastructure; INV: area with an inventory of biodiversity and having a strong natural heritage interest but not protected by regulatory tools (ZNIEFF, type I and II).

Cat.	Group	Protect. level	Code name	UICN class	Protection approaches	Level	French denomination
PA	1	+++	RNN	I-IV	Regulatory	National	National Nature Reserve
			SIN,SCL	III	Regulatory	National	Classified site and Registred site
			RNR,RB, SCL,APB	I-V	Regulatory essentially	National or Regional	Biological Reserve, Conservatoire du littoral Site, Regional Nature Reserve, Biotope Protection Order
	2		SCEN, ENS	I-VI	Regulatory, Contractual management or Land purchase and management	Regional	Area managed by Natural Area Conservation Societies (CENs)
			PNR	V	Contractual management	Regional	Regional Nature Park
	3	+	N2000	/	Contractual management	European	Natura 2000 sites (N2000)
GI	4	0	BOCAGE	/	Non-regulatory Advisory / informative	National to local	Traditional hedgerow mosaic landscapes Infrastructure
		0	WOOD	/			Woodlands infrastructure
		0	WET	/			Wetlands infrastructure
		0	COAST	/			Coastlands infrastructure
		0	OPEN	/			Natural open fields infrastructure
		0	Corridor	/			Ecological cooridors
INV	5	0	ZNIEFF	/	Biodiversity inventories only	National	Natural areas of special ecological, faunal and floral value

Table 3. Accuracy of stacked species distribution models for three species groups according to the nine studied amphibian species. Results obtained with external data with 100 permutations. Community AUC were obtained using species diversity results for each studied species group and for all species inventoried in the external dataset (n=17). Individual AUC and SEDI are from a previous study in Matutini, Baudry, Fortin, et al. (2021).

Species group	R²	cAUC	Min individual AUC / SEDI	Max individual AUC / SEDI	R^2 all
All nine species	0.31	0.86	0.73 / 0.67	0.86 / 0.89	0.07
Forest species (n=3)	0.23	0.76	0.78 / 0.68	0.86 / 0.84	0.14
Generalist species (n=5)	0.30	0.87	0.73 / 0.67	0.81 / 0.89	0.02

Table 4. Proportion of the conservation area covered by different habitat suitability classes (1. conservation efficiency) and proportion of different habitat suitability classes covered by the conservation area (2. conservation coverage) for different species and species groups. Classes of conservation area: Protected (groups 1 to 3) – classes I to V derived from the UICN classification; EURO: European conservation network from Natura 2000; Non-protected: GI (Green infrastructure) and INV (inventoried sites define as "high naturalist interest"). Suitability class for each groups or species: "+++" = highly suitable (>0.80) ; "++" = suitable (>0.65) ; +/- = "moderately or not suitable" (<=0.55). See Table 2 for "code names" details. Names with * show sites with point data only (surface area is not available).

1) AnB / A — CONSERVATION EFFICIENCY

	Group	Name	Class	Area (ha)	All studied species (n=9) +++	++	+/-	Forest species (n=3) +++	++	+/-	Generalist species (n=5) +++	++	+/-	Triturus marmoratus +++	++	+/-	Rana temporaria +++	++	+/-	Pelodytes punctatus +++	++	+/-
Protected (PA)	1	RNN	I-IV	3183	0	23	26	0	19	26	0	21	24	0	3	21	0	4	12	12	1	4
	1	SIN,SCL	III	80912	7	33	36	7	11	18	12	35	29	8	13	19	4	2	3	24	18	20
	1	RNR,RB,SCL,APB	I-V	4902	17	30	28	20	12	8	24	32	28	3	12	17	22	4	2	14	11	15
	2	SCEN,ENS*	I-VI	/	12	50	26	12	24	23	26	44	19	8	21	20	6	7	7	6	10	15
	2	PNR	V	358668	11	28	27	15	15	10	13	32	24	5	10	13	15	8	4	10	12	24
	3	N2000	EURO	144070	10	33	36	10	16	12	19	34	26	2	10	23	10	8	6	24	14	13
Non protected (GI & INV)	4.1	BOCAGE	GI	424568	20	63	17	21	26	29	42	43	13	12	27	30	15	11	9	0	3	10
	4.1	WOOD	GI	176157	26	55	18	38	25	23	38	38	20	6	20	22	31	12	11	0	1	4
	4.1	WET	GI	270266	19	56	22	21	25	25	38	44	15	5	18	26	15	11	11	1	3	8
	4.1	COAST	GI	9376	2	51	35	0	12	30	4	59	26	12	34	25	0	0	0	13	20	26
	4.1	OPEN	GI	19445	15	53	28	11	22	20	42	41	13	1	5	11	13	11	12	1	5	17
	4.1	Total core hab.	GI	650572	20	60	19	24	25	27	40	43	15	9	24	26	18	10	9	0	3	9
	4.2	Corridor	GI	378391	17	52	25	13	21	26	39	41	16	5	14	20	9	8	9	0	4	14
	5	ZNIEFF	INV	279266	20	57	20	29	24	25	34	44	18	11	28	25	21	9	10	1	4	10
REGION				3236407	12	49	28	11	17	23	33	40	17	5	13	19	9	7	7	3	6	15

2) AnB / B — CONSERVATION COVERAGE

	Group	Name	Class	All studied species (n=9) +++	++	+/-	Forest species (n=3) +++	++	+/-	Generalist species (n=5) +++	++	+/-	Triturus marmoratus +++	++	+/-	Rana temporaria +++	++	+/-	Pelodytes punctatus +++	++	+/-
Protected	1	RNN	I-IV	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1	SIN,SCL	III	1	2	3	1	2	2	1	2	4	4	2	2	1	1	1	19	8	3
	1	RNR,RB,SCL,APB	I-V	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
	2	SCEN,ENS*	I-VI	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/
	2	PNR	V	10	6	10	14	10	5	4	9	15	11	9	8	20	13	6	35	23	18
	3	N2000	EURO	4	3	6	4	4	2	3	4	7	2	3	5	5	5	4	35	11	4
Non protected	4.1	BOCAGE	GI	21	17	8	24	20	17	16	14	10	33	27	20	23	21	17	2	8	9
	4.1	WOOD	GI	11	6	3	18	8	5	6	5	6	7	8	6	20	9	9	1	1	2
	4.1	WET	GI	13	10	7	16	12	9	10	9	7	8	12	11	15	14	14	2	4	5
	4.1	COAST	GI	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	1	1	1
	4.1	OPEN	GI	1	1	1	1	1	1	1	1	0	0	0	0	1	1	1	0	0	1
	4.1	Total core hab.	GI	33	25	13	42	30	24	24	21	17	39	36	28	43	32	28	3	11	13
	4.2	Corridor	GI	16	13	10	14	15	13	14	12	10	12	12	12	12	15	16	2	9	11
	5	ZNIEFF	INV	14	10	6	22	12	10	9	9	9	21	18	11	21	12	12	2	6	6