Experimental infection of Mexican free-tailed bats (*Tadarida brasiliensis)* with SARS-CoV-2.

Hall JS[1,3], Hofmeister E[1], Ip HS[1], Nashold SW[1], Leon AE[1], Malavé CM[1], Falendysz EA[1], Rocke

TE[1], Carossino M[2], Balasuriya U[2], Knowles S[1]

[1]U.S. Geological Survey, National Wildlife Health Center, Madison, Wisconsin

[2]Louisiana Animal Disease Diagnostic Laboratory and Department of Pathobiological

Sciences, School of Veterinary Medicine, Louisiana State University, Baton Rouge, Louisiana

[3]Corresponding author

Abstract.

The severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) virus originated in wild

bats from Asia, and as the resulting pandemic continues into its third year, concerns have been

raised that the virus will expand its host range and infect North American wildlife species,

including bats. Mexican free-tailed bats (*Tadarida brasiliensis*: TABR) live in large colonies in

the southern United States, often in urban areas, and as such, could be exposed to the virus from

infected humans. We experimentally challenged wild TABR with SARS-CoV-2 to determine the

susceptibility, reservoir potential, and population impacts of infection in this species. Of nine

bats oronasally inoculated with SARS-CoV-2, five became infected and orally excreted moderate

amounts of virus for up to 18 days post inoculation. These five subjects all seroconverted and

cleared the virus before the end of the study with no obvious clinical signs of disease. We

additionally found no evidence of viral transmission to uninoculated subjects. These results

indicate that while TABR are susceptible to SARS-CoV-2 infection, infection of wild

populations of TABR would not likely cause mortality. However, the transmission of SARS-

CoV-2 from TABR to or from humans, or to other animal species, is a distinct possibility

requiring further investigation to better define.

Introduction.

As we enter the third year of the severe acute respiratory syndrome coronavirus-2

(SARS-CoV-2) pandemic, many unanswered questions remain regarding the ecology of the

virus. How the virus interacts with wild species is critical knowledge to obtain, including

whether: (1) North American wildlife can act as reservoirs of the virus, (2) the virus can adapt

genetically to new hosts and become more virulent, and (3) the virus can affect the health of wild

populations, particularly threatened or endangered species.

SARS-CoV-2 naturally infects several wild species including captive wild animals. For

example, American mink (*Neovison vison*) were infected with SARS-CoV-2 in the United States

in proximity to domestic mink farm operations (Shriner et al. 2021). Antibodies to SARS-CoV-2

were detected in white-tailed deer (*Odocoileus virginianus*) indicating exposure to the virus

(Chandler et al. 2021), and infection was later confirmed in this species (Hale et al. 2022).

Captive wild animals in zoos, particularly members of the Felidae family, were infected with the

virus (McAloose et al. 2020), and several North American species have experimentally been

shown susceptible to the virus, including deer mice (*Peromyscus maniculatus*), striped skunks

(*Mephitis mephitis*), and bushy-tailed woodrats (*Neotoma cinerea*) (Bosco-Lauth et al. 2021).

Because evidence indicates that SARS-CoV-2 originated in Asian bats (Zhou et al. 2020;

Zhou et al. 2021; Wacharapluesadee et al. 2021), concern that the virus could infect North

American bat species has been raised, particularly for bat populations under severe threat from

another pathogen, *Pseudogymnoascus destructans* (Cheng et al. 2021; Hoyt et al. 2021). Whether

North American bats could provide a reservoir of the virus and additional routes of transmission

to humans and other susceptible species, as well as any effects of SARS-CoV-2 on populations

are important to determine, as are any management measures that could be used to help protect

these populations.

Big brown bats (*Eptesicus fuscus*) were previously challenged with SARS-CoV-2 and

demonstrated resistance to infection (Hall et al. 2021). This species often encounters humans, as

they frequently reside in anthropogenic structures, including occupied homes and other

buildings. Another common North American bat species, the Mexican free-tailed bat, (TABR:

Tadarida brasiliensis), resides in very large colonies in the southern United States, often in urban

areas. This species is migratory and if susceptible to SARS-CoV-2, could transport the virus

to/from Central and South America on their migratory routes. In this study we challenged TABR

with SARS-CoV-2 to determine their susceptibility to infection, reservoir potential, the

adaptability of the virus to a new potential host, and potential effects of the virus on their

populations.

Materials and methods.

Virus acquisition and propagation. We obtained the SARS-CoV-2 virus (2019-nCoV/USA-

WA1/2020) from BEI Resources, National Institute of Allergy and Infectious Diseases (NIAID),

National Institutes of Health (NIH) (Manassas, Virginia)The virus was isolated from the first

confirmed patient with coronavirus disease 2019 (COVID-19) in the United States (Harcourt et

al. 2020)**.** We propagated and quantified the virus in Vero E6 cell culture using standard

techniques.

Animal acquisition and husbandry. All husbandry and experimental protocols were

approved by the U.S. Geological Survey (USGS) National Wildlife Health Center (NWHC)

Institutional Animal Care and Use Committee. Wild TABR were captured in Williamson

County, Texas in August 2021. A mixture of adult and juvenile male bats was collected and

immediately placed into a temperature-controlled chamber maintained at approximately 20°C.

This temperature induced the bats to enter torpor during transport to the NWHC, Madison,

Wisconsin.

On arrival at the NWHC, the bats were given veterinary examinations and treated

topically with selamectin for parasites (Zoetis, Florham Park, New Jersey). The bats were hand

fed mealworms (*Tenebrio molito*) supplemented with a vitamin and mineral mixture, and water

was provided *ad libitum.* Bats underwent a quarantine and acclimatization period of 30 days

prior to commencement of this study during which time the bats learned to feed themselves.

Pre-inoculation fecal sampling and coronavirus analysis. During the acclimatization period, we

collected fecal samples from the individual bats to determine the presence of other coronaviruses

in these subjects. Each fecal sample was suspended 10% (w/v) in viral transport medium (VTM;

Hanks Balanced Salt Solution, 0.05% gelatin, 5% glycerin, 1500 units/ml penicillin, 1500 mg/ml

streptomycin, 0.1 mg/ml gentamicin,1 mg/ml fungizone). Viral RNA was extracted using the

MagMax Pathogen RNA/DNA kit (Applied Biosystems, Forest City, California) on a Kingfisher

Flex magnetic particle processor according to the manufacturer's instructions. The presence of

coronaviruses was determined using methods previously described (Decaro and Larusso, 2020)**.**

Virus inoculation. Experimental inoculations were performed under Biosafety Level-3

conditions at the NWHC. We utilized 21 male Mexican free-tailed bats after the acclimatization

period and pairs of bats were cohoused in mesh cages. One bat from each of nine bat pairs was

inoculated with SARS-CoV-2, and its cagemate was left uninoculated to determine if the virus

could be horizontally transmitted between bats. One bat was inoculated and housed individually.

The SARS-CoV-2 inoculum dose was 10^5 TCID$_{50}$/bat and was administered nasally (4 µl) and

orally (6 µl) using a micropipette. One bat pair was sham inoculated with the same volume of

VTM. This technique has been used to inoculate other species with SARS-CoV-2 (Munster et al.

2020; Schlottau et al. 2020; Shi et al. 2020; Hall et al. 2021). The inoculum titer was verified by

quantitative reverse transcription-polymerase chain reaction (qRT-PCR) as described below and

virus viability confirmed in cell culture using Vero E6 cells.

Animal monitoring and sampling. Bats were observed at least twice daily to monitor health

status and document development of clinical signs. Just prior to inoculation and every other day

thereafter, each bat was weighed, and oropharyngeal and rectal swabs (Puritan Medical Products,

Guilford, Maine) were collected and placed in 0.5 ml VTM. On day post-inoculation (DPI) 7 and

on DPI 14, bats from one cage (one inoculated bat, one uninoculated) were euthanized, a

postmortem examination was conducted, and tissues and blood collected. At the end of the study

(DPI 20), all remaining bats were euthanized and postmortem examinations were completed for

the control bats and an additional cage pair. Blood was collected for serological analyses from all

euthanized bats.

qRT-PCR analyses. RNA extractions of swab material were performed in 96-well plates using

Mag Max-96 AI/ND Viral RNA Isolation Kit (Applied Biosystems, Foster City, California)

following the manufacturer's instructions. A positive control sample consisting of a 1:100

dilution of the SARS-CoV-2 inoculum used in the study was included with each extraction series

to validate successful RNA extraction. qRT-PCR analyses were conducted utilizing the Centers

for Disease Control 2019-nCoV N1 primers and probe (https://www.cdc.gov/coronavirus/2019-

ncov/lab/rt-pcr-panel-primer-probes.html) and AgPath-ID One-Step RT-PCR reagents

(Ambion/ThermoFisher, Waltham, Massachusetts). We included a standard curve of serial

dilutions of RNA extracted from SARS-CoV-2 virus stock (10^7 $TCID_{50}$/ml) in each qRT-PCR

assay to quantify viral amounts.

Rabies Diagnostics. Brain tissue was assessed for rabies infection using the direct fluorescent

antibody test (DFA). After brain impressions were fixed in acetone, slides were stained with a

FITC-labelled monoclonal antibody (mAB) conjugate (Fujirebio U.S. Inc., Malvern,

Pennsylvania, USA) and visualized under a fluorescent microscope (Dean et al.1996).

Necropsy and histopathology. Two animals (inoculated and uninoculated cagemates) were

euthanized at DPI 7 (bats 127, 128) and DPI 14 (bats 103, 104), and an additional 2 sets of

cagemates at DPI 20 (bats 109, 110, 111, 117), using an overdose of isoflurane with subsequent

decapitation. Two uninoculated control animals (102, 108) were also euthanized at DPI 20.

These subjects were immediately necropsied after euthanasia and body condition and gross

observations were recorded. Portions of the nares, caudal lung, cranial lung, heart, liver, spleen,

kidney, small intestine, colon and brain were collected and saved frozen at -80°C for virological

analyses. Additional tissue portions were fixed in 10% neutral buffered formalin for histological

analysis. For histopathological examination, fixed tissues were processed routinely, sectioned at

approximately 5 µm and stained with hematoxylin and eosin at the Wisconsin Veterinary

Diagnostic Laboratory (Madison, Wisconsin). At DPI 20, all remaining bats were euthanized,

serum was collected, and all bat carcasses were saved frozen. Three bats (118, 123, 124) that

were saved frozen and later shown to be infected by swab analysis were subsequently necropsied

and sampled.

SARS-CoV-2-specific immunohistochemistry (IHC). For IHC, 4 µm sections of formalin-fixed

paraffin-embedded tissue were mounted on positively charged Superfrost® Plus slides and

subjected to IHC using an anti-nucleocapsid rabbit monoclonal antibody (HL344, Cell Signaling

Technology, Danvers, Massachusetts). IHC was performed using the automated BOND-RXm

platform and the Polymer Refine Red Detection kit (Leica Biosystems, Wetzlar, Germany).

Following automated deparaffinization, heat-induced epitope retrieval (HIER) was performed

using a ready-to-use citrate-based solution (pH 6.0; Leica Biosystems) at 100°C for 20 min.

Sections were then incubated with the primary antibody (diluted at 1:1,600 in primary antibody

diluent [Leica Biosystems]) for 30 min at room temperature, followed by a polymer-labeled goat

anti-rabbit IgG coupled with alkaline phosphatase (30 min). Fast Red was used as the chromogen

(15 minutes), and counterstaining was performed with hematoxylin for 5 min. Slides were dried

in a 60 °C oven for 30 min and mounted with a permanent mounting medium (Micromount®,

Leica Biosystems). Lung sections from a SARS-CoV-2-infected hamster were used as positive

assay controls.

***Virus RNA extraction and qRT-PCR from bat tissues*.** Approximately 10 mg of each tissue was

macerated in extraction buffer and RNA extracted using the ZYMO Research Quick DNA/RNA

Pathogen Miniprep kit (ZYMO Research, Irvine, California) according to the manufacturer's

directions. qRT-PCR analyses were performed as described above.

Antibody detection. To detect neutralizing antibodies to SARS-CoV-2, bat sera were screened at

a 1:10 dilution using a competitive enzyme linked immunosorbent assay (SARS-CoV-2 sVNT,

GenScript, Piscataway, New Jersey) according to the manufacturer's instructions. As directed, a

reduction in optical density (OD) of $\geq 30\%$ compared to the mean OD of the negative control

was considered positive for the presence of neutralizing antibodies. In addition to the positive

control provided in the kit, we used positive guinea pig serum obtained through BEI Resources,

NIAID, NIH: Polyclonal Anti-SARS Coronavirus antiserum (Guinea Pig, NR-10361). To

determine neutralizing antibody titers from positive sera, samples were two-fold serially diluted

and titers recorded as the reciprocal of the end-point dilution where the serum was no longer

considered positive.

Virus recovery and whole genome sequencing. Oral swab VTM from Bat 118 DPI 8 was

inoculated into Vero E6 cells and incubated at 37 °C, 5% CO_2 for 7 days. The flasks were

examined daily for cytopathic effects. Cell lysates collected after three cycles of freezing and

thawing were used for RNA extraction and serial passage. Extracted RNA was converted to

cDNA with SuperScript IV (ThermoFisher, Waltham, Massachusetts) or Maxima H minus

(ThermoFisher, Waltham, Massachusetts) reverse transcriptase according to manufacturer's

instructions. Tiled amplicon sequencing by the ARTIC method (Tyson et al. 2020) was

performed using Oxford Nanopore Technology's MinION running on a MK1C

instrument. Bioinformatic analysis was performed using the CLC Genomics Workbench v22

(Qiagen, Redwood City, California) using a publicly available workflow

(https://storage.googleapis.com/theiagen-resources/qiagen/SARS-CoV-2_Tutorial.zip).

200
Results.

Presence of coronaviruses in Mexican free-tailed bats prior to inoculation. RT-PCR analyses

of fecal material collected from the TABR prior to virus challenge revealed no evidence of

alpha- or betacoronavirus infection, in any subject (Supplemental Table 1).

Rabies virus infection. Prior to SARS-CoV-2 inoculation, one bat exhibited loss of appetite,

aggressive behavior towards its cagemate and weight loss over 3-4 days. We euthanized this bat

and tested for rabies. It was positive for the presence of rabies virus by DFA, therefore we tested

all the remaining TABR after euthanasia and all were negative for the presence of rabies virus

(Supplemental Table 1).

SARS-CoV-2 excretion after experimental inoculation. qRT-PCR analysis of oral swabs is

shown in Table 1. Of the 10 SARS-CoV-2 inoculated TABR, five (Bats 104, 110, 118, 124, 123)

excreted detectable viral RNA. Two additional bats (111, 129) had high cycle threshold (C_t)

readings only on the first sampling after inoculation (DPI 2) and likely was detection of residual

inoculum. The duration of oral excretion ranged from DPI 6 (Bat 123), up to 18 DPI (Bat 118).

The maximum amount of virus detected on an oral swab was 4.82×10^3 $TCID_{50}$ equivalent/ml (bat

118, DPI 4). In contrast to oral excretion, no virus was detected in rectal swabs from any TABR (Table

2). The effect of the bat's age on susceptibility was inconclusive as four of the infected bats were adults

and one was a juvenile, but the numbers were too small to be reliable.

An oral swab from a positive bat (Bat 118 DPI 8) was inoculated into Vero E6 tissue culture to

verify viral viability. This resulted in positive virus isolates from the first and third passage that were

confirmed to be SARS-CoV-2 by qRT-PCR analyses as described above. Whole genome sequencing of

these isolates revealed that they shared the same genetic changes as the WA-1 isolate used as inoculum,

when compared to the original Wuhan SARS-CoV-2 isolate. Two additional, consistent changes in both

passages of the Bat 118 DPI 8 isolate were found in noncoding regions that were not in the WA-1

inoculum isolate. These are shown in Supplemental Table 2.

Serological analyses. SARS-CoV-2 antibodies were detected by competitive Enzyme Linked

Immunosorbent Assay (cELISA) in five TABR with percent inhibitions ranging from about 34 –

79% (Table 3). These five bats were the same five that were orally excreting virus detected by

qRT-PCR (Table 1). These seropositive bat sera were subsequently tested at dilutions from 1:20

– 1:80. One serum (110) was positive only at the original 1:10 dilution. Two sera (123, 124)

were positive at 1:20 whereas two sera (104, 118) remained positive at 1:40 dilutions.

Clinical signs of infection, postmortem examination and histopathology. Over the three-week

course of this study, no overt clinical signs of SARS-CoV-2 disease were observed in 18/21 bats,

including the five bats that became infected and were orally excreting virus. These bats

maintained or gained weight during the study (Table 4) and appeared healthy. One inoculated bat

(127) presented with lethargy, decreased respiratory rate, and increased respiratory effort; and

was euthanized on DPI 6, along with its cagemate (128). An additional bat (112) presented with

lethargy, obtundation, weight loss, hypersalivation, inability to swallow, and respiratory distress;

and was euthanized at DPI 10.

Upon further examination, the bat that presented with respiratory difficulty (Bat 127) and

its cagemate (128) were in poor or emaciated body condition, respectively, with minimal or no

fat stores. Bat 112, euthanized due to respiratory distress and hypersalivation, was in fair body

condition. All other bats were in good body condition evidenced by moderate to abundant fat

stores. Gross findings included clear nasal discharge (112, 127), red foci or mottling in the lungs

(all cases except 127), pulmonary congestion (102-104, 008-011, 117, 118, 128), intestinal pallor

(102, 108, 109, 111, 117, 123, 124, 128), small foci of depigmentation of the patagia (103, 108,

111, 118, 123, 124, 127) and patagial tears (112, 128).

Histopathologic findings included pulmonary congestion, hemorrhage and alveolar

collapse (all non-frozen cases), meningeal hemorrhage (103, 104, 111, 112, 127), pulmonary

vascular thrombosis (112), large numbers of bacterial rods in bronchial epithelium and cardiac

valve and parenchyma (127), pale cardiomyofibers (110, 112, 127), nasal cavity hemorrhage

(102, 104, 109, 110, 111, 117, 128), necroulcerative and suppurative pharyngitis with

intralesional bacteria (112), exocytosis of neutrophils in nasal turbinate epithelium (104), non-

suppurative dermatitis (102, 108, 109, 110, 117), non-suppurative dermal myositis (102),

mononuclear cells in hepatic sinusoids (108, 110, 111, 128), trematodes in the lumen of the bile

ducts (110, 118) or gall bladder (118, 127), bile duct hyperplasia (110, 118), intestinal

nematodiasis (104, 118, 127), intestinal cestodiasis (117, 127), intestinal trematodiasis (118), and

intestinal luminal hemorrhage (127). The three additional bats (118, 123, 124) that excreted virus

were thawed, necropsies were performed, and tissues were collected. The histologic evaluation

of the lungs in these cases was severely hindered by freeze artifact.

These histopathologic findings observed for all examined bats were consistent with the

euthanasia procedures utilized, parasitism, or bacterial infections, and were not consistent with

findings observed in other animals experimentally infected with SARS-CoV-2 (Munster et al.

2020; Schlottau et al. 2020; Shi et al. 2020).

Immunohistochemistry for detection of SARS-CoV-2 antigen. Sections from the rostral nasal

cavity, lung, heart, spleen, liver, pancreas, stomach, small and large intestine, and brain from a

total of 14 experimentally and mock inoculated bats were subjected to immunohistochemistry for

the detection of SARS-CoV-2 antigen. No viral antigen was detected in any of the tissues

examined from these bats (Supplemental Table 3).

SARS-CoV-2 in bat tissues. qRT-PCR analyses of tissues collected from the infected and control

bats did not detect viral RNA in any tissue from any of the bats examined (Table 5).

Discussion.

We experimentally challenged Mexican free-tailed bats with SARS-CoV-2. Of the ten

inoculated bats, five orally excreted virus for 6-18 days post challenge. These five also mounted

an immune response and cleared the virus before the end of the study. Based on these findings

we concluded that Mexican free-tailed bats are susceptible to infection by SARS-CoV-2.

We found no evidence of transmission between cohoused TABR. Five of the ten (50%)

inoculated bats became infected indicating that our inoculum titer (10^5 $TCID_{50/}$ dose) was

apparently near the 50% infectious dose for this species. However, the largest amount of virus

excreted by infected bats was between 10^3 and 10^4 $TCID_{50}$ equivalents/ml. Thus, based on these

data, contact transmission between TABR would be unlikely.

Another possible reason for no transmission between TABR was the way we housed the

bats during the challenge study. In the wild, TABR roost in very dense colonies that aggregate in

large numbers in natural and anthropogenic structures such as bridges, caves, and culverts. In our

study we cohoused two bats, one inoculated and one uninoculated, in cages of approximately 1

meter^3. This relatively large space did not force the bats to congregate as closely as occurs in

natural roosts and thus "social distancing" may have impeded viral transmission. In future

challenge studies we plan to adjust the bat housing to help take this factor into account.

In addition to the apparent lack of transmission, another observation from this study is

that the virus was not excreted via the digestive system. Given the sizes of TABR colonies, large

amounts of fecal material accumulate that could potentially become a source of infectious virus.

We found no evidence of virus in the digestive tracts or in rectal swabs of any bat, including the

infected bats.

Another finding was that TABR showed no obvious adverse health effects from SARS-

CoV-2 infection. While it is unknown if infected wild bats would have diminished capacity to

forage for food, perform maternal care, or other life functions, our findings indicate that TABR

populations are likely not at risk from the pandemic.

Regardless, an accurate determination of the infectious dose of SARS-CoV-2 in TABR

would be an important next study. We do not know if the amounts of virus excreted are enough

to infect other mammalian species, including humans or if TABR can be infected with SARS-

CoV-2 by exposure to sick humans. Because this species often resides in urban settings, these are

important public health and pandemic ecological concerns.

We were unable to conduct necropsies and collect tissues from actively infected animals.

All bats euthanized for these purposes during the study were not among the infected cohort, and

by the end of the study, all infected bats had cleared the virus. Therefore, the qRT-PCR,

pathology and histochemistry results from bat tissues were inconclusive.

It is generally accepted that SARS-CoV-2 originated in wild horseshoe bats (*Rhinolophus*

sp.) from China, subsequently transmitted to other host species, and ultimately infected humans,

leading to a pandemic (Zhou et al. 2020). The virus also infects at least one other bat species,

Egyptian fruit bats, *Rousettus aegyptiacus* (Schlottau et al. 2020). This has raised concerns that

the virus could infect North American bats, some of whose populations are under severe pressure

from other diseases and from habitat degradation. Hall et al. 2021 previously showed that big

brown bats, a species commonly encountered by humans, is resistant to infection by SARS-CoV-

2. In this study, however, we demonstrated that TABR, a migratory bat that resides in large

colonies, often in urban areas, is susceptible to infection by this virus. Based on the comparative

structure of the SARS-CoV-2 cellular receptor, Damas et al. (2020) predicted that TABR were

unlikely to become infected with SARS-CoV-2. Thus, it is likely that other factors are involved

in mediating susceptibility to this virus. The genetic changes we detected in viral isolates from

the swab of one infected bat indicates that mutations can occur rapidly in a new host and that

genetic analyses of recovered viral isolates may further inform our understanding of the

emergence of novel viral variants. Our findings also indicate that susceptibility of each species to

SARS-CoV-2 is independent, and each species would benefit from being examined individually.

These results have implications for bat rehabilitators, wildlife biologists, cave recreationists, and

the public at large, if they contact Mexican free-tailed bats or enter caves or other environments

where bats are roosting.

Supporting data are available in Hall et al. (2022) https://doi.org/10.5066/P9RDA1H6.

Acknowledgements.

This work was funded by the U.S. Geological Survey's Ecosystems Management Area.

M. Carossino received support from the Center for Lung Biology and Disease (CLBD), Center of

Biomedical Research Excellence, National Institute of General Medical Sciences of the National

Institutes of Health under P20GM130555. We thank Katy Griffin, Jeffrey Messer, Harrison

Lamb, Lauren Dycee-Holtz, Rachel Lambert, Carrie Allison Smith, and Casey Hall, for technical

contributions. We also thank members of the histopathology and immunohistochemistry section

at the Louisiana Animal Disease Diagnostic Laboratory for their assistance. We are particularly

indebted to the NWHC Animal Services staff and volunteers, without whose assistance,

diligence and hard work this study could not have been accomplished. We are also indebted to

the Texas Department of Wildlife for coordinating live bat acquisition. The use of trade, firm, or

product names is for descriptive purposes only and does not imply endorsement by the U.S.

Government. The authors acknowledge they have no personal financial interests or conflicts of

interest with this research article.

341
342
343 **References.**
344
Bosco-Lauth AM, Root JJ, Porter SM, Walker AE, Guilbert L, Hawvermale D, Pepper A,
Maison RM, Hartwig AE, Gordy P, Bielefeldt-Ohmann H, Bowen RA. Peridomestic mammal
susceptibility to severe acute respiratory syndrome coronavirus 2 infection. *Emerg Infect Dis*.
2021 Aug;27(8):2073-2080. doi: 10.3201/eid2708.210180.

Chandler JC, Bevins SN, Ellis JW, Linder TJ, Tell RM, Jenkins-Moore M, Root JJ, Lenoch JB,
Robbe-Austerman S, DeLiberto TJ, Gidlewski T, Kim Torchetti M, Shriner SA. SARS-CoV-2
exposure in wild white-tailed deer (*Odocoileus virginianus*). *Proc Natl Acad Sci* 2021;
118:e2114828118.

Cheng TL, Reichard JD, Coleman JTH, Weller TJ, Thogmartin WE, Reichert BE, Bennett AB,
Broders HG, Campbell J, Etchison K, Feller DJ, Geboy R, Hemberger T, Herzog C, Hicks AC,
Houghton S, Humber J, Kath JA, King RA, Loeb SC, Massé A, Morris KM, Niederriter H,
Nordquist G, Perry RW, Reynolds RJ, Sasse DB, Scafini MR, Stark RC, Stihler CW, Thomas
SC, Turner GG, Webb S, Westrich BJ, Frick WF. The scope and severity of white-nose
syndrome on hibernating bats in North America. *Conserv Biol* 2021; 5, 1586-1597. doi:
10.1111/cobi.13739

Damas J, Hughes GM, Keough KC, Painter CA, Persky NS, Corbo M, Hiller M, Koepfli KP,
Pfenning AR, Zhao H, Genereux DP, Swofford R, Pollard KS, Ryder OA, Nweeia MT,
Lindblad-Toh K, Teeling EC, Karlsson EK, Lewin HA. Broad host range of SARS-CoV-2
predicted by comparative and structural analysis of ACE2 in vertebrates. *Proc Natl Acad Sci*
USA. 2020 Sep 8;117:22311-22322. doi: 10.1073/pnas.2010146117.

Dean, D.J.; Abelseth, M.K.; Atanasiu, P. *The flourescent antibody test. Laboratory Techniques in Rabies*; Meslin, F.X., Kaplan, M.M., Koprowski, H., Eds.; World Health Organization: Geneva, Switzerland, 1996; pp. 83–93.

Decaro N, and Larusso A. Novel human coronavirus (SARS-CoV-2): A lesson from animal coronaviruses. *Veterinary Microb.* 2020; 108693.

Hale VL, Dennis PM, McBride DS, Nolting JM, Madden C, Huey D, Ehrlich M, Grieser J, Winston J, Lombardi D, Gibson S, Saif L, Killian ML, Lantz K, Tell RM, Torchetti M, Robbe-Austerman S, Nelson MI, Faith SA, Bowman AS. SARS-CoV-2 infection in free-ranging white-tailed deer. *Nature.* 2022;602:481-486. doi:10.1038/s41586-021-04353-x.

Hall JS, Knowles S, Nashold SW, Ip H, Leon A, Rocke T, Hofmeister E. Experimental challenge of a North American bat species, big brown bat (*Eptesicus fuscus*), with SARS-CoV-2. *Transboundary Emerg. Dis.* 2021; 68:3443-3452. doi:10.1111/TBED.13949.

Hall, JS., Hofmeister, EK., Ip, HS., Nashold, SW., Leon, AE., Malavé, CM., Falendysz, EA., Rocke, TE., Carossino, M., Balasuriya, U., Knowles, S. 2022. Experimental infection of Mexican free-tailed bats (*Tadarida brasiliensis*) with SARS-CoV-2. U.S. Geological Survey data release, https://doi.org/10.5066/P9RDA1H6.

Harcourt J, Tamin A, Lu X, Kamili S, Sakthivel SK, Murray J, **et al.** Severe acute respiratory syndrome coronavirus-2 from patient with 2019 novel coronavirus disease, United States. *Emerg. Infect. Dis*. 2020; 26:1266-1273.

Hoyt, J.R., Kilpatrick, A.M. & Langwig, K.E. Ecology and impacts of white-nose syndrome on bats. *Nat Rev Microbiol* 2021; 19**,** 196–210. https://doi.org/10.1038/s41579-020-00493-5

McAloose D, Laverack M, Wang L, Killian ML, Caserta LC, Yuan F, Mitchell PK, Queen K, Mauldin MR, Cronk BD, Bartlett SL, Sykes JM, Zec S, Stokol T, Ingerman K, Delaney MA, Fredrickson R, Ivančić M, Jenkins-Moore M, Mozingo K, Franzen K, Bergeson NH, Goodman L, Wang H, Fang Y, Olmstead C, McCann C, Thomas P, Goodrich E, Elvinger F, Smith DC, Tong S, Slavinski S, Calle PP, Terio K, Torchetti MK, Diel DG From people to *Panthera*: natural SARS-CoV-2 infection in tigers and lions at the Bronx Zoo. *MBio*. 2020;11:e02220–20. doi: 10.1128/mBio.02220-20.

Munster VJ, Feldmann F, Williamson BN, van Doremalen N, Pérez-Pérez L, Schulz J, **et al.** Respiratory disease and virus shedding in rhesus macaques inoculated with SARS-CoV-2. *Nature*. 2020; doi: 10.1038/s41586-020-2324-7.

Shi J, Wen Z, Zhong G, Yang H, Wang C, Huang B, et al. Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS-coronavirus 2. *Science* 2020; 368:1016-1020. doi: 10.1126/science.abb7015.

414   Schlottau K, Rissmann M, Graaf A, Schön J, Sehl J, Wylezich C, Höper D, Mettenleiter TC,
415   Balkema-Buschmann A, Harder T, Grund C, Hoffmann D, Breithaupt A, Beer M. SARS-CoV-2
416   in fruit bats, ferrets, pigs, and chickens: an experimental transmission study. *Lancet Microbe*
417   2020; 1:e218-25. doi: 10.1016/S2666-5247(20)30089-6.
418
419   Shriner SA, Ellis JW, Root J, Roug A, Stopak SR, Wiscomb GW, Zierenber, JR, Ip, HS,
420   Torchetti, MK, Deliberto, TJ. SARS-CoV-2 Exposure in Escaped Mink, Utah, USA. *Emerg*
421   *Infect Dis*. 2021; 27:988- 990. doi.org/10.3201/eid2703.204444
422
423   Tyson JR, James P, Stoddart D, Sparks N, Wickenhagen A, Hall G, Choi JH, Lapointe H,
424   Kamelian K, Smith AD, Prystajecky N, Goodfellow I, Wilson SJ, Harrigan R, Snutch TP,
425   Loman NJ, Quick J. Improvements to the ARTIC multiplex PCR method for SARS-CoV-2
426   genome sequencing using nanopore. bioRxiv 2020.09.04.283077;
427   doi: 10.1101/2020.09.04.283077
428
Wacharapluesadee S, Tan CW, Maneeorn P, Duengkae P, Zhu F, Joyjinda Y, Kaewpom T, Chia
WN, Ampoot W, Lim BL, Worachotsueptrakun K, Chen VC, Sirichan N, Ruchisrisarod C,
Rodpan A, Noradechanon K, Phaichana T, Jantarat N, Thongnumchaima B, Tu C, Crameri G,
Stokes MM, Hemachudha T, Wang LF. Evidence for SARS-CoV-2 related coronaviruses
circulating in bats and pangolins in Southeast Asia. *Nat Commun* 2021; 12(1):972. doi:
10.1038/s41467-021-21240-1.
435
436   Zhou P, Yang XL, Wang XG, et al. A pneumonia outbreak associated with a new coronavirus of
437   probable bat origin. *Nature* 2020; 579:270-73. doi: 10.1038/s41586-020-2012-7.
438
439   Zhou H, Ji J, Chen X, Bi Y, Li J, Wang Q, Hu T, Song H, Zhao R, Chen Y, Cui M, Zhang Y,
440   Hughes AC, Holmes EC, Shi W. Identification of novel bat coronaviruses sheds light on the
441   evolutionary origins of SARS-CoV-2 and related viruses. *Cell* 184 (17), 4380-4391 (2021). doi:
442   10.1016/j.cell.2021.06.
443

444 Table 1. Quantitative RT-PCR analyses of oral swabs from Mexican free-tailed bats inoculated with SARS-CoV-2.

Bat ID	Treatment[1]	Age[2]	DPI[3] 0	DPI 2	DPI 4	DPI 6	DPI 8	DPI 10	DPI 12	DPI 14	DPI 16	DPI 18	DPI 20
102	Control	j	No Ct[4]	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
108	Control	j	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
104	Inoc.	a	No Ct	7.53	2.8×10^1	2.32	2.15	1.97	No Ct	No Ct	x[5]	x	x
103	Trans.	j	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	x	x	x
112	Inoc.	j	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	x	x	x	x	x
113	Trans.	a	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
110	Inoc.	a	No Ct	2.96×10^3	3.14×10^3	2.51×10^2	1.07×10^3	3.10×10^2	9.08×10^1	3.44	1.37×10^1	No Ct	No Ct
117	Trans.	a	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
118	Inoc.	a	No Ct	8.63×10^2	4.82×10^3	1.82×10^3	2.18×10^3	5.99×10^1	4.74×10^1	2.08×10^2	4.74×10^2	4.30×10^1	No Ct
114	Trans.	j	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
120	Inoc.	j	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
121	Trans.	j	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
124	Inoc.	j	No Ct	6.80×10^2	2.52×10^2	6.56×10^2	1.00×10^3	2.18×10^1	No Ct	No Ct	No Ct	No Ct	No Ct
125	Trans.	a	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
111	Inoc.	j	No Ct	6.24	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
109	Trans.	j	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
123	Inoc.	a	No Ct	1.94×10^2	1.39×10^2	6.00×10^1	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
122	Trans.	j	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
127	Inoc.	a	No Ct	No Ct	No Ct	No Ct	x	x	x	x	x	x	x
128	Trans.	a	No Ct	No Ct	No Ct	No Ct	x	x	x	x	x	x	x
129	Inoc.	j	No Ct	1.38×10^1	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct

[1]Inoc- Bat oronasally inoculated with $10^5 TCID_{50}$ SARS-CoV-2; Trans.-Uninoculated bat cohoused with inoculated bat to determine virus
transmission
[2]a-adult; j-juvenile
[3]Day post-inoculation
[4]No Ct- no detectable real-time PCR amplification. Quantities of virus based on standard curve of known virus concentrations and expressed as
$TCID_{50}$ equivalent/ml
[5]x-Bat euthanized after previous sampling DPI

Table 2. Quantitative RT-PCR analyses of rectal swabs from Mexican free-tailed bats inoculated with SARS-CoV-2.

Bat ID	Treatment[1]	DPI[2] 0	DPI 2	DPI 4	DPI 6	DPI 8	DPI 10	DPI 12	DPI 14	DPI 16	DPI 18	DPI 20
102	Control	No Ct[3]	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
108	Control	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
104	Inoc.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	x[4]	x	x
103	Trans.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	x	x	x
112	Inoc.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	x	x	x	x	x
113	Trans.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
110	Inoc.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
117	Trans.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
118	Inoc.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
114	Trans.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
120	Inoc.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
121	Trans.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
124	Inoc.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
125	Trans.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
111	Inoc.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
109	Trans.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
123	Inoc.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
122	Trans.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct
127	Inoc.	No Ct	No Ct	No Ct	No Ct	x	x	x	x	x	x	x
128	Trans.	No Ct	No Ct	No Ct	No Ct	x	x	x	x	x	x	x
129	Inoc.	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct	No Ct

[1]Inoc- Bat oronasally inoculated with 10^5TCID$_{50}$ SARS-CoV-2; Trans.-Uninoculated bat cohoused with inoculated bat to determine virus
transmission
[2]Day post-inoculation
[3]No Ct- no detectable real-time PCR amplification. Quantities of virus based on standard curve of known virus concentrations and expressed as
TCID$_{50}$ equivalents/ml.
[4]x-Bat euthanized after previous sampling DPI

Table 3. Competitive ELISA[1] results of Mexican free-tailed bat sera tested after experimental challenge with SARS-CoV-2.

		Results[3]			
Bat ID	**% Inhibition[2]**	**1:10**	**1:20**	**1:40**	**1:80**
102*	-2.656	Neg			
103	14.73	Neg			
104	62.75	Pos	Pos	Pos	Neg
105[4]	-157.1	Neg			
108*	12.66	Neg			
109	13.49	Neg			
110	34.32	Pos	Neg	Neg	Neg
111	1.690	Neg			
112	25.28	Neg			
113	15.21	Neg			
114	34.25	Neg			
117	17.42	Neg			
118	79.30	Pos	Pos	Pos	Neg
119[4]	14.45	Neg			
120	0.862	Neg			
121	14.25	Neg			
122	-24.39	Neg			
123	65.44	Pos	Pos	Neg	Neg
124	57.30	Pos	Pos	Neg	Neg
125	-31.63	Neg			
127	-15.07	Neg			
128	28.04	Neg			
129	7.899	Neg			

[1]SARS-CoV-2 sVNT, GenScript, Piscataway, New Jersey

[2]Inhibition of >30% is positive.

[3] Two-fold serial dilutions of positive sera

[4] Bats euthanized prior to the study

*Uninoculated controls

489   Table 4. Body weights (g) of SARS-CoV-2 inoculated Mexican free-tailed bats.

Bat ID	Treatment	DPI[1] 0	DPI 2	DPI 4	DPI 6	DPI 8	DPI 10	DPI 12	DPI 14	DPI 16	DPI 18	DPI 20
102	Control	10.19	10.11	11.23	11.27	11.8	12.13	11.78	11.7	13.45	13.13	13.24
108	Control	11.97	11.42	12.1	11.86	12.34	12.48	12.3	12.93	12.31	14.38	14.46
104	Inoculated	11.99	10.56	11.85	12.13	12.9	12.76	12.81	13.11	x^2	x	x
103	Transmission	11.81	10.31	11.3	11.28	11.56	12.12	11.88	12.81	x	x	x
112	Inoculated	11.75	11.97	11.31	9.72	9.26	8.4	x	x	x	x	x
113	Transmission	14.37	13.25	12.84	13.81	14.13	15.08	14.83	15.01	14.67	14.43	14.43
110	Inoculated	16.19	14.34	15.39	15.14	15.36	15.6	16.03	15.09	16.04	16.46	16.66
117	Transmission	12.45	11.26	11.84	11.53	11.83	12.87	12.86	12.18	12.66	13.09	13.13
118	Inoculated	13.45	12.34	12.87	12.56	12.9	13.57	13.7	13.72	13.84	13.91	13.75
114	Transmission	10.85	10.2	10.92	10.66	11.31	11.61	11.7	11.39	11.29	11.27	13.68
120	Inoculated	12.1	10.73	11.71	11.5	11.63	12.14	12.27	12.48	12.9	12.47	12.37
121	Transmission	10.33	11.1	12	11.53	11.55	12.05	11.97	12.04	12.54	12.49	13.11
124	Inoculated	13.25	11.95	12.87	12.52	13.33	13.73	14.05	14.49	14.44	14.09	14.9
125	Transmission	13.07	12.2	12.45	12.16	11.57	12.05	12.19	12.41	12.36	12.56	12.84
111	Inoculated	14.02	13.11	13.15	13.73	13.62	13.92	14.25	13.95	13.98	13.64	13.93
109	Transmission	10.86	11.29	11.14	10.56	10.64	10.99	11.39	12.05	12.14	11.97	12.23
123	Inoculated	11.09	10.45	10.83	10.31	10.62	11.19	11.5	11.78	11.48	11.84	12.12
122	Transmission	9.23	9.45	9.78	10.33	10.6	11.37	11.72	11.57	11.65	11.53	11.71
127	Inoculated	9.13	8.86	9.37	8.69	x	x	x	x	x	x	x
128	Transmission	8.39	8.12	8.07	7.95	x	x	x	x	x	x	x
129	Inoculated	10.4	10.96	10	11.51	10.92	10.7	11.48	12.02	10.74	10.76	11.39

492   [1]Day post-inoculation

493   [2]x-Bat euthanized after previous sampling DPI

497 Table 5. qRT-PCR analyses of selected tissues from SARS-CoV-2 infected and uninoculated

498 control Mexican free-tailed bats.

	Bat ID						
Tissue	102^1	108^1	104	110	118	123	124
Brain	NoCt[2]	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt
Nares	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt
Lung cranial	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt
Lung, caudal	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt
Heart	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt
Liver	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt
Kidney	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt
Spleen	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt
Small intestine	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt
Colon	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt	NoCt

500 [1]Uninoculated control

501 [2]NoCt- no detectable RT-PCR reaction

502 Tissues collected from bats euthanized and necropsied on DPI 20

Supplemental Table 1. Rabies diagnostic results from Mexican free-tailed bats by direct fluorescent antibody (DFA) and the presence of alpha- or betacoronaviruses prior to initiation of the SARS-CoV-2 challenge study.

Bat ID	Age	Treatment	DFA[2]	Coronavirus
101	j	None[1]	Pos	Neg
105	j	None[1]	Neg	Neg
107	j	None[1]	Neg	Neg
115	j	None[1]	Neg	Neg
119	j	None[1]	Neg	Neg
102	j	Control	Neg	Neg
108	j	Control	Neg	Neg
104	a	Inoculated	Neg	Neg
103	j	Transmission	Neg	Neg
112	j	Inoculated	Neg	Neg
113	a	Transmission	Neg	Neg
110	a	Inoculated	Neg	Neg
117	a	Transmission	Neg	Neg
118	a	Inoculated	Neg	Neg
114	j	Transmission	Neg	Neg
120	j	Inoculated	Neg	Neg
121	j	Transmission	Neg	Neg
124	j	Inoculated	Neg	Neg
125	a	Transmission	Neg	Neg
111	j	Inoculated	Neg	Neg
109	j	Transmission	Neg	Neg
123	a	Inoculated	Neg	Neg
122	j	Transmission	Neg	Neg
127	a	Inoculated	Neg	Neg
128	a	Transmission	Neg	Neg
129	j	Inoculated	Neg	Neg

[1]None - Bat was euthanized prior to SARS-CoV-2 challenge
[2]Pos – rabies virus presence in brain tissue; Neg – rabies virus abscent in brain tissue or coronavirus abscent in fecal material

Supplemental Table 2

Variant detection in SARS-CoV-2 isolated from Bat 118, DPI 8. Genetic changes in the
SARS-CoV-2 inoculum strain (USA-WA1/2020; Genbank MN985325) and Bat 118 DPI 8
(Genbank OM995890) after one passage in Vero cells compared with the reference strain
(Wuhan HU-1; Genbank MN908947). Sample ID- sample name. Reference position- location of
genetic variant on Wuhan HU-1 genome. Reference sequence- nucleotide sequence in HU-1.
Variant- nucleotide in WA-1 and bat isolate. Count- number of reads covering the variant.
Frequency- proportions of reads with variant. AA change- change in amino acid coding affected
by variant. Syn.- synonymous
552

Sample ID	Reference Position	Reference Sequence	Variant	Count/frequency	AA Change
WA-1	8782	C	T	3628/98.45	Syn.
	28144	T	C	282/96.58	Syn.
Bat 118 DPI 8	8782	C	T	3514/98.46	Syn.
	28144	T	C	311/97.49	Syn.
	28253	C	T	1509/94.85	Syn.
	28603	C	T	3928/90.05	Syn.

561 Supplemental Table 3. SARS-CoV-2 antigen distribution in selected tissues from SARS-CoV-2 inoculated and uninoculated control
562 Mexican free-tailed bats. Bats euthanized, necropsied and tissue samples collected on day post-inoculation (DPI) 7 (bats 127, 128));
563 DPI 10 (bat 112); DPI 14 (bats 103, 104); and DPI 20 for the remaining bats.

Bat ID

Tissue	102[1]	108[1]	103	104[2]	109	110[2]	111	112	117	118[2]	123[2]	124[2]	127	128
Brain	None	None	None	None	None	None	None	None	None	None	None	None	None	None
Nares	None	None	None	None	None	None	None	None	None	None	None	None	None	None
Lung	None	None	None	None	None	None	None	None	None	None	None	None	None	None
Heart	None	None	None	None	None	None	None	None	None	None	None	None	None	None
Liver	None	None	None	None	None	None	None	None	None	None	None	None	None	None
Kidney	None	None	None	None	None	None	None	None	None	None	None	None	None	None
Spleen	None	None	None	None	None	None	None	None	None	None	None	None	None	None
Stomach	None	None	None	None	None	None	None	None	None	None	None	None	None	None
Pancreas	None	None	None	None	None	None	None	None	None	None	None	None	None	None
Small intestine	None	None	None	None	None	None	None	None	None	None	None	None	None	None
Colon	None	None	None	None	None	None	None	None	None	None	None	None	None	None

565 [1]Uninoculated control

566 [2]SARS-CoV-2 infected bats

567 None: no detectable SARS-CoV-2 nucleocapsid antigen