Spindle-dependent memory consolidation in healthy adults: A meta-analysis

iD **Deniz Kumral**[a,b,c,✉], **Alina Matzerath**[a], **Rainer Leonhart**[a], **and** iD **Monika Schönauer**[a]

[a]Institute of Psychology, Neuropsychology, University of Freiburg, Freiburg im Breisgau, Germany
[b]Institute of Psychology, Clinical Psychology and Psychotherapy Unit, University of Freiburg, Freiburg im Breisgau, Germany
[c]Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany
[d]Institute of Psychology, Social Psychology and Methodology, University of Freiburg, Freiburg im Breisgau, Germany
[e]Bernstein Center Freiburg, Freiburg im Breisgau, Germany

Accumulating evidence suggests a central role for sleep spindles in the consolidation of new memories. However, no meta-analysis of the association between sleep spindles and memory performance has been conducted so far. Here, we report meta-analytical evidence for spindle-memory associations and investigate how multiple factors, including memory type, spindle type, spindle characteristics, and EEG topography affect this relationship. The literature search yielded 53 studies reporting 1427 effect sizes, resulting in a small to moderate effect for the average association. We further found that spindle-memory associations were significantly stronger for procedural memory than for declarative memory. Neither spindle types nor EEG scalp topography had an impact on the strength of the spindle-memory relation, but we observed a distinct functional role of global and fast sleep spindles, especially for procedural memory. We also found a moderation effect of spindle characteristics, with power showing the largest effect sizes. Collectively, our findings suggest that sleep spindles are involved in learning, thereby representing a general physiological mechanism for memory consolidation.

sleep spindles, sleep, memory consolidation, procedural memory, declarative memory, EEG, meta-analysis

✉ *deniz.kumral@psychologie.uni-freiburg.de*

Highlights

- Spindle measures showed a small to medium-sized association with memory performance.

- This relationship was stronger for procedural memory than declarative memory.

- No moderation effects of spindle type and EEG scalp topography have been observed.

- Spindle power emerged as the strongest predictors.

- Naps showed similar spindle-related consolidation mechanisms to whole-night sleep.

1.Introduction

A wealth of research, much of it conducted over the past 30 years, has shown that we retain memories better over a period of sleep relative to wakefulness. With the positive effect of sleep on memory being well-established, current research efforts have been directed at uncovering the mechanisms that mediate this relationship. Accumulating evidence suggests that sleep does not merely passively protect memories from interference and decay, but actively fosters their consolidation. Thus, longer sleep times lead to greater memory benefits (Diekelmann et al., 2012; Schönauer et al., 2014), the occurrence of sleep-specific brain oscillatory activity has been linked with memory consolidation (Fogel and Smith, 2006; Gais et al., 2002; Girardeau et al., 2009; Marshall et al., 2006; Perrault et al., 2019; Schmidt et al., 2006), and it has been discovered that learning-related brain activity is reactivated during non-rapid eye movement (NREM) sleep (Peigneux et al., 2003; Schönauer et al., 2017; Schreiner et al., 2021; Wilson and McNaughton, 1994). Sleep-dependent memory reactivation, in particular, is a promising candidate mechanism to underlie behavioral benefits of sleep on memory performance (Antony et al., 2012; Rasch et al., 2007; Rudoy et al., 2009). Memory reactivation has been linked to oscillations characteristic of NREM sleep, such as slow oscillations (Cousins et al., 2014; Ngo and Staresina, 2022; Schönauer et al., 2017; Schreiner et al., 2021), sleep spindles (Antony et al., 2018; Bergmann et al., 2012; Cairney et al., 2018; Fogel et al., 2017; Jegou et al., 2019; Laventure et al., 2018; Schönauer et al., 2017; Wang et al., 2019), and sharp-wave ripples (SWR) (Fernández-Ruiz et al., 2019; Rothschild et al., 2017; Zhang et al., 2018). These oscillations are coordinated within a network of brain regions including the thalamus, neocortex, and hippocampus. Research in both animals and humans has shown that the precise orchestration of these brain rhythms contributes to memory consolidation during sleep (Maingret et al., 2016; Mölle et al., 2009; Staresina et al., 2015). Sleep spindles, in particular, have emerged as a promising candidate marker for memory reactivation in the human brain and have been linked to behavioral benefits of sleep for memory. Spindle activity during sleep increases after an intense learning experience and is related to how well the memory is retained (Gais et al., 2002; Morin et al., 2008; Schabus et al., 2008), and artificially boosting sleep spindle occurrence by pharmacological interventions benefits memory consolidation (Mednick et al., 2013). Moreover, the presentation of learning-related sounds during sleep elicits stronger activity in the spindle frequency band than the presentation of control sounds (Cairney et al.,

2018), and brain activity during sleep spindles is informative about the kind of learning material studied before sleep (Cairney et al., 2018; Schönauer et al., 2017). Because recent findings have shown that effect sizes for associations between sleep and memory performance have been overestimated (Ackermann et al., 2015; Berres and Erdfelder, 2021; Cordi and Rasch, 2021; Pöhlchen et al., 2021; Reverberi et al., 2020), we should quantitively evaluate the influence of specific sleep parameters on memory in a similar way. In this meta-analysis, we therefore investigated whether spindle activity shows a consistent link with memory performance, and overnight memory consolidation in particular.

1.1. The role of sleep spindles in memory consolidation and reprocessing

Sleep spindles are a physiological hallmark of NREM sleep, where they appear as brief oscillatory events with a burst-like sequence of 10–16 Hz sinusoidal cycles, lasting for at least 0.5 sec, but no longer than 2-3 sec (Fernandez and Lüthi, 2020; Lüthi, 2014). They are generated through interactions between inhibitory cells in the thalamic reticular nucleus and bursting thalamocortical relay neurons (Contreras and Steriade, 1996; Steriade, 2006). Cortical inputs to the thalamus may regulate spindle initiation and termination (Bonjean et al., 2011).Importantly, spindles are thought to play a major role in mediating neural plasticity (Peyrache and Seibt, 2020; Steriade and Timofeev, 2003) and memory replay (Siapas and Wilson, 1998).

A direct link between spindle activity and plasticity has first been proposed by Timofeev and colleagues (Timofeev et al., 2002), who showed that spindle activity is associated with long-term changes in responsiveness of cortical neurons. Similarly, input firing patterns during spindles have been found to induce long-term potentiation (LTP) effectively in neocortical pyramidal cells (Rosanova and Ulrich, 2005), probably by increasing intracellular calcium concentrations. Recent calcium imaging studies have also provided evidence that spindles may trigger cell- and layer-specific activation (Niethard et al., 2017), which could support plasticity underlying memory consolidation. It has been shown that the calcium activity (Ca2+) in neocortical dendrites is increased and synchronized during oscillations in the spindle range (Seibt et al., 2017), which may be enhanced through coupling with slow oscillation (SO) up-states (Niethard et al., 2018), thereby strengthening memory consolidation.

Sleep spindles are further thought to play a pivotal role in memory reactivation. Memory reactivation has mainly been studied in the rodent hippocampus, where it predominantly occurs during SWRs, which are transient high-frequency oscillations (140–200 Hz) during rest and NREM sleep. It has been reported that hippocampal replay during SWR can be accompanied by the cortical replay of the same event (Ji and Wilson, 2007). Interestingly, optogenetic stimulation of SWRs enhances replay and results in memory improvement (Fernández-Ruiz et al., 2019). Intracranial EEG data recorded in humans has shown that SWRs in the hippocampus are nested into the troughs of the spindle oscillations (Ngo et al., 2020; Staresina et al., 2015), suggesting that spindles might facilitate the interaction, and thus communication between neural ensembles of the hippocampus and the neocortex. Notably, this memory reactivation may strengthen long-term memory representations in the neocortex such that they can support memory retrieval independent of hippocampal engagement, a process that has been proposed to serve active systems memory consolidation in sleep (Diekelmann and Born, 2010; Klinzing et al., 2019).

Converging evidence further suggests that the content of memory reactivation can be decoded from brain activity during sleep spindles (Cairney et al., 2018; Wang et al., 2019), underlining their role in reactivation-related consolidation during sleep. Indeed, targeted memory reactivation (TMR), where sensory cues related to previous learning material are used to trigger reactivation of these learning contents during sleep, is only effective if the reactivation cues are followed by a sleep spindle, demonstrating their functional role in sleep-dependent memory consolidation (Antony et al., 2018; Schönauer, 2018).

1.2. Sleep spindles and different types of memory

It is well-established that sleep fosters learning and retention of both declarative and procedural memories (Diekelmann and Born, 2010; Klinzing et al., 2019), and sleep spindles have been linked to these behavioral benefits. For instance, several studies showed that sleep spindles strengthen memory for a broad range of declarative tasks including wordlist and vocabulary learning (Blaskovich et al., 2017; Gais et al., 2002; Schabus et al., 2004; Schmidt et al., 2006; Studte et al., 2017, 2015; Tamminen et al., 2010), but also visual memory for pictures (Cox et al., 2012; Ward et al., 2014), and visuospatial tasks (Clemens et al., 2006; Wilhelm et al., 2011). Likewise, spindles have been shown to facilitate procedural memory, such as finger tapping, motor sequential learning, and motor adaptation (Barakat et al., 2011a; Boutin et al., 2018; Fogel and Smith, 2006; Nishida and Walker, 2007; Simor et al., 2019; Thürer et al., 2018) and mirror tracing (Holz et al., 2012; Tamaki et al., 2008; van Schalkwijk et al., 2019). However, recent findings and reviews have started to question the robustness, replicability, and generalizability of some of these findings (Ackermann et al., 2015; Cordi and Rasch, 2021). In support of this, Ackermann and colleagues (2015) observed no significant association between sleep spindles and overnight memory retention in both declarative and procedural memory measures in over 900 healthy young participants.

If spindles are a mechanism that fosters brain plasticity and if they are directly related to sleep-dependent memory reactivation, then spindle activity during sleep should be positively related to memory consolidation. In this meta-analysis, we quantified evidence for the effect of sleep spindles on memory consolidation and assessed how strongly the extant literature is subject to publication bias. We further investigated whether the consolidation of both declarative and procedu-

ral memory is equally sleep spindle-dependent. Finally, we assessed how strongly the extant literature is subject to *publication bias* and discuss to what extent this could affect the interpretation of the obtained results.

1.3. Spindle characteristics and their relation to memory performance

1.3.1. Spindle type and topography
Studies in the 1990s hinted at the existence of two distinct sleep spindles types, observed at 10-13 Hz (slow spindles) or 13-16 Hz (fast spindles), and peaking at frontal and centroparietal cortical sites, respectively (McCormick et al., 1997; Werth et al., 1997). The idea that sleep spindles in humans are a diverse rather than prototypical phenomenon was further supported over the last few decades in studies using functional magnetic resonance imaging (fMRI; (Schabus et al., 2007), intracranial EEG (Andrillon et al., 2011; Gonzalez et al., 2022; Piantoni et al., 2017) or M/EEG source estimation techniques (Dehghani et al., 2010; Urakami, 2008). Further, a functional differentiation between slow and fast spindles has been suggested: For instance, it has been observed that both types preferentially occur at different times of the SO cycle (Mölle et al., 2011; Staresina et al., 2015), respond differentially to pharmacological interventions (Ayoub et al., 2013), and have different phenotypic and genetic profiles (Purcell et al., 2017). Spindle topography might be directly related to the spindle type that is studied (fast or slow). Additionally, it has recently been shown that the topography of spindle amplitude can also depend on the prior learning task (Petzka et al., 2022) since spindle topographies during sleep recapitulate spatial patterns of brain activation during memory encoding, a process that benefits memory consolidation. It is currently unclear whether slow and fast spindles serve distinct functional roles in memory, and how spindle topographies are related to memory consolidation. To investigate this question further, in the present meta-analysis, we assessed whether spindle type and spindle topography modulate the strength of spindle-memory association for both declarative and procedural memory.

1.3.2. Spindle measures
A final important factor to consider when investigating the effect of sleep spindles on memory is the spindle measures used. Different aspects, such as count, power, or duration (see Figure 1) may reflect distinct underlying physiological processes in the brain and may be differentially associated with behavioral outcomes. Although many studies report the relationship between sleep spindles and memory, it is currently unknown if the magnitude of these associations depends on different spindle measures. The present meta-analysis will therefore assess such differences in the strength of the relationship between sleep spindles and memory consolidation across various spindle measures. (**Fig. 1**).

Fig. 1. Schematic representation of how spindles appear in a non-rapid eye movement (NREM) sleep in electroencephalogram (EEG) and overview of the most frequently studied measures. *Count*: number of spindles; *Duration*: mean time (in sec) of spindle events; *Density*: number of spindles per unit of time (per minute or per 30 sec); *Power*: mean amount of activity in spindle frequency; *Amplitude*: the absolute value (in µV) of the maximum negative peak, measured as the voltage difference between the largest trough and the largest peak; *Frequency*: Number of occurrences of a repeating event per unit of time (cycle number/duration) or maximal spectral peak amplitude (in Hz); *Spindle Activity (SpA)*: mean duration x amplitude.

1.4. Objectives of the meta-analysis
Although there are meta-analyses investigating the relation of sleep spindles with trait-like aspects like intelligence, cognitive performance, and general cognitive ability in healthy adults (e.g. Ujma, 2018), in the present study, we aim to contribute a more comprehensive overview of the relationship between sleep spindles and memory. We review the relevant methodological aspects that vary across studies and report meta-analytical evidence for the relationship between sleep spindles measured with EEG and memory. More precisely, we aimed to address the following main research questions systematically:

1. What is the overall strength of the association between sleep spindles and memory performance?

2. Does the sleep spindle-memory association depend on the type of memory that is consolidated (e.g., procedural vs. declarative)?

3. Do certain spindle types and spindle scalp locations result in stronger spindle-memory associations?

4. Does the strength of the sleep spindle-memory relationship depend on the spindle measures?

To investigate the general effect of sleep spindles on memory, we included studies reporting different kinds of memory measures. Both memory retention and post-sleep memory measures were categorized as reflecting the process of memory consolidation, and thus constitute a *state* measure of memory influenced by the specific learning experience.

Trait-like measures of memory, such as recall performance on a control day or pre-sleep memory performance were categorized as learning ability, reflecting a general cognitive trait that is largely independent of the specific learning experience (Clemens et al., 2006, 2005; Lustenberger et al., 2012; Schabus et al., 2008; Sopp et al., 2018). In the main part of the meta-analysis, we focused on the association between sleep spindles and memory consolidation as a *state* measure and assessed how various spindle characteristics influence this relationship, but we also evaluated the overall strength of the spindle-memory association for the *trait* measure. We finally explored the strength of the spindle-memory association depending on the sleep protocol and sleep duration, as the included studies investigated sleep both during the whole night and nap sleep protocols.

2.Methods

2.1. Search strategy
The present meta-analysis was implemented according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines for designing, implementing, and reporting systematic reviews and meta-analyses (Page et al., 2021). The systematic literature search was conducted on electronic databases including PubMed and PsychINFO in May 2021 using the following terms: sleep spindles, memory, EEG as well as database-specific search terms (including Medical Subject Heading (MeSH) terms). Additionally, further sources including reference lists of included studies and previous systematic reviews on the study topic as well as two key journals (Neuroscience & Biobehavioral Reviews and Sleep Medicine Reviews) were searched to increase the likelihood of retrieving relevant empirical studies.

2.2. Study selection
Studies were included if they **1)** reported results from a cohort, case-control, cross-sectional or experimental study and **2)** assessed the relationship between memory (procedural and/or declarative) and sleep spindles measured with EEG in healthy individuals between 18 and 65 years of age. Additionally, studies met the inclusion criteria only if they were published in peer-reviewed journals in English. On top of the inclusion criteria, we excluded studies with **1)** non-human samples (e.g., rodents, primates), **2)** samples from clinical populations (e.g., neurological or psychiatric disorders), **3)** the use of a reactivation protocol including targeted memory reactivation (TMR), **4)** applying pharmacological (e.g., SSRIs) or physiological interventions (e.g., transcranial direct current stimulation (tDCS), transcranial magnetic stimulation (TMS), auditory simulations), **5)** the absence of a long-term memory measure (e.g., reporting only working memory or measures of executive function), or **6)** absence of sleep, EEG, or spindle measures. Other types of studies including systematic reviews, meta-analyses, letters to editors, opinions, case reports, case series, and clinical trials were also excluded.

2.3. Study screening and data extraction
Identified titles and abstracts were screened for relevance, separately by the two reviewers (DK, AM). Full texts of identified articles were retrieved and read in full to assess eligibility for inclusion. A standardized data coding form was developed to extract the following information from each study: **(a)** authors, publication year, and DOI; **(b)** characteristics of the study sample (age, gender, number of participants); **(c)** spindle characteristics (type and measure); **(d)** sleep information (duration, protocol, stage); **(e)** EEG information (scalp location, channel); **(f)** memory type and specific outcome measure and **(g)** statistics (e.g., the correlation between memory and sleep spindles).

In cases of relevant studies not reporting extractable statistics with regard to the association between sleep spindles and memory performance, authors were contacted via email to request the minimum required data for the meta-analysis. E-mails were sent at least twice at approximately fourteen-day intervals.

Importantly, we aim to make this meta-analysis transparent by publicly sharing data files, associated codebooks, and analysis at Open Science Framework https://osf.io/wu6d7.

2.4. Statistical Analysis
The meta-analysis was conducted using the *metafor* package (Viechtbauer, 2010) with the function of *rma.mv* in the *R* statistical software (version 3.5; R Core Team, 2017), allowing to fit multilevel meta-analytic models.

Before computing the meta-analyses, correlation coefficients were transformed to the normally distributed Fisher's z (*escalc*). The Fisher's z score and its variance were used in the meta-analyses, and afterwards back-transformed to Pearson's correlation coefficients to facilitate interpretation. If Pearson correlations were not reported, Fisher's z was estimated using statistics such as partial r and Spearman's rank correlation.

Since the studies included in this meta-analysis have dependent effect sizes, with several effect sizes reported from the same sample, statistical analysis based on traditional approaches such as averaging all effect sizes, selecting one effect size per study, or conducting separate meta-analyses in independent subsets may lead to inaccurate or false-positive results (Borenstein et al., 2009; Cheung, 2019). The extant literature has proposed several procedures to account for these dependencies (Borenstein et al., 2009). In the current study, we opted for a multilevel random-effects model using a restricted maximum likelihood method (REML) to estimate overall effects and account for the heterogeneity across included studies (Cheung, 2019; Fernández-Castilla et al., 2020; Konstantopoulos, 2011). The three-level random-effects model quantifies the sampling variance (level 1), the variance between effect sizes within studies (level 2), and the variance of effect sizes between studies (level 3). To examine variance and heterogeneity among effect sizes of included studies, we computed Q statistics, the significance of Q, and

I^2, respectively. I^2 values of ~25%, 50%, and 75% were interpreted as low, moderate, and high, respectively (Higgins and Thompson, 2002; Quintana, 2015; Thompson and Higgins, 2002). Lastly, the presence of publication bias was examined by investigating the association between sleep spindles and memory consolidation with the modified version of the Egger test using standard error as a moderator. To estimate how publication bias may influence the estimated effect sizes between sleep spindles and memory, we used information on whether an effect size was originally reported in the literature or whether it was obtained only later, upon asking the authors (effect size reporting status: reported vs. unreported) as a categorical moderator to assess whether reported correlation values had significantly larger effect sizes than unreported ones (for a similar approach, see (Hu et al., 2020)).

2.5. Moderators

A first moderator analysis was conducted to determine the overall strength of the association between sleep spindles and memory performance depending on state consolidation and trait learning ability measures. With regard to memory consolidation measures, we then assessed possible effect size differences between memory retention and post-sleep memory measures. We further investigated differences in the magnitude of sleep-memory associations for declarative compared with procedural memory. As sleep may contribute differentially to the memory consolidation depending on the procedural memory task (Schendan et al., 2003; Schönauer et al., 2015; Walker et al., 2003), we also explored effect size differences between hippocampal (e.g., finger tapping) vs. non-hippocampal procedural tasks (e.g., mirror-tracing or motor-adaptation). We next assessed the effect of spindle type and EEG scalp topography on memory consolidation. Spindle type was defined as slow, fast, and global spindle-frequency range. For investigating scalp topography, we grouped EEG channels into three coarser brain regions (frontal, central, and parietal), as the EEG montage system, electrode positioning, and the number of electrodes varied between studies. We additionally explored the effect of EEG scalp topography depending on spindle type, as it was previously demonstrated that slow spindles prevail over anterior and fast spindles over posterior brain areas (Cox et al., 2017). We finally examined differences in the spindle-memory association depending on spindle measures, which were categorized as power, duration, count, amplitude, density, oscillatory frequency, and spindle activity (SpA) (**Fig. 1**). We defined these variables based on the definition or calculation reported in the original studies. In addition to these planned analyses, we also explored the possible effects of sleep protocol (whole night vs. napping), sleep duration (in min), and a possible effect of age (in years), as these variables may impact sleep-behavior correlates (Gui et al., 2017; Hokett et al., 2021; Lo et al., 2016; Qin et al., 2022; Schmid et al., 2020). In all analyses, moderator effects for categorical variables were only considered if at least six to seven effects sizes were available per category (Tipton et al., 2019). In all analyses, moderator effects for

categorical variables were only considered if at least six to seven effects sizes were available per category (Tipton et al., 2019).

2.6. Control Analyses

Two control analyses were conducted to compare meta-analytic results obtained from analyses with and without outliers, as the presence of outliers may affect the validity and robustness of the conclusions from a meta-analysis (Viechtbauer and Cheung, 2010). We determined statistical outliers using standardized residuals and Cook's distances (D_i) (Cook, 1977). D_i indicates the relative influence of each effect size on the summary estimate. A standard rule of thumb for potential outliers is a Di value greater than three times the mean D_i (Viechtbauer and Cheung, 2010). Similarly, as an outlier criterion, we excluded effect sizes that have standardized residual greater than 3 in absolute magnitude (Hedges and Olkin, 2014).

As a second control analysis, we excluded studies with the largest sample size (Ackermann et al., 2015) and the smallest sample size (Fogel and Smith, 2006) and also on studies that reported a substantially larger number of estimates than the average (Blaskovich et al., 2017; Thürer et al., 2018; k > 130) and repeated the meta-analysis for the association between sleep spindles and memory consolidation.

3.Results

3.1. Characteristics of the included studies

After reviewing the 421 titles and abstracts, 140 studies were retained for full-text review and screened for eligibility. Eighty-seven of these studies were excluded for the following reasons: Not providing a spindle characteristics or type diagnosis (n = 12), no clear measure of memory (n = 35), no sleep (n = 2) or EEG acquisition (n = 4), lacking necessary data to compute effect sizes (n = 25), investigating dream-recall (n = 2), working memory/intelligence (n = 6) or other (n = 1) as shown in **Fig. 2**. A total of 53 studies were included in the final analysis published between 2002 (Gais et al., 2002) and 2021 (Lutz et al., 2021) (**Fig. 3**). The included studies were conducted in the following countries: Germany (n = 14), United States of America (n = 8), Canada (n = 7), Austria (n = 6), Switzerland (n = 4), United Kingdom (n = 4), Hungary (n = 4), The Netherlands (n = 2), Japan (n = 2), Iran (n = 1), and Cuba (n = 1). There were 33 studies investigating declarative memory, 16 studies assessing procedural memory, and 4 studies reporting results from both (Ackermann et al., 2015; Genzel et al., 2014; Griessenberger et al., 2013; Holz et al., 2012). Further, there were 6 putatively non-hippocampal procedural tasks like mirror-tracing (e.g., Tamaki et al., 2008; van Schalkwijk et al., 2020) and 14 procedural tasks that have been shown to rely on hippocampal contributions, like finger sequence tapping (e.g., Nishida et al., 2016; Schönauer et al., 2014).

The number of estimates for each measure of interest across 53 studies in 1427 effect sizes can be found in Supplementary

Table 1.

3.2. The association between sleep spindles and memory performance

We first tested the overall strength of association between sleep spindles and memory performance including both state consolidation and trait learning ability measures in 1427 unique associations (k) across 53 independent studies. Using a three-level meta-analytic model, we observed that the pooled correlation between sleep spindles and memory performance was r = 0.236; CI: [0.161–0.308]; $p < 0.001$, reflecting a small- to moderate effect size for the spindle-memory association. Moderation analyses indicated that there were no differences in effect sizes of the spindle-memory association between state and trait measures of memory ($Q_m(1) = 0.560$, $p = 0.454$), but we observed a numerically higher effect size for state consolidation measures (r = 0.238, k = 1204) than for trait-like measures (r = 0.218, k = 223).

3.3. Average association between sleep spindles and memory consolidation

Since our main objective was to assess spindle-dependent memory consolidation and how various spindle characteristics, including spindle type, spindle topography, and different spindle measures, influence the spindle-memory relationship, we focused on state consolidation measures from this point onwards.

We first tested the strength of association between sleep spindles and memory consolidation and observed a moderate correlation (r = 0.243; CI: [0.166–0.317]; $p < .001$; Supplementary Fig 1). There was substantial heterogeneity among effect sizes ($Q(1203) = 2828.262$, $p < .001$; $I^2(2) = 21.43$ and $I^2(3) = 52.75$ especially moderate to high between-study heterogeneity. We assessed publication bias using the modified version of the Egger test using standard error as moderator and detected significant publication bias ($\beta = 1.64$, $SE = 0.53$, $p = 0.0020$, Supplementary Fig 2, Supplementary Fig 3), indicating that significant results are more likely to be published than small or non-significant findings. We then tested whether the effect sizes were larger if reported in published manuscripts than if they were not reported. When effect size reporting status (yes vs. no) was examined in the moderator analysis, we observed no significant differences between reported and unreported effect sizes ($Q_m(1) = 0.445$, $p = 0.505$; reported; r = 0.249, k = 793, unreported; r = 0.222, k = 441) across all studies. Interestingly, when we conducted this moderation analyses separately for studies assessing declarative or procedural memory, we observed significant differences ($Q_m(1) = 6.094$, $p = 0.014$ in declarative; ($Q_m(1) = 13.645$, $p < 0.01$ in procedural memory): While in studies with declarative memory we found the effect size of reported studies (r = 0.254, k = 517) were higher than unreported ones (r = 0.131, k = 295), we found opposite pattern such that the effect size of reported results (r = 0.169, k = 276) were higher than unreported ones (r = 0.421, k = 116) in studies with as-

sessing procedural memory.

Subsequent moderation analyses further revealed no differences between memory retention and post-sleep measures ($Q_m(1) = 0.341$, $p = 0.560$; memory retention; r = 0.249, k = 592, post-sleep; r = 0.231, k = 612), suggesting that both measures show a similar effect size for the spindle-memory association.

3.4. Spindle-memory association depending on memory type

To address the second research question, memory type (declarative vs. procedural) was examined as a moderator for the mean effect size between sleep spindles and memory consolidation. We found that memory type is a significant moderator of the spindle-memory association ($Q_m(1) = 7.589$, $p = 0.006$), with a stronger spindle-memory association for procedural memory (r = 0.319, $p < .001$, k = 392) than for declarative memory (r = 0.210, $p < .001$, k = 812), as shown in (**Fig. 4A**. Although studies employing hippocampal procedural memory tasks (r = 0.263, k = 195) showed higher effect sizes than those using non-hippocampal tasks (r = 0.244, k = 197), this was not a significant moderator ($Q_m(1) = 0.008$, $p = 0.928$).

3.5. Spindle-memory association depending on spindle characteristics

3.5.1. Spindle types

We further assessed whether spindle type, i.e., slow or fast spindles, acted as a moderator for spindle-memory associations across all studies. There was a no moderation effect of spindle type ($Q_m(2) = 1.831$, $p = 0.400$; fast; r = 0.264, k = 552, slow; r = 0.238, k = 327, global; r = 0.233, k = 325, **Fig. 4B**). The subsequent moderation analysis separately performed in studies on declarative memory was also non-significant ($Q_m(2) = 0.675$, $p = 0.714$, k = 812), suggesting no notable difference between the spindle types in declarative memory (fast; r = 0.244, k = 308, slow; r = 0.250, k = 226, global; r = 0.216, k = 278). Nevertheless, we found a significant effect of spindle type ($Q_m(2) = 8.610$, $p = 0.013$, k = 392) for procedural memory measures, where global and fast, but not slow frequency spindles contributed to the spindle-memory relationship (fast; r = 0.247, $p = 0.01$, k = 244, slow; r = 0.141, $p = 0.186$, k = 101, global; r = 0.328, $p = 0.005$, k = 47). Follow up post-hoc comparisons with Bonferroni correction indicated a correlation difference between fast and slow (fast-slow; rdiff = 0.110, ($p_{adj} = 0.007$), but not between fast and global or slow and global spindle frequencies in procedural memory ($p_{adj} > 0.05$).

3.5.2. EEG scalp topography

Next, moderator analyses were conducted to determine if scalp topography (frontal (k = 350), central (k = 438), or parietal (k = 222) modulated the strength of the spindle-memory relation. There was no significant moderation effect of scalp

Fig. 2. PRISMA flow diagram of the literature search, screening, and inclusion processes; n = number of studies; k = number of estimates.

topography across all studies ($Q_m(2) = 0.885$, $p = 0.659$, k = 1010), meaning that there was no influence of recording location of the spindles on the strength of the spindle-memory associations (**Fig. 4C**). Moreover, we calculated separate meta-analyses across studies reporting slow and fast spindles, respectively and observed no moderator effect of scalp topography (slow; ($Q_m(2) = 0.186$, $p = 0.911$, k = 297, fast; ($Q_m(2) = 0.137$, $p = 0.934$, k = 488), suggesting that the effect sizes of EEG clusters do not differ in studies reporting either slow or fast spindle frequencies (r = 0.125 – 0.175).

3.5.3. Spindle measures
We further investigated effect size differences in the sleep-memory association depending on different spindle measures (amplitude; k = 142, count; k = 235, density; k = 321, duration; k =145, frequency; k = 27, power; k = 204, and SpA; k = 130). We found a significant moderation effect of spindle measure ($Q_m(6) = 69.692$, $p < 0.001$, k = 1204) with the largest effect sizes for frequency and power (**Fig. 4D**). All of the spindle measures showed significant memory associations (all $p < 0.05$). The results of the post-hoc tests (number of comparisons = 21) indicated larger effects for spindle frequency compared to all spindle measures except for spindle power ($p = 0.06$) and amplitude compared to duration ($p_{adj} < 0.05$, Supplementary Table 2), while we did not observe any other significant effect size differences between the rest of the spindle measures after Bonferroni correction (all $p_{adj} > 0.05$, Supplementary Table 2).

Because only two studies reported the relation of memory consolidation (procedural memory) with oscillatory peak frequency (Boutin et al., 2018; Thürer et al., 2018), defined as maximal spectral peak amplitude (in Hz), we repeated the analysis after excluding the frequency measure from these studies (n = 27). After this exclusion, we confirmed the observed findings of the moderator effect of spindle measures

Fig. 3. Overall number of estimates and sample size per study extracted for the meta-analyses (k=1427). To increase the interpretability of the graph, we visualized one study (Ackermann et al., 2015) with the largest sample size (N = 929) as dark brown. While the average sample size (N) across 53 studies was 35.66, after excluding this study the average sample size became 18.59 (ranging from N = 6 to N = 44), shown in vertical dot line. The total sample size across 53 studies was 1896.

Fig. 4. The average effect size (r) of sleep-memory associations by A. memory types, **B**. spindle types, **C**. EEG topography, and **D**. spindle measures. Error bars indicate 95% confidence intervals. While we observed significant effect size differences for memory types and spindle measures, there were no significant moderation of spindle type and scalp topography on spindle-memory consolidation.

$(Q_m(5) = 14.968, p = 0.01, k = 1177)$ with the largest effect size for the spindle power and the smallest effect size for SpA. Further post-hoc comparisons (number of comparisons = 15) similarly indicated that there were significant effect size differences between amplitude and duration, but not between other measures $(p_{adj} < 0.05$, Supplementary Table 3).

3.6 Exploratory Analyses

3.6.1. Sleep Protocol and Duration
Moderator analyses were conducted to determine if sleep protocol (nap vs. whole-night) or sleep duration (as a continuous variable) modulated the strength of the sleep-memory association. There was no significant difference $(Q_m(1) = 2.149, p = 0.143, k = 1169)$ in sleep-memory associations for nap sleep (r = 0.167, k = 543) and whole night sleep (r = 0.280, k = 626). Confirming this result, we also observed non-significant findings in studies reporting either declarative $(Q_m(1) = 0.471, p = 0.493, k = 812)$ or procedural memory measures $(Q_m(1) = 2.089, p = 0.148, k = 357)$. Similarly, we found no significant moderation effect of sleep duration (M = 242.41, SD = 169 min) on spindle-memory association neither across all studies $(Q_m(1) = 1.357, p = 0.244, k = 1204, r$

= 0.181), nor separately in studies reporting either declarative $(Q_m(1) = 1.032, p = 0.310, k = 812, r = 0.174)$ or procedural memory $(Q_m(1) = 0.251, p = 0.617, k = 392, r = 0.210)$.

3.6.2. Age
Moderator analyses were conducted to determine if age (as a continuous variable) modulated the strength of the sleep-memory association. We observed that continuous age (in years) was not a significant moderator for the association between sleep spindles and memory $(Q_m(1) = 0.235, p = 0.628, r = 0.355)$. We observed similar results when separately analyzing studies reporting declarative $(Q_m(1) = 0.204, p = 0.652, r = 0.138)$ and procedural memory $(Q_m(1) = 1.100, p = 0.294, r = 0.744)$.

3.7. Control Analyses
After computing standardized residuals and Cook's distance (Di), 50 effect sizes were excluded. We observed similar findings for the summary estimate of the spindle-dependent memory consolidation and no major changes occurred to the results of moderator analyses.

After excluding studies with the largest (Ackermann et al., 2015) and smallest (Fogel and Smith, 2006) sample size and a higher number of estimates (Blaskovich et al., 2017;

Thürer et al., 2018), no substantial changes occurred to the main summary statistics with memory consolidation measures. However, we observed a major shift in the moderation results. Specifically, the effect of scalp topography was statistically significant in these analyses, whereas we no longer found a significant effect of spindle measure on memory consolidation. Given that we removed one-third of the estimates and sample size, such statistical differences are to be expected. The details of both analyses can be found online (https://osf.io/wu6d7). In this article, we focus on the initial findings for the spindle-dependent memory consolidation using all effect sizes (k = 1204).

4. Discussion

The present meta-analysis aimed to **i)** determine if there is sufficient evidence to support the claim that sleep spindles facilitate memory processing and **ii)** assess how multiple factors, including memory type, spindle characteristics, and EEG scalp topography affect the sleep spindle-memory association. The literature search yielded 53 studies that were quantitatively analyzed. We found a small to moderate effect size for the association between sleep spindle and memory measures. Moderation analyses revealed that the association between sleep spindles and memory consolidation was stronger for procedural memory than for declarative memory. Neither spindle type (i.e. fast vs. slow) nor EEG scalp topography impacted the strength of the spindle-memory association, but we found a moderator effect of spindle measures, with the largest effect sizes for spindle frequency and spindle power. Exploratory analyses indicated no differences in effect size for the spindle-memory association between sleep protocols (nap vs. whole-night sleep) or sleep duration, indicating that sleep spindles may be a general mechanism supporting memory consolidation during sleep.

4.1. Sleep spindles and memory consolidation
The studies included in this meta-analysis reported both associations of trait learning abilities with spindle measures, and how sleep spindles are related to state-dependent consolidation of newly learnt material. We found that sleep spindles show a consistent relationship with both trait-like and state-like measures of memory, with a numerically higher effect size for state-dependent consolidation measures. We focused our analyses on a systematic investigation of sleep spindles on the consolidation of newly learnt information. Theoretical accounts of sleep-dependent memory consolidation postulate an important role of sleep spindles in memory consolidation. Mechanistically, the consolidation of declarative memories in NREM sleep is thought to depend on the interaction of characteristic oscillatory signatures, including sleep spindles. As proposed in the model of active systems consolidation during sleep, brain-wide reactivation and integration of hippocampal memories requires global synchronization and communication between cortical and subcortical neural networks (Diekelmann and Born, 2010; Klinzing

et al., 2019; Maingret et al., 2016). Sleep spindles, which are generated in thalamo-cortical circuits, may facilitate this dialogue. Cortical neurons are excited by thalamocortical neurons which generate spindle activity in the cortical field potential, while the neocortex in turn drives spindle onset and regulates their termination via projections to the thalamus (Timofeev et al., 2001; Timofeev and Chauvette, 2013), indicating a coordinated communication between cortical and subcortical structures. This dialogue between the thalamus and neocortex is instrumental to memory functions: optogenetic stimulation of thalamic sleep spindles drives cross-regional spindle co-occurrence and supports hippocampus-dependent learning (Latchoumane et al., 2017). Sleep spindles may thus be actively involved in the dialogue between segregated brain areas in the service of memory consolidation. Our finding that sleep spindles support memory performance, and the consolidation of newly learnt memories in particular, is thus in line with theoretical accounts of sleep-dependent memory consolidation and findings that implicate spindles in nighttime reactivation of memories. Although we could not directly determine the neural mechanisms by which sleep spindles exert their influence on memory functions, we systematically investigated potential electrophysiological and behavioral moderators to further characterize the spindle-memory relationship.

4.2. Spindle-dependent memory consolidation in different types of memory
In the current study, we observed that EEG sleep spindles are associated with memory consolidation in both major memory domains, consistent with the idea that sleep spindles are not only implicated in declarative memory (Peyrache and Seibt, 2020; Schabus et al., 2004) but also in procedural memory and skill learning (Boutin and Doyon, 2020; King et al., 2017; Walker et al., 2005). Interestingly, the strength of spindle-memory association was stronger for procedural than for declarative memory. Previous research on sequential motor learning tasks demonstrated that increases in BOLD signal in the cortico-striatal system during learning were correlated positively with both fast and slow spindle activity during later sleep (Barakat et al., 2013). As these tasks rely on hippocampal activity, next to the striatum, precuneus, prefrontal cortex, and primary motor cortex (Albouy et al., 2015, 2008; Boutin et al., 2018; King et al., 2017), a network of brain regions that overlaps with areas involved in declarative memory, the role of sleep spindles for the consolidation of sequential motor learning may be functionally similar to consolidation in the declarative memory system. Indeed, systems memory consolidation has recently been observed in the procedural domain (Boutin and Doyon, 2020; King et al., 2017; Schmid et al., 2020). Importantly, sleep spindles have been shown to foster changes in the neural substrate supporting memories and to promote brain-wide integration after learning in both declarative (Bergmann et al., 2012; Cowan et al., 2020; Jegou et al., 2019) and procedural memory tasks (Boutin et al., 2018; Fogel et al., 2017). While historically the role of spindles has mainly been discussed for declarative

memory, future theoretical accounts should thus consider that similar mechanisms may apply in the procedural domain.

Because hippocampal coordination is a central factor in theoretical accounts of sleep-dependent consolidation in the declarative domain, yet many procedural tasks do not require hippocampal activity during acquisition, we further tested whether sleep spindles equally contribute to procedural memory tasks with and without hippocampal involvement (Schendan et al., 2003; Walker et al., 2003). We observed no moderator effect of procedural memory task (non-hippocampal vs. hippocampal) on spindle-memory associations. This is in line with a recent meta-analysis that similarly reported a small effect of sleep on motor memory consolidation with no notable differences between various motor tasks including finger tapping and mirror tracing (Schmid et al., 2020). This is surprising, given an elaborate theoretical and experimental foundation that links spindle-related consolidation to a cross-regional dialogue between the hippocampus, thalamus, and neocortex (Antony et al., 2019; Latchoumane et al., 2017; Staresina et al., 2015) and stresses the role of the hippocampus reactivation-dependent consolidation. It has been suggested that the hippocampus could play a broader role in tasks traditionally considered hippocampal-independent, where it may help to reinstate extra-hippocampal memory traces (Sawangjit et al., 2018). Therefore, both hippocampal and non-hippocampal procedural memory tasks may indeed benefit similarly from spindle activity during sleep. Alternatively, sleep spindles may benefit memory via mechanisms independent of hippocampally coordinated reinstatement of learning-related activity. Indeed, it has been found that the thalamus can instruct plasticity in the primary visual cortex during a sleep period following visual discrimination learning and that sleep spindles mediate this effect (Durkin et al., 2017). Spindles' role in memory may thus be two-fold: In supporting thalamo-cortical and thalamo-hippocampal communication, they may aid systems-level consolidation, but their plasticity-promoting properties can also foster synaptic consolidation, more generally. Because also memory tasks with no hippocampal contribution showed a spindle-related consolidation benefit, we would argue that our results support this more global role for spindles in mnemonic processes such as consolidation.

4.3. Spindle characteristics and their relation to memory consolidation

4.3.1. Spindle Types and Topography

As mentioned in the introduction, many authors segregate sleep spindles into two main types according to their oscillatory frequency and topography, with slow spindles occurring over frontal regions and fast spindles dominating over parietal and central sites (Andrillon et al., 2011; Cox et al., 2017; Schabus et al., 2007; Urakami, 2008). It has further been hypothesized that slow frontal spindles may be preferentially generated by cortico-cortical or intracortical mechanisms, whereas faster parietal spindle activity is more closely re-

lated to thalamo-cortical communication (Ayoub et al., 2013; Timofeev and Chauvette, 2013). However, the distinct functional roles of these spindle types remain largely unknown. Given that spindle oscillations are widely distributed across the brain, it may seem plausible to assume that sleep spindles foster the strengthening of memory traces in different memory systems depending on spindle type (fast vs. slow), their location, or both. In the present meta-analysis, we assessed this question by examining the moderator effect of spindle types and of EEG scalp topography. Although spindle type and scalp topography did not impact the strength of the overall spindle-memory association, we observed that the procedural memory-spindle association was moderated by spindle types, with global and fast sleep spindles, showing the strongest effect. This result is in line with previous studies, which demonstrated that global and fast sleep spindles are related to memory improvement in motor procedural memory tasks (Barakat et al., 2013, 2011b; Fogel and Smith, 2006; Nishida and Walker, 2007; Simor et al., 2019; Tamaki et al., 2008). As fast sleep spindles dominate over parietal and central sites including motor regions, we believe that our results may reflect learning-related patterns of brain activation underlying localized spindle expression during sleep (for a recent study demonstrating this in declarative learning, see Petzka et al., 2022).

Importantly, in the literature, there is no clear consensus on the spectral distinction between fast and slow sleep spindles (De Gennaro and Ferrara, 2003). While most studies have highlighted a distinct scalp topography for fast and slow spindles (Cox et al., 2017; Schabus et al., 2007), several experiments reported overlapping M/EEG sources for faster and slower spindle components (Dehghani et al., 2010; Gumenyuk et al., 2009). Moreover, unlike in humans, a distinction between slow frontal and fast parietal spindles in mean frequency is not found in mice (Kim et al., 2015).

The way in which spindles manifest on the scalp might be linked to the structural anatomy of the generating network (Piantoni et al., 2016). In other words, the frequency variability might be caused by differences in anatomical structure, like the length or degree of myelination of thalamo-cortical white-matter tracts (Piantoni et al., 2017, 2013), which affect conduction velocity (Fields, 2015) and axonal propagation delays (O'Reilly and Nielsen, 2014). Alternatively, task-specific localization of neuronal activity may drive functional differences between spindle subtypes and differences in spindle topography. Speaking against the idea that slow and fast spindles rely on a distinct neural circuitry, a recent electrocorticography study with laminar profiling in humans suggested that slow and fast spindle subtypes are generated in cortical and thalamic circuits that have similar projections to cortical regions (Ujma et al., 2021). Although in the present meta-analysis, we did not observe effect size differences depending on EEG scalp location, our results do not contradict the notion that the topography of sleep spindles can reflect or even bear relevance to memory consolidation. A recent study suggests that participant-specific topographies of sleep

spindles are related to regional activation during memory encoding, and spindle activity during later sleep may be relevant for expressing plasticity in these regions (Petzka et al., 2022). Our topographical analyses do not reflect such subtle spatial patterns. Given what we know about spindle generation and their role in plasticity, determining their regional expression, however, appears a promising area for future research.

4.3.2. Spindle Measures

Spindle-memory associations can be measured using various sleep spindles properties including amplitude, count, density, duration, frequency, power, and SpA. However, it is still unknown whether the magnitude depends on specific spindle measures, which may reflect both similar or different biological underpinnings. In the present meta-analysis, we assessed a moderation effect of spindle measure and expected that the spindle-memory association would show a similar strength for different spindle measures, given their high intercorrelation (O'Reilly and Nielsen, 2014). Yet, we observed that spindle measures moderated the spindle-memory association with the strongest effect for spindle frequency and spindle power.

Our research found a strong effect of spindle peak frequency on memory consolidation. However, it is important to note that this finding should be interpreted with caution as there have been only two studies (k=27) reporting a correlation between spindle peak frequency and memory consolidation. Individual spindle peak frequency is known to vary broadly, in a trait-like manner across individuals (Werth et al., 1997), constant over different nights within a person, influenced by genetic factors, and resistant to experimental perturbations (De Gennaro et al., 2005, 2008; Purcell et al., 2017). Given that spindle frequency is associated with specific anatomical differences in brain structure (Saletin et al., 2013), the measure of spindle frequency may reflect individual differences in functional brain anatomy rather than sleep-specific mechanisms. Thus, participants with faster spindle frequency might generally show larger sleep-dependent consolidation benefits routed in anatomical advantages, rather than one individual's spindle peak frequency changing with more efficient processing of specific memory content during sleep. Despite the strong effect of spindle peak frequency on memory consolidation found in our meta-analysis, more research is needed to confirm these findings. Further studies are needed to provide a more robust understanding of the relationship between spindle peak frequency and memory consolidation, as has been done for alpha peak frequency regarding its role in cognitive performance (Cecere et al., 2015; Haegens et al., 2014; Mahjoory et al., 2019; Mierau et al., 2017).

The role of spindle power in brain plasticity has been previously shown in research on both animals and humans. In rodents, it has been demonstrated that spindle power is related to the degree of synchrony in neuronal activity, network synchronization, the rate of neural recruitment (Vyazovskiy et al., 2007), and also Ca2+ dendritic activity synchronization (Seibt et al., 2017). After lesions to corticothalamic projec-

tions one may expect that the synchrony of spindles is reduced, leading to a decrease in spindle power. Such a power reduction has indeed been observed in patients with unilateral hemispheric stroke (Gottselig et al., 2002) and with damage to the left thalamus (Mensen et al., 2018). Moreover, GABA reuptake inhibitors decrease spindle power and impair sleep-dependent consolidation of sequential motor learning (Feld et al., 2013). Spindle amplitude, which is proportional to spindle power, has also been shown to correlate with the spatial extent of spindle regional recruitment, reflecting a stronger degree of modulation in neuronal firing (Nir et al., 2011). We therefore argue that spindle power, which reflects neural synchrony and the degree of neuronal recruitment, could be directly related to sleep-dependent consolidation.

4.4. Exploratory Analyses

The vast majority of sleep and memory research has been conducted in either whole night sleep or nap sleep designs and several studies directly investigated whether the chosen sleep protocol directly affects sleep-memory associations (Cousins et al., 2021; Lo et al., 2014; Mednick et al., 2003; Pöhlchen et al., 2021; Sugawara et al., 2018), especially with regards to the role of sleep spindles in memory consolidation (Cousins et al., 2021; van Schalkwijk et al., 2019). A recent study on procedural memory observed naps and whole nights of sleep may both yield similar memory benefits, suggesting that daytime naps protect memories from deterioration, whereas whole night sleep improves performance, especially in procedural memory (van Schalkwijk et al., 2019). As spindle count and the number of reactivation events may increase with a full night of sleep, and the timing of sleep may have an impact on spindle density (van Schalkwijk et al., 2019), one could expect that nocturnal sleep would result in larger sleep-memory associations as compared to napping. In the current meta-analysis, however, no significant differences between sleep protocols were found, neither across all studies, nor separately in studies reporting declarative or procedural memory. We further confirmed this finding by moderation analyses with sleep duration, showing that sleep duration has no effect on the size of the observed spindle-memory association. Our results align with five previous meta-analyses reporting nearly the same effect size for short and long sleep durations in motor memory (Schmid et al., 2020) and also reporting no moderating effect of sleep protocol in episodic (Hokett et al., 2021) or emotional memory (Schäfer et al., 2020). Similarly, total sleep time was not found to be a significant predictor for an increase in the sleep benefit in episodic memory (Berres and Erdfelder, 2021) and was also not associated with memory and cognition in healthy older adults (Qin et al., 2022). It thus seems that spindles are a general mechanism for memory consolidation, regardless of sleep duration or the timing of sleep.

In the sleep literature, age-related brain alterations have been related to sleep physiology, and sleep spindles, in particular (Fogel et al., 2012, 2014; Martin et al., 2013; Muehlroth et al., 2020, 2019). As older adults also do not show sleep

benefits in episodic memory performance (Gui et al., 2017), it is intriguing to ask whether age plays a moderating role in the strength of the association between sleep spindles and memory. In our meta-analyses, however, we did not observe this modulatory effect of age, which aligns with other recent meta-analyses reporting that young and older adults maintain similarly strong sleep-memory relationships (Hokett et al., 2021). This finding is expected, as our sample is relatively young (M = 22.48, age range: 19-37 years), which limits our ability to examine age-related differences for some of the variables of interest. Since sleep studies are predominantly carried out with younger participants (Hokett et al., 2021), future research should more often include older individuals.

4.5. Statistical Considerations

In sleep and memory studies, methodological flexibility has often been combined with a shotgun analytical approach, in which a high number of statistical analyses were performed based on several M/EEG scalp locations, spindle types, or spindle measures (see: recent technical review by (Cox and Fell, 2020), with any significant effect interpreted as meaningful. While the studies reviewed here performed on average more than 20 statistical tests, only less than half of the studies used a correction for multiple comparisons. One way to diminish the risk of false positives in high-dimensional brain imaging data, such as multi-channel EEG recordings, is nonparametric permutation testing, which can take into account the spatial or temporal distribution of M/EEG data to reduce the number of comparisons or to adequately control for them (Maris and Oostenveld, 2007). However, only one study reviewed here implemented such cluster-level statistics on the correlation between sleep spindles and behavioral data (Dehnavi et al., 2019). It would be desirable that in the future, more studies looking at spindle-memory associations accounted for multiple comparisons in a similar way.

While any individual researcher has the freedom of defining a statistical threshold and the number of statistical tests they perform, we suggest that the field of sleep and memory research as a whole would benefit if these analytical approaches were chosen prior to statistical testing even before data collection has begun (Cox and Fell, 2020), as implemented in study preregistrations. Preregistrations motivate researchers to formulate hypotheses before seeing the data and can help research transparency and reproducibility by reducing biases (Gorgolewski and Poldrack, 2016; Ioannidis et al., 2014).

Another important statistical consideration is low statistical power because of small sample size (n) that poses a particular threat to the validity of sleep studies. It is known that studies with a small n can lead to an estimation of inflated effect sizes (Schäfer and Schwarz, 2019). In the current meta-analysis, we similarly observed this inflated effect size with smaller sample sizes. Correlating effect size and sample size across studies resulted in a significant association (Supplementary Fig 3), suggesting a clear indication of bias: larger effects result from smaller samples, while smaller effects were observed in the larger samples. We therefore suggest future

studies should aim for larger sample sizes and complete an a priori sample size estimation to assure sufficient statistical power (Cordi and Rasch, 2021). This would reduce the risk of obtaining spuriously high effect size measures due to the higher variability in these estimates at a smaller sample size. Since sleep studies have high monetary and time demands, collaborative networks and data-sharing initiatives may also be a promising future avenue for the advancement of sleep science.

Lastly, we observed substantial publication bias in our meta-analysis, similar to other recent meta-analyses of sleep and episodic memory performance (Hokett et al., 2021), complex associative memory processing (Chatburn et al., 2014) or explicit motor sequence learning (Pan and Rickard, 2015; Rickard et al., 2022; but see: Schmid et al., 2020). The results of meta-analyses may overestimate the true effect if publication bias is observed (Ferguson and Brannick, 2012). Several techniques have been developed to correct for publication bias, including trim-and-fill analyses (Duval and Tweedie, 2000a, 2000b) or precision-effect test and precision-effect estimate with standard errors (PET-PEESE; Bartoš et al., 2022). In the current meta-analysis, we employed a moderation analysis testing whether reported effect sizes in published articles were significantly larger than unreported effect sizes. We observed no differences between reported vs. unreported effect sizes across all studies. Unexpectedly, when performing this analysis separately for studies of declarative and procedural memory, we found that reporting status had significant yet opposite effects on the size of the spindle-memory association. While correlation coefficients for declarative memory were higher when reported in the literature, indicating that publication bias led to an overestimation of the spindle-memory association in these studies, for procedural memory, the unreported correlation coefficients showed higher values. This suggests that the spindle-memory association may actually have been underestimated for procedural tasks.

Using our publicly available datasets in OSF (https://osf.io/wu6d7), Ujma (2022) have recently computed the association between sleep spindles and memory by adjusting publication bias using a non-standard multivariate implementation of the above-mentioned PET-PEESE methods. He observed that correcting for publication bias in this way reduces the correlation between sleep spindles and memory function to insignificance. Although this analysis was only performed on all effect size measures we obtained without considering whether general learning ability, memory consolidation, procedural memory consolidation or declarative memory consolidation was investigated, this finding still raises an important question: if spindles were not related to memory consolidation, why do they appear so strongly associated with memory reactivation during sleep (Cairney et al., 2018; Schönauer et al., 2017; Schreiner et al., 2021; Wang et al., 2019)? More importantly, how could they have a causal role in relaying the behavioral effects of targeted memory reactivation (Antony et al., 2018)? We would argue that rather than refraining from studying the role

of sleep spindles in human memory because it may not yield significant results, research needs to more directly address the functional relevance of spindles in memory formation. Lastly, given the significant heterogeneity in effect sizes and the low sample sizes reported in the studies included in this meta-analysis, we urge a continued evaluation of true effect sizes and publication bias as the field progresses.

5. Limitations

This meta-analysis aimed to assess the strength of the correlation between sleep spindles and memory performance. Given this objective, we have not assessed learning-induced changes in sleep spindles. Therefore, investigating the impact of new learning on subsequent sleep seems to offer a promising direction for future meta-analytic research on sleep spindles and memory consolidation. We further excluded several studies, in particular, studies using manipulations (e.g., targeted memory reactivation, TMR). In TMR, the timing of cue presentation relative to sleep spindles and to the phase of ongoing SOs determines their efficacy in reactivation and memory consolidation (Antony et al., 2019, 2018; Cox et al., 2014). More research is needed to determine how strongly TMR impacts the relationship between sleep oscillatory events and memory performance. A focus on coupling between SO and sleep spindles could provide valuable additional insights into the nature of the relationship between sleep spindles and memory consolidation. However, this question was beyond the scope of the current meta-analysis. While a strength of the current study is a large participant sample and a high number of estimates, most of the included individual experiments had relatively small sample sizes, which might result in an imprecise estimation of the respective effect sizes. Future studies in the field of sleep and memory research should thus aim for larger sample sizes. Lastly, we broadly clustered EEG channels for the moderator analyses into frontal, central, and parietal regions, due to a large variability in recording methods between studies. Using this broad clustering, we found no moderating effect of spindle location on the spindle-memory association. Finer regions of interest, or other topographical approaches would allow investigation of this relationship with better spatial precision. This avenue seems highly promising in light of recent findings that spindle topographies track regional engagement during memory encoding (Petzka et al., 2022). In this regard, it may be fruitful to pursue the question of spindle topography and memory association further depending on different learning tasks.

6. Conclusions

In the present study, we examined meta-analytical evidence for an association between sleep spindles and memory performance. We assessed how multiple factors, including memory type, spindle type, different spindle measures, and EEG scalp topography affect this relationship. The results of this meta-analysis have demonstrated a medium but robust association between sleep spindles and memory that was largely independent of spindle type or spindle topography. Collectively, our findings suggest that sleep spindles are involved in learning, and may present a general physiological mechanism for sleep-dependent memory consolidation. Spindle power, in particular, may be a promising predictor of sleep-dependent memory consolidation. We hope future studies can take advantage of our findings to highlight the critical gaps in the understanding of the complex interaction between sleep spindles and memory.

7. Data availability statement

All data files, associated codebooks, and analysis scripts are publicly available on the study's page on the Open Science Framework: (https://osf.io/wu6d7/).

8. Acknowledgement

We thank all authors, who shared their data (of the non-significant results for the correlation values between spindles and memory) openly with us, which extensively improved the manuscript. We further thank Eduard Stroukov for his help on data assessment and Dorothee Pöhlchen and Nora Roüast for their insightful comments on a previous draft of this manuscript.

9. Funding

This study was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project number: 426865207, and the BMBF Germany.

10. References

1. Ackermann, S., Hartmann, F., Papassotiropoulos, A., De Quervain, D.J.F., Rasch, B., 2015. No associations between interindividual differences in sleep parameters and episodic memory consolidation. Sleep 38, 951–959. https://doi.org/10.5665/sleep.4748

2. Albouy, G., Fogel, S., King, B.R., Laventure, S., Benali, H., Karni, A., Carrier, J., Robertson, E.M., Doyon, J., 2015. Maintaining vs. enhancing motor sequence memories: Respective roles of striatal and hippocampal systems. Neuroimage 108, 423–434. https://doi.org/10.1016/j.neuroimage.2014.12.049

3. Albouy, G., Sterpenich, V., Balteau, E., Vandewalle, G., Desseilles, M., Dang-Vu, T., Darsaud, A., Ruby, P., Luppi, P.H., Degueldre, C., Peigneux, P., Luxen, A., Maquet, P., 2008. Both the Hippocampus and Striatum Are Involved in Consolidation of Motor Sequence Memory. Neuron 58, 261–272. https://doi.org/10.1016/J.NEURON.2008.02.008

4. Andrillon, T., Nir, Y., Staba, R.J., Ferrarelli, F., Cirelli, C., Tononi, G., Fried, I., 2011. Sleep spindles in humans: Insights from intracranial EEG and unit recordings. Journal of Neuroscience 31, 17821–17834. https://doi.org/10.1523/JNEUROSCI.2604-11.2011

5. Antony, J.W., Gobel, E.W., O'Hare, J.K., Reber, P.J., Paller, K.A., 2012. Cued memory reactivation during sleep influences skill learning. Nat Neurosci 15, 1114–1116. https://doi.org/10.1038/nn.3152

6. Antony, J.W., Piloto, L., Wang, M., Pacheco, P., Norman, K.A., Paller, K.A., 2018. Sleep Spindle Refractoriness Segregates Periods of Memory Reactivation. Current Biology 28, 1736-1743.e4. https://doi.org/10.1016/j.cub.2018.04.020

7. Antony, J.W., Schönauer, M., Staresina, B.P., Cairney, S.A., 2019. Sleep Spindles and Memory Reprocessing. Trends Neurosci 42, 1–3. https://doi.org/10.1016/j.tins.2018.09.012

8. Ayoub, A., Aumann, D., Hörschelmann, A., Kouchekmanesch, A., Paul, P., Born, J., Marshall, L., 2013. Differential effects on fast and slow spindle activity, and the sleep slow oscillation in humans with carbamazepine and flunarizine to antagonize voltage-dependent Na+ and Ca2+ channel activity. Sleep 36, 905–911. https://doi.org/10.5665/sleep.2722

9. Barakat, M., Carrier, J., Debas, K., Lungu, O., Fogel, S., Vandewalle, G., Hoge, R.D., Bellec, P., Karni, A., Ungerleider, L.G., Benali, H., Doyon, J., 2013. Sleep spindles predict neural and behavioral changes in motor sequence consolidation. Hum Brain Mapp 34, 2918–2928. https://doi.org/10.1002/hbm.22116

10. Barakat, M., Doyon, J., Debas, K., Vandewalle, G., Morin, A., Poirier, G., Martin, N., Lafortune, M., Karni, A., Ungerleider, L.G., Benali, H., Carrier, J., 2011a. Fast and slow spindle involvement in the consolidation of a new motor sequence. Behavioural Brain Research 217, 117–121. https://doi.org/10.1016/j.bbr.2010.10.019

11. Barakat, M., Doyon, J., Debas, K., Vandewalle, G., Morin, A., Poirier, G., Martin, N., Lafortune, M., Karni, A., Ungerleider, L.G., Benali, H., Carrier, J., 2011b. Fast and slow spindle involvement in the consolidation of a new motor sequence. Behavioural Brain Research 217, 117–121. https://doi.org/10.1016/j.bbr.2010.10.019

12. Bartoš, F., Maier, M., Quintana, D.S., Wagenmakers, E.J., 2022. Adjusting for Publication Bias in JASP and R: Selection Models, PET-PEESE, and Robust Bayesian Meta-Analysis. Adv Methods Pract Psychol Sci 5. https://doi.org/10.1177/25152459221109259

13. Bergmann, T.O., Mölle, M., Diedrichs, J., Born, J., Siebner, H.R., 2012. Sleep spindle-related reactivation of category-specific cortical regions after learning face-scene associations. Neuroimage 59, 2733–2742.

14. Berres, S., Erdfelder, E., 2021. The sleep benefit in episodic memory: An integrative review and a meta-analysis. Psychol Bull 147, 1309–1353. https://doi.org/10.1037/bul0000350

15. Blaskovich, B., Szollosi, Á., Gombos, F., Racsmány, M., Simor, P., 2017. The benefit of directed forgetting persists after a daytime nap: The role of spindles and rapid eye movement sleep in the consolidation of relevant memories. Sleep 40. https://doi.org/10.1093/sleep/zsw076

16. Bonjean, M., Baker, T., Lemieux, M., Timofeev, I., Sejnowski, T., Bazhenov, M., 2011. Corticothalamic feedback controls sleep spindle duration in vivo. Journal of Neuroscience 31, 9124–9134. https://doi.org/10.1523/JNEUROSCI.0077-11.2011

17. Borenstein, M., Hedges, L. V, Higgins, J.P.T., Rothstein, H.R., 2009. Complex Data Structures Introduction to Meta-Analysis. Overview.

18. Boutin, A., Doyon, J., 2020. A sleep spindle framework for motor memory consolidation. Philosophical Transactions of the Royal Society B: Biological Sciences 375, 1–7. https://doi.org/10.1098/rstb.2019.0232

19. Boutin, A., Pinsard, B., Boré, A., Carrier, J., Fogel, S.M., Doyon, J., 2018. Transient synchronization of hippocampo-striato-thalamo-cortical networks during sleep spindle oscillations induces motor memory consolidation. Neuroimage 169, 419–430. https://doi.org/10.1016/j.neuroimage.2017.12.066

20. Cairney, S.A., Guttesen, A.á. V., El Marj, N., Staresina, B.P., 2018. Memory Consolidation Is Linked to Spindle-Mediated Information Processing during Sleep. Current Biology 28, 948-954.e4. https://doi.org/10.1016/j.cub.2018.01.087

21. Cecere, R., Rees, G., Romei, V., 2015. Individual differences in alpha frequency drive crossmodal illusory perception. Current Biology 25, 231–235. https://doi.org/10.1016/J.CUB.2014.11.034

22. Chatburn, A., Lushington, K., Kohler, M.J., 2014. Complex associative memory processing and sleep: A systematic review and meta-analysis of behavioural evidence and underlying EEG mechanisms. Neurosci Biobehav Rev 47, 646–655. https://doi.org/10.1016/j.neubiorev.2014.10.018

23. Cheung, M.W.L., 2019. A Guide to Conducting a Meta-Analysis with Non-Independent Effect Sizes. Neuropsychol Rev 29, 387–396. https://doi.org/10.1007/s11065-019-09415-6

24. Clemens, Z., Fabó, D., Halász, P., 2006. Twenty-four hours retention of visuospatial memory correlates with the number of parietal sleep spindles. Neurosci Lett 403, 52–56. https://doi.org/10.1016/j.neulet.2006.04.035

25. Clemens, Z., Fabó, D., Halász, P., 2005. Overnight verbal memory retention correlates with the number of sleep spindles. Neuroscience 132, 529–535. https://doi.org/10.1016/j.neuroscience.2005.01.011

26. Contreras, D., Steriade, M., 1996. Spindle oscillation in cats: The role of corticothalamic feedback in a thalamically generated rhythm. Journal of Physiology 490, 159–179. https://doi.org/10.1113/JPHYSIOL.1996.SP021133

27. Cook, R.D., 1977. Detection of Influential Observation in Linear Regression. Technometrics 19, 15–18. https://doi.org/10.1080/00401706.1977.10489493

28. Cordi, M.J., Rasch, B., 2021. How robust are sleep-mediated memory benefits? Curr Opin Neurobiol 67, 1–7. https://doi.org/10.1016/j.conb.2020.06.002

29. Cousins, J.N., El-Deredy, W., Parkes, L.M., Hennies, N., Lewis, P.A., 2014. Cued memory reactivation during slow-wave sleep promotes explicit knowledge of a motor sequence. Journal of Neuroscience 34, 15870–15876. https://doi.org/10.1523/JNEUROSCI.1011-14.2014

30. Cousins, J.N., Leong, R.L.F., Jamaluddin, S.A., Ng, A.S.C., Ong, J.L., Chee, M.W.L., 2021. Splitting sleep between the night and a daytime nap reduces homeostatic sleep pressure and enhances long-term memory. Sci Rep 11. https://doi.org/10.1038/s41598-021-84625-8

31. Cowan, E., Liu, A., Henin, S., Kothare, S., Devinsky, O., Davachi, X.L., 2020. Sleep spindles promote the restructuring of memory representations in ventromedial prefrontal cortex through enhanced hippocampal–cortical functional connectivity. Journal of Neuroscience 40, 1909–1919. https://doi.org/10.1523/JNEUROSCI.1946-19.2020

32. Cox, R., Fell, J., 2020. Analyzing human sleep EEG: A methodological primer with code implementation. Sleep Med Rev 54, 101353. https://doi.org/10.1016/j.smrv.2020.101353

33. Cox, R., Hofman, W.F., Talamini, L.M., 2012. Involvement of spindles in memory consolidation is slow wave sleep-specific. Learning and Memory 19, 264–267. https://doi.org/10.1101/lm.026252.112

34. Cox, R., Korjoukov, I., De Boer, M., Talamini, L.M., 2014. Sound Asleep: Processing and Retention of Slow Oscillation Phase-Targeted Stimuli. PLoS One 9, e101567. https://doi.org/10.1371/JOURNAL.PONE.0101567

35. Cox, R., Schapiro, A.C., Manoach, D.S., Stickgold, R., 2017. Individual differences in frequency and topography of slow and fast sleep spindles. Front Hum Neurosci 11, 433. https://doi.org/10.3389/FNHUM.2017.00433/BIBTEX

36. De Gennaro, L., Ferrara, M., 2003. Sleep spindles: An overview. Sleep Med Rev 7, 423–440. https://doi.org/10.1053/smrv.2002.0252

37. Dehghani, N., Cash, S.S., Chen, C.C., Hagler, D.J., Huang, M., Dale, A.M., Halgren, E., 2010. Divergent Cortical Generators of MEG and EEG during Human Sleep Spindles Suggested by Distributed Source Modeling. PLoS One 5, e11454. https://doi.org/10.1371/JOURNAL.PONE.0011454

38. Dehnavi, F., Moghimi, S., Sadrabadi Haghighi, S., Safaie, M., Ghorbani, M., 2019. Opposite effect of motivated forgetting on sleep spindles during stage 2 and slow wave sleep. Sleep 42. https://doi.org/10.1093/sleep/zsz085

39. Diekelmann, S., Biggel, S., Rasch, B., Born, J., 2012. Offline consolidation of memory varies with time in slow wave sleep and can be accelerated by cuing memory reactivations. Neurobiol Learn Mem 98, 103–111. https://doi.org/10.1016/j.nlm.2012.07.002

40. Diekelmann, S., Born, J., 2010. The memory function of sleep. Nat Rev Neurosci 11, 114–126. https://doi.org/10.1038/nrn2762

41. Durkin, J., Suresh, A.K., Colbath, J., Broussard, C., Wu, J., Zochowski, M., Aton, S.J., 2017. Cortically coordinated NREM thalamocortical oscillations play an essential, instructive role in visual system plasticity. Proc Natl Acad Sci U S A 114, 10485–10490. https://doi.org/10.1073/pnas.1710613114

42. Duval, S., Tweedie, R., 2000a. Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. Biometrics 56, 455–463. https://doi.org/10.1111/j.0006-341X.2000.00455.x

43. Duval, S., Tweedie, R., 2000b. A Nonparametric "Trim and Fill" Method of Accounting for Publication Bias in Meta-Analysis. J Am Stat Assoc 95, 89–98. https://doi.org/10.1080/01621459.2000.10473905

44. Feld, G.B., Wilhelm, I., Ma, Y., Groch, S., Binkofski, F., Mölle, M., Born, J., 2013. Slow wave sleep induced by GABA agonist tiagabine fails to benefit memory consolidation. Sleep 36, 1317–1326. https://doi.org/10.5665/sleep.2954

45. Ferguson, C.J., Brannick, M.T., 2012. Publication Bias in Psychological Science: Prevalence, Methods for Identifying and Controlling, and Implications for the Use of Meta-Analyses. Psychol Methods. https://doi.org/10.1037/a0024445.supp

46. Fernandez, L.M.J., Lüthi, A., 2020. Sleep spindles: Mechanisms and functions. Physiol Rev 100, 805–868. https://doi.org/10.1152/physrev.00042.2018

47. Fernández-Castilla, B., Jamshidi, L., Declercq, L., Beretvas, S.N., Onghena, P., Van den Noortgate, W., 2020. The application of meta-analytic (multi-level) models with multiple random effects: A systematic review. Behav Res Methods 52, 2031–2052. https://doi.org/10.3758/s13428-020-01373-9

48. Fernández-Ruiz, A., Oliva, A., de Oliveira, E.F., Rocha-Almeida, F., Tingley, D., Buzsáki, G., 2019. Long-duration hippocampal sharp wave ripples improve memory. Science (1979) 364, 1082–1086. https://doi.org/10.1126/science.aax0758

49. Fields, R.D., 2015. A new mechanism of nervous system plasticity: Activity-dependent myelination. Nat Rev Neurosci 16, 756–767. https://doi.org/10.1038/nrn4023

50. Fogel, S., Albouy, G., King, B.R., Lungu, O., Vien, C., Bore, A., Pinsard, B., Benali, H., Carrier, J., Doyon, J., 2017. Reactivation or transformation? Motor memory consolidation associated with cerebral activation time-locked to sleep spindles. PLoS One 12. https://doi.org/10.1371/journal.pone.0174755

51. Fogel, S., Martin, N., Lafortune, M., Barakat, M., Debas, K., Laventure, S., Latreille, V., Gagnon, J.F., Doyon, J., Carrier, J., 2012. NREM sleep oscillations and brain plasticity in aging. Front Neurol. https://doi.org/10.3389/fneur.2012.00176

52. Fogel, S.M., Albouy, G., Vien, C., Popovicci, R., King, B.R., Hoge, R., Jbabdi, S., Benali, H., Karni, A., Maquet, P., Carrier, J., Doyon, J., 2014. fMRI and sleep correlates of the age-related impairment in motor memory consolidation. Hum Brain Mapp 35, 3625–3645. https://doi.org/10.1002/hbm.22426

53. Fogel, S.M., Smith, C.T., 2006. Learning-dependent changes in sleep spindles and Stage 2 sleep. J Sleep Res 15, 250–255. https://doi.org/10.1111/j.1365-2869.2006.00522.x

54. Gais, S., Mölle, M., Helms, K., Born, J., 2002. Learning-dependent increases in sleep spindle density. Journal of Neuroscience 22, 6830–6834. https://doi.org/10.1523/jneurosci.22-15-06830.2002

55. Genzel, L., Bäurle, A., Potyka, A., Wehrle, R., Adamczyk, M., Friess, E., Steiger, A., Dresler, M., 2014. Diminished nap effects on memory consolidation are seen under oral contraceptive use. Neuropsychobiology 70, 253–261. https://doi.org/10.1159/000369022

56. Girardeau, G., Benchenane, K., Wiener, S.I., Buzsáki, G., Zugaro, M.B., 2009. Selective suppression of hippocampal ripples impairs spatial memory. Nat Neurosci 12, 1222–1223. https://doi.org/10.1038/nn.2384

57. Gonzalez, C., Jiang, X., Gonzalez-Martinez, J., Halgren, E., 2022. Human Spindle Variability. J Neurosci 42, 4517–4537. https://doi.org/10.1523/JNEUROSCI.1786-21.2022

58. Gorgolewski, K.J., Poldrack, R.A., 2016. A Practical Guide for Improving Transparency and Reproducibility in Neuroimaging Research. PLoS Biol 14, 1–13. https://doi.org/10.1371/journal.pbio.1002506

59. Gottselig, J.M., Bassetti, C.L., Achermann, P., 2002. Power and coherence of sleep spindle frequency activity following hemispheric stroke, Brain. https://doi.org/10.1093/brain/awf021

60. Griessenberger, H., Heib, D.P.J., Lechinger, J., Luketina, N., Petzka, M., Moeckel, T., Hoedlmoser, K., Schabus, M., 2013. Susceptibility to Declarative Memory Interference is Pronounced in Primary Insomnia. PLoS One 8. https://doi.org/10.1371/journal.pone.0057394

61. Gui, W.-J., Li, H.-J., Guo, Y.-H., Peng, P., Lei, X., Yu, J., 2017. Age-related differences in sleep-based memory consolidation: A meta-analysis. https://doi.org/10.1016/j.neuropsychologia.2017.02.001

62. Gumenyuk, V., Roth, T., Moran, J.E., Jefferson, C., Bowyer, S.M., Tepley, N., Drake, C.L., 2009. Cortical locations of maximal spindle activity: Magnetoencephalography (MEG) study. J Sleep Res 18, 245–253. https://doi.org/10.1111/J.1365-2869.2008.00717.X

63. Haegens, S., Cousijn, H., Wallis, G., Harrison, P.J., Nobre, A.C., 2014. Inter- and intra-individual variability in alpha peak frequency. Neuroimage 92, 46–55. https://doi.org/10.1016/j.neuroimage.2014.01.049

64. Hedges, L., Olkin, I., 2014. Statistical methods for meta-analysis.

65. Higgins, J.P.T., Thompson, S.G., 2002. Quantifying heterogeneity in a meta-analysis. Stat Med 21, 1539–1558. https://doi.org/10.1002/sim.1186

66. Hokett, E., Arunmozhi, A., Campbell, J., Verhaeghen, P., Duarte, A., 2021. A systematic review and meta-analysis of individual differences in naturalistic sleep quality and episodic memory performance in young and older adults. Neurosci Biobehav Rev. https://doi.org/10.1016/j.neubiorev.2021.05.010

67. Holz, J., Piosczyk, H., Feige, B., Spiegelhalder, K., Baglioni, C., Riemann, D., Nissen, C., 2012. EEG sigma and slow-wave activity during NREM sleep correlate with overnight declarative and procedural memory consolidation. J Sleep Res 21, 612–619. https://doi.org/10.1111/j.1365-2869.2012.01017.x

68. Hu, X., Cheng, L.Y., Chiu, M.H., Paller, K.A., 2020. Promoting memory consolidation during sleep: A meta-analysis of targeted memory reactivation. Psychol Bull 146, 218–244. https://doi.org/10.1037/bul0000223

69. Ioannidis, J.P.A., Munafò, M.R., Fusar-Poli, P., Nosek, B.A., David, S.P., 2014. Publication and other reporting biases in cognitive sciences: Detection, prevalence, and prevention. Trends Cogn Sci 18, 235–241. https://doi.org/10.1016/j.tics.2014.02.010

70. Jegou, A., Schabus, M., Gosseries, O., Dahmen, B., Albouy, G., Desseilles, M., Sterpenich, V., Phillips, C., Maquet, P., Grova, C., Dang-Vu, T.T., 2019. Cortical reactivations during sleep spindles following declarative learning. Neuroimage 195, 104–112. https://doi.org/10.1016/j.neuroimage.2019.03.051

71. Ji, D., Wilson, M.A., 2007. Coordinated memory replay in the visual cortex and hippocampus during sleep. Nat Neurosci 10, 100–107. https://doi.org/10.1038/nn1825

72. Kim, D., Hwang, E., Lee, M., Sung, H., Choi, J.H., 2015. Characterization of topographically specific sleep spindles in mice. Sleep 38, 85–96. https://doi.org/10.5665/sleep.4330

73. King, B.R., Hoedlmoser, K., Hirschauer, F., Dolfen, N., Albouy, G., 2017. Sleeping on the motor engram: The multifaceted nature of sleep-related motor memory consolidation. Neurosci Biobehav Rev 80, 1–22. https://doi.org/10.1016/j.neubiorev.2017.04.026

74. Klinzing, J.G., Niethard, N., Born, J., 2019. Mechanisms of systems memory consolidation during sleep. Nat Neurosci 22. https://doi.org/10.1038/s41593-019-0467-3

75. Konstantopoulos, S., 2011. Fixed effects and variance components estimation in three-level meta-analysis. Res Synth Methods 2, 61–76. https://doi.org/10.1002/JRSM.35

76. Latchoumane, C.F. V., Ngo, H.V. V., Born, J., Shin, H.S., 2017. Thalamic Spindles Promote Memory Formation during Sleep through Triple Phase-Locking of Cortical, Thalamic, and Hippocampal Rhythms. Neuron 95, 424-435.e6. https://doi.org/10.1016/j.neuron.2017.06.025

77. Laventure, S., Pinsard, B., Lungu, O., Carrier, J., Fogel, S., Benali, H., Lina, J.M., Boutin, A., Doyon, J., 2018. Beyond spindles: Interactions between sleep spindles and boundary frequencies during cued reactivation of motor memory representations. Sleep 41. https://doi.org/10.1093/sleep/zsy142

78. Lo, J.C., Dijk, D.-J., Groeger, J.A., 2014. Comparing the Effects of Nocturnal Sleep and Daytime Napping on Declarative Memory Consolidation. PLoS One 9, 108100. https://doi.org/10.1371/journal.pone.0108100

79. Lo, J.C., Groeger, J.A., Cheng, G.H., Dijk, D.J., Chee, M.W.L., 2016. Self-reported sleep duration and cognitive performance in older adults: A systematic review and meta-analysis. Sleep Med 17, 87–98. https://doi.org/10.1016/J.SLEEP.2015.08.021

80. Lustenberger, C., Maric, A., Dürr, R., Achermann, P., Huber, R., 2012. Triangular Relationship between Sleep Spindle Activity, General Cognitive Ability and the Efficiency of Declarative Learning. PLoS One 7. https://doi.org/10.1371/journal.pone.0049561

81. Lüthi, A., 2014. Sleep spindles: Where they come from, what they do. Neuroscientist. https://doi.org/10.1177/1073858413500854

82. Lutz, N.D., Admard, M., Genzoni, E., Born, J., Rauss, K., 2021. Occipital sleep spindles predict sequence learning in a visuo-motor task. Sleep 44. https://doi.org/10.1093/sleep/zsab056

83. Mahjoory, K., Cesnaite, E., Hohlefeld, F.U., Villringer, A., Nikulin, V. v., 2019. Power and temporal dynamics of alpha oscillations at rest differentiate cognitive performance involving sustained and phasic cognitive control. Neuroimage 188, 135–144. https://doi.org/10.1016/j.neuroimage.2018.12.001

84. Maingret, N., Girardeau, G., Todorova, R., Goutierre, M., Zugaro, M., 2016. Hippocampo-cortical coupling mediates memory consolidation during sleep. Nat Neurosci 19, 959–964. https://doi.org/10.1038/nn.4304

85. Maris, E., Oostenveld, R., 2007. Nonparametric statistical testing of EEG- and MEG-data. J Neurosci Methods 164, 177–190. https://doi.org/10.1016/j.jneumeth.2007.03.024

86. Marshall, L., Helgadóttir, H., Mölle, M., Born, J., 2006. Boosting slow oscillations during sleep potentiates memory. Nature 444, 610–613. https://doi.org/10.1038/nature05278

87. Martin, N., Lafortune, M., Godbout, J., Barakat, M., Robillard, R., Poirier, G., Bastien, C., Carrier, J., 2013. Topography of age-related changes in sleep spindles. Neurobiol Aging 34, 468–476. https://doi.org/10.1016/j.neurobiolaging.2012.05.020

88. McCormick, L., Nielsen, T., Nicolas, A., Ptito, M., Montplaisir, J., 1997. Topographical distribution of spindles and K-complexes in normal subjects. Sleep 20, 939–941. https://doi.org/10.1093/sleep/20.11.939

89. Mednick, S., Nakayama, K., Stickgold, R., 2003. Sleep-dependent learning: A nap is as good as a night. Nat Neurosci 6, 697–698. https://doi.org/10.1038/nn1078

90. Mednick, S.C., McDevitt, E.A., Walsh, J.K., Wamsley, E., Paulus, M., Kanady, J.C., Drummond, S.P.A., 2013. The critical role of sleep spindles in hippocampal-dependent memory: A pharmacology study. Journal of Neuroscience 33, 4494–4504. https://doi.org/10.1523/JNEUROSCI.3127-12.2013

91. Mensen, A., Poryazova, R., Huber, R., Bassetti, C.L., 2018. Individual spindle detection and analysis in high-density recordings across the night and in thalamic stroke. Sci Rep 8, 1–11. https://doi.org/10.1038/s41598-018-36327-x

92. Mierau, A., Klimesch, W., Lefebvre, J., 2017. State-dependent alpha peak frequency shifts: Experimental evidence, potential mechanisms and functional implications. Neuroscience 360, 146–154. https://doi.org/10.1016/j.neuroscience.2017.07.037

93. Mölle, M., Bergmann, T.O., Marshall, L., Born, J., 2011. Fast and slow spindles during the sleep slow oscillation: Disparate coalescence and engagement in memory processing. Sleep 34, 1411–1421. https://doi.org/10.5665/SLEEP.1290

94. Mölle, M., Eschenko, O., Gais, S., Sara, S.J., Born, J., 2009. The influence of learning on sleep slow oscillations and associated spindles and ripples in humans and rats. European Journal of Neuroscience 29, 1071–1081. https://doi.org/10.1111/j.1460-9568.2009.06654.x

95. Morin, A., Doyon, J., Dostie, V., Barakat, M., Tahar, A.H., Korman, M., Benali, H., Karni, A., Ungerleider, L.G., Carrier, J., 2008. Motor sequence learning increases sleep spindles and fast frequencies in post-training sleep. Sleep 31, 1149–1156. https://doi.org/10.5665/sleep/31.8.1149

96. Muehlroth, B.E., Sander, M.C., Fandakova, Y., Grandy, T.H., Rasch, B., Lee Shing, Y., Werkle-Bergner, M., 2020. Memory quality modulates the effect of aging on memory consolidation during sleep: Reduced maintenance but intact gain. Neuroimage 209. https://doi.org/10.1016/j.neuroimage.2019.116490

97. Muehlroth, B.E., Sander, M.C., Fandakova, Y., Grandy, T.H., Rasch, B., Shing, Y.L., Werkle-Bergner, M., 2019. Precise Slow Oscillation–Spindle Coupling Promotes Memory Consolidation in Younger and Older Adults. Sci Rep 9. https://doi.org/10.1038/s41598-018-36557-z

98. Ngo, H.V., Fell, J., Staresina, B., 2020. Sleep spindles mediate hippocampal-neocortical coupling during long-duration ripples. Elife 9, 1–18. https://doi.org/10.7554/eLife.57011

99. Ngo, H.-V. v, Staresina, B.P., 2022. Shaping overnight consolidation via slow-oscillation closed-loop targeted memory reactivation. https://doi.org/10.1073/pnas

100. Niethard, N., Burgalossi, A., Born, J., 2017. Plasticity during sleep is linked to specific regulation of cortical circuit activity. Front Neural Circuits 11. https://doi.org/10.3389/fncir.2017.00065

101. Niethard, N., Ngo, H.V. V., Ehrlich, I., Born, J., 2018. Cortical circuit activity underlying sleep slow oscillations and spindles. Proc Natl Acad Sci U S A 115, E9220–E9229. https://doi.org/10.1073/pnas.1805517115

102. Nir, Y., Staba, R.J., Andrillon, T., Vyazovskiy, V. V., Cirelli, C., Fried, I., Tononi, G., 2011. Regional Slow Waves and Spindles in Human Sleep. Neuron 70, 153–169. https://doi.org/10.1016/j.neuron.2011.02.043

103. Nishida, M., Nakashima, Y., Nishikawa, T., 2016. Slow sleep spindle and procedural memory consolidation in patients with major depressive disorder. Nat Sci Sleep 8, 63–72. https://doi.org/10.2147/NSS.S100337

104. Nishida, M., Walker, M.P., 2007. Daytime naps, motor memory consolidation and regionally specific sleep spindles. PLoS One 2. https://doi.org/10.1371/journal.pone.0000341

105. O'Reilly, C., Nielsen, T., 2014. Assessing EEG sleep spindle propagation. Part 2: Experimental characterization. J Neurosci Methods 221, 215–227. https://doi.org/10.1016/j.jneumeth.2013.08.014

106. Page, M.J., McKenzie, J.E., Bossuyt, P.M., Boutron, I., Hoffmann, T.C., Mulrow, C.D., Shamseer, L., Tetzlaff, J.M., Akl, E.A., Brennan, S.E., Chou, R., Glanville, J., Grimshaw, J.M., Hróbjartsson, A., Lalu, M.M., Li, T., Loder, E.W., Mayo-Wilson, E., McDonald, S., McGuinness, L.A., Stewart, L.A., Thomas, J., Tricco, A.C., Welch, V.A., Whiting, P., Moher, D., 2021. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. The BMJ. https://doi.org/10.1136/bmj.n71

107. Pan, S.C., Rickard, T.C., 2015. Sleep and motor learning: Is there room for consolidation? Psychol Bull 141, 812–834. https://doi.org/10.1037/BUL0000009

108. Peigneux, P., Laureys, S., Fuchs, S., Destrebecqz, A., Collette, F., Delbeuck, X., Phillips, C., Aerts, J., del Fiore, G., Degueldre, C., Luxen, A., Cleeremans, A., Maquet, P., 2003. Learned material content and acquisition level modulate cerebral reactivation during posttraining rapid-eye-movements sleep. Neuroimage 20, 125–134. https://doi.org/10.1016/S1053-8119(03)00278-7

109. Perrault, A.A., Khani, A., Quairiaux, C., Kompotis, K., Franken, P., Muhlethaler, M., Schwartz, S., Bayer, L., 2019. Whole-Night Continuous Rocking Entrains Spontaneous Neural Oscillations with Benefits for Sleep and Memory. Current Biology 29, 402-411.e3. https://doi.org/10.1016/j.cub.2018.12.028

110. Petzka, M., Chatburn, A., Charest, I., Balanos, G.M., Staresina Correspondence, B.P., Staresina, B.P., 2022. Sleep spindles track cortical learning patterns for memory consolidation. Current Biology 0. https://doi.org/10.1016/J.CUB.2022.04.045

111. Peyrache, A., Seibt, J., 2020. A mechanism for learning with sleep spindles. Philosophical Transactions of the Royal Society B: Biological Sciences 375, 22–24. https://doi.org/10.1098/rstb.2019.0230

112. Piantoni, G., Halgren, E., Cash, S.S., 2017. Spatiotemporal characteristics of sleep spindles depend on cortical location. Neuroimage 146, 236–245. https://doi.org/10.1016/J.NEUROIMAGE.2016.11.010

113. Piantoni, G., Halgren, E., Cash, S.S., 2016. The contribution of thalamocortical core and matrix pathways to sleep spindles. Neural Plast 2016. https://doi.org/10.1155/2016/3024342

114. Piantoni, G., Poil, S.S., Linkenkaer-Hansen, K., Verweij, I.M., Ramautar, J.R., Van Someren, E.J.W., Van Der Werf, Y.D., 2013. Individual differences in white matter diffusion affect sleep oscillations. Journal of Neuroscience 33, 227–233. https://doi.org/10.1523/JNEUROSCI.2030-12.2013

115. Pöhlchen, D., Pawlizki, A., Gais, S., Schönauer, M., 2021. Evidence against a large effect of sleep in protecting verbal memories from interference. J Sleep Res 30, e13042. https://doi.org/10.1111/JSR.13042

116. Purcell, S.M., Manoach, D.S., Demanuele, C., Cade, B.E., Mariani, S., Cox, R., Panagiotaropoulou, G., Saxena, R., Pan, J.Q., Smoller, J.W., Redline, S., Stickgold, R., 2017. Characterizing sleep spindles in 11,630 individuals from the National Sleep Research Resource. Nat Commun 8. https://doi.org/10.1038/ncomms15930

117. Qin, S., Leong, R.L.F., Ong, J.L., Chee, M.W.L., 2022. Associations between objectively measured sleep parameters and cognition in healthy older adults: A meta-analysis. Sleep Med Rev 67, 101734. https://doi.org/10.1016/J.SMRV.2022.101734

118. Quintana, D.S., 2015. From pre-registration to publication: A non-technical primer for conducting a meta-analysis to synthesize correlational data. Front Psychol 6, 1–9. https://doi.org/10.3389/fpsyg.2015.01549

119. Rasch, B., Büchel, C., Gais, S., Born, J., 2007. Odor cues during slow-wave sleep prompt declarative memory consolidation. Science (1979) 315, 1426–1429. https://doi.org/10.1126/science.1138581

120. Reverberi, S., Kohn, N., Fernández, G., 2020. No evidence for an effect of explicit relevance instruction on consolidation of associative memories. Neuropsychologia 143. https://doi.org/10.1016/j.neuropsychologia.2020.107491

121. Rickard, T.C., Pan, S.C., Gupta, M.W., 2022. Severe publication bias contributes to illusory sleep consolidation in the motor sequence learning literature. J Exp Psychol Learn Mem Cogn. https://doi.org/10.1037/XLM0001090

122. Rothschild, G., Eban, E., Frank, L.M., 2017. A cortical-hippocampal-cortical loop of information processing during memory consolidation. Nat Neurosci 20, 251–259. https://doi.org/10.1038/nn.4457

123. Rudoy, J.D., Voss, J.L., Westerberg, C.E., Paller, K.A., 2009. Strengthening individual memories by reactivating them during sleep. Science (1979) 326, 1079. https://doi.org/10.1126/science.1179013

124. Sawangjit, A., Oyanedel, C.N., Niethard, N., Salazar, C., Born, J., Inostroza, M., 2018. The hippocampus is crucial for forming non-hippocampal long-term memory during sleep. Nature 564, 109–113. https://doi.org/10.1038/s41586-018-0716-8

125. Schabus, M., Dang-Vu, T.T., Albouy, G., Balteau, E., Boly, M., Carrier, J., Darsaud, A., Degueldre, C., Desseilles, M., Gais, S., Phillips, C., Rauchs, G., Schnakers, C., Sterpenich, V., Vandewalle, G., Luxen, A., Maquet, P., 2007. Hemodynamic cerebral correlates of sleep spindles during human non-rapid eye movement sleep. Proceedings of the National Academy of Sciences 104, 13164–13169. https://doi.org/10.1073/PNAS.0703084104

126. Schabus, M., Gruber, G., Parapatics, S., Sauter, C., Klösch, G., Anderer, P., Klimesch, W., Saletu, B., Zeitlhofer, J., 2004. Sleep spindles and their significance for declarative memory consolidation. Sleep 27, 1479–1485. https://doi.org/10.1093/sleep/27.7.1479

127. Schabus, M., Hoedlmoser, K., Pecherstorfer, T., Anderer, P., Gruber, G., Parapatics, S., Sauter, C., Kloesch, G., Klimesch, W., Saletu, B., Zeitlhofer, J., 2008. Interindividual sleep spindle differences and their relation to learning-related enhancements. Brain Res 1191, 127–135. https://doi.org/10.1016/j.brainres.2007.10.106

128. Schäfer, S.K., Wirth, B.E., Staginnus, M., Becker, N., Michael, T., Sopp, M.R., 2020. Sleep's impact on emotional recognition memory: A meta-analysis of whole-night, nap, and REM sleep effects. Sleep Med Rev 51, 101280. https://doi.org/10.1016/J.SMRV.2020.101284

129. Schäfer, T., Schwarz, M.A., 2019. The meaningfulness of effect sizes in psychological research: Differences between subdisciplines and the impact of potential biases. Front Psychol 10, 813. https://doi.org/10.3389/FPSYG.2019.00813/BIBTEX

130. Schendan, H.E., Searl, M.M., Melrose, R.J., Stern, C.E., 2003. An fMRI study of the role of the medial temporal lobe in implicit and explicit sequence learning. Neuron 37, 1013–1025. https://doi.org/10.1016/S0896-6273(03)00123-5

131. Schmid, D., Erlacher, D., Klostermann, A., Kredel, R., Hossner, E.J., 2020. Sleep-dependent motor memory consolidation in healthy adults: A meta-analysis. Neurosci Biobehav Rev 118, 270–281. https://doi.org/10.1016/J.NEUBIOREV.2020.07.028

132. Schmidt, C., Peigneux, P., Muto, V., Schenkel, M., Knoblauch, V., Münch, M., De Quervain, D.J.F., Wirz-Justice, A., Cajochen, C., 2006. Encoding difficulty promotes postlearning changes in sleep spindle activity during napping. Journal of Neuroscience 26, 8976–8982. https://doi.org/10.1523/JNEUROSCI.2464-06.2006

133. Schönauer, M., 2018. Sleep Spindles: Timed for Memory Consolidation. Current Biology 28, R656–R658. https://doi.org/10.1016/j.cub.2018.03.046

134. Schönauer, M., Alizadeh, S., Jamalabadi, H., Abraham, A., Pawlizki, A., Gais, S., 2017. Decoding material-specific memory reprocessing during sleep in humans. Nat Commun 8. https://doi.org/10.1038/ncomms15404

135. Schönauer, M., Geisler, T., Gais, S., 2014. Strengthening procedural memories by reactivation in sleep. J Cogn Neurosci 26, 143–153.

136. Schönauer, M., Grätsch, M., Gais, S., 2015. Evidence for two distinct sleep-related long-term memory consolidation processes. Cortex 63, 68–78. https://doi.org/10.1016/j.cortex.2014.08.005

137. Schreiner, T., Petzka, M., Staudigl, T., Staresina, B.P., 2021. Endogenous memory reactivation during sleep in humans is clocked by slow oscillation-spindle complexes. Nat Commun 12. https://doi.org/10.1038/s41467-021-23520-2

138. Seibt, J., Richard, C.J., Sigl-Glöckner, J., Takahashi, N., Kaplan, D.I., Doron, G., De Limoges, D., Bocklisch, C., Larkum, M.E., 2017. Cortical dendritic activity correlates with spindle-rich oscillations during sleep in rodents. Nat Commun 8. https://doi.org/10.1038/s41467-017-00735-w

139. Siapas, A.G., Wilson, M.A., 1998. Coordinated interactions between hippocampal ripples and cortical spindles during slow-wave sleep. Neuron 21, 1123–1128. https://doi.org/10.1016/S0896-6273(00)80629-7

140. Simor, P., Zavecz, Z., Horváth, K., Élteto, N., Török, C., Pesthy, O., Gombos, F., Janacsek, K., Nemeth, D., 2019. Deconstructing procedural memory: Different learning trajectories and consolidation of sequence and statistical learning. Front Psychol 9. https://doi.org/10.3389/fpsyg.2018.02708

141. Sopp, M.R., Michael, T., Mecklinger, A., 2018. Effects of early morning nap sleep on associative memory for neutral and emotional stimuli. Brain Res 1698, 29–42. https://doi.org/10.1016/j.brainres.2018.06.020

142. Staresina, B.P., Bergmann, T.O., Bonnefond, M., Van Der Meij, R., Jensen, O., Deuker, L., Elger, C.E., Axmacher, N., Fell, J., 2015. Hierarchical nesting of slow oscillations, spindles and ripples in the human hippocampus during sleep. Nat Neurosci 18, 1679–1686. https://doi.org/10.1038/nn.4119

143. Steriade, M., 2006. Grouping of brain rhythms in corticothalamic systems. Neuroscience 137, 1087–1106. https://doi.org/10.1016/j.neuroscience.2005.10.029

144. Steriade, M., Timofeev, I., 2003. Neuronal plasticity in thalamocortical networks during sleep and waking oscillations. Neuron 37, 563–576. https://doi.org/10.1016/S0896-6273(03)00065-5

145. Studte, S., Bridger, E., Mecklinger, A., 2015. Nap sleep preserves associative but not item memory performance. Neurobiol Learn Mem 120, 84–93. https://doi.org/10.1016/j.nlm.2015.02.012

146. Sugawara, S.K., Koike, T., Kawamichi, H., Makita, K., Hamano, Y.H., Takahashi, H.K., Nakagawa, E., Sadato, N., 2018. Qualitative differences in offline improvement of procedural memory by daytime napping and overnight sleep: An fMRI study. Neurosci Res 132, 37–45. https://doi.org/10.1016/J.NEURES.2017.09.006

147. Tamaki, M., Matsuoka, T., Nittono, H., Hori, T., 2008. Fast sleep spindle (13-15 Hz) activity correlates with sleep-dependent improvement in visuomotor performance. Sleep 31, 204–211. https://doi.org/10.1093/sleep/31.2.204

148. Tamminen, J., Payne, J.D., Stickgold, R., Wamsley, E.J., Gaskell, M.G., 2010. Sleep spindle activity is associated with the integration of new memories and existing knowledge. Journal of Neuroscience 30, 14356–14360. https://doi.org/10.1523/JNEUROSCI.3028-10.2010

149. Thompson, S.G., Higgins, J.P.T., 2002. How should meta-regression analyses be undertaken and interpreted? Stat Med 21, 1559–1573. https://doi.org/10.1002/sim.1187

150. Thürer, B., Weber, F.D., Born, J., Stein, T., 2018. Variable training but not sleep improves consolidation of motor adaptation. Sci Rep 8. https://doi.org/10.1038/s41598-018-34225-w

151. Timofeev, I., Bazhenov, M., Sejnowski, T.J., Steriade, M., 2001. Contribution of intrinsic and synaptic factors in the desynchronization of thalamic oscillatory activity. Thalamus Relat Syst 1, 53–69. https://doi.org/10.1016/S1472-9288(01)00004-8

152. Timofeev, I., Chauvette, S., 2013. The Spindles: Are They Still Thalamic? Commentary on Ayoub et al. Differential effects on fast and slow spindle activity, and the sleep slow oscillation in humans with carbamazepine and flunarizine to antagonize voltage-dependent Na + and Ca 2+ channel activity. SLEEP 2013;36:905-911. Sleep 36. https://doi.org/10.5665/sleep.2702

153. Timofeev, I., Grenier, F., Bazhenov, M., Houweling, A.R., Sejnowski, T.J., Steriade, M., 2002. Short-and medium-term plasticity associated with augmenting responses in cortical slabs and spindles in intact cortex of cats in vivo. Journal of Physiology 542, 583–598. https://doi.org/10.1113/jphysiol.2001.013479

154. Tipton, E., Pustejovsky, J.E., Ahmadi, H., Elizabeth Tipton, C., Professor, A., 2019. Current practices in meta-regression in psychology, education, and medicine. Wiley Online Library 10, 180–194. https://doi.org/10.1002/jrsm.1339

155. Ujma, P., 2022. Meta-analytic evidence suggests no correlation between sleep spindles and memory. 10.31234/osf.io/hjdex

156. Ujma, P.P., 2018. Sleep spindles and general cognitive ability – A meta-analysis. Sleep Spindles Cortical Up States 2, 1–17. https://doi.org/10.1556/2053.2.2018.01

157. Ujma, P.P., Hajnal, B., Bódizs, R., Gombos, F., Erőss, L., Wittner, L., Halgren, E., Cash, S.S., Ulbert, I., Fabó, D., 2021. The laminar profile of sleep spindles in humans. Neuroimage 226. https://doi.org/10.1016/j.neuroimage.2020.117587

158. Urakami, Y., 2008. Relationships between sleep spindles and activities of cerebral cortex as determined by simultaneous EEG and MEG recording. Journal of Clinical Neurophysiology 25, 13–24. https://doi.org/10.1097/WNP.0B013E318162A8A4

159. van Schalkwijk, F.J., Hauser, T., Hoedlmoser, K., Ameen, M.S., Wilhelm, F.H., Sauter, C., Klösch, G., Moser, D., Gruber, G., Anderer, P., Saletu, B., Parapatics, S., Zeitlhofer, J., Schabus, M., 2020. Procedural memory consolidation is associated with heart rate variability and sleep spindles. J Sleep Res 29, 1–8. https://doi.org/10.1111/jsr.12910

160. van Schalkwijk, F.J., Sauter, C., Hoedlmoser, K., Heib, D.P.J., Klösch, G., Moser, D., Gruber, G., Anderer, P., Zeitlhofer, J., Schabus, M., 2019. The effect of daytime napping and full-night sleep on the consolidation of declarative and procedural information. J Sleep Res 28, e12649. https://doi.org/10.1111/jsr.12649

161. Viechtbauer, W., 2010. Conducting meta-analyses in R with the metafor. J Stat Softw 36, 1–48. https://doi.org/10.18637/jss.v036.i03

162. Viechtbauer, W., Cheung, M.W.-L., 2010. Outlier and influence diagnostics for meta-analysis. Res Synth Methods 1, 112–125. https://doi.org/10.1002/jrsm.11

163. Vyazovskiy, V. V., Riedner, B.A., Cirelli, C., Tononi, G., 2007. Sleep homeostasis and cortical synchronization: II. A local field potential study of sleep slow waves in the rat. Sleep 30, 1631–1642. https://doi.org/10.1093/sleep/30.12.1631

164. Walker, M.P., Brakefield, T., Seidman, J., Morgan, A., Hobson, J.A., Stickgold, R., 2003. Sleep and the time course of motor skill learning. Learning and Memory 10, 275–284. https://doi.org/10.1101/lm.58503

165. Walker, M.P., Stickgold, R., Jolesz, F.A., Yoo, S.S., 2005. The functional anatomy of sleep-dependent visual skill learning. Cerebral Cortex 15, 1666–1675. https://doi.org/10.1093/cercor/bhi043

166. Wang, B., Antony, J.W., Lurie, S., Brooks, P.P., Paller, K.A., Norman, K.A., 2019. Targeted Memory Reactivation during Sleep Elicits Neural Signals Related to Learning Content. J Neurosci 39, 6728–6736. https://doi.org/10.1523/JNEUROSCI.2798-18.2019

167. Ward, M.P., Peters, K.R., Smith, C.T., 2014. Effect of emotional and neutral declarative memory consolidation on sleep architecture. Exp Brain Res 232, 1525–1534. https://doi.org/10.1007/s00221-013-3781-0

168. Werth, E., Achermann, P., Dijk, D.J., Borbély, A.A., 1997. Spindle frequency activity in the sleep EEG: Individual differences and topographic distribution. Electroencephalogr Clin Neurophysiol 103, 535–542. https://doi.org/10.1016/S0013-4694(97)00070-9

169. Wilhelm, I., Diekelmann, S., Molzow, I., Ayoub, A., Mölle, M., Born, J., 2011. Sleep selectively enhances memory expected to be of future relevance. Journal of Neuroscience 31, 1563–1569. https://doi.org/10.1523/JNEUROSCI.3575-10.2011

170. Wilson, M.A., McNaughton, B.L., 1994. Reactivation of hippocampal ensemble memories during sleep. Science (1979) 265, 676–679. https://doi.org/10.1126/science.8036517

171. Zhang, H., Fell, J., Axmacher, N., 2018. Electrophysiological mechanisms of human memory consolidation. Nat Commun 9. https://doi.org/10.1038/s41467-018-06553-y

Other articles which are included in the meta-analysis:

172. Adamczyk, M., Genzel, L., Dresler, M., Steiger, A., Friess, E. (2015). Automatic sleep spindle detection and genetic influence estimation using continuous wavelet transform. Frontiers in Human Neuroscience, Vol. 9. https://doi.org/10.3389/fnhum.2015.00624

173. Albouy, G., Fogel, S., Pottiez, H., Nguyen, V. A., Ray, L., Lungu, O., … Doyon, J. (2013). Daytime Sleep Enhances Consolidation of the Spatial but Not Motoric Representation of Motor Sequence Memory. PLoS ONE, 8(1), 11–13. https://doi.org/10.1371/journal.pone.0052805

174. Alger, S. E., Chen, S., Payne, J. D. (2019). Do Different Salience Cues Compete for Dominance in Memory over a Daytime Nap? https://doi.org/10.1016/j.nlm.2018.06.005

175. Cairney, S. A., Durrant, S. J., Jackson, R., Lewis, P. A. (2014). Sleep spindles provide indirect support to the consolidation of emotional encoding contexts. Neuropsychologia, 63, 285–292. https://doi.org/10.1016/j.neuropsychologia.2014.09.016

176. Casey, S. J., Solomons, L. C., Steier, J., Kabra, N., Burnside, A., Pengo, M. F., … Kopelman, M. D. (2016). Slow wave and rem sleep deprivation effects on explicit and implicit memory during sleep. Neuropsychology, 30(8), 931–945. https://doi.org/10.1037/neu0000314

177. Fogel, S. M., Ray, L. B., Binnie, L., Owen, A. M. (2015). How to become an expert: A new perspective on the role of sleep in the mastery of procedural skills. Neurobiology of Learning and Memory, 125, 236–248. https://doi.org/10.1016/j.nlm.2015.10.004

178. Göder, R., Graf, A., Ballhausen, F., Weinhold, S., Baier, P. C., Junghanns, K., Prehn-Kristensen, A. (2015). Impairment of sleep-related memory consolidation in schizophrenia: Relevance of sleep spindles? Sleep Medicine, 16(5), 564–569. https://doi.org/10.1016/j.sleep.2014.12.022

179. Hanert, A., Weber, F. D., Pedersen, A., Born, J., Bartsch, T. (2017). Sleep in humans stabilizes pattern separation performance. Journal of Neuroscience, 37(50), 12238–12246. https://doi.org/10.1523/JNEUROSCI.1189-17.2017

180. Heib, D. P. J., Gruber, G., Hoedlmoser, K., Zeitlhofer, J. (2015). Oscillatory Theta Activity during Memory Formation and Its Impact on Overnight Consolidation: A Missing Link? Washington, DC: APA, Guideline Development Panel for the Treatment of Posttraumatic Stress Disorder in Adults., (March), 139. https://doi.org/10.1162/jocn

181. Hennies, N., Ralph, M. A. L., Kempkes, M., Cousins, J. N., Lewis, P. A. (2016). Sleep spindle density predicts the effect of prior knowledge on memory consolidation. Journal of Neuroscience, 36(13), 3799–3810. https://doi.org/10.1523/JNEUROSCI.3162-15.2016

182. Rodríguez-Labrada, R., Galicia-Polo, L., Canales-Ochoa, N., Voss, U., Tuin, I., Peña-Acosta, A., … Velázquez-Pérez, L. (2019). Sleep spindles and K-complex activities are decreased in spinocerebellar ataxia type 2: relationship to memory and motor performances. Sleep Medicine, 60, 188–196. https://doi.org/10.1016/j.sleep.2019.04.005

183. Ruch, S., Markes, O., Duss, S. B., Oppliger, D., Reber, T. P., Koenig, T., … Henke, K. (2012). Sleep stage II contributes to the consolidation of declarative memories. Neuropsychologia, 50(10), 2389–2396. https://doi.org/10.1016/j.neuropsychologia.2012.06.008

184. Saletin, J. M., Goldstein, A. N., Walker, M. P. (2011). The role of sleep in directed forgetting and remembering of human memories. Cerebral Cortex, 21(11), 2534–2541. https://doi.org/10.1093/cercor/bhr034

185. Sattari, N., McDevitt, E. A., Panas, D., Niknazar, M., Ahmadi, M., Naji, M., … Mednick, S. C. (2017). The effect of sex and menstrual phase on memory formation during a nap. Neurobiology of Learning and Memory, 145(June), 119–128. https://doi.org/10.1016/j.nlm.2017.09.007

186. Shaw, J. J., Monaghan, P. (2017). Lateralised sleep spindles relate to false memory generation. Neuropsychologia, 107(October), 60–67. https://doi.org/10.1016/j.neuropsychologia.2017.11.002

187. Sopp, M. R., Michael, T., Weeß, H. G., Mecklinger, A. (2017). Remembering specific features of emotional events across time: The role of REM sleep and prefrontal theta oscillations. Cognitive, Affective and Behavioral Neuroscience, 17(6), 1186–1209. https://doi.org/10.3758/s13415-017-0542-8

188. Tucker, M. A., Fishbein, W. (2009). The impact of sleep duration and subject intelligence on declarative and motor memory performance: How much is enough? Journal of Sleep Research, 18(3), 304–312. https://doi.org/10.1111/j.1365-2869.2009.00740.x

189. van der Helm, E., Gujar, N., Nishida, M., Walker, M. P. (2011). Sleep-Dependent Facilitation of Episodic Memory Details. PLoS ONE, 6(11). https://doi.org/10.1371/journal.pone.0027421

190. Weber, F. D., Wang, J. Y., Born, J., Inostroza, M. (2014). Sleep benefits in parallel implicit and explicit measures of episodic memory. Learning and Memory, 21(4), 190–198. https://doi.org/10.1101/lm.033530.113

191. Wilhelm, I., Diekelmann, S., Molzow, I., Ayoub, A., Mölle, M., Born, J. (2011). Sleep selectively enhances memory expected to be of future relevance. Journal of Neuroscience, 31(5), 1563–1569. https://doi.org/10.1523/JNEUROSCI.3575-10.2011

11. Supplementary Material

Supplementary Figure 1. Forest plot for the (r) between levels of sleep spindles and long-term memory, reflecting memory consolidation (k = 1204). Correlations are shown for all effects in the meta-analysis. The size of each dot reflects the weight given to the observed effect during model fitting. The diamond at the bottom shows the meta-analytically weighted mean correlation (with 95% CI, z = 0.240, r = 0.236). The dotted vertical line represents an effect size of 0. Multiple measures were adjusted for dependency using multilevel random effect models (See: **Method section**).

Supplementary Figure 2. Funnel plot showing the standard errors of effect sizes between sleep spindles and memory consolidation across all studies with 1204 unique associations. The key areas of statistical significance have been superimposed on the funnel plot. While the red zones show effects between $p = 0.1$ and $p = 0.05$, and the orange zones show effects between $p = 0.05$ and $p = 0.01$.

Supplementary Figure 3. The scatter plot shows the relationship between sample size and effect size. Pearson's correlation across all studies, reporting memory consolidation resulted in a significant association (r = -0.175, $p < 0.001$, k = 1204) suggesting a clear indication of a publication bias taking the form that smaller effects would result from larger samples and vice versa. Note that to increase the interpretability of the graph, we visualized the correlation plot by excluding one study (Ackermann et al., 2015) with a higher sample size (N = 929 and r = -0.377, $p < 0.001$)).

Supplementary Table 1. The number of estimates for each measure of interests across extracted 53 studies with 1424 effect sizes.

Variables	N
total estimates	1427
learning ability (trait)	223
memory consolidation (state)	1204
memory retention	592
post-sleep	612
Memory types	
declarative memory	939
procedural memory	488
hippocampal tasks	216
non-hippocampal tasks	272
Spindle features	
fast	610
slow	348
global	469
EEG scalp topography	
frontal	428
central	485
parietal	275
others (e.g., temporal, occipital)	239
Spindle measures	
amplitude	164
count	340
density	372
duration	172
frequency	36
power	207
spindle activity	136
Sleep protocol	
sleep	779
nap	580
others (e.g., half-night, cycle)	68
Effect size status	
Reported	984
Unreported	443

Supplementary Table 2. Post-hoc comparisons for the moderation analysis of spindle measures on spindle-memory associations.
Note that p-values are adjusted according to Bonferroni correction.
*** < 0.001, ** < 0.01, *< 0.05, . < 0.1

comparisons	Estimate (difference)	p_{adj}	
count-amplitude	-0.12	0.225	
density-amplitude	-0.052	1	
duration-amplitude	-0.118	0.0145	*
frequency-amplitude	0.344	<0.001	***
power-amplitude	0.05	1	
SpA-amplitude	-0.133	1	
density-count	0.068	1	
duration-count	0.001	1	
frequency-count	0.464	<0.001	***
power-count	0.169	0.961	
SpA-count	-0.014	1	
duration-density	-0.067	0.659	
frequency-density	0.396	<0.001	***
power-density	0.102	1	
SpA-density	-0.081	1	
frequency-duration	0.463	<0.001	***
power-duration	0.168	0.827	
SpA-duration	-0.015	1	
power-frequency	-0.295	0.06	.
SpA-frequency	-0.478	<0.001	***
SpA-power	-0.183	1	

Supplementary Table 3. Post-hoc comparisons for the moderation analysis of spindle measures on spindle-memory associations after excluding 2 studies reporting frequency measure (N = 27).
Note that p-values are adjusted according to Bonferroni correction. *** < 0.001, ** < 0.01, *< 0.05, . < 0.1

comparisons	Estimate	p_{adj}
count-amplitude	-0.109	0.2964
density-amplitude	-0.046	1
duration-amplitude	-0.111	0.0225 *
power-amplitude	0.055	1
SpA-amplitude	-0.128	0.9742
density-count	0.063	1
duration-count	-0.002	1
power-count	0.164	0.7887
SpA-count	-0.019	1
duration-density	-0.065	0.5365
power-density	0.101	1
SpA-density	-0.082	1
power-duration	0.166	0.6319
SpA-duration	-0.017	1
SpA-power	-0.183	0.9867