Reduced parasite burden in feral honeybee colonies

Patrick L. Kohl[1,*], Paul D'Alvise[2,§], Benjamin Rutschmann[1,3], Sebastian Roth[3], Felix Remter[3,4], Ingolf Steffan-Dewenter[1] & Martin Hasselmann[2]

[1] Department of Animal Ecology and Tropical Biology, Biocenter, University of Würzburg, Würzburg, Germany

[2] Department of Livestock Population Genomics, Institute of Animal Science, University of Hohenheim, Stuttgart, Germany

[3] BEEtree-Monitor, Munich, Germany

[4] Katholische Stiftungshochschule München

[§] Current address: Institute for Medical Microbiology and Hygiene, University Hospital Tübingen, Tübingen, Germany

[*] Correspondence; email: patrick.kohl@uni-wuerzburg.de

Abstract

Bee parasites are the main threat to apiculture, and since many parasite taxa can spill over from honeybees (*Apis mellifera*) to other bee species, honeybee disease management is important for pollinator conservation in general. It is unknown whether honeybees that escaped from apiaries (i.e., feral colonies) benefit from natural parasite-reducing mechanisms like swarming or suffer from high parasite pressure due to the lack of medical treatment. In the latter case, they could function as parasite reservoirs and pose a risk to the health of managed honeybees (spillback) and wild bees (spillover). We compared the occurrence of 18 microparasites among managed (N=74) and feral (N=64) honeybee colony samples from four regions in Germany using qPCR. We distinguished five colony types representing differences in colony age and management histories, two variables potentially modulating parasite

prevalence. Besides strong regional variation in parasite communities, parasite burden was consistently lower in feral than in managed colonies. The overall number of detected parasite taxa per colony was lower, and Trypanosomatidae, chronic bee paralysis virus, and deformed wing viruses A and B were less prevalent and abundant in feral colonies than in managed colonies. Parasite burden was lowest in newly founded feral colonies, intermediate in overwintered feral colonies and managed nucleus colonies, and highest in overwintered managed colonies and hived swarms. Our study confirms the hypothesis that the natural mode of colony reproduction and dispersal by swarming temporally reduces parasite pressure in honeybees. We conclude that feral colonies are unlikely to contribute significantly to the spread of bee diseases.

Keywords

Bee viruses, disease management, horizontal transmission, livestock-wildlife interactions, pathogen spillover, swarming, wild honeybees, *Varroa destructor*.

Introduction

Disease transfer between managed and wild animals is a potential source of conflict between the livestock sector and nature conservation (Mysterud & Rolandsen, 2019). Populations managed in intensive animal husbandry can be vulnerable to diseases transmitted from wild populations because high animal densities and low genetic variance increase the risk of epidemics (Gortázar et al., 2007). Conversely, diseases of livestock can become a threat to wildlife, e.g., when managed species introduce diseases into non-native areas (Ayala et al., 2020; Costanzi et al., 2021; Schommer & Woolever, 2008). An interesting example of potentially negative livestock-wildlife interactions is that of the disease associations between managed western honeybees (*Apis mellifera*), wild-living honeybees, and other bee species (Fürst et al., 2014; Ravoet et al., 2014). The Western honeybee is native to Africa, Europe,

and Western Asia, and is an important pollinator of wild plants both in its native and

introduced range (Dick, 2001; Hung et al., 2018). The species is managed by beekeepers

globally to pollinate the flowers of crops and to produce honey and other hive products

(Crane, 1990; Rollin & Garibaldi, 2019). Where self-sustaining wild populations have gone

extinct, as in many regions of Europe and Western Asia (Pirk et al., 2017; Requier et al.,

2019), apiculture also serves to protect the honeybee as a species. However, modern

rationalized beekeeping can conflict with conservation (Geldmann & González-Varo, 2018;

Iwasaki & Hogendoorn, 2022; Panziera et al., 2022). By preventing swarming and

maintaining unnaturally large, continuously breeding colonies; by crowding hives in apiaries,

and by seasonally moving colonies between bee yards, apiculture promotes the reproduction

and spread of bee parasites (Brosi et al., 2017; Loftus et al., 2016; Martínez-López et al.,

2022; Nolan & Delaplane, 2017; Peck & Seeley, 2019; Seeley & Smith, 2015). On a global

scale, the transport of hives and their products can expose the bees to entirely novel parasites

(Goulson & Hughes, 2015). This was famously demonstrated by the worldwide invasions of

western honeybee populations by the ectoparasitic mite *Varroa destructor,* whose natural host

was the eastern honeybee *A. cerana* (Kraus & Page, 1995; Traynor et al., 2020; Wilfert et al.,

2016). Unfortunately, many bee parasites are not restricted to a single species. Since

honeybees inevitably share floral resources with other pollinators, their parasites can spill

over to, and harm, populations of other bee species (Burnham et al., 2021; Fürst et al., 2014;

Piot et al., 2022; Tehel et al., 2022). Therefore, managing honeybee diseases is important for

the conservation of bee pollination services in general, both in the contexts of crop production

and ecosystem functioning (Bartlett, 2022; Brosi et al., 2017).

Honeybees are social insects that live in large perennial nests, and as such, they are naturally

attractive hosts for a range of parasites including arthropods, fungi, bacteria, and viruses

(Schmid-Hempel, 1998). However, parasites are not considered to be a limiting factor for

wild honeybee populations under natural conditions (Bailey, 1958; Fries et al., 2006; Fries &

Camazine, 2001; Ratnieks & Nowakowski, 1989). An important reason seems to be the

regulation of parasite populations within colonies as a side-effect of the bees' natural cycle of

colony reproduction and dispersal (DeGrandi-Hoffman et al., 2017; Loftus et al., 2016).

Honeybee colonies reproduce via fission when the old queen and approximately 70% of the

workers leave as a swarm to build a new nest in another cavity (Seeley, 2010). The swarming

bees do not transfer brood to the new nesting site, and it takes around three weeks until a

young queen resumes egg-laying in the old nest; hence, brood production is interrupted in

both colony parts and the reproduction of parasites infecting the brood or young workers is

halted (Loftus et al., 2016; Royce et al., 1991). Furthermore, the rate at which new parasite

species enter wild colonies is most probably lower compared to the situation at apiaries, since

their nests are typically widely dispersed in the landscape (Lindström et al., 2008; Nolan &

Delaplane, 2017; Seeley & Smith, 2015). Besides these ecological factors, wild honeybee

populations are more resilient against disease than managed populations because they are

more likely to evolve defences against parasites via natural selection (Neumann & Blacquière,

2017; Pirk et al., 2017). A famous example is a population of non-native wild honeybees,

inhabiting forests in the species' introduced range in the north-eastern USA, which was

subjected to rapid evolution upon the arrival of *Varroa destructor* in the 1980s and remains

viable (Mikheyev et al., 2016; Seeley, 2017). Not only do wild honeybee populations deliver

free pollination services (Chang & Hoopingarner, 1991), but they can also benefit the

beekeeping sector when genetic adaptations to (novel) stressors, including parasites, are

passed on to the managed population (Pirk et al., 2017). Self-sustaining wild honeybee

populations exist both in the species' native range in Africa (Dietemann et al., 2009) and in its

introduced range in Australia (Oldroyd et al., 1997), and parts of the Americas (Seeley, 2007;

Winston, 1992).

The situation might differ for wild-living honeybee colonies which are recent escapees from

apiaries. For example, in Germany, despite active swarm prevention by beekeepers, thousands

of honeybee swarms emigrate from bee yards each spring and colonize tree cavities in

managed forests. However, their annual survival rate is far below the threshold required to

maintain a self-sustaining population (Kohl et al., 2022). We here refer to such honeybees as

"feral" (Daniels & Bekoff, 1989), as opposed to "wild", regardless of whether the native or

introduced range of the species is considered. Feral colonies exist wherever apiculture is

practised, and they are probably also widespread in Europe (Browne et al., 2020; Dubaić et

al., 2021; Kohl et al., 2022; Kohl & Rutschmann, 2018; Oleksa et al., 2013; Rutschmann et

al., 2022; Thompson, 2012). On the one hand, feral colonies might have a lower parasite

burden than managed colonies due to the natural parasite-reducing effects of swarming and

the dispersal of nest sites. On the other hand, with an evolutionary background of artificial

selection and a history of care by beekeepers, feral colonies are likely to be stressed by

multiple environmental factors and might develop high parasite loads without medical

treatment (Thompson et al., 2014).

Understanding the disease ecology of recently feralised honeybee colonies is important

because they create a potential management conflict. Honeybees are usually considered

livestock animals and beekeepers are obligated to register their hives, regularly apply miticide

treatments, and report infectious diseases to prevent epidemics. If feral colonies carry high

loads of parasites, however, they might reinfect managed colonies (spillback), undermining

the veterinary measures undertaken to combat disease (Frey & Rosenkranz, 2014; Thompson

et al., 2014). When highly infected feral colonies disperse into natural areas, they might also

function as vectors for parasites that can spill over to non-*Apis* wild bees (Burnham et al.,

2021; Fürst et al., 2014; Piot et al., 2022; Tehel et al., 2022). This situation would suggest

management options aiming at preventing feralisation or eradicating feral colonies (Taylor et

al., 2007). Conversely, where the Western honeybee is a native species, promoting or re-

establishing populations of wild-living honeybees is a legitimate conservation goal (Panziera

et al., 2022; Requier et al., 2019). It would then be inappropriate to combat feral colonies

since they can be the source of future wild populations.

To assess whether feral honeybees might pose a risk to the health of managed and wild bee

populations, we compared the occurrence of 18 microparasite taxa in feral and managed

honeybee colonies using qPCR (D'Alvise et al., 2019). We collected colony samples from

four regions in southern Germany encompassing both rural and urban landscapes, to cover

cases from different environments with potentially different parasite communities

(Youngsteadt et al., 2015). By taking all samples within a four-week time window in July, we

made sure that variation in parasite communities could not be attributable to seasonal

variation (D'Alvise et al., 2019; Faurot-Daniels et al., 2020). We chose this point in time

because it is epidemiologically relevant: a high frequency of between-colony robbing

behaviour due to nectar scarcity in summer (Garbuzov et al., 2020) increases the risks of

parasites transmissions, and parasite loads in summer affect subsequent colony winter

survival (Ravoet et al., 2013). In addition to the difference between managed and feral, we

distinguished five colony types representing different colony age classes and management

histories since these factors potentially modulate parasite prevalence. The collected data

allowed us to make a nuanced assessment of the effect of environmental differences between

managed and feral honeybee colonies on parasite prevalence.

Material and Methods

Sample collection

We collected worker bees from managed colonies (N=74, from 73 apiaries) and feral colonies

(N=64, from 63 sites) in four regions in southern Germany (Swabian Alb, counties Coburg

and Lichtenfels, county Weilheim-Schongau, and the city of Munich; Fig. 1) in July 2020. In

the first three study regions, feral colonies were found during a population-monitoring study

that was based on systematic censuses of tree cavities made by the black woodpecker, which

are the primary nesting sites of honeybee colonies in managed forests in Germany (Kohl et

al., 2022; Kohl & Rutschmann, 2018). We sampled from all detected feral colonies whose

recent history was known to us (see next paragraph). In the fourth study region (Munich), we

were able to select 15 out of approx. 80 feral honeybee nest sites that had previously been

mapped by two of us using a combination of private search and citizen reports collected via

the BEEtree-Monitor network (S. Roth and F. Remter, unpublished). We selected the 15 nest

sites for sampling based on the criteria of accessibility and knowledge of the recent

occupation history, and to assure an even distribution over the city and an equal representation

of nests in trees and buildings. To obtain locations of managed colonies, we contacted

beekeepers via the local beekeeping organizations and asked for permission to sample bees

from their hives. This resulted in a sample of managed colonies that roughly matched the

sample of feral colonies in terms of size and spatial distribution (Fig. 1). The sampled feral

colonies nested in tree cavities or building walls with entrance heights between 2–18 m above

the ground. Except for four of the 64 colonies, which had neighbouring colonies in the same

tree or building wall, feral colonies were spatially separated from others. The sampled

managed colonies were kept in hives and were usually placed next to other colonies in

apiaries. The median number of colonies at the apiaries was 6 (range: 1–28 colonies), which

well represents the typical apiary size of German beekeepers, who mostly practice beekeeping

as a hobby or sideline occupation (Deutscher Imkerbund e.V., 2020).

A factor potentially affecting parasite burden is colony age. Therefore, we noted for each

colony whether it had overwintered at least once (age > 1 year) or had been newly founded in

the year of sampling (age < 1 year). Amongst feral colonies 72% (N=46) were recent founders

(swarms that moved into the cavity and founded a colony in spring), and 28% (N=18) had

overwintered at least once. Amongst managed colonies we sampled similar numbers of

overwintered (N=38; 51%) and newly founded colonies (N=36; 49%), since we assumed the

ratio of old versus young colonies at apiaries in summer is about 50:50. In the group of young

managed colonies, we further distinguished between nucleus colonies, i.e., daughter colonies

created by taking brood frames and bees from an established colony (N=27) and hived

swarms (N=9). The latter were either natural swarms captured by beekeepers (N=7) or man-

made swarms ("packages") created by brushing bees off the combs of a mother colony and

transferring them into a new hive together with a queen (N=2).

Al least 20 bees per colony were captured by placing a white butterfly net in front of the hive

or cavity entrance (see supplementary information in (Kohl et al., 2022) for details of the

sampling method). With this method, we primarily sampled foragers on their outbound

flights. The sampled bees did not show any overt disease symptoms. After capture, they were

directly freeze-killed and collected in 50 mL vials. They were permanently kept on dry ice

during transport and stored in freezers at -80°C until the analyses.

Figure 1. Geographic locations of the sampled managed (N = 74) and feral (N = 64) honeybee colonies in four regions in southern Germany (1: Swabian Alb, 2: Counties Coburg and Lichtenfels, 3: County Weilheim-Schongau, 4: City of Munich). Note that the map of Munich has a different scale. Land cover data are from (Weigand et al., 2020).

Parasite testing

We assessed the colony-level occurrence of 18 microparasites using high-throughput qPCR on a Biomark HD system (Standard BioTools, San Francisco, CA) with parasite-specific primers (Budge et al., 2010; Cepero et al., 2015; D'Alvise et al., 2019; Evans, 2006; Locke et al., 2012; Lourenço et al., 2008; Martínez et al., 2010; Papp et al., 2014) closely following an established protocol (D'Alvise et al., 2019), with the exception that RNA was extracted from homogenates of 20 workers per colony, not from individual bees (see supplementary information for details on RNA extraction and Table S1 for a list of the parasites and reference genes assayed). We ran three qPCR replicates for each assay and considered a parasite taxon as present in a colony if target molecules were detected in at least two. If a colony sample was positive, we averaged the number of target molecules detected in all three

203 assay replicates to obtain a mean number of target molecules per colony sample. As a

204 measure of colony-level parasite abundance, we calculated for each parasite the number of

205 detected target molecules per 100 ng of extracted RNA and reported the logarithm of this

206 ratio, $\log_{10}(n/100 \text{ ng RNA}+1)$, as the "parasite load".

Statistical analyses

All statistical analyses were performed with R version 4.0.5 (R Core Team, 2022) and figures

were created using "ggplot2" (Wickham, 2016). We analysed whether the number of detected

parasite taxa per colony (univariate analysis) and the microparasite community composition

(multivariate analysis) differed depending on "management" (two levels: "managed" versus

"feral") and "colony type" (five levels: "managed: overwintered", "managed: nucleus

colonies", "managed: hived swarms", "feral: overwintered" and "feral: founders"). In all four

analyses, we accounted for the potential effect of spatial location on microparasite

communities.

We analysed the number of parasite taxa detected (count data) using generalized linear

models (function "glmmTMB"; (Brooks et al., 2017, 2019)). Besides the main factors of

interest, "management" or "colony" type", we included "region" (four levels) as well as the

interaction between "region" and "management" or "colony type" as additional predictors to

account for spatial differences and for potential variation in the effect of management or

colony type between regions. Considering "region" to describe the spatial component was

reasonable since most spatial variation in parasite assemblages was attributable to variation

between regions rather than variation between sites within regions (see below). Count data

can be analysed with a range of model types assuming different probability distributions

(Brooks et al., 2017). We therefore first created five models with the same predictor formula

but different family functions in "glmmTMB" (family = "poisson", "nbinom1", "nbinom2",

"compois" or "genpois") and selected the best models using the Akaike information criterion

for small sample sizes (AICc, function "AICctab" from the "bbmle" package (Bolker, 2022)).

In both comparisons (factor "management" or "colony type") models with a generalized

Poisson distribution (family= "genpois") had the lowest AICc (ΔAICc $\geq$ 12) so we used these

for the analyses. We then tested for significant deviations from model assumptions using the

functions "simulateResiduals", "testResiduals" and "testCategorical" from the "DHARMa"

package (Hartig, 2022). No significant deviations were found in tests for uniformity,

dispersion, and outliers, but Levene's test detected significant differences in variance of

parasite counts between the four study regions. This problem could be fixed by adding a

formula for dispersion with "region" as the single fixed effect. The final formulations of the

full models were "glmmTMB(*Number of parasites* ~ X + region + X : region, dispformula =

~ region, family = genpois)" with "X" denoting either "management" or "colony type".

Predictions of the mean number of parasite taxa and 95% confidence limits were produced

using the function "emmeans" (Lenth, 2022). To test whether management and colony type or

their interactions with region significantly affected parasite counts, we compared pairs of

nested models (where one model contained, and the other missed, the predictor of interest)

using likelihood ratio tests (LRT, function "anova"). We assessed the effect of management or

colony type while accounting for the effect of region (see supplementary information, Tables

S2–S4, S11 and S12 for the specifications of the nested models). In the case of the five-level

factor "colony type", we used the function "glht" from the "multcomp" package (Hothorn et

al., 2008) for Tukey post hoc tests of pair-wise differences in parasite numbers between

colony types.

To analyse differences in parasite community composition, we used distance-based

redundancy analyses with "management" or "colony type" as the constraining factor (dbRDA,

function "db.rda" from the "vegan" package, (Oksanen et al., 2022)). Redundancy analysis

summarizes multi-factor variation so that dissimilarities between samples can be graphically

displayed in a two-dimensional coordinate system whose axes best separate the data based on

predefined factors of interest ("constraints"). We chose to analyse parasite community

compositions based on Jaccard distances (and thus based on presence/absence of parasite

taxa) as opposed to Euclidean distance or Bray-Curtis dissimilarities (based on colony-level

parasite abundance), because abundance data contained many zeroes and were non-normally

distributed, and because it is hard to compare target molecule abundance variation of several

orders of magnitude between parasites as different as arthropods and viruses. To partial out

spatial structure, we conditioned our dbRDAs on principal coordinates of neighbour matrices

(function "pcnm" from the "vegan" package; (Borcard & Legendre, 2002; Oksanen et al.,

2022)). The model formulations were "db.rda(*Jaccard distance matrix of parasite*

communities ~ X + Condition(scores(pcnm(*distance matrix of sample locations*))))", with "X"

denoting either "management" or "colony type". This revealed that spatial location explained

17.7 % of the variation in parasite communities. Interestingly, a test with "study region" (four

levels) as a constraining factor (formulation: "db.rda(*Jaccard distance matrix of parasite*

communities ~ region)") showed that 14.9 % of the variation could be attributed to

differences between the four regions (see Fig. S1). This means that the spatial structure in

parasite communities was mostly caused by large-scale (between regions) rather than fine-

scale (between sites within one region) spatial variation. To infer the statistical significance of

the constraining factors we used permutation tests with 99999 permutations ("anova.cca"

function).

We did not perform separate statistical tests for each microparasite since potential interactions

between taxa lead to non-independent data, and since the high number of individual tests

would introduce statistical problems related to multiple testing. However, we graphically

compared for each tested microparasite the prevalence and the associated 95% confidence

intervals (Blaker, 2000; Stevenson et al., 2022) as well as the mean colony-level loads.

Results

The number of microparasite taxa detected per colony ranged between 1 and 9, with an overall average of 5.8 taxa. Parasite counts differed significantly between regions (LRT: D.f.=6, χ^2= 26.819, P=0.00016, Table S2); they were lowest in Coburg and Lichtenfels (mean: 4.9), intermediate on the Swabian Alb (mean: 5.7), and highest in Weilheim-Schongau (mean: 6.3) and Munich (mean: 6.2). On top of regional differences, management significantly affected the number of parasites per colony (LRT: D.f.=1, χ^2= 14.677, P=0.00013, Table S3). Feral colonies had, on average, one parasite less (median: 5, mean: 5.4, range 1–8) than managed colonies (median: 6, mean: 6.2, range 4–9) (Fig. 2a). The difference between feral and managed colonies was consistent across study regions (no significant interaction between management and region; LRT: D.f.=3, χ^2= 0.947, P=0.814, Table S4).

The distance-based redundancy analysis revealed a marginal difference in the composition of parasite taxa between managed and feral colonies (permutational anova: P=0.074), with management explaining 1.1% of the variation in microparasite communities (Fig. 2b). A direct inspection of each of the 18 microparasites showed that prevalence was either similar in managed and feral colonies or lower in feral colonies (Fig. 3). A clear reduction in prevalence of more than 10 percentage points was found in four microparasite taxa: *Crithidia/Lotmaria* (Managed: 90.5%, Feral: 78.1%), Chronic bee paralysis virus (Managed: 14.9%, Feral: 1.6%), Deformed wing virus A (Managed: 18.9%, Feral: 3.1%), and Deformed wing virus B (Managed: 47.3%, Feral: 35.9%). Comparing mean colony-level parasite loads yielded a similar result, with abundances tending to be lower in feral colonies (Fig. 4a). Considering only colonies which were tested positive showed that parasite loads of infected feral and managed colonies were generally similar (Fig. 4b).

Figure 2. Comparison of parasite burden between managed (N=74) and feral (N=64) honeybee colonies. **(a)** Number of microparasite taxa detected among the 18 taxa assayed for each of the four study regions. Dots are raw data; diamond symbols and vertical lines give model-estimated means and 95% confidence intervals. See Table S5 for an overview of model predictions. **(b)** Representation of dissimilarities in microparasite communities as created by a distance-based redundancy analysis with management as the constraining factor (spatial structure partialled out). Managed and feral colonies are separated along the dbRDA1-axis (1.1% of total variation). The first unconstrained axis (MDS1) explains 16.7% of the variation. Dots represent locations of individual colonies and diamonds are mean locations.

Figure 3. Prevalence (percentage of colonies tested positive; bars) and 95% confidence intervals (black lines) for 18 microparasite taxa among managed (N=74) and feral (N=64) honeybee colonies. See Table S6 for an overview of prevalence values and Tables S7–S10 for overviews of parasite prevalence divided by study region.

Figure 4. Parasite loads of managed and feral honeybee colonies. Only microparasite taxa detected at least once in each group are shown. **(a)** Parasite loads of individual colonies (dots; random jitter is added to reduce overplotting) and mean parasite loads (diamonds) based on all colonies tested (managed: N=74, feral: N=64). **(b)** Parasite loads based on positive cases only. Dots are parasite loads of individual colonies, diamonds are means, and numbers denote sample sizes. See Table S6 for an overview of parasite load values and Tables S7–S10 for overviews divided by study region.

An analysis of the number of parasites in relation to five colony types gave a more nuanced picture of differences in parasite burden (LRT, factor "colony type" after "region": D.f.: 4, χ^2=23.23, P=0.0001, Table S11). Parasite counts were lowest in newly founded feral colonies (mean: 5.3, range: 1–7), intermediate in overwintered feral colonies (mean: 5.7, range: 4–8) and nucleus colonies (mean: 5.7, range: 4–8), and highest in hived swarms (mean: 6.6, range: 4–8) and overwintered managed colonies (mean: 6.3, range: 4–9) (Fig. 5a). These differences were largely consistent across study regions (no significant interaction between colony type and region; LRT: D.f.: 12, χ^2=9.546, P=0.656, Table S12; see Figure S2 and Table S13 for

parasite counts divided by colony type and study region). A pairwise comparison revealed

that the difference between feral founders and overwintered managed colonies (P<0.001) and

between feral founders and hived swarms (P=0.002) were statistically significant. There were

also differences in the colony-level composition of parasite taxa between the five colony types

(colony type explained 3.3% of parasite community variation according to dbRDA;

permutational anova: P=0.099; Fig. 5b). The differences in parasite communities, albeit

marginal, resembled the differences in parasite counts: along the first dbRDA-axis, parasite

assemblages of feral founders were separated from those of hived swarms and overwintered

managed colonies, while the parasite communities of overwintered feral colonies took an

intermediate position, and the parasite communities of managed nucleus colonies were closer

to those of feral colonies than to those of other managed colonies (Fig. 5b). Accordingly,

mean colony-level loads of individual parasite taxa tended to be lower in feral founders

compared to overwintered feral colonies, and among the three types of managed colonies,

mean parasite loads were most often lowest in nucleus colonies (Fig. 5c).

Figure 5. Comparison of parasite burden between different types of managed and feral
honeybee colonies **(a)** Number of microparasite taxa detected among the 18 taxa assayed.
Dots are raw data; large symbols and vertical lines give model-estimated means and 95%

confidence intervals. Pairs that do not share a letter differ significantly (P<0.05). **(b)** Relative
differences in microparasite community composition between the five colony types as
revealed by a distance-based redundancy analysis with colony type as the constraining factor
(spatial structure partialled out). The primary dbRDA-axis separating the colony types (1.9%
of total variation) is shown along the first unconstrained axis (MDS1, 16.4% of the variation).
Dots represent locations of individual colonies and diamonds are mean locations. **(c)** Parasite
loads in relation to colony type. Dots are parasite loads of individual colonies and diamonds
are means. Only microparasite taxa detected at least once among managed colonies and
among feral colonies are shown (same selection as in Figure 4). See Table S14 for an
overview of parasite prevalence and parasite loads by colony type.

Discussion

We compared the parasite burden of honeybee colonies in managed and feral conditions to

evaluate whether colonies that escaped from apiaries pose a risk to managed and wild bee

health by acting as reservoirs of disease-causing agents. The number of microparasites

detected per colony and the prevalence of four important taxa were clearly lower in feral

compared to managed colonies. This was explained by differences in population demography,

with most feral colonies being recent founders with very low parasite loads, and by

environmental differences between managed and feral colonies. We conclude that feral

honeybee colonies are unlikely to contribute disproportionately to the spread of bee parasites.

We considered parasite burden based on the prevalence of 18 microparasites determined using

qPCR. This means that we did not cover all known bee parasites. We also cannot exclude the

possibility of some false non-detections of targeted RNA viruses due to their high

evolutionary rates. However, our conclusions about relative differences in parasite burden

between groups are robust because all colonies were subjected to the same qPCR-assays and

because groups were evenly distributed across study regions.

Comparing colonies from four regions in southern Germany revealed strong geographic

differences in parasite numbers and parasite community composition. These differences might

be explained by a combination of factors including managed colony density, land use,

climate, and colonization histories of individual parasites. The management implication is that

moving managed hives, even over moderate distances (the distance between our study regions

Weilheim-Schongau and Munich is approx. 50 km), bears the risk of introducing bee parasites

that would otherwise not be present locally. Considering the potential negative effects on

local wild bee communities and on honeybees managed by non-migratory beekeepers

(Martínez-López et al., 2022), apicultural disease management needs to pay more attention to

regional differences in parasite communities.

Although region of sampling had a stronger effect on parasite communities than management,

we believe that the differences found between feral and managed colonies are ecologically

relevant. With about six parasite taxa detected per colony on the overall average, the reduced

count of one parasite per colony from managed to feral colonies is noteworthy, especially

since this pattern was consistent across study regions. Higher numbers of parasites have been

linked to higher colony mortality (vanEngelsdorp et al., 2009), although this relationship

might be non-linear (Ravoet et al., 2013). Analysing the associations between 18 parasite taxa

and colony mortality, Ravoet et al. (2013) found that winter mortality steeply increased with

the number of parasite species rising from 3 to 5, but additional parasites had no further

effect. In our study, the percentage of colonies with more than 5 detected parasite taxa was

only 47 % in the group of feral colonies but 65 % in the group of managed colonies, hence the

difference was likely within a relevant range.

Importantly, it was not entirely random which parasites were less prevalent in feral colonies:

Trypanosomatidae, chronic bee paralysis virus and deformed wing virus strains A and B were

less frequently detected in feral colonies. These four parasites are all important since they can

induce host mortality. Furthermore, they have been detected in other bee species, so they are

also relevant in the context of honeybee-wild bee interactions (Strobl et al., 2019; Tehel et al.,

2016). The two trypanosomatid species *Chrithidia mellificae* and *Lotmaria passim* (among

which we did not distinguish with our PCR assay) are unicellular parasites that colonize the

bees' hindgut (Schwarz et al., 2015). They have been experimentally shown to reduce the

lifespan of individual workers (Strobl et al., 2019) and their presence is associated with

colony-level winter mortality (Ravoet et al., 2013). Chronic bee paralysis virus (CBPV)

causes the bee paralysis disease in adult workers, which involves symptoms like hair loss,

undirected trembling walks, and loss of flight ability. The virus can spread in a colony via

worker contact and eventually lead to its collapse (Ribière et al., 2010). Interestingly, CBPV

seems to be an emerging threat, as indicated by a rapid rise of cases among Britain's apiaries

during the last decade (Budge et al., 2020). Given that in this study, its prevalence was 14.9%

in managed colonies but only 1.6% in feral colonies, the latter are unlikely to represent an

important dispersal route for the virus.

The reduction in the prevalence of deformed wing viruses (DWV) A and B in feral compared

to managed colonies is, ecologically, the most important detected difference since DWV is

regarded as one of the main causes of winter colony losses (Dainat et al., 2012a; de Miranda

& Genersch, 2010; Genersch et al., 2010). When transmitted during development, DWV

directly kill their host at the pupal stage or cause body deformations of the emerging bee,

leading to premature death (de Miranda & Genersch, 2010). Even infected bees that do not

show overt symptoms have their life expectancy reduced by DWV, which can result in mass

losses of bees on the colony level and its collapse during winter (Dainat et al., 2012a).

Importantly, the abundance of DWV in honeybee colonies and the severity of the resulting

symptoms are tightly linked to the co-occurrence with *V. destructor* (de Miranda & Genersch,

2010; Paxton et al., 2022). While DWV were unproblematic before the invasion of

V. destructor, the mite represented a new vector not only aiding the spread but also fuelling

the evolution of the viruses, as the emergence and ongoing replacement of DWV-A by the

presumably more virulent DWV-B indicates (McMahon et al., 2016; Paxton et al., 2022). In

turn, the mite's reproduction is enhanced by the presence of DWV, making the mite-virus pair

a deadly symbiosis (Di et al., 2016). Unfortunately, it was not possible in our study to

investigate the levels of *V. destructor* infestation, because the numbers of mites on foragers

are generally too low to make a reasonable analysis based on bees captured at the colony

entrance. It would have required sampling pupae or young bees from the brood nest or

assessing the rate at which dead mites naturally fall to the bottom of the nest cavity

(Dietemann et al., 2013), but this was not possible in the case of the feral colonies. However,

since DWV and *Varroa* correlate (Dainat et al., 2012b; Norton et al., 2021), the lower

prevalence of DWV in feral colonies suggests that mite infestation levels were also reduced.

Our findings are in seeming conflict with a previous study from England which concluded

that parasite pressure on feral colonies is relatively high. Thompson et al. (2014) tested for 10

parasites and found that these were equally prevalent among managed and feral colonies, but

the colony-level parasite abundance of one parasite, deformed wing virus, was significantly

higher in feral colonies (no difference was made between DWV-A and DWV-B). However,

the sample of feral colonies analysed in their study only included colonies aged at least one

year. We also found that the parasite prevalence in feral colonies aged at least one year

(overwintered feral colonies) did not differ significantly from managed colonies and that they

had relatively high loads of DWV-B (although not of DWV-A; Fig. 5c). Therefore, had we

only considered old feral colonies, we might have come to similar conclusions. However, we

would have then seriously overestimated the parasite burden in the feral population as a whole

because only about 10% of the feral colonies present in summer are older than one year (Kohl

et al., 2022), and especially the young, newly founded feral colonies had their parasite burden

reduced compared to managed colonies. These considerations demonstrate that it is important

to know the population demography of the feral honeybees under consideration when asking

questions about their ecological impact, e.g., their contribution to the spread of bee diseases.

The feral honeybees investigated in this study are known to be recent descendants from

colonies managed in apiaries (Kohl et al., 2022), and therefore, their reduced parasite burden

needs to be explained by ecological/environmental, rather than genetic, differences from

managed honeybees. By considering five colony types representing differences in colony age

and management history (Table 1), we gained some insights into why parasite burden is

reduced in feral colonies. We found that newly founded feral colonies had a lower parasite

burden than both overwintered feral colonies and overwintered managed colonies (from

which many of the young feral colonies directly descend). This supports the hypothesis that

the natural process of reproductive swarming – the abandonment of the old nest, the pause of

brood production, and the construction of fresh comb at a new dwelling place – leads to a

temporal release from parasite pressure (DeGrandi-Hoffman et al., 2017; Loftus et al., 2016;

Royce et al., 1991). The high proportion of young, swarm-founded colonies in the feral

population is one of the reasons for the overall significant difference in parasite burden

between feral and managed colonies. In the managed population, the proportion of young

colonies is much lower (about 50 %), and most young colonies are so-called nucleus colonies,

created by transferring several combs with brood from the mother colony into a new hive – a

completely different founding mechanism. Since nucleus colonies directly inherit the parasites

residing in the old combs and the brood, their parasite communities should resemble those of

established (old) colonies. Indeed, parasite counts did not differ significantly between nucleus

colonies and overwintered managed colonies, nor between nucleus colonies and overwintered

feral colonies, albeit overwintered managed colonies had the highest numbers and a different

community composition of parasites. The latter might be explained by the fact that beekeepers

usually manage established colonies in such a way that brood is continuously produced, while

managed nucleus colonies and overwintered feral colonies typically experience a brood pause

in spring and thus a temporal reduction of the breeding ground for parasites (Table 1) (Loftus

et al., 2016). Importantly, young managed colonies founded by swarms ("hived swarms")

were infested by a significantly higher number of microparasite taxa than young feral

colonies. This contrast needs to be regarded with care since our sample size of hived swarms

was low (N=9), but it suggests that not only differences in population age structure between

managed and feral colonies, but also in the environment, contribute to the difference in

parasite burden.

The most obvious environmental difference that has been demonstrated to affect parasite

pressure is that managed hives are typically clustered in apiaries, while feral colonies are

spatially dispersed (Lindström et al., 2008; Nolan & Delaplane, 2017; Seeley & Smith, 2015).

The crowding of colonies promotes random drifting and robbing behaviour, and both drifters

and robbers can transfer parasites between hives (Lindström et al., 2008; Peck & Seeley,

2019; Seeley & Smith, 2015). Another highly consistent difference is that beekeepers

conventionally keep their hives at ground level, while feral colonies typically nest in cavities

several meters above the ground. The bees' preference for aerial cavities is thought to serve as

a protection against predators, but a positive side effect might be that colonies dispose of sick

bees more easily. For example, both CBPV and DWV cause the loss of flight ability, and

flightless bees that fall out of the nest might struggle to find their way back when the entrance

is high above the ground. There are more differences which might directly or indirectly affect

the colonies' likelihood of becoming infected by parasites and of developing disease

symptoms, e.g., the size of the nest entrance, the presence of landing boards to aid bees

entering the hive, the insulation of the hive, the microclimate at the nest site, the frequency of

disturbance, or the type of food consumed. Given the role parasites play in limiting the

survival of managed colonies in apiculture, it seems worth experimentally investigating these

factors in more detail. For example, an interesting new question raised by our study is whether

it is possible to restore the natural parasite-reducing effect of swarm-founding under

apicultural management.

Table 1: Nonexclusive list of factors that might affect parasite burden, and their typical
manifestation in different types of managed and feral honeybee colonies.

Factor	Managed colonies			Feral colonies		Assumed effect on parasite prevalence
	Overwintered colonies	Nucleus colonies	Hived swarms	Overwintered colonies	Founder colonies	
Time since acaricide treatment	<1 year	<1 year	<1 year	>1 year	>1 year / <1 year*	Treatment reduces pressure from mites and associated viruses
Pause of brood production in spring	no	yes / no§	yes	yes	yes	Brood pause cuts off reproduction of brood parasites
Presence of old comb in nest	yes	yes	no	yes	yes / no#	Old comb can be a source of parasites
Spatial arrangement of colonies	clustered			dispersed		Parasite transmission rates are increased when colonies are closer to each other
Height of nest entrance	low			high		Colonies might get rid of sick, flightless bees more quickly when nest entrance is high above the ground

* Newly founded feral colonies that directly descend from managed hives were usually treated against mites < 1
year before founding.

§ Beekeepers can either produce a nucleus colony by taking combs with worker bees and brood from an
established colony and let the new colony raise a queen from existing female larvae, in which case they create a
brood pause, or directly introduce a mature queen to the new colony, in which case the nucleus colony does not
experience a significant brood pause.

Swarms prefer to occupy cavities that have been used by bees before, so they sometimes move into cavities still
containing old comb.

Feral honeybee colonies carrying many parasites and/or high parasite abundances would be a

nuisance to apicultural disease management and would pose a risk to the health of non-*Apis*

wild bees. However, we found that feral colonies have on average fewer parasites and lower

parasite loads than managed colonies. This is partly explained by the effect of natural swarm

reproduction and dispersal – a mechanism of parasite reduction that might also be effective in

other swarm-founding social insects (McGlynn, 2012). Given that feral colonies have a

relatively low parasite burden and that they make up a small fraction of the overall honeybee

population (in Germany feral colonies make up about 5% of the whole honeybee population

in summer), we conclude that it is unlikely that they significantly contribute to the spread of

bee parasites. On the contrary, new disease agents are probably primarily propagated by

managed colonies, as indicated by the higher prevalence of two emerging viruses, CBPV and

DWV-B, in the sample of managed hives. The management implication of this work is that

the prevention of epidemics is no suitable argument for the often-practised removal or

destruction of feral honeybee nests (Taylor et al., 2007). In fact, our data suggest that there is

no conflict between the promotion of wild-living honeybee populations and the management

of bee diseases in apiculture. What remains unclear is how the various environmental

differences between wild nests and hives at apiaries contribute to the reduced parasite burden

in feral colonies. Some known natural parasite-reducing factors, e.g., the spatial separation of

colonies and the periodic interruption of brood production, can readily be adopted by

beekeepers to increase the health of managed honeybees and to reduce the risk of disease

spread by apiculture (Büchler et al., 2020; Dynes et al., 2019; Loftus et al., 2016).

Authors' contributions

P.L.K.: conceptualization, formal analysis, funding acquisition, investigation, methodology,

visualization, writing–original draft, writing–review & editing; P.D..: formal analysis,

investigation, methodology, validation, writing–review & editing; B.R.: conceptualization,

investigation, methodology, writing–review & editing; S.R.: conceptualization, investigation,

methodology, writing–review & editing; F.R.: conceptualization, investigation, methodology,

writing–review & editing; I.S.D.: conceptualization, supervision, writing–review & editing;

M.H.: resources, supervision, writing–review & editing.

Acknowledgements

545   We thank Luis Sikora, Norbert Wimmer and Kurt Zeimentz for sharing locations of

546   woodpecker cavities which serve as nest sites for feral honeybees in German forests. We also

547   thank all beekeepers who granted us permission to sample bees from their hives. This study

548   was supported German Federal Environmental Foundation (DBU).

References

Ayala, A. J., Yabsley, M. J., & Hernandez, S. M. (2020). A Review of Pathogen Transmission

at the Backyard Chicken–Wild Bird Interface. *Frontiers in Veterinary Science*,

7(September), 1–15. https://doi.org/10.3389/fvets.2020.539925

Bailey, L. (1958). Wild Honeybees and Disease. *Bee World*, *39*(4), 92–95.

https://doi.org/10.1080/0005772X.1958.11095044

Bartlett, L. J. (2022). Frontiers in effective control of problem parasites in beekeeping.

International Journal for Parasitology: Parasites and Wildlife, *17*(February), 263–272.

https://doi.org/10.1016/j.ijppaw.2022.03.003

Blaker, H. (2000). Confidence curves and improved exact confidence intervals for discrete

distributions. *Canadian Journal of Statistics*, *28*, 783–798.

Bolker, B. and R. D. C. T. (2022). *bbmle: Tools for General Maximum Likelihood Estimation.*

R package version 1.0.25. https://cran.r-project.org/package=bbmle

Borcard, D., & Legendre, P. (2002). All-scale spatial analysis of ecological data by means of

principal coordinates of neighbour matrices Daniel. *Ecological Modelling*, *153*, 51–68.

Brooks, M. E., Kristensen, K., Darrigo, M. R., Rubim, P., Uriarte, M., Bruna, E., & Bolker, B.

M. (2019). Statistical modeling of patterns in annual reproductive rates. *Ecology*, *100*(7),

1–7. https://doi.org/10.1002/ecy.2706

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A.,

Skaug, H. J., Mächler, M., & Bolker, B. M. (2017). glmmTMB balances speed and

flexibility among packages for zero-inflated generalized linear mixed modeling. *R

Journal*, *9*(2), 378–400. https://doi.org/10.32614/rj-2017-066

Brosi, B. J., Delaplane, K. S., Boots, M., & De Roode, J. C. (2017). Ecological and

evolutionary approaches to managing honeybee disease. *Nature Ecology and Evolution*,

1(9), 1250–1262. https://doi.org/10.1038/s41559-017-0246-z

Browne, K. A., Hassett, J., Geary, M., Moore, E., Soland-reckeweg, G., Ferrari, R., Loughlin,

E. Mac, Brien, E. O., Driscoll, S. O., Young, P., Pinto, M. A., Grace, P., Browne, K. A.,

Hassett, J., Geary, M., Moore, E., Soland-reckeweg, G., Ferrari, R., Loughlin, E. Mac, …

Mccormack, G. P. (2020). Investigation of free-living honey bee colonies in Ireland.

Journal of Apicultural Research, *0*(0), 1–12.

https://doi.org/10.1080/00218839.2020.1837530

Büchler, R., Uzunov, A., Kovačić, M., Prešern, J., Pietropaoli, M., Hatjina, F., Pavlov, B.,

Charistos, L., Formato, G., Galarza, E., Gerula, D., Gregorc, A., Malagnini, V., Meixner,

M., Nedić, N., Puškadija, Z., Rivera-Gomis, J., Rogelj Jenko, M., Smodiš Škerl, M. I., …

Nanetti, A. (2020). Summer brood interruption as integrated management strategy for

effective Varroa control in Europe. *Journal of Apicultural Research*, *59*(5), 764–773.

https://doi.org/10.1080/00218839.2020.1793278

Budge, G. E., Barrett, B., Jones, B., Pietravalle, S., Marris, G., Chantawannakul, P., Thwaites,

R., Hall, J., Cuthbertson, A. G. S., & Brown, M. A. (2010). The occurrence of

Melissococcus plutonius in healthy colonies of Apis mellifera and the efficacy of

European foulbrood control measures. *Journal of Invertebrate Pathology*, *105*(2), 164–

170. https://doi.org/10.1016/j.jip.2010.06.004

Budge, G. E., Simcock, N. K., Holder, P. J., Shirley, M. D. F., Brown, M. A., Van Weymers,

P. S. M., Evans, D. J., & Rushton, S. P. (2020). Chronic bee paralysis as a serious

emerging threat to honey bees. *Nature Communications*, *11*(1).

https://doi.org/10.1038/s41467-020-15919-0

Burnham, P. A., Alger, S. A., Case, B., Boncristiani, H., Hébert-Dufresne, L., & Brody, A. K.

(2021). Flowers as dirty doorknobs: Deformed wing virus transmitted between Apis

mellifera and Bombus impatiens through shared flowers. *Journal of Applied Ecology*,

58(10), 2065–2074. https://doi.org/10.1111/1365-2664.13962

Cepero, A., Martín-Hernández, R., Prieto, L., Gómez-Moracho, T., Martínez-Salvador, A.,

Bartolomé, C., Maside, X., Meana, A., & Higes, M. (2015). Is Acarapis woodi a single

species? A new PCR protocol to evaluate its prevalence. *Parasitology Research*, *114*(2),

651–658. https://doi.org/10.1007/s00436-014-4229-6

Chang, S. Y., & Hoopingarner, R. A. (1991). Relative Importance of Feral Honey Bees in

Apple Pollination. In *Acta Horticulturae* (Issue 288, pp. 239–243).

https://doi.org/10.17660/actahortic.1991.288.35

Costanzi, L., Brambilla, A., Di Blasio, A., Dondo, A., Goria, M., Masoero, L., Gennero, M.

S., & Bassano, B. (2021). Beware of dogs! Domestic animals as a threat for wildlife

conservation in Alpine protected areas. *European Journal of Wildlife Research*, *67*(4).

https://doi.org/10.1007/s10344-021-01510-5

Crane, E. (1990). *Bees and beekeeping: science, practice and world resources*. Heinemann

Newnes.

D'Alvise, P., Seeburger, V., Gihring, K., Kieboom, M., & Hasselmann, M. (2019). Seasonal

dynamics and co - occurrence patterns of honey bee pathogens revealed by high -

throughput RT - qPCR analysis. *Ecology and Evolution*, *July*, 1–12.

https://doi.org/10.1002/ece3.5544

Dainat, B., Evans, J. D., Chen, Y. P., Gauthier, L., & Neumann, P. (2012a). Dead or alive:

Deformed wing virus and varroa destructor reduce the life span of winter honeybees.

Applied and Environmental Microbiology, *78*(4), 981–987.

https://doi.org/10.1128/AEM.06537-11

Dainat, B., Evans, J. D., Chen, Y. P., Gauthier, L., & Neumann, P. (2012b). Predictive

markers of honey bee colony collapse. *PLoS ONE*, *7*(2).

https://doi.org/10.1371/journal.pone.0032151

Daniels, T. J., & Bekoff, M. (1989). Feralization: The making of wild domestic animals.

Behavioural Processes, *19*(1–3), 79–94. https://doi.org/10.1016/0376-6357(89)90032-6

de Miranda, J. R., & Genersch, E. (2010). Deformed wing virus. *Journal of Invertebrate*

Pathology, *103*(SUPPL. 1), S48–S61. https://doi.org/10.1016/j.jip.2009.06.012

DeGrandi-Hoffman, G., Ahumada, F., & Graham, H. (2017). Are dispersal mechanisms

changing the host-parasite relationship and increasing the virulence of varroa destructor

(mesostigmata: Varroidae) in managed honey bee (hymenoptera: Apidae) colonies?

Environmental Entomology, *46*(4), 737–746. https://doi.org/10.1093/ee/nvx077

Deutscher Imkerbund e.V. (2020). *Jahresbericht 2020*. https://deutscherimkerbund.de/304-

Taetigkeitsberichte

Di, G., Annoscia, D., Margiotta, M., Ferrara, R., Varricchio, P., & Zanni, V. (2016). A

mutualistic symbiosis between a parasitic mite and a pathogenic virus undermines honey

bee immunity and health. *Proceedings of the National Academy of Sciences of the United*

States of America, *113*(12), 3203–3208. https://doi.org/10.1073/pnas.1523515113

Dick, C. W. (2001). Genetic rescue of remnant tropical trees by an alien pollinator.

Proceedings of the Royal Society B: Biological Sciences, *268*(1483), 2391–2396.

https://doi.org/10.1098/rspb.2001.1781

Dietemann, V., Nazzi, F., Martin, S. J., Anderson, D. L., Locke, B., Delaplane, K. S.,

Wauquiez, Q., Tannahill, C., Frey, E., Ziegelmann, B., Rosenkranz, P., & Ellis, J. D.

(2013). Standard methods for varroa research. *Journal of Apicultural Research*, *52*(1), 1–

54. https://doi.org/10.3896/IBRA.1.52.1.09

Dietemann, V., Pirk, C. W. W., & Crewe, R. (2009). Is there a need for conservation of

honeybees in Africa? *Apidologie*, *40*(3), 285–295. https://doi.org/10.1051/apido/2009013

Dubaić, J. B., Simonović, S., Plećaš, M., Stanisavljević, L., Davidović, S., Tanasković, M., &

Ćetković, A. (2021). Unprecedented density and persistence of feral honey bees in urban

environments of a large SE-European city (Belgrade, Serbia). *Insects*, *12*, 1127.

https://doi.org/10.3390/insects12121127

Dynes, T. L., Berry, J. A., Delaplane, K. S., Brosi, B. J., & De Roode, J. C. (2019). Reduced

density and visually complex apiaries reduce parasite load and promote honey

production and overwintering survival in honey bees. *PLoS ONE*, *14*(5), 1–16.

https://doi.org/10.1371/journal.pone.0216286

Evans, J. D. (2006). Beepath: An ordered quantitative-PCR array for exploring honey bee

immunity and disease. *Journal of Invertebrate Pathology*, *93*(2), 135–139.

https://doi.org/10.1016/j.jip.2006.04.004

Faurot-Daniels, C., Glenny, W., Daughenbaugh, K. F., McMenamin, A. J., Burkle, L. A., &

Flenniken, M. L. (2020). Longitudinal monitoring of honey bee colonies reveals dynamic

nature of virus abundance and indicates a negative impact of Lake Sinai virus 2 on

colony health. In *PLoS ONE* (Vol. 15, Issue 9 September).

https://doi.org/10.1371/journal.pone.0237544

Frey, E., & Rosenkranz, P. (2014). Autumn invasion rates of Varroa destructor

(Mesostigmata: Varroidae) into honey bee (Hymenoptera: Apidae) colonies and the

resulting increase in mite populations. *Journal of Economic Entomology*, *107*(2), 508–

515.

Fries, I., & Camazine, S. (2001). Implications of horizontal and vertical pathogen

transmission for honey bee epidemiology. *Apidologie*, *32*(3), 199–214.

https://doi.org/10.1051/apido:2001122

Fries, I., Lindström, A., & Korpela, S. (2006). Vertical transmission of American foulbrood

(Paenibacillus larvae) in honey bees (Apis mellifera). *Veterinary Microbiology*, *114*(3–

4), 269–274. https://doi.org/10.1016/j.vetmic.2005.11.068

Fürst, M. a., McMahon, D. P., Osborne, J. L., Paxton, R. J., & Brown, M. J. F. (2014).

Disease associations between honeybees and bumblebees as a threat to wild pollinators.

Nature, *506*(7488), 364–366. https://doi.org/10.1038/nature12977

Garbuzov, M., Balfour, N. J., Shackleton, K., Al Toufailia, H., Scandian, L., & Ratnieks, F. L.

W. (2020). Multiple methods of assessing nectar foraging conditions indicate peak

foraging difficulty in late season. *Insect Conservation and Diversity*, *13*(6), 532–542.

https://doi.org/10.1111/icad.12420

Geldmann, J., & González-Varo, J. P. (2018). Conserving honey bees does not help wildlife.

Science, *359*(6374), 392–393. https://doi.org/10.1126/science.aar2269

Genersch, E., von der Ohe, W., Kaatz, H., Schroeder, A., Otten, C., Büchler, R., Berg, S.,

Ritter, W., Mühlen, W., Gisder, S., Meixner, M., Liebig, G., & Rosenkranz, P. (2010).

The German bee monitoring project: a long term study to understand periodically high

winter losses of honey bee colonies. *Apidologie*, *41*(3), 332–352.

https://doi.org/10.1051/apido/2010014

Gortázar, C., Ferroglio, E., Höfle, U., Frölich, K., & Vicente, J. (2007). Diseases shared

between wildlife and livestock: A European perspective. *European Journal of Wildlife*

Research, *53*(4), 241–256. https://doi.org/10.1007/s10344-007-0098-y

Goulson, D., & Hughes, W. O. H. (2015). Mitigating the anthropogenic spread of bee

parasites to protect wild pollinators. *Biological Conservation*, *191*, 10–19.

https://doi.org/10.1016/j.biocon.2015.06.023

Hartig, F. (2022). *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level/Mixed)*

Regression Models. R package version 0.4.5. https://cran.r-

project.org/package=DHARMa

Hothorn, T., Bretz, F., & Westfall, P. (2008). Simultaneous Inference in General Parametric

Models. *Biometrical Journal*, *50*(3), 346–363.

Hung, K.-L. J., Kingston, J. M., Albrecht, M., Holway, D. A., & Kohn, J. R. (2018). The

worldwide importance of honey bees as pollinators in natural habitats. *Proceedings of*

the Royal Society B, *285*, 20172140. https://doi.org/doi:10.1086/303379

Iwasaki, J. M., & Hogendoorn, K. (2022). Mounting evidence that managed and introduced

bees have negative impacts on wild bees : an updated review. *Current Research in Insect*

Science, *2*, 100043. https://doi.org/10.1016/j.cris.2022.100043

Kohl, P. L., & Rutschmann, B. (2018). The neglected bee trees: European beech forests as a

home for feral honey bee colonies. *PeerJ*, *6*, e4602. https://doi.org/10.7717/peerj.4602

Kohl, P. L., Rutschmann, B., & Steffan-Dewenter, I. (2022). Population demography of feral

honeybee colonies in Central European forests. *Royal Society Open Science*, *9*, 220565.

https://doi.org/10.1098/rsos.220565

Kraus, B., & Page, R. E. R. (1995). Effect of Varroa jacobsoni (Mesostigmata: Varroidae) on

feral Apis mellifera (Hymenoptera: Apidae) in California. *Environmental Entomology*,

24(6), 1473–1480. https://doi.org/10.1093/ee/24.6.1473

Lenth, R. V. (2022). *emmeans: Estimated Marginal Means, aka Least-Squares Means* (R

package version 1.7.3). https://cran.r-project.org/package=emmeans

Lindström, A., Korpela, S., & Fries, I. (2008). Horizontal transmission of Paenibacillus larvae

spores between honey bee (Apis mellifera) colonies through robbing. *Apidologie*, *39*(5),

515–522. https://doi.org/10.1051/apido:2008032

Locke, B., Forsgren, E., Fries, I., & de Miranda, J. R. (2012). Acaricide treatment affects viral

dynamics in Varroa destructor-infested honey bee colonies via both host physiology and

mite control. *Applied and Environmental Microbiology*, *78*(1), 227–235.

https://doi.org/10.1128/AEM.06094-11

Loftus, J. C., Smith, M. L., & Seeley, T. D. (2016). *How Honey Bee Colonies Survive in the*

Wild : Testing the Importance of Small Nests and Frequent Swarming. 1–11.

https://doi.org/10.1371/journal.pone.0150362

Lourenço, A. P., Mackert, A., Cristino, A. D. S., & Simões, Z. L. P. (2008). Validation of

reference genes for gene expression studies in the honey bee, Apis mellifera, by

quantitative real-time RT-PCR. *Apidologie*, *39*(3), 372–385.

https://doi.org/10.1051/apido:2008015

Martínez-López, V., Ruiz, C., & De la Rúa, P. (2022). Migratory beekeeping and its influence

on the prevalence and dispersal of pathogens to managed and wild bees. *International*

Journal for Parasitology: Parasites and Wildlife, *18*(February), 184–193.

https://doi.org/10.1016/j.ijppaw.2022.05.004

Martínez, J., Simon, V., Gonzalez, B., & Conget, P. (2010). A real-time PCR-based strategy

for the detection of Paenibacillus larvae vegetative cells and spores to improve the

diagnosis and the screening of American foulbrood. *Letters in Applied Microbiology*,

50(6), 603–610. https://doi.org/10.1111/j.1472-765X.2010.02840.x

McGlynn, T. P. (2012). The ecology of nest movement in social insects. *Annual Review of*

Entomology, *57*, 291–308. https://doi.org/10.1146/annurev-ento-120710-100708

McMahon, D. P., Natsopoulou, M. E., Doublet, V., Fürst, M., Weging, S., Brown, M. J. F.,

Gogol-Döring, A., & Paxton, R. J. (2016). Elevated virulence of an emerging viral

genotype as a driver of honeybee loss. *Proceedings of the Royal Society B: Biological*

Sciences, *283*(1833). https://doi.org/10.1098/rspb.2016.0811

Mikheyev, A. S., Tin, M. M. Y., Arora, J., & Seeley, T. (2016). Museum samples reveal rapid

evolution by wild honey bees exposed to a novel parasite. *Nature Communications*,

6(AUGUST 2015), 1–8. https://doi.org/10.1038/ncomms8991

Mysterud, A., & Rolandsen, C. M. (2019). Fencing for wildlife disease control. *Journal of*

Applied Ecology, *56*(3), 519–525. https://doi.org/10.1111/1365-2664.13301

Neumann, P., & Blacquière, T. (2017). The Darwin cure for apiculture? Natural selection and

managed honeybee health. *Evolutionary Applications*, *10*(3), 226–230.

https://doi.org/10.1111/eva.12448

Nolan, M. P., & Delaplane, K. S. (2017). Distance between honey bee Apis mellifera colonies

regulates populations of Varroa destructor at a landscape scale. *Apidologie*, *48*(1), 8–16.

https://doi.org/10.1007/s13592-016-0443-9

Norton, A. M., Remnant, E. J., Tom, J., Buchmann, G., Blacquiere, T., & Beekman, M.

(2021). Adaptation to vector-based transmission in a honeybee virus. *Journal of Animal*

Ecology, *90*(10), 2254–2267. https://doi.org/10.1111/1365-2656.13493

Oksanen, J., Simpson, G., Blanchet, F., Kindt, R., Legendre, P., Minchin, P., O'Hara, R.,

Solymos, P., Stevens, M., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker,

B., Borcard, D., Carvalho, G., Chirico, M., De Caceres, M., Durand, S., … Weedon, J.

(2022). *vegan: Community Ecology Package* (R package version 2.6-2).

Oldroyd, B. P., Thexton, E. G., Lawler, S. H., & Crozier, R. H. (1997). Population

demography of australian feral bees (Apis mellifera). *Oecologia*, *111*, 381–387.

https://doi.org/10.1007/s004420050249

Oleksa, A., Gawroński, R., & Tofilski, A. (2013). Rural avenues as a refuge for feral honey

bee population. *Journal of Insect Conservation*, *17*(3), 465–472.

https://doi.org/10.1007/s10841-012-9528-6

Panziera, D., Requier, F., Chantawannakul, P., Pirk, C. W. W., & Blacquière, T. (2022). The

Diversity Decline in Wild and Managed Honey Bee Populations Urges for an Integrated

Conservation Approach. *Frontiers in Ecology and Evolution*, *10*(March).

https://doi.org/10.3389/fevo.2022.767950

Papp, T., Spann, D., & Marschang, R. E. (2014). Development and use of a real-time

polymerase chain reaction for the detection of group II invertebrate iridoviruses in pet

lizards and prey insects. J. Zoo Wildl. Med., 45(2), 219–27./stable/24551086. *Journal of*

Zoo and Wildlife Medicine, *45*(2), 219–227.

Paxton, R. J., Schäfer, M. O., Nazzi, F., Zanni, V., Annoscia, D., Marroni, F., Bigot, D.,

Laws-Quinn, E. R., Panziera, D., Jenkins, C., & Shafiey, H. (2022). Epidemiology of a

major honey bee pathogen, deformed wing virus: potential worldwide replacement of

genotype A by genotype B. *International Journal for Parasitology: Parasites and*

Wildlife, *18*(May), 157–171. https://doi.org/10.1016/j.ijppaw.2022.04.013

Peck, D. T., & Seeley, T. D. (2019). Mite bombs or robber lures ? The roles of drifting and

robbing in Varroa destructor transmission from collapsing honey bee colonies to their

neighbors. *PLoS ONE*, *14*(6), e0218392.

Piot, N., Schweiger, O., Meeus, I., Yañez, O., Straub, L., Villamar-Bouza, L., De la Rúa, P.,

Jara, L., Ruiz, C., Malmstrøm, M., Mustafa, S., Nielsen, A., Mänd, M., Karise, R., Tlak-

Gajger, I., Özgör, E., Keskin, N., Diévart, V., Dalmon, A., … de Miranda, J. R. (2022).

Honey bees and climate explain viral prevalence in wild bee communities on a

continental scale. *Scientific Reports*, *12*(1), 1–12. https://doi.org/10.1038/s41598-022-

05603-2

Pirk, C. W. W., Crewe, R. M., & Moritz, R. F. A. (2017). Risks and benefits of the biological

interface between managed and wild bee pollinators. *Functional Ecology*, *31*(1), 47–55.

https://doi.org/10.1111/1365-2435.12768

R Core Team. (2022). *R: A Language and Environment for Statistical Computing*. R

Foundation for Statistical Computing; R Foundation for Statistical Computing.

https://www.r-project.org/

Ratnieks, F. W., & Nowakowski, J. (1989). Honeybee swarms accept bait hives contaminated

with American foulbrood disease. *Ecological Entomology*, *14*(4), 475–478.

https://doi.org/10.1111/j.1365-2311.1989.tb00951.x

Ravoet, J., De Smet, L., Meeus, I., Smagghe, G., Wenseleers, T., & de Graaf, D. C. (2014).

Widespread occurrence of honey bee pathogens in solitary bees. *Journal of Invertebrate*

Pathology, *122*, 55–58. https://doi.org/10.1016/j.jip.2014.08.007

Ravoet, J., Maharramov, J., Meeus, I., De Smet, L., Wenseleers, T., Smagghe, G., & de Graaf,

D. C. (2013). Comprehensive Bee Pathogen Screening in Belgium Reveals Crithidia

mellificae as a New Contributory Factor to Winter Mortality. *PLoS ONE*, *8*(8).

https://doi.org/10.1371/journal.pone.0072443

Requier, F., Garnery, L., Kohl, P. L., Njovu, H. K., Pirk, C. W. W., Crewe, R. M., & Steffan-

dewenter, I. (2019). The conservation of native honey bees is crucial. *Trends in Ecology*

& Evolution, *34*(9), 789–798. https://doi.org/10.1016/j.tree.2019.04.008

Ribière, M., Olivier, V., & Blanchard, P. (2010). Chronic bee paralysis: A disease and a virus

like no other? *Journal of Invertebrate Pathology*, *103*(SUPPL. 1), S120–S131.

https://doi.org/10.1016/j.jip.2009.06.013

Rollin, O., & Garibaldi, L. A. (2019). Impacts of honeybee density on crop yield: A meta-

analysis. *Journal of Applied Ecology*, *56*(5), 1152–1163. https://doi.org/10.1111/1365-

2664.13355

Royce, L. A., Rossignol, P. A., Burgett, D. M., & Stringer, B. A. (1991). Reduction of

tracheal mite parasitism of honey bees by swarming. *Philosophical Transactions - Royal*

Society of London, B, *331*(1260), 123–129. https://doi.org/10.1098/rstb.1991.0003

Rutschmann, B., Kohl, P. L., Machado, A., & Steffan-Dewenter, I. (2022). Semi-natural

habitats promote winter survival of wild-living honeybees in an agricultural landscape.

Biological Conservation, *266*, 109450. https://doi.org/10.1016/j.biocon.2022.109450

Schmid-Hempel, P. (1998). *Parasites in social insects*. Princeton University Press.

Schommer, T. J., & Woolever, M. M. (2008). A review of disease related conflicts between

domestic sheep and goats and bighorn sheep. In *USDA Forest Service - General*

Technical Report RMRS-GTR-209. U.S. Department of Agriculture, Forest Service,

Rocky Mountain Research Station.

Schwarz, R. S., Bauchan, G. R., Murphy, C. A., Ravoet, J., De Graaf, D. C., & Evans, J. D.

(2015). Characterization of two species of trypanosomatidae from the Honey Bee Apis

mellifera: Crithidia mellificae Langridge and McGhee, and Lotmaria passim n. gen., n.

sp. *Journal of Eukaryotic Microbiology*, *62*(5), 567–583.

https://doi.org/10.1111/jeu.12209

Seeley, T. D. (2007). Honey bees of the Arnot Forest: a population of feral colonies persisting

with Varroa destructor in the northeastern United States. *Apidologie*, *38*, 19–29.

Seeley, T. D. (2010). *Honeybee Democracy*. Princeton University Press.

Seeley, T. D. (2017). Life-history traits of wild honey bee colonies living in forests around

Ithaca, NY, USA. *Apidologie*, *48*, 743–754. https://doi.org/10.1007/s13592-017-0519-1

Seeley, T. D., & Smith, M. L. (2015). Crowding honeybee colonies in apiaries can increase

their vulnerability to the deadly ectoparasite Varroa destructor. *Apidologie*, *46*(6), 716–

727. https://doi.org/10.1007/s13592-015-0361-2

Stevenson, M., Sergeant, E., Nunes, T., Heuer, C., Marshall, J., Sanchez, J., Thornton, R.,

Reiczigel, J., Robison-Cox, J., Sebastiani, P., Solymos, P., Yoshida, K., Jones, G.,

Pirikahu, S., Firestone, S., Kyle, R., Popp, J., Jay, M., Reynard, C., … Rabiee, A. (2022).

epiR: Tools for the Analysis of Epidemiological Data (R package version 2.0.46).

https://cran.r-project.org/package=epiR

Strobl, V., Yañez, O., Straub, L., Albrecht, M., & Neumann, P. (2019). Trypanosomatid

parasites infecting managed honeybees and wild solitary bees. *International Journal for

Parasitology*, *49*(8), 605–613. https://doi.org/10.1016/j.ijpara.2019.03.006

Taylor, M. A., Goodwin, R. M., McBrydie, H. M., & Cox, H. M. (2007). Destroying managed

and feral honey bee (Apis mellifera) colonies to eradicate honey bee pests. *New Zealand

Journal of Crop and Horticultural Science*, *35*(3), 313–323.

https://doi.org/10.1080/01140670709510197

Tehel, A., Brown, M. J. F., & Paxton, R. J. (2016). Impact of managed honey bee viruses on

wild bees. *Current Opinion in Virology*, *19*, 16–22.

https://doi.org/10.1016/j.coviro.2016.06.006

Tehel, A., Streicher, T., Tragust, S., & Paxton, R. J. (2022). Experimental cross species

transmission of a major viral pathogen in bees is predominantly from honeybees to

bumblebees. *Proceedings of the Royal Society B: Biological Sciences*, *289*(1969).

https://doi.org/10.1098/rspb.2021.2255

Thompson, C E. (2012). *The health and status of the feral honeybee (Apis mellifera sp) and*

Apis mellifera mellifera population of the UK (Issue November). The Unversity of

Leeds.

Thompson, Catherine E, Biesmeijer, J. C., Allnutt, T. R., Pietravalle, S., & Budge, G. E.

(2014). Parasite pressures on feral honey bees (Apis mellifera sp.). *PloS One*, *9*(8),

e105164. https://doi.org/10.1371/journal.pone.0105164

Traynor, K. S., Mondet, F., de Miranda, J. R., Techer, M., Kowallik, V., Oddie, M. A. Y.,

Chantawannakul, P., & McAfee, A. (2020). Varroa destructor: A Complex Parasite,

Crippling Honey Bees Worldwide. *Trends in Parasitology*, *36*(7), 592–606.

https://doi.org/10.1016/j.pt.2020.04.004

vanEngelsdorp, D., Evans, J. D., Saegerman, C., Mullin, C., Haubruge, E., Nguyen, B. K.,

Frazier, M., Frazier, J., Cox-Foster, D., Chen, Y., Underwood, R., Tarpy, D. R., & Pettis,

J. S. (2009). Colony collapse disorder: A descriptive study. *PLoS ONE*, *4*(8).

https://doi.org/10.1371/journal.pone.0006481

Weigand, M., Staab, J., Wurm, M., & Taubenböck, H. (2020). Spatial and semantic effects of

LUCAS samples on fully automated land use/land cover classification in high-resolution

Sentinel-2 data. *International Journal of Applied Earth Observation and*

Geoinformation, *88*(February), 102065. https://doi.org/10.1016/j.jag.2020.102065

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New

York.

Wilfert, L., Long, G., Leggett, H. C., Schmid-Hempel, P., Butlin, R., Martin, S. J. M., &

Boots, M. (2016). Honeybee disease: Deformed wing virus is a recent global epidemic in

honeybees driven by Varroa mites. *Science*, *351*(6273), 594–597.

https://doi.org/10.1126/science.aac9976

Winston, M. L. (1992). The biology and management of africanized honey bees. *Annual*

Review of Entomology, *37*, 173–193.

Youngsteadt, E., Appler, R. H., López-Uribe, M. M., Tarpy, D. R., & Frank, S. D. (2015).

Urbanization increases pathogen pressure on feral and managed honey bees. *PLoS ONE*,

10(11), 1–16. https://doi.org/10.1371/journal.pone.0142031