Differential effect of climate on structural and biochemical plant traits

Dinesh Thakur[1*], Věroslava Hadincová[1], Renáta Schnablová[1], Helena Synková[3], Daniel Haisel[3],

Nada Wilhelmová[3], Tomáš Dostálek[1,2], Zuzana Münzbergová[1,2*]

*Corresponding author e-mail: 552.dinesh@gmail.com and zuzmun@natur.cuni.cz

[1] Department of Population Ecology, Institute of Botany, Czech Academy of Sciences, Prague,

Czechia. [2] Department of Botany, Faculty of Science, Charles University, Prague, Czechia.

[3] Institute of Experimental Botany, Czech Academy of Sciences, Prague, Czechia.

ABSTRACT

Exploring patterns and causes of intraspecific trait variation is crucial for a better

understanding of the effects of climate change on plant populations and ecosystems. However,

our current understanding of the intraspecific trait variation is mainly based on structural

(morphological) traits, and we have limited knowledge on patterns and causes of variation in

biochemical traits (e.g., leaf pigments) which are also crucial for plant adaptation. As a result, we

also do not know how similar the climatic effects on structural versus biochemical traits are.

Using plant traits from 110 genotypes representing 11 populations of *Festuca rubra*

grown in 4 different climates, we studied trait covariation among structural traits (linked to

fitness, resource use, gas exchange and reproduction) and biochemical traits (linked to

photosynthesis, photoprotection and oxidative stress). We also disentangled the relative role of

climate of origin and cultivation climate on structural versus biochemical traits and tested for

adaptive plasticity in the traits.

We found that 1) biochemical traits did not covary with structural traits and represent

'photoharvesting – photoprotection' strategy dimension of functional variation; 2) the interactive

effects of climate of origin and cultivation were more pronounced for biochemical than structural

traits. 3) Trait plasticity was affected by the climate of origin (precipitation and temperature as

well as their interaction); 4) *F. rubra* showed both adaptive as well as mal-adaptive plasticity and

adaptiveness depends upon trait type, cultivation climate and climate of origin.

Overall, our results suggest that structural and biochemical plant traits respond

differentially to climate and response of one group of traits thus cannot be predicted from the

other. The responses are also strongly determined by interactions between the climate of origin

and cultivation. More studies on variation in biochemical traits, their correspondence to other

traits and their variation with climate are thus needed.

Key Words: climate change effects; adaptive plasticity; clonal plant; adaptation strategies;

intraspecific trait variation; variance decomposition

INTRODUCTION

Plant traits determine various plant functions, define ecological strategies, describe responses to environmental change and also the effects of plants on ecosystem functioning (Westoby et al. 2002, McGill et al. 2006, Violle et al. 2007, Enquist et al. 2015). Therefore, establishing links between functional traits and environmental factors is crucial for understanding how plant populations and communities respond to climate change.

Traits vary both within and between species, however, our trait-based understanding is mainly based on interspecific trait variability and only recently, ecologists have focused on the implications and consequences of intraspecific trait variation (ITV) (Bolnick et al. 2011, Siefert et al. 2015, Westerband et al. 2021). Lately, ITV has been suggested to be equally important as interspecific variation for understanding, predicting, and managing large ecological processes (Bennett et al. 2016, Martin et al. 2017, Henn et al. 2018, Mitchell et al. 2020, Westerband et al. 2021).

ITV arises from both genetic differences and phenotypic plasticity (Matesanz et al. 2012). Intraspecific variation in traits due to genetic differences is achieved very slowly by the process of natural selection while plasticity allows for rapid responses. Many studies elucidated genetic versus plastic responses in various plant traits in various species (Blanquart et al. 2013, Münzbergová et al. 2017, Datta et al. 2017, Manzanedo et al. 2019, Kosová et al. 2022). However, most of these studies deal only with structural traits (i.e., morphological traits e.g., specific leaf area, plant height, plant biomass etc.). Variation in biochemical traits, such as contents of leaf pigments, is an important mechanism through which plants can increase mean fitness in a variable environment (Taiz et al. 2015), but variation in such traits is much less studied. Unlike structural traits, biochemical traits are easy to construct and thus easy to modify

with changing conditions within a growing season and could thus be more plastic than the
structural traits.

Biochemical traits are closely related to the physiological activity of plants under variable
environments (Taiz et al. 2015) and should be closely linked to structural traits as all the
structures in plants are formed as a result of biochemical processes (e.g., high content of
photosynthetic pigments should lead to increased plant biomass). The available ecological
studies exploring biochemical traits study them separately of structural traits e.g. (Alonso-
Amelot et al. 2007, Martin et al. 2007). Due to this, we do not know if the genetic versus plastic
responses differ between structural and biochemical traits. Identifying specific plant functions
that are more genetically controlled or are more plastic, is crucial for our ability to understand
and predict species responses to the ongoing climate change more accurately (Henn et al. 2018,
Pérez-Ramos et al. 2019).

Phenotypic plasticity is considered a major mechanism of response to environmental
variability (Valladares et al. 2006, Nicotra et al. 2010, Henn et al. 2018). Despite a range of
studies on the plasticity of various traits, we know little if the plasticity is adaptive. Plasticity is
advantageous only if it increases plant fitness of an individual in a new environment (Nicotra et
al. 2010). In such a case, it can be considered as adaptive (van Kleunen and Fisher 2001, Enquist
et al. 2015). Recently, Enquist et al. (2015) suggested that we can determine whether phenotypic
plasticity is adaptive by exploring if the trait changes toward its optimum in a specific
environment (i.e., trait values in the home environment). If the traits of plants grown in a foreign
climate converge towards traits of plants which are at home in that climate, the trait shift is
considered adaptive. However, only very few studies have aCtempted to explore this so far
(Henn et al. 2018, Bush et al. 2021).

The extent and adaptive nature of plasticity may be limited by genetic basis as well as by

new environmental conditions (Valladares et al. 2006). For instance, plants from one climate of

origin may be more plastic and plastic responses may be stronger to one environmental factor

than to another (Münzbergová et al. 2017). Similarly, plants from a specific climate of origin

may show adaptive plasticity in some climates but not in others. However, such knowledge is

still largely lacking. Therefore, we also need to elucidate if the adaptive or mal-adaptive

plasticity is consistent among populations from different climates of origin shifted to different

novel climates. We also need to test if adaptive plasticity is consistent among different trait types

(e.g., structural versus biochemical traits). This will enable us to better understand the adaptive

potential of plant populations under climate change.

Both at intraspecific and interspecific levels, traits do not vary independently but covary

with other traits (Westoby et al. 2002, Fajardo and Piper 2011, Walker et al. 2017, Yang et al.

2019). The covariation among traits is due to multiple traits having common functional roles or

developmental pathways, or due to genetic linkages and pleiotropy (Armbruster and Schwaegerle

1996). Examining the trait covariation enhances understanding of plant ecological strategies and

fundamental tradeoffs (e.g., acquisitive – conservative resource use strategy) and can also help to

improve the prediction of one set of traits from others (e.g., root morphology from leaf

morphology) (Westoby et al. 2002, Hu et al. 2019, Yang et al. 2019). However, we don't know

about trait covariation patterns (and plant adaptation strategies) at biochemical levels and if the

structural and biochemical traits covary in the same dimensions.

Considering the above gaps, we used plant genotypes sampled from 11 populations of

Festuca rubra occurring in contrasting climates (climates of origin) and grown in 4 different

growth chambers (cultivation climates) and measured wide range of biochemical and structural

traits of the plants to address the following questions:

1.   What are the major dimensions of variation in structural and biochemical traits (trait co-

variation)?

2.   Do the relative role of climate of origin and cultivation climate and their interactions in

trait variation differ among structural and biochemical traits?

3.   What is the magnitude of variation in structural and biochemical traits in plants from

different climate of origin in response to contrasting cultivation climates?

4.   Is the plasticity in both structural and biochemical traits adaptive, and does the

adaptiveness depend upon climate of origin and cultivation climate?

We hypothesized that 1) some of the structural traits co-vary with biochemical traits (e.g.,

plant biomass and content of photosynthetic pigments), because plant structural traits are the

result of processes occurring at biochemical levels; 2) biochemical traits should be more plastic

(and less genetically controlled) than the structural traits and thus less affected by the climate

of origin than structural traits because of their flexible nature and ease in construction; 3) both

structural and biochemical traits show adaptive plasticity, but because flexible nature and ease

of construction, biochemical traits show a higher degree of adaptive plasticity.

METHODOLOGY

Study species and sampled localities for plant material

In the experiment, we used *Festuca rubra* ssp. *rubra*, a widespread perennial grass

species occurring in the temperate grasslands in Europe. This species can reproduce both by

seeds as well as clonally by rhizomes. The experimental plants of this species were collected in

2014 from 12 natural localities representing factorially crossed precipitation and temperature

gradients corresponding to the SeedClim Grid (Meineri et al. 2014, Tingstad et al. 2015) in

Western Norway. This grid represents an independent combination of three levels of mean

summer temperature (4 warmest months, 6.5 °C, 8.5 °C and 10.5 °C) with four levels of annual

precipitation (600, 1300, 2000 and 2700 mm).

The plant material used in the current study has been collected for previous studies, that

have set up the experiments and provided data also for this study (Münzbergová et al. 2017).

From each locality, at least 40 plants of *F. rubra* (at least 1 m apart) were collected in the

growing season of 2014 and transported to the experimental garden of the Institute of Botany,

Czech Academy of Sciences in Průhonice, Czech Republic (49°59′38.972″N, 14°33′57.637″E)

and planted into pots. To ensure that each of the collected plants corresponded to a single

genotype, the plants were reduced to a single ramet and planted into $16 \times 16 \times 16$ cm pots in 1:2

soil:sand ratio in August 2014. Later, the plants were checked by flow cytometry and confirmed

that all belonged to the hexaploid cytotype of *F. rubra* ssp. *rubra* (Šurinová et al. 2019). In one

of the localities (i.e., 6.5 °C temperature and 1300 mm precipitation), *F. rubra* ssp. *rubra* was

absent. So, finally, we worked with plants from only 11 localities.

In November 2014, 25 genotypes from each locality were transferred to a greenhouse

with temperature between 5 and 10 °C. In the greenhouse, the genotypes were multiplied (for

details see (Münzbergová et al. 2017)). At the end of February 2015, four ramets of each of the

25 genotypes from all the 11 localities were planted in separate pots, resulting in a total of 1100

ramets. One ramet of each genotype has been shifted to one of each 4 different cultivation

climates (see next paragraph) so that each genotype is present in each cultivation climate.

Cultivation climates

The experimental plants were grown in 4 different growth chambers (Votch 1014) under conditions simulating four extreme localities in the SeedClim grid (wettest/driest combined with warmest/coldest) for the spring to summer climate in the field. The temperature and moisture in each of the growth chambers followed the course as in the simulated natural locality (for details, see Münzbergová et al., 2017). The moisture level in the growth chambers was monitored using TMS dataloggers (Wild et al. 2019) (three inside each chamber) and then adding the necessary amount of water to mimic the soil moisture of the natural localities (also monitored using the TMS dataloggers). Briefly, in the climate chambers with dry conditions, plants were watered with about 20 ml of tap water per plant, applied to the trays if the soil moisture was lower than 15%. In the wet regime, plants were cultivated under full soil saturation with ~1.5 cm water level in the tray. By manipulating the soil moisture in the growth chambers, soil moisture at localities with a certain precipitation level was mimicked. Thus, the moisture conditions in the growth chamber were referred to as precipitation values of the simulated localities. Day length and radiation levels inside the growth chambers were also controlled in a manner to mimic the conditions of the original localities (Münzbergová et al. 2017).

Due to the limitation of resources to take measurements of the biochemical traits from all the grown plants (see below), only 10 genotypes per locality were used for the purpose of this study. The other genotypes along with those considered in this study were used in other studies exploring the growth, fitness, and physiological traits of the plants (Münzbergová et al. 2017, Stojanova et al. 2018, Kosová et al. 2022, Thakur and Münzbergová 2022). The dataset used in this study comprised of samples collected from 440 plants (i.e., 11 localities $\times$ 10 genotypes $\times$ 4 cultivation climates = 440 individuals).

Plant trait measurements

From each genotype in each of the cultivation climate, we measured 11 biochemical traits and 9 structural (i.e., morphological) traits. The biochemical traits were represented by the content of antheraxanthin, β-carotene, chlorophyll a, chlorophyll b, phenolic substances, lutein, neoxanthin, superoxide dismutase (SOD), violaxanthin and zeaxanthin and xanthophyll cycle de-epoxidation state (DEPS). These biochemical traits have been estimated specifically for this study. The considered biochemical traits are related to plant photosynthesis, photoprotection and oxidative stress response (Table 1).

Structural traits included aboveground biomass, plant height, rhizome weight, root weight, stomatal density, stomatal size, and number of ramets. Additionally, specific leaf area (SLA) was measured from the 10 genotypes originating from each of the four most extreme climates of the climatic grid. All these structural traits have been also used in our previous studies to address totally different hypotheses (Münzbergová et al. 2017, Kosová et al. 2022, Thakur and Münzbergová 2022). The considered structural traits are linked to plant resource use, competitive ability, fitness, growth and gas exchange (Table 1).

During the experiment, in mid-June, the biomass of the plants has been cut at 3 cm to simulate biomass removal during regular management in the field sites. This biomass was used for trait measurements as described below. The number of ramets was counted and the length (cm) of the longest ramet (hereafter referred to as plant height) was measured at the end of August 2015 and the plants were subsequently harvested, sorted into aboveground parts, roots and rhizomes, dried to a constant mass at 60 °C and weighed.

The remaining traits were measured during biomass removal in mid-June 2015. Stomatal

density and stomatal size (length) were measured on epidermal imprints generated using nail

polish (Gitz & Baker, 2009). The impressions were mounted on microscope slides with

transparent adhesive tape and stomata were counted from three different regions on a leaf (each

$500 \times 500\mu m$). From each of these three areas, three stomata were randomly chosen, and their

length was measured, resulting in nine measured stomata per sample. SLA was estimated as a

ratio of leaf area (mm^2) to dry mass (mg). Leaf area was calculated on scanned folded leaves (as

Festuca leaves are naturally folded and are often hard to unfold) multiplied by two to acquire the

real area of the leaves.

The analyses of chlorophylls and carotenoids followed the protocol described in

(Münzbergová and Haisel 2019). The freshly sampled leaves (collected from the plants in mid-

June 2015) were frozen in liquid nitrogen, lyophilized for 24 h and stored in a freezer (-80 °C)

until the analyses. The contents of chlorophyll (chlorophyll a and chlorophyll b) and carotenoids

(b-carotene, lutein, neoxanthin, violaxanthin, antheraxanthin, and zeaxanthin) in all the samples

were analyzed by High-performance liquid chromatography (HPLC; ECOM, Prague, Czech

Republic) using a reversed-phase column (Watrex Nucleosil 120-5-C18, 5 mm particle size, 125

x 4 mm, ECOM, Prague, Czech Republic). Prior to HPLC injection, pigments were extracted

from the leaves with acetone. In HPLC, elution was carried out for 25 min using a gradient

solvent system acetonitrile/methanol/water (80:12:6) followed by methanol:ethylacetate (9:1),

the gradient was run at the time of 2–5 min. The flow rate was 1 ml min[-1] and the detection

wavelength was 445 nm. Each sample was analyzed twice, and the results were averaged. The

contents of all the pigments were expressed as mg/g of dry weight. The de-epoxidation state

(DEPS) was derived from xanthophyll pigments following (Pospíšilová et al. 2000).

Total content of phenolic compounds was quantified spectrophotometrically from 80%

methanolic extract of dried leaves using modified Folin-Ciocalteu method (Folin and Ciocalteu

1927). The absorbance was measured in 1 cm cuvette at 765 nm. The total phenolic compound

concentration in the analyzed extract was expressed in gallic acid equivalents.

SOD was extracted from frozen leaves homogenized with mortar and pestle in 2.5 ml

buffer (0.1 M Tris-HCl, 1 mM dithiothreitol, 1 mM Na2EDTA, 1% Triton X-100, 5 mM

ascorbic acid, pH 7.8) and a small amount of PVP (polyvinylpyrrolidone). Samples were

incubated on ice in the dark for 30 min and centrifuged (16,000 g, 10 min, 4 °C). Pellets were

discarded, and supernatants were divided into Eppendorf tubes, frozen in liquid nitrogen, and

stored at -70 °C (Lubovska et al., 2014). Later, supernatants were used for gel electrophoresis

and SOD was stained on the gel (Fridovich 1986). These stained gels were scanned and

densitograms were created and analyzed in the ImageJ program (Schneider et al. 2012). The

relative activity of total SOD was estimated as the sum of intensities of bands expressed as peak

areas.

Data analysis

All statistical analyses were performed using R version 4.0.5 (R Development Core Team

2019). Our data were having some missing values (less than 1 % for any trait). Therefore, prior

to analysis, missing values in the data were imputed using the 'missMDA' package (Josse and

Husson 2016). This imputation of missing values was important specifically for doing

multivariate analyses. SLA was estimated only for genotypes from 4 localities (corners of

SeedClim grid), and therefore it was only used in analyses involving traits from genotypes from

only these localities and not used in any of the following analyses involving genotypes from all

the localities.

Trait covariation analysis and dependence of traits on climate

We used principal component analysis (PCA) ['vegan' package (Oksanen et al. 2019)] to

characterize trait covariation patterns and to identify the major dimensions of functional

variation. The number of principal components to retain after PCA was determined on the basis

of randomization-based procedure (Peres-Neto et al. 2005) using rndLambdaF function of

'PCDimension' package (Wang et al. 2018). To analyze the dependence of traits on temperature

and precipitation of original and cultivation climate and their interaction, we used redundancy

analysis (RDA, vegan package) and tested their effects using permutation tests. In this analysis,

all the variables and their interactions had significant effect on traits, but some had only very low

explanatory power. Therefore, in the final model, we only retained the variables which were

significant and explained at least 1% variation in the data. A table showing the change in R^2

upon addition of the given predictor to the model along with significance is shown in

Supplementary Table 1.

Relative role of different factors in variation of structural versus biochemical traits

259 To investigate the relative role of each factor (temperature and precipitation of origin and

260 cultivation climate and their interactions) in trait variation, we determined what fraction of the

261 total variance was explained by each factor independently. This was done for each trait

262 individually by performing variance decomposition analysis using linear mixed-effects model in

263 'lme4' package. In these models, the intercept was the only fixed effect and all the factors (i.e.,

temperature and precipitation of origin and cultivation climate) and their interactions, as well as genotype, were entered as random effects. The variance attributable to each factor was then expressed as a percentage of the total variance. Such models for partitioning the total variance into various components have been widely used (Messier et al. 2017, Isaac et al. 2017, Anderegg et al. 2018, Henn et al. 2018). Before analysis, the traits with non-normal distributions were transformed appropriately (log or cube root).

The variance partitioning results obtained from the analyses above were then used to test if the effect of climate of origin, cultivation climate and interactions between the climate of origin and cultivation differ among trait types. The variance due to climate of origin was taken as the sum of variation explained by original temperature, original precipitation and their interaction (Otemp, Oprec and Otemp $\times$ Oprec in Supplementary Table 1), due to the cultivation climate as variance explained by cultivation temperature, cultivation precipitation and their interaction (Ctemp, Cprec and Ctemp $\times$ Cprec in Supplementary Table 1). The variance due to all the remaining interactions represents the variance due to the interaction of climate of origin and climate of cultivation. The significant effect of trait type on explained variance was analysed using the Mann-Whitney test with explained variance as a response variable.

We also estimated the plasticity of each trait for each genotype across all the four cultivation climates as

Plasticity = (maximum value – minimum value) / maximum value

These plasticity values were then used to test the effect of climate of origin on trait plasticity. The relationship between the climate of origin and plasticity of all the traits was analyzed using RDA with plasticity values as the response variable and climate of origin (temperature and precipitation and their interaction) as explanatory variables.

Trait adaptive potential

To estimate trait adaptive potential, we first performed a PCA analysis of the

standardized trait values separately for biochemical and structural traits and expressed the

position of each genotype (in each growth chamber separately) as its PC scores. For structural

traits, we used scores of the first 3 axes and for biochemical traits, we used scores of the first 2

axes. The number of PCs to be retained was again determined based on randomization-based

method as mentioned earlier. In structural traits, the first three axes of the PCA explained 32%,

18% and 15% of the total variance, respectively. For biochemical traits, the first two axes

explained 65% and 13% of the total variance, respectively. These scores later served as

summarised biochemical and structural traits. We then tested the effects of climate of origin and

cultivation climate and their interaction on these scores using linear mixed-effects models with

genotype as a random factor in 'lme4' package (Bates et al. 2015) in R. For model statistics see

supplementary Table 1. This test has only been done for the 4 extreme populations

corresponding to the climates simulated in the growth chambers and thus also included SLA

values.

Based on these models, we calculated marginal means of the summarised traits for plants

of each origin in each cultivation climate. We then estimated standardized effect size among

pairs of means and tested which pairs of means differed significantly from each other. Both

marginal means and standardized effect sizes were estimated using 'emmeans' package (Searle

et al. 2022) in R. We used this information to make inferences if traits converged or diverged or

none of the two in response to cultivation climate. For example, let us consider two climates A

and B. In case of convergence, when plants originating from A will be grown in B climate their

traits will be more similar to plants of origin B grown in B climate than the traits of plants

originating in A and grown in A. Divergence means the opposite. Therefore, in case of trait

convergence among plants (from different climates of origin) grown in a common climate, there

will be significant shift in the estimated marginal mean of plants grown in a foreign climate

towards that of plants that are grown in their home climate (decreased standardized effect-size).

According to Enquist et al. (2015), phenotypic plasticity may be adaptive if it results in the shift

of trait values toward an optimum that is adaptive for a specific environment, i.e., a trait can be

considered as adaptative if we detect trait convergence, while trait divergence indicates mal-

adaptation.

RESULTS

Trait covariation

The PCA showed that the studied traits are not independent but show covariation

patterns. Our analysis revealed 5 independent dimensions of trait variation and structural traits

varied independently of biochemical traits. The first 5 axes of the PCA accounted for 66% of the

total trait variation (Table 2 and Supplementary Figure 1). The PC 1 alone explained more

variation than the other 4 axes combined and was dominated mainly by biochemical traits related

to photosynthetic machinery (photosynthetic traits as opposed to photoprotective traits). Root

weight also varied along PC1 but orthogonally to biochemical traits (Figure 1). The remaining

PC axes represented traits related to plant fitness and biomass allocation (PC2), gas exchange

(PC3), low wavelength light harvesting (PC4) and plant size (PC5) (details in Table 2 and

Supplementary Figure 1).

Effect of original and cultivation climate on traits

The RDA analysis showed that climatic factors have a significant effect on traits (Supplementary

Table 2A) and explained 37% of their total variation. The first two axes jointly explained 31 %

of the variation with axis 1 explaining 8 times more than axis 2. Precipitation of cultivation

climate (Cprec) was the key variable along with the interaction of Cprec with precipitation of

origin (Oprec) and temperature of cultivation conditions (Ctemp) associated with axis 1 (Figure

2). Ctemp along with Cprec × Ctemp were the variables associated with axis 2. Traits related to

photosynthetic machinery increased towards wetter cultivation climate, whereas the traits related

to photoprotection and biomass of belowground parts tend to have higher values in plants

cultivated in drier climates. The key variable that tends to increase with increasing temperature

of cultivation climate were plant height and aboveground biomass.

Climate change [difference between precipitation and temperature of cultivation and

original climate)] had a significant effect on traits. In the RDA with change in precipitation and

temperature as predictors, 24% of the total variance in traits was explained (Figure 3). Axis 1

explained a lot more (10×) variation than axis 2. Change in precipitation was associated with axis

1 and change in temperature was associated with axis 2. Interaction between change in

precipitation and change in temperature was not associated with any of the first two axes. From

the RDA plot, it is evident that photoprotective pigments and allocation to belowground parts

(roots and rhizomes) increased while photo-harvesting pigments decreased in plants shifted to

drier conditions (Figure 3). Plant height increased in plants shifted to warmer conditions.

Drivers of trait variation in structural versus biochemical traits

The relative role of cultivation climate and its interactions with the climate of origin on

traits was specific to trait type (P<0.05 in the Mann-Whitney test). The variation in structural

traits was significantly higher due to the interaction of climate of origin with cultivation climate

but was significantly lower due to cultivation climate (Figure 4). However, the effect of trait type

on variance explained by the climate of origin was non-significant.

Genotypes from different climates of origin differed in their plasticity. In the RDA, a

total of 20 % of the variability in the plasticity of traits was explained by the climate of origin.

Both the precipitation and temperature of origin as well as their interaction significantly affected

trait plasticity (P<0.05). Biochemical traits (except phenol and SOD) of plants originating from

wet or wet and warm climate (axis 1) tend to be more plastic than biochemical traits of plants

originating from drier climates (Figure 5).

Trait adaptive potential

In structural traits, the overall tendency was to converge (i.e., the traits showed adaptive

potential). PC1 which was loaded with plant fitness traits (i.e., number of ramets and plant

biomass traits) showed the highest degree of convergence (Table 3, Supplementary Figure 2)

while PC2 scores (representing gas exchange traits) mainly showed no change. The results based

on PC3 (representing plant size) also showed either convergence or no change. Biochemical

traits representing PC1 (photoharvesting and photoprotection related traits) tend to diverge

except convergence in cold and wet cultivation climates (Table 3, Supplementary Figure 3) while

those representing PC2 (related to low wavelength light harvesting) tend to converge. Overall,

these results suggest that *F. rubra* shows adaptive plasticity as well as mal-adaptive plasticity in

response to different cultivation climates. However, adaptive plasticity depends upon trait type,

cultivation climate and climate of origin.

DISCUSSION

Independent covariation of biochemical traits and *photoharvesting – photoprotection*

strategy dimension

One of the key findings (contrary to our expectation) of trait covariation analysis was

independent variation of structural and biochemical traits in the multivariate trait space. The

possible reason for the observed patterns (i.e., for decoupling in trait variation among structural

and biochemical traits) could be that plants have distinct strategies at biochemical and structural

levels and differential effects of climatic factors on morphological traits than on biochemical

traits. This lack of covariation among biochemical and structural traits limits our ability to

predict biochemical plant functions from morphological plant functions or *vice-versa*. Our

results, therefore, highlight the importance of incorporating biochemical plant traits (which are

largely ignored) in ecological studies to better understand plant functioning in response to

environmental change.

The second key finding emanating from covariation analysis was covariation between

photosynthesis and photoprotective biochemical traits along the first axis. At the positive end of

this axis, there were individuals with high photoprotective and low photosynthetic pigments, a

syndrome that can be considered as the '*photoharvesting – photoprotection*' strategy dimension

of functional variation. This strategy dimension is inferred following Wright et al (2002) who

proposed the famous leaf economics spectrum ('*conservative-exploitative*' resource use strategy

based on the leaf trait covariation).

The strategy that we propose represents trade-off among leaf biochemical traits related to

photosynthetic machinery and is characterized by an increase in photoharvesting pigments with a

decrease in photoprotective pigments or *vice versa*. Higher amounts of photoprotective pigments

are important in high light conditions for the protection of photosynthetic machinery (Ramírez-

Valiente et al. 2015, Munné-Bosch et al. 2016) and high amounts of photoharvesting pigments

are important in low light conditions to make maximum use of available solar radiation. In this

study, all genotypes were grown under the same light conditions in each cultivation climate

simulating the original light conditions in Norway, hence, light is not the direct driving factor

here. Variation in leaf pigments in this study may be determined by temperature and

precipitation (of both origin and cultivation climates) in addition to light. For instance, plants

growing in higher temperatures and drier conditions have higher chance of photodamage (Taiz et

al. 2015, Szymańska et al. 2017) and hence higher amounts of photoprotective pigments are

needed even if light is unchanged.

Another trade-off evident in our analysis was between stomatal size and stomatal

number. This corroborated with earlier studies (Lawson and Blatt 2014, Kardiman and Raebild

2018). Trade-off between stomatal size and number is linked to plant gas exchange strategy.

Plants with large and few stomata tend to have higher water-use efficiency, but are compromised

with reduced photosynthetic rate, compared with plants that have many but smaller stomata. The

higher gas exchange efficiency of smaller but higher number of stomata per unit area is because

they are easy to regulate (Franks and Beerling 2009, Taiz et al. 2015). Considering this, higher

number of smaller stomata per unit area may be important in drier and warmer climates for

efficiently using water to fix carbon. In this study, stomatal size and plant aboveground biomass

also covaried closely in same direction suggesting a close link between stomatal size and

biomass production. However, currently much less is known about linkage between biomass and

stomatal traits. Smaller and greater number of stomata cause loss of excessive water through

transpiration leading to sub-optimal gas exchange as well as lost opportunities for photosynthesis

(Lawson and Blatt 2014) and could be the possible reason for lower plant biomass (when smaller

but dense stomata are present).

Climate change effects on structural and biochemical traits

Temperature is often considered more important than precipitation in climate change

context (Pepin et al. 2015, Bjorkman et al. 2018, Vitasse et al. 2018), however, our results

suggested that change in precipitation (cultivation – original climate) had stronger effect on plant

traits than change in temperature. Additionally, structural traits are the most used to understand

climate change impacts (Valverde-Barrantes et al. 2017, Bjorkman et al. 2018, Henn et al. 2018,

Yang et al. 2019, Shen et al. 2019) [biochemical in case of a few crop or experimental species

(Renaut et al. 2005, Lei et al. 2006, Hekneby et al. 2006, Pradhan et al. 2019)], and the effects

were stronger in biochemical traits. Especially, photosynthesis related traits were the most

affected by climate change (cultivation – original climate) suggesting that the most important

process leading to primary production in the Biosphere is the most sensitive to climate change.

The greater effect of change in precipitation on photosynthesis related traits could be because

photosynthesis is closely related to gas exchange through stomata and the stomatal opening and

closing depends upon water availability (Taiz et al. 2015, Kardiman and Raebild 2018). Plant

height and plant aboveground biomass increased and root mass decreased with increase in

temperature of cultivation conditions. This can be attributed to lower rate of photosynthesis

(carbon shortage) or direct thermal restriction of tissue formation (Hoch et al. 2002). The

possible reason for decreased allocation to root mass in warmer and wetter cultivation climates

could be linked to increased resource availability [water and nutrient availability in soil, because

of increased mineralization] (Pugnaire et al. 2019). The increased root mass was also linked to

decrease in precipitation of cultivation. This is the expected result as more roots are needed to

meet the water demand in drier climates. Overall, these findings highlight the importance of

studying change in precipitation and incorporating biochemical traits in studies to better

understand climate change impacts on plants.

Interactive effects of climate of origin and cultivation climate are stronger on biochemical

traits

High intraspecific trait variation is considered important in population persistence (Jung

et al. 2010, Bolnick et al. 2011) and can be achieved through genetic differentiation and/or

plasticity (Matesanz et al. 2012). We expected structural traits to be more genetically

differentiated than biochemical traits. In contrast to this, we found that biochemical traits were

more strongly affected by interaction of climate of origin and cultivation climate (i.e., genetic

differentiation in plasticity) than structural traits. This implies that individuals from some plant

populations will be better able to persist under novel climates than others. Genetic controls over

trait variability in plants in both structural (Münzbergová et al. 2017, Laitinen and Nikoloski

2019) and biochemical traits (Benomar et al. 2016, Kreyling et al. 2019) has been already

reported. However, evidence for higher genetic differentiation in plasticity in biochemical traits

than structural traits is not reported previously up to the best of our knowledge. We theorize that

the biochemical traits are directly downstream of gene expression, enable quick response and are

also easy to construct, and therefore, could have higher genetic differentiation in plasticity. On

the other hand, the structural traits are one more step downstream than biochemical traits (i.e.,

structures are product of biochemicals), require more time and lot of energy, thereby leading to

lower genetic differentiation in plasticity in these traits.

Plants originating from climates with higher precipitation or a combination of higher

temperature and precipitation, tend to have higher plasticity in biochemical traits compared to

structural traits. The possible reason is that plants originating from drier and warmer climates

always tend to have high levels of photoprotective pigments (because of higher risk of

photodamage) (Ramírez-Valiente et al. 2015) causing lower plasticity. While in climates with

non-limiting conditions, there is no such requirement and plants show more plastic responses.

However, this needs to be validated using gene regulation studies. The other possible reason

could be attributed to intraspecific trait convergence at stressful climates due to which plants

originating from more limiting climates should have more similar trait values (i.e., lower

plasticity). However, why it is more evident in biochemical traits is still an open question.

Adaptiveness of plasticity depends upon climate (origin and cultivation) and trait type

Adaptive plasticity may facilitate or even speed up the process of adaptive evolution

(Ghalambor et al. 2007). We did find evidence for adaptive as well as mal-adaptive plasticity in

this study. However, we also report that, structural traits showed adaptive plasticity while

biochemical traits largely showed mal-adaptive plasticity. These findings are in line with studies

of (van Kleunen and Fisher 2001, Henn et al. 2018, Kreyling et al. 2019, Laitinen and Nikoloski

2019) suggesting that adaptive as well as mal-adaptive responses in plasticity might be quite

common. Our findings suggest that the adaptiveness is also dependent upon the type of plant

function studied. The possible reason could be that when plants are growing outside their home

climates (i.e., non-optimal climates), they are under stress (Lichtenthaler 1998) which is causing

divergence in biochemical traits. We also believe that greater genetic differentiation in plasticity

in biochemical traits (see Figure 4) might also be limiting adaptive responses in biochemical

traits.

Another important question is, if plants are showing adaptive plasticity in climates that

are predicted for the future. In the context of climate change scenarios of the studied populations

(i.e., in Norway), warming and increased precipitation are predicted (Hanssen-Bauer et al. 2017).

By looking into our findings (related to adaptive plasticity in populations originating from cold

and dry climates and cultivated in warm and wet climates), the species is likely to show adaptive

plasticity in structural traits and will likely show mal-adaptive plasticity in photosynthesis related

traits. Even if the climate change will be in the other direction (i.e., warmer and drier), the

adaptive responses of populations will remain the same.

This combination of adaptive plasticity in structural and mal-adaptive plasticity in

biochemical traits will likely lead to incomplete overall adaptive responses in plants (Ghalambor

et al. 2007) and may have negative consequences to some extent on plant population persistence

in the future (Lande 2009, Chevin et al. 2010). Furthermore, plasticity is also affected by climate

of origin (Münzbergová et al. 2017), which possibly have resulted in plants from different

climates of origin showing adaptive plasticity in some cultivation climates and neutral response

or non-adaptive in others. Considering this, some populations are expected to adapt better than

others to climate change through mechanism of adaptive plasticity. The ability of this species to

adapt to the novel climate is also supported by Münzbergová et al. (2021), demonstrating that

Festuca rubra (i.e., the same model species in the same study system as used here) is capable of

real rapid evolution by selection of genotypes best adapted to novel conditions.

CONCLUSIONS

We conclude that the structural and biochemical plant traits respond differentially to climate. Biochemical traits also represent an important but independent plant strategy linked to plant photosynthetic machinery i.e., *'photoharvesting – photoprotection'* strategy dimension of functional variation. The responses of each trait type to climatic factors are complex and interactive effects of climate of origin and cultivation are very important (different populations will respond in a different way to novel climates). The traits showed adaptive as well as mal-adaptive plasticity, however, adaptiveness also depends upon traits under consideration along with climate (of origin and cultivation). Therefore, studying biochemical traits is important to better understand climate change effects on plants and predict the ability of plants to persist under climate change.

ACKNOWLEDGEMENTS: We thank the POPEKOL discussion group for useful comments on the manuscript draft. The study was supported by the project GACR 19-00522S and institutional research projects RVO 67985939 and MSMT. We also thank Vigdis Vandvik and her team for granting access to the SEEDCLIM climate grid from where the initial material had been obtained. DT also acknowledges "European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant agreement No 101038052" for providing research fellowship.

AUTHOR CONTRIBUTIONS: DT and ZM conceptualised the study. ZM supervised all the work including experiments and secured funding to carry out all the experimental work. DT

analysed the data with extensive comments and suggestions from ZM. VH, RS, HS, DH and NW

collected the experimental data. DT wrote the draft of the paper with extensive suggestions,

inputs and editing from ZM. TD helped in some of the data analysis and provided comments

during manuscript preparation. All the authors commented on and approved the final version of

the manuscript.

CONFLICT OF INTEREST: Nothing to declare

REFERENCES

Anderegg, L. D. L., L. T. Berner, G. Badgley, M. L. Sethi, B. E. Law, and J. HilleRisLambers.
2018. Within-species patterns challenge our understanding of the leaf economics spectrum.
Ecology Letters 21:734–744.

Armbruster, W. S., and K. E. Schwaegerle. 1996. Causes of covariation of phenotypic traits
among populations. Journal of Evolutionary Biology 9:261–276.

Bates, D., M. Machler, B. M. Bolker, and S. C. Walker. 2015. Fitting Linear Mixed-Effects
Models using lme4. Journal of Statistical Software 67:1–48.

Bennett, J. A., K. Riibak, R. Tamme, R. J. Lewis, and M. Pärtel. 2016. The reciprocal
relationship between competition and intraspecific trait variation. Journal of Ecology
104:1410–1420.

Benomar, L., M. S. Lamhamedi, A. Rainville, J. Beaulieu, J. Bousquet, and H. A. Margolis.
2016. Genetic Adaptation vs. Ecophysiological Plasticity of Photosynthetic-Related Traits
in Young Picea glauca Trees along a Regional Climatic Gradient. Frontiers in Plant Science
7:48.

Bjorkman, A. D., I. H. Myers-Smith, S. C. Elmendorf, S. Normand, N. Rüger, P. S. A. Beck, A.
Blach-Overgaard, D. Blok, J. H. C. Cornelissen, B. C. Forbes, D. Georges, S. J. Goetz, K.
C. Guay, G. H. R. Henry, J. HilleRisLambers, R. D. Hollister, D. N. Karger, J. Kattge, P.
Manning, J. S. Prevéy, C. Rixen, G. Schaepman-Strub, H. J. D. Thomas, M. Vellend, M.
Wilmking, S. Wipf, M. Carbognani, L. Hermanutz, E. Lévesque, U. Molau, A. Petraglia, N.
A. Soudzilovskaia, M. J. Spasojevic, M. Tomaselli, T. Vowles, J. M. Alatalo, H. D.
Alexander, A. Anadon-Rosell, S. Angers-Blondin, M. te Beest, L. Berner, R. G. Björk, A.
Buchwal, A. Buras, K. Christie, E. J. Cooper, S. Dullinger, B. Elberling, A. Eskelinen, E. R.
Frei, O. Grau, P. Grogan, M. Hallinger, K. A. Harper, M. M. P. D. Heijmans, J. Hudson, K.
Hülber, M. Iturrate-Garcia, C. M. Iversen, F. Jaroszynska, J. F. Johnstone, R. H. Jørgensen,
E. Kaarlejärvi, R. Klady, S. Kuleza, A. Kulonen, L. J. Lamarque, T. Lantz, C. J. Little, J. D.
M. Speed, A. Michelsen, A. Milbau, J. Nabe-Nielsen, S. S. Nielsen, J. M. Ninot, S. F.
Oberbauer, J. Olofsson, V. G. Onipchenko, S. B. Rumpf, P. Semenchuk, R. Shetti, L. S.

Collier, L. E. Street, K. N. Suding, K. D. Tape, A. Trant, U. A. Treier, J. P. Tremblay, M.
Tremblay, S. Venn, S. Weijers, T. Zamin, N. Boulanger-Lapointe, W. A. Gould, D. S. Hik,
A. Hofgaard, I. S. Jónsdóttir, J. Jorgenson, J. Klein, B. Magnusson, C. Tweedie, P. A.
Wookey, M. Bahn, B. Blonder, P. M. van Bodegom, B. Bond-Lamberty, G. Campetella, B.
E. L. Cerabolini, F. S. Chapin, W. K. Cornwell, J. Craine, M. Dainese, F. T. de Vries, S.
Díaz, B. J. Enquist, W. Green, R. Milla, Ü. Niinemets, Y. Onoda, J. C. Ordoñez, W. A.
Ozinga, J. Penuelas, H. Poorter, P. Poschlod, P. B. Reich, B. Sandel, B. Schamp, S.
Sheremetev, and E. Weiher. 2018. Plant functional trait change across a warming tundra
biome. Nature 562:57–62.

Blanquart, F., O. Kaltz, S. L. Nuismer, and S. Gandon. 2013. A practical guide to measuring
local adaptation. Ecology Letters 16:1195–1205.

Bolnick, D. I., P. Amarasekare, M. S. Araújo, R. Bürger, J. M. Levine, M. Novak, V. H. Rudolf,
S. J. Schreiber, M. C. Urban, and D. Vasseur. 2011. Why intraspecific trait variation matters
in community ecology. Trends Ecol Evol 26:183–192.

Bush, S. E., J. S. Guo, D. Dehn, K. C. Grady, J. B. Hull, E. Johnson, D. F. Koepke, R. W. Long,
D. L. Potts, and K. R. Hultine. 2021. Adaptive versus non-adaptive responses to drought in
a non-native riparian tree / shrub, Tamarix spp. Agricultural and Forest Meteorology 301–
302:108342.

Chevin, L. M., R. Lande, and G. M. Mace. 2010. Adaptation, Plasticity, and Extinction in a
Changing Environment: Towards a Predictive Theory. PLOS Biology 8:e1000357.

Datta, A., I. Kühn, M. Ahmad, S. Michalski, and H. Auge. 2017. Processes affecting altitudinal
distribution of invasive Ageratina adenophora in western Himalaya: The role of local
adaptation and the importance of different life-cycle stages. PLOS ONE 12:e0187708.

Enquist, B. J., J. Norberg, S. P. Bonser, C. Violle, C. T. Webb, A. Henderson, L. L. Sloat, and V.
M. Savage. 2015. Scaling from Traits to Ecosystems: Developing a General Trait Driver
Theory via Integrating Trait-Based and Metabolic Scaling Theories. Advances in Ecological
Research 52:249–318.

Fajardo, A., and F. I. Piper. 2011. Intraspecific trait variation and covariation in a widespread
tree species (Nothofagus pumilio) in southern Chile. New Phytologist 189:259–271.

Folin, O., and V. Ciocalteu. 1927. tyrosine. The Journal of Biological Chemistry 2:627–650.

Franks, P. J., and D. J. Beerling. 2009. Maximum leaf conductance driven by CO_2 effects on
stomatal size and density over geologic time. Proceedings of the National Academy of
Sciences of the United States of America 106:10343–10347.

Fridovich, I. 1986. Superoxide dismutases. Advances in enzymology and related areas of
molecular biology 58:61–97.

Ghalambor, C. K., J. K. Mckay, S. P. Carroll, and D. N. Reznick. 2007. Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. Functional Ecology 21:394–407.

Hanssen-Bauer, I., E. J. Førland, I. Haddeland, H. Hisdal, D. Lawrence, S. Mayer, A. Nesje, S. Sandven, A. B. Sandø, and A. Sorteberg. 2017. Climate in Norway 2100 – a knowledge base for climate adaptation.

Hekneby, M., M. C. Antolín, and M. Sánchez-Díaz. 2006. Frost resistance and biochemical changes during cold acclimation in different annual legumes. Environmental and Experimental Botany 55:305–314.

Henn, J. J., V. Buzzard, B. J. Enquist, A. H. Halbritter, K. Klanderud, B. S. Maitner, S. T. Michaletz, C. Pötsch, L. Seltzer, R. J. Telford, Y. Yang, L. Zhang, and V. Vandvik. 2018. Intraspecific trait variation and phenotypic plasticity mediate alpine plant species response to climate change. Frontiers in Plant Science 871:1548.

Hoch, G., M. Popp, and C. Körner. 2002. Altitudinal increase of mobile carbon pools in Pinus cembra suggests sink limitation of growth at the Swiss treeline. Oikos 98:361–374.

Hu, Y. K., X. Pan, X. J. Yang, G. F. Liu, X. Y. Liu, Y. bin Song, M. Y. Zhang, L. J. Cui, and M. Dong. 2019. Is there coordination of leaf and fine root traits at local scales? A test in temperate forest swamps. Ecology and Evolution 9:8714–8723.

Isaac, M. E., A. R. Martin, E. de M. V. Filho, B. Rapidel, O. Roupsard, and K. van den Meersche. 2017. Intraspecific trait variation and coordination: Root and leaf economics spectra in coffee across environmental gradients. Frontiers in Plant Science 8:1196.

Josse, J., and F. Husson. 2016. missMDA: A package for handling missing values in multivariate data analysis. Journal of Statistical Software 70:1–31.

Jung, V., C. Violle, C. Mondy, L. Hoffmann, and S. Muller. 2010. Intraspecific variability and trait-based community assembly. Journal of Ecology 98:1134–1140.

Kardiman, R., and A. Raebild. 2018. Relationship between stomatal density, size and speed of opening in Sumatran rainforest species. Tree Physiology 38:696–705.

van Kleunen, M., and M. Fisher. 2001. Adaptive evolution of plastic foraging responses in a clonal plant. Ecology 82:3309–3319.

Kosová, V., T. Hájek, V. Hadincová, and Z. Münzbergová. 2022. The importance of ecophysiological traits in response of Festuca rubra to changing climate. Physiologia Plantarum 174.

Kreyling, J., S. J. Puechmaille, A. v. Malyshev, and F. Valladares. 2019. Phenotypic plasticity closely linked to climate at origin and resulting in increased mortality under warming and frost stress in a common grass. Ecology and Evolution 9:1344–1352.

Laitinen, R. A. E., and Z. Nikoloski. 2019. Genetic basis of plasticity in plants. Journal of
Experimental Botany 70:739–745.

Lande, R. 2009. Adaptation to an extraordinary environment by evolution of phenotypic
plasticity and genetic assimilation. Journal of Evolutionary Biology 22:1435–1446.

Lawson, T., and M. R. Blatt. 2014. Stomatal Size, Speed, and Responsiveness Impact on
Photosynthesis and Water Use Efficiency. Plant Physiology 164:1556–1570.

Lei, Y., C. Yin, and C. Li. 2006. Differences in some morphological, physiological, and
biochemical responses to drought stress in two contrasting populations of Populus
przewalskii. Physiologia Plantarum 127:182–191.

Lichtenthaler, H. K. 1998. The stress concept in plants: an introduction. Annals of the New York
Academy of Sciences 851:187–198.

Manzanedo, R. D., M. Fischer, R. María Navarro☐Cerrillo, and E. Allan. 2019. A new approach
to study local adaptation in long☐lived woody species: Virtual transplant experiments.
Methods in Ecology and Evolution 10:1761–1772.

Martin, A. R., B. Rapidel, O. Roupsard, K. van den Meersche, E. de Melo Virginio Filho, M.
Barrios, and M. E. Isaac. 2017. Intraspecific trait variation across multiple scales: the leaf
economics spectrum in coffee. Functional Ecology 31:604–612.

Matesanz, S., T. Horgan-Kobelski, and S. E. Sultan. 2012. Phenotypic Plasticity and Population
Differentiation in an Ongoing Species Invasion. PLoS ONE 7:44955.

McGill, B. J., B. J. Enquist, E. Weiher, and M. Westoby. 2006. Rebuilding community ecology
from functional traits. Trends in Ecology and Evolution 21:178–185.

Meineri, E., O. Skarpaas, J. Spindelböck, T. Bargmann, and V. Vandvik. 2014. Direct and size-
dependent effects of climate on flowering performance in alpine and lowland herbaceous
species. Journal of Vegetation Science 25:275–286.

Messier, J., B. J. McGill, B. J. Enquist, and M. J. Lechowicz. 2017. Trait variation and
integration across scales: is the leaf economic spectrum present at local scales? Ecography
40:685–697.

Mitchell, R. M., G. M. Ames, and J. P. Wright. 2020. Intraspecific trait variability shapes leaf
trait response to altered fire regimes. Annals of Botany:1–10.

Munné-Bosch, S., A. Cotado, M. Morales, E. Fleta-Soriano, J. Villellas, and M. B. Garcia. 2016.
Adaptation of the long-lived monocarpic perennial Saxifraga longifolia to high altitude.
Plant Physiology 172:765–775.

Münzbergová, Z., V. Hadincová, H. Skálová, and V. Vandvik. 2017. Genetic differentiation and
plasticity interact along temperature and precipitation gradients to determine plant
performance under climate change. Journal of Ecology 105:1358–1373.

Münzbergová, Z., and D. Haisel. 2019. Effects of polyploidization on the contents of
photosynthetic pigments are largely population-specific. Photosynthesis Research 140:289–
299.

Münzbergová, Z., V. Vandvik, and V. Hadincová. 2021. Evolutionary Rescue as a Mechanism
Allowing a Clonal Grass to Adapt to Novel Climates. Frontiers in Plant Science 12:706.

Nicotra, A. B., O. K. Atkin, S. P. Bonser, A. M. Davidson, E. J. Finnegan, U. Mathesius, P. Poot,
M. D. Purugganan, C. L. Richards, F. Valladares, and M. van Kleunen. 2010. Plant
phenotypic plasticity in a changing climate. Trends in Plant Science 15:684–692.

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. Mcglinn, P. R. Minchin, R.
B. O'Hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, and H. Wagner. 2019.
vegan: Community Ecology Package. R package version 2.4-2.

Pepin, N., R. S. Bradley, H. F. Diaz, M. Baraer, E. B. Caceres, N. Forsythe, H. Fowler, G.
Greenwood, M. Z. Hashmi, X. D. Liu, J. R. Miller, L. Ning, A. Ohmura, E. Palazzi, I.
Rangwala, W. Schöner, I. Severskiy, M. Shahgedanova, M. B. Wang, S. N. Williamson,
and D. Q. Yang. 2015. Elevation-dependent warming in mountain regions of the world.
Nature Climate Change 2015 5:5 5:424–430.

Peres-Neto, P. R., D. A. Jackson, and K. M. Somers. 2005. How many principal components?
stopping rules for determining the number of non-trivial axes revisited. Computational
Statistics and Data Analysis 49:974–997.

Pérez-Ramos, I. M., L. Matías, L. Gómez-Aparicio, and Ó. Godoy. 2019. Functional traits and
phenotypic plasticity modulate species coexistence across contrasting climatic conditions.
Nature Communications 10:2555.

Pospíšilová, J., D. Haisel, H. Synková, J. Čatský, N. Wilhelmová, Š. Plzáková, D. Procházková,
and F. Šrámek. 2000. Photosynthetic pigments and gas exchange during ex vitro
acclimation of tobacco plants as affected by CO_2 supply and abscisic acid. Plant Cell,
Tissue and Organ Culture 61:125–133.

Pradhan, S., A. K. Goswami, S. K. Singh, J. Prakash, S. Goswami, V. Chinnusamy, A. Talukdar,
and N. K. Maurya. 2019. Low temperature stress induced physiological and biochemical
alterations in papaya genotypes. South African Journal of Botany 123:133–141.

Pugnaire, F. I., J. A. Morillo, J. Peñuelas, P. B. Reich, R. D. Bardgett, A. Gaxiola, D. A. Wardle,
and W. H. van der Putten. 2019. Climate change effects on plant-soil feedbacks and
consequences for biodiversity and functioning of terrestrial ecosystems. Science Advances
5:eaaz1834.

R Development Core Team. 2019. R: A language and environment for statistical computing. R
Foundation for Statistical Computing 1:409.

Ramírez-Valiente, J. A., K. Koehler, and J. Cavender-Bares. 2015. Climatic origins predict
variation in photoprotective leaf pigments in response to drought and low temperatures in
live oaks (Quercus series Virentes). Tree Physiology 35:521–534.

Renaut, J., L. Hoffmann, and J.-F. Hausman. 2005. Biochemical and physiological mechanisms
related to cold acclimation and enhanced freezing tolerance in poplar plantlets. Physiologia
Plantarum 125:82–94.

Schneider, C. A., W. S. Rasband, and K. W. Eliceiri. 2012. NIH Image to ImageJ: 25 years of
image analysis.

Searle, S. R., F. M. Speed, and G. A. Milliken. 2022. Estimated Marginal Means, aka Least-
Squares Means [R package emmeans version 1.7.2]. American Statistician 34:216–221.

Shen, Y., G. S. Gilbert, W. Li, M. Fang, H. Lu, and S. Yu. 2019. Linking Aboveground Traits to
Root Traits and Local Environment: Implications of the Plant Economics Spectrum.
Frontiers in Plant Science 10:1412.

Siefert, A., C. Violle, L. Chalmandrier, C. H. Albert, A. Taudiere, A. Fajardo, L. W. Aarssen, C.
Baraloto, M. B. Carlucci, M. v. Cianciaruso, V. de L. Dantas, F. de Bello, L. D. S. Duarte,
C. R. Fonseca, G. T. Freschet, S. Gaucherand, N. Gross, K. Hikosaka, B. Jackson, V. Jung,
C. Kamiyama, M. Katabuchi, S. W. Kembel, E. Kichenin, N. J. B. Kraft, A. Lagerström, Y.
le Bagousse-Pinguet, Y. Li, N. Mason, J. Messier, T. Nakashizuka, J. M. Overton, D. A.
Peltzer, I. M. Pérez-Ramos, V. D. Pillar, H. C. Prentice, S. Richardson, T. Sasaki, B. S.
Schamp, C. Schöb, B. Shipley, M. Sundqvist, M. T. Sykes, M. Vandewalle, and D. A.
Wardle. 2015. A global meta-analysis of the relative extent of intraspecific trait variation in
plant communities. Ecology Letters 18:1406–1419.

Stojanova, B., M. Šurinová, J. Klápště, V. Koláříková, V. Hadincová, and Z. Münzbergová.
2018. Adaptive differentiation of Festuca rubra along a climate gradient revealed by
molecular markers and quantitative traits. PLoS ONE 13:1–20.

Šurinová, M., V. Hadincová, V. Vandvik, and Z. Münzbergová. 2019. Temperature and
precipitation, but not geographic distance, explain genetic relatedness among populations in
the perennial grass Festuca rubra. Journal of Plant Ecology 12:730–741.

Szymańska, R., I. Ślesak, A. Orzechowska, and J. Kruk. 2017. Physiological and biochemical
responses to high light and temperature stress in plants. Environmental and Experimental
Botany 139:165–177.

Taiz, L., E. Zeiger, I. M. Møller, and A. Murphy. 2015. Plant physiology and development. Page
(L. Taiz and E. Zeiger, Eds.). 6th edition. Sinauer Associates, Sunderland pp. 761.

Thakur, D., and Z. Münzbergová. 2022. Rhizome trait scaling relationships are modulated by
growth conditions and are linked to plant fitness. Annals of Botany 129:529–540.

Tingstad, L., S. L. Olsen, K. Klanderud, V. Vandvik, and M. Ohlson. 2015. Temperature,
precipitation and biotic interactions as determinants of tree seedling recruitment across the
tree line ecotone. Oecologia 179:599–608.

Valladares, F., D. Sanchez-Gomez, and M. A. Zavala. 2006. Quantitative estimation of
phenotypic plasticity: Bridging the gap between the evolutionary concept and its ecological
applications. Journal of Ecology 94:1103–1116.

Valverde-Barrantes, O. J., G. T. Freschet, C. Roumet, and C. B. Blackwood. 2017. A worldview
of root traits: the influence of ancestry, growth form, climate and mycorrhizal association
on the functional trait variation of fine-root tissues in seed plants. New Phytologist
215:1562–1573.

Violle, C., M.-L. Navas, D. Vile, E. Kazakou, C. Fortunel, I. Hummel, and E. Garnier. 2007. Let
the concept of trait be functional! Oikos 116:882–892.

Vitasse, Y., C. Signarbieux, and Y. H. Fu. 2018. Global warming leads to more uniform spring
phenology across elevations. Proceedings of the National Academy of Sciences of the
United States of America 115:1004–1008.

Walker, A. P., M. L. McCormack, J. Messier, I. H. Myers-Smith, and S. D. Wullschleger. 2017.
Trait covariance: the functional warp of plant diversity? New Phytologist 216:976–980.

Wang, M., S. M. Kornblau, and K. R. Coombes. 2018. Decomposing the Apoptosis Pathway Into
Biologically Interpretable Principal Components. Cancer Informatics 17.

Westerband, A. C., J. L. Funk, and K. E. Barton. 2021. Review: Intraspecific trait variation in
plants: a renewed focus on its role in ecological processes. Annals of Botany 127:397–410.

Westoby, M., D. S. Falster, A. T. Moles, P. A. Vesk, and I. J. Wright. 2002. Plant ecological
strategies: Some leading dimensions of variation between species. Annu. Rev. Ecol. Syst
33:125–59.

Wild, J., M. Kopecký, M. Macek, M. Šanda, J. Jankovec, and T. Haase. 2019. Climate at
ecologically relevant scales: A new temperature and soil moisture logger for long-term
microclimate measurement. Agricultural and Forest Meteorology 268:40–47.

Yang, Y., H. Wang, S. P. Harrison, I. C. Prentice, I. J. Wright, C. Peng, and G. Lin. 2019.
Quantifying leaf-trait covariation and its controls across climates and biomes. New
Phytologist 221:155–168.

Figures and tables

Figure 1: Principal component (PC) showing covariation patterns of studied functional traits. Structural traits are underlined. Trait abbreviations are as: Ag.Biom = Aboveground biomass; P.Ht = Plant height; Rhizome.wt = Rhizome weight; Rt.wt = Root weight; Sto.Density = Stomatal density; Sto.Size = Stomatal size; No.Ramets = Number of ramets; Anth. = Antheraxanthin; b-car = β-carotene Chl.a = Chlorophyll a; Chl.b = Chlorophyll b; DEPS = xanthophyll cycle de-epoxidation state; Neox. = Neoxanthin; SOD = Superoxide dismutase; Viox. = Violaxanthin; Zeax. = Zeaxanthin; Phenol = phenolic compounds.

Figure 2: RDA plot representing the effects of cultivation climate and climate of origin on various traits. The names of structural trait are underlined. Ag.Biom = Aboveground biomass; P.Ht = Plant height (cm); Rhizome.wt = Rhizome weight ; Rt.wt = Root weight; Sto.Density = Stomatal density; Sto.Size = Stomatal size; No.Ramets = Number of ramets; Anth. = Antheraxanthin; b-car = β-carotene; Chl.a = Chlorophyll a; Chl.b = Chlorophyll b; DEPS = xanthophyll cycle de-epoxidation state; Neox. = Neoxanthin; SOD = Superoxide dismutase; Viox. = Violaxanthin; Zeax. = Zeaxanthin; Phenol = phenolic compounds. Oprec and Otemp represent precipitation and temperature of climate of origin; Cprec and Ctemp represent precipitation and temperature of cultivation climate.

Figure 3: RDA plot representing effect of climate change (cultivation - origin) on various traits. The names of structural trait are underlined. Ag.Biom = Aboveground biomass; P.Ht = Plant height (cm); Rhizome.wt = Rhizome weight; Rt.wt = Root weight ; Sto.Density = Stomatal density; Sto.Size = Stomatal size; No.Ramets = Number of ramets; Anth. = Antheraxanthin; b-car = β-carotene; Chl.a = Chlorophyll a; Chl.b = Chlorophyll b; DEPS = xanthophyll cycle de-epoxidation state; Neox. = Neoxanthin; SOD = Superoxide dismutase; Viox. = Violaxanthin; Zeax. = Zeaxanthin; Phenol = phenolic compounds. ChangeP and ChangeT represent difference in precipitation and moisture between origin and cultivation climate (cultivation-origin).

Figure 4: Boxplots showing the fraction of total variation explained by climate of origin, cultivation climate and their interaction in biochemical (BC) versus structural (ST) traits. Dark horizontal lines in the plots represent the median value and the grey part in each boxplot represent the interquartile range.

Figure 5: RDA plot representing the effect of climate of plant origin on plasticity in plant traits. The names of structural trait are underlined. Ag.Biom = Aboveground biomass; P.Ht = Plant height (cm); Rhizome.wt = Rhizome weight; Rt.wt = Root weight; Sto.Density = Stomatal density; Sto.Size = Stomatal size; No.Ramets = Number of ramets; Anth. = Antheraxanthin; b-car = β-carotene; Chl.a = Chlorophyll a; Chl.b = Chlorophyll b; DEPS = xanthophyll cycle de-epoxidation state; Neox. = Neoxanthin; Phenol = phenolic compounds; SOD = Superoxide dismutase; Viox. = Violaxanthin; Zeax. = Zeaxanthin. Oprec and Otemp represent precipitation and temperature of climate of origin.

Table 1: List of traits used in this study and their significance. Trait names that are underlined represent structural traits and the remaining represent biochemical traits

Trait	Significance
Aboveground biomass	Biomass allocation to above ground resource acquisition, is related to plant size and is also indicator of plant fitness
Plant height	Plant competitive ability for light
Rhizome weight	Investment into clonal reproduction
Root weight	Biomass allocation to soil resource acquisition
Specific leaf area	Per unit leaf construction cost; related to plant resource use strategy; indicator of plant growth rate
Stomatal density & Stomatal size	Gas exchange and allow plants to adjust their stomatal pore area in response to the surrounding environment.
Number of ramets	Measure of plant fitness and related to plant resource use strategy
Chlorophyll a & b	Key photosynthetic pigments in plants. Chlorophyll a absorb both high (~662 nm) and low (~430nm) wavelength light and chlorophyll b mainly absorbs low wavelength (~453 nm) light
β-carotene	Carotenoids are responsible for the quenching of light and protection of cells from damages caused by light and superoxide radicales
Xanthophylls (Antheraxanthin, Neoxanthin, Violaxanthin, Zeaxanthin and Lutein)	Accessory light-harvesting pigments and plays important role in the protection of photosynthetic machinery from the potentially toxic effects of light
Xanthophyll cycle de-epoxidation state	De-epoxidation state provides information on xanthophyll cycle activity and the degree of violaxanthin to zeaxanthin conversion
Phenolic compounds	Protect against UV radiation by scavenging the reactive oxygen species generated due to light stress.
Superoxide dismutase	It forms the first line of defence against oxidative stresses in plants and plays vital role is scavenging the reactive oxygen species produced during metabolic processes and under stress conditions.

Table 2: Trait loading, eigenvalues and percentage of trait variation explained by the first five principal components (PCs).

	PC1	PC2	PC3	PC4	PC5
Aboveground biomass	-0.34	**1.40**	-0.01	0.72	**1.18**
Plant height	-0.59	0.83	0.33	-0.36	**1.59**
Rhizome weight	**-1.02**	0.10	-0.88	-0.07	-0.32
Root weight	**-1.23**	0.92	-0.67	0.47	-0.36
Stomatal density	-0.15	**-1.27**	-0.96	0.06	0.87
Stomatal size	-0.21	**1.28**	0.66	-0.10	**-1.00**
Number of ramets	-0.32	0.51	**-1.58**	0.77	-0.35
Antheraxanthin	**-1.83**	0.01	-0.18	-0.50	0.07
β-carotene	-0.26	0.41	-0.04	**-1.59**	-0.04
Chlorophyll *a*	**1.97**	0.25	-0.42	-0.34	0.07
Chlorophyll *b*	**1.43**	0.11	-0.62	-0.87	0.22
Xanthophyll cycle de-epoxidation state	**-2.09**	-0.17	-0.09	-0.14	0.00
Phenolic compounds	-0.79	0.41	-0.37	-0.97	-0.14
Lutein	**1.95**	0.17	-0.15	-0.08	-0.01
Neoxanthin	**1.61**	0.44	-0.54	-0.06	0.08
SOD	-0.66	0.05	-0.70	-0.55	0.01
Violaxanthin	**1.73**	0.41	-0.45	-0.12	-0.14
Zeaxanthin	**-2.00**	-0.10	-0.18	-0.19	-0.03
Explained variation (%)	*35.3*	*8.8*	*7.8*	*7.2*	*6.9*

Loadings > 1 in magnitude are in bold

Table 3: Trait convergence or divergence in different populations of *Festuca rubra* when grown in different foreign climates. Inferences were made based upon the significant differences between estimated marginal means between plants in climate of origin (home) and cultivation (foreign) climate. Convergence refers to significant shift in estimated marginal mean (based on PCA scores generated using traits) of plants grown in a cultivation (foreign) climate towards that of plants in climate of origin (home). Conv. = convergence; Div. = divergence; None = neither convergence nor divergence. Cells with '-' represent those plants were in their home climates.

		Cultivation climate			
		Cold and dry	Warm and dry	Cold and wet	Warm and wet
Climate of Origin		**Structural traits (PC1)**			
	Cold and dry	-	None	Conv.	Conv.
	Warm and dry	None	-	Conv.	Conv.
	Cold and wet	Conv.	Conv.	-	Conv.
	Warm and wet	None	Conv.	None	-
		Structural traits (PC2)			
	Cold and dry	-	None	None	None
	Warm and dry	None	-	None	None
	Cold and wet	None	None	-	None
	Warm and wet	Conv.	None	None	-
		Structural traits (PC3)			
	Cold and dry	-	None	Conv.	None
	Warm and dry	None	-	None	None
	Cold and wet	Conv.	Conv.	-	Conv.
	Warm and wet	Conv.	None	None	-
		Biochemical traits (PC1)			
	Cold and dry	-	None	Conv.	Div.
	Warm and dry	Div.	-	Conv.	Div.
	Cold and wet	Div.	Div.	-	None
	Warm and wet	Div.	Div.	Conv.	-
		Biochemical traits (PC2)			
	Cold and dry	-	None	Conv.	Conv.
	Warm and dry	Conv.	-	Conv.	Conv.
	Cold and wet	Conv.	Conv.	-	Conv.
	Warm and wet	None	Conv.	None	-

Supplementary Figure 1: Principal component (PC) patterns of studied functional traits. The axes 3 to 5 are plotted against axis 1. Trait abbreviations are as: Ag.Biom = Aboveground biomass; P.Ht = Plant height; Rhizome.wt = Rhizome weight ; Rt.wt = Root weight ; Sto.Density = Stomatal density; Sto.Size = Stomatal size; No.Ramets = Number of ramets; Anth. = Antheraxanthin; b-car = β-carotene Chl.a = Chlorophyll a; Chl.b = Chlorophyll b; DEPS = xanthophyll cycle de-epoxidation state; Neox. = Neoxanthin; SOD = Superoxide dismutase; Viox. = Violaxanthin; Zeax. = Zeaxanthin; Phenol = phenolic compounds.

Supplementary Figure 2: Estimated marginal means of PC axis 1 to 3 (based on trait values of structural traits) for different populations cultivated in contrasting climates. The shaded areas are 95% confidence intervals. Means in large blue dots are for populations grown in home climate. The arrows on shaded areas represent direction of shift in traits (from home climate) along with convergence (blue arrows).

Supplementary Figure 3: Estimated marginal means of PCA axis 1 and 2 scores (based on trait values of biochemical traits) for different populations cultivated in contrasting climates. The shaded areas are 95% confidence intervals. Means in large blue dots are for populations grown in home climate. The arrows on shaded areas represent direction of shift in traits (from home climate) along with convergence (blue arrows) and divergence (red arrows).

Supplementary Table 1: Mixed effects model statistics showing the effect of climate of origin and cultivation climate on PC scores based structural and biochemical traits separately. Significant effects are shown in bold. Oprec and Otemp represent precipitation and temperature of climate of origin; Cprec and Ctemp represent precipitation and temperature of cultivation climate.

Predictor	Structural traits						Biochemical traits			
	PC1		PC2		PC3		PC1		PC2	
	F value	P value	F value	P value	F value	P value	F value	P value	F value	P value
Otemp	8.30	**0.007**	14.15	**<0.001**	10.98	**0.002**	628	**<0.001**	91.6	**<0.001**
Oprec	1.90	0.177	3.27	0.079	0.06	0.812	1.6	0.207	0.3	0.614
Ctemp	31.24	**<0.001**	23.1	**<0.001**	115.3	**<0.001**	628	**<0.001**	91.6	**<0.001**
Cprec	308.2	**<0.001**	6.07	**0.015**	1.03	0.313	25.5	**<0.001**	13.6	**<0.001**
Otemp:Oprec	0.02	0.881	23.9	**<0.001**	0.12	0.735	0.6	0.425	0.4	0.542
Otemp:Ctemp	0.17	0.683	0.10	0.749	0.01	0.942	628.1	**<0.001**	91.6	**<0.001**
Otemp:Cprec	3.43	0.067	0.31	0.577	0.56	0.457	26.3	**<0.001**	13.6	**<0.001**
Oprec:Ctemp	0.63	0.431	2.88	0.093	1.89	0.172	7.4	**0.008**	5.1	**0.026**
Oprec:Cprec	0.02	0.884	0.15	0.695	1.26	0.265	24.5	**<0.001**	5.7	**0.019**
Ctemp:Cprec	11.97	**<0.001**	0.38	0.540	34.10	**<0.001**	30.7	**<0.001**	5.9	**0.017**
Otemp:Oprec:Ctemp	4.54	0.035	0.88	0.351	3.88	0.052	0	0.939	2.4	0.126
Otemp:Oprec:Cprec	1.46	0.230	0.14	0.704	0.22	0.641	19.5	**<0.001**	4.7	**0.032**
Otemp:Ctemp:Cprec	0.00	0.963	1.48	0.227	3.76	0.055	53.7	**<0.001**	14.3	**<0.001**
Oprec:Ctemp:Cprec	0.49	0.487	0.38	0.539	0.56	0.454	10.4	**0.002**	9.9	**0.002**
Otemp:Oprec:Ctemp:Cprec	8.01	**0.006**	0.01	0.940	0.03	0.872	79.8	**<0.001**	16.1	**<0.001**