1
2
3
Differential effects of ankle constraints on foot placement control between normal and split belt treadmills

Mayra Hos, Lieke van Iersel, Moira van Leeuwen, Sjoerd M. Bruijn

Corresponding author at:

Moira van Leeuwen

moiravleeuwen@gmail.com

Keywords: gait stability, Splitbelt, foot placement control, step width

Abstract

Mediolateral ankle moment control contributes to gait stability. Therefore, constraining ankle

moment control might affect other stability mechanisms, including foot placement control. Ankle

moments can be constrained by walking with a shoe with a ridge underneath the sole, narrowing the

mediolateral support surface. Based on the literature, it seems that constraining ankle moment control

in this fashion results in an increased step width and a decrease in the degree of foot placement

control, as defined by the percentage of variance in foot placement that can be explained by CoM

state. However, since that study was performed on a splitbelt treadmill and the narrow ridge could fit

inside the gap between the belts, it is not evident whether these effects can be attributed to the

constrained ankle moment control or to avoidance of this gap. Therefore, we investigated if the effects

of ankle moment constraints are dependent on whether participants walk on a normal treadmill or a

splitbelt treadmill. We included fourteen healthy young adults. Walking with constrained ankle

moment control resulted in a wider step width on both treadmills. Yet, the increase in step width was

larger on the splitbelt treadmill compared to on the normal treadmill. Moreover, we only found a

decreased degree of foot placement control on the splitbelt, whilst the degree of foot placement

control increased on the normal treadmill. We conclude that the effects of ankle moment constraints

reported in our previous study were confounded by the use of a splitbelt. For future research, we

recommend using a normal treadmill whenever possible, because the gap in a splitbelt treadmill might

affect gait parameters.

1. Introduction

In bipedal locomotion, the Centre of Mass (CoM) is relatively high above a relatively small and continuously changing Base of Support (BoS). This makes stability control complex, as a perturbation can rapidly result in instability (Bruijn & Van Dieën, 2018; Reimann et al., 2018). The vertical projection of the CoM remains medial from the BoS during single-legged support. To prevent falling, the legs must be placed accurately to accommodate changes in the CoM kinematic state (Reimann et al., 2018; Winter et al., 1990).

The changing of the base of support via foot placement has been coined the foot placement strategy (Bruijn & Van Dieën, 2018; Hof et al., 2007; Reimann et al., 2018). When increasing step width, larger changes of CoM movement can be accommodated (Lencioni et al., 2020; Reimann et al., 2018). Moreover, maintaining a high degree of step-by-step foot placement control with respect to variations in CoM kinematic state ensures stability, also at narrower step widths (Mahaki et al., 2019; Perry & Srinivasan, 2017). In addition to the foot placement strategy, individuals can control their ankle moment so as to shift the center of pressure (CoP) under the foot during the stance phase. Through the displacement of the CoP, the ground reaction force (GRF) provides a more beneficial torque with respect to the CoM, compensating for "errors" in foot placement (Bruijn & Van Dieën, 2018; Hof et al., 2007; Reimann et al., 2018; van Leeuwen et al., 2021, 2022).

In recent work, we (Van Leeuwen et al., 2020) investigated the effects of constraining ankle moment control by a shoe (LesSchuh) with a narrow, yet flexible, ridge underneath the shoes' soles (somewhat like skates). Constraining ankle moment control with LesSchuh resulted in an increased step width and a decreased degree of foot placement control (as defined by the percentage of foot placement that can be explained from the CoM state). However, along with the constrained ankle moment control, these results might have been affected by fear of stepping in the gap between the belts of the splitbelt the study was performed on.

In the current study, we investigated whether the effects we reported earlier were confounded by the use of a splitbelt. Therefore, we compared conditions on a normal treadmill and on a splitbelt treadmill. Based on the literature, we expected that constraining ankle moment control would result in an increased step width and a diminished degree of foot placement control, both when walking on a normal and on a splitbelt treadmill.

2. Methods

2.1. participants

63   Fifteen young adults, without injuries or other constraints which might affect gait, participated

64 in this study. Data of one participant were not included in the statistical analyses, due to measurement

65 errors. The remaining participants (3 men, 11 women, age 22.4 (2.9) years (mean(sd)), weight 67.3

66 (12) kg, length 1.72 (0.08) m) are included in the statistical analysis. All participants signed informed

67 consent, and the protocol was approved by the local ethical review board (VCWE-2019-108).

68 2.2. Research design

69   Participants walked on two different treadmills (Splitbelt, C-Mill; Motek-Forcelink,

70 Amsterdam, Netherland) with two types of shoes (regular, LesSchuh), making for a total of four

71 conditions (see table 1). The Splitbelt is a treadmill with a small gap between the two belts. The C-mill

72 is a regular treadmill. The Regular shoe is a sneaker with a flat sole. LesSchuh is the same kind of

73 sneaker, however, a narrow ridge (one centimeter) is attached to the sole, see Figure 1 (van Leeuwen

74 et al., 2020, p. 5). This ridge constrains ankle moment control during walking.

76 *Table 1.*

77 Conditions.

	Description	Symbol
Condition 1	Splitbelt + Regular shoe	▌▌◖
Condition 2	Splitbelt + LesSchuh	▌▌◗
Condition 3	C-mill + Regular shoe	▌ ◖
Condition 4	C-mill + LesSchuh	▌ ◗

Figure 1. LesSchuh. The narrow ridge has a width of one centimeter and bends along with the sole, allowing for normal anterior-posterior roll-off, yet constraining mediolateral center-of-pressure modulation. Reprinted from "Active foot placement control ensures stable gait: Effect of constraints on foot placement and ankle moments", by A.M. van Leeuwen, J.H. van Dieën, A. Daffertshofer and S.M. Bruijn (2020), *PLOS ONE, 15*(12), p. 5.

2.3. Procedure

First, the preferred step frequency was determined while walking at a speed of 1.25 × √(leg length [m]) m/s (Hof, 1996). During all subsequent trials, subjects walked at this step frequency, as dictated by a metronome.

For both treadmills we used the same set-up (see Figure 2), consisting of the treadmill and two Optotrak camera arrays (Northern Digital Inc, Waterloo Ontario, Canada), which recorded kinematics of three cluster markers attached to the thorax and the back of both shoes (sampling rate 50 Hz). Before the measurements, anatomical landmarks were defined with a pointer.

Figure 2. Experimental Set-up Splitbelt. The red circles in this figure point out the Optotrak camera arrays.

Each condition lasted twelve minutes, of which the first two minutes were regarded as warming-up. We used counterbalanced randomization for the conditions. On both treadmills

participants started on the same shoes. During the measurement period, we informed the participant
of the remaining walking time at five and nine minutes. Between every condition, a rest period was
included. During this period, participants were asked to change their shoes.

2.4. Data processing

Using the heel markers' vertical position and velocity, we determined the moments of heel
strike and toe-off (Pijnappels et al., 2001). Midstance was defined as the moment between heel strike
and toe-off. At midstance, we calculated step width as the distance between the calcanei of the two
feet.

The degree of foot placement control was calculated as the relative explained variance of a
regression model which relates mediolateral foot placement to mediolateral CoM position and
velocity, in accordance with (Bruijn & van Leeuwen, 2020; van Leeuwen et al., 2020; Wang &
Srinivasan, 2014). The data and code for analysis will be published on Zenodo if the manuscript is
accepted, and is now available at https://surfdrive.surf.nl/files/index.php/s/SyoYsw2IspBSdmn.

2.5. Statistics

A power analysis (α = .05; statistical power = .80) indicated that a minimum of nine participants
were required for this study (effect size for step width in data of (van Leeuwen et al., 2020a)= 0.83).

We conducted repeated measures ANOVAs with factors Shoe [Regular, LesSchuh] and
Treadmill [Splitbelt, C-mill] for step width and foot placement control. All statistical analyses were
performed with Jamovi version 1.6.23.0 (The jamovi project, 2021), and $p<0.05$ was considered
significant.

3. Results

For step width, we found significant effects of Treadmill, Shoe as well as their interaction (see
Table 2). The interaction showed a greater effect of LesSchuh on the Splitbelt, compared to on the C-
mill (see Figure 3).

For R^2, we found a significant main effect for Treadmill and a significant interaction effect (see
table 2). The interaction shows opposite effects of ankle moment constraint for the Splitbelt and the
C-Mill. On the Splitbelt, LesSchuh decreased the degree of foot placement control compared to walking
on regular shoes ($p<0.05$), whereas on the C-mill the degree of foot placement control increased when
walking with LesSchuh ($p<0.01$) (see Figure 4).

Results of statistical tests. Significant effects are displayed in **bold**.

		Treadmill	Shoe	Treadmill*Shoe
Step width	$F(1,13)$	108.9	27.6	15.3
	p	**< 0.001**	**< 0.001**	**0.002**
R^2	$F(1,13)$	49.969	0.492	0.496
	p	**< 0.001**	0.496	**< 0.001**

Figure 3. Effects of Treadmill and Shoe on step width. The grey dots and lines represent the individual data, bars represent

averages.

Figure 4. Effects of Treadmill and Shoe on R^2. The grey dots and lines represent the individual data, bars represent averages.

4. Discussion

In a recent study, we concluded that constrained ankle moment control resulted in a greater
step width and a decreased degree of foot placement control (Van Leeuwen et al., 2020). However,
that study was performed on a Splitbelt, and thus, the observed effect might be partially explained by
the gap between the belts. To investigate whether constraining ankle moment control affects step
width and foot placement control, regardless of treadmill type, we conducted our measurements on
a Splitbelt and a C-mill (normal treadmill).

We hypothesized that constraining ankle moments, by walking on LesSchuh, results in an
increased step width and a decreased degree of foot placement control, when compared to walking
on regular shoes, regardless of the treadmill.

We found a significant wider step width while walking on LesSchuh, compared to walking on
the regular shoe. This suggests that maintaining stability during walking is more difficult with
constrained ankle moment control. As errors in foot placement control cannot be corrected for by
ankle moment control anymore, increasing step width could serve as a compensation strategy (van
Leeuwen et al., 2021). This is in line with our hypothesis. We also found a greater effect of LesSchuh
on the Splitbelt compared to the C-mill. This might be explained by fear of stepping in the gap while
walking on LesSchuh on the Splitbelt. Stepping in the gap with LesSchuh leads to a significant
perturbation. To compensate, the walking strategy of the participants might have been adjusted,
resulting in the greater step width. After the experiment, participants sometimes reported that they
were afraid to step in the gap (note that we did not explicitly assess fear).

As expected from previous work, we also found a decreased degree of foot placement control
when walking with LesSchuh, but only on the Splitbelt. On the C-mill, walking on LesSchuh resulted in
an increased degree of foot placement control. These results are not in line with our hypothesis,
because we predicted a decreased degree of foot placement control, regardless of the treadmill (Van
Leeuwen et al., 2020). In the previous study, we explained the lower degree of foot placement control
as indicative of how ankle moment control may not only complement foot placement by correcting
errors after foot placement, but also is an intrinsic part of foot placement control itself (Zhang et al.,
2020). Thus, we attributed the decrease in foot placement control to perturbed ankle moments
effectively perturbing foot placement control. However, even in our previous study, we had actually
hypothesized an increased degree of foot placement control when walking with LesSchuh. We
reasoned that without ankle moment control to correct for foot placement errors, participants would

compensate by tightening their degree of foot placement control. The latter seemed to have occurred for our current C-mill results, showing that constrained ankle moment control increases the degree of foot placement control. Clearly, the effects we previously reported were confounded by the gap between the belts. The additional task constraint of avoiding the gap, is likely to have led to the decreased foot placement control. This is in line with a lower degree of foot placement control when subjects are instructed to walk on lines projected on a treadmill (Perry & Srinivasan, 2017). Here we provide further evidence that when an additional task constraints are imposed on foot placement, the degree of foot placement control with respect to the CoM is compromised.

4.1. Limitations

A limitation of this study is that, apart from adjusting step width and foot placement control, other compensation strategies may have been used to maintain stability while walking on LesSchuh. One such strategy may have been increasing stride frequency, and indeed walking on LesSchuh resulted in a small but significant increased stride frequency (on average 0.004 Hz) (see Supplementary figure 1, supplementary table 1) (Hak et al., 2013; van Leeuwen et al., 2020b). Additionally, walking on the splitbelt resulted in a significant decrease in stride frequency (on average 0.005 Hz). Toeing out may have been another strategy that subjects may have used. Indeed, walking on LesSchuh resulted in significantly more toeing-out (on average 1.9˚) (see Supplementary figure 2, supplementary table 1) (Hoogstad et al., 2022; Rebula et al., 2017). Also, walking on the splitbelt resulted in a significant increase in toeing-out (on average 2.9˚). Because of these small differences, we expect that the adjustments in stride frequency and toeing-out do not significantly influence our results. Lastly, despite the instructions to walk on the beam of LesSchuh, this might not have happened continuously.

4.2. Practical relevance & recommendations

Previous work has suggested that LesSchuh may be used as a perturbation of foot placement control, and that the aftereffects after walking with such a perturbation can be used as training (Heitkamp et al., 2019; Hoogstad et al., 2022; Reimold et al., 2020). However, the current results show that walking with LesSchuh can only be regarded as a perturbation to foot placement control if subjects walk on a split belt treadmill. During walking on a normal treadmill, foot placement control during walking with LesSchuh actually improved (which was the original hypothesis in our earlier work). It remains to be seen whether the latter adaptation maintains as an aftereffect upon returning to normal shoes and as such whether LesSchuh has a training potential on a normal treadmill. Like LesSchuh on a normal treadmill, an assistive force field improved the degree of foot placement control during application (Heitkamp et al., 2019). Yet, an aftereffect of walking in this assistive force field was a diminished degree of foot placement control compared to normal walking. Still, LesSchuh may have a

training potential on a normal treadmill. Walking with LesSchuh on a normal treadmill would promote
increased active foot placement control by constraining a correction mechanism for foot placement
errors, rather than decreased active control due to external assistance. Therefore, if any aftereffects
after walking with LesSchuh on a normal treadmill would occur, we speculate them to be more similar
to those after application of a perturbing force field (i.e. a higher degree of foot placement control as
an aftereffect) (Reimold et al., 2020).

For future research investigating mechanisms stabilizing gait, we recommend using a normal
treadmill whenever possible. Our study showed that walking on the splitbelt affected step width and
foot placement control.

5. Conclusion

Walking with constrained ankle moment control resulted in a wider step width on both
treadmills and a decreased foot placement control on the splitbelt and an increased foot placement
control on the normal treadmill. The increase in step width was larger on the splitbelt, compared to a
normal treadmill. We conclude that the effect reported in our previous study were confounded by the
use of a splitbelt (Van Leeuwen et al., 2020).

206 6. References

Bruijn, S. M., & van Leeuwen, A. M. (2020). SjoerdBruijn/FootPlacement: Initial Release (V1.0). *Zenodo*.
https://doi.org/https://doi.org/10.5281/zenodo.4292212

Bruijn, Sjoerd M., & Van Dieën, J. H. (2018). Control of human gait stability through foot placement.
Journal of the Royal Society Interface, *15*(143), 1–11. https://doi.org/10.1098/rsif.2017.0816

Hak, L., Houdijk, H., Beek, P. J., & Van Dieën, J. H. (2013). Steps to take to enhance gait stability: The
effect of stride frequency, stride length, and walking speed on local dynamic stability and margins
of stability. *PLoS ONE*, *8*(12), e82842. https://doi.org/10.1371/journal.pone.0082842

Heitkamp, L. N., Stimpson, K. H., & Dean, J. C. (2019). Application of a Novel Force-Field to Manipulate
the Relationship between Pelvis Motion and Step Width in Human Walking. *IEEE Transactions on
Neural   Systems   and   Rehabilitation   Engineering*,   *27*(10),   2051–2058.
https://doi.org/10.1109/TNSRE.2019.2941372

Hof, A. L. (1996). Scaling gait data to body size. *Gait and Posture*, *3*(4), 222–223.

Hof, A. L., van Bockel, R. M., Schoppen, T., & Postema, K. (2007). Control of lateral balance in walking.
Experimental findings in normal subjects and above-knee amputees. *Gait and Posture*, *25*(2),

250–258. https://doi.org/10.1016/j.gaitpost.2006.04.013

Hoogstad, L. A., van Leeuwen, A. M., van Dieën, J. H., & Bruijn, S. M. (2022). Can foot placement during
gait be trained? Adaptations in stability control when ankle moments are constrained. *Journal of*
Biomechanics, *134*, 110990. https://doi.org/10.1016/j.jbiomech.2022.110990

Lencioni, T., Carpinella, I., Rabuffetti, M., Cattaneo, D., & Ferrarin, M. (2020). Measures of dynamic
balance during level walking in healthy adult subjects: Relationship with age, anthropometry and
spatio-temporal gait parameters. *Proceedings of the Institution of Mechanical Engineers, Part H:*
Journal of Engineering in Medicine, *234*(2), 131–140.
https://doi.org/10.1177/0954411919889237

Mahaki, M., Bruijn, S. M., & Van Dieën, J. H. (2019). The effect of external lateral stabilization on the
use of foot placement to control mediolateral stability in walking and running. *PeerJ*, *7*, e7939.
https://doi.org/10.7717/peerj.7939

Perry, J. A., & Srinivasan, M. (2017). Walking with wider steps changes foot placement control,
increases kinematic variability and does not improve linear stability. *Royal Society Open Science*,
4(9), 160627. https://doi.org/10.1098/rsos.160627

Pijnappels, M., Bobbert, M. F., & Van Dieën, J. H. (2001). Changes in walking pattern caused by the
possibility of a tripping reaction. *Gait and Posture*, *14*(1), 11–18. https://doi.org/10.1016/S0966-
6362(01)00110-2

Rebula, J. R., Ojeda, L. V., Adamczyk, P. G., & Kuo, A. D. (2017). The stabilizing properties of foot yaw
in human walking. *Journal of Biomechanics*, *53*, 1–8.
https://doi.org/10.1016/j.jbiomech.2016.11.059

Reimann, H., Fettrow, T., & Jeka, J. J. (2018). Strategies for the control of balance during locomotion.
Kinesiology Review, *7*(1), 18–25. https://doi.org/10.1123/kr.2017-0053

Reimold, N. K., Knapp, H. A., Henderson, R. E., Wilson, L., Chesnutt, A. N., & Dean, J. C. (2020). Altered
active control of step width in response to mediolateral leg perturbations while walking. *Scientific*
Reports, *10*(1), 1–15. https://doi.org/10.1038/s41598-020-69052-5

van Leeuwen, A. M., van Dieën, J. H., & Bruijn, S. M. (2022). The effect of external lateral stabilization
on the control of. *BioRxiv*. https://doi.org/https://doi.org/10.1101/2022.04.29.490037

van Leeuwen, A. M., van Dieën, J. H., Daffertshofer, A., & Bruijn, S. M. (2020a). Dataset and analyses
for: Active foot placement control ensures stable gait: Effect of constraints on foot placement

and ankle moments [Data set]. *Zenodo*.
https://doi.org/https://doi.org/10.5281/zenodo.4229851

van Leeuwen, A. M., van Dieën, J. H., Daffertshofer, A., & Bruijn, S. M. (2020b). Active foot placement
control ensures stable gait: Effect of constraints on foot placement and ankle moments. *PLOS*
ONE, *15*(12), 1–19. https://doi.org/10.1371/journal.pone.0242215

van Leeuwen, A. M., van Dieën, J. H., Daffertshofer, A., & Bruijn, S. M. (2021). Ankle muscles drive
mediolateral center of pressure control to ensure stable steady state gait. *Scientific Reports*,
11(1), 1–14. https://doi.org/10.1038/s41598-021-00463-8

Wang, Y., & Srinivasan, M. (2014). Stepping in the direction of the fall: The next foot placement can be
predicted from current upper body state in steady-state walking. *Biology Letters*, *10*(9), 1–5.
https://doi.org/10.1098/rsbl.2014.0405

Winter, D. A., Patla, A. E., & Frank, J. S. (1990). Assessment of balance control in humans. *Medical*
Progress through Technology, *16*(1–2), 31–51.

Zhang, Y., Smeets, J. B. J., Duysens, J., Brenner, E., & Verschueren, S. (2020). Fast responses to stepping-
target displacements when walking. *The Journal of Physiology*, *598*(10), 1987–2000.
https://doi.org/10.1113/JP278986

Supplementary material

Supplementary Figure 1. Stride frequency. The grey dots and lines represent the individual data, bars represent the average.

Supplementary Figure 2. Toeing out. The grey dots and lines represent the individual data, bars represent the average.

Supplementary Table 1.

Results of statistical tests. Significant effects are displayed in **bold**.

		Treadmill	Shoe	Treadmill*Shoe
Stride frequency	F(1,13)	5.416	7.870	0.676
	p	**0.015**	**< 0.05**	0.426

Toeing out	F(1,13)	28.835	27.921	0.379
	p	**< 0.001**	**< 0.001**	0.549