Community evenness and sample size affect estimates of predation intensity and prey

selection: A model-based validation.

Madhura Bhattacherjee, Devapriya Chattopadhyay*

RRH: COMMUNITY STRUCTURE AFFECTS INFERRED PREDATION

LRH: BHATTACHERJEE AND CHATTOPADHYAY

Keywords: Drilling predation, durophagous predation, evenness, sampling, selective

predation

* Corresponding author

Email: devapriya@iiserpune.ac.in

Abstract. –

Predation estimates inferred from the preserved records of predation traces are

essential in evaluating the evolutionary effect of ecological interactions. It is, however,

crucial to establish how sampling intensity and community composition of an assemblage

influence the reliability of these measures.

Using a resampling technique, we evaluated the effect of sampling intensity and a

community's evenness on the inferred predation estimates. We theoretically simulated

model communities representing different levels of evenness, predation intensity, and

predatory behavior (selective, non-selective). We calculated the total predation intensity and

the number of prey species for each community. We then resampled each community

without replacement and noted variations in the inferred measure from the accurate estimate

as the sampling intensity increased. Our results demonstrate that the evenness of a

community does not influence the inferred predation intensity for non-selective predation.

However, communities with highly selective predation are sensitive to evenness and

sampling intensity; inferred predation intensity of these assemblages can substantially differ

from the actual value. The inferred number of prey species is also influenced by the

community's original evenness, predation selectivity, and predation intensity. When

predation is selective, sampling intensity heavily influences communities with low evenness

and low predation intensity; inferred predation intensity is underrepresented in smaller

sample sizes. For communities of low evenness and predation intensity where rare species

are attacked preferentially, the inferred prey richness differs significantly at a small sample

size.

We proposed a post-facto standardization method for comparing predation estimates

of discrete communities that differ in the sample size. We validated its utility using the

published predation data of the Plio-Pleistocene molluscan fossil assemblage. The present

approach attempts to provide critical insight into the reliability of predation estimates and

may help in comparing predation patterns across time and space. Several factors, including

preservation bias, might impact the final predation signature of an assemblage. It warrants a

future research direction to develop a comprehensive framework of post-hoc standardization

of assemblages with differing predation styles and preservation history.

Madhura Bhattacherjee, Department of Earth and Climate Science, IISER Pune, Dr. Homi

Bhaba Road, Pashan, Pune-411008, INDIA

Devapriya Chattopadhyay, Department of Earth and Climate Science, IISER Pune, Dr.

Homi Bhaba Road, Pashan, Pune-411008, INDIA, Email: devapriya@iiserpune.ac.in

Introduction

The role of predation in shaping the marine ecosystems through time has been a common

theme of study (Vermeij 1977; Vermeij et al. 1981; Signor and Brett 1984; Langerhans

2007; Stanley 2008; Barnes et al. 2010; Gorzelak et al. 2012; Kotta et al. 2018; Petsios et al.

2021). The relationship between the prey and predator is complex in theoretical terms

posing a challenge in predicting the evolutionary outcome of predation (DeAngelis et al.

1975; Berryman 1992; Haque 2012; Abrams 2015). For evaluating the evolutionary effects

of predation, researchers rely on the deep time record of predation (Kitchell and Kitchell

1980; Vermeij et al. 1981; Kelley and Hansen 1993; Vermeij 1993; McNamara 1994;

Kowalewski et al. 2005; Huntley and Kowalewski 2007; Baumiller et al. 2010; Klompmaker

et al. 2017; Bicknell and Paterson 2018). Therefore, the accurate estimation of predation

measures is of primary importance to studies of predator-prey systems.

For establishing predation events and inferring predation intensities, ecological

studies use direct observations or indirect measures such as compositional characterization

of digested food and fecal matter (Nilsen et al. 2012; Pringle et al. 2019). Although it is

possible to recover direct observational evidence of predation events in past ecosystems by

studying "caught-in-the-act" occurrences (Ehret et al. 2009; Ebert et al. 2015),

paleoecological studies primarily rely on preserved predation traces, such as drill holes and

repair scars (DeAngelis et al. 1985; Kelley and Hansen 1993; Dietl and Alexander 2000;

Dietl et al. 2004; Alexander and Dietl 2005; Klompmaker and Kelley 2015). Based on the

neontological experiments and field observations, complete drill holes and repair scars are

interpreted as a successful attack by carnivorous gastropod (Carriker 1951; Kitchell et al.

1981; Kowalewski 2004; Hutchings and Herbert 2013; Chattopadhyay et al. 2014*a*; Mondal

et al. 2014) and an unsuccessful predation attempt by durophagous predator respectively

(Carriker 1951; Blundon and Kennedy 1982; Dietl and Alexander 2009). These traces

recording the predation attempts on the prey's hard shells, are some of the best quantifiable

proxies for inferring predation from the fossil record (for review see (Alexander and Dietl

2003; Kelley and Hansen 2003; Klompmaker et al. 2019)). The frequency of repair scar

(RF) and complete drill holes (DF) are used for evaluating various aspects of predation in

deep time, including predation intensity and prey selection (Kitchell et al. 1981; Kelley and

Hansen 1993; Kowalewski et al. 1998; Dietl 2003; Kase and Ishikawa 2003; Chattopadhyay

and Baumiller 2010; Chattopadhyay and Dutta 2013; Tyler et al. 2013).

Inferences about interactions from predation traces have their limitations. The

implicit assumption for such interpretation is that other processes do not alter the

quantitative data provided by predation traces. It is recognized, however, that biases

introduced through taphonomy may influence the biological reliability of these measures

affecting the overall frequency of traces, site stereotypy, prey selection, and size selection

(Roy et al. 1994; Nebelsick 1999; Zuschin et al. 2003; Kosloski 2011; Gorzelak et al. 2013;

Chattopadhyay et al. 2014*b*; Chojnacki and Leighton 2014; Sime and Kelley 2016; Dyer et

al. 2018; Pruden et al. 2018; Smith et al. 2019; Salamon et al. 2020). Apart from taphonomy,

methods of collection and subsequent analyses may also influence the interpretation of

predation patterns. In contrast to the bulk collection, targeted sampling of specific size class

or taxon impacts inferred predation intensities (Kowalewski and Hoffmeister 2003; Kosloski

et al. 2008; Ottens et al. 2012; Hattori et al. 2014; Chattopadhyay et al. 2016; Hausmann et

al. 2018). Theoretical investigations also demonstrated the effect of sample size on inferred

predation intensity (Smith et al. 2018, 2022). Analytical techniques to evaluate and compare

predation measures across groups often impact the inferences (Kowalewski 2002; Leighton

2002; Grey et al. 2006; Stafford and Leighton 2011; Dietl and Kosloski 2013; Smith et al.

2018; Budd and Mann 2019).

Aspects of a specific community, such as evenness, selectivity of predation, and

sampling intensity, may influence predation inferences drawn at the community level, such

as predation intensity, prey selection. Such influences are crucial for studies that combine

predation data from discrete samples and reconstruct temporal/spatial changes in predation

patterns. Using theoretical simulation based on a resampling technique, we develop a

methodological framework to understand the effect of community evenness, sampling

intensity, and the nature of predation selectivity on inferred predation estimates. We attempt

to estimate these effects on the inferred predation intensity and the number of prey species.

The inferred number of prey species provides an insight into the choice of prey by the

predator. We also propose a method of post-facto standardization and validate our approach

using predation data from four Plio-Pleistocene fossil assemblages of Florida.

113   **Materials and method**

We created several hypothetical live assemblages of molluscs that a specific group of

predators attacks with differing probabilities. We use a resampling method to compare the

predation patterns inferred from these assemblages. We assumed that all individuals are

finally represented in the death assemblage, each predation attempt leaves a distinct mark on

the prey, and all predators demonstrate the same prey-selection behavior in specific

situations. We acknowledge that some of the particular values of species richness, evenness,

predation intensity, and predation selectivity might be rare in nature. Our attempt, however,

is to design and test a general framework applicable to a large spectrum of community

structures and predation patterns, even if some end-member scenarios do not have a natural

representation. It is also true that predation patterns observed in fossil assemblages may

differ from that of the death assemblage due to taphonomic factors which have not been

considered in the present study.

Model assemblages:

We created 30 hypothetical model assemblages, each with 30 species and 3000

individuals with varying evenness, predation intensity, and prey preferences (Table 1). Each

model assemblage had a unique combination of evenness, predation intensity, and prey

preference. To evaluate evenness, we used Pielou's evenness index, one of the commonly

used measures of evenness. We calculated the evenness of an assemblage (E_T) as

133                          $$E_T = H/ \ln(S_T)$$

134   Where,

135   H = Shannon's diversity index

136   S_T = Total number of species in the assemblage

The evenness in these models ranged from a theoretical minimum of 0.1 to a

theoretical maximum of 1. Model assemblages with maximum evenness of one had 100

individuals for 30 species. Assemblages with intermediate evenness of 0.7 had five common

species with 500 individuals each and 25 rare species with 20 individuals each (Table 1).

Assemblages with low evenness of 0.5 had 910 individuals in each of the three common

species and ten individuals in each of the 27 rare species. For assemblages with a very low

evenness of 0.2, there are only one common species with 2710 individuals, and the

remaining 29 rare species consist of 10 individuals each.

We calculated the predation intensity at the assemblage level (PI_T) and for prey

species (PI_{prey}). The total number of prey species is S_{prey}. PI_T is calculated as

147                                   $PI_T = N_P/N$

148   Where,

149   N_P = Number of individuals with predation mark

150   N = Total number of individuals in the assemblage

PI_{prey} denotes predation intensity in the species that have been attacked. The predation

intensity of the total assemblage (PI_T) was categorized into three levels: low (0.2), medium

(0.5), and high (0.8) (Table 1). A certain number of individuals from specific species would

be considered prey with predation marks as dictated by the (PI_T). The prey-preference of the

predator can either be non-selective or selective. In the case of non-selective predation (Case

1), all species have an equal probability of being attacked irrespective of their abundance

(Fig 1). Selective predation represents assemblages where prey species have an unequal

chance of being attacked relative to their abundance. In model assemblages with selective

predation, we constructed two cases; the predator can either attack the common species

(Case 2) (Fig 1) or the rare species (Case 3) (Fig 1).

For all models, the probability of an attack is determined by the PI_T which can be

0.2, 0.5 or 0.8. For scenarios of selective predation, only certain species are available as prey

and we assign the probability of attack as 0 to the rest of the species. In the case of selective

predation on abundant species with low predation intensity, for instance, the probability of

an attack is assigned as 0.2 for all the individuals of common species and 0.0 for all the

individuals of rare species. Selective predation has not been considered for assemblages with

maximum evenness because all the species are equally abundant and share an equal

probability of attack.

Simulation design

We performed a simulation to evaluate the effect of sample size on inferred

predation intensity ($PI_{T.inf}$) and the number of prey species ($S_{prey.inf}$) for all the model

assemblages. In the simulation, 100 individuals were drawn randomly from a model

assemblage. The number of attacked individuals (N_P) and the number of prey species (S_{prey})

represented by the attacked individuals were counted in those 100 individuals. Inferred

predation intensity ($PI_{T.inf}$) for the drawn sample is calculated as a ratio of the number of

attacked individuals and the total number of individuals (i.e., 100 in the first draw). We kept

the step size as 100 to gain an accurate representation of predation intensity and to avoid the

issues related to insufficient sample size (Kosloski et al. 2008; Dietl and Kosloski 2013;

Smith et al. 2022). The exact process is repeated 30 times without replacement until all the

individuals from the assemblage are sampled. Following the principles of rarefaction

analysis that are helpful when attempting to standardize sampling effort, we chose to use

subsampling without replacement (Kowalewski and Novack-Gottshall 2010). This entire

process was iterated 1000 times. The mean and standard deviation are calculated for inferred

predation intensity ($PI_{T.inf}$) and prey species richness ($S_{prey.inf}$) over 1000 iterations for a

specific model assemblage. The difference in predation intensity ($Diff_{PI}$) is calculated as the

difference between PI_T and $PI_{T.inf}$ for an assemblage. Similarly, the difference between S_{prey}

and $S_{prey.inf}$ is taken as the difference of prey species richness (Diff$_S$). The same technique is

applied to all the model assemblages.

Predation dataset:

We used data on predation records of molluscs from four Pleistocene localities in

Florida (Chattopadhyay and Baumiller 2010) to validate the proposed technique. The goal is

to quantitatively evaluate if we can compare the predation estimates of discrete communities

characterized by different evenness, predation style and sample size. The dataset consists of

abundance, drilling frequency, and repair scar frequency of 14 molluscan species. We drew

samples without replacement from each locality with increasing sample size. The sample

size for each draw was a hundred until the last draw; in the last draw, where the remaining

sample size is less than 200, all are drawn. For Punta Gorda (total=2418 individuals), 100

individuals were drawn 23 times, and 118 individuals were drawn for the last (24th) draw. A

similar procedure is followed for Miami Canal (total = 4794 individuals), Mc Queens pit

(total=659 individuals), and Chiquita (total=894 individuals).

We used a sampling standardization protocol to compare these assemblages and

assess the sensitivity of the inferred predation intensity (PI$_{T.inf}$) and inferred prey-species

richness (S$_{prey.inf}$) on sampling intensity. The sample size of Mc Queens pit (659) is

considered a reference as it has the smallest sample size among all four locations. The

distribution of inferred predation intensity (PI$_T$) is compared for all assemblages at a sample

size of 500 by a pairwise comparison using Kolmogorov-Smirnov (K-S) test. If the pairwise

K-S test shows significant differences between all pairs of assemblages, then the variation

between assemblages is not caused by sampling and community evenness. If some

assemblages show non-significant differences in the pairwise K-S test, then small sample

size might influence inferred predation intensity and prey richness. Larger sample size is

considered a reference, and the pairwise comparison using the K-S test is repeated again for

those pairs of assemblage. The process is repeated until the maximum number of

assemblage pairs shows significant differences. Following a similar protocol, the

distribution of inferred prey-species richness ($S_{prey.inf}$) is also compared.

All simulations and statistical analyses were performed in R (version 4.2.0) (R Core

Development Team, 2012).

220   **Results**

Inferred predation intensity

The inferred predation intensity ($PI_{T.inf}$) may vary substantially from the actual value

of overall predation intensity (PI_T) and predation intensity of prey groups (PI_{prey}), especially

at smaller sample sizes (Fig 2). For non-selective predation (Case 1), $Diff_{PI}$ is affected by the

sample size, although not by evenness. At a smaller sample size, the difference is higher

($Diff_{PI=}0.04$) implying a lower $PI_{T.inf}$ than the actual PI_T value. $PI_{T.inf}$ converges to PI_T with

increasing sample size (Fig 3, Table 5).

Evenness influences inferred predation intensity ($PI_{T.inf}$) when the predation is non-

selective (Case 2 and 3) (Fig 2). When the common species are preferentially attacked (Case

2), $Diff_{PI}$ is low (mean = -0.0003, standard deviation = 0.0124) for communities with lower

evenness ($E_T=0.2$) and low original predation intensity ($PI_T =0.2$) implying good

correspondence between $PI_{T.inf}$ and PI_T (Fig 4, Table 5). Communities with higher evenness

($E_T>0.2$) showed high $Diff_{PI}$ (Table 5) even at a higher sample size implying that $PI_{T.inf}$ will

be different from PI_T (Fig 4). Except for one specific model assemblage (E_T=0.5, PI_T =0.2),

all assemblages show a lower $PI_{T.inf}$ than the original PI_T (Table 5).

When rare species are attacked (Case 3), the $Diff_{PI}$ varies depending on the

combination of evenness and predation intensity. The $Diff_{PI}$ is positive for all communities

with low evenness (E_T=0.2) irrespective of the predation intensity (Fig 5A-C, Table 5),

implying, a lower value of $PI_{T.inf}$ compared to PI_T. Communities with high evenness

(E_T=0.7) showed negative $Diff_{PI}$, indicating a higher $PI_{T.inf}$ than PI_T (Fig 5G-I). The $Diff_{PI}$

value in communities with medium evenness (E_T=0.5) depends on predation intensity; in

those communities, $PI_{T.inf}$ is lower compared to PI_T for lower predation intensity ($PI_T < 0.8$)

(Fig 5D-E) and higher in for higher predation intensity (Fig 5F, Table 5). However, the

variation in $Diff_{PI}$ is lower for Case 3 compared to similar communities in Case 2 (Table 5).

Inferred number of prey species

The inferred number of prey species ($S_{prey.inf}$) follows a rarefaction curve where

$S_{prey.inf}$ increases with increasing sample size before plateauing and converging to the actual

value of S_{prey} (Fig 6). The required sample size for convergence depends on evenness and

selectivity of predation. In the case of non-selective predation (Case 1), the $Diff_S$ decreases

exponentially with increasing sample size and the converges to zero at a range of sample

sizes depending on the evenness and predation intensity (Fig 7, Table 6). At a given

predation intensity, convergence occurs at a smaller sample size with increasing evenness.

For example, at low predation intensity (PI_T=0.2), the required sample size for $Diff_S$ to

converge to 0 is 3000 when evenness is low (E_T=0.2) (Fig 7A) and 1200 when evenness is

high (E_T=1) (Fig 7J, Table 6).

In selective predation when common species are preyed upon (Case 2), $Diff_S$ does

not reflect any sensitivity to the sample size (Fig 8). This is due to the low value of S_{prey} that

converges to its actual value within the first few draws (Fig 8, Table 6). However, when the

rare species are attacked (Case 3), $S_{prey.inf}$ is highly sensitive to the sample size because the

required sample size for convergence of $Diff_S$ depends on evenness and predation intensity

(Fig 9). In general, communities with higher evenness require small sample size for

convergence of $Diff_S$ implying less sensitivity of $S_{prey.inf}$ on sample size at a given predation

intensity (Fig 9). For example, at low predation intensity ($PI_T=0.2$), the required sample size

for $Diff_S$ to converge to 0 is 3000 when evenness is low ($E_T=0.2$) (Fig 9A) and 2800 when

evenness is high ($E_T=0.7$) (Fig 9G, Table 6).

Inferred predation estimates from Florida

The assemblages from the four localities of Florida are different in terms of their

evenness and sample size (Table 2). Except for Miami Canal, all the localities show a case

of non-selective predation represented by the lack of correlation between the relative

abundance of the prey and prey-specific predation intensity (PI_{prey}) for drilling and

durophagous predation (Table 3). The significant positive correlation in Miami Canal

implies that this is not non-selective predation (Case 1). Because the drilling predators attack

all the species, it does not represent Case 2 or Case 3 for drilling predation. For durophagous

predation where the most abundant species are attacked, it approximates Case 2.

For inferred predation intensity ($PI_{T.inf}$), there is substantial overlap between three

localities (Punta Gorda, Miami Canal, and Mc Queens pit) for both drilling and durophagy

(Fig 10). For inferred prey species richness ($S_{prey.inf}$), the assemblages show slightly different

patterns between drilling and durophagous predation. For drilling predation, all the

assemblages show a substantial overlap (Fig 10). The durophagous predation record,

however, shows a separation between communities with low evenness (Punta Gorda) and

high-evenness (Mc Queens pit, Chiquita) (Fig 10).

For inferred predation intensity ($PI_{T.inf}$) and species richness ($S_{prey.inf}$), the sample

size-standardized resampling protocol (described before) shows a significant difference (*p*

<0.005) in all pairwise comparison (Table 4). This implies that the difference across

assemblages cannot be explained by the sampling intensity or the evenness of the

assemblage.

290   **Discussion**

Paleontological research on predation has expanded rapidly in scope, methods, and

goals. In the recent years, many studies focused on documenting the evidence of predation

from times, geographic areas, and taxa that are poorly known for their predation record

(Rojas et al. 2014; Randle and Sansom 2019; Bicknell and Holland 2020; Gordillo and

Malvé 2021; Klompmaker and Landman 2021; Gordillo et al. 2022) and using predation

records for testing evolutionary hypotheses (Klompmaker et al. 2017; Gehling and Droser

2018; Harper et al. 2018; Lerosey-Aubril and Peel 2018; Petsios et al. 2021). In contrast, a

relatively small number of studies focused on the analytical methods to evaluate the

reliability of predation measures in recent years (Smith et al. 2018, 2019, 2022; Budd and

Mann 2019). Our model demonstrates how the inferred predation intensities may vary with

community structures, predation selectivity, and sampling intensities. It highlighted the

importance of these factors in influencing the predation estimates of live and death/fossil

assemblages; it also underscores why it is necessary to develop a methodological framework

of sample standardization before comparing predation estimates of assemblages separated

by time and space.

Effect on the inferred intensity

Our simulation results show that communities' evenness does not significantly

change the inferred predation intensity when random encounters between predator and prey

guide predation. However, it is uncommon to find predation events completely random in

the natural world. Predators select prey species to maximize net energy gain within the

constraints of many factors, including reproductive demands, predator interference,

predation risk, avoidance of prey, deterrents, and predator behavior (Seitz et al. 2001;

Stephens and Krebs 2019). The inferred predation intensity may differ significantly from the

original predation intensity in such selective predation. Following the considerations of

optimal foraging theory (Hughes 1980; Pyke 1984; Burrows and Hughes 1991; Stephens

and Krebs 2019), two aspects make predation selective. The first is the relative ease with

which a predator encounters prey. Encounter in the marine ecosystem is determined by

several things, including the abundance of the prey, accessibility of the prey, landscape

heterogeneity, predator abundance, abundance of secondary predators, and habitat type

(Ryer and Olla 1995; Seitz et al. 2001; Sims et al. 2006; Casey and Chattopadhyay 2008;

Martinelli et al. 2015). Keeping the other factors constant, the probability of encounter

increases with the relative abundance of a prey species (Vermeij 1983; Leighton 2002;

Leonard-Pingel and Jackson 2013); this decreases the foraging time and increases the net

energy gain of the predator. The second aspect is the traits (morphological, ecological,

behavioral) of the prey that dictate the net energy gain of the predator. The final selection by

the predator is often a combination of these factors. A higher attack rate may be found in an

abundant prey species due to its higher encounter rate than a rarer species. This would lead

to scenarios similar to Case 2, where the inferred predation intensity of low-evenness

communities would be higher than the actual predation intensity. This inflated measure

results from the over-representation of common species in smaller samples that are primarily

attacked.

Most often than not, the encounter frequency does not dictate the attack frequency,

and the selection of prey is guided by the prey traits such as size, morphology, behavior

(Kitchell et al. 1981; Palmqvist et al. 1996; Leighton 2001; Zlotnik and Ceranka 2005;

Chattopadhyay and Dutta 2013; Chattopadhyay et al. 2014*a*, 2015, 2020; Martinelli et al.

2015; Chandroth and Chattopadhyay 2022). These would be similar to Case 3, where the

most dominant groups are not preyed upon. The inferred predation intensity of low-evenness

communities would be lower than the actual predation intensity. This apparent drop in

predation intensity results from the lack of representation of rare species in smaller samples

that are never attacked. It is especially problematic because this difference is substantial for

all evenness. This observation is consistent with the findings by Smith et al. (2021) where

they demonstrated the effects of overdispersion and zero inflation using count data of

predation traces. They concluded that the crucial element underlying these effects was

sample size. Their results support our findings that predation measures lack reliability in a

small sample size.

Effect on inferred selectivity

Predation is known to impact the structure of a community, including the overall

richness, distribution and evenness (Schemske et al. 2009; Freestone et al. 2011, 2020). It is

therefore important to evaluate the inherent dependence of the predation inferences on the

community structure before considering the evolutionary impact of predation on shaping the

community structure in deep time. Our models demonstrate that the inferred number of prey

species may depend on the evenness of the live community. Communities with low

evenness differ significantly from the original prey species and yield fewer inferred prey

species even when the predation is non-selective (Case 1). This may lead to the development

of an artificial selectivity, primarily driven by the preferential counting of the dominant

species and not by the biological preference demonstrated by the predators. Therefore, any

community with low evenness suffers from the high likelihood of underrepresenting the

number of prey species. The deviation from the true prey-species richness is higher for

smaller sample size and lower predation intensity. Communities with higher predation

intensity will provide the true prey-species richness at a smaller sample size than

communities with lower predation intensity. Selective predation (as indicated by Case 2 and

3) also creates similar deviations.

The sensitivity of inferred prey species richness on sample size, evenness, and

original predation selectivity makes the comparison of prey species richness in spatially or

temporally distinct assemblages somewhat unreliable unless they are normalized. This is

especially important when comparing predation estimates from assemblages representing

different time-bins or environments likely to show varying diversity/evenness.

371   **Paleontological case study**

The assemblages from the four Florida localities have been used to interpret the

relationship between durophagy and drilling predation (Chattopadhyay and Baumiller,

2010). However, the study's conclusions did not consider sample size or community

structure. The assemblages at these localities differ in their evenness and sample size (Table

2). Only in Miami Canal, predatory attacks (durophagous and drilling) are guided by the

relative abundance of prey species and hence deviates from non-selective predation. The

sample size-standardized resampling protocol revealed a significant difference in pairwise

comparison for all inferred predation intensity and prey species richness estimates. This

implies that the differences in predation measure across assemblages are largely independent

of sampling intensity or the evenness of the assemblage.

It is essential to recognize that several factors played a role in this particular case,

making these assemblages less susceptible to community evenness and sampling intensity.

Because three localities (Punta Gorda, Mc Queens pit, Chiquita) are showing non-selective

predation, they are less likely to be affected by sample size. Moreover, they have medium to

high evenness, making them less sensitive to sample size. Miami Canal, however, is

characterized by low evenness (0.31), shows evidence of selective predation and low

predation intensity ($PI_T < 0.2$). Assemblages with these characteristics are more prone to

differ significantly from actual predation measures at small sample sizes (Fig 4A). Because

Miami Canal has the largest sample size among the localities, makes it less likely to be

affected by these factors. Hence, the observed $S_{prey.drill}$ and $S_{prey.repair}$ are least likely to be

affected by these factors.

Proposed protocol of post-facto standardization of predation data

The following protocol may be followed to compare predation intensity and

selectivity of spatially/temporally distinct assemblages to avoid misinterpretation. Using the

protocol in the simulation model, inferred predation intensity ($PI_{T.inf}$) and inferred prey-

species richness ($S_{prey.inf}$) need to be calculated at a specific step size of 100 for all

assemblages. The step size of all assemblages should be equal till the last step when the

remaining number of individuals in that assemblage are drawn. The step size can be lowered

till 30 if the total assemblage size is small. Reducing the step size any further may create

erroneous results due to a smaller sample size (Kosloski et al. 2008; Dietl and Kosloski

2013; Smith et al. 2022). To understand the sensitivity of inferred predation intensity (PI_T)

on sample size, the assemblage with the smallest sample size should be considered as the

reference. The distribution of inferred predation intensity (PI_T) for all assemblages should be

compared at that sample size by a pairwise comparison using Kolmogorov-Smirnov (K-S)

test. Suppose the pair-wise comparison yields a significant difference between two

assemblages. In that case the differences in inferred predation intensity (PI_T) cannot be

explained by sample size alone and hence, likely to represent the actual variation. These

pairs would be considered comparable at that sample size. Suppose assemblages show non-

significant difference in the pairwise K-S test. In that case, we cannot reject the possibility

of small sample size influencing the inferred predation intensity and hence, should not be

considered for further comparative analysis at that pre-selected sample size (Fig 11). The

same process can be repeated to understand the sensitivity of the inferred prey-species

richness ($S_{prey.inf}$) on sample size.

The next step is to compare the remaining pairs with non-significant differences at

different sample sizes. If any pair contains the assemblage with previously set reference

sample size (smallest sample size), then we cannot include the pair for further analysis. For

other pairs where the smallest sample size is larger than the previously set reference, the

entire process described above should be repeated. The number of pairs that yield statistical

significance can be considered comparable at the new reference sample size. This iteration

should be performed with increasing sample sizes till the maximum number of assemblage

pairs show significant differences in distribution of inferred predation intensity (PI_T) and

prey species richness ($S_{prey.inf}$). Finally, the pairs that show non-significant differences even

at the largest sample size, we cannot reject the influence of sampling intensity and inherent

community structure in shaping the predation measures. They should be excluded from

comparative analyses of predation signals. Estimating PI_{prey} is difficult, especially for cases

where rare species are attacked; excluding species without any predation trace while

calculating PI_T may give us some insight.

431   **Caveats and implications**

The fossil record of predation has shaped our understanding of how the nature of

biotic interaction changed over time and its role as an evolutionary mechanism. Preserved

traces, such as drill holes and repair scars, are some of the best quantifiable predation

proxies and are often used to assess the evolutionary impact of predation in deep time

(Vermeij et al. 1981; Alexander and Dietl 2003; Kelley and Hansen 2003). However, studies

aiming to evaluate the predation trend through time are often forced to use predation data

from discrete assemblages that differ in sample size, inherent community evenness, and the

type of predation selectivity (Harper 2016). Our study demonstrates the effect of such

factors on the inferred predation intensity and the recognized prey richness. Comparison

between temporally separated collections, such as Paleozoic and Cenozoic predation records

that are known to be different in the sample size (and probably predatory behavior), are

susceptible to such factors.

Our proposed method of post-facto standardization will be essential for such

comparisons and to establish the true nature of biotic interaction through time. It is

important to recognize that the proposed protocol is a preliminary attempt toward

standardization without considering several complexities. The simulations are primarily

developed for communities that are preserving the community structure of the live

communities. A number of actualistic studies revealed, however, that the death/fossil

assemblage can substantially differ from the live assemblage because they are typically

time-averaged, representing a mix of multiple generations (Kidwell et al. 1991; Kidwell and

Flessa 1995; Kidwell 2007; Olszewski and Kidwell 2007; Tomašových and Kidwell 2009;

Kidwell and Tomasovych 2013; Bhattacherjee et al. 2021). If a specific section of the live

community is preferentially lost due to preservation and if the predation signature of those

specimens differs from the remaining assemblage, the proposed standardization method will

fail to detect that. For example, some predation attempts are size-selective, and larger size

classes often show higher predation resistance and lower predation intensity. Because the

preservation potential of smaller size classes is lower than larger ones (Cooper et al. 2006),

the elective absence of small size class in the fossils would result in a low inferred predation

intensity compared to the original value. Multiple interactions during the lifetime or after the

death of the prey may change the frequency of the overall assemblage (Kosloski 2011;

Gordillo and Archuby 2014). A molluscan community affected by drilling predation may

also be subjected to crushing predation; because the durophagous predators only go after the

live prey (non-drilled), the relative proportion of drilled shells increase if the predators

successfully destroy the shells as part of the predation process (Smith et al. 2019). Predation

style and resulting predation trace also differ among predators. Two of the most common

types of predations studied in the fossil, drilling and durophagy, are quite different in several

aspects. It is possible to identify successful and unsuccessful predation by observing the

completeness of the drill holes, successful attacks by durophagous predators often result in

unrecognizable fragmentation (Kosloski 2011; Leighton et al. 2016; Dyer et al. 2018).

Repair scars represent a failed durophagous attack. Comparing drill holes and repair scars,

therefore, are not without limitations. Although our study attempts to recognize the possible

source of analytical bias and to recognize them in the observed database, it clearly glosses

over the full complexities of predation style, post-mortem alteration and time-averaging.

Following the direction of reconstructing fossil assemblages from live data using the

modeling approach (Olszewski 2004, 2012), we plan to develop more inclusive frameworks

in future to address such complexities.

479   **Conclusions**

The effect of community structure and sampling intensity on the inferred predation

estimates is rarely explored. Using a resampling technique, our study demonstrates the

impact of these aspects on the estimates of predation intensity and the number of prey

species. Our results show that the communities with highly selective predation are the most

sensitive to sampling intensity, and the inferred predation intensity of these assemblages can

substantially deviate from the actual value. In contrast, predation intensity for non-selective

predation tends to be unaffected by sampling intensity. Inferred prey-species richness is also

influenced by the nature of community evenness, predation selectivity, and actual predation

intensity. For non-selective predation, communities with low evenness and low predation

intensity are highly sensitive to sample size. The inferred prey-species richness can be

underrepresented significantly at smaller sample size. For selective predation, the sensitivity

depends on the nature of selection. The inferred prey-species richness deviates significantly

when rare species are attacked preferentially. Our study also provides a framework of post-

facto standardization of the predation data to remove the effect of sample size/evenness

during comparison. The proposed method, although simple, will provide a fundamental

framework for comparison of discrete assemblages as they are often characterized by a

difference in sample size, community structure and predation selectivity.

Acknowledgments

This project was supported by the SERB core research grant (CRG/2018/002604). The

IISER Pune Doctoral fellowship supported MB. We would like to thank Satyaki Mazumdar

for his help in designing the R code. We are grateful to the associate editor, Erin Saupe and

reviewer, Jansen Smith for their detailed comments that substantially improved the quality

of the manuscript.

References

Abrams, P. A. 2015: Why ratio dependence is (still) a bad model of predation. Biological

Reviews 90:794–814.

Alexander, R. R., and G. P. Dietl. 2003: The Fossil Record of Shell-Breaking Predation on

Marine Bivalves and Gastropods. Pp.141–176 *in* P. H. Kelley, M. Kowalewski, and

T. A. Hansen, eds. Predator—Prey Interactions in the Fossil Record. Springer US,

Boston, MA.

———. 2005: Non-Predatory Shell Damage in Neogene Western Atlantic Deep-Burrowing

Bivalves. PALAIOS 20:280–295.

Barnes, C., D. Maxwell, D. C. Reuman, and S. Jennings. 2010: Global patterns in predator–

prey size relationships reveal size dependency of trophic transfer efficiency. Ecology

91:222–232.

Baumiller, T. K., M. A. Salamon, P. Gorzelak, R. Mooi, C. G. Messing, and F. J. Gahn.

2010: Post-Paleozoic crinoid radiation in response to benthic predation preceded the

Mesozoic marine revolution. Proceedings of the National Academy of Sciences

107:5893–5896.

Berryman, A. A. 1992: The Orgins and Evolution of Predator-Prey Theory. Ecology

73:1530–1535.

Bhattacherjee, M., D. Chattopadhyay, B. Som, A. S. Sankar, and S. Mazumder. 2021:

Molluscan live-dead fidelity of a storm-dominated shallow-marine setting and its

implications. PALAIOS 36:77–93.

Bicknell, R. D. C., and J. R. Paterson. 2018: Reappraising the early evidence of durophagy

and drilling predation in the fossil record: implications for escalation and the

Cambrian Explosion: Reappraising the early evidence of predation. Biological

Reviews 93:754–784.

Bicknell, R. D. C., and B. Holland. 2020: Injured trilobites within a collection of dinosaurs:

Using the Royal Tyrrell Museum of Palaeontology to document Cambrian predation.

Palaeontologia Electronica 23:a33.

Blundon, J. A., and V. S. Kennedy. 1982: Refuges for infaunal bivalves from blue crab,

Callinectes sapidus (Rathbun), predation in Chesapeake Bay. Journal of

Experimental Marine Biology and Ecology 65:67–81.

Budd, G. E., and R. P. Mann. 2019: Modeling durophagous predation and mortality rates

from the fossil record of gastropods. Paleobiology 45:246–264.

Burrows, M. T., and R. N. Hughes. 1991: Optimal Foraging Decisions by Dogwhelks,

Nucella lapillus (L.): Influences of Mortality Risk and Rate-Constrained Digestion.

Functional Ecology 5:461.

Carriker, M. R. 1951: Observations on the Penetration of Tightly Closing Bivalves by

Busycon and Other Predators. Ecology 32:73–83.

Casey, M. M., and D. Chattopadhyay. 2008: Clumping behavior as a strategy against drilling

predation: Implications for the fossil record. Journal of Experimental Marine Biology

and Ecology 367:174–179.

Chandroth, A., and D. Chattopadhyay. 2022: Micromorphy Offers Effective Defense

Against Predation: Insights from Cost-Benefit Analyses of the Miocene

Microgastropod Predation Record from Kerala, India. Contributions from the

Museum of Paleontology, University of Michigan 34:63–81.

Chattopadhyay, D., and T. K. Baumiller. 2010: Effect of durophagy on drilling predation: a

case study of Cenozoic molluscs from North America. Historical Biology 22:367–

379.

Chattopadhyay, D., and S. Dutta. 2013: Prey selection by drilling predators: A case study

from Miocene of Kutch, India. Palaeogeography, Palaeoclimatology, Palaeoecology

374:187–196.

Chattopadhyay, D., M. Zuschin, and A. Tomašových. 2015: How effective are ecological

traits against drilling predation? Insights from recent bivalve assemblages of the

northern Red Sea. Palaeogeography, Palaeoclimatology, Palaeoecology 440:659–

670.

Chattopadhyay, D., V. G. S. Kella, and D. Chattopadhyay. 2020: Effectiveness of small size

against drilling predation: Insights from lower Miocene faunal assemblage of Quilon

Limestone, India. Palaeogeography, Palaeoclimatology, Palaeoecology 551:109742.

Chattopadhyay, D., D. Sarkar, S. Dutta, and S. R. Prasanjit. 2014a: What controls

cannibalism in drilling gastropods? A case study on Natica tigrina. Palaeogeography,

Palaeoclimatology, Palaeoecology 410:126–133.

Chattopadhyay, D., A. Rathie, D. J. Miller, and T. K. Baumiller. 2014b: Hydrodynamic

effects of drill holes on postmortem transportation of bivalve shells and its

taphonomic implications. PALAIOS 28:875–884.

Chattopadhyay, D., M. Zuschin, S. Dominici, and J. A. Sawyer. 2016: Patterns of drilling

predation in relation to stratigraphy, locality and sieve size: Insights from the Eocene

molluscan fauna of the Paris Basin. Palaeogeography, Palaeoclimatology,

Palaeoecology 459:86–98.

Chojnacki, N. C., and L. R. Leighton. 2014: Comparing predatory drillholes to taphonomic

damage from simulated wave action on a modern gastropod. Historical Biology

26:69–79.

Cooper, R. A., P. A. Maxwell, J. S. Crampton, A. G. Beu, C. M. Jones, and B. A. Marshall.

2006: Completeness of the fossil record: Estimating losses due to small body size.

Geology 34:241–244.

DeAngelis, D. L., R. A. Goldstein, and R. V. O'Neill. 1975: A Model for Tropic Interaction.

Ecology 56:881–892.

DeAngelis, D. L., J. A. Kitchell, and W. M. Post. 1985: The Influence of Naticid Predation

on Evolutionary Strategies of Bivalve Prey: Conclusions from a Model. The

American Naturalist 126:817–842.

Dietl, G. P. 2003: Coevolution of a marine gastropod predator and its dangerous bivalve

prey. Biological Journal of the Linnean Society 80:409–436.

Dietl, G. P., and R. R. Alexander. 2000: Post-Miocene Shift in Stereotypic Naticid Predation

on Confamilial Prey from the Mid-Atlantic Shelf: Coevolution with Dangerous Prey.

PALAIOS 15:414–429.

Dietl, G. P., and R. R. Alexander. 2009: Patterns of unsuccessful shell-crushing predation

along a tidal gradient in two geographically separated salt marshes. Marine Ecology

30:116–124.

Dietl, G. P., and M. Kosloski. 2013: On the measurement of repair frequency: how

important is data standardization? PALAIOS 28:394–402.

Dietl, G. P., G. S. Herbert, and G. J. Vermeij. 2004: Reduced Competition and Altered

Feeding Behavior Among Marine Snails After a Mass Extinction. Science 306:2229–

2231.

Dyer, A. D., E. R. Ellis, D. J. Molinaro, and L. R. Leighton. 2018: Experimental

fragmentation of gastropod shells by sediment compaction: Implications for

interpreting drilling predation intensities in the fossil record. Palaeogeography,

Palaeoclimatology, Palaeoecology 511:332–340.

Ebert, M., M. Kölbl-Ebert, and J. A. Lane. 2015: Fauna and Predator-Prey Relationships of

Ettling, an Actinopterygian Fish-Dominated Konservat-Lagerstätte from the Late

Jurassic of Southern Germany. PLOS ONE 10:e0116140.

Ehret, D. J., B. J. Macfadden, and R. Salas-Gismondi. 2009: Caught in the act: trophic

interactions between a 4-million-year-old white shark (Carcharodon) and mysticete

whale from Peru. PALAIOS 24:329–333.

Freestone, A. L., R. W. Osman, G. M. Ruiz, and M. E. Torchin. 2011: Stronger predation in

the tropics shapes species richness patterns in marine communities. Ecology 92:983–

993.

Freestone, A. L., E. W. Carroll, K. J. Papacostas, G. M. Ruiz, M. E. Torchin, and B. J.

Sewall. 2020: Predation shapes invertebrate diversity in tropical but not temperate

seagrass communities. Journal of Animal Ecology 89:323–333.

Gehling, J. G., and M. L. Droser. 2018: Ediacaran scavenging as a prelude to predation.

Emerging Topics in Life Sciences 2:213–222.

Gordillo, S., and F. Archuby. 2014: Live-Live and Live-Dead Interactions in Marine Death

Assemblages: The Case of the Patagonian Clam Venus antiqua. Acta

Palaeontologica Polonica 59:429–442.

Gordillo, S., and M. E. Malvé. 2021: Drilling predation on Antarctic tusk shells: first records

on Recent scaphopods from the Southern Hemisphere. Antarctic Science 33:344–

348.

Gordillo, S., G. A. Morán, and M. E. Malvé. 2022: Octopuses and drilling snails as the main

suspects of predation traces on shelled molluscs in West Antarctica. Polar Biology

45:127–141.

Gorzelak, P., M. A. Salamon, and T. K. Baumiller. 2012: Predator-induced

macroevolutionary trends in Mesozoic crinoids. Proceedings of the National

Academy of Sciences 109:7004–7007.

Gorzelak, P., M. A. Salamon, D. Trzęsiok, and R. Niedźwiedzki. 2013: Drill Holes and

Predation Traces versus Abrasion-Induced Artifacts Revealed by Tumbling

Experiments. PLoS ONE 8:e58528.

Grey, M., E. G. Boulding, and M. E. Brookfield. 2006: Estimating multivariate selection

gradients in the fossil record: a naticid gastropod case study. Paleobiology 32:100–

108.

Haque, M. 2012: Existence of complex patterns in the Beddington–DeAngelis predator–prey

model. Mathematical Biosciences 239:179–190.

Harper, E. M. 2016: Uncovering the holes and cracks: from anecdote to testable hypotheses

in predation studies. Palaeontology 59:597–609.

Harper, E. M., J. A. Crame, and C. E. Sogot. 2018: "Business as usual": Drilling predation

637   across the K-Pg mass extinction event in Antarctica. Palaeogeography,

638   Palaeoclimatology, Palaeoecology 498:115–126.

639   Hattori, K. E., P. H. Kelley, G. P. Dietl, N. O. Moore, S. L. Simpson, A. M. Zappulla, K. J.

640   Ottens, and C. C. Visaggi. 2014: Validation of taxon-specific sampling by novice

641   collectors for studying drilling predation in fossil bivalves. Palaeogeography,

642   Palaeoclimatology, Palaeoecology 412:199–207.

643   Hausmann, I. M., H. Domanski, and M. Zuschin. 2018: Influence of setting, sieve size, and

644   sediment depth on multivariate and univariate assemblage attributes of coral reef-

645   associated mollusc death assemblages from the Gulf of Aqaba. Facies 64:20.

646   Hughes, R. N. 1980: Optimal foraging theory in the marine context. Oceanography and

647   Marine Biology: An Annual Review 18:423–481.

648   Huntley, J. W., and M. Kowalewski. 2007: Strong coupling of predation intensity and

649   diversity in the Phanerozoic fossil record. Proceedings of the National Academy of

650   Sciences 104:15006–15010.

651   Hutchings, J. A., and G. S. Herbert. 2013: No honor among snails: Conspecific competition

652   leads to incomplete drill holes by a naticid gastropod. Palaeogeography,

653   Palaeoclimatology, Palaeoecology 379–380:32–38.

654   Kase, T., and M. Ishikawa. 2003: Mystery of naticid predation history solved: Evidence

655   from a "living fossil" species. Geology 31:403.

656   Kelley, P. H., and T. A. Hansen. 1993: Evolution of the Naticid Gastropod Predator-Prey

657   System: An Evaluation of the Hypothesis of Escalation. PALAIOS 8:358.

———. 2003: The Fossil Record of Drilling Predation on Bivalves and Gastropods.

Pp.113–139 *in* P. H. Kelley, M. Kowalewski, and T. A. Hansen, eds. Predator—Prey

Interactions in the Fossil Record. Springer US, Boston, MA.

Kidwell, S., and A. Tomasovych. 2013: Implications of Time-Averaged Death Assemblages

for Ecology and Conservation Biology. Annual Review of Ecology Evolution and

Systematics 44:539–563.

Kidwell, S. M. 2007: Discordance between living and death assemblages as evidence for

anthropogenic ecological change. Proceedings of the National Academy of Sciences

104:17701–17706.

Kidwell, S. M., and K. W. Flessa. 1995: The Quality of the Fossil Record: Populations,

Species, and Communities. Annual Review of Ecology and Systematics 26:269–299.

Kidwell, S. M., D. W. Bosence, P. A. Allison, and D. E. G. Briggs. 1991: Taphonomy and

time-averaging of marine shelly faunas. Pp.115–209 *in* Taphonomy: Releasing the

Data Locked in the Fossil Record. Plenum, New York, US.

Kitchell, J. A., and J. F. Kitchell. 1980: Size-selective predation, light transmission, and

oxygen stratification: Evidence from the recent sediments of manipulated lakes1:

Predation and visibility. Limnology and Oceanography 25:389–402.

Kitchell, J. A., C. H. Boggs, J. F. Kitchell, and J. A. Rice. 1981: Prey Selection by Naticid

Gastropods: Experimental Tests and Application to Application to the Fossil Record.

Paleobiology 7:533–552.

Klompmaker, A. A., and P. H. Kelley. 2015: Shell ornamentation as a likely exaptation:

evidence from predatory drilling on Cenozoic bivalves. Paleobiology 41:187–201.

Klompmaker, A. A., and N. H. Landman. 2021: Octopodoidea as predators near the end of

the Mesozoic Marine Revolution. Biological Journal of the Linnean Society

132:894–899.

Klompmaker, A. A., M. Kowalewski, J. W. Huntley, and S. Finnegan. 2017: Increase in

predator-prey size ratios throughout the Phanerozoic history of marine ecosystems.

Science 356:1178–1180.

Klompmaker, A. A., P. H. Kelley, D. Chattopadhyay, J. C. Clements, J. W. Huntley, and M.

Kowalewski. 2019: Predation in the marine fossil record: Studies, data, recognition,

environmental factors, and behavior. Earth-Science Reviews 194:472–520.

Kosloski, M. 2011: Recognizing biotic breakage of the hard clam, Mercenaria mercenaria

caused by the stone crab, Menippe mercenaria: An experimental taphonomic

approach. Journal of Experimental Marine Biology and Ecology 396:115–121.

Kosloski, M. E., G. P. Dietl, and G. S. Herbert. 2008: Are museum collections adequate to

test the escalation hypothesis?: A preliminary case study using the Plio-Pleistocene

Strombus alatus species complex from Florida. Geological Society of America,

Abstracts with Program 40:373.

Kotta, J., T. Wernberg, H. Jänes, I. Kotta, K. Nurkse, M. Pärnoja, and H. Orav-Kotta. 2018:

Novel crab predator causes marine ecosystem regime shift. Scientific Reports

8:4956.

Kowalewski, M. 2002: The Fossil Record of Predation: An Overview of Analytical

Methods. The Paleontological Society Papers 8:3–42.

Kowalewski, M. 2004: Drill holes produced by the predatory gastropod Nucella lamellosa

(Muricidae): Palaeobiological and ecological implications. Journal Molluscan

Studies 70:359–370.

Kowalewski, M., and A. P. Hoffmeister. 2003: Sieves and Fossils: Effects of Mesh Size on

Paleontological Patterns. PALAIOS 18:460–469.

Kowalewski, M., and P. Novack-Gottshall. 2010: Resampling Methods in Paleontology. The

Paleontological Society Papers 16:19–54.

Kowalewski, M., A. Dulai, and F. T. Fürsich. 1998: A fossil record full of holes: The

Phanerozoic history of drilling predation. Geology 26:1091.

Kowalewski, M., A. P. Hoffmeister, T. K. Baumiller, and R. K. Bambach. 2005: Secondary

Evolutionary Escalation Between Brachiopods and Enemies of Other Prey. Science

308:1774–1777.

Langerhans, R. B. 2007: Evolutionary consequences of predation: avoidance, escape,

reproduction, and diversification. Pp.177–220 *in* A. M. T. Elewa, ed. Predation in

Organisms: A Distinct Phenomenon. Springer, Berlin, Heidelberg.

Leighton, L., N. Chojnacki, E. Stafford, C. Tyler, and C. Schneider. 2016: Categorization of

shell fragments provides a proxy for environmental energy and predation intensity.

Journal of the Geological Society 173:jgs2015-086.

Leighton, L. R. 2001: New example of Devonian predatory boreholes and the influence of

brachiopod spines on predator success. Palaeogeography, Palaeoclimatology,

Palaeoecology 165:53–69.

———. 2002: Inferring predation intensity in the marine fossil record. Paleobiology

28:328–342.

Leonard-Pingel, J. S., and J. B. Jackson. 2013: Drilling Intensity Varies Among Neogene

Tropical American Bivalvia in Relation to Shell Form and Life Habit. Bulletin of

Marine Science 89:905–919.

Lerosey-Aubril, R., and J. S. Peel. 2018: Gut evolution in early Cambrian trilobites and the

origin of predation on infaunal macroinvertebrates: evidence from muscle scars in

Mesolenellus. Palaeontology 61:747–760.

Martinelli, J. C., M. A. Kosnik, and J. S. Madin. 2015: Encounter frequency does not predict

predation frequency in tropical dead-shell assemblages. PALAIOS 30:818–826.

McNamara, K. J. 1994: The significance of gastropod predation to patterns of evolution and

extinction in Australian Tertiary echinoids. Pp.*in* Echinoderms through Time. CRC

Press.

Mondal, S., J. A. Hutchings, and G. S. Herbert. 2014: A note on edge drilling predation by

naticid gastropods. Journal of Molluscan Studies 80:206–212.

Nebelsick, J. 1999: Taphonomic legacy of predation on echinoids. Pp.347–352 .

Nilsen, E., D. Christianson, J.-M. Gaillard, D. Halley, J. Linnell, M. Odden, M. Panzacchi,

C. Toïgo, and B. Zimmermann. 2012: Describing food habits and predation: field

methods and statistical considerations. Carnivore Ecology and Conservation - a

handbook of techniques:256–272.

Olszewski, T. D. 2004: Modeling the Influence of Taphonomic Destruction, Reworking, and

Burial on Time-Averaging in Fossil Accumulations. PALAIOS 19:39–50.

———. 2012: Remembrance of things past: modelling the relationship between species'

abundances in living communities and death assemblages. Biology Letters 8:131–

134.

Olszewski, T. D., and S. M. Kidwell. 2007: The preservational fidelity of evenness in

molluscan death assemblages. Paleobiology 33:1–23.

Ottens, K. J., G. P. Dietl, P. H. Kelley, and S. D. Stanford. 2012: A comparison of analyses

of drilling predation on fossil bivalves: Bulk- vs. taxon-specific sampling and the

role of collector experience. Palaeogeography, Palaeoclimatology, Palaeoecology

319–320:84–92.

Palmqvist, P., B. Martínez-Navarro, and A. Arribas. 1996: Prey selection by terrestrial

carnivores in a lower Pleistocene paleocommunity. Paleobiology 22:514–534.

Petsios, E., R. W. Portell, L. Farrar, S. Tennakoon, T. B. Grun, M. Kowalewski, and C. L.

Tyler. 2021: An asynchronous Mesozoic marine revolution: the Cenozoic

intensification of predation on echinoids. Proceedings of the Royal Society B:

Biological Sciences 288:20210400.

Pringle, R. M., T. R. Kartzinel, T. M. Palmer, T. J. Thurman, K. Fox-Dobbs, C. C. Y. Xu,

M. C. Hutchinson, T. C. Coverdale, J. H. Daskin, D. A. Evangelista, K. M. Gotanda,

N. A. Man in 't Veld, J. E. Wegener, J. J. Kolbe, T. W. Schoener, D. A. Spiller, J. B.

Losos, and R. D. H. Barrett. 2019: Predator-induced collapse of niche structure and

species coexistence. Nature 570:58–64.

Pruden, M. J., S. E. Mendonca, and L. R. Leighton. 2018: The effects of predation on the

preservation of ontogenetically young individuals. Palaeogeography,

Palaeoclimatology, Palaeoecology 490:404–414.

Pyke, G. H. 1984: Optimal Foraging Theory: A Critical Review. Annual Review of Ecology

and Systematics 15:523–575.

Randle, E., and R. S. Sansom. 2019: Bite marks and predation of fossil jawless fish during

the rise of jawed vertebrates. Proceedings of the Royal Society B: Biological

Sciences 286:20191596.

Rojas, A., Cabrera, Fernanda, M. Verde, D. Urteaga, F. Scarabino, and S. Martínez. 2014:

The first predatory drillhole on a fossil chiton plate: an occasional prey item or an

erroneous attack? PALAIOS 29:414–419.

Roy, K., D. J. Miller, and M. Labarbera. 1994: Taphonomic Bias in Analyses of Drilling

Predation: Effects of Gastropod Drill Holes on Bivalve Shell Strength. PALAIOS

9:413.

Ryer, C. H., and B. L. Olla. 1995: Influences of food distribution on fish foraging behaviour.

Animal Behaviour 49:411–418.

Salamon, M. A., T. Brachaniec, and P. Gorzelak. 2020: Durophagous fish predation traces

versus tumbling-induced shell damage—a paleobiological perspective. PALAIOS

35:37–47.

Schemske, D. W., G. G. Mittelbach, H. V. Cornell, J. M. Sobel, and K. Roy. 2009: Is There

a Latitudinal Gradient in the Importance of Biotic Interactions? Annual Review of

Ecology, Evolution, and Systematics 40:245–269.

Seitz, R. D., R. N. Lipcius, A. H. Hines, and D. B. Eggleston. 2001: Density-Dependent

Predation, Habitat Variation, and the Persistence of Marine Bivalve Prey. Ecology

82:2435–2451.

Signor, P. W., and C. E. Brett. 1984: The Mid-Paleozoic Precursor to the Mesozoic Marine

Revolution. Paleobiology 10:229–245.

Sime, J. A., and P. H. Kelley. 2016: Common mollusk genera indicate interactions with their

predators were ecologically stable across the Plio-Pleistocene extinction.

Palaeogeography, Palaeoclimatology, Palaeoecology 463:216–229.

Sims, D. W., M. J. Witt, A. J. Richardson, E. J. Southall, and J. D. Metcalfe. 2006:

Encounter success of free-ranging marine predator movements across a dynamic

prey landscape. Proceedings of the Royal Society B: Biological Sciences 273:1195–

1201.

Smith, J. A., J. C. Handley, and G. P. Dietl. 2018: On drilling frequency and Manly's alpha:

towards a null model for predator preference in paleoecology. PALAIOS 33:61–68.

Smith, J. A., G. P. Dietl, and J. C. Handley. 2019: Durophagy bias: The effect of shell

destruction by crushing predators on drilling frequency. Palaeogeography,

Palaeoclimatology, Palaeoecology 514:690–694.

Smith, J. A., J. C. Handley, and G. P. Dietl. 2022: Accounting for uncertainty from zero

inflation and overdispersion in paleoecological studies of predation using a

hierarchical Bayesian framework. Paleobiology 48:65–82.

Stafford, E. S., and L. R. Leighton. 2011: Vermeij Crushing Analysis: A new old technique

for estimating crushing predation in gastropod assemblages. Palaeogeography,

Palaeoclimatology, Palaeoecology 305:123–137.

Stanley, S. M. 2008: Predation defeats competition on the seafloor. Paleobiology 34:1–21.

Stephens, D. W., and J. R. Krebs. 2019: Foraging Theory. Princeton University Press, p.

Tomašových, A., and S. M. Kidwell. 2009: Fidelity of variation in species composition and

diversity partitioning by death assemblages: time-averaging transfers diversity from

beta to alpha levels. Paleobiology 35:94–118.

Tyler, C. L., L. R. Leighton, S. J. Carlson, J. W. Huntley, and M. Kowalewski. 2013:

Predation on Modern and Fossil Brachiopods: Assessing Chemical Defenses and

Palatability. PALAIOS 28:724–735.

Vermeij, G. J. 1977: The Mesozoic Marine Revolution: Evidence from Snails, Predators and

Grazers. Paleobiology 3:245–258.

———. 1983: Shell-Breaking Predation through Time. Pp.649–669 *in* M. J. S. Tevesz and

P. L. McCall, eds. Biotic Interactions in Recent and Fossil Benthic Communities.

Springer US, Boston, MA.

———. 1993: Evolution and Escalation: An Ecological History of Life. Princeton

University Press, p.

Vermeij, G. J., D. E. Schindel, and E. Zipser. 1981: Predation Through Geological Time:

Evidence from Gastropod Shell Repair. Science 214:1024–1026.

Zlotnik, M., and T. Ceranka. 2005: Patterns of drilling predation of cassid gastropods

preying on echinoids from the Middle Miocene of Poland. Acta Palaeontologica

Polonica 50.

Zuschin, M., M. Stachowitsch, and R. J. Stanton. 2003: Patterns and processes of shell

fragmentation in modern and ancient marine environments. Earth-Science Reviews

63:33–82.

833   Table caption

Table 1. A summary of the model assemblages used for this study with varying evenness,

predation intensity and predator preference.

Table 2. A summary of the predation data from four Plio-Pleistocene fossil assemblages of

Florida (Chattopadhyay and Baumiller, 2010).

Table 3. The result of Spearman rank order correlation test for proportional abundance and

PI_{prey} for the predation estimates across four Plio-Pleistocene fossil assemblages of

Florida (Chattopadhyay and Baumiller, 2010). The statistically significant ($p<0.05$)

results are marked in bold.

Table 4. The test-statistic (D) of Kolmogorov–Smirnov test comparing the predation

estimates across four Plio-Pleistocene fossil assemblages of Florida using sample-

standardization protocol. All the results are statistically significant ($p<0.05$).

Table 5. A summary of the difference in inferred predation intensity from the original value

for the model assemblages. Each cell contains information about the mean value and

standard deviation of $Diff_{PI}$; the first two represents the sign and magnitude of the

mean value. A positive mean value of $Diff_{PI}$ indicates a larger value of original than

inferred predation intensity ($PI_T > PI_{T.inf}$).

Table 6. A summary of the difference in inferred prey species richness from the original

value for the model assemblages. Each cell contains the minimum sample size

required for $Diff_S$ to converge to zero for each model assemblages. A smaller number

indicates that the inferred prey species richness converges to the original value

($S_{prey.inf} = S_{prey}$) at smaller sample size.

Figure caption

Figure 1. An illustrative diagram of model assemblages with varying degrees of evenness,

predation intensity, and predation style (selective and non-selective). Mollusc

drawings are from publicdomainpictures.net with subsequent modifications. [Figure

1. Two-column; Grayscale]

Figure 2. Plot showing variation in inferred predation intensity ($PI_{T.inf}$) with varying sample

sizes for different model assemblages. The rows indicate the different degrees of the

selectiveness of predation, and the columns indicate predation intensity in the

original assemblage (PI_T). The warmer colors represent assemblages with higher

evenness. [Figure 2. Two columns; Color]

Figure 3. Plot showing the difference between the original (PI_T) and inferred predation

intensity ($PI_{T.inf}$) at varying sample size for non-selective predation (Case 1). The

rows indicate evenness and the columns represent original predation intensity. The

red line represents the zero line where overall and inferred predation intensities are

the same ($PI_{T.inf} = PI_T$). The grey dots and bars represent the mean and standard

deviation of the simulated differences for specific model assemblages. [Figure 3.

Two columns; Color]

Figure 4. Plot showing the difference between the original (PI_T) and inferred predation

intensity ($PI_{T.inf}$) at varying sample size for selective predation of the common

species (Case 2). The rows indicate evenness and the columns represent original

883   predation intensity. The red line represents the zero line where overall and inferred

884   predation intensities are the same ($PI_{T.inf} = PI_T$). The grey dots and bars represent the

885   mean and standard deviation of the simulated differences for specific model

886   assemblages. [Figure 4. Two columns; Color]

888 Figure 5. Plot showing the difference between the original (PI_T) and inferred predation

889   intensity ($PI_{T.inf}$) at varying sample size for selective predation of the rare species

890   (Case 3). The rows indicate evenness and the columns represent original predation

891   intensity. The red line represents the zero line where overall and inferred predation

892   intensities are the same ($PI_{T.inf} = PI_T$). The grey dots and bars represent the mean and

893   standard deviation of the simulated differences for specific model assemblages.

894   [Figure 5. Two columns; Color]

896 Figure 6. Plot showing variation in inferred prey species richness ($S_{prey.inf}$) with varying

897   sample sizes for different model assemblages. The rows indicate the different

898   degrees of the selectiveness of predation, and the columns indicate prey species

899   richness in the original assemblage (S_{prey}). The warmer colors represent assemblages

900   with higher evenness. [Figure 6. Two columns; Color]

902 Figure 7. Plot showing the difference between the original (S_{prey}) and prey species richness

903   ($S_{prey.inf}$) at varying sample size for non-selective predation (Case 1). The rows

904   indicate evenness and the columns represent original predation intensity. The red line

905   represents the zero line where overall and inferred prey species richness are the same

906   ($S_{prey.inf} = S_{prey}$). The grey dots and bars represent the mean and standard deviation of

907   the simulated differences for specific model assemblages [Figure 7. Two columns;

908   Color]

910   Figure 8. Plot showing the difference between the original (S_{prey}) and prey species richness

911   ($S_{prey.inf}$) at varying sample size for selective predation of common species (Case 2).

912   The rows indicate evenness and the columns represent original predation intensity.

913   The red line represents the zero line where overall and inferred prey species richness

914   are the same ($S_{prey.inf} = S_{prey}$). The grey dots and bars represent the mean and standard

915   deviation of the simulated differences for specific model assemblages [Figure 8. Two

916   columns; Color]

918   Figure 9. Plot showing the difference between the original (S_{prey}) and prey species richness

919   ($S_{prey.inf}$) at varying sample size for selective predation of rare species (Case 3). The

920   rows indicate evenness and the columns represent original predation intensity. The

921   red line represents the zero line where overall and inferred prey species richness are

922   the same ($S_{prey.inf} = S_{prey}$). The grey dots and bars represent the mean and standard

923   deviation of the simulated differences for specific model assemblages [Figure 9. Two

924   columns; Color]

926   Figure 10. Plot showing variation in inferred estimates of drilling and durophagous

927   predation with varying degrees of sampling for four Pleistocene molluscan

928   assemblages of Florida with different evenness (E_T). The top row represents the

sample size variation in inferred predation intensity (PI_{inf}). The bottom row shows

the inferred number of prey species ($S_{prey.inf}$) with varying sample sizes. The warmer

colours represent assemblages of higher evenness. [Figure 10. Two columns; Color]

Figure 11. Flowchart of the general framework of proposed method of the post-hoc

standardization. [Figure 11. Two columns; Grayscale]

Supplementary materials

Figure S1. The plot showing variation in inferred predation intensity ($PI_{T.inf}$) and inferred the

number of prey species ($S_{prey.inf}$) with specific sample sizes for different model

assemblages. The rows represent different degrees of selectivity of predation and the

columns indicate predation intensity in the original assemblage (PI_T). The warmer

colors represent higher evenness. [Figure S1. Two columns; Color]

S2. This is a compiled datasheet used for analyzing the predation estimates across four Plio-

Pleistocene fossil assemblages of Florida (Chattopadhyay and Baumiller, 2010)

S3. This is the R-script used for the present study. The required data is available at S2.

Table 1. A summary of the model assemblages used for this study with varying evenness, predation intensity and predator preference.

Evenness	Structure	PI_T for Case 1 (Preys of all species are attacked with a probability of 0.2, 0.5 and 0.8 for low, medium and high PI_{prey}, respectively)			PI_T for Case 2 (Preys of only common species are attacked with a probability of 0.2, 0.5 and 0.8 for low, medium and high PI_{prey} respectively)			PI_T for Case 3 (Preys of only rare species are attacked with a probability of 0.2, 0.5 and 0.8 for low, medium and high PI_{prey} respectively)		
		Low	Medium	High	Low	Medium	High	Low	Medium	High
$E_T = 0.2$	$N_{(S=1:29)} = 10$, $N_{(S=30)} = 2710$ [1*2710 + 29*10] = 3000	0.2	0.5	0.8	0.18	0.45	0.72	0.02	0.05	0.08
$E_T = 0.5$	$N_{(S=1:3)} = 910$, $N_{(S=4:30)} = 10$ [3*910 + 27*10] = 3000	0.2	0.5	0.8	0.18	0.46	0.73	0.02	0.05	0.07
$E_T = 0.7$	$N_{(S=1:5)} = 500$, $N_{(S=6:30)} = 20$ [5*500 + 25*20] = 3000	0.2	0.5	0.8	0.17	0.42	0.67	0.03	0.08	0.13
$E_T = 1$	$N_{(S=1:30)} = 100$ [30*100] = 3000	0.2	0.5	0.8	NA	NA	NA	NA	NA	NA

Table 2. A summary of the predation data from four Plio-Pleistocene fossil assemblages of Florida.

Locality	Evenness (E_T)	Sample size	S_T	Drilling frequency	$S_{prey.drill}$	Repair scar frequency	$S_{prey.repair}$
Miami Canal	0.31	4794	7	0.16	7	0.01	6
Punta Gorda	0.47	2417	5	0.18	5	0.01	4
Chiquita	0.74	894	7	0.04	6	0.14	7
McQueen's pit	0.87	657	6	0.15	5	0.03	6

Table 3. The result of Spearman rank order correlation test for proportional abundance and PI_{prey} for the predation estimates across four Plio-Pleistocene fossil assemblages of Florida (Chattopadhyay and Baumiller, 2010). The statistically significant ($p<0.05$) results are marked in bold.

Predation	Location	rho	p	Inferred scenario
Drilling	Punta Gorda	0.87	0.05	Case 2
	McQueen's pit	0.83	0.06	Case 1
	Chiquita	0.68	0.08	Case 1
	Miami canal	**0.99**	**<0.001**	-
Durophagy	Punta Gorda	0.21	0.74	Case 1
	McQueen's pit	0.46	0.35	Case 1
	Chiquita	0.24	0.61	Case 1
	Miami canal	**0.79**	**0.03**	Case 2

Table 4. The test-statistic (D) of Kolmogorov–Smirnov test comparing the predation estimates across four Plio-Pleistocene fossil assemblages of Florida using sample-standardization protocol. All the results are statistically significant ($p < 0.05$).

Estimate	Predation	Location	McQueen's pit	Chiquita	Miami Canal
Predation intensity	Drilling	Punta Gorda	**0.8**	**1**	**0.53**
		McQueen's pit		**0.24**	**0.31**
		Chiquita			**1**
	Durophagy	Punta Gorda	**0.9**	**1**	**0.19**
		McQueen's pit		**0.96**	**0.85**
		Chiquita			**1**
Prey species richness	Drilling	Punta Gorda	**0.38**	**0.31**	**0.13**
		McQueen's pit		**0.24**	**0.36**
		Chiquita			**0.29**
	Durophagy	Punta Gorda	**0.84**	**1**	**0.41**
		McQueen's pit		**1**	**0.95**
		Chiquita			**1**

Table 5. A summary of the difference in inferred predation intensity from the original value for the model assemblages. Each cell contains information about the mean value and standard deviation of $Diff_{PI}$; the first two represents the sign and magnitude of the mean value. A positive mean value of $Diff_{PI}$ indicates a larger value of original than inferred predation intensity ($PI_T > PI_{T.inf}$).

Evenness	$Diff_{PI}$ for Case1			$Diff_{PI}$ for Case2			$Diff_{PI}$ for Case3		
	Low	Medium	High	Low	Medium	High	Low	Medium	High
$E_T = 0.2$	+ve, <0.001, 0.013	+ve, <0.001, 0.016	+ve, <0.001, 0.013	-ve, <0.001, 0.012	-ve, 0.001, 0.015	-ve, 0.003, 0.014	+ve, <0.001, 0.004	+ve, 0.002, 0.006	+ve, 0.002, 0.008
$E_T = 0.5$	+ve, <0.001, 0.013	+ve, <0.001, 0.016	+ve, <0.001, 0.013	-ve, 0.002, 0.012	+ve, 0.005, 0.015	+ve, 0.002, 0.014	+ve, 0.002, 0.004	+ve, 0.005, 0.006	-ve, 0.002, 0.008
$E_T = 0.7$	+ve, <0.001, 0.012	-ve, <0.001, 0.015	-ve, <0.001, 0.012	+ve, 0.003, 0.012	+ve, 0.004, 0.016	+ve, 0.003, 0.016	-ve, 0.003, 0.006	-ve, 0.003, 0.009	-ve, 0.003, 0.011
$E_T = 1$	-ve, <0.001, 0.013	+ve, <0.001, 0.015	+ve, <0.001, 0.013	NA	NA	NA	NA	NA	NA

Table 6. A summary of the difference in inferred prey species richness from the original value for the model assemblages. Each cell contains the minimum sample size required for Diff_S to converge to zero for each model assemblages. A smaller number indicates that the inferred prey species richness converges to the original value ($S_{prey.inf} = S_{prey}$) at smaller sample size.

Evenness	Required sample size for convergence of Diff_S for Case 1			Required sample size for convergence of Diff_S for Case 2			Required sample size for convergence of Diff_S for Case 3		
	Low	Medium	High	Low	Medium	High	Low	Medium	High
$E_T = 0.2$	3000	2800	2100	100	100	100	3000	2700	2200
$E_T = 0.5$	3000	2700	2100	100	100	100	3000	2700	2200
$E_T = 0.7$	2900	1900	1400	300	100	100	2800	1900	1400
$E_T = 1$	1200	700	400	NA	NA	NA	NA	NA	NA