**Title:**

Sustained spinal motor activity triggered by direct mechanosensory stimulation in adult

*Drosophila melanogaster*

Alexandra M. Medeiros[1], Anna F. Hobbiss[1,2], Gonçalo Borges[1], Marta Moita[2], César S.

Mendes[1, 3]

1. iNOVA4Health, NOVA Medical School|Faculdade de Ciências Médicas, NMS|FCM,

Universidade Nova de Lisboa; Lisbon, Portugal

2. Champalimaud Research, Champalimaud Center for the Unknown, Lisbon, Portugal

3. Lead contact

Correspondence: cesar.mendes@nms.unl.pt

**Abstract**

Most vertebrates and invertebrates such as *Drosophila melanogaster* are able to move in

complex environments due to their ability to integrate sensory information along with

motor commands. Mechanosensory structures exist along each leg to assist in motor

coordination by transmitting external cues or proprioceptive information to motor centers

in the central nervous system.

Nevertheless, how different mechanosensory structures engage these locomotor centers

and their underlying circuits remains poorly understood. Here, we tested the role of

mechanosensory structures in movement initiation by optogenetically stimulating specific

classes of leg sensory structures. We found that stimulation of leg Mechanosensory

Bristles (MsB) and femoral Chordotonal Organ (ChO) is sufficient to initiate forward

movement in immobile animals. We also found that while the stimulation of the ChO

depends on the presence of the central brain to trigger forward movement, stimulation of

leg MsB could trigger cyclic motor activity independently of central brain circuits. Leg

MsB-triggered movement lacks inter-leg and intra-leg coordination, whilst antagonistic

muscle activity within joints is preserved. Moreover, we found that the duration of the

MsB-induced movement is dependent on the number of excited cells and is specific

to leg afferents, since stimulation of MsB in other segments lead to different motor

outcomes. This study sheds light on the ability of specific sensory circuits to modulate

motor control, including initiation of movement, and how this information can be

automatically translated to a motor command by spinal neuronal pathways.

## Introduction

A remarkable feature of moving organisms is their ability to adapt to external environmental conditions, optimizing speed, stability and energy consumption (Alexander, 2003; Dickinson et al., 2000; Hoyt and Taylor, 1981). Key for this flexibility is mechanosensory feedback, including proprioception and exteroception, enabled by specialized neurons that relay internal and external physical features such as muscle extension, tissue compression or gravitational orientation to motor centers. Lack of sensory feedback renders animals highly uncoordinated, largely determined by the number and class of inactivated cells (Akay et al., 2014; Mendes et al., 2013). Sensory structures vary in shape and properties depending on the type of mechanical feature being transduced (Smith and Shepherd, 1996). For example, in insects, the Chordotonal organ (ChO) relays the mechanical features of joint angles, a structure analogous to mammalian muscle spindles (Tuthill and Wilson, 2016b), whereas Campaniform Sensilla (CS), dome-like structures in the cuticle of the fly leg, sense deformation caused by load (Akay et al., 2001; Tuthill and Wilson, 2016b).

While dendrites of proprioceptors are located in the leg segment containing the sensory cell, axons project into the central nervous system, mostly into the ventral-most layer of the leg neuropil within the Ventral Nerve Chord (VNC), the insect equivalent of the mammalian spinal cord, where sensory information is processed (Court et al., 2020; Merritt and Murphey, 1992; Murphey et al., 1989; Tsubouchi et al., 2017; Tuthill and Wilson, 2016a). Sensory structures interact with local circuits in the VNC in order to modulate and adjust locomotor patterns. However, to generate fully functional and coordinated locomotion, communication between motor circuits and higher order centers is required (Ritzmann and Buschges, 2007). Evidence shows that locally these structures can directly trigger muscle contraction to correct movement and posture. For instance, ChO receptors that sense the position of the femur-tibia joint react to an imposed extension of the tibia by inducing excitation of flexor tibiae motor neurons and at the same time activate local interneurons that inhibit the antagonistic extensor tibiae motor neurons (Burrows et al., 1988). Another example of local modulation by sensory inputs comes from the stick insect, where it was shown that sensory feedback, notably from the

ChO and CS, assists in phase transition from stance to swing phase, a process termed
"active reaction", which influences the activation threshold by flexion and extensor motor
pools providing a positive reinforcement of movement (Akay and Büschges, 2006;
Bassler and Buschges, 1998). Further studies suggest the existence of a local neuronal
architecture within the VNC linking particular sensory structures to Central Pattern
Generators (CPGs) and leg motor neurons to produce inter-segmental coordination
(Borgmann et al., 2011; Tuthill and Wilson, 2016b). These features allow the generation
of a continuous, stable and coordinated walking pattern. In addition, this model depicts
the role of sensory information not just as correcting or inducing movement via supra
spinal motor centers, but also suggests that sensory input can directly engage local
circuits triggering or sustaining step cycling.
Another crucial mechanosensory structure are the Mechanosensory Bristles (MsB). These
structures are involved in detecting external cues, including dust particles, parasites and
wind intensity, triggering varied motor outcomes such as grooming, defensive kicking or
locomotion arrest (Li et al., 2016; Mamiya et al., 2018; Seeds et al., 2014; Tsubouchi et
al., 2017). In the context of collective behavior, mechanosensory bristles play a role in
conspecific avoidance by initiating walking responses triggered by another fly's touch
(Ramdya et al., 2015). An avoidance response was also seen after the optogenetic
stimulation of mechanosensory bristles in specific legs during walking (DeAngelis et al.,
2020). Given that self-generated movement also results in bristle deflection, these
structures may also play a role in proprioception, as well as exteroception (Tuthill and
Wilson, 2016a).
However, despite their ethological importance, the downstream circuits that integrate
information from mechanosensory bristles, and transform it into the appropriate motor
response, are unknown. Towards this end, we aimed to understand mechanosensory
integration within downstream motor circuits by asking if a particular class of leg sensory
neurons, the Mechanosensory Bristles (MsB), could elicit movement, and characterizing
the response in comparison with the ChO. For this we took advantage of the rich
*Drosophila* genetic tool-kit and kinematic quantitative tools, to address these questions
with cellular resolution and detail.

Here we show that stimulation of MsB or ChO is sufficient to initiate forward movement in an immobile fly. Moreover, we found that while leg ChO stimulation depends on the presence of a central brain for flies to engage in forward movement, leg MsB stimulation could trigger sustained motor activity independently of central brain activity. Leg MsB-triggered forward movement lacks inter and intra-leg coordination although antagonistic muscle activity is nevertheless preserved. Moreover, we found that the MsB-evoked response can be triggered from different leg segments and its intensity depends on the number of stimulated bristles. This response is also specific to leg MsB since stimulation in the wing or head lead to different motor outcomes.

This study sheds light on the ability of specific sensory circuits to modulate motor control, including initiation of movement, and how this information can automatically be translated to spinal motor programs. Moreover, it suggests how and what parameters the central brain can control during forward coordinated locomotion.

**Results**

**Activation of leg mechanosensory bristles evokes central brain independent movement**

To understand the role of leg sensory afferents in movement control and initiation we identified fly lines that targeted the different types of mechanosensory neurons present in the leg. For that, we performed an expression screen for GAL4 lines from the Janelia Flylight Collection that labelled specific sensory structures in the fly leg (Jenett et al., 2012; Pfeiffer et al., 2010). In order to restrict GAL4 expression to the legs, we used a combinatorial approach with flipase under the control of the Dac$^{RE}$ enhancer fragment (Mendes et al., 2013). Among the relevant lines and based on the expression pattern we selected 2 lines, one labelling Mechanosensory Bristles (hereafter termed MsB1) and one labelling the femoral Chordotonal Organ (termed ChO1) (Fig. 1A, B, and S1). Although normally classified as exteroceptors, MsB can additionally be described as proprioceptors since these afferents can also be activated during self-movement, which is supported by the observation that their inactivation causes perturbations in walking patterns (Tuthill and Wilson, 2016a) (Medeiros and Mendes, unpublished). Moreover, we chose ChO lines due to the large body of work focusing on this structure (Agrawal et al., 2020; Chen et al., 2021; Cheng et al., 2010; Jarman et al., 1993; Mamiya et al., 2018; Mendes et al., 2014; Shanbhag et al., 1992; Smith and Shepherd, 1996; Tuthill and Wilson, 2016b).

We tested the expression patterns of the selected lines in sensory neurons by visualization of mVenus tagged to the light-gated channel CsChrimson used in subsequent experiments (Wu et al., 2016). We observed that MsB1 neurons project to the ventral-most layer of the three leg Neuropils, the ventral association center (VAC) (Court et al., 2020) (Fig. S1). Conversely, the medial ventral association layer (mVAC) and the intermediate Neuropil (IntNP) received projections from ChO1 line neurons (Court et al., 2020) (Fig. 1 and S1). In the central brain we identified ChO1 axon terminals in the Wedge (WED) and AMMC consistent with the previously identified ChO primary neuron projections to the central brain (Fig. S1). The MsB1 line showed some labelling in the Antennal lobes and GNG, possibly axon terminals from head appendages (Gao et al., 2000; Kain and Dahanukar, 2015; Tsubouchi et al., 2017). After close inspection of the head appendages

we found cell labelling in the maxillary palps, in the proboscis and in the 3$^{rd}$ antennal

segment (Fig. S1 and data not shown).

Next we measured the behavioral responses of immobile flies to optogenetic stimulation

of these 2 classes of sensory neurons. For this we drove the expression of the light-gated

CsChrimson channel in the respective structures and analyzed the flies' behavior after a

single 100 ms stimulation in an open field arena (see Methods). While control Empty-

GAL4 flies remained immobile immediately after stimulation, eventually resuming motor

activity, the stimulation of leg mechanosensory Bristles or the leg femoral ChO led to

movement (Fig. 1D and Video S1). Motor responses differed between these two lines,

with MsB1 stimulation inducing a very short latency response, reaching a maximum

speed of 7.38 ± 1.17 mm/s, 60 ms after stimulus onset, while ChO1 stimulation led to a

more progressive response reaching maximum speed of 6.98 ± 1.99 mm/s after 1.6

seconds (Fig. 1D' and D'', upper panels). Moreover, MsB1 stimulation triggered a fast

startle response followed by sustained walking (Fig. 1D').

The different responses to optogenetic stimulation between ChO and MsB stimulation

suggested that these neurons promote movement by different neuronal circuits. We then

asked if movement initiation was dependent on descending information, by decapitating

the animals before optogenetic stimulation (Fig. 1E). Decapitated Drosophila are capable

of maintaining an upright posture and display innate behaviors such as grooming

(Hampel et al., 2017; Vandervorst and Ghysen, 1980).

Control animals (empty-GAL4) did not exhibit any optogenetic response, maintaining an

upright posture with occasional grooming behavior (Fig. 1E and data not shown).

Strikingly, MsB1 stimulation triggered a startle response and sustained motor activity for

approximately 2.9 seconds (Fig. 1E' and Video S1). Moreover, under these conditions,

animals reached a peak speed of 6.85 ± 1.27 mm/s, 60 ms after stimulus onset, similar to

the one observed in head-intact animals, indicating that the initial response to

mechanosensory stimulation is mediated by circuits located within the VNC, independent

of descending circuits. In contrast, ChO1 stimulation in beheaded flies did not induce

movement initiation (Fig. 1E''), indicating that these sensory neurons activate central

brain circuits that in turn are responsible for walking initiation. Optogenetic stimulation

of ChO1 in decapitated flies did however lead to a reflexive leg twitch, which induced a

small increase in the perceived instantaneous speed (arrow in Fig. 1E''and Video S**1**).

This reflex is most likely the consequence of direct synaptic connections between the

ChO and leg motor neurons, as described in locusts (Burrows, 1996; Burrows et al.,

1988; Kittmann and Schmitz, 1992; Tuthill and Wilson, 2016a).

These results indicate that stimulating mechanosensory structures in the leg is sufficient

to induce walking initiation, which in the case of MsB only depends on local circuits

within the VNC. However, sustained walking after MsB stimulation requires descending

activity from the head.

**MsB stimulation evokes movement encompassing a non-coordinated phase**

Having shown that MsB stimulation triggers walking response in head intact flies, we

next investigated the kinematic features associated with this motor response. For this we

used the FlyWalker system that allows the quantification of kinematic parameters with

high spatial and temporal resolution generating a large set of outputs including step,

spatial and gait parameters (Cabrita et al., 2022; Mendes et al., 2013; Mendes et al.,

2014)**.** The walking chamber was modified in order to accommodate a far-red LED array

to perform optogenetic stimulation where each fly received one pulse of 100 ms red light

stimulation and the resulting behavior was quantified.

Control animals display a typical gait pattern showing an alternating tripod configuration

(Fig 2A) and consistent stance traces (Fig 2C), which mark the tarsal contacts relative to

the body axis during stance phases, indicating high levels of leg coordination under these

conditions (Burrows, 1996; Mendes et al., 2013; Strauss and Heisenberg, 1990; Wosnitza

et al., 2013)**.** Forward movement induced by leg MsB stimulation can be classified into

two distinct phases using visual inspection of the step patterns: an early *uncoordinated*

*phase* followed by a late *coordinated phase* (Fig. 2B/D and B'/E, respectively; and Movie

S2). The *uncoordinated phase* immediately follows optogenetic stimulation, displaying

an apparent random step pattern without any clear coupling between the six legs, leading

to an inefficient force transduction that results in a slow and non-cyclic movement (Fig.

2B). We further analyzed the step configurations, a proxy for inter-leg coordination, and

found a large proportion of frames that do not match any of the normal gaits or

configurations described by control flies, such as tripod or tetrapod (Mendes et al., 2013;

Strauss and Heisenberg, 1990; Szczecinski et al., 2018; Wosnitza et al., 2013) (Fig. 2B,

lower panel). We describe these configurations as '*non-compliant*' since these do not

conform with the rules of coordinated movement described by Durr et al. (Durr et al.,

2004). Non-compliant configurations include, for example, two contralateral or two

consecutive ipsilateral legs lifted simultaneously, which creates static instability. We

observed that non-compliant configurations are used to a striking degree during the

uncoordinated phase, ranging from 20% to 50% (average of 29.6±4.4%) of the frames

compared to being nearly absent in control animals (Fig. 2F and H). This shift in gait

pattern is matched by a reduction in the use of the canonical tripod and tetrapod gaits

(Fig. 2H). These data suggest that stimulation of leg MsB induces activation of motor

activity lacking inter-leg coordination. Moreover, stance traces display an extremely

uncoordinated pattern with highly variable and inconsistent leg contacts, highlighting a

lack of control and coordination during this phase of locomotion (Fig. 2B and D), even

though leg retraction shows the typical anterior to posterior movement (Fig. 2D).

Nevertheless, the high degree of uncoordinated movement suggests that in addition to a

lack of inter-leg coordination, the proper coupling between leg segments (or intra-leg

coordination) is lost during this initial phase of locomotion. We evaluated the whole

kinematic profile and found that virtually all kinematic parameters were altered, with the

notable exception of the average swing duration (Fig. 2H). Interestingly, we found that

animals shifted from an uncoordinated phase to a distinct coordinated phase without

requiring a transitional stop (Fig. 2B and B'). Nevertheless, during the transition most

flies displayed a period during which all legs were in a stance phase, contacting the

ground (19 out of 21, corresponding to 90.4%) (Fig. 2B, ~1.4s). From a total of 21 flies

tested, 10 (48%) shifted to coordinated walking without stopping (see for example, Fig.

2B to 2B'), 4 (19%) shifted to coordinated walking after a small pause of 200 to 400 ms

and 6 (29 %) of the flies initiated grooming for a significant period of time before

resuming walking. During the coordinated phase the fly adopts mainly tripod and

tetrapod gaits while non-compliance configurations are absent (Fig. 2B', lower panel).

Once animals have returned to a coordinated phase, both gait patterns and stance traces

display a pattern similar to control animals (Fig. 2H, 2B' and E). Kinematic parameters in

the coordinated phase are more concordant with the standard walking pattern seen in

control animals although the recovery is not total, indicating that transient sensory
stimulation can have sustained effects on locomotion (Fig.2H, 2B').
In summary, stimulation of leg MsB results in a reflexive forward cyclic movement,
during which flies display an uncoordinated phase, lacking inter-leg coordination,
followed by a coordinated phase presenting the stereotypical properties of walking
animals.
244
**Kinematic features after leg MsB stimulation in headless flies**
We next investigated the kinematic features of headless flies upon leg MsB stimulation
(Fig. 3), which we previously found initiates locomotor behavior in the open arena setup
(Fig. 1 E). As described for Figure 2, we optogenetically stimulated leg MsB1 in an
enclosed arena and extracted kinematic features using the FlyWalker system (see
Methods). Prior to stimulation, we decapitated the flies and glued the heads back to the
thorax to allow proper body tracking by the FlyWalker software.
Stimulated flies display cyclic leg movements with a highly uncoordinated gait pattern
and stance traces (Fig. 3A, B). Moreover, kinematic properties also display a starkly
different profile compared to control animals (Fig. 3C). Noticeably, there is a similarity
between the walking pattern of the uncoordinated phase of the head intact flies and the
headless flies after stimulation (Fig. 3C). In fact, the stance straightness index is
indistinguishable between these two conditions (Fig. 3D, compare Fig. 2B' with 3B).
However, we observed that the time headless flies spend moving after the stimulus is
longer compared to the uncoordinated phase in head intact flies (Fig. 3E), suggesting that
descending circuits can interrupt the uncoordinated cyclic activity triggered by leg MsB
and mediate a transition to a coordinated walking pattern.
We then asked if descending control could influence the response to leg MsB stimulation
by comparing the early response in head intact animals to the response in headless flies.
First, we noticed a larger fraction of frames spent in a non-compliant configuration (Fig.
3F), which could be explained by a sustained swing phase during the initial stages of the
motor response (Fig. 3A, below ~1 sec, compare lower section, Fig. 2B with 3A).
Second, we subjected all the kinematic parameters to a Principal Component Analysis
(PCA) selecting the top 3 axes that explained most variance, thus yielding a three-

dimensional space representation (Fig. 3G). PC1, which explained the highest variance of

the data (50.2%), allowed a statistical discrimination between headless animals and the

uncoordinated phase in head-intact animals, with the cloud of points of headless animals

lying further away from control animals in the behavioral space (Fig. 3G' and G''),

suggesting that higher order circuits can influence the cyclic motor activity triggered by

leg MsB.

These data indicate that leg MsB can trigger sustained cyclic motor activity using solely

circuits within the VNC, albeit with severely impaired coordination between legs and leg

segments. This cyclic activity is nevertheless modulated and ultimately restrained by

central brain circuits and descending information.

**Inter-joint coordination is partially disrupted after MsB stimulation**

To further explore the disruption of coordinated locomotion following leg MsB

stimulation, we examined intra-leg, or alternatively called inter-joint coordination, i.e. the

correct modulation of joint angles within a single limb in relation to one another, required

to produce efficient walking (Cruse and Bartling, 1995). To extract measures of joint

angles we tethered flies expressing CsChrimson in leg MsB to a wire while they walked

freely on a polystyrene ball (Barrios et al., 2021; Buchner, 1976; Götz and Wenking,

1973; Seelig et al., 2010) (Fig 4A). As above, one 100ms light stimulation pulse was

delivered while the flies were immobile. Fly behavior was imaged using a lateral camera

at 120 fps and videos were tracked using DeepLabCut (Mathis et al., 2018), which

allowed us to extract 3 joint angles: 1) Coxa-Femur joint, 2) Femur-Tibia joint, 3) Tibia-

Tarsus joint (Fig. 4A, B and Video S3). We only considered angles from the T1 leg, since

this leg is predominantly at a perpendicular angle to the imaging path and thus the angles

could be accurately calculated in a 2D video. As expected from our earlier results, MsB

stimulation induced movement in both head-intact and headless flies (Fig. 4B, Video S3).

To determine whether inter-joint coordination was perturbed by the stimulation protocol

we compared 4 conditions: walking flies (*pre-stimulation*) (Fig. 4B, top panel); head-

intact flies for the initial 1s after stimulation onset (termed "*early response*") (Fig. 4B,

middle panel, red section); head-intact flies 3-5s after stimulation onset (termed "*late*

*response*") (Fig. 4B, middle panel, green section); and headless flies for the first 1s after

stimulation onset (termed "*headless response*") (Fig. 4B, lower panel, purple section). We chose these time periods to capture the 'uncoordinated' and 'coordinated' phases of post-stimulation movement previously described (Figs. 2 and 3).

As expected, *pre-stimulation* walking flies showed stereotyped periodic patterns of joint angles (Feng et al., 2020) (Fig 4B). The Coxa-Femur joint and the Femur-Tibia joint were strongly modulated throughout the step-cycle in a correlated fashion, whilst the Tibia-Tarsus joint did not show an obvious correlation to the step cycle. We performed a Fast Fourier Transform (FFT) to extract component frequencies from the data (Fig. 4C). In *pre-stimulation* walking flies, the Coxa-Femur and Femur-Tibia joints showed the highest power at 7.0 Hz, corresponding to the average step frequency (Fig. 4C and Fig. S3A), while the Tibia-Tarsus joint was not periodically modulated (Fig. S3A). FFT analysis of the *early-response*, *late-response* and *headless response* also showed that Coxa-Femur and Femur-Tibia joints were more strongly modulated than the Tibia-Tarsal joint (Fig. S3A). We thus restricted our subsequent analysis to only Coxa-Femur and Femur-Tibia joints. To understand the regularity of motor behavior following stimulation we compared the FFT of our 4 conditions for the Coxa-Femur joint (Fig. 4C). Whilst in the intact-early stage the strong periodic modulation of joint angles seen in walking flies is lost, by the late stage, a peak has re-emerged (albeit around 5.2 Hz, a lower frequency than in the walking condition), showing the angles traces are once more becoming structured. Strikingly in headless flies, frequencies were highly variable with no clear peaks (Fig. 4C).

To further investigate the relationship between joints, we generated angle-angle plots between the Coxa-Femur joint and the Femur-Tibia joint for our 4 conditions (Fig. 4D). Angle-angle plots for walking flies showed a half moon-shaped 'ring' trajectory (Fig. 4D), corresponding to the stance and swing phases of the step cycle (Fig. S3B). Angle correlations were altered in the *early response* following stimulation (Fig. 4D). Whilst the half-moon outline was still visible, indicating that the fundamental relationship between joints has not been totally lost, the variability of the movement was greatly increased. Under *late response* conditions (i.e. 3-5 seconds after the stimulation) traces once more display a regular 'walking-like' ring (Fig. 4D). However, during *headless response* conditions, very distorted angle-angle traces are visible with an undetectable ring pattern.

To quantify dispersion from the average walking cycle, we calculated the mean trace of
the walking angles (Fig. 4D, black lines and Fig. S3B). We then analyzed the minimum
distance for each measured point to the mean trace (Fig. 4E, F). Distances were
significantly increased for all stimulated conditions as compared to pre-stimulus
condition (Fig. 4F), indicating a significant disruption of the co-ordination between joints
after MsB stimulation ($P_{walking-early\ response} < 0.001$, $P_{walking-late\ response} < 0.001$, $P_{walking-decapitated}$
$< 0.001$, Dunn's test). Distances were reduced from the *early response* to the *late*
*response*, indicating a partial recovery of coordination ($P_{early\ response-late\ response} < 0.001$,
Dunn's test). In contrast, distances were greatly increased in the headless condition,
indicating that loss of brain control leads to wildly disrupted relationships between joint
angles ($P_{early\ response-decapitated} < 0.001$, $P_{late\ response-decapitated} < 0.001$, Dunn's test). To
understand which aspect of the relationship between angles was disrupted, we divided the
angle-angle space into 4 quadrants (described in Fig. S3C). The positions of the lines
were chosen to split the walking traces into 4 components of the step cycle – stance
onset/quadrant 1 (corresponding generally to high Coxa-Femur angles); stance
offset/quadrant 2 (corresponding to low Femur-Tibia angles); swing onset /quadrant 3
(corresponding to low Coxa-Femur angles); and swing offset/quadrant 4 (corresponding
to high Femur-Tibia angles). We calculated the proportions of data points which fell in
each quadrant for each condition (Fig. 4G). Pre-stimulus walking flies spend more time in
quadrants 1 and 2, corresponding to the slow stance phase, and less time in quadrants 3
and 4 which correspond to the faster swing phase. Flies immediately after stimulation
show an increase in occupancy of quadrant 3, likely corresponding to aborted swing
initiations. Strikingly, headless flies show a large over-representation of points in
quadrant 4, corresponding to extended Femur-Tibia angles, indicating that without
descending control, extension is the default response, indeed to an extent where the flies
almost invariably let the ball fall (Video S3). This effect was also observed in the gait-
analysis experiments (Fig. 3 and Video S2).
Overall, our data show that inter-joint coordination is partially lost immediately after
stimulation in head-intact flies, but has begun to recover by the period of 3 – 5 seconds
after stimulation. In contrast, joint angles are highly uncoordinated following stimulation
in headless flies, which show large overextensions of the Femur-Tibia joint, suggesting a

role for descending control to appropriately regulate the relationships of joint angles to

one another.

**Intra joint muscle coordination is maintained during movement triggered by MsB**

Key for coherent locomotor activity is the reciprocal inhibition of opposing flexor and

extensor muscles, each one controlled by their respective excitatory motor neuron pool

(Eccles et al., 1956; Grillner, 1975; Sherrington, 1893). This implies that contraction of

one joint muscle is concurrently matched by relaxation of its antagonistic muscle within

the same joint. We asked if sensory-triggered movement also follows this muscle pattern,

which we refer as intra-joint coordination. For this we monitored muscle contraction at

the femur-tibia joint by measuring GCaMP fluorescence in leg muscles while

optogenetically stimulating Leg MsB (Fig. 5A) (Heredia et al., 2021; Soler et al., 2004,

see Methods). The fly's body and legs were glued to a coverslip (Kamikouchi et al.,

2010), and muscle activity in the femur was recorded using a spinning disc confocal

while the fly was stimulated with a 100ms light stimulus (see Methods). We identified the

femur extensor and flexor muscles (Fig. 5B) and quantified the percentage of

fluorescence in each frame, normalized to the maximum and minimum of the

fluorescence for each video (Fig. 5B-D) (see Methods). In addition, we generated a

kymograph covering both extensor and flexor muscles (Fig. S4). As expected in head

intact flies, both controls groups (flies without stimulation and stimulated without all-

trans-retinol (ATR)-supplemented food) display alternating muscle activity, meaning that

antagonistic muscles were not active simultaneously (Fig. 5C, left and middle panel and

kymograph on Fig. S4B). We calculated the Spearman correlation between the two

muscle signals over time, which generated negative values confirming their anti-phasic

relation (Fig. 5E). Optogenetic stimulation of leg MsB showed a fast muscle response in

accordance with the locomotor behavior previously observed. Interestingly, the observed

muscle activity maintained the antagonistic alternation seen in control flies (Fig. 5C right

panel, E and Fig. S4B). Stimulation of headless flies also led to an immediate muscle

contraction showing an alternating antagonistic muscle activity, albeit with a higher

variability (Fig. 5D, E). Notably, both in head-intact and headless animals, we found no

flexor or extensor muscle bias upon sensory stimulation (Data not shown), suggesting

that no particular muscle was more responsive to sensory stimulation and that indeed a

reciprocal motor pool inhibition mechanism is present to prevent muscle co-contraction

upon stimulation (Brown, 1911 ; Eccles et al., 1956). We also compared the GCaMP

fluorescence profile during muscle contraction by measuring peaks intensities and widths

(Fig. 5F-H, see Methods). Noticeably, stimulation of leg MsB in headless flies led to a

weaker muscle contraction pattern compared to head intact conditions (Fig. 5G),

suggesting that higher order centers may act as modulators of motor neuron recruitment

and activity. However, no differences were observed in the width of the muscle

contraction peaks between head intact and headless flies (Fig. 5H).

Overall, our results show that leg MsB stimulation evoked movement largely maintains

intra-joint muscle coordination properties in head-intact and headless flies.

**Cyclic movement triggered by MsB is leg specific**

Full body stimulation of mechanosensory bristles (using dust particles or genetically) has

been shown to trigger grooming behavior (Hampel et al., 2017; Seeds et al., 2014), while

in our study where we limit activation to just leg MsB, we found mostly uncoordinated

walking to be the main response. It has been suggested that compartmentalized sensory

connections mediate different behavioral responses (Siegler and Burrows, 1986;

Vandervorst and Ghysen, 1980). We thus asked if the motor response could be

determined by the body structure from which the stimulus originates. To target

expression of CsChrimson to MsB in other body segments, we used the same GAL4

driver but instead combined with a head capsule or wing-specific flp line using the

eyeless (ey) and vestigial (vg) fragments, respectably (Fig. 6A, see methods). We

confirmed the effectiveness of this approach based on the presence of positive cells in the

respective cuticle and axonal projections (Fig. S1). We then cataloged the repertoire of

motor behaviors in response to optogenetic stimulation using manual and automatic

classifiers (see Methods). These included uncoordinated and coordinated movement, leg

cycling and grooming.

As previously shown, leg MsB activation triggered uncoordinated displacement,

immediately followed by coordinated walking and grooming in some animals (12 out of

20) (Fig. 6B). Notably, wing and head stimulation did not induce uncoordinated

movement, with animals instead displaying coordinated walking (Fig. 6C-E), showing
that the uncoordinated activation of leg motor centers is specific to leg afferents.
Moreover, stimulation of wing MsB led only a small fraction of the animals to groom (2
out of 20) compared to leg or head stimulation (11 and 10 out of 20, respectively) (Fig.
6B-E), further highlighting the specificity of sensory stimulation for motor selection. This
suggests that behavior selection (in this case grooming) is dependent on the place where
the stimulus is received.

We then asked if the aforementioned behaviors are dependent on the presence of higher-
order brain commands. As observed previously, leg MsB stimulation in headless animals
caused uncoordinated body displacement (Fig. 6F and H). Additionally, close inspection
of the movies revealed an extra behavior, which is interestingly absent in head intact
animals: sustained random leg movements (Figs. 6F, brown sections and Video S4).
These movements do not display any synchrony between legs, thus rendering the fly
immobile.

Our data also show that the walking behavior triggered by the stimulation of wing MsB is
dependent on the presence of higher-order brain commands. Stimulation of wing MsB
afferents in headless flies did not induce walking or uncoordinated movement (Fig. 6G,
H), but instead grooming, (a behavior which is absent in head intact conditions (Fig. 6C),
suggesting a descending inhibitory command that blocks the grooming behavior). Video
analysis of headless flies showed that the observed leg movement bouts resembled the
preparatory pose for grooming (data not shown), or in some cases a defensive kick
behavior described previously to remove fictitious parasites (Li et al., 2016) (Video S4).

Overall, our data show that the motor response triggered by sensory stimulation depends
not only on the class of proprioceptors (MsB), but also on the source of the stimulus.
These data also highlight the role of the central brain controlling VNC pre-motor centers,
while executing particular leg motor commands such as leg twitching or grooming.

**Mechanosensory-triggered movement is dependent on the number of cells**

Leg mechanosensory bristles are present on the surface of the entire leg of the fly, with
neurons present in the most distal part of the leg projecting their axons to the central part

of the leg neuropil while neurons occupying proximal positions project to more

peripheral areas (Tsubouchi et al., 2017). Leg MsB display a concentric organization in

the leg neuropil instead of a concentrated topographical distribution within the VNC

(Tsubouchi et al., 2017). Since we showed that uncoordinated movement is evoked

specifically by leg MsB, we conjectured the possibility that a discrete subpopulation of

leg cells could be sufficient to activate the downstream motor program that triggers cyclic

movement, or alternatively, that the response results from the contribution of all leg MsB

being activated. To test these possibilities, we used 6 additional GAL4 lines to drive the

expression of CsChrimson in different bristles subpopulations in the fly leg, which we

called MsB2 to 7 (with our original line termed MsB1), ordered by the number of cells

labeled, with MsB1 the most and MsB7 the least. For each line we quantified number of

MsB labeled and looked at their localization in specific leg segments (Fig. 7A, B and

Table S1). We also measured the response of the different lines to optogenetic

stimulation by registering the duration of motor activity and categorizing the different

motor outputs in an open field arena (Fig. 7C and D, respectively). As previously

described (Fig. 1) the MsB1 line, which expressed GAL4 in 169 bristles in all leg

segments, had a strong reaction to the stimulus, showing motor activity for up to 1.5

minutes (Fig. 7C). Interestingly, stimulation of a lower number of cells in MsB2-6, which

label between 57-26 neurons, still triggered a motor response but with a shorter time span

(Fig. 7C), whilst MsB7, which drives expression in just 4 cells in the tarsal segments,

triggered minimal motor activity, suggesting that the duration of movement is influenced

by the number of activated cells. To further analyze the type of behaviors triggered by

mechanosensory bristles stimulation, we used the same behavioral classifier previously

used to identify immobility, body displacement, grooming, and sporadic leg movement

(Fig. 7D, E and S5). We observed that most lines show uncoordinated walking behavior

followed by sporadic leg movement without displacement. Some flies also displayed

grooming behavior, consistent with previous observations (Hampel et al., 2017) (Fig. 7D,

E). We also considered if particular leg segments are more responsive to the optogenetic

stimulus. Although we could not find lines specific to any particular segment at

comparable numbers, we noticed that activating the 57 tibial MsB in line MsB2 triggered

a shorter and less robust response compared to other lines, which labelled fewer, but

more proximally located, cells such as MsB3 and MsB5 (Fig. 7C-E, S5C), hinting at the
possibility that cells in proximal segments can trigger a more robust response to
mechanosensory stimulation.
These results suggest that no particular population of MsB is uniquely responsible for
inducing VNC-dependent leg movement but instead the cyclic motor response is due to
the cumulative stimulation of multiple MsB broadly distributed circuit in the VNC.

**Discussion**

In this study we took advantage of the sophisticated *Drosophila* genetic toolkit and kinematic analysis tools to understand mechanosensory evoked motor responses in different body regions. We found that a short 100-millisecond (ms) optogenetic stimulation of two different leg mechanosensory structures is sufficient to initiate forward movement in immobile animals (Fig. 1). In particular, MsB stimulation led a two-phase motor response comprised by an immediate *uncoordinated phase* followed by a late *coordinated phase* (Fig. 2; Video2). While the coordinated phase shows all the fundamental features of normal locomotion, the uncoordinated phase lacks both inter and intra-leg coordination whilst sustaining antagonistic antiphasic muscle activity (Fig.2, 4 and 5). Moreover, the uncoordinated phase is mostly dependent on spinal circuits as leg MsB stimulated headless flies largely phenocopy the *uncoordinated phase* present in head-intact animals (Fig.3, 4 and 5). Finally, we show that the motor response to MsB stimulation is dependent on the segment and number of cells being stimulated (Fig. 6 and 7).

**Motor activity by targeting of Spinal Motor Circuits**

This study unveils a new circuit for direct activation of VNC locomotor neural networks. The stimulation of MsB leads to the direct engagement of spinal circuits without interference of the central brain, resulting in a forward movement that lacks inter-joint and inter-leg coordination but still maintains intra-joint coupling.

Previous studies have shown the generation of motor programs through the stimulation of sensory structures. For example, the mechanical and optogenetic activation of bristles in the entire body of the fly are able to elicit grooming movements (Hampel et al., 2017), whereas the activation of specific bristles on the wing margin, which could be caused by an invading mite, leads to the initiation of a directed kick towards the stimulated area (Li et al., 2016). Importantly, these sensory-triggered behaviors are still maintained in headless flies. These data highlight that the circuits required for the processing and execution of ethologically relevant behaviors are present in the VNC, which can be executed with remarkable speed and precision. Likewise, our data present evidence that

neural networks in the VNC are capable of processing leg mechanosensory feedback to
execute the appropriate cyclic motor program. Moreover, this response is not exclusively
due to the activation of MsB but also depends on the origin of the stimulus. Notably,
stimulation of wing MsB did not induce uncoordinated walking but mostly grooming and
kicking as previously described (Hampel et al., 2017; Li et al., 2016) (Fig. 6). These data
support that the location of the sensory input plays a major role in the selection of the
motor outcome, instead of solely the class of the mechanosensory structures conveying
the stimulus. Since the leg and wing MsB target distinct neuropil regions in the VNC,
namely the leg and wing neuropils (Tsubouchi et al., 2017) (Fig. S1), respectively, it is
thus likely that distinct and possibly non-overlapping motor circuits become active.
While the grooming and kicking behavior after bristle stimulation are important for the
fly to rid itself of invading parasites and debris, the uncoordinated forward movement
here does not follow the same purpose. Instead, this behavior could be an escape response
caused by inter-fly contacts or other imminent threats (Ramdya et al., 2015). This
response is analog to the jump reflex, a behavior triggered by visual stimuli that activates
the giant fiber – a large descending interneuron synapsing directly on the mesothoracic
neuromere (T2) motor neurons leading to a jump and flight takeoff (King and Wyman,
1980; Tanouye and Wyman, 1980; Trimarchi and Schneiderman, 1995)}. However, we
should point out an important distinction between giant fiber and leg MsB stimulation:
while stimulation of the giant fiber induces a reflexive tonic response from the ipsilateral
tergotrochanteral muscle (TTM) leading to a jump reflex (King and Wyman, 1980); MsB
stimulation leads to a motor program comprised of sustained alternating antagonistic
muscle activity.
545

**Sustained activity and segmental coupling**
546

In this study, the movement evoked by the stimulation of MsB shows characteristic CPG-
like activation, where antagonistic muscles maintain anti-phasic contraction leading to
joint movement (Büschges et al., 1995; Fuchs et al., 2011; Pearson, 1972; Ryckebusch
and Laurent, 1993) (Fig 5). The same behavior was observed in headless flies, implying a
lack of interference of ascending or descending neurons in the sensory-evoked
movement. These results are similar to the pharmacological stimulation of decapitated

flies with stimulatory amines, notably serotonin, dopamine, and octopamine, which lead
to locomotor bouts (Yellman et al., 1997). Moreover, studies in deafferented VNCs using
the muscarinic cholinergic agonist pilocarpine, have shown alternating and rhythmic
activity in antagonistic leg motor neuron pools driven by the CPGs (Bidaye et al., 2018;
Büschges et al., 1995; David et al., 2016; Fuchs et al., 2011; Goldammer et al., 2018;
Pearson, 1972; Ryckebusch and Laurent, 1993). Despite this large body of work, the
identity, location and architecture of the CPG population remains largely elusive. The
reason for this gap in knowledge is the difficulty in accessing the neuronal VNC
population, adding to the problems in separating the outputs from CPG and the activity in
neurons that process sensory information during stepping. Nevertheless, some
interneurons were identified as being part of this network in the cockroach and in the
stick insect (Bassler and Buschges, 1998; Pearson and Fourtner, 1975). The use of
mechanosensory stimulation in the fly provides an attractive port of entry in order to
further identify and perform physiological characterization of leg CPGs taking advantage
of the expanding Drosophila genetic toolkit (Meissner et al., 2022). The second-order
neurons from the mechanosensory fields remain elusive, although data from the locust
suggest that these afferents target spiking local interneurons, some of which synapse on
motor neurons (Siegler and Burrows, 1986). The reconstruction of serial-section electron
microscopy (EM) datasets from the *Drosophila* VNC will provide a detailed connectome
profile of the leg MsB (Phelps et al., 2021).

**Cyclic activity and intersegmental coordination**
Remarkably, we found reduced intersegmental coordination between hemi segments and
leg joints immediately after MsB stimulation (Figs 2-4), albeit with antagonistic muscle
contraction still being maintained (Fig. 5). These data resemble the effects of the
muscarinic agonist pilocarpine in deafferented insect VNC preparations, which generate
rhythmic activity of alternating motor neuron pools controlling joint movement (Baudoux
et al., 1998; Chrachri and Clarac, 1987; David et al., 2016; Goldammer et al., 2018;
Rillich et al., 2013; Ryckebusch and Laurent, 1993). Moreover, this rhythmic activity is
independent within the three leg joints, as these rarely exhibit coordinated activity
between them, meaning that different CPGs control the antagonistic activity of motor

neuron pools of each leg joint (Bidaye et al., 2018; Büschges et al., 1995; Daun et al.,
2019; Ryckebusch and Laurent, 1994). Given that in our experimental design, animals
still maintain their proprioceptive features that assist in segment coupling, we can
speculate that the MsB-dependent cyclic activity overrides or is insensitive to sensory
feedback or to any other coupling mechanism. Our data also suggest a role for
supraspinal feedback promoting segmental coordination since decapitation decreased the
degree of inter- and intra-leg coordination (Figs. 3F, G and 4F). Work on
hemimetabolous insects has shown a role for descending neurons (DNs) that traverse the
neck controlling the onset and features of walking patterns (reviewed by (Bidaye et al.,
2018; Strausfeld et al., 1984)). Moreover, optogenetic manipulation of defined
populations of DNs can trigger stereotyped behaviors and change walking patterns
(Cande et al., 2018; Namiki et al., 2018). Multisegmental coupling is provided either by
sensory feedback, descending inputs or centrally by nonspiking interneurons (Buschges,
1995; Juvin et al., 2012). The relevance of each component depends largely on the speed
the animal typically assumes, with slow-walking animals - such as the stick insect -
relying heavily on sensory feedback (Akay and Büschges, 2006; Akay et al., 2007;
Mantziaris et al., 2017). Regardless, our results suggest that, as in other insects,
alternating activity of antagonistic motor pools in *Drosophila* are uncoupled between
different joints and hemisegments, further reinforcing the use of *Drosophila* as a unique
model system to unravel the neuronal architecture responsible for coordinated motor
patterns.

**Implications for vertebrate systems /Clinical Applications**
Our study demonstrates that the stimulation of a specific sensory class has a direct effect
on the engagement of locomotor circuits in the VNC. These findings highlight the
possibility of using mechanosensory feedback circuits as a rehabilitation tool after spinal
cord injury. Regaining locomotor function after a spinal cord lesion requires the
reconnection of downstream locomotor circuits below the lesion and descending neurons
(Ballermann and Fouad, 2006; Bareyre et al., 2004; Rosenbaum et al., 2010). This can
happen through the growth and synaptic arborization of supraspinal and spinal projection
neurons leading to spontaneous recovery after incomplete spinal cord injury (Takeoka et

al., 2014). Post lesion studies show that mice deprived from sensory feedback exhibit poor reconnection of the descending circuits, leading to a lack of recovery and strong impairments in locomotion (Bareyre et al., 2004; Courtine et al., 2008; Takeoka et al., 2014). These loss-of-function results suggest that sensory feedback support the stabilization and rewiring of circuits deprived of upstream stimulation. Conversely, our data show that activation of specific sensory afferents are able to directly engage spinal motor circuits. Considering the evolutionary conservation of motor circuits, our results support the hypothesis that direct sensory electrical stimulation can foster the rewiring of injured circuits enhancing recovery outcomes (Capogrosso et al., 2013; Takeoka et al., 2014). In humans, targeted spinal cord stimulation improves locomotor performance in individuals with severe motor deficits or even leg paralysis (Angeli et al., 2014; Formento et al., 2018; Wagner et al., 2018). The direct stimulation of dorsal roots, where proprioception structures reside thus represents a more surgically accessible approach for the activation and recovery of paralyzed structures.

**Methods**

**Fly strains**

All flies used in this study were non-virgin females between 1 and 7 days post-eclosion.

Flies were housed in a 25° incubator with 50-70% humidity. Prior to experimentation, all

flies were manipulated under cold anesthesia and allowed to recover for at least 24h. The

GAL4 lines from the Janelia Flylight Collection were: R65D12-Gal4(attP2), R77E09-

Gal4(attP2), R85A07-Gal4(attP2), R45D07-Gal4(attP2), R54A11-Gal4(attP2), R39A11-

Gal4(attP2) and R64D07-Gal4(attP2), R50A05-Gal4(attP2), R22E04-Gal4(attP2),

R79E02-gal4(attP2) and pBDP-GAL4UW (w[1118];P{GAL4.1Uw}attp2) (EmptyGal4).

Ato1.9-flp and $Vg^{BE}$ -flp were generated by TOPO cloning into an entry clone the 1.9kb

enhancer fragment from the regulatory region of atonal (zur Lage et al., 2004), and the

boundary enhancer of vestigial ($Vg^{BE}$) (Williams et al., 1994; Zecca and Struhl, 2007) (a

gift from Mirian Zecca), respectively. Both entry clones were subjected to a Gateway

reaction into a flp destination plasmid. Transgenic lines were generated by standard

procedures in a *yw* background. The dac$^{RE}$-flp was used in previously (Mendes et al.,

2013). UAS-FRT-stop-FRT- CsChrimson-mVenus was a gift from Gerry Rubin (Wu et

al., 2016). UAS-FRT-stop-FRT-GFP, Lexop-GCaMP6f and ey-flp lines were obtained

from the Bloomington Stock Center. For the calcium imaging in leg muscles we used

lexO-GCamp6f/;mhc-lexA,lexO-Cherry/SM6ˆTm6b (Heredia et al., 2021) and the

recombinant line Dac$^{RE}$-flp;Uas-FRT-stop-FRT-CsChrimson-mVenus (VK5), R65D12-

Gal4 (attP2).

**Behavior Setup and optogenetic activation of mechanosensory structures**

The open field behavior arena for optogenetic stimulation was custom built from 3 mm

opaque white and black acrylic sheets using a laser cutter. The circular arena had a

diameter of 54mm, and was surrounded by 6 deep 655 nm red Leds (LUXEON Rebel

Led's) evenly distributed providing a stimulation power of 0.016 mW/mm$^2$ (Fig. 1C),

The edge of the arena was surrounded by water to prevent flies accessing the walls. The

top of the arena (transparent acrylic) was covered with Sigmacote (Sigma-Aldrich)**.** A

white LED array was placed under the arena to serve as backlight. Flies were imaged

from above using a camera **(**Flea3-USB3, Point Grey, Teledyne FLIR LLC**)** equipped

with a 12 mm Computar (Commack, NY) lens covered with a 600nm, OD 4.0 Shortpass

Filter (Edmund optics, Barrington, NJ) to block optogenetic stimulation light.

For the experiments in Figure 1, we used 1-7 day old females, comprising 20 control flies

(empty-GAL4), and 20 flies expressing CsChrimson in leg mechanosensory bristles. Flies

were bred and kept in food infused with all trans-Retinal (Final Concentration 0.4 mM

;R2500, Sigma-Aldrich). Single flies were aspirated into the chamber. We waited for a

spontaneous pause in behavior (no walking or grooming) and manually delivered a single

1-pulse stimulation of 100ms. Imaging continued for at least 20 seconds after stimulation.

For decapitated fly experiments, flies were anesthetized on a cold plate and their head

was cut off. After the surgery flies were kept in an empty vial for 1-5 minutes prior to

testing.

**Video acquisition and Tracking**

The video image was acquired at 30 Hz (1200 x 1210) using Fly capture software. Image

segmentation was performed by custom software in python using OpenCV. Two main

features were extracted from the videos: fly position (X and Y coordinates) and pixel

change around the fly. Positions were calculated from the centroid of an ellipse fitted to

the fly by background subtraction. The pixel change was quantified by calculating the

number of active pixels in a 100x100 pixel region surrounding the centroid. A pixel was

considered to be active if it recorded a change higher than 10 intensity levels.

The fly distance and speed were calculated for each frame using fly position x and y

coordinates with the formula:

$$distance = \sqrt{(x - x0)^2 + (y - y0)^2}$$

$$speed = \frac{distance}{time}$$

**Behavioral Classifiers**

We used velocity and pixel change features to automatically classify behavioral states.

We classified fly behavior into 4 categories: Walking, Grooming, Stopping and Leg

Displacement. A fly is considered to be walking if its average speed is higher than

1.9mm/sec. Grooming is detected when the velocity of the fly is lower than 1.9 mm/s and

average pixel change of a section of 8 frames is above 180 pixel/s. We classified as leg

displacement the movement in which velocity is lower than 1.9 mm/s and average pixel

change of a section of 8 frames is between 30 and 180 pixel/s. Finally, a fly is considered

to be stopping when velocity is lower than 1.9 and pixel change is lower than 30 pixel/s.

Thresholds were determined and confirmed by manual annotations of fly behavior (% of

efficacy). Later walking behavior was either classified into coordinated or uncoordinated

walking by visual inspection (Fig. 6 and 7).

**fTIR imaging and FlyWalker software**

For the analysis of walking behavior upon stimulation of sensory structures a new arena

comprising 12 LEDs (Deep red 655nm, Luxeon Rebel LED) was incorporated in the

previously described fTIR setup (Mendes et al., 2013), rendering a stimulation power of

0.014 mW/mm$^2$. Flies were placed in the FlyWalker arena and a 100 ms pulse of red light

was given. Each fly only received 1 pulse. The walking kinematics of flies after

stimulation were quantified using FlyWalker software package. One more parameter

called Non-Compliance Index was added to the previously published Kinematics

parameters. Non-Compliance Index is defined as the fraction of frames in a video in

which the fly displays non-compliant leg combinations. Non-canonical stance

combinations correspond to patterns that do not fit either of the idealized tripod, tetrapod

and wave gaits (Mendes et al., 2013; Mendes et al., 2014). In the experiments where we

used headless flies, we glued the head back onto the thorax using UV curable glue

(Loctite 3104, Henkel Adhesives) in order to avoid alterations in the body weight that

could potentially change walking patterns of the fly and to facilitate tracking.

**Tethered fly imaging**

For imaging of leg angles in Figure 4 we used 3-7 days-old females expressing

CsChrimson in mechanosensory bristles, as above. Adult flies were kept in food infused

with all trans-Retinal since the day of eclosion (Final Concentration 0.4 mM;

R2500,Sigma-Aldrich). Flies were anaesthetised on ice, and either decapitated with

scissors (Fine Scientific Tools) or left with head intact. They were positioned on a cold

plate with the thorax upwards, and a wire was glued to the thorax with UV curable glue

(Norland Optical Adhesives #81). Flies were fixed in the tethered set-up using the wire

above a porexpam ball (4.5 mm diameter, painted with black paint (Posca Uni)), held in a

custom made holder (the glass funnel of a Pasteur pipette). Flies could freely pick up and

walk on the ball, and if dropped could regain it again. Each fly was allowed to recover for

5 minutes on the set-up. After 5 minutes we waited for a spontaneous pause in behavior

and delivered a 100ms light stimulus via a single deep red LED (wavelength 627nm),

positioned 3 cm away from the fly. Light power at the fly was 13.8 nm/cm$^2$, calibrated to

produce a motor response in CsChrimson flies whilst not affecting flies in food lacking

all trans-Retinal (data not shown). Animals were illuminated with an InfraRed LED

(Thorlabs) and imaged with a PointGrey Camera (Flea3) with a Computar Macro-zoom

lens and a long-pass filter, at a frame rate of 120 fps. Video capture and LED stimulation

were controlled using a custom built Bonsai workflow (Lopes et al., 2015).

Walking bouts were extracted from head-intact non-stimulated flies (either before

stimulation or in sessions where no stimulation was delivered), and were manually

classified as satisfying straight walking with at least 6 step cycles. In total 8 walking

bouts were used from 3 flies. In the Head-intact and headless stimulation conditions for

the 'early response' and 'late response' timepoint, we visually inspected videos and

excluded timepoints where the flies were either still or performing grooming, as being

non-informative for our question of limb coordination while walking. Total N was 11

(head intact early response), 6 (head intact late response) and 10 (decapitated early

response). Videos were analyzed using DeepLabCut (Mathis et al., 2018). 18 points were

tracked in total; 3 body points (head, thorax and tail) and 5 points per leg (Body-Coxa;

Coxa-Femur; Femur-Tibia; Tibia-Tarsus; and Tarsus-end); however only T1 limb

tracking was used for the final analysis. Limb angles for T1 legs were calculated from

tracked positions using python.

**Leg and VNC dissection and Mounting**

To observe the morphology of leg sensory neurons, legs were dissected from the body in

PBS. The legs were then fixed in PFA 4% overnight at 4°C. After fixation legs were

washed 3x in 0.3% Triton-X in PBS for 20 minutes at R.T and mounted in 70% glycerol.

Brains and VNCs were fixed in PFA 4% for 20 minutes and dissected in PBS. Tissues

were labeled with a primary antibody raised against brp (mouse nc82, 1:10; DSHB)

followed by a secondary anti-mouse 647 (1:500; Jackson immunologicals). Tissues were

mounted in in 70% glycerol and scanned in a confocal microscope Zeiss (Oberkochen,

Germany) LSM710 confocal microscope with a 40× objective.

### *In vivo* calcium imaging during leg stimulation

Flies containing the following genotype: lexO-GCamp6f;mhc-lexA,lexO-Cherry and

dacRE-flp;Uas-FRT-stop-FRT-CsChrimson-mVenus(VK5), R65D12-Gal4(attP2) were

anesthetized with ice for 3-5 min. Three ipsilateral legs were glued to a glass slide with

UV-Curable glue. The body of the fly was immobilized using beeswax. GCaMP imaging

was performed at 25°C on a Nikon/Andor Revolution XD spinning-disk confocal

microscope with an EMCCD camera (iXon 897) using iQ 3.5 software and using a 20×

CFI Super Fluor 20X 0.75 NA dry objective. Images were taken at the scan speed of

100ms per frame. A set of 6 deep red LEDs was paced around the fly to provide the

100ms red light pulse.

### Data processing and Analysis

We extracted motor parameters using FlyWalker Software. Since many of the measured

gait parameters vary with speed, we analyzed the data for these parameters by

determining the best-fit regression model of each individual parameter with speed for the

control experiment and then determining the residual values for each experimental group

in relation to this regression model. Data was then expressed as the difference to the

residual normalized line. Normalized data was previously tested for normality and

homoscedasticity with Shapiro-Wilk and Levene tests. Statistical differences between

experimental groups were determined using Kruskal-Wallis analysis of variance followed

by Dunn's post hoc test (for non-normal distributions) or one-way-Anova followed by

Tukey's post hoc test (for normal distributions). Data analysis was performed using

custom Python scripts. Boxplots represent the median as the middle line, with the lower

and upper edges of the boxes representing the 25% and 75% quartiles, respectively.

Whiskers represent the range of the full data set, excluding outliers. Outliers are defined

as any value that is 1.5 times the interquartile range below or above the 25% and 75%

quartiles, respectively. For the boxplots in fig 2 and 3 we decided to use raw data not

normalized to the control since Non-Compliance index and Stance straightness

parameters do not show a correlation with velocity (data not shown).

To have a more succinct representation of the data we used Principal Component

Analysis (PCA), an unsupervised dimensionality reduction method. PCA finds a linear

projection of the data from a high-dimensional space onto a lower-dimensional subspace,

while maximizing variance of the projected data, and thus retains meaningful

information, resulting in a description of the data as a function of a smaller set of

uncorrelated variables (or principal components). The data were first pre-processed

by centering and scaling, after which the PCA algorithm computed the covariance matrix

in order to determine the correlation between variables and calculated the eigenvectors

and eigenvalues of the covariance matrix in order to identify the principal components.

We chose the first three principal components to visualize the data. The first two

components were chosen to generate a two-dimensional plot. Each dot in the plots

corresponds to a video, representing these new abstract variables. Color-coded dots were

used to distinguish specific groups. As such, clusters of dots reflect similar walking

patterns, shared by the corresponding flies (Cabrita et al., 2022).

For the analysis of the muscle fluorescence, we used Fiji software. To delineate the ROI

Area of the depressor and levator muscle we used the threshold tool and measured the

mean fluorescence value for each frame using ROI Manager, multimeasure tool.

Fluorescence values for each muscle were normalized by subtracting the minimum value

of each video and dividing by the maximum value, giving contraction values between 0

and 100%. Muscle contraction peaks were detected with a peak finding algorithm in

Python. Peak width was calculated by subtracting fluorescence 20 percentage points from

each peak value, and measuring the width of the signal that exceeded this value around

each peak (Fig 4F).

**Figures**

Figure 1

A          MsB1- Mechanosensory Bristles

B          ChO - Chordotonal Organ

C

Red LEDS

D  Control- Empty Gal 4    D′  MsB1- Mechanosensrory Bristles    D″  ChO- Chordotonal Organ

Head-Intact

10mm

E  Empty Gal 4    E′  MsB1- Mechanosensory Bristles    E″  ChO- Chordotonal Organ

Headless

10mm

**Figure 1. Optogenetic stimulation of leg mechanosensory bristles triggers movement in stationary flies**

(A) Leg and VNC expression pattern of leg MsB. mVenus expression under combinatorial control of R65D12 and dac$^{RE}$-flp. Genotype: R65D12-GAL44, dac$^{RE}$-flp, UAS-FRT-stop-FRT-Chrimson-mVenus. Cell bodies are located in the legs (left panel) and send axonal projections to the VNC (right panel). Bar, 100 μm. (B) Leg and VNC expression pattern in femoral ChO neurons. mVenus expression under combinatorial control of R79E02 and ato-flp. Genotype: R79E02-Gal4, ato-flp, UAS-FRT-stop-FRT-Chrimson-mVenus. Cell bodies are located in the legs (left panel) and send axonal projections to the VNC (right panel). Bar, 100 μm. (C) Schematic of the behavioral arena. Arena is surrounded by 6 deep red LEDs. See Methods for details. (D, E) Instantaneous speed and trajectories over time after optogenetic stimulation. Upper panels show instantaneous speed over time with red bar representing 100 ms red light stimulus. Dark line and light blue shadows indicate the average and SEM for all replicates, respectively. Lower panels show representative trajectories for 4 seconds after optogenetic stimulation. Trajectory plots are color coded according to the speed of the fly with black arrows and red dots indicating the initial and final positions, respectively. (D) Head intact and (E) headless animals. Red arrow indicates a brief speed increase due to reflexive leg twitch. At least 18 animals were used for each condition.

# Figure 2

**Figure 2. Kinematic analysis of walking patterns in MSB1 flies after**
**mechanosensory bristles stimulation**

(A-B) Representative gait features after MsB stimulation. Upper panel shows the flies'
step patterns across time. For each leg, swing phases are represented in black and stance
phases in white. Right hind (RH); right mid (RM); right front (RF); left hind (LH); left
middle (LM); left front (LF). Middle panel shows the instantaneous speed across time for
the same video. Thick and thin lines correspond to integration times of 25 or 12.5 ms,
respectively. Lower panel represents the gaits adopted by the fly during the period after
stimulation. For each frame the corresponding gait was color-coded as follows: Yellow
(tripod), blue (tetrapod), green (non-canonical), red (non-compliant configuration). See
Methods. (A) Empty GAL4 control (Genotype: empty-GAL44, dac$^{RE}$-flp, UAS-FRT-
stop-FRT-Chrimson-mVenus). (B) Post-stimulus response of MSB1 flies (Genotype:
R65D12-GAL4, dac$^{RE}$-flp, UAS-FRT-stop-FRT-Chrimson-mVenus). 100ms optogenetic
stimulation is represented by a red shadow. Sequence is divided in early response
(Uncoordinated Phase) and late response (Coordinated Phase) (B and B', respectively).
(C-E) Representative *stance traces*. Traces are generated by the position of the stance
phase footprints relative to the body center (set at 0.0,0.0). For each leg, stance onset
corresponds to the *Anterior Extreme Position* (*AEP*) (colored circle) while stance offset is
termed *Posterior Extreme Position* (*PEP*). Experimental conditions are color matched
with panels (A-B). (C) Control Empty GAL4 control. (D) Post-stimulus of MSB1 flies,
early response corresponding to the uncoordinated phase. (E) Post-stimulus of MSB1
flies, late response corresponding to the coordinated phase. (F-H) Kinematic
quantification between the empty GAL4 control and the early and late response post
stimulated (n=20 for each condition). In (F-G) boxplots represent the median as the
middle line, with the lower and upper edges of the boxes representing the 25% and 75%
quartiles, respectively, whiskers represent the range of the full data set, excluding outliers
represented by diamonds. Filled dots represent individual flies. Values are normalized for
body size. Statistical analysis with Kruskal–Wallis-ANOVA followed by Dunn's *post*
*hoc* analysis, **p<0.01, ***p<0.001. (F) *Non-Compliance Index* (G) *Stance straightness*
*index*. (H) Heat map of kinematic parameters comparing empty GAL4 control with early
response/ Uncoordinated Phase (first Column) and late response/Coordinated Phase

(second Column) of experimental group, with each line representing a motor parameter.

Data were residual normalized and outliers were excluded. Statistical analysis with one-

way ANOVA followed by Tukey's post hoc test (for normal distributions) or Kruskal–

Wallis-ANOVA followed by Dunn's post hoc test (for non-normal distribution). p values

are represented by a color code with red and blue shades indicate an increase or decrease

or relative to control, respectively. *P<0.05; **P<0.01; ***P<0.001. White indicates no

statistically significant variation.

# Figure 3

**A**

Tripod Gait · Tetrapod Gait · Non-Canonical · Non-Compliant · Stance Phase · Swing Phase

**C**

| | Head-Intact Flies | | Headless Resp |
| --- | --- | --- | --- |
| | Early Resp | Late Resp | |
| Average Speed | | | |
| Step Frequency | | | |
| Swing Speed(AVR) | | | |
| Swing Speed(F) | | | |
| Swing Speed(M) | | | |
| Swing Speed(H) | | | |
| Step Length(AVR | | | |
| Step Length(F) | | | |
| Step Length(M) | | | |
| Step Length(H) | | | |
| Swing Duration(AVR) | | | |
| Swing Duration(F) | | | |
| Swing Duration(M) | | | |
| Swing Duration(H) | | | |
| Stance Duration(AVR) | | | |
| Stance Duration(F) | | | |
| Stance Duration(M) | | | |
| Stance Duration(H) | | | |
| Stance Linearity | | | |
| AEP Footprint Clustering | | | |
| PEP Footprint Clustering | | | |
| Stance Straightness(AVR) | | | |
| Tripod Index | | | |
| Tetrapod Index | | | |
| Non-canonical configuration | | | |
| Wave Index | | | |
| Body displacement ratio | | | |
| Non-compliance Index | | | |

*p*
*** / ** / * / n.s. / * / ** / ***

**Figure 3. Mechanosensory Bristles evoked movement is independent of the central**

**Brain.**

(A) Representative gait features after MsB stimulation of headless flies. Genotype:

R65D12-GAL4, dac$^{RE}$-flp, UAS-FRT-stop-FRT-Chrimson-mVenus. Upper panel shows

the flies' step patterns across time. For each leg swing phases are represented in black

and stance phases in white. Right hind (RH); right mid (RM); right front (RF); left hind

(LH); left middle (LM); left front (LF). Middle panel shows the instantaneous speed

across time for the same video. Thick and thin lines correspond to integration times of 25

or 12.5 ms, respectively. Lower panel represents the gaits adopted by the fly during the

period after stimulation. For each frame the corresponding gait was color-coded as

follows: Yellow (tripod), blue (tetrapod), green (non-canonical), red (non-compliant

configurations). 100ms optogenetic stimulation is represented by a red shadow.

(B) Representative *stance traces* following optogenetic stimulation. Traces are generated

by the position of the stance phase footprints relative to the body center (set at 0.0,0.0).

For each leg, stance onset corresponds to the *Anterior Extreme Position* (*AEP*) (colored

circle) while stance offset is termed *Posterior Extreme Position* (*PEP*).

(C) Heat map of kinematic parameters comparing empty GAL4 control of head-intact

animals with the early/uncoordinated and late/coordinated responses (first and second

columns, respectively) of post-stimulated head-intact animals, and post-stimulated

headless flies (third column). Statistical analysis with one-way ANOVA followed by

Tukey's post hoc test (for normal distributions) or Kruskal–Wallis-ANOVA followed by

Dunn's post hoc test (for non-normal distribution). p values are represented by a color

code with red and blue shades indicate a decrease or increase relative to control,

respectively. *P<0.05; **P<0.01; ***P<0.001. White indicates no variation. (D) Stance

straightness index in head-intact empty GAL4 controls, early and late response of post-

stimulated head-intact animals, and post-stimulated headless flies. (E) Post stimulation

activity between head-intact (early response) and headless flies. (F) Non-Compliance

Index in head-intact empty GAL4 controls, early and late response of post-stimulated

head-intact animals, and post-stimulated headless flies. (G-G'') PCA of all kinematic

parameters in head-intact empty GAL4 controls, early and late response of post-

stimulated head-intact animals, and post-stimulated headless flies. (G) Tridimensional

representation of three-component PCA analysis. Each individual small dot represents

one video while larger dots represent the average point (n=20 for each condition).

Contribution of each component is indicated in each axis. (G') 2D representation of PC1-

PC2 with ellipses delimiting 50% of variance of the data. (G'') Comparison of each PC1

coordinates. In (D-F, G'') boxplots represent the median as the middle line, with the

lower and upper edges of the boxes representing the 25% and 75% quartiles, respectively,

whiskers represent the range of the full data set, excluding outliers represented by

diamonds. Filled dots represent individual flies. Values are normalized for body size.

Statistical analysis with Kruskal–Wallis-ANOVA followed by Dunn's *post hoc* analysis,

*P < 0.05; **P < 0.01; ***P < 0.001. (D) PC1 coordinates. (E) PC1 coordinates. See

Methods for details.

# Figure 4

**Figure 4. Mechanosensory Bristles evoked movement shows partially disrupted**
**inter-joint coordination.**
(A) Tethered fly set-up showing tracked joint positions (blue/green circles). *Inset*
schematic of Fly T1 leg showing 5 segments and 3 measured joint angles (Brown - Coxa-
Femur angle, Bright Orange – Femur-Tibia angle, Light Orange – Tibia-Tarsus angle). B)
Modulation of joint angles throughout time for our 4 experimental conditions: Walking
flies (blue horizontal bar), Head-Intact Early Response (red horizontal bar), Head-Intact
Late Response (green horizontal bar) and Headless Early Response (purple horizontal
bar). Red transparent vertical bars in the mid and lower panel indicate the time of
optogenetic stimulation. C) Fast Fourier Transform of leg angles for the Coxa-Femur
Joint. D) Angle-angle plots of Coxa-Femur vs Femur-Tibia angles over time. Different
colors represent individual trials. Black line indicates the mean trace of the 'walking'
condition over the step cycle. E) Histogram of distances of points to the mean walking
trace. (F) Violin plots for distances to mean walking trace. P values calculated using
Dunns post-hoc test. ***$P < 0.001$ G) Frequency of occupancy for quadrants of the Coxa-
Femur angle vs Femur-Tibia angle space.
932

# Figure 5

**Figure 5. Muscle activity alternation is maintained during sensory-triggered movement**

(A) Leg schematic showing tibia levator (blue) and depressor (orange) muscles in the femur. Adapted from (Venkatasubramanian et al., 2019). Dashed rectangle represents scanned region of the genotype: lexO-GCaMP6f; mhc-lexA:LHV2, dacRE-flp;Uas-FRT-stop-FRT-CsChrimson-mVenus, R65D12-Gal4 (B) Representative images of a femur showing levator (top panel) or depressor muscle (lower panel) contraction, resulting in GCaMP fluorescence inside the respective muscles. Yellow lines depict the ROIs used for fluorescence calculations. (C-D) Representative plots showing the fluctuation of fluorescence for levator (blue) and depressor (orange) muscle across time. Red bar represents the 100 ms red light stimulation. Values are normalized to maximum signal measured in each ROI throughout the experiment, corresponding to 100%. (C) Head-intact animals. Left panel: Non-stimulated animals in the presence of All-Trans Retinal (ATR), N=11; Middle Panel: Stimulated animals in the absence of ATR, N=13; Right Panel: Stimulated animals in the presence of ATR. N=13 (D) Headless animals. Left panel: Stimulated animals in the absence of ATR, N=12; Right Panel: Stimulated animals in the presence of ATR, N=13. (E) Spearman correlation between levator and depressor muscle activity for each experimental condition. Positive values indicate that muscles are in-phase, while Negative values correspond to out of phase (antagonistic) activity. Boxplots represent the median as the middle line, with the lower and upper edges of the boxes representing the 25% and 75% quartiles, respectively, whiskers represent the range of the full data set, excluding outliers represented by diamonds. Filled dots represent individual flies. Statistical analysis with Kruskal–Wallis-ANOVA followed by Dunn's *post hoc* analysis, no significant difference between the groups was detected (F) Schematic showing muscle contraction curve analysis shown in (G) and (H). Red dot indicates the maximum fluorescence value for the peak. Red line indicates the peak width at 20% below peak level. (G) Violin plot showing the maximum value of fluorescence for each contraction peak for each experimental condition. Each dot corresponds to a single muscle contraction peak. Horizontal line represents the median. (H) Violin plot showing the peak width for each contraction peak and each experimental condition. Each dot corresponds to the width of a single muscle contraction peak. Horizontal line represents

965    the median. P values in (G, H) were calculated using Dunns post-hoc test. *$P < 0.05$; **P

966    $< 0.01$; ***$P < 0.001$

# Figure 6

**Figure 6. Cyclic movement is specific to leg MS bristle stimulation**

(A) Schematic showing the combinatorial genetic tools used to restrict the expression of

the MsB GAL4 driver R65D12 to the legs (DAC$^{RE}$-flp), wings (Vg$^{BE}$-flp) and head (ey-

flp) regions. Each combination was crossed with UAS>>CsChrimson. (B-D, F-G) Raster

plot showing the type of movement performed by the flies before and after the

stimulation of leg MsB. X axis represents the time in seconds where 0 is the moment of

the red light pulse stimulation (red Bar). Each line corresponds to one individual.

Movement categories: Immobility (beige); Uncoordinated walking (orange); Coordinated

walking (yellow); Leg movement (brown); and Grooming (green). Upper Panels shows

Empty GAL4 behavior while lower panel shows MSB1 GAL4 (R65D12) behavior (B-D)

Head-intact animals. (B) Leg domain (with DAC$^{RE}$-flp). (C) Wing domain (with Vg$^{BE}$-

flp). (D) Head domain (with ey-flp). (E, H) Percentage of time spent doing one of the

behaviors described for (B-D,F-G). Boxplots represent the median as the middle line,

with the lower and upper edges of the boxes representing the 25% and 75% quartiles,

respectively, whiskers represent the range of the full data set, excluding outliers

represented by diamonds. Statistical analysis with Kruskal–Wallis-ANOVA followed by

Dunn's *post hoc* analysis, $*p<0.05$ $**p<0.01$, $***p<0.001$. (E) Quantification for head-

intact animals as shown in (B-D). (F-G) Headless flies. (F) Leg domain (with DAC$^{RE}$-

flp). (G) Wing domain (with Vg$^{BE}$-flp). (H) Quantification for head-intact animals as

shown in (F-G).

Figure 7

A

Coxa
Trochanter
Femur
Tibia
Tarsus

B

Cell Number

Coxa
Trochanter
Femur
Tibia
tarsus

MsB1  MsB3  MsB5  MsB4  MsB6  MsB2  MsB7

C

# cells

EmptyGAL4    0
MsB1    169±15
MsB2    57±10
MsB3    48±7
MsB4    43±12
MsB5    43±8
MsB6    26±4
MsB7    4±0.4

0    20    40    60    80    100
Duration of Movement (Sec)

D

Immobility    Uncoordinated Walking    Leg movement    Grooming

EmptyGAL4

MsB1

MsB3

MsB5

MsB4

MsB6

MsB2

MsB7

Time (s)

E

% Time

***    ***    ***  ***

EmptyGAL4    MsB1    MsB3    MsB5    MsB4    MsB6    MsB2    MsB7

**Figure 7. Movement pattern triggered by distinct leg MsB populations in headless animals**

(A) Schematic depicting the leg segments of the of *Drosophila melanogaster*. (B) Cell labelling distribution amongst leg segments for each line. Error bar represent standard deviation. MSB1: R65D12[leg], MSB2:R77E09[leg], MSB3:R39A11[leg], MSB4: R54A11[leg], MSB5: R64D07[leg], MSB6: R85A07[leg], MSB7: R45D07[leg]. (C) Duration of movement after one pulse of red light stimulation (100ms) for each MSB line. Lines are organized according to the number of labeled cells in an ascending order. (D) Raster plots showing the type of movement performed by the flies in each group, 1 second before and up to 10 seconds after the stimulation. X axis represent the time in seconds being 0 the time of optogenetic stimulation (red Bar). Each line corresponds to one individual. The categorization of movements includes Immobility (beige), Uncoordinated Walking (orange), Leg Displacement (brown); and Grooming (green). Lines are ordered in descending order according to the length of uncoordinated walking response (E) Boxplots showing for each MSB line the percentage of time spent doing a specific behavior. Behaviors are color-coded as in panel D. Boxplots represent the median as the middle line, with the lower and upper edges of the boxes representing the 25% and 75% quartiles, respectively, whiskers represent the range of the full data set, excluding outliers represented by diamonds. Statistical analysis with Kruskal–Wallis-ANOVA followed by Dunn's *post hoc* analysis, *p<0.05, **p<0.01, ***p<0.001.

**Lead Contact and Material Availability**

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, César S. Mendes (cesar.mendes@nms.unl.pt). The fly lines generated in this study are available from the Lead Contact without restrictions.

**Acknowledgements**

We thank Mendes lab, Rita Teodoro and her lab for comments and suggestions during the execution of this project, Allan Mancoo for feedback on data analysis, Inês Fernandes for experimental assistance, and Luísa Vasconcelos, Turgay Akay and Daniela Pereira for comments on the manuscript. We thank Mirian Zecca and Richard Mann for support generating the transgenic lines used in this study. We also thank CONGENTO: consortium for genetically tractable organisms, the Fly Platform and the Scientific Hardware Platform at the Champalimaud Centre for the Unknown, and the Bloomington Drosophila Stock Center for fly stocks. This work was supported by Fundação para a Ciência e a Tecnologia (FCT) (PTDC/BIA-COM/0151/2020), iNOVA4Health (UIDB/04462/2020 and UIDP/04462/2020), and LS4FUTURE (LA/P/0087/2020) to CSM. AM was supported by a doctoral fellowship from FCT (PD/BD/128445/2017).

**Author contributions**

AMM, AFH and CM contributed to conception and design of the study. AMM performed the experiments and analyzed the raw data; AFH performed data acquisition and analysis of Figure 4, and contributed to the analysis of Figure 5 data. GB contributed for analysis and data acquisition of Figure 7. AMM, AFH, MM and CSM wrote the manuscript. All authors contributed to manuscript revision, read, and approved the submitted version.

**Declaration of interests**

1043    The authors declare no competing interests.

1044

1045    **References**

1046

Agrawal, S., Dickinson, E.S., Sustar, A., Gurung, P., Shepherd, D., Truman, J.W., and
Tuthill, J.C. (2020). Central processing of leg proprioception in Drosophila. Elife *9*.

Akay, T., Bässler, U., Gerharz, P., and Büschges, A. (2001). The role of sensory signals
from the insect coxa-trochanteral joint in controlling motor activity of the femur-
tibia joint. Journal of neurophysiology *85*, 594-604.

Akay, T., and Büschges, A. (2006). Load signals assist the generation of movement-
dependent reflex reversal in the femur-tibia joint of stick insects. Journal of
neurophysiology *96*, 3532-3537.

Akay, T., Ludwar, B., Goritz, M.L., Schmitz, J., and Buschges, A. (2007). Segment
specificity of load signal processing depends on walking direction in the stick insect
leg muscle control system. J Neurosci *27*, 3285-3294.

Akay, T., Tourtellotte, W.G., Arber, S., and Jessell, T.M. (2014). Degradation of mouse
locomotor pattern in the absence of proprioceptive sensory feedback. Proc Natl
Acad Sci U S A *111*, 16877-16882.

Alexander, R.M. (2003). Principles of Animal Locomotion, STU - Student edition edn
(Princeton University Press).

Angeli, C.A., Edgerton, V.R., Gerasimenko, Y.P., and Harkema, S.J. (2014). Altering spinal
cord excitability enables voluntary movements after chronic complete paralysis in
humans. Brain *137*, 1394-1409.

Ballermann, M., and Fouad, K. (2006). Spontaneous locomotor recovery in spinal cord
injured rats is accompanied by anatomical plasticity of reticulospinal fibers. Eur J
Neurosci *23*, 1988-1996.

Bareyre, F.M., Kerschensteiner, M., Raineteau, O., Mettenleiter, T.C., Weinmann, O., and
Schwab, M.E. (2004). The injured spinal cord spontaneously forms a new intraspinal
circuit in adult rats. Nat Neurosci *7*, 269-277.

Barrios, N., Farias, M., and Moita, M.A. (2021). Threat induces cardiac and metabolic
changes that negatively impact survival in flies. Current biology : CB *31*, 5462-
5472.e5464.

Bassler, U., and Buschges, A. (1998). Pattern generation for stick insect walking
movements--multisensory control of a locomotor program. Brain Res Brain Res Rev
*27*, 65-88.

Baudoux, S., Duch, C., and Morris, O.T. (1998). Coupling of efferent neuromodulatory
neurons to rhythmical leg motor activity in the locust. J Neurophysiol *79*, 361-370.

Bidaye, S.S., Bockemühl, T., and Büschges, A. (2018). Six-legged walking in insects: how
CPGs, peripheral feedback, and descending signals generate coordinated and
adaptive motor rhythms. Journal of neurophysiology *119*, 459-475.

Borgmann, A., Toth, T.I., Gruhn, M., Daun-Gruhn, S., and Büschges, A. (2011).
Dominance of local sensory signals over inter-segmental effects in a motor system:
experiments. Biol Cybern *105*, 399-411.

Brown, T.G. (1911 ). The intrinsic factors in the act of progression in the mammal.
Proceedings of the Royal Society of London Series B, containing papers of a
biological character *84*, 308-319.

Buchner, E. (1976). Elementary movement detectors in an insect visual system.
Biological Cybernetics *24*, 85-101.

Burrows, M. (1996). The Neurobiology of an Insect Brain (Oxford University Press).

Burrows, M., Laurent, G.J., and Field, L.H. (1988). Proprioceptive inputs to nonspiking
local interneurons contribute to local reflexes of a locust hindleg. J Neurosci *8*, 3085-
3093.

Buschges, A. (1995). Role of local nonspiking interneurons in the generation of
rhythmic motor activity in the stick insect. J Neurobiol *27*, 488-512.

Büschges, A., Schmitz, J., and Bässler, U. (1995). Rhythmic patterns in the thoracic
nerve cord of the stick insect induced by pilocarpine. Journal of Experimental
Biology *198*, 435-456.

Cabrita, A., Medeiros, A.M., Pereira, T., Rodrigues, A.S., Kranendonk, M., and Mendes,
C.S. (2022). Motor dysfunction in Drosophila melanogaster as a biomarker for
developmental neurotoxicity. iScience *25*, 104541.

Cande, J., Namiki, S., Qiu, J., Korff, W., Card, G.M., Shaevitz, J.W., Stern, D.L., and Berman,
G.J. (2018). Optogenetic dissection of descending behavioral control in Drosophila.
eLife *7*, e34275.

Capogrosso, M., Wenger, N., Raspopovic, S., Musienko, P., Beauparlant, J., Bassi Luciani,
L., Courtine, G., and Micera, S. (2013). A computational model for epidural electrical
stimulation of spinal sensorimotor circuits. J Neurosci *33*, 19326-19340.

Chen, C., Agrawal, S., Mark, B., Mamiya, A., Sustar, A., Phelps, J.S., Lee, W.-C.A., Dickson,
B.J., Card, G.M., and Tuthill, J.C. (2021). Functional architecture of neural circuits for
leg proprioception in Drosophila. Current Biology *31*, 5163-5175.e5167.

Cheng, L.E., Song, W., Looger, L.L., Jan, L.Y., and Jan, Y.N. (2010). The role of the TRP
channel NompC in Drosophila larval and adult locomotion. Neuron *67*, 373-380.

Chrachri, A., and Clarac, F. (1987). Induction of rhythmic activity in motoneurons of
crayfish thoracic ganglia by cholinergic agonists. Neurosci Lett *77*, 49-54.

Court, R., Namiki, S., Armstrong, J.D., Börner, J., Card, G., Costa, M., Dickinson, M., Duch,
C., Korff, W., Mann, R., *et al.* (2020). A Systematic Nomenclature for the Drosophila
Ventral Nerve Cord. Neuron *107*, 1071-1079.e1072. doi:
1010.1016/j.neuron.2020.1008.1005. Epub 2020 Sep 1014.

Courtine, G., Song, B., Roy, R.R., Zhong, H., Herrmann, J.E., Ao, Y., Qi, J., Edgerton, V.R.,
and Sofroniew, M.V. (2008). Recovery of supraspinal control of stepping via indirect
propriospinal relay connections after spinal cord injury. Nat Med *14*, 69-74.

Cruse, H., and Bartling, C. (1995). Movement of Joint Angles in the Legs of a Walking
Insect, Carausius morosus. J Insecr Physiol *41*, 761-771.

Daun, S., Mantziaris, C., Tóth, T., Büschges, A., and Rosjat, N. (2019). Unravelling intra-
and intersegmental neuronal connectivity between central pattern generating
networks in a multi-legged locomotor system. PLoS One *14*, e0220767.

David, I., Holmes, P., and Ayali, A. (2016). Endogenous rhythm and pattern-generating
circuit interactions in cockroach motor centres. Biol Open *5*, 1229-1240.

DeAngelis, B.D., Zavatone-Veth, J.A., Gonzalez-Suarez, A.D., and Clark, D.A. (2020).
Spatiotemporally precise optogenetic activation of sensory neurons in freely
walking Drosophila. eLife *9*, e54183.

Dickinson, M.H., Farley, C.T., Full, R.J., Koehl, M.A., Kram, R., and Lehman, S. (2000). How
animals move: an integrative view. Science *288*, 100-106.

Durr, V., Schmitz, J., and Cruse, H. (2004). Behaviour-based modelling of hexapod
locomotion: linking biology and technical application. Arthropod Struct Dev *33*, 237-
250.

Eccles, J.C., Fatt, P., and Landgren, S. (1956). Central pathway for direct inhibitory
action of impulses in largest afferent nerve fibres to muscle. J Neurophysiol *19*, 75-
98.

Feng, K., Sen, R., Minegishi, R., Dubbert, M., Bockemuhl, T., Buschges, A., and Dickson,
B.J. (2020). Distributed control of motor circuits for backward walking in
Drosophila. Nat Commun *11*, 6166.

Formento, E., Minassian, K., Wagner, F., Mignardot, J.B., Le Goff-Mignardot, C.G., Rowald,
A., Bloch, J., Micera, S., Capogrosso, M., and Courtine, G. (2018). Electrical spinal cord
stimulation must preserve proprioception to enable locomotion in humans with
spinal cord injury. Nat Neurosci *21*, 1728-1741.

Fuchs, E., Holmes, P., Kiemel, T., and Ayali, A. (2011). Intersegmental coordination of
cockroach locomotion: adaptive control of centrally coupled pattern generator
circuits. Front Neural Circuits *4*, 125.

Gao, Q., Yuan, B., and Chess, A. (2000). Convergent projections of Drosophila olfactory
neurons to specific glomeruli in the antennal lobe. Nat Neurosci *3*, 780-785.

Goldammer, J., Mantziaris, C., Büschges, A., and Schmidt, J. (2018). Calcium imaging of
CPG-evoked activity in efferent neurons of the stick insect. PLoS One *13*, e0202822.

Götz, K.G., and Wenking, H. (1973). Visual control of locomotion in the walking
fruitflyDrosophila. Journal of comparative physiology *85*, 235-266.

Grillner, S. (1975). Locomotion in vertebrates: central mechanisms and reflex
interaction. Physiol Rev *55*, 247-304.

Hampel, S., McKellar, C.E., Simpson, J.H., and Seeds, A.M. (2017). Simultaneous
activation of parallel sensory pathways promotes a grooming sequence in
Drosophila. Elife *6*.

Heredia, F., Volonté, Y., Pereirinha, J., Fernandez-Acosta, M., Casimiro, A.P., Belém, C.G.,
Viegas, F., Tanaka, K., Menezes, J., Arana, M.*, et al.* (2021). The steroid-hormone
ecdysone coordinates parallel pupariation neuromotor and morphogenetic
subprograms via epidermis-to-neuron Dilp8-Lgr3 signal induction. Nat Commun *12*,
3328.

Hoyt, D.F., and Taylor, R. (1981). Gait and the energetics of locomotion in horses.
Nature *292*, 239-240.

Jarman, A.P., Grau, Y., Jan, L.Y., and Jan, Y.N. (1993). atonal is a proneural gene that
directs chordotonal organ formation in the Drosophila peripheral nervous system.
Cell *73*, 1307-1321.

Jenett, A., Rubin, G.M., Ngo, T.T., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer, B.D.,
Cavallaro, A., Hall, D., Jeter, J.*, et al.* (2012). A GAL4-driver line resource for
Drosophila neurobiology. Cell Rep *2*, 991-1001.

Juvin, L., Le Gal, J.-P., Simmers, J., and Morin, D. (2012). Cervicolumbar Coordination in
Mammalian Quadrupedal Locomotion: Role of Spinal Thoracic Circuitry and Limb
Sensory Inputs. The Journal of Neuroscience *32*, 953.

Kain, P., and Dahanukar, A. (2015). Secondary taste neurons that convey sweet taste
and starvation in the Drosophila brain. Neuron *85*, 819-832.

Kamikouchi, A., Wiek, R., Effertz, T., Göpfert, M.C., and Fiala, A. (2010). Transcuticular
optical imaging of stimulus-evoked neural activities in the Drosophila peripheral
nervous system. Nat Protoc *5*, 1229-1235.

King, D.G., and Wyman, R.J. (1980). Anatomy of the giant fibre pathway in Drosophila. I.
Three thoracic components of the pathway. Journal of Neurocytology *9*, 753-770.

Kittmann, R., and Schmitz, J. (1992). FUNCTIONAL SPECIALIZATION OF THE
SCOLOPARIA OF THE FEMORAL CHORDOTONAL ORGAN IN STICK INSECTS. Journal
of Experimental Biology *173*, 91-108.

Li, J., Zhang, W., Guo, Z., Wu, S., Jan, L.Y., and Jan, Y.N. (2016). A Defensive Kicking
Behavior in Response to Mechanical Stimuli Mediated by Drosophila Wing Margin
Bristles. J Neurosci *36*, 11275-11282. doi: 11210.11523/JNEUROSCI.11416-
11216.12016.

Lopes, G., Bonacchi, N., Frazao, J., Neto, J.P., Atallah, B.V., Soares, S., Moreira, L., Matias,
S., Itskov, P.M., Correia, P.A.*, et al.* (2015). Bonsai: an event-based framework for
processing and controlling data streams. Front Neuroinform *9*, 7.

Mamiya, A., Gurung, P., and Tuthill, J.C. (2018). Neural Coding of Leg Proprioception in
Drosophila. Neuron *100*, 636-650.e636.

Mantziaris, C., Bockemühl, T., Holmes, P., Borgmann, A., Daun, S., and Büschges, A.
(2017). Intra- and intersegmental influences among central pattern generating

networks in the walking system of the stick insect. Journal of neurophysiology *118*,
2296-2310.

Mathis, A., Mamidanna, P., Cury, K.M., Abe, T., Murthy, V.N., Mathis, M.W., and Bethge, M.
(2018). DeepLabCut: markerless pose estimation of user-defined body parts with
deep learning. Nature neuroscience *21*, 1281-1289.

Meissner, G.W., Nern, A., Dorman, Z., DePasquale, G.M., Forster, K., Gibney, T.,
Hausenfluck, J.H., He, Y., Iyer, N., Jeter, J., *et al.* (2022). A searchable image resource
of <em>Drosophila</em> GAL4-driver expression patterns with single
neuron resolution. bioRxiv, 2020.2005.2029.080473.

Mendes, C.S., Bartos, I., Akay, T., Marka, S., and Mann, R.S. (2013). Quantification of gait
parameters in freely walking wild type and sensory deprived Drosophila
melanogaster. Elife *2*, e00231.

Mendes, C.S., Rajendren, S.V., Bartos, I., Marka, S., and Mann, R.S. (2014). Kinematic
responses to changes in walking orientation and gravitational load in Drosophila
melanogaster. PLoS One *9*, e109204.

Merritt, D.J., and Murphey, R.K. (1992). Projections of leg proprioceptors within the
CNS of the fly Phormia in relation to the generalized insect ganglion. J Comp Neurol
*322*, 16-34.

Murphey, R.K., Possidente, D.R., Vandervorst, P., and Ghysen, A. (1989). Compartments
and the topography of leg afferent projections in Drosophila. J Neurosci *9*, 3209-
3217.

Namiki, S., Dickinson, M.H., Wong, A.M., Korff, W., and Card, G.M. (2018). The functional
organization of descending sensory-motor pathways in Drosophila. eLife *7*, e34272.

Pearson, K.G. (1972). Central Programming and Reflex Control of Walking in the
Cockroach. Journal of Experimental Biology *56*, 173-193.

Pearson, K.G., and Fourtner, C.R. (1975). Nonspiking interneurons in walking system of
the cockroach. J Neurophysiol *38*, 33-52.

Pfeiffer, B.D., Ngo, T.T., Hibbard, K.L., Murphy, C., Jenett, A., Truman, J.W., and Rubin,
G.M. (2010). Refinement of tools for targeted gene expression in Drosophila.
Genetics *186*, 735-755.

Phelps, J.S., Hildebrand, D.G.C., Graham, B.J., Kuan, A.T., Thomas, L.A., Nguyen, T.M.,
Buhmann, J., Azevedo, A.W., Sustar, A., Agrawal, S.*, et al.* (2021). Reconstruction of
motor control circuits in adult Drosophila using automated transmission electron
microscopy. Cell *184*, 759-774.e718. doi: 710.1016/j.cell.2020.1012.1013. Epub
2021 Jan 1014.

Ramdya, P., Lichocki, P., Cruchet, S., Frisch, L., Tse, W., Floreano, D., and Benton, R.
(2015). Mechanosensory interactions drive collective behaviour in Drosophila.
Nature *519*, 233-236.

Rillich, J., Stevenson, P.A., and Pflueger, H.J. (2013). Flight and walking in locusts-
cholinergic co-activation, temporal coupling and its modulation by biogenic amines.
PLoS One *8*, e62899.

Ritzmann, R.E., and Buschges, A. (2007). Adaptive motor behavior in insects. Curr Opin
Neurobiol *17*, 629-636.

Rosenbaum, P., Wosnitza, A., Buschges, A., and Gruhn, M. (2010). Activity patterns and
timing of muscle activity in the forward walking and backward walking stick insect
Carausius morosus. Journal of neurophysiology *104*, 1681-1695.

Ryckebusch, S., and Laurent, G. (1993). Rhythmic patterns evoked in locust leg motor
neurons by the muscarinic agonist pilocarpine. J Neurophysiol *69*, 1583-1595.

Ryckebusch, S., and Laurent, G. (1994). Interactions between segmental leg central
pattern generators during fictive rhythms in the locust. J Neurophysiol *72*, 2771-
2785.

Seeds, A.M., Ravbar, P., Chung, P., Hampel, S., Midgley, F.M., Jr., Mensh, B.D., and
Simpson, J.H. (2014). A suppression hierarchy among competing motor programs
drives sequential grooming in Drosophila. Elife *3*, e02951.

Seelig, J.D., Chiappe, M.E., Lott, G.K., Dutta, A., Osborne, J.E., Reiser, M.B., and Jayaraman,
V. (2010). Two-photon calcium imaging from head-fixed Drosophila during
optomotor walking behavior. Nature methods *7*, 535-540.

Shanbhag, S.R., Singh, K., and Naresh Singh, R. (1992). Ultrastructure of the femoral
chordotonal organs and their novel synaptic organization in the legs of Drosophila
melanogaster Meigen (Diptera : Drosophilidae). International Journal of Insect
Morphology and Embryology *21*, 311-322.

Sherrington, C.S. (1893). Further Experimental Note on the Correlation of Action of
Antagonistic Muscles. Br Med J *1*, 1218.

Siegler, M.V., and Burrows, M. (1986). Receptive fields of motor neurons underlying
local tactile reflexes in the locust. J Neurosci *6*, 507-513.

Smith, S.A., and Shepherd, D. (1996). Central afferent projections of proprioceptive
sensory neurons in Drosophila revealed with the enhancer-trap technique. J Comp
Neurol *364*, 311-323.

Soler, C., Daczewska, M., Da Ponte, J.P., Dastugue, B., and Jagla, K. (2004). Coordinated
development of muscles and tendons of the Drosophila leg. Development *131*, 6041-
6051.

Strausfeld, N.J., Bassemir, U., Singh, R.N., and Bacon, J.P. (1984). Organizational
principles of outputs from Dipteran brains. Journal of Insect Physiology *30*, 73-93.

Strauss, R., and Heisenberg, M. (1990). Coordination of legs during straight walking
and turning in Drosophila melanogaster. J Comp Physiol A *167*, 403-412.

Szczecinski, N.S., Bockemuhl, T., Chockley, A.S., and Buschges, A. (2018). Static stability
predicts the continuum of interleg coordination patterns in Drosophila. J Exp Biol
*221*.

Takeoka, A., Vollenweider, I., Courtine, G., and Arber, S. (2014). Muscle spindle
feedback directs locomotor recovery and circuit reorganization after spinal cord
injury. Cell *159*, 1626-1639.

Tanouye, M.A., and Wyman, R.J. (1980). Motor outputs of giant nerve fiber in
Drosophila. Journal of Neurophysiology *44*, 405-421.

Trimarchi, J.R., and Schneiderman, A.M. (1995). Flight initiations in Drosophila
melanogaster are mediated by several distinct motor patterns. Journal of
Comparative Physiology A *176*, 355-364.

Tsubouchi, A., Yano, T., Yokoyama, T.K., Murtin, C., Otsuna, H., and Ito, K. (2017).
Topological and modality-specific representation of somatosensory information in
the fly brain. Science *358*, 615-623.

Tuthill, J.C., and Wilson, R.I. (2016a). Mechanosensation and Adaptive Motor Control in
Insects. Curr Biol *26*, R1022-R1038.

Tuthill, J.C., and Wilson, R.I. (2016b). Parallel Transformation of Tactile Signals in
Central Circuits of Drosophila. Cell *164*, 1046-1059.

Vandervorst, P., and Ghysen, A. (1980). Genetic control of sensory connections in
Drosophila. Nature *286*, 65-67.

Venkatasubramanian, L., Guo, Z., Xu, S., Tan, L., Xiao, Q., Nagarkar-Jaiswal, S., and Mann,
R.S. (2019). Stereotyped terminal axon branching of leg motor neurons mediated by
IgSF proteins DIP-α and Dpr10. eLife *8*, e42692.

Wagner, F.B., Mignardot, J.B., Le Goff-Mignardot, C.G., Demesmaeker, R., Komi, S.,
Capogrosso, M., Rowald, A., Seanez, I., Caban, M., Pirondini, E., *et al.* (2018). Targeted
neurotechnology restores walking in humans with spinal cord injury. Nature *563*,
65-71.

Williams, J.A., Paddock, S.W., Vorwerk, K., and Carroll, S.B. (1994). Organization of wing
formation and induction of a wing-patterning gene at the dorsal/ventral
compartment boundary. Nature *368*, 299-305.

Wosnitza, A., Bockemuhl, T., Dubbert, M., Scholz, H., and Buschges, A. (2013). Inter-leg
coordination in the control of walking speed in Drosophila. J Exp Biol *216*, 480-491.

Wu, M., Nern, A., Williamson, W.R., Morimoto, M.M., Reiser, M.B., Card, G.M., and Rubin,
G.M. (2016). Visual projection neurons in the Drosophila lobula link feature
detection to distinct behavioral programs. Elife *5*.

Yellman, C., Tao, H., He, B., and Hirsh, J. (1997). Conserved and sexually dimorphic
behavioral responses to biogenic amines in decapitated Drosophila. Proc Natl Acad
Sci U S A *94*, 4131-4136.

Zecca, M., and Struhl, G. (2007). Control of Drosophila wing growth by the vestigial
quadrant enhancer. Development (Cambridge, England) *134*, 3011-3020.

zur Lage, P.I., Powell, L.M., Prentice, D.R., McLaughlin, P., and Jarman, A.P. (2004). EGF
receptor signaling triggers recruitment of Drosophila sense organ precursors by
stimulating proneural gene autoregulation. Dev Cell *7*, 687-696.

1317