Relationship between salmon egg subsidy and the distribution of an avian predator

Taihei Yamada[1,*], Hirotaka Katahira[2], Kazuki Miura[3,a], Futoshi Nakamura[1]

[1]Graduate School of Agriculture, Hokkaido University, N9W9, Sapporo, Hokkaido 060-8589, Japan

[2]Department of Environmental Science, School of Life and Environmental Science, Azabu University, 1-17-71 Fuchinobe, Chuo-ku, Sagamihara, Kanagawa 252-5201, Japan

[3]Shiretoko Museum, Honmachi 49, Shari-gun, Shari, Hokkaido, 099-4113, Japan

[a]**Present address**: Research Institute of Energy, Environment, and Geology, Hokkaido Research Organization, N19W12, Sapporo, Hokkaido 060-0819, Japan

*Corresponding author: Taihei Yamada; E-mail, thymd8@gmail.com; TEL, +818039994855

Graduate School of Agriculture, Hokkaido University, N9W9, Sapporo, Hokkaido 060-8589, Japan

ABSTRACT

As a spatial subsidy, which is the phenomenon of transferring resources from a donor system to a recipient system, anadromous salmonids contribute to the supply of marine-derived nutrients to freshwater and terrestrial systems. Live salmon and salmon carcasses and eggs are utilized by various organisms and affect their abundance and distribution. However, the evaluation of the effect of salmon subsidies on the abundance and distribution of terrestrial animals is biased towards predators or scavengers that utilize spawning adults and carcasses, and few studies have focused on the effect of salmon eggs as a subsidy. To avoid underestimating the function of salmon subsidies, the response to the availability of salmon eggs in various systems should be investigated. Here, we investigated the abundance and feeding behaviour of the brown dipper *Cinclus pallasii*, as salmon egg a consumer, based on the hypothesis that the availability of salmon eggs affects the diet composition and stream distribution of this small predator. In addition, to test whether changes in the abundance of brown dippers are determined by salmon spawning, their abundance was compared upstream and downstream of the dam. Brown dippers used salmon eggs during the spawning season (53.7% of diet composition), and their abundance increased as the number of spawning redds increased. In contrast, this pattern was not observed upstream of the dam. These results suggested that the abundance and stream distribution of brown dippers vary according to the variation in the spatiotemporal availability of salmon eggs.

1. INTRODUCTION

Spatial subsidies are a phenomenon in which resources are transferred from a donor system to a recipient system (Polis et al., 1997). Spatial subsidies play a crucial role in biological communities because they affect the abundance and distribution patterns of organisms and the food web structure in the recipient systems by affecting the availability of basal resources (Hocking et al., 2013; Kawaguchi et al., 2003; Nakano & Murakami, 2001; Spiller et al., 2010; Terui et al., 2018).

Anadromous salmonids, well-known spatial subsidy representatives, transport marine-derived nutrients and energy to freshwater and terrestrial ecosystems through their migrations (Gende et al., 2002; Hocking & Reynolds, 2011; Koshino et al., 2013; Schindler et al., 2003). Salmon subsidies contribute to increasing aquatic invertebrate biomass and freshwater fish abundance in rivers (Denton et al., 2009; Wipfli et al., 1998, 1999). Spawning adults and carcasses are also used as food by terrestrial animals not only in the underwater ecosystem but also in surrounding riparian ecosystems and eventually affect the abundance and distribution of terrestrial scavengers and top predators (Boulanger et al., 2004; Christie & Reimchen, 2005; Field & Reynolds, 2013; Levi et al., 2012; Walters et al., 2021). Salmon subsidies thus provide insights into how multiple ecosystems are tangled together.

Past investigations on the effects of salmon subsidies on terrestrial organism abundance and distribution have thus far been biased towards predators or scavengers that utilize spawning adults and carcasses (e.g., Boulanger et al., 2004; Christie & Reimchen, 2005; Field & Reynolds, 2013; Levi et al., 2012; Walters et al., 2021). Considering that terrestrial organisms are provided multiple resources by salmon runs, such as eggs and fry, as well as spawning adults and carcasses (Munro, 1941; Willson & Halupka, 1995), salmon subsidies may have even more unexpected far-reaching effects. Therefore, a comprehensive understanding of the effect of

62   subsidies on the population and distribution of organisms is important. However, few studies

63   have focused on the effect of salmon eggs as a subsidy on the abundance and distribution of

64   terrestrial animals. It is necessary to clarify the response to the availability of salmon eggs in

65   various systems to avoid underestimating the function of salmon subsidies.

66   Dippers (Aves: Cinclidae) are riparian birds that mainly feed on aquatic invertebrates by

diving into the water (Eguchi, 1990; Taylor & O'Halloran, 1997, 2001) and which are known to

use salmon eggs in available rivers and seasons (Goodge, 1959; Obermeyer et al., 1999, 2006;

Reimchen, 2017; Whitehorne, 2010). For example, the American dipper *Cinclus mexicanus* can

achieve higher reproductive success (as measured by fecundity and juvenile growth) in reaches

where *Oncorhynchus* swim upstream than in reaches where it does not (Obermeyer et al., 2006;

Tonra et al., 2016). The population size of the white-throated dipper *C. cinclus* in Norway may

also benefit from eating salmon fry because it was correlated with the annual density of salmon

fry (Nilsson et al., 2018). Because dippers, which are not scavengers, are not affected by the

amount of carcasses – in addition to their well-studied relationship with salmon, as noted above –

they are a suitable model species for examining the effect of salmon egg subsidies on the

abundance and distribution of terrestrial animals. The brown dipper *C. pallasii* (Figure 1), which

is distributed in Asia (Hong et al., 2019), preys on salmon eggs and juvenile salmon (Murata,

1900). However, its actual status has never been evaluated quantitatively, and the relationship

with salmon subsidies has long been overlooked.

81   We therefore investigated the abundance and diet composition of the brown dipper in

82   the Shiretoko Peninsula of Hokkaido, northern Japan, where the spawning migrations of salmon

83   are well observed, based on the hypothesis that the availability of salmon subsidies drives the

84   diet composition and stream distribution of this small predator. In addition, by comparing the

85   abundance of brown dippers above and below dams where salmon cannot run upstream during

peak salmon spawning runs, we tested whether changes in the abundance of brown dippers are

determined by salmon spawning. More specifically, it was predicted that salmon spawning would

cause a shift in the diet of brown dippers to salmon eggs and an increase in the abundance of

brown dippers by altering the distribution of food resources, while no such pattern occurs

upstream of the dam.

2. MATERIALS AND METHODS

2-1. Study site

Four streams located in the Shiretoko Peninsula were selected for the present survey (Figure 2).

Natural spawning sustains pink salmon *Oncorhynchus gorbuscha* and chum salmon *O. keta*

populations in these streams, and the former is dominant (T. Yamada, unpublished data). The

release of juvenile chum salmon has been conducted only in the Mosekarubetsu stream, and in-

stream harvesting does not occur in all streams. The central part of the Shiretoko Peninsula has

been designated as a World Natural Heritage site since 2005, partially because of the close

relationships between the marine and terrestrial ecosystems sustained by the anadromous

migration of pink salmon and chum salmon (IUCN, 2005). The upper reaches of the studied

streams are included in the Shiretoko World Natural Heritage site. Rivers and streams in the

Shiretoko Peninsula are highly fragmented by more than 330 artificial dams (Takahashi et al.,

2005), which is no exception in all selected streams. Study sections were set up in each stream

from the mouth of the stream to the proximal dam. The Chienbetsu stream was surveyed up to

the second proximal dam because many salmon pass through the first proximal dam. The length

of the study section was 289.4, 348.2, 154.9, and 211.3 m in the Chienbetsu stream, Funbe

stream, Mosekarubetsu stream, and Shoji stream, respectively.

2-2. Field survey

Temporal changes in the abundance and diet of brown dippers were evaluated from mid-August to early November 2021, the spawning period of pink salmon. Field observations were conducted in one or two streams per day for a total effort of eight or nine days at every 9- to 11-day interval in each stream. In the observation protocol, one investigator (T. Yamada) walked along the study section from the lower to the upper reach, counting the number of brown dippers. To avoid recounting birds, the investigator checked where the flying individuals stopped and ignored individuals who flew ahead of the investigator, expecting territorial individuals to characteristically 'double-back' when pushed to the ends of their territory (Chiu et al., 2008).

The date of the count survey was the same as that of the diet survey, and the count survey was conducted before the diet survey was conducted. In the diet survey, when the investigator found an individual, they approached it at an observable distance and recorded the diet composition and age category (adult or juvenile) using binoculars (MONARCH 10 × 42; Nikon, Tokyo, Japan) (Obermeyer et al., 1999). The age category was classified by the presence or absence of juvenal plumage. The contents of the dipper's diet were classified into four categories: aquatic insects, terrestrial insects, algae, and salmon eggs. If no observable individuals were found, no observations were made. The diet survey was conducted only once per individual at each observation cycle; the mean ± sd observation time was 4.29 ± 3.16 minutes.

Salmon spawning redds were also visually counted on the same day as the above observation procedures to obtain an index of the availability of salmon eggs. Pink salmon exhibit "probing", a periodic and short-term migration behaviour between the sea and multiple drainages (Morita, 2021; Thedinga et al., 2000). If many individuals exhibit probing, salmon abundance

cannot be a direct indicator of the number of spawners; therefore, we used the number of

spawning redds as an indicator of the number of spawners. Spawning redds were visually judged

as the area of disturbed gravel or bright (denuded) areas among the periphyton-covered gravel

(Ortlepp & Mürle, 2003; Pedersen et al., 2009). We also measured the stream surface area of the

study section only once in each section during the study period.

The abundance of brown dippers was also surveyed upstream of dams in the selected

stream at the end of September 2021 during the peak spawning period of pink salmon, except in

the Chienbetsu stream, where the dam has a fishway allowing migration to the upper reaches.

Additional study sections were set up at approximately 400 m from the dam in each case. The

distance between the end of the below dam section and the start of the above dam section in each

stream was 1 to 2 m. An investigator walked from the dam to the upstream end of the study

section counting the number of dippers, as was the case in the lower reach survey. This survey

was conducted on the same day in the fifth cycle of the count survey mentioned above. We also

measured the stream surface area of the study section only once in each stream.

2-3. Statistical analysis

The dependence on salmon eggs in brown dipper diets was evaluated by fitting a generalized

linear mixed model (GLMM) to the individual diet data per. In the analysis, the dominance ratio

of salmon eggs in the diet composition was used as a response variable and it was assumed to

follow a binomial distribution. Age category and number of spawning redds were used as the

candidate explanatory variables, considering stream ID and observation cycle ID as a nested

random intercept. To avoid multicollinearity, variance inflation factors (VIFs) were calculated

before the analysis; all variables had values less than 2.5, the threshold indicative of troubling

collinearity for regressions (Johnston et al., 2018). Akaike's information criterion (AIC) was

used for model selection (Burnham & Anderson, 2002). If several plausible models had ΔAIC ≤

2, the optimal model was selected according to the principle of parsimony (Burnham &

Anderson, 2002).

A relationship between the availability of salmon eggs (represented as the number of

spawning redds) and the brown dipper abundance was also estimated by fitting GLMMs to the

dipper count data as a response variable assuming a Poisson distribution. The number of

spawning redds and time in the day when the survey was started were used as the candidate

explanatory variables, considering log-transformed stream surface area as an offset term and

stream ID as a random intercept. VIF of all variables were less than 2.5. Akaike's information

criterion (AIC) was used for model selection (Burnham & Anderson, 2002). If several plausible

models had ΔAIC ≤ 2, the optimal model was selected according to the principle of parsimony

(Burnham & Anderson, 2002).

Since resource availability may affect organism distribution (Dingle, 2014; e.g., Dingle

& Drake, 2007), it was also expected that the dipper abundance differed between the upper and

lower reaches of the dam. Brown dipper abundance in the fifth observation cycle was compared

between the lower reaches and the upper reaches of the dam by fitting the count data to a GLMM

considering log-transformed stream surface area as an offset term and the stream ID as a random

intercept. When the 95% confidence intervals (95% CI) of the estimated coefficient values did

not overlap between the study sections, we considered the differences significant.

All data analyses were conducted with R v. 4.2.0 (R Core Team, 2022) using lme4 v.

1.1.30 (Bates et al., 2015) for GLMMs.

3. RESULTS

A total of 108 brown dipper individuals, 631 redds, 1257 pink salmon individuals, and 118 chum

salmon individuals were observed during our survey. Feeding behaviour was monitored in 4

individuals (3 adults and 1 juvenile) in the pre-spawning period and in 24 individuals (15 adults

and 9 juveniles) in the spawning period from the three streams, except for the Mosekarubetsu

stream, where close observation could not be made.

The diet composition changed between the salmon pre-spawning and spawning periods

(Figure 3a). The percentage of salmon eggs in the diet was up to 53.7% (Figure 3a) during the

latter period. The brown dippers basically ingested only small food in the water and did not peck

salmon carcasses. As a result of model selection based on AIC, the model including the number

of spawning redds as the explanatory variable was selected (Table1), indicating that the number

of spawning redds had a positive effect on the salmon egg ratio in the diet (Figure 3b; Table 2).

The abundance survey showed that brown dipper abundance tended to be relatively high

during the salmon spawning period in each stream (Figure 4). In the model selection process, the

model including the number of spawning redds as the explanatory variable was selected (Table

1). The brown dipper abundance positively correlated with the number of spawning redds in the

selected model (Figure 5; Table 2). The comparison between abundances in the upper and lower

sections of the dam showed that the abundance in the lower section was clearly higher than that

in the upper section (Marginal R^2 = 0.792, Conditional R^2 = 0.821; Figure 6; Table 2).

4. DISCUSSION

203   This study demonstrates for the first time that salmon eggs are the dominant dietary item for the

204   brown dipper during the salmon spawning season and that the abundance and stream distribution

205   of terrestrial vertebrate species can be predicted by the number of spawning redds used to

represent the availability of salmon eggs. In addition, brown dipper abundance during the peak spawning season differed significantly between upstream and downstream of the dam, with the downstream abundance being higher. It is therefore indicated that the distribution of brown dippers varies according to variation in the spatiotemporal availability of salmon subsidies. One of the typical theories of the spatial distribution of animal populations is the ideal free distribution, which assumes that individuals are free to move among sites (Fretwell & Lucas, 1969), and our results may be explained by this theory. The reason is that brown dippers did not use the site upstream of the dam (without salmon subsidies) even when their density was high downstream of the dam.

While salmon eggs serve as an important food source for brown dippers, the salmon spawning behaviour of digging up the riverbed leads to a reduction in the abundance of aquatic invertebrates that are prey for brown dippers (Minakawa & Gara, 2003; Moore & Schindler, 2008). Since the energy value per salmon egg is higher than that per individual aquatic invertebrate (Obermeyer et al., 2006; Whitehorne, 2010), the positive effect of egg-eating may outweigh the negative effect of reduced eating of aquatic invertebrates. In fact, juvenile weight and mortality in the American dipper in salmon spawning reaches are known to be higher and lower than those in the non-spawning reaches (Obermeyer et al., 2006). Further verification is required to clarify whether these findings may be supported in the present system.

The abundance of aquatic invertebrates varies greatly with the season (Rundio & Lindley, 2008) and declines with flooding (Chiu et al., 2008; McMullen & Lytle, 2012). Accordingly, the decrease in aquatic invertebrate abundance leads to a decline in brown dipper abundance and survival (Chiu et al., 2008, 2013). Pink salmon, chum salmon, and masu salmon running up Japanese rivers spawn during the summer, fall and winter (Iida et al., 2021; Kovach et al., 2012; Kuzishchin et al., 2009; Quinn, 2018). Since summer and fall are typhoon seasons in

East Asia, salmon subsidies may compensate for the decline in aquatic invertebrates. In addition,

dippers sometimes prey on salmon fry (Obermeyer et al., 2006; Ormerod, 1985; Ormerod &

Tyler, 1986, 1991). Since most salmon fry mainly emerge during spring and summer (Kirillov et

al., 2018; Pavlov et al., 2008; Yamada et al., 2022), salmon fry may be used as a food resource

by dippers during this period. Therefore, spawning by anadromous salmonids may compensate

for declines in the abundance of aquatic invertebrates during various seasons.

Salmon spawning abundance is disturbed by several human activities, such as dam

construction and fisheries (Finney et al., 2000; Nakamura & Komiyama, 2010; Romakkaniemi et

al., 2003). This study shows for the first time that the distribution patterns of small terrestrial

predators are determined by the supply of salmon eggs, indicating that the disruption of natural

spawning may have unexpected effects on the abundance and distribution of terrestrial salmon

egg consumers. Future studies are needed to examine the effects of these anthropogenic

restrictions of the salmon egg subsidy on the abundance and distribution of terrestrial consumers.

COMPETING INTERESTS STATEMENT

None declared.

ACKNOWLEDGMENTS

We thank M. Kitazawa at Hokkaido University for his valuable comments. This study was

supported by JSPS KAKENHI (Grant Numbers 21H03647, 22J11475) and JST SPRING (Grant

Number JPMJSP2119).

REFERENCES

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/jss.v067.i01

Boulanger, J., Himmer, S., & Swan, C. (2004). Monitoring of grizzly bear population trends and demography using DNA mark–recapture methods in the Owikeno Lake area of British Columbia. *Canadian Journal of Zoology*, *82*(8), 1267–1277. https://doi.org/10.1139/z04-100

Burnham, K. P., & Anderson, D. R. (2002). *Model Selection and Multimodel Inference* (K. P. Burnham & D. R. Anderson (eds.)). Springer New York. https://doi.org/10.1007/b97636

Chiu, M. C., Kuo, M. H., Hong, S. Y., & Sun, Y. H. (2013). Impact of extreme flooding on the annual survival of a riparian predator, the Brown Dipper *Cinclus pallasii*. *Ibis*, *155*(2), 377–383. https://doi.org/10.1111/ibi.12035

Chiu, M. C., Kuo, M. H., Sun, Y. H., Hong, S. Y., & Kuo, H. C. (2008). Effects of flooding on avian top-predators and their invertebrate prey in a monsoonal Taiwan stream. *Freshwater Biology*, *53*(7), 1335–1344. https://doi.org/10.1111/j.1365-2427.2008.01968.x

Christie, K. S., & Reimchen, T. E. (2005). Post-reproductive Pacific salmon, *Oncorhynchus* spp., as a major nutrient source for large aggregations of gulls, Larus spp. *Canadian Field-Naturalist*, *119*(2), 202–207. https://doi.org/10.22621/cfn.v119i2.107

Denton, K. P., Rich, H. B., & Quinn, T. P. (2009). Diet, Movement, and Growth of Dolly Varden in Response to Sockeye Salmon Subsidies. *Transactions of the American Fisheries Society*, *138*(6), 1207–1219. https://doi.org/10.1577/t09-006.1

Dingle, H. (2014). *Migration: The Biology of Life on the Move*. Oxford University Press. https://doi.org/10.1093/acprof:oso/9780199640386.001.0001

Dingle, H., & Drake, V. A. (2007). What Is Migration? *BioScience*, *57*(2), 113–121.

https://doi.org/10.1641/B570206

Eguchi, K. (1990). The choice of foraging methods of the Brown Dipper,*Cinclus pallasii* (Aves:

Cinclidae). *Journal of Ethology*, *8*(2), 121–127. https://doi.org/10.1007/BF02350282

Field, R. D., & Reynolds, J. D. (2013). Ecological links between salmon, large carnivore

predation, and scavenging birds. *Journal of Avian Biology*, *44*(1), 009–016.

https://doi.org/10.1111/j.1600-048X.2012.05601.x

Finney, B. P., Gregory-Eaves, I., Sweetman, J., Douglas, M. S. V., & Smol, J. P. (2000). Impacts

of Climatic Change and Fishing on Pacific Salmon Abundance Over the Past 300 Years.

Science, *290*(5492), 795–799. https://doi.org/10.1126/science.290.5492.795

Fretwell, S. D., & Lucas, H. L. (1969). On territorial behavior and other factors influencing

habitat distribution in birds. *Acta Biotheoretica*, *19*(1), 16–36.

https://doi.org/10.1007/BF01601953

Gende, S. M., Edwards, R. T., Willson, M. F., & Wipfli, M. S. (2002). Pacific salmon in aquatic

and terrestrial ecosystems. *BioScience*, *52*(10), 917–928. https://doi.org/10.1641/0006-

3568(2002)052[0917:PSIAAT]2.0.CO;2

Goodge, W. R. (1959). Locomotion and Other Behavior of the Dipper. *The Condor*, *61*(1), 4–17.

https://doi.org/10.2307/1365341

Hocking, M. D., Dulvy, N. K., Reynolds, J. D., Ring, R. A., & Reimchen, T. E. (2013). Salmon

subsidize an escape from a size spectrum. *Proceedings of the Royal Society B: Biological*

Sciences, *280*(1753). https://doi.org/10.1098/rspb.2012.2433

Hocking, M. D., & Reynolds, J. D. (2011). Impacts of salmon on riparian plant diversity.

Science, *331*(6024), 1609–1612. https://doi.org/10.1126/science.1201079

Hong, S. Y., Wang, T. W., Sun, Y. H., Chiu, M. C., Kuo, M. H., & Chen, C. C. (2019). Stream

type influences food abundance and reproductive performance of a stream specialist: the

Brown Dipper (*Cinclus pallasii*). *Journal of Ornithology*, *160*(1), 105–115.

https://doi.org/10.1007/s10336-018-1604-6

Iida, M., Yagi, Y., & Iseki, T. (2021). Occurrence of wild chum salmon fry in the surf zone, and

spawning and emergence timing in the adjacent nonstocked river in Niigata Prefecture,

Japan. *Fisheries Science*, *87*(4), 549–557. https://doi.org/10.1007/s12562-021-01535-4

IUCN. (2005). *World Heritage Nomination – IUCN Technical Evaluation, Shiretoko (Japan)*.

https://whc.unesco.org/document/152000

Johnston, R., Jones, K., & Manley, D. (2018). Confounding and collinearity in regression

analysis: a cautionary tale and an alternative procedure, illustrated by studies of British

voting behaviour. *Quality & Quantity*, *52*(4), 1957–1976. https://doi.org/10.1007/s11135-

017-0584-6

Kawaguchi, Y., Taniguchi, Y., & Nakano, S. (2003). Terrestrial invertebrate inputs determine

the local abundance of stream fishes in a forested stream. *Ecology*, *84*(3), 701–708.

https://doi.org/10.1890/0012-9658(2003)084[0701:TIIDTL]2.0.CO;2

Kirillov, P. I., Kirillova, E. A., & Pavlov, D. S. (2018). Patterns of Downstream Migration of

Pink Salmon *Oncorhynchus gorbuscha* in the Malaya Khusi River (Sakhalin Oblast).

Journal of Ichthyology, *58*(6), 889–901. https://doi.org/10.1134/S0032945218060085

Koshino, Y., Kudo, H., & Kaeriyama, M. (2013). Stable isotope evidence indicates the

incorporation into Japanese catchments of marine-derived nutrients transported by

spawning Pacific Salmon. *Freshwater Biology*, *58*(9), 1864–1877.

https://doi.org/10.1111/fwb.12175

Kovach, R. P., Gharrett, A. J., & Tallmon, D. A. (2012). Genetic change for earlier migration

timing in a pink salmon population. *Proceedings of the Royal Society B: Biological*

Sciences, *279*(1743), 3870–3878. https://doi.org/10.1098/rspb.2012.1158

Kuzishchin, K. V., Malyutina, A. M., Gruzdeva, M. A., Savvaitova, K. A., & Pavlov, D. S.

(2009). Reproduction ecology of masu salmon *Oncorhynchus masou* in the Kol basin

(Western Kamchatka). *Journal of Ichthyology*, *49*(6), 441–453.

https://doi.org/10.1134/S0032945209060034

Levi, T., Darimont, C. T., MacDuffee, M., Mangel, M., Paquet, P., & Wilmers, C. C. (2012).

Using grizzly bears to assess harvest-ecosystem tradeoffs in salmon fisheries. *PLoS*

Biology, *10*(4). https://doi.org/10.1371/journal.pbio.1001303

McMullen, L. E., & Lytle, D. A. (2012). Quantifying invertebrate resistance to floods: a global-

scale meta-analysis. *Ecological Applications*, *22*(8), 2164–2175. https://doi.org/10.1890/11-

1650.1

Minakawa, N., & Gara, R. I. (2003). Effects of Chum Salmon Redd Excavation on Benthic

Communities in a Stream in the Pacific Northwest. *Transactions of the American Fisheries*

Society, *132*(3), 598–604. https://doi.org/10.1577/1548-

8659(2003)132<0598:eocsre>2.0.co;2

Moore, J. W., & Schindler, D. E. (2008). Biotic disturbance and benthic community dynamics in

salmon-bearing streams. *Journal of Animal Ecology*, *77*(2), 275–284.

https://doi.org/10.1111/j.1365-2656.2007.01336.x

Morita, K. (2021). Reverse migration of adult pink salmon (*Oncorhynchus gorbuscha*) to the sea

after their return to fresh water. *Environmental Biology of Fishes*, *0123456789*.

https://doi.org/10.1007/s10641-021-01139-y

Munro, J. A. (1941). Studies of waterfowl in British Columbia: Greater scaup duck, lesser scaup

duck. *Canadian Journal of Research*, *19*(4), 113–138. https://doi.org/10.1139/cjr41d-010

Murata, S. (1900). First Group of Hokkaido Birds (2) (in Japanese). *Zoological Magazine*,

12(144), 365–369.

Nakamura, F., & Komiyama, E. (2010). A challenge to dam improvement for the protection of

both salmon and human livelihood in Shiretoko, Japan's third Natural Heritage Site.

Landscape and Ecological Engineering, *6*(1), 143–152. https://doi.org/10.1007/s11355-

009-0083-6

Nakano, S., & Murakami, M. (2001). Reciprocal subsidies: Dynamic interdependence between

terrestrial and aquatic food webs. *Proceedings of the National Academy of Sciences*, *98*(1),

166–170. https://doi.org/10.1073/pnas.98.1.166

Nilsson, A. L. K., L'Abée-Lund, J. H., Vøllestad, L. A., Jerstad, K., Larsen, B. M., Røstad, O.

W., Saltveit, S. J., Skaugen, T., Stenseth, N. C., & Walseng, B. (2018). The potential

influence of Atlantic salmon *Salmo salar* and brown trout *Salmo trutta* on density and

breeding of the white-throated dipper *Cinclus cinclus*. *Ecology and Evolution*, *8*(8), 4065–

4073. https://doi.org/10.1002/ece3.3958

Obermeyer, K. E., Hodgson, A., & Willson, M. F. (1999). American Dipper, *Cinclus mexicanus*,

foraging on Pacific salmon, *Oncorhynchus* sp., eggs. *Canadian Field-Naturalist*, *113*(2),

288–290.

Obermeyer, K. E., White, K. S., & Willson, M. F. (2006). Influence of salmon on the nesting

ecology of American dippers in southeastern Alaska. *Northwest Science*, *80*(1), 26–33.

Ormerod, S. J. (1985). The diet of breeding Dippers *Cinclus cinclus* and their nestlings in the

catchment of the River Wye, mid-Wales: a preliminary study by faecal analysis. *Ibis*,

127(3), 316–331. https://doi.org/10.1111/j.1474-919X.1985.tb05073.x

Ormerod, S. J., & Tyler, S. J. (1986). The diet of dippers *Cinclus cinclus* wintering in the

catchment of the River Wye, Wales. *Bird Study*, *33*(1), 36–45.

https://doi.org/10.1080/00063658609476888

Ormerod, S. J., & Tyler, S. J. (1991). Exploitation of prey by a river bird, the dipper *Cinclus*

cinclus (L.), along acidic and circumneutral streams in upland Wales. *Freshwater Biology*,

25(1), 105–116. https://doi.org/10.1111/j.1365-2427.1991.tb00477.x

Ortlepp, J., & Mürle, U. (2003). Effects of experimental flooding on brown trout (*Salmo trutta*

fario L.): The River Spöl, Swiss National Park. *Aquatic Sciences*, *65*(3), 232–238.

https://doi.org/10.1007/s00027-003-0666-5

Pavlov, D. S., Kirillova, E. A., & Kirillov, P. I. (2008). Patterns and some mechanisms of

downstream migration of juvenile salmonids (with reference to the Utkholok and

Kalkaveyem rivers in Northwestern Kamchatka). *Journal of Ichthyology*, *48*(11), 937–980.

https://doi.org/10.1134/S0032945208110027

Pedersen, M. L., Kristensen, E. A., Kronvang, B., & Thodsen, H. (2009). Ecological effects of

re-introduction of salmonid spawning gravel in lowland Danish streams. *River Research*

and Applications, *25*(5), 626–638. https://doi.org/10.1002/rra.1232

Polis, G. A., Anderson, W. B., & Holt, R. D. (1997). Toward an integration of landscape and

food web ecology: The dynamics of spatially subsidized food webs. *Annual Review of*

Ecology and Systematics, *28*, 289–316. https://doi.org/10.1146/annurev.ecolsys.28.1.289

Quinn, T. P. (2018). *The Behavior and Ecology of Pacific Salmon and Trout* (Second Edi).

University of Washington Press.

R Core Team. (2022). *R: A Language and Environment for Statistical Computing*. https://www.r-

project.org/

Reimchen, T. E. (2017). Diverse ecological pathways of salmon nutrients through an intact

marine-terrestrial interface. *Canadian Field-Naturalist*, *131*(4), 350–368.

https://doi.org/10.22621/cfn.v131i4.1965

Romakkaniemi, A., Perä, I., Karlsson, L., Jutila, E., Carlsson, U., & Pakarinen, T. (2003).

Development of wild Atlantic salmon stocks in the rivers of the northern Baltic Sea in

response to management measures. *ICES Journal of Marine Science*, *60*(2), 329–342.

https://doi.org/10.1016/S1054-3139(03)00020-1

Rundio, D. E., & Lindley, S. T. (2008). Seasonal Patterns of Terrestrial and Aquatic Prey

Abundance and Use by *Oncorhynchus mykiss* in a California Coastal Basin with a

Mediterranean Climate. *Transactions of the American Fisheries Society*, *137*(2), 467–480.

https://doi.org/10.1577/t07-076.1

Schindler, D. E., Scheuerell, M. D., Moore, J. W., Gende, S. M., Francis, T. B., & Palen, W. J.

(2003). Pacific Salmon and the Ecology of Coastal Ecosystems. *Frontiers in Ecology and

the Environment*, *1*(1), 31. https://doi.org/10.2307/3867962

Spiller, D. A., Piovia-Scott, J., Wright, A. N., Yang, L. H., Takimoto, G., Schoener, T. W., &

Iwata, T. (2010). Marine subsidies have multiple effects on coastal food webs. *Ecology*,

91(5), 1424–1434. https://doi.org/10.1890/09-0715.1

Takahashi, G., Kuwahara, T., & Yamanaka, M. (2005). Dams in the Shiretoko Peninsula: Issues

in river management and environmental conservation (in Japanese with English abstract).

Japanese Journal of Conservation Ecology, *10*(2), 139–149.

https://doi.org/https://doi.org/10.18960/hozen.10.2_139

Taylor, A. J., & O'Halloran, J. (1997). The diet of the Dipper *Cinclus cinclus* as represented by

faecal and regurgitate pellets: A comparison. *Bird Study*, *44*(3), 338–347.

https://doi.org/10.1080/00063659709461069

Taylor, A. J., & O'Halloran, J. (2001). Diet of dippers cinclus cinclus during an early winter

spate and the possible implications for dipper populations subjected to climate change. *Bird

Study*, *48*(2), 173–179. https://doi.org/10.1080/00063650109461215

Terui, A., Negishi, J. N., Watanabe, N., & Nakamura, F. (2018). Stream Resource Gradients

Drive Consumption Rates of Supplemental Prey in the Adjacent Riparian Zone.

Ecosystems, *21*(4), 772–781. https://doi.org/10.1007/s10021-017-0183-3

Thedinga, J. F., Wertheimer, A. C., Heintz, R. A., Maselko, J. M., & Rice, S. D. (2000). Effect of

stock, coded-wire tagging, and transplant on straying of pink salmon (*Oncorhynchus*

gorbuscha) in southeastern Alaska. *Canadian Journal of Fisheries and Aquatic Sciences*,

57(10), 2076–2085. https://doi.org/10.1139/f00-163

Tonra, C. M., Sager-Fradkin, K., & Marra, P. P. (2016). Barriers to salmon migration impact

body condition, offspring size, and life history variation in an avian consumer. *Ecography*,

39(11), 1056–1065. https://doi.org/10.1111/ecog.02014

Walters, K. E., Reynolds, J. D., & Ydenberg, R. C. (2021). Ideal free eagles: Bald Eagle

(*Haliaeetus leucocephalus*) distribution in relation to Pacific salmon (*Oncorhynchus* spp.)

availability on four spawning rivers. *Canadian Journal of Zoology*, *99*(9), 792–800.

https://doi.org/10.1139/cjz-2020-0191

Whitehorne, I. (2010). Wintering behavior, physiology and site fidelity in a partial migrant, the

American Dipper (*Cinclus mexicanus*). *Waterbirds*, *33*(4), 461–470.

https://doi.org/10.1675/063.033.0405

Willson, M. F., & Halupka, K. C. (1995). Anadromous Fish as Keystone Species in Vertebrate

Communities. *Conservation Biology*, *9*(3), 489–497. https://doi.org/10.1046/j.1523-

1739.1995.09030489.x

Wipfli, M. S., Hudson, J., & Caouette, J. (1998). Influence of salmon carcasses on stream

productivity: Response of biofilm and benthic macroinvertebrates in southeastern Alaska,

U.S.A. *Canadian Journal of Fisheries and Aquatic Sciences*, *55*(6), 1503–1511.

https://doi.org/10.1139/f98-031

Wipfli, M. S., Hudson, J. P., Chaloner, D. T., & Caouette, J. P. (1999). Influence of salmon

spawner densities on stream productivity in Southeast Alaska. *Canadian Journal of*

Fisheries and Aquatic Sciences, *56*(9), 1600–1611. https://doi.org/10.1139/f99-087

Yamada, T., Urabe, H., & Nakamura, F. (2022). Diel migration pattern of pink salmon fry in

small streams. *Journal of Fish Biology*, *100*(4), 1088–1092.

https://doi.org/10.1111/jfb.15007

TABLES

Table 1. Results of model selection for each response variable (i.e., salmon egg ratio in brown

dipper diets or brown dipper abundance). AIC corresponds to Akaike information criteria; ΔAIC

is the difference between the AIC of that model and that of the model with the lowest AIC. Bold

indicates the best model. Redds, number of spawning redds; Time, survey start time; Age, age

category.

Model	logLik	AIC	ΔAIC	Marginal R^2	Conditional R^2
Salmon egg ratio					
Redds	**-45.98**	**99.97**	**0.00**	**0.319**	**0.494**
Redds + Age	-45.71	101.42	1.45	0.324	0.483
Null	-48.91	103.82	3.85	–	–
Age	-48.59	105.17	5.20	0.090	0.313
Dipper abundance					
Redds	**-63.11**	**132.23**	**0.00**	**0.396**	**0.528**
Redds + Time	-62.66	133.32	1.09	0.410	0.571
Time	-71.60	149.20	16.97	0.188	0.311
Null	-72.85	149.71	17.48	–	–

461   **Table 2.** Results of GLMMs testing effects of number of redds on the salmon egg ratio in brown

462   dipper diets and brown dipper abundance, and the presence/absence of salmon (below/above

463   dam) on brown dipper abundance during peak spawning period.

Response variable	Fixed effect	Estimate	Std. Error	z value
Salmon egg ratio in dipper diets	Intercept	-2.07	1.36	-1.53
	Number of redds	0.05	0.02	2.70
Dipper abundance	Intercept	-6.56	0.20	-32.47
	Number of redds	0.02	0.00	4.40
Dipper abundance during peak spawning period	Intercept	-8.32	0.72	-11.52
	Section -Below dam	2.63	0.75	3.50

FIGURE CAPTIONS

Figure 1. The brown dipper *Cinclus pallasii*. Photo by Yuya Eguchi.

Figure 2. Map of the study area located on the Shiretoko Peninsula. The blue lines indicate the

streams surveyed.

Figure 3. (a) Composition of brown dipper diets during the salmon pre-spawning and spawning

periods. AI: aquatic invertebrate, TI: terrestrial invertebrate, A: algae, and SE: salmon egg. (b)

Mean predicted marginal effects of the number of spawning redds on the salmon egg ratio in

brown dipper diets. The shaded area indicates 95% CI.

Figure 4. Observed brown dipper abundance and number of spawning redds in relation to

surveyed date in each stream.

Figure 5. Mean predicted marginal effects of the number of spawning redds on brown dipper

abundance. The 95% CI is denoted by shaded area.

Figure 6. Mean predicted marginal effects of salmon occurrence on brown dipper abundance

during the peak spawning season. The translucent boxes indicate 95% CIs.

FIGURES

Figure 1.

Sea of Okhotsk

Shoji stream

Chienbetsu stream

Mosekarubetsu stream

Nemuro Channel

Funbe stream

145.12

Hokkaido Is.

44.07

N

0   3   6 km

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.