Concise Communications

Title: Human Galectin-9 Promotes the Expansion of HIV Reservoirs *in vivo* in Humanized Mice

Running title: Galectin-9 and HIV Reservoirs in *vivo*

Authors: Zhe YUAN[1*], Leila B. GIRON[1*], Colin HART, Akwasi GYAMPOH[1], Jane KOSHY[1], Kai Ying HONG[1], Toshiro NIKI[2], Thomas A. PREMEAUX[3], Lishomwa C. NDHLOVU[3], Luis J MONTANER[1#], Mohamed ABDEL-MOHSEN[1#]

Affiliations: [1]The Wistar Institute, Philadelphia, PA, USA; [2]Department of Immunology and Immunopathology, Kagawa University, Kagawa, Japan; [3]Department of Medicine, Division of Infectious Diseases, Weill Cornell Medicine, New York, NY, USA.

* Authors contributed equally to this work

co-corresponding authors

Abstract word count: 249 words; Text word count: 1796 words.

Authors have no competing interests.

FUNDING

This work was primarily funded by a grant from Mizutani Foundation for Glycoscience to M.A-M. Additional funding was provided by the NIH-funded BEAT-HIV Martin Delaney Collaboratory to cure HIV-1 infection (1UM1Al126620), Kean Family Professorship, the Philadelphia Foundation (Roberts I. Jacobs Fund), and NIH awards R01MH112457 and UM1AI164559 to L.C.N.

ABSTRACT [249 words]:

Objective: The human endogenous β-galactoside-binding protein Galectin-9 (Gal-9) reactivates latently HIV-infected cells, which may allow for immune-mediated clearance of these cells. However, Gal-9 also activates T cell Receptor (TCR) signaling pathways, which could negatively affect HIV persistence by promoting T cell expansion and chronic activation/exhaustion. This potential "double-edged sword" effect of Gal-9 during HIV infection raises the question of the overall beneficial versus detrimental impact of Gal-9 on HIV persistence *in vivo*.

Design: We used the BLT (bone marrow, liver, thymus) humanized mouse model to evaluate the overall impact of Gal-9 on HIV persistence *in vivo* during antiretroviral therapy (ART).

Methods: Two independent cohorts of BLT mice with high human immune reconstitution were infected with HIV, placed on ART, and then treated with either recombinant human Gal-9 or PBS during ART suppression. Plasma viral loads and levels of tissue-associated HIV DNA and RNA were measured by qPCR. Markers of T cell activation/exhaustion were measured by flow cytometry, and plasma markers of inflammation were measured by multiplex cytokine arrays.

Results: Gal-9 treatment was tolerable in ART-suppressed humanized mice and did not significantly induce plasma markers of inflammation or T cell markers of activation/exhaustion. However, Gal-9 treatment during ART significantly increased levels of tissue-associated HIV DNA and RNA compared to controls (P=0.0007 and P=0.011, respectively, for cohort I and P=0.002 and P=0.005, respectively, for cohort II).

Conclusions: Our study highlights the overall adverse effects of Gal-9 on HIV persistence and the potential need to block Gal-9 interactions during ART-suppressed HIV infection.

Key Words: Galectin-9; HIV; humanized mice; BLT mice; HIV reservoirs.

TEXT [1796 words]

INTRODUCTION

The persistence of HIV latently-infected cells in blood and tissues despite effective antiretroviral

drug therapy (ART) remains a barrier to a cure for HIV infection [1]. Persistent latently-infected

cells do not express enough viral antigens to be eliminated by the immune system, and different

strategies to reactivate these cells have been proposed, including the use of the human lectin

Galectin-9 (Gal-9) [2, 3]. However, Gal-9 modulates HIV transcription through activating the T-

cell receptor (TCR)-downstream ERK and CREB signaling pathways [4] and this Gal-9-mediated

activation can lead to several undesirable effects due to T cell expansion, activation, and/or

exhaustion. Indeed, recent studies showed that Gal-9, which is rapidly and sustainably elevated

during HIV infection [5], may contribute to the state of chronic immune activation and

inflammation during HIV infection [6-8]. This potential "double-edged sword" of Gal-9 during

HIV infection raises the question on how the potentiation of reactivating latent infection and the

undesirable outcomes on T cell effector function due to Gal-9 effects HIV persistence *in vivo*.

Thus, evaluating the overall impact of Gal-9 on HIV persistence requires a relevant *in vivo*

animal model of HIV infection during ART suppression. In this short report, we investigate the

impact of Gal-9 in a humanized mouse model of HIV infection (the bone marrow-liver-thymus

humanized (BLT) mouse model of HIV latency) to determine the overall beneficial versus

detrimental impact of Gal-9 on HIV persistence *in vivo*.

METHODS

Generation of the BLT mice. BLT mice were generated as previously described [9, 10], in

accordance with The Wistar Institute Animal Care and Research Committee regulations. Briefly,

6-8 weeks old female NSG (NOD.Cg-Prkdcscid Il2rgtm1Wjl/SzJ, Jackson Laboratory) mice

were pretreated with busulfan at 30mg/kg and were then implanted with human fetal thymic

tissue fragments and fetal liver tissue fragments under the murine renal capsule. Following the

surgery, mice were injected via the tail vein with CD34$^+$ hematopoietic stem cells isolated from

human fetal liver tissues. Human fetal liver and thymus tissues were procured from Advanced

Bioscience Resources (Alameda, CA). Twelve weeks post-surgery, human immune cell

reconstitution in peripheral blood was determined by Symphony flow cytometer (BD
Biosciences, San Jose, CA) using the following antibodies: mCD45-AF700, hCD45-FITC,
hCD3-BUV805, hCD4-BUV395, hCD8-PerCP-Cy5.5 and Fixable Viability Stain 510 (BD
Biosciences, San Jose, CA). Data were analyzed with FlowJo (FlowJo LLC, Ashland, OR).

HIV infection, ART suppression, and Gal-9 treatment. BLT mice were randomly divided into
two groups and were infected intravenously (IV) with 1×10^4 $TCID_{50}$ of HIV_{SUMA}. Peripheral
blood was collected weekly for plasma viral load assay. Two weeks post-infection, mice were
placed on a diet combined with ART (1,500mg/kg Emtricitabine, 1,560mg/kg Tenofovir-
Disoproxil-Fumarate, and 600mg/kg Raltegravir). Five weeks post-ART, mice were treated with
phosphate-buffered saline (PBS) control or 2 mg/kg recombinant Gal-9 for two weeks
(intraperitoneal (IP) injections every other day; seven doses) during ART suppression. Mice
were then euthanized, and blood and tissues were collected.

Measuring plasma viral load by qPCR. Plasma viral loads were measured as previously
described [9, 10]. Briefly, viral RNA was extracted using the QIAamp Viral RNA Mini kit
(Qiagen). Plasma viral load was measured using qRT-PCR on a C1000 Thermal Cycler and the
CFX96 Real-Time system (Bio-Rad) with 100 copies/ml limit of detection.

Measuring cell-associated HIV DNA and RNA by qPCR. Single cells are isolated with
gentleMACS™ Octo Dissociator (San Diego, CA). DNA and RNA were extracted using AllPrep
DNA/RNA/miRNA Universal Kit (Qiagen, catalog # 80224). Cell-associated HIV DNA was
measured by qPCR using LTR-specific primers F522-43 (5' GCC TCA ATA AAG CTT GCC
TTG A 3'; HXB2522–543) and R626-43 (5' GGG CGC CAC TGC TAG AGA 3'; 626–643)
coupled with a FAM-BQ probe (5' CCA GAG TCA CAC AAC AGA CGG GCA CA 3). Cell
count was estimated by qPCR using human copy number reference assay TERT (Applied
Biosystems, cat# 4403316). Cell-associated HIV RNA was measured using the same primers in a
one-step qPCR using TaqMan® RNA to Ct™ 1 Step kit (Applied Biosystems, catalog #
4392656). Cell counts were determined by qPCR using human RPLP0 (Applied Biosystems,
catalog # 4310879E).

Measuring markers of T cell activation and exhaustion by flow cytometry. Cell suspension was
stained with the following antibodies: CD45-AL700, CD8-FITC, CD38-APC, CD25-PerCP-
Cy5.5, HLA-DR-APC-H7, CD4-V450, PD-1-PE, CD3-PE-CF594, and CD69-PE-Cy7. Data

were collected on a BD Biosciences LSRII flow cytometer (gating strategy is in **Supplementary**

Figure 1).

Measuring plasma markers of inflammation. Markers of inflammation were measured using U-

PLEX Biomarker Group 1 Assay from Meso Scale Discovery (MSD catalog # K15067L-2).

Statistical analysis. Mann–Whitney tests were used for the two-group comparisons in Figures

1B, G, H, and unpaired t-tests were used for the two-group comparisons in Figure 2. Statistical

analyses were performed using Prism 9.0 (GraphPad).

RESULTS

Gal-9 treatment is tolerable *in vivo* but increases levels of tissue-associated HIV DNA and

RNA during ART. We generated two independent cohorts of humanized mice (n=9 for cohort I

and n=6 for cohort II) (**Fig 1A**) and achieved high levels of immune reconstitution as measured

by the percentage of human $CD45^+$ cells in blood (**Fig. 1B**). BLT mice were infected with HIV

(HIV_{SUMA} transmitted/founder virus) and then placed on ART two weeks post-infection. During

ART suppression, mice were treated with either PBS or 2 mg/kg recombinant human Gal-9 for

two weeks (every other day; seven doses). Mice were then euthanized, and blood and tissues

(liver, spleen, and lung) were collected (**Fig. 1A**). Gal-9 treatment did not affect the weight of the

mice from both cohorts compared to controls **(Fig. 2C-D)**, suggesting this concentration was

generally tolerable in ART-suppressed HIV-infected BLT mice. Plasma HIV viral loads shown

in **Fig. 1E-F** indicate that, in both cohorts, the infection was successful and resulted in high viral

loads by the second week post-infection and that ART suppressed the virus to below the limit of

detection. We have not observed viral blips in the Gal-9 treated mice to suggest any significant

viral reversal effects. We next examined cell-associated HIV RNA and DNA levels in the liver,

lung, and spleen. We found that Gal-9 treatment increased the levels of tissue-associated HIV

DNA (*P*=0.0007) and RNA (*P*=0.0106) (**Fig. 1G**) in cohort I. Consistently, in cohort II, Gal-9

increased tissue-associated HIV DNA (*P*=0.0019) and RNA (*P*=0.0047) (**Fig. 1H**). Collectively,

these data suggest that the overall impact of Gal-9 on HIV persistence is negative, and that Gal-9

expands the levels of tissue-associated HIV reservoirs during ART.

Gal-9 treatment does not significantly induce markers of inflammation of T cell activation.
Next, we examined whether Gal-9 exhibits its adverse effects on HIV persistence by inducing T cell activation/exhaustion or systemic inflammation. We first examined markers of $CD4^+$ and $CD8^+$ T cell activation/exhaustion in blood and spleen (collected at the time of mice euthanization from cohort I). Gal-9 treatment did not significantly induce T cell activation (as measured by the co-expression of CD38 and HLA-DR or the co-expression of CD69 and CD25 activation markers) or exhaustion (as measured by PD-1 expression) in blood (**Fig. 2A**) or tissues (**Fig. 2B**). However, a trend toward an increase in $CD8^+$ T cell expression of CD38 and HLA-DR was observed in the blood in the Gal-9 treated mice compared to controls (p=0.0767, **Fig. 2A**). Next, we examined the plasma levels of several cytokines. While Gal-9 treatment reduced the levels of IL-33 (p=0.0321), it did not significantly alter the systemic levels of several cytokines involved in inflammation (**Fig. 2C**). These data suggest that Gal-9 does not induce levels of cell-associated HIV DNA and RNA by inducing generalized T cell activation or systemic inflammation.

DISCUSSION

A comprehensive understanding of the overall impact of viral and host factors that modulate HIV persistence is critical to develop curative strategies for HIV. Gal-9 is one of the endogenous host immune-modulatory factors that has been recently associated with opposing effects on HIV infection [5, 11, 12]. Several studies have highlighted the potential beneficial effects of Gal-9 during HIV infection. First, Gal-9 renders $CD4^+$ T cells less susceptible to HIV infection via induction of the host restriction factor cyclin-dependent kinase inhibitor 1 (p21) [13]. Second, recombinant Gal-9 induces HIV transcription and reverses HIV latency *in vitro* and *ex vivo*. This ability of Gal-9 to induce latent HIV transcription suggested that it could be considered within the "shock and kill" HIV eradication framework [2, 3]. However, on the other hand, several other studies have highlighted the detrimental effects of Gal-9 during HIV infection. First, endogenous Gal-9 is rapidly increased after HIV infection [5] and do not return to normal after suppressive ART [5]. These persistent elevated Gal-9 levels are associated with higher HIV transcription *in vivo* in plasma of HIV-infected ART-suppressed individuals [2], and several studies suggest may contribute to the state of chronic immune activation and inflammation during HIV infection [6-8].

Second, Gal-9 increases HIV entry by inducing the CD4[+] T cell-surface concentration of protein

disulfide isomerase (PDI) [14]. Lastly, Gal-9 modulates HIV transcription by activating the TCR-

downstream signaling pathways *in vitro* [4]. Due to these pleiotropic effects of Gal-9 on HIV, we

sought to evaluate whether the overall impact of Gal-9 on HIV persistence is beneficial or

detrimental *in vivo*.

Using the BLT humanized mouse model of HIV infection, we found that Gal-9 can directly

expand HIV reservoirs and exhibit overall adverse effects on HIV persistence. These effects

would limit the potential use of Gal-9 to reduce viral reservoirs by itself. However, future studies

will be needed to investigate whether Gal-9 together with added immune effector strategies

would positively affect HIV persistence. Future studies will also be needed using larger animal

models with a more intact immune system compared to the BLT humanized mouse model. Our

data also show that the Gal-9-mediated expansion of HIV reservoirs is not linked to elevated T

cell activation or systemic inflammation. However, Gal-9 can induce T cell proliferation [2],

which might be linked to its modulation of HIV persistence. Furthermore, Gal-9 can impact

many aspects of immune responses relevant to persistence. For example, Gal-9 exhibits several

immunosuppressive activities [15-23], including the ability to increase the function of regulatory T

cells (T-regs)[23] and to impair natural killer (NK) cells' cytotoxicity [24]. The potential link

between Gal-9-mediated impact on T cell proliferation or immune functions and its ability to

impact HIV reservoir size warrants a broader investigation.

Endogenous Gal-9 is highly abundant *in vivo*, especially during HIV infection[5], and its

sustained levels have been associated, during ART-suppressed HIV infection, with the state of

chronic inflammation and immune activation that is central to the development of several HIV-

associated co-morbidities [25-32]. Therefore, clarifying the mechanistic underpinnings of the

overall adverse effects of elevated Gal-9 during ART-suppressed HIV infection may lead to the

development of interventions to target Gal-9 (such as anti-Gal-9 antibodies and small molecule

inhibitors targeting Gal-9 [33-35]) to improve immune functionality, reduce inflammation-

associated co-morbidities, and reduce levels of HIV persistence, in the setting of viral

suppression by ART.

REFERENCES:

1. Deeks SG, Archin N, Cannon P, Collins S, Jones RB, de Jong M, et al. **Research priorities for an HIV cure: International AIDS Society Global Scientific Strategy 2021**. *Nat Med* 2021; 27(12):2085-2098.

2. Abdel-Mohsen M, Chavez L, Tandon R, Chew GM, Deng X, Danesh A, et al. **Human Galectin-9 Is a Potent Mediator of HIV Transcription and Reactivation**. *PLoS Pathog* 2016; 12(6):e1005677.

3. Sanz M, Madrid-Elena N, Serrano-Villar S, Vallejo A, Gutierrez C, Moreno S. **Effect of the use of Galectin-9 and blockade of TIM-3 receptor in the latent cellular reservoir of HIV-1**. *J Virol* 2020.

4. Colomb F, Giron LB, Premeaux TA, Mitchell BI, Niki T, Papasavvas E, et al. **Galectin-9 Mediates HIV Transcription by Inducing TCR-Dependent ERK Signaling**. *Front Immunol* 2019; 10:267.

5. Tandon R, Chew GM, Byron MM, Borrow P, Niki T, Hirashima M, et al. **Galectin-9 is rapidly released during acute HIV-1 infection and remains sustained at high levels despite viral suppression even in elite controllers**. *AIDS Res Hum Retroviruses* 2014; 30(7):654-664.

6. Dunsmore G, Rosero EP, Shahbaz S, Santer DM, Jovel J, Lacy P, et al. **Neutrophils promote T-cell activation through the regulated release of CD44-bound Galectin-9 from the cell surface during HIV infection**. *PLoS Biol* 2021; 19(8):e3001387.

7. Premeaux TA, D'Antoni ML, Abdel-Mohsen M, Pillai SK, Kallianpur KJ, Nakamoto BK, et al. **Elevated cerebrospinal fluid Galectin-9 is associated with central nervous system immune activation and poor cognitive performance in older HIV-infected individuals**. *J Neurovirol* 2019; 25(2):150-161.

8. Premeaux TA, Moser CB, McKhann A, Hoenigl M, Laws EI, Aquino DL, et al. **Plasma galectin-9 as a predictor of adverse non-AIDS events in persons with chronic HIV during suppressive antiretroviral therapy**. *AIDS* 2021; 35(15):2489-2495.

9. Yuan Z, Kang G, Daharsh L, Fan W, Li Q. **SIVcpz closely related to the ancestral HIV-1 is less or non-pathogenic to humans in a hu-BLT mouse model**. *Emerg Microbes Infect* 2018; 7(1):59.

10. Yuan Z, Kang G, Ma F, Lu W, Fan W, Fennessey CM, et al. **Recapitulating Cross-Species Transmission of Simian Immunodeficiency Virus SIVcpz to Humans by Using Humanized BLT Mice**. *J Virol* 2016; 90(17):7728-7739.

11. Sehrawat S, Reddy PB, Rajasagi N, Suryawanshi A, Hirashima M, Rouse BT. **Galectin-9/TIM-3 interaction regulates virus-specific primary and memory CD8 T cell response**. *PLoS pathogens* 2010; 6(5):e1000882.

12. Jost S, Moreno-Nieves UY, Garcia-Beltran WF, Rands K, Reardon J, Toth I, et al. **Dysregulated Tim-3 expression on natural killer cells is associated with increased Galectin-9 levels in HIV-1 infection**. *Retrovirology* 2013; 10:74.

13. Elahi S, Niki T, Hirashima M, Horton H. **Galectin-9 binding to Tim-3 renders activated human CD4+ T cells less susceptible to HIV-1 infection**. *Blood* 2012; 119(18):4192-4204.

14. Schaefer K, Webb NE, Pang M, Hernandez-Davies JE, Lee KP, Gonzalez P, et al. **Galectin-9 binds to O-glycans on protein disulfide isomerase**. *Glycobiology* 2017; 27(9):878-887.

15. Seki M, Oomizu S, Sakata KM, Sakata A, Arikawa T, Watanabe K, et al. **Galectin-9 suppresses the generation of Th17, promotes the induction of regulatory T cells, and regulates experimental autoimmune arthritis**. *Clinical immunology* 2008; 127(1):78-88.

16. Wang F, He W, Zhou H, Yuan J, Wu K, Xu L, et al. **The Tim-3 ligand galectin-9 negatively regulates CD8+ alloreactive T cell and prolongs survival of skin graft**. *Cellular immunology* 2007; 250(1-2):68-74.

17. Zhu C, Anderson AC, Schubart A, Xiong H, Imitola J, Khoury SJ, et al. **The Tim-3 ligand galectin-9 negatively regulates T helper type 1 immunity**. *Nat Immunol* 2005; 6(12):1245-1252.

18. Lv K, Zhang Y, Zhang M, Zhong M, Suo Q. **Galectin-9 promotes TGF-beta1-dependent induction of regulatory T cells via the TGF-beta/Smad signaling pathway**. *Mol Med Rep* 2013; 7(1):205-210.

19. Yang S, Wang J, Chen F, Liu G, Weng Z, Chen J. **Elevated Galectin-9 Suppresses Th1 Effector Function and Induces Apoptosis of Activated CD4(+) T Cells in Osteoarthritis**. *Inflammation* 2017; 40(3):1062-1071.

20. Panda SK, Facchinetti V, Voynova E, Hanabuchi S, Karnell JL, Hanna RN, et al. **Galectin-9 inhibits TLR7-mediated autoimmunity in murine lupus models**. *J Clin Invest* 2018; 128(5):1873-1887.

21. Ungerer C, Quade-Lyssy P, Radeke HH, Henschler R, Konigs C, Kohl U, et al. **Galectin-9 is a suppressor of T and B cells and predicts the immune modulatory potential of mesenchymal stromal cell preparations**. *Stem Cells Dev* 2014; 23(7):755-766.

22. Kim SN, Lee HJ, Jeon MS, Yi T, Song SU. **Galectin-9 is Involved in Immunosuppression Mediated by Human Bone Marrow-derived Clonal Mesenchymal Stem Cells**. *Immune Netw* 2015; 15(5):241-251.

23. Wu C, Thalhamer T, Franca RF, Xiao S, Wang C, Hotta C, et al. **Galectin-9-CD44 interaction enhances stability and function of adaptive regulatory T cells**. *Immunity* 2014; 41(2):270-282.

24. Golden-Mason L, McMahan RH, Strong M, Reisdorph R, Mahaffey S, Palmer BE, et al. **Galectin-9 functionally impairs natural killer cells in humans and mice**. *J Virol* 2013; 87(9):4835-4845.

25. Hunt PW, Martin JN, Sinclair E, Bredt B, Hagos E, Lampiris H, et al. **T cell activation is associated with lower CD4+ T cell gains in human immunodeficiency virus-infected patients with sustained viral suppression during antiretroviral therapy**. *J Infect Dis* 2003; 187(10):1534-1543.

26. Hunt PW, Cao HL, Muzoora C, Ssewanyana I, Bennett J, Emenyonu N, et al. **Impact of CD8+ T-cell activation on CD4+ T-cell recovery and mortality in HIV-infected Ugandans initiating antiretroviral therapy**. *Aids* 2011; 25(17):2123-2131.

27. Kaplan RC, Sinclair E, Landay AL, Lurain N, Sharrett AR, Gange SJ, et al. **T cell activation and senescence predict subclinical carotid artery disease in HIV-infected women**. *The Journal of infectious diseases* 2011; 203(4):452-463.

28. Kaplan RC, Sinclair E, Landay AL, Lurain N, Sharrett AR, Gange SJ, et al. **T cell activation predicts carotid artery stiffness among HIV-infected women**. *Atherosclerosis* 2011; 217(1):207-213.

29. Chomont N, El-Far M, Ancuta P, Trautmann L, Procopio FA, Yassine-Diab B, et al. **HIV reservoir size and persistence are driven by T cell survival and homeostatic proliferation**. *Nature medicine* 2009; 15(8):893-900.

30. Chun TW, Justement JS, Pandya P, Hallahan CW, McLaughlin M, Liu S, et al. **Relationship between the size of the human immunodeficiency virus type 1 (HIV-1) reservoir in peripheral blood CD4+ T cells and CD4+:CD8+ T cell ratios in aviremic HIV-1-infected**

individuals receiving long-term highly active antiretroviral therapy. *J Infect Dis* 2002; 185(11):1672-1676.

31. Hatano H, Jain V, Hunt PW, Lee TH, Sinclair E, Do TD, et al. **Cell-based measures of viral persistence are associated with immune activation and programmed cell death protein 1 (PD-1)-expressing CD4+ T cells**. *J Infect Dis* 2013; 208(1):50-56.

32. Khoury G, Fromentin R, Solomon A, Hartogensis W, Killian M, Hoh R, et al. **Human Immunodeficiency Virus Persistence and T-Cell Activation in Blood, Rectal, and Lymph Node Tissue in Human Immunodeficiency Virus-Infected Individuals Receiving Suppressive Antiretroviral Therapy**. *J Infect Dis* 2017; 215(6):911-919.

33. Di Gaetano S, Pirone L, Galdadas I, Traboni S, Iadonisi A, Pedone E, et al. **Design, Synthesis, and Anticancer Activity of a Selenium-Containing Galectin-3 and Galectin-9N Inhibitor**. *Int J Mol Sci* 2022; 23(5).

34. Tanino Y, Hashimoto T, Ojima T, Mizuno M. **F-fucoidan from Saccharina japonica is a novel inducer of galectin-9 and exhibits anti-allergic activity**. *J Clin Biochem Nutr* 2016; 59(1):25-30.

35. Yang R, Sun L, Li CF, Wang YH, Xia W, Liu B, et al. **Development and characterization of anti-galectin-9 antibodies that protect T cells from galectin-9-induced cell death**. *J Biol Chem* 2022; 298(4):101821.

FIGURE LEGENDS:

Figure 1. Gal-9 treatment is tolerable *in vivo* but increases levels of tissue-associated HIV DNA and RNA during ART-suppressed HIV infection. (**A**) a schematic overview of the study design. (**B**) percentage of human CD45[+] cells measured in the peripheral blood of BLT mice. No difference in levels of human immune reconstitution between mice treated with Gal-9 versus mice treated with PBS control. Mann Whitney t test. Median and interquartile range (IQR) are displayed. (**C-D**) mice weight over time in cohort I (C) and cohort II (D). (**E-F**) plasma viral load measured by ddPCR over time in cohort I (E) and cohort II (F). (**G-H**) Gal-9 treatment-induced levels of tissue-associated HIV DNA and HIV RNA (measured by qPCR) than controls in both cohort I (G) and cohort II (H). Data shown are only from tissues with detectable levels of cell-associated HIV DNA or RNA. Mann Whitney t-tests. Median and interquartile range (IQR) are displayed.

Figure 2. Gal-9 treatment does not significantly induce markers of inflammation of T cell activation. (**A-B**) impact of Gal-9 treatment on markers of CD4[+] and CD8[+] T cell activation (co-expression of CD38 and HLA-DR or the co-expression of CD69 and CD25 activation markers) and exhaustion (PD-1 expression) measured by flow cytometry in the blood (A) and spleen (B). Unpaired t-tests. The mean and standard mean of error (SEM) are displayed. (**C**) Impact of Gal-9 treatment on plasma levels of markers of systemic inflammation measured by multiplex arrays. The mean and standard mean of error (SEM) are displayed.

SUPPLEMENTARY MATERIALS:

Supplementary Figure 1. A gating strategy [Leila]. A representative example of the gating strategy used for the phenotyping of markers of T cell activation and exhaustion in Figure 2.

AUTHOR CONTRIBUTIONS

Z.Y, L.B.G, T.N, L.C.N, L.J.M, and M.A-M designed the experiments. Z.Y, L.B.G, C.H, A.G, J.K, and K.Y.H carried out the experiments. Z.Y, L.B.G, L.J.M, M.A-M analyzed and interpreted data. Z.Y, L.B.G, L.J.M, M.A-M wrote the manuscript, and all authors edited it.

COMPETING INTERESTS STATEMENT

Authors have no competing interests.

CORRESPONDING AUTHOR CONTACT INFORMATION

Luis J. Montaner, D.V.M., M.Sc., D.Phil.

Professor, Vaccine and Immunotherapy Center, The Wistar Institute

3601 Spruce St., Philadelphia, PA 19104

Phone: 215-898-9143

Email: montaner@wistar.org

Mohamed Abdel-Mohsen, Ph.D.

Associate Professor, Vaccine and Immunotherapy Center, The Wistar Institute

3601 Spruce St., Philadelphia, PA 19104

Phone: 215-898-6008

Email: mmohsen@Wistar.org

Figure 1

Figure 2