Spherical harmonics representation of the steady-state membrane potential shift induced by tDCS in realistic neuron models

Adrià Galan-Gadea[1]**, Ricardo Salvador**[1]**, Fabrice Bartolomei**[2]**, Fabrice Wendling**[3] **and Giulio Ruffini**[1]‡

[1] Neuroelectrics, Av. Tibidabo 47b, 08035 Barcelona, Spain
[2] Clinical Physiology Department, INSERM, UMR 1106 and Timone University Hospital, Aix-Marseille Université, Marseille, France
[3] Univ Rennes, INSERM, LTSI (Laboratoire de Traitement du Signal et de l'Image) U1099, 35000 Rennes, France

E-mail: `giulio.ruffini@neuroelectrics.com`

Abstract. *Objective:* We provide a systematic framework for quantifying the effect of externally applied weak electric fields on realistic neuron compartment models as captured by physiologically relevant quantities such as the membrane potential or transmembrane current as a function of the orientation of the field. *Approach:* We define a response function as the steady-state change of the membrane potential induced by a canonical external field of 1 V/m as a function of its orientation. We estimate the function values through simulations employing reconstructions of the rat somatosensory cortex from the Blue Brain Project. The response of different cell types is simulated using the NEURON simulation environment. We represent and analyze the angular response as an expansion in spherical harmonics. *Main results:* We report membrane perturbation values comparable to those in the literature, extend them to different cell types, and provide their profiles as spherical harmonic coefficients. We show that at rest, responses are dominated by their dipole terms ($\ell = 1$), in agreement with experimental findings and compartment theory. Indeed, we show analytically that for a passive cell, only the dipole term is nonzero. However, while minor, other terms are relevant for states different from resting. In particular, we show how $\ell = 0$ and $\ell = 2$ terms can modify the function to induce asymmetries in the response. *Significance:* This work provides a practical framework for the realistic representation of the effects of weak electric fields on different neuron types and their main regions— an important milestone for developing micro- and mesoscale models and optimizing brain stimulation solutions.

Keywords: Trancranial Direct Current Stimulation (tDCS), spherical harmonics, Non-Invasive Brain Stimulation (NIBS), compartment models, electric field effects

---

‡ giulio.ruffini(at)neuroelectrics.com

1. Introduction

Transcranial electrical stimulation (tES) is a non-invasive neuromodulatory technique based on accurate control of weak currents injected from multiple scalp electrodes and the resulting electric fields induced in the brain. tES, pioneered by Nitsche and Paulus [1], includes direct and alternating current variants known as tDCS and tACS. Low intensity, controlled currents (typically ∼1 mA but ≤ 4 mA) are applied through scalp electrodes in repeated 20–60 min sessions. The subtle but persistent modulation of neuronal activity is believed to lead to plastic effects deriving from Hebbian mechanisms [2].

The current evolution of this and related techniques such as transcranial magnetic stimulation or deep brain stimulation is veering towards model-driven approaches and optimization principles based on a mechanistic understanding of pathophysiology, the interaction of the electric field with neuron populations and the resulting dynamical, and then plastic, effects [3, 4]. In this work, we aim to quantify in a systematic way the response of different cell types and cell parts to be used in future approaches to model-driven stimulation.

The concurrent effects of tES currents on the brain are thought to be mediated by the generated electric field's action on cortical neurons [5, 6]. These electric fields are weak (∼1 V/m) [7], relatively large-scale (with spatial correlation scales of the order of a few cm), and induce small changes in the membrane potential of cells (sub-mV) [8, 9]. Polarization happens in a compartment-specific manner [10, 11, 12], hyperpolarizing compartments close to the virtual anode and depolarizing those close to the virtual cathode. Furthermore, it is expected from cable theory that changes in the electric field along neuron fibers cause local polarizations as well. Then, the effect in a cell under the same stimulation can have very different responses locally. The effects are significantly larger where fibers bend and mostly at terminations [13]. Thus, the overall morphology is key to explaining effects observed locally. To take all effects into account, we use multi-compartment models that allow us to locally couple the field to realistic morphology models and make predictions of the induced perturbations [14].

Pyramidal cells have been of particular interest in the field for two reasons. First, they are elongated, which leads to the generation of a larger membrane perturbation when the field and neuron are aligned [9]. Secondly, they are spatially organized. Because of this and the spatial homogeneity of the electric fields, the effects on such cells are spatially coherent and thought to lead to stronger, network-enhanced effects [15, 16]. In contrast, effects on interneurons are usually disregarded due to their small size and non-elongated branching profiles [9].

So-called hybrid models combining physics and physiology represented by computational models of brain networks can be used to represent pathophysiology and the effects of electric fields [17]. A widely used example of physiology models at the mesoscale is neural mass models (NMM) [18, 19, 20] and their more recent extensions (NMM2) [21, 22, 23]. To be used in model-driven stimulation, they all require a

description of the electric field in the cortex — which can be provided using realistic finite element modeling [6, 24] — as well as an interaction model for the effects of the field on neuron populations. In the case of classical neural mass models (NMM), this can be provided from an estimation of the field-induced changes in membrane potential or transmembrane current in relevant compartments such as the soma at the population level. A practical starting point for this is modeling the effects at the single cell level.

In this paper, we focus on characterizing the effects of the weak electric field on the stable equilibrium of single neuron models using expansions in terms of spherical harmonics. With this approach, we aim to provide (i) a systematized method for describing the angular dependence of the response through simulations, (ii) a framework that allows for the averaging of the angular response of many compartments to describe cell parts and types, and (iii) a representation in meaningful terms that represent the dipolar behavior on its own but accounts for other angular dependence profiles if necessary. We validate the common dipolar approach and that it is accurate for small perturbations induced at neurons at rest (far from the threshold) and show that it is exact for the passive membrane. Nevertheless, we observe that corrections to pure dipolar responses may need to be applied closer to the threshold in realistic neurons.

2. Methods

2.1. Formalism for steady-state changes induced by an electric field

We will be studying the response of a neuron to a constant, spatially uniform, weak electric field as typically generated by non-invasive brain stimulation methods such as tDCS — at least away from tissue boundaries, since these can introduce strong local changes in the field distribution due to the change in electrical properties between tissues. The field is characterized by a magnitude $E \equiv ||\mathbf{E}||$ and an orientation that can be expressed with two spherical angles, the polar angle θ and the azimuth φ. Thus, in spherical coordinates, the electric field is given by

$$\mathbf{E} = E \left(\sin \theta \cos \varphi \, \mathbf{e}_x + \sin \theta \sin \varphi \, \mathbf{e}_y + \cos \theta \, \mathbf{e}_z \right) \equiv E \, \hat{\mathbf{e}} \tag{1}$$

Let us consider a variable v that characterizes a physiologically relevant magnitude affected by an external electric field. Our study, focuses on the membrane potential, but the formalism can be extended to other variables, e.g., transmembrane currents. If the variable v has a stable equilibrium, i.e., resting potential in our case, we expect that the applied electric field will shift that equilibrium to a new value v'. We aim to quantify that induced change δv of the steady-state solution for different realistic models of cell types. In our case, this shift is commonly referred to as polarization.

Let us assume that two independent terms can express the effect of the magnitude and orientation. On the one hand, the polarization δv scales linearly with E, which has been observed experimentally for small fields [9]. Regarding the dependency of the shift to the orientation of the field, we propose a function defined on a sphere $\Phi(\theta, \varphi)$, i.e., that has a value for any direction in space. We refer to Φ as the *response function*, which is a function that outputs the shift δv under an electric field of 1 V/m in a given direction. Under these assumptions §, δv can be expressed as shown in Equation 2.

$$\delta v = E \, \Phi(\hat{\mathbf{e}}) = E \, \Phi(\theta, \phi) \tag{2}$$

2.2. Expansion in spherical harmonics

Spherical harmonics form a complete and orthonormal set of functions $Y_{\ell m}$ defined on the surface of the sphere (parametrized by two spherical angles, θ and φ), where ℓ and m refer to the degree and the order of the harmonic, respectively. Squared-integrable functions defined on the sphere can be expressed as a sum of spherical harmonic terms [25, 26], much as functions on the plane can be expressed as a Fourier series. In fact, both spherical harmonics and complex exponential are eigenfunctions of equations associated with the Laplace partial differential equation in different manifolds (the plane and the sphere). As the response function $\Phi(\theta, \varphi)$ is a function on the sphere, it can be expanded

§ More generally, we can start from $\delta v(\mathbf{E}) = \delta v(E, \hat{\mathbf{e}}) = \Phi_0(\hat{\mathbf{e}}) + E\Phi_1(\hat{\mathbf{e}}) + \frac{1}{2}E^2\Phi_2(\hat{\mathbf{e}}) + ...$ The first term is zero here by definition.

in terms of spherical harmonics, meaning that it can be expressed as in Equation 3 and fully characterized by a set of coefficients $f_{\ell m}$,

$$\Phi(\theta, \varphi) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} f_{\ell m} Y_{\ell m}(\theta, \varphi) \tag{3}$$

Spherical harmonics are characterized by a degree and order pair (ℓ, m); see Appendix A for a definition. They are tightly linked to the multipole expansion, so one can think of the $\ell = 0$ term as a *monopole* behavior, a term that does not depend on orientation. The $\ell = 1$ terms represent a dipolar behavior, $\ell = 2$ a quadrupole, and so on. Finally, under a parity transformation, where a point with coordinates $\{\theta, \phi\}$ changes to $\{\pi - \theta, \pi + \phi\}$ (corresponding to a reversal of the orientation of the electric field in our case),

$$Y_{\ell m}(\theta, \varphi) \rightarrow Y_{\ell m}(\pi - \theta, \pi + \varphi) = (-1)^{\ell} Y_{\ell m}(\theta, \varphi) \tag{4}$$

This means that if we expect the membrane response to change sign under electric field reversal (e.g., if it is linear in $\mathbf{E}$), only odd ℓ terms should contribute to the expansion.

The l-2 norm of the coefficients is an influenceability measure. The l-2 norm of the coefficients quantifies the maximum of the response function regardless of the orientation. The actual values are subject to the specific normalization used. We name this measure the influenceability ξ of the response function,

$$\xi \equiv \|f_{\ell m}\| = \sqrt{\sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} f_{\ell m}^2}, \tag{5}$$

and serves as a metric to compare how affected in magnitude by an electric field each model is. In other words, for a set of response functions $f_{\ell m}$, we can compare how susceptible they are to being influenced given the same electric field.

The λE linear model is contained in the $\ell = 1$ terms of the expansion. The f_{1m} terms correspond to the λ (dipole) in the λE model, where the induced polarization is represented as the dot product of the electric field vector $\mathbf{E}$ and a vector $\boldsymbol{\lambda}$ of magnitude the length constant of the membrane and in the orthodromic direction of the cell [27]. We can rewrite the λE model as

$$\delta v = \boldsymbol{\lambda} \cdot \boldsymbol{E} = E \boldsymbol{\lambda} \cdot \boldsymbol{e} = E \left(\lambda_x \sin\theta \cos\varphi + \lambda_y \sin\theta \sin\varphi + \lambda_z \cos\theta \right), \tag{6}$$

which is seen to correspond to the $\ell = 1$ terms in the expansion, with λ_z is represented by f_{10}, λ_x by f_{11} and λ_y by f_{1-1}. Thus, the λE model predicts symmetric polarizations. In particular, inverting the direction of $\mathbf{E}$ inverts the polarization pattern. This translates to polarizations equal in magnitude but opposite in sign in anodal and cathodal conditions, as shown in Figure 1.

2.3. Cell models

We use the multi-compartment conductance-based models from the cortical microcircuit reconstruction by the Blue Brain Project [28]. There, model morphologies are based on reconstructions from real acquisitions from juvenile rats. This work, considers all the morphology types in the database except the layer I interneurons. That includes nine types of interneurons: bipolar, bitufted, chandelier, double bouquet, Martinotti, neurogliaform, and basket cells (including large, small, and nest). These nine types make 36 morphology types, one of each for each layer (II/III, IV, V, and VI). Regarding excitatory cells, there are 13 morphology types (12 types of pyramidal cells and spiny stellate cells). All morphologies are made of connected cables, as required by the NEURON simulation environment (v8.0.0). Each cable is discretized in evenly spaced compartments. All cables were assigned a minimum of one compartment and, two compartments were added every 40 µm of length — as in the original work. The biophysical membrane models contain 13 different known Hodgkin-Huxley type ion channel models distributed using different criteria along the morphology. The Blue Brain Project team defined 11 ion channel profiles or *electrical types* exhibiting different spiking features. The options unfold into ten inhibitory and one excitatory electric profile, where the latter is the one representing all excitatory cell models of the dataset (continuous adapting, cAD). Regarding inhibitory cells, they included bursting (b), continuous (c), and delayed (d) models, later subdivided into accommodating (AC), non-accommodating (NAC), stuttering (STUT), and irregular (IR). Irregular spiking models contain stochastic potassium channels that introduce stochasticity into their behavior. For this study, we suppressed these because of incompatibility with our framework. Along the same lines, we discarded electric types generating spontaneous spiking at rest: cNAC, cSTUT, dSTUT, and dNAC. Given that the same morphology types were found to show different electrical types, they created models for each morphoelectrical combination. Then, the model dataset has five statistical clones of each morphoelectrical type.

2.3.1. Uniform electric field coupling The coupling of the external electric field to the neuron models is implemented via the extracellular potential in NEURON 8.0.0. We use the `extracellular` mechanism to access the extracellular potential in each compartment. To attach the value induced by the external electric field, we tweak the `xtra` mechanism for extracellular recording and stimulation based on the approach by McIntyre and Grill [29]. Recall that the extracellular potential is related to the external uniform field by $\mathbf{E} = -\boldsymbol{\nabla} V_{\text{ext}}$, with

$$V_{\text{ext}}(x, y, z) = -E\big(x \sin\theta \cos\varphi + y \sin\theta \sin\varphi + z \cos\theta\big) \tag{7}$$

Here (x, y, z) refers to the spatial coordinates in $\mathbb{R}^3$ of the center of a compartment of the neuron model.

2.4. Response function estimation

Let $\hat{\Phi}(\theta, \varphi)$ be the discretized estimation of the response function. $\hat{\Phi}(\theta, \varphi)$ can be defined as a collection of changes δv observed for different orientations, where the particular orientation set is determined by the sampling method and the chosen resolution. We obtain $\hat{\Phi}(\theta, \varphi)$ performing simulations applying electric fields of 1 V/m of magnitude in different orientations to a resting neuron model using the NEURON simulation environment (v8.0.0) [30]. All experiments are performed at temperatures of $T = 36.9°$C to mimic the average physiological temperatures of living human brains [31].

2.4.1. Temporal design of the simulation experiment
The simulation pipeline to estimate the steady-state response of any neuron model that has a stable equilibrium under an electric field consists of two phases: (1) computing the resting state and (2) applying an electric field to estimate the change in the induced change in the steady-state condition. The resting state is defined as the steady-state solution of the system when no external stimulation is present. It is assessed by letting the system converge until it experiences no further change in its state variables. Computationally this requirement is met when the membrane potential varies less than 1 nV anywhere in the cell without further external inputs. The second phase assesses the steady-state solution's displacement under an external perturbation; see Figure 1.D. The system is initialized in the pre-computed resting state. Then, the shift in the steady-state solution is retrieved as the change experienced after 500 ms of 1 V/m uniform electric field application in a given condition. We chose 500 ms as a relatively large fixed simulation time at which we can assume the system has reached a stable solution. The induced change, i.e., δv, is computed with respect to the control condition with no field applied. All simulations are integrated using the fully implicit backward Euler method with an integration time step of 25 µs.

2.4.2. Experimental conditions
To characterize the response function, we assess the induced changes for different orientations. We use the Type I Driscoll-Healy sampling of the sphere (DH1), using N=12 to reconstruct harmonics until a maximum degree of $\ell_{\max} = 5$. The sampling theorem gives the relationship among both: $\ell_{\max} = N/2 - 1 = 5$. The DH1 sampling consists of sampling both θ and φ in N points each, building an $N \times N$ grid covering the sphere [32]. Because the full span of both angular dimensions is used, the polar resolution ($\Delta\theta$) is half the azimuthal one ($\Delta\varphi$). For $N = 12$, the angular distance among consecutive sampled points is $\Delta\varphi = 2\pi/13$ and $\Delta\theta = \pi/13$ rad. The discretization is shown in Figure 1.B.

2.4.3. Spherical harmonics-based representation
Following the expansion introduced in Equation 2, once $\hat{\Phi}(\theta, \varphi)$ has been estimated for each compartment, it is represented by a set of spherical harmonics coefficients. The calculation is performed with the `pyshtools` Python package, using a 4π-normalization and the Condon-Shortley phase

Figure 1. Experimental design for estimation of the response function.
A. Temporal profile of the simulation of different experimental conditions. The first part of the experiment computes the resting condition of the model and is common for all experimental conditions. The latter consists in applying a constant external electric field and is repeated of all the desired orientations of the electric field.
B. Experimental conditions. Anodal (red), cathodal (blue) and control (black) conditions in the context of the DH1 Sampling of the sphere. The grid represents the $N \times N$ sampling of the sphere with $N = 12$, with points separated by $\Delta\theta = \pi/13$ and $\Delta\varphi = 2\pi/13$ [32]. **C. Coordinate system** definition with respect to the cell. The vertical direction represents the component normal to the cortical surface and corresponds to the z Cartesian coordinate. The spherical coordinates represent with θ the inclination and with φ the azimuth. The radius is the magnitude of the field E. For all cell models, the origin of coordinates O is set at the center of the soma. **D. Stimulation as an attractor shift.** A constant applied field changes the steady-state condition. In the soma of a prototypical pyramidal cell, the state is shifted towards a depolarized membrane potential.

factor convention [33]. The coefficients $f_{\ell m}$ fully represent the response function $\Phi(\theta, \varphi)$, which can be assessed by integrating the response function over the sphere. Using the 4π normalization, it reads as a double integral over the sphere for a continuous $\Phi(\theta, \varphi)$. In our case, it translates to a discrete sum, as in Equation 8.

$$f_{\ell m} \approx \frac{1}{4\pi} \sum_{\varphi_j \in \mathcal{D}} \sum_{\theta_k \in \mathcal{D}} \sin\theta_k \, \Delta\varphi \, \Delta\theta \, \hat{\Phi}(\theta_k, \varphi_j) Y_{\ell m}(\theta_k, \varphi_j) \tag{8}$$

where $\mathcal{D}$ is the domain of sampled points. Modern implementations of these expansions use fast Fourier transforms over latitude bands, among other methods, to improve

computational efficiency [34, 33].

2.5. Averaging response functions

A response function can be derived for each compartment of a cell model through the methodology already described above, v. Figure 1. However, in the context of model-driven stimulation, the interest lies in simplifications of the complex morphology. Due to linearity, *lumped* response functions can be obtained by averaging the coefficients. We exploit this to create two models: response functions of (i) cell parts and (ii) cell types. The multi-compartment models here have hundreds of compartments for each cell along their morphology. Each compartment can be tagged to belong to a specific region or part of the cell. We use the model's original classification of axon, soma, and dendrites, with a further distinction of apical and basal dendrites in the case of pyramidal cells. In the case of apical dendrites, we make a further distinction between those found in LI with respect to the rest. To build models of cell types, we use the different statistical clones available in the database belonging to the same cell type. We averaged the models of the different cell parts of the clones of the same type and, this way built a canonical model for that type.

2.6. State-dependence

Response functions are expected to vary depending on the morphology and biophysical models of the cells. The ion channels models are functions of the membrane potential and hence depend on the state of the cell. Thus, we expect that response functions are also state-dependent, and we set out to investigate how it may vary for the soma of the canonical LV thick-tufted pyramidal and the size of these effects. Given our steady-state analysis framework, we reconstruct a response function for different membrane potential values using a current clamp at the same study compartment, i.e., the soma. We apply injected currents from -0.2 nA until the system loses its stable equilibrium and stops converging — i.e., the neuron model starts firing. The membrane potential also affects the time constants of ion channel models, and they are expected to increase when moving closer to spiking regimes. For this reason, we provide the system for longer times in experiment phase 2 (10 s) to assess the steady-state change. Currents are injected prior to the application of the electric field and kept during the whole simulation. In other words, we change the location of the stable equilibrium through an external current and explore the perturbations induced by a 1 V/m electric field in that new vicinity.

3. Results

3.1. The response function at rest is dipolar

Consistent with the analysis of the passive cell (see Appendix B), the response function is dipolar under weak fields at rest. In other words, only $\ell = 1$ terms are nonzero. All coefficients of all cell models analyzed are shown in Figure D1. During stimulation, each cell compartment reaches a new steady-state solution. The maximum observed somatic polarization corresponds to fields parallel to the somatodendritic axis at 0.14 mV per 1 V/m in layer V early-branching thick-tufted pyramidal cells. These values are in line with the literature as with results reported in CA1 pyramidal cells [35] and comparable to the typically assumed value of 0.2 mV per V/m of applied electric field [9].

3.2. Somatic polarizations in excitatory cells

Consistent with previous findings, the most affected cells are pyramidals of subgranular layers (V and VI), with an increased effect on those with an apical tuft. The most considerable effect is observed in the early-branching thick tufted population due to the asymmetry of the processes distribution with respect to the soma. Different responses also accompany the variety of layer VI pyramidal cells. Somatic polarization in the LVI tufted cell models is at 0.10 mV, i.e., 58% larger than in the untufted, at 0.07 mV. Somata of inverted pyramidals show an effect of size comparable to that of tufted cells, i.e., 0.10 mV but with reversed polarity. In supragranular layers, effects on pyramidal cells were minimal and irrelevant on spiny stellate and star pyramidals. All maximum somatic polarizations found in the different averaged pyramidal cell models are compiled in Table 1.

3.3. Somatic polarizations in interneurons

Due to its central position with respect to other cell processes, effects in the soma of interneurons are rather small compared to pyramidal cells. Restricting the electrical type to cAC models, we observe a large variety of responses across morphologies and layers—see Figure 4. In bipolar, bitufted, chandelier, and double bouquet cells, we observe a reversal of the polarity of the effect at the soma between supragranular and infragranular layers. Martinotti cell models show the lowest variability between layers. Basket cells are the group that displays the smallest response at the level of the soma consistently in their different types (large, nest, and small) and layers. On the other hand, the cell type model displaying the largest somatic effect is bipolar cells. In this particular cell morphology, the size of the effect strongly depends on its electrical type (see Figure 5). However, we do not observe such dependence in the other types of interneurons.

Figure 2. Polarization dependence on the electric field orientation. A. Polarization of the membrane under a uniform electric field of 1 V/m. The direction of the external electric field has a direct impact on how the charges are redistributed along the cell morphology. In this example, the effects on a layer V pyramidal cell model are displayed for an applied external electric field of 1 V/m in the direction of the background arrows displayed below each cell. **B. Somatic response function estimate.** Estimated polarization of the soma under a 1 V/m electric field for each orientation of the DH1 sampling as described in Figure 1.B. The 2D projection is used with the Lambert method, where the azimuthal angle matches the horizontal axis. **C. Spherical harmonics spectrum.** Representation of each spherical harmonic coefficient squared. Only the $\ell = 1$ terms are present, i.e., the response is dipolar. The description in spherical harmonics terms displays a strong predominance of the $(\ell = 1, m = 0)$ indicating an alignment with the z-axis (see Figure 1.C) slightly tilted towards the x-axis as determined by the presence of a small $(\ell = 1, m = 1)$ term. **D. Reconstructed response function.** Continuous version of the response function represented by the coefficients. It represents all the information contained in the sampled estimate shown in B. **E. Spherical harmonics.** Polar representation of all spherical harmonics up to $\ell = 3$ in the same layout as in C for an intuitive interpretation of each of the coefficients. The magnitude of the function for each orientation sizes the radius and the color codes the sign, warm and cold being positive and negative respectively.

3.4. Effects on different cell parts can be represented with different response functions

As expected from the pre-existing literature on the field, different parts of pyramidal cells respond differently to the presence of the field. Analyzing the effect locally, i.e., compartment-wise, bends and fiber terminations increase the effect of the field on membrane polarizations. The overall size of the this effect is captured by the influenceability metric as defined in Equation 5. In a pyramidal cell, we observe how

Figure 3. Polarization of soma of pyramidal cells under 1 V/m electric field.
Each row represents a neocortical layer and each column a pyramidal cell type. Each
plot displays the modeled change of the membrane potential (mV) in the soma due
to an external electric field of 1 V/m as a function of the relative orientation of the
electric field. Orientation dependence is represented using a Lambert projection. The
vertical axis codes the polar angle θ which ranges from $\theta = 0$ in the top (indicating
a vertically aligned electric field pointing up) to $\theta = \pi$ (down) in the bottom. The
horizontal axis is the azimuthal angle φ, where the left end is $\varphi = 0$ and the right one
is $\varphi = 2\pi$. "*Early*" and "*late*" in layer V (LV) thick-tufted pyramidal cells refer to
when the apical dendrites branching of the tuft begins. In LVI tufted pyramidal cells,
the layer in parenthesis is the layer in which the dendritic tuft terminates.

the most susceptible areas are the tuft of the apical dendrites and terminations at the
axon and the basal dendrites, see Figure 6.

On the other hand, we provide functions for the major regions of the cell. The
estimated response functions for the different cell parts, obtained through averaging the
responses of the individual compartments belonging to each region, display the classical
response of the apical dendrites with a reversed polarity compared to the one observed
in the soma and the basal dendrites, e.g., hyperpolarized under anodal stimulation,
see Figure 7. The largest response among the explored regions is found in the apical
dendrites contained in layer I. In the example oñf the figure, the polarization in the
direction where maximum effect reaches 0.24 mV per V/m in layer I apical dendrites,
whereas it is only 0.03 mV at the trunk, defined as the compartments from the apical
tree found from layer II down. When averaging the whole of the apical dendritic tree,
the response function is reduced to a maximum of 0.07 mV (per V/m). The soma and
the basal dendrites display a similar pattern, with a maximum polarization of 0.16 in

Figure 4. Polarization of soma of interneurons under 1 V/m electric field.
Simulations of the average membrane potential change (mV) in the soma in continuous
accomodating (cAC) models of interneurons. Rows and columns represent cortical
layers and morphology types respectively. The color range is set to match that of
Figure 3 to ease comparison with pyramidal cells. Bipolar cell models of layers II-III
appear saturated because their polarization values exceed the 0.15 mV. Large basket
cell models for LVI are not available in the original set of models.

the soma and 0.18 mV in the basal dendrites. In the axon, including the descendant
and the collaterals, the average is smaller at 0.04 mV per V/m.

3.5. The direction and magnitude of the effect depend on the cell state

We observe that the response function changes by injecting hyperpolarizing and
depolarizing currents in the soma of the layer V pyramidal cell model. Recall that
the injection starts before applying the electric field, and the system is already in its
corresponding steady state, given the injected current. It keeps displaying a dipolar
behavior for most values, purely captured by the $\ell = 1$ coefficients. Moreover, the most
relevant term in all the states is the dipole in the direction parallel to the normal to the
cortical surface, i.e., aligned with the z-axis ($\ell = 1, m = 0$) as in Figure 1.C. Under a
hyperpolarizing current, we observe a slight increase of the effect saturating at 0.17 mV
per V/m. Under depolarizing currents, we observe two different types of behavior.
For currents up to 0.25 nA (corresponding to a membrane potential of ~-62.50 mV,
10 mV above resting), we observe a decrease in the magnitude of the effect reaching
a minimum of 0.12 mV per V/m in the direction of maximum effect. We observe a
different behavior for currents above 0.25 nA until the system enters a spiking regime
for currents above 0.34 nA, corresponding to potentials over -58.13 mV. In this window,
we observe a rapidly increasing effect and the emergence of a non-dipolar response. The
response becomes asymmetric, i.e., the magnitude of the effect of anodal and cathodal

Figure 5. Effect differs across electrical types of bipolar cells. A. Morphology examples of bipolar cells of different layers. **B.** Somatic responses of the different cell models under anodal stimulation of 1 V/m. Each plot provides the polarization curves as a function of time of the models of bipolar cells of a given layer. The stimulation onset is marked as a small arrow over the horizontal axis. The total stimulation duration is of 500 ms. At the right of each plot, the plot is extended with the distribution of the end-polarization states.

stimulations of equal field magnitude is not the same. Still, fields perpendicular to the axis of the *dipole* have negligible effects. In spherical harmonics terms, we observe it as a pair of $\ell = 0$ and $\ell = 2$ terms that increases with the size of the asymmetry, being largest at the verge of the bifurcation. Right before the bifurcation, we report an effect of anodal stimulation of 0.35 mV, whereas cathodal stimulation shifts the membrane potential by -0.26 mV. The inducible changes thus span a range of 0.61 mV with an asymmetry of 0.09 mV.

Figure 6. Influenceability values of a pyramidal cell. A. Variation of the influenceability measure throughout the morphology of an early-branching LV thick-tufted pyramidal cell. Scale: 250 µm. **B.** Distributions across compartments associated with different cell parts. The distribution of the apical dendrites displays bimodality, distinguishing the difference in values along the apical trunk from the values observed in the apical tuft.

Cortical layer	Cell Morphology type	δV (µV)	δI_m (fA)	Ratio δV	Ratio δI_m
LII/III	Pyramidal	36	26	0.26	0.11
LIV	Pyramidal	52	27	0.38	0.11
LIV	Star pyramidal	19	5	0.14	0.02
LIV	Spiny stellate	25	14	0.18	0.06
LV	Untufted pyramidal	49	28	0.35	0.12
LV	Thick-tufted pyramidal (late)	89	122	0.64	0.52
LV	**Thick-tufted pyramidal (early)**	**138**	**233**	**1.00**	**1.00**
LV	Slender-tufted pyramidal	56	65	0.41	0.28
LVI	Untufted pyramidal	68	21	0.49	0.09
LVI	Tufted pyramidal (LI)	96	36	0.70	0.16
LVI	Tufted pyramidal (LIV)	119	30	0.87	0.13
LVI	Inverted pyramidal	104	27	0.76	0.11
LVI	Bipolar pyramidal	25	9	0.18	0.04

Table 1. Maximum observed somatic polarizations δV and transmembrane currents δI_m rounded to the µV and fA, respectively, and their respective ratio referenced to the largest. The cell type with the largest effect is the LV thick-tufted cell with an early branching of the apical tuft (bold).

Figure 7. Models of different sections of a cell. Average response functions of different cell parts of a layer V (LV) early-branching thick-tufted pyramidal cell model and their 2D spectrums. In the morphology, the different cell parts considered are shown in different shades of gray. Models of apical dendrites are split in compartments belonging to L1, to the rest of layers and to the whole of the apical arborization. Coefficients are reported in Table D1.

Figure 8. State dependence of somatic response to the electric field. **A.** Solid lines display the membrane polarization induced by an electric field of 1 V/m in anodal (red) and cathodal (blue) conditions. The control condition, i.e., zero-field, is displayed in black. The asymmetry between anodal and cathodal responses is displayed as the sum of both polarizations (dashed). The 2D spectrum of the response function is shown for hyperpolarization (**B**), resting (**C**), a depolarized state (**D**) and at the edge of the bifurcation (**E**). **F.** The response function at the edge of the bifurcation associated with E and its decomposition into dipolar (**G**, $\ell = 1$) and non-dipolar terms (**H**, $\ell \neq 1$).

4. Discussion

One of this study's primary goals is to develop a simple and intuitive framework that quantifies the effects of weak fields that can be used to build large-scale models. The values shown here for the juvenile rat reconstructions serve as a proof of concept that can be extrapolated to cell models of other species or even the same ones adapted to match dimensions and other physiological features to that of adult rats or humans [14]. The approach applies to any multi-compartment model with a precise enough spatial resolution in its morphology.

Modeled effects from a dynamical systems perspective Equation 2 quantifies the displacement of the stable equilibrium of a neuron model under a weak electric field. As previously discussed, the model requires this type of solution in the analyzed regime. However, neurons are excitable because they operate close to bifurcations that change the dynamical landscape, generally to give rise to periodic solutions, e.g., sustained firing. For instance, integrator models have a stable node (resting) and a saddle node. Short excitatory inputs push the system closer to the saddle-node from the stable node, and if the system is pushed over the stable manifold of the saddle (threshold), it elicits an action potential on its excursion back to the stable point. A constant electric field works by persistently displacing the stable solution, and that is what the framework proposed quantifies, the shift of the stable node. In the example of an integrator model, this shift can be towards (excitatory) or away from (inhibitory) the spiking threshold. However, if the system is in an oscillatory regime, our formalism enters a problem of definition and cannot be strictly used anymore. A formalism in similar lines should be developed to describe changes in oscillatory solutions.

The leading terms in the spherical harmonic expansion are the dipolar ones. The expansion in spherical harmonics of the angular dependence of the effect provides a general framework containing different elements of interest in its different terms. Relevant to stimulation models based on head models, in which one cannot know the azimuthal orientation of a cell with respect to the electric field induced, it becomes convenient that the expansion already collects in the $m = 0$ terms the behavior that does not depend on the azimuth. In fact, by definition, the $m = 0$ terms alone describe the *average* behavior across all azimuths. Another relevant feature is that the method quantifies the dipolar part of the response in its $\ell = 1$ terms, which we find to be the only present for most cases, in agreement with theoretical arguments (Appendix B).This suggests that after the quantification, the steady-state shift can be represented by a dot product as proposed in the literature under the λ E model framework [27]. Furthermore, in realistic settings where azimuthal information of single neurons is unavailable and for cells near rest, the only relevant term is thus the (1,0).

However, we have observed how dipoles are not the only type of response, and we observe asymmetries among anodal and cathodal responses while keeping a neutral

response to perpendicular fields in cells closer to the threshold. This behavior needs other terms that modify the dipolar ones accordingly. For dipoles aligned with the z-axis (1,0), we observed the emergence of a pair of equal values of (0,0) and (2,0). This pair shifts the values of the 1-degree terms at the poles of the dipole. Equivalent expressions involving other terms can modify dipoles oriented in other directions.

State-dependence In the neighborhood of the resting condition, we observed an increase in the effect size if currents were hyperpolarizing and a decrease if depolarizing, as seen in Figure 8.A. This effect aligns with the fact that the size of the polarization induced by the external electric field depends on the membrane resistance. If the resistance increases, the effect's magnitude should also increase [36]. Nevertheless, there are opposite views in this regard since it has also been pointed out that active networks are preferentially modulated in tDCS [37]. The latter, however, seems to be more cohesive with the increased excitability we observe close to the bifurcation. The window of increased excitability right before the spike is also consistent with the typical effect of tDCS affecting spike times [38]. The emergence of other terms, i.e., asymmetries and increased excitability towards the spiking threshold, cannot be explained just by a passive-like membrane argument and requires a different interpretation. We suggest the effects' origin is different and comes from the asymmetric ion channel distributions and their nonlinear behavior.

Reduced compartment models. The average response of a cell part composed of different compartments can be obtained by averaging the coefficients of all the compartments. In other words, averaging the coefficients provides a model of the average response of selected segments or cell types. This strategy can build an averaged response model of the axon, the apical dendrites, or other regions of interest. Regional averaged response models can be translated into multi-compartment models with just a few compartments representing the local membrane dynamics of a few parts of a cell, like the apical tuft, the soma, or the initial axon segment. Reducing the models here to just a few compartments allows us to use this knowledge in models at higher scales, like models of cortical columns made of multi-compartmental cells. Reducing cell models to just a few compartments significantly reduces the computation cost. The approach can be narrowed down to one-compartment models as long as they are assumed to be modeling a specific part of the cell. Then, it poses a proxy for stimulating commonly used models such as leaky or quadratic integrate and fire models [39].

Translation to mesoscale models. Computational models provide a powerful tool for understanding the pathophysiology underlying neurological disorders such as epileptic activity. In [40] and more recently in [41], neural mass models (NMM) have been used to represent epileptic seizure transitions: changes from interictal to ictal state were achieved by increasing the excitatory synaptic gain at the level of glutamatergic pyramidal cells and varying the inhibitory synaptic gains of GABAergic interneurons. There is growing

interest in using non-invasive brain stimulation techniques as a potential treatment for drug-resistant patients and using computational models to optimize these treatments [42]. However, such models require the representation of the effects of electric fields on neuronal populations.

The described methods are a starting point to derive the effects of weak, uniform electric fields at the population level (cortical column or small cortical patch scale). However, the translation from single cells to the mesoscale is not trivial, given that network effects are arising at the population level [15, 16]. A first approach can be to computationally obtain the response of a population by building models of cortical columns or networks of cell models and then extracting the effects at the population level. One could use a full cortical column model with multi-compartment models such as the one developed by the Blue Brain Project [28] and include the effect of the field at each compartment of each cell. A less computationally expensive approach could be to build a similar cortical column model made of neuron models with reduced numbers of compartments, as introduced in the previous paragraph [43]. An alternative to computationally assessing the population-scale effects is to work in a theoretical framework that connects the microscale to the mesoscale, starting from simplifying assumptions. This kind of theoretical connection has recently grown in interest in the field of NMMs since the work by [21], who derived an exact mean field theory from the quadratic integrate and fire neuron model. In this framework, values for the induced transmembrane currents by weak fields are especially relevant since there is no ambiguity in translating the effects from the microscale to the population level.

5. Conclusions

In this work, we have provided a framework for analysis of the effects of weak, uniform electric fields in neuron models in terms of changes in their steady-state conditions. We quantify the induced perturbation dependence on the orientation using spherical harmonic coefficients. Through this analysis, we have validated numerically the commonly used dipole approximation for the studied models in their resting state (far from the threshold) and weak electric field regime. We have also provided a theoretical justification for this in a generic passive membrane model. We have shown, however, that there are conditions under which this approximation is less accurate—particularly close to the spiking threshold. We have also seen that although pyramidal cells generally display the most substantial effects, some interneurons can also be strongly affected but with a more diverse set of responses. Although these effects' impact on neuron networks is left for future research, our work provides the basis for further computational studies studying the relation of electric field effects at microscale and mesoscales. This can benefit the development of brain models using neural mass frameworks and derived clinical applications relying on model-driven brain stimulation with weak electric fields, such as tES.

Appendix A. Real spherical harmonics

A square-integrable function $f : S^2 \to \mathbb{R}$ can be expanded in terms of the real harmonics $Y_{\ell m} : S^2 \to \mathbb{R}$ like:

$$f(\theta, \varphi) = \sum_{l=0}^{\infty} \sum_{m=-\ell}^{\ell} f_{\ell m} Y_{\ell m}(\theta, \varphi) \tag{A.1}$$

The general definition of each spherical harmonic function $Y_{\ell m}$ can be cumbersome, and its derivation is out of the scope of this text. We summarize how they can be stated for completeness, but it does not serve as an introduction to the field. Here it is worthwhile to recall the notation:

- θ stands for the polar angle or inclination.
- φ for the azimuth or longitude.
- ℓ for the spherical harmonic degree.
- m for the order.

Under this convention, the real spherical harmonics are defined as [33]:

$$Y_{\ell m}(\theta, \varphi) = \begin{cases} \bar{P}_{\ell m}(\cos \theta) \cos m\varphi & \text{if } m \geq 0 \\ \bar{P}_{\ell |m|}(\cos \theta) \sin |m| \varphi & \text{if } m < 0 \end{cases}$$

where $\bar{P}_{\ell m}(\mu)$ are the normalized associated Legendre functions. For 4π-normalized spherical harmonic functions, they can be written as:

$$\bar{P}_{\ell m}(\mu) = \sqrt{(2 - \delta_{m0})(2\ell + 1)\frac{(\ell - m)!}{(\ell + m)!}} P_{\ell m}(\mu)$$

Note that δ_{ij} refers to the Kronecker delta and $P_{\ell m}(\mu)$ refers to the unnormalized associated Legendre functions. They can be derived from the standard Legendre polynomials using the relations shown below:

$$P_{\ell m}(\mu) = \left(1 - \mu^2\right)^{m/2} \frac{d^m}{d\mu^m} P_\ell(\mu)$$

$$P_\ell(\mu) = \frac{1}{2^l \ell!} \frac{d^\ell}{d\mu^\ell} \left(\mu^2 - 1\right)^\ell$$

As a set of basis functions, they need to be orthogonal. Note that the associated Legendre functions are indeed orthogonal for any given m, given that:

$$\int_{-1}^{1} \bar{P}_{\ell m}(\mu) P_{\ell m}(\mu) = 2 \left(2 - \delta_{0m}\right) \delta_{\ell\ell}$$

Appendix B. Dipole nature of electric field effect in compartment model of a neuron in the passive, linear membrane regime

Here we show that for a passive membrane, the effect of an electric field on a neuron compartment, or its average over a set of compartments (uniform field), can be written in the form $\delta V = \mathbf{P} \cdot \mathbf{E}$, where $\mathbf{P}$ is a vector. Note that the behavior far from the threshold and under weak electric fields is considered comparable to that of a passive membrane since the active properties of membranes (nonlinear) require more significant effects to come into play. This is the so-called "λE" model used in transcranial electrical stimulation models [27, 44].

The effect of an external field can be represented in a neuron compartment model by an axial current that results from the potential difference induced by the field along the fiber associated with a compartment

$$I = g^a \int_x^{x+\Delta x} E_x(s)\, ds \tag{B.1}$$

where g^a is the axial conductivity and E_x is the field along the fiber. We can rewrite this as $I = g^a \int_f \mathbf{E} \cdot d\mathbf{l}$ with the line integral along the fiber f compartment. If the field is constant along the fiber we can express this simply as $I = g^a\, \mathbf{E} \cdot \mathbf{u}$. The vector $\mathbf{u}$ is a vector defined from one center of compartment to another center of a connected compartment, and pointing into the compartment of interest (see Figure B1). If the E field is aligned with this direction, we get a positive current into the compartment. If the compartment of interest is k and the connected compartment is j, we refer to the vector $\mathbf{u}_{k,j}$ parallel to $\mathbf{x}_k - \mathbf{x}_j$ (the coordinates of the compartments).

Figure B1. Compartments and vectors for cable equation in E field. Here we represent a fiber by a node and a set of directed edges.

From conservation of charge (the sum over j is over connected compartments, and i over membrane currents), and assuming that the field is uniform and thus independent of the compartment k, we obtain

$$C_k \frac{dV_k}{dt} + \sum_i I_{i,k} = \sum_{j \leftrightsquigarrow k} g_j^a\, (V_j - V_k) + \sum_{j \leftrightsquigarrow k} g_j^a\, \mathbf{E} \cdot \mathbf{u}_{k,j} \tag{B.2}$$

where the sum on the left hand side is over ionic membrane currents out of the compartment. The notation $j \leftrightsquigarrow k$ indicates a sum over all compartments j connected

to the kth one. For an arbitrary compartment-dependent quantity Q_j, sums over $j \leftrightsquigarrow k$ can be rewritten via a compartment connectivity matrix c_{kj}, with value of 1 if compartment j is connected to k, zero otherwise. That is, $\sum_{j \leftrightsquigarrow k} Q_j \equiv \sum_j c_{kj} Q_j$.

We can simplify this and also pull the constant electric field out the sum,

$$C_k \frac{dV_k}{dt} + I_k = \sum_{j \leftrightsquigarrow k} g_j^a (V_j - V_k) + \mathbf{p}_k \cdot \mathbf{E} \tag{B.3}$$

where $I_k = \sum_i I_{i,k}$ and where

$$\mathbf{p}_k = \sum_{j \leftrightsquigarrow k} g_j^a \mathbf{u}_{k,j} = \sum_j c_{kj} g_j^a \mathbf{u}_{k,j} \tag{B.4}$$

The equation for a single compartment in steady state is thus of the form

$$C_k \frac{dV_k}{dt} - \mathbf{p}_k \cdot \mathbf{E} + g_k^r V_k + I_k^0 + \sum_{j \leftrightsquigarrow k} g_j^a (V_k - V_j) = 0 \tag{B.5}$$

where $\mathbf{p}_k$ is the dipole vector corresponding to the kth compartment, and $I_k = V_k g_k^r + I_k^0$, with I_k^0 the sum of currents associated to the reversal potentials, and g_k^r the radial (transmembrane) conductivity in the linear regime.

Thus, we see that the behavior of each compartment is characterized by its own dipole and the interaction with other compartments.

Equation B.5 is linear and can be expressed in matrix form, with the compartment array notation $\mathbf{V} = (V_k) = (V_1, V_2, \ldots, V_N)^T$, $C = \mathrm{diag}(C_k) = \mathrm{diag}(C_1, C_2, \ldots, C_N)$, and $P = (\mathbf{p}_1, \ldots, \mathbf{p}_N)^T$.

$$C \frac{\partial \mathbf{V}}{\partial t} + A\mathbf{V} = \mathbf{I^0} - P \cdot \mathbf{E} \tag{B.6}$$

and with the matrix A given by

$$A_{kj} = \delta_{kj} g_j^r - g_j^a c_{k,j} + \delta_{kj} \sum_l g_l^a c_{l,j} = \delta_{kj} \left(g_j^r + \sum_l g_l^a c_{l,j} \right) - g_j^a c_{k,j} \tag{B.7}$$

where we split the diagonal and off diagonal elements in the second equality.

We recall here that a matrix is strictly diagonally dominant if, for every row of the matrix, the magnitude of the diagonal entry in a row is larger than or equal to the sum of the magnitudes of all the other (non-diagonal) entries in that row, i.e.,

$$|a_{ii}| > \sum_{j \neq i} |a_{ii}|$$

Now, the Gershgorin circle theorem implies that a strictly diagonally dominant matrix (or an irreducibly diagonally dominant matrix) is non-singular [45] [Th 6.2.27]. We can observe that this is the case for A, since (all g's are taken to be positive, and recall the c's are 1s or 0s, with $c_{ii} = 0$)

$$|A_{ii}| = g_i^r + \sum_l g_l^a c_{l,i}$$

and

$$\sum_{j\neq i}|A_{ij}| = \sum_{j\neq i} g_j^a c_{i,j} < |A_{ii}|$$

This means that the matrix A has an inverse.

In steady state ($\dot{\mathbf{V}} = 0$), the solution to this equation is

$$\mathbf{V} = A^{-1}[\mathbf{I^0} - P \cdot \mathbf{E}] \tag{B.8}$$

or

$$V_k = \sum_j A_{kj}^{-1}[I_k^0 - \mathbf{p}_k \cdot \mathbf{E}] = \sum_j A_{kj}^{-1} - \mathbf{E} \cdot \sum_j A_{kj}^{-1} \mathbf{p}_j \tag{B.9}$$

where the second equality is a result of linearity of the A matrix, i.e., the fact that we are dealing with a linear equation. Finally, we can write the solution as

$$V_k = \sum_j A_{kj}^{-1} I_j^0 + \tilde{\mathbf{p}}_\mathbf{k} \cdot \mathbf{E} \tag{B.10}$$

This equation shows that the response of the compartment to an electric field is of dipole form, with the total dipole $\tilde{\mathbf{p}}_\mathbf{k}$ resulting from a superposition of dipoles,

$$\tilde{\mathbf{p}}_\mathbf{k} = -\sum_j A_{kj}^{-1} \mathbf{p}_j = -\sum_{j,l} A_{kj}^{-1} c_{jl} g_j^a \mathbf{u}_{j,l} \tag{B.11}$$

Furthermore, we observe that any linear function of compartment potentials will also display, by further superposition, a dipole response. This includes, for example, the average membrane perturbation for apical dendrite compartments, or soma.

What happens in the **nonlinear case**, when the neuron is not at baseline? We can write (steady state)

$$-\mathbf{p}_k \cdot \mathbf{E} + I_k^r(V_k) + \sum_{j\rightsquigarrow k} g_j^a(V_k - V_j) = 0 \tag{B.12}$$

with $I_k^r(V_k)$ a nonlinear function, or, in a fashion analogous to the discussion above, in matrix form

$$-\mathbf{p}_k \cdot \mathbf{E} + B\mathbf{V} + I(\mathbf{V}) = 0$$

or

$$F_k(V) = -\mathbf{p}_\mathbf{k} \cdot \mathbf{E} \tag{B.13}$$

for a nonlinear operator F. This means we cannot pull out the E field out of the inverse. However, if the electric field is very weak, we can carry out the analysis above with the linearized version of Equation B.13 by expanding it around the potential with zero field, with the same conclusion.

Appendix C. Realistic montage exploration

Context   Based on the λE approach, one logical strategy is to optimize for the normal component of the electric field with respect to the cortical surface. The quantification of the performance of the montage is carried out by averaging the electric field on the target area. Expressions like the λE model, which captures the main response of neurons to the fields can be used to develop metrics that are more representative of the relevant physiological effects of the stimulation. As an example of this, we performed a montage optimization to stimulate the dorsolateral prefrontal cortex (dlPFC), which is a common area for stimulation in tDCS, in a real head model - see Figure C1.A.

Montage creation   We ran the Stimweaver algorithm [24] with a maximum of 8 channels, and a total injected current of 4.0 mA, under the restriction of 2.0 mA maximum per channel. The montage was optimized to achieve a normal component to the cortical surface of the electric field of 0.25 V/m in the dlPFC, using weights of 10 and 2 to on- and off-target nodes, respectively. We computed the predicted average membrane polarization of LV pyramidal cells somata on- and off-target as a proof of concept (Figure C1).

Figure C1. Predicted effects in a realistic montage Representation of a pipeline from a target definition to a predicted physiological effect. **A: Target definition.** Overlay in red of the cortical area marked as the target of the stimulation. In gray, the rest of the cortex. **B: Optimized montage.** Anodal (blue) and cathodal (red) electrodes location over the cortical surface. The predicted electric field magnitude is displayed color coded over the cortical surface, being the color code the one displayed in the colorbar on the left (V/m). **C: Predicted polarization.** Prediction of the somatic polarization of LV thick-tufted pyramidal cells. **D: Quantification of effects on and off target.** Predicted values of steady-state somatic polarization induced in a LV thick-tufted pyramidal cell using the optimized montage shown in B. The figure shows how effects are induced in the cortex selectively to the original target.

Results   The target map of the montage is shown in Figure C1.A. The optimized montage is shown in Figure C1.B, together with the predicted electric fields on the cortex—overlaid on the brain morphology. The normal component on-target predicted with the optimized montage has a median value across nodes of 0.11 V/m and a mode of 0.17 V/m computed as the maximum of the kernel density estimate. Through the formalism in Equation 2 and the computationally derived response functions, that map can be translated to a variable with actual physiological meaning, e.g., the somatic membrane potential perturbation of LV early branching thick-tufted pyramidal cells, Figure C1.C. We observe a significant effect on-target (Mann-Whitney U, p< 1×10^{-3}, see Figure C1.D).

Appendix D. Supplementary tables and figures

Figure D1. Soma coefficients for all cell models at rest. *A*. Distributions of soma coefficients for excitatory and inhibitory cell models. The central box plot (in black) displays the overall distribution while the split distributions between excitatory and inhibitory populations are at the sides. Dashed lines separate different degrees (ℓ) of the coefficients. *B,C*. Values of the soma coefficients of all excitatory (B) and inhibitory (C) cell models. Dashed lines separate degrees as in A. Models are also split according to the layer where the cell model belongs.

Table D1. Spherical harmonic coefficients of the response observed in different cell parts of a layer V thick-tufted pyramidal reconstructed cell model.

	Apical	Axon	Basal	Soma
(0, 0)	0.000269004	-0.000260209	-7.28E-05	-8.07E-05
(1, -1)	-0.003474872	0.006142342	-0.000209493	0.002328097
(1, 0)	0.041506581	-0.025128212	-0.103012672	-0.090822117
(1, 1)	-0.002554922	0.002155694	0.002506737	-0.010029683
(2, -2)	8.11E-07	-1.26E-05	-2.85E-07	-2.22E-07
(2, -1)	6.38E-07	2.56E-06	-2.28E-06	-2.88E-06
(2, 0)	0.000201417	-0.000156987	-8.87E-05	-9.50E-05
(2, 1)	8.95E-06	8.25E-06	-6.00E-06	-5.83E-06
(2, 2)	-9.89E-06	1.65E-05	-5.27E-06	-5.19E-06
(3, -3)	-6.71E-09	-6.81E-09	-5.59E-09	-5.32E-09
(3, -2)	-6.96E-09	-7.92E-09	-1.56E-08	-1.54E-08
(3, -1)	1.56E-08	6.54E-08	4.13E-08	3.78E-08
(3, 0)	7.77E-07	-1.84E-08	5.53E-07	5.50E-07
(3, 1)	8.62E-08	3.90E-08	4.62E-08	4.91E-08
(3, 2)	-3.19E-08	-4.24E-09	-3.13E-08	-3.02E-08
(3, 3)	-3.07E-09	-4.22E-09	-3.02E-09	-2.75E-09
(4, -4)	-9.39E-12	-4.23E-11	-3.46E-13	-4.16E-13
(4, -3)	-8.21E-11	-3.38E-11	-4.45E-11	-4.37E-11
(4, -2)	-1.71E-10	-8.52E-11	-9.84E-11	-9.90E-11
(4, -1)	1.71E-10	5.51E-10	6.73E-10	6.73E-10
(4, 0)	5.44E-10	3.64E-09	5.16E-09	5.28E-09
(4, 1)	1.19E-10	1.48E-10	1.82E-10	1.93E-10
(4, 2)	2.91E-10	4.35E-11	-1.95E-11	-1.11E-11
(4, 3)	6.25E-11	1.46E-11	2.51E-13	2.41E-12
(4, 4)	3.27E-13	-6.69E-12	4.28E-12	3.95E-12
(5, -5)	9.75E-15	4.67E-14	1.11E-14	1.19E-14
(5, -4)	-2.80E-15	-5.73E-14	-1.85E-14	-1.80E-14
(5, -3)	-3.45E-13	-3.11E-13	-1.60E-13	-1.62E-13
(5, -2)	-1.51E-12	-5.01E-14	-2.66E-13	-2.80E-13
(5, -1)	-1.76E-12	-1.18E-12	-1.26E-12	-1.28E-12
(5, 0)	-4.01E-11	2.43E-12	-1.90E-11	-1.92E-11
(5, 1)	-4.20E-12	-8.62E-14	-1.30E-12	-1.33E-12
(5, 2)	5.01E-12	1.03E-12	1.57E-12	1.63E-12
(5, 3)	7.95E-13	-7.65E-14	1.57E-13	1.69E-13
(5, 4)	-4.13E-14	-1.27E-14	-5.17E-15	-6.70E-15
(5, 5)	-1.08E-14	-6.81E-14	-6.35E-15	-5.73E-15

Acknowledgments

This work has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 855109; ERC-SyG 2019 Galvani) and from FET under the European Union's Horizon 2020 research and innovation programme (grant agreement No 101017716, Neurotwin).

References

[1] M. A. Nitsche and W. Paulus. Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *J Physiol*, 527 Pt 3:633–9, 2000.

[2] Hebb, d. o. the organization of behavior: A neuropsychological theory. *Science Education*, 34(5):336–337, December 1949.

[3] D. Liebetanz, F. Fregni, K. K. Monte-Silva, M. B. Oliveira, A. Amancio-dos Santos, M. A. Nitsche, and R. C. Guedes. After-effects of transcranial direct current stimulation (tDCS) on cortical spreading depression. *Neurosci Lett*, 398(1-2):85–90, 2006.

[4] D. Liebetanz, M. A. Nitsche, F. Tergau, and W. Paulus. Pharmacological approach to the mechanisms of transcranial DC-stimulation-induced after-effects of human motor cortex excitability. *Brain*, 125(2238-2247), 2002.

[5] Giulio Ruffini, Fabrice Wendling, Isabelle Merlet, Behnam Molaee-Ardekani, Abeye Mekonnen, Ricardo Salvador, Aureli Soria-Frisch, Carles Grau, Stephen Dunne, and Pedro C. Miranda. Transcranial current brain stimulation (tCS): Models and technologies. *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, 21(3):333–345, May 2013.

[6] Pedro Cavaleiro Miranda, Abeye Mekonnen, Ricardo Salvador, and Giulio Ruffini. The electric field in the cortex during transcranial current stimulation. *Neuroimage*, 70:45–58, 2013.

[7] P. C. Miranda, M. Hallett, and P. J. Basser. The electric field induced in the brain by magnetic stimulation: a 3-D finite-element analysis of the effect of tissue heterogeneity and anisotropy. *IEEE Trans Biomed Eng*, 50(9):1074–85, 2003.

[8] Frank Rattay. Analysis of models for external stimulation of axons. *IEEE Transactions on Biomedical Engineering*, 33(10):974–977, 1986.

[9] Thomas Radman, Raddy L. Ramos, Joshua C. Brumberg, and Marom Bikson. Role of cortical cell type and morphology in subthreshold and suprathreshold uniform electric field stimulation in vitro. *Brain Stimulation*, 2(215):28, 2009.

[10] W. Rall. Core conductor theory and cable properties of neurons. pages 39–97. American Physiological Society, 1977.

[11] D. Tranchina and C. Nicholson. A model for the polarization of neurons by extrinsically applied electric fields. *Biophys J*, 50(6):1139–56, 1986.

[12] Frank Rattay. Analysis of models for extracellular fiber stimulation. *IEEE Transactions on Biomedical Engineering*, 36(7):676–682, 1989.

[13] B. J. Roth. Mechanisms for electrical stimulation of excitable tissue. *Crit Rev Biomed Eng*, 22(3-4):253–305, 1994.

[14] Aman S Aberra, Angel V Peterchev, and Warren M Grill. Biophysically realistic neuron models for simulation of cortical stimulation. *Journal of Neural Engineering*, 15(6):066023, October 2018.

[15] F Fröhlich and D. A. McCormick. Endogenous Electric Fields May Guide Neocortical Network Activity. *Neuron*, 67:129–143, 2010.

[16] D. Reato, A. Rahman, M. Bikson, and L. C. Parra. Low-intensity electrical stimulation affects network dynamics by modulating population rate and spike timing. *Journal of Neuroscience*, 30(45):15067–15079, November 2010.

[17] Giulio Ruffini, Ricardo Salvador, Ehsan Tadayon, Roser Sanchez-Todo, Alvaro Pascual-Leone, and

Emiliano Santarnecchi. Realistic modeling of mesoscopic ephaptic coupling in the human brain. *PLoS Comput. Biol.*, 2020.

[18] B. H. Jansen, G. Zouridakis, and M. E. Brandt. A neurophysiologically-based mathematical model of flash visual evoked potentials. *Biol Cybern*, 68(3):275–83, 1993.

[19] B. H. Jansen and V. G. Rit. Electroencephalogram and visual evoked potential generation in a mathematical model of coupled cortical columns. *Biol Cybern*, 73(4):357–66, 1995.

[20] F. Grimbert and Olivier Faugeras. Analysis of Jansen's model of a single cortical column. *INRIA*, RR-5597:34, June 2006.

[21] Ernest Montbrió, Diego Pazó, and Alex Roxin. Macroscopic description for networks of spiking neurons. *Phys. Rev. X*, 021028, 2015.

[22] Federico Devalle, Alex Roxin, and Ernest Montbrió. Firing rate equations require a spike synchrony mechanism to correctly describe fast oscillations in inhibitory networks. *PLOS Computational Biology*, 13(12), 2017.

[23] Giulio Ruffini. Analysis and extension of exact mean-field theory with dynamic synaptic currents. *bioRxiv*, (https://www.biorxiv.org/content/10.1101/2021.09.01.458563v6), 2021.

[24] Giulio Ruffini, Michael D. Fox, Oscar Ripolles, Pedro Cavaleiro Miranda, and Alvaro Pascual-Leone. Optimization of multifocal transcranial current stimulation for weighted cortical pattern targeting from realistic modeling of electric fields. *NeuroImage*, 89:216 – 225, 2014.

[25] Pierre Simon Laplace, Nathaniel Bowditch, and N. I Bowditch. *Mécanique céleste*. Hillard, Gray, Little, and Wilkins, 1829.

[26] W.T.B. Kelvin and P.G. Tait. *Treatise on Natural Philosophy*. Number v. 1 in Clarendon Press series. Clarendon Press, 1867.

[27] Giulio Ruffini, Fabrice Wendling, Isabelle Merlet, Behnam Molaee-Ardekani, Abeye Mekkonen, Ricardo Salvador, Aureli Soria-Frisch, Carles Grau, Stephen Dunne, and Pedro Miranda. Transcranial Current Brain Stimulation (tCS): Models and Technologies. *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, 21(3):333–345, May 2013.

[28] Henry Markram, Eilif Muller, Srikanth Ramaswamy, Michael W. Reimann, Marwan Abdellah, Carlos Aguado Sanchez, Anastasia Ailamaki, Lidia Alonso-Nanclares, Nicolas Antille, Selim Arsever, Guy Antoine Atenekeng Kahou, Thomas K. Berger, Ahmet Bilgili, Nenad Buncic, Athanassia Chalimourda, Giuseppe Chindemi, Jean-Denis Courcol, Fabien Delalondre, Vincent Delattre, Shaul Druckmann, Raphael Dumusc, James Dynes, Stefan Eilemann, Eyal Gal, Michael Emiel Gevaert, Jean-Pierre Ghobril, Albert Gidon, Joe W. Graham, Anirudh Gupta, Valentin Haenel, Etay Hay, Thomas Heinis, Juan B. Hernando, Michael Hines, Lida Kanari, Daniel Keller, John Kenyon, Georges Khazen, Yihwa Kim, James G. King, Zoltan Kisvarday, Pramod Kumbhar, Sébastien Lasserre, Jean-Vincent Le Bé, Bruno R.C. Magalhães, Angel Merchán-Pérez, Julie Meystre, Benjamin Roy Morrice, Jeffrey Muller, Alberto Muñoz-Céspedes, Shruti Muralidhar, Keerthan Muthurasa, Daniel Nachbaur, Taylor H. Newton, Max Nolte, Aleksandr Ovcharenko, Juan Palacios, Luis Pastor, Rodrigo Perin, Rajnish Ranjan, Imad Riachi, José-Rodrigo Rodríguez, Juan Luis Riquelme, Christian Rössert, Konstantinos Sfyrakis, Ying Shi, Julian C. Shillcock, Gilad Silberberg, Ricardo Silva, Farhan Tauheed, Martin Telefont, Maria Toledo-Rodriguez, Thomas Tränkler, Werner Van Geit, Jafet Villafranca Díaz, Richard Walker, Yun Wang, Stefano M. Zaninetta, Javier DeFelipe, Sean L. Hill, Idan Segev, and Felix Schürmann. Reconstruction and simulation of neocortical microcircuitry. *Cell*, 163(2):456–492, October 2015.

[29] Cameron C. McIntyre and Warren M. Grill. Extracellular stimulation of central neurons: Influence of stimulus waveform and frequency on neuronal output. *Journal of Neurophysiology*, 88(4):1592–1604, October 2002.

[30] M. L. Hines and N. T. Carnevale. NEURON: a tool for neuroscientists. *Neuroscientist*, 7(2):123–35, 2001.

[31] Huan Wang, Bonnie Wang, Kieran P. Normoyle, Kevin Jackson, Kevin Spitler, Matthew F. Sharrock, Claire M. Miller, Catherine Best, Daniel Llano, and Rose Du. Brain temperature

and its fundamental properties: a review for clinical neuroscientists. *Frontiers in Neuroscience*, 8, October 2014.

[32] J.R. Driscoll and D.M. Healy. Computing fourier transforms and convolutions on the 2-sphere. *Advances in Applied Mathematics*, 15(2):202–250, June 1994.

[33] Mark A. Wieczorek and Matthias Meschede. SHTools: Tools for working with spherical harmonics. *Geochemistry, Geophysics, Geosystems*, 19(8):2574–2592, August 2018.

[34] J. A. Rod Blais. Discrete spherical harmonic transforms: Numerical preconditioning and optimization. In *Computational Science – ICCS 2008*, pages 638–645. Springer Berlin Heidelberg, 2008.

[35] M. Bikson, M. Inoue, H. Akiyama, J. K. Deans, J. E. Fox, H. Miyakawa, and J. G. Jefferys. Effects of uniform extracellular DC electric fields on excitability in rat hippocampal slices in vitro. *J Physiol*, 557(Pt 1):175–90, 2004.

[36] Walter Paulus and John C. Rothwell. Membrane resistance and shunting inhibition: where biophysics meets state-dependent human neurophysiology. *The Journal of Physiology*, 594(10):2719–2728, May 2016.

[37] Marom Bikson and Asif Rahman. Origins of specificity during tDCS: anatomical, activity-selective, and input-bias mechanisms. *Frontiers in Human Neuroscience*, 7, 2013.

[38] T. Radman, Y. Su, J. H. An, L. C. Parra, and M. Bikson. Spike timing amplifies the effect of electric fields on neurons: implications for endogenous field effects. *J Neurosci*, 27(11):3030–6, 2007.

[39] Peter Dayan and L F Abbott. *Theoretical neuroscience*. Computational neuroscience. MIT Press, London, England, December 2001.

[40] F. Wendling, F. Bartolomei, J. J. Bellanger, and P. Chauvel. Epileptic fast activity can be explained by a model of impaired GABAergic dendritic inhibition. *Eur J Neurosci*, 15(9):1499–508, 2002.

[41] Edmundo Lopez-Sola, Roser Sanchez-Todo, Èlia Lleal, Elif Köksal Ersöz, Maxime Yochum, Julia Makhalova, Borja Mercadal, Maria Guasch, Ricardo Salvador, Diego Lozano-Soldevilla, Julien Modolo, Fabrice Bartolomei, Fabrice Wendling, Pascal Benquet, and Giulio Ruffini. A personalizable autonomous neural mass model of epileptic seizures. *Journal of Neural Engineering*, 2022.

[42] Giulio Ruffini, Fabrice Wendling, Roser Sanchez-Todo, and Emiliano Santarnecchi. Targeting brain networks with multichannel transcranial current stimulation (tCS). *Current Opinion in Biomedical Engineering*, 2018.

[43] Yves Denoyer, Isabelle Merlet, Fabrice Wendling, and Pascal Benquet. Modelling acute and lasting effects of tDCS on epileptic activity. *Journal of Computational Neuroscience*, 48(2):161–176, May 2020.

[44] Giulio Ruffini, Michael D Fox, Oscar Ripolles, Pedro Cavaleiro Miranda, and Alvaro Pascual-Leone. Optimization of multifocal transcranial current stimulation for weighted cortical pattern targeting from realistic modeling of electric fields. *Neuroimage*, 89:216–225, 2014.

[45] Roger A Horn and Charles R Johnson. *Matrix Analysis*. Cambridge University Press, Cambridge, England, February 1990.