Deep mutational scanning and massively parallel kinetics of plasminogen activator-1

functional stability

Laura M. Haynes[1], Zachary M. Huttinger[1,2,#], Andrew Yee[3], Colin A. Kretz[4], David R.

Siemieniak[1,5], Daniel A. Lawrence[6,7], and David Ginsburg[1,2,5,6 8,†]

[1]Life Sciences Institute, University of Michigan, Ann Arbor, MI

[2] Cellular and Molecular Biology Program, University of Michigan Medical School, Ann Arbor, MI

[3]Department of Pediatrics, Baylor College of Medicine, Houston, TX

[4]Department of Medicine, McMaster University and the Thrombosis and Atherosclerosis

Research Institute, Hamilton, ON, Canada

[5]Howard Hughes Medical Institute

[6]Department of Internal Medicine, University of Michigan Medical School, Ann Arbor, MI

[7]Department of Pathology, University of Michigan Medical School, Ann Arbor, MI

[8]Departments of Human Genetics and Pediatrics, University of Michigan, Ann Arbor, MI

[#]Current Address: Department of Otolaryngology, The Ohio State University College of

Medicine, Columbus, OH

[†]Corresponding author: David Ginsburg

E-mail: ginsburg@umich.edu

Running Title: PAI-1 functional stability

Keywords: high-throughput sequencing (HTS), phage display, serpin, enzyme inhibitor,

protease inhibitor, fibrinolysis, kinetics

ABSTRACT

Plasminogen activator inhibitor-1 (PAI-1), a member of the serine protease inhibitor (SERPIN)

superfamily of proteins, is unique among SERPINs for exhibiting a spontaneous conformational

change to a latent or inactive state. The functional half-life for this transition at physiologic

temperature and pH is ~1-2 h. To better understand the molecular mechanisms underlying this

transition, we now report on the analysis of a comprehensive PAI-1 variant library expressed on

filamentous phage and selected for functional stability after 48 h at 37 $^{\circ}$C. Of the 7,201 possible

single amino acid substitutions in PAI-1, we identify 439 that increase the functional stability of

PAI-1 beyond that of the wild-type protein and 1,549 that retain inhibitory activity toward PAI-1's

canonical protease (urokinase-like plasminogen activator, uPA), while exhibiting functional

stability less than or equal to that of wild-type PAI-1. Missense mutations that increase PAI-1

functional stability are concentrated in highly flexible regions within the PAI-1 structure. Finally,

we developed a method for simultaneously measuring the functional half-lives of hundreds of

PAI-1 variants in a multiplexed, massively parallel manner, quantifying the functional half-lives

for 697 single missense variants of PAI-1 by this approach. Overall, these findings provide novel

insight into the mechanisms underlying PAI-1's latency transition, and provide a database for

the interpretation of human PAI-1 genetic variants.

INRODUCTION

Serine protease inhibitors, or SERPINs, comprise a protein superfamily with over 30 members in humans, the majority of which display specific inhibitory activity towards a subset of serine proteases (1,2). Despite their high degree of target specificity, all SERPINs share a common structure that consists of a globular domain containing a central β-sheet (β-sheet A), with a reactive center loop (RCL) that extends above the core of the protein when the SERPIN is in its metastable active state. The RCL contains a consensus sequence that is the primary recognition site for its unique set of target serine protease(s) (3). When the serine protease encounters its target sequence in the RCL, proteolysis between the P1 and P1' positions is initiated by formation of an acyl intermediate. Inhibition of the target serine protease results when the RCL inserts into β-sheet A prior to hydrolysis of the acyl intermediate (4),with the resulting conformational changes rendering both the serine protease and SERPIN no longer functional. If the acyl-intermediate is hydrolyzed prior to insertion of the RCL into β-sheet A, the serine protease will remain active, while the proteolyzed RCL will render the cleaved SERPIN non-inhibitory (1,5).

Plasminogen activator inhibitor-1 (PAI-1, *SERPINE1*) is a 379 amino acid, prototypic member of the SERPIN superfamily composed of nine α-helices and three β-sheets. PAI-1's canonical function is to inhibit the serine proteases urokinase-like and tissue-type plasminogen activator (uPA and tPA, respectfully)—the primary proteolytic activators of the zymogen plasminogen to the serine protease plasmin. Plasmin is the key enzyme responsible for fibrinolysis, the proteolytic degradation of fibrin clots. Although the major function of PAI-1 appears to be in hemostasis through regulation of the fibrinolytic pathway, other functional roles have been proposed (6). *In vivo*, PAI-1 circulates in an 1:1 complex with its cofactor vitronectin (7,8), which also enhances PAI-1's ability to inhibit other proteases in the coagulation pathway including thrombin and activated protein C (9). PAI-1 has also been demonstrated to inhibit other coagulation factors including coagulation factor IXa, coagulation factor XIIa, plasmin, and kallikrein (8,10), as well as non-coagulation proteases including furin and TMPRSS2—both of which have been implicated in the priming of viruses for cellular entry, including influenza A and SARS-CoV-2 (11-15).

Mice deficient in PAI-1 are viable without hemostatic abnormalities (16,17), and PAI-1 deficiency has been reported to be protective in animal models for a number of disease states (18). Complete PAI-1 deficiency in humans results in a mild to moderate bleeding diathesis (19,20),

with heterozygous PAI-1 deficiency linked to increased longevity (21). Elevated levels of

plasma PAI-1 have also been associated with a number of disorders including myocardial

infarction, atherosclerosis, fibrosis, obesity, type 2 diabetes, metabolic syndrome, and cancer

(6,22,23).

PAI-1 is unique among the SERPIN protein superfamily in undergoing a spontaneous transition

to a non-inhibitory but more thermodynamically stable, latent conformation when, in the absence

of a proteolytic event, the RCL spontaneously inserts into β-sheet A (24). The metastable PAI-1

active conformation exists in equilibrium with a prelatent state in which the RCL is hypothesized

to be partially inserted into β-sheet A (25-30). Full insertion of the RCL is conformationally

blocked by the Val[157]-loop between helix F and β-strand 3A; however, partial unfolding of helix F

permits full insertion of the RCL into β-sheet A, while the return of helix F to the native state

traps PAI-1 in its latent conformation (27,28).

The capacity of PAI-1 to undergo this latency transition is conserved across vertebrates (31-33),

with human PAI-1 exhibiting a half-life ($t_{1/2}$) for the active conformation of approximately one-two

hours (30,34-36), at physiologic temperature and pH, although this half-life is prolonged in the

presence of vitronectin (30,37), which stabilizes PAI-1 in its metastable active conformation.

While the physiologic function of the PAI-1 latency transition is unknown, it may serve to clear

plasma PAI-1 activity from circulation that is distal from a site of fibrinolytic activity (38).

A limited number of PAI-1 variants have been engineered with extended half-lives, using both

unbiased and rational design approaches (39-43); We previously reported the use of deep

mutational scanning (DMS) of PAI-1 to characterize the effects of missense mutations on PAI-

1's inhibitory activity toward uPA (44). We now extend this approach to broadly profile the

impact of amino acid substitutions throughout the PAI-1 coding sequence on functional stability.

We also describe a novel approach to simultaneously measure the half-lives of multiple PAI-1

variants in a massively parallel fashion.

RESULTS

Massively parallel determination of PAI-1 variant functional stability

The fraction of active phage displayed PAI-1 (phPAI-1) (44) was determined by

immunoprecipitation of phPAI-1:uPA complexes with an α-uPA antibody at the indicated time

points of incubation at 37°C over a 48 h time course (Fig. 1). Using this approach, the half-life

($t_{1/2}$) of WT phPAI-1 was determined to be 2.1 h, consistent with published values for WT

recombinant PAI-1 (rPAI-1) (30,34,35). In parallel experiments, immunoprecipitation by the N-

terminal *myc*-tag of phPAI-1 further confirmed that the loss of phPAI-1 functional activity is due

to its transition to latency and not degradation or proteolysis (Fig. 1). When the previously

characterized phPAI-1 mutant library (44) was analyzed over the same 48 h time course, the

apparent aggregate half-life was determined to be 11.2 h (Fig. 1). This prolonged half-life is

likely explained by the presence within the library of multiple phage carrying mutations that

prolong the functional stability of PAI-1.

We previously reported construction of a phPAI-1 mutant library containing 8.04 x 10^6

independent clones with an average of 4.2 ± 1.8 missense mutations per fusion protein (44). To

assess the relative functional stability of phPAI-1 variants that retained inhibitory activity at 48 h,

the phPAI-1 mutant library was incubated at 37°C for 48 h, prior to selection for complex

formation with uPA (SI Data 1). Selected and unselected phage were analyzed by high

throughput DNA sequencing. Variants that were non-functional at 0 h (i.e., depleted following

selection for complex formation without heat-induced latency) were removed from subsequent

analyses as these variants were assumed to either possess no inhibitory activity or have an

extremely short functional half-life (44). The relative functional stability scores for missense

mutations compared to the WT amino acid at each position are illustrated in Fig. 2. Of the 1,988

missense mutations that were characterized for functional stability relative to WT PAI-1, 439

amino acid substitutions generated PAI-1 variants with increased functional stability while 1,549

resulted in PAI-1 variants that were functionally active yet less functionally stable than WT PAI-

1. The missense mutations with the top 35 functional stability scores are listed in Table 1.

Although PAI-1 mutations that increase or decrease the rate of uPA inhibition may potentially

contribute to relative functional stability scores, this effect should be negligible given the

concentrations of uPA and phPAI-1, as well as the relatively long reaction time (30 min).

Massively parallel kinetic measurement of PAI-1 functional half-lives

Leveraging the power of next generation sequencing coupled with our phage display system,

we developed an approach to accurately extract latency transition half-lives for individual

missense PAI-1 mutations from the aggregate half-life data for the entire phPAI-1 mutational

library. As described in more detail in *Experimental Procedures*, uPA selection of the library was

performed at multiple time points over a 48 h time course (Fig. 1), subjected to HTS, and these

data used to determine the half-lives of individual missense mutations. By this approach, we

calculated the functional half-lives of 697 individual PAI-1 missense mutations (Fig. 3A, SI Data 2), including 21 of the 35 variants listed in Table 1 (the latter ranging from 2 to 13.1 hours). The mean and median half-lives of the library were determined to be 2.9 and 2.6 h respectively and exhibit strong correlation with functional stability scores (R = 0.6; SI Data 1 and 2). Additionally, excellent correlation (slope = 0.5, R = 0.8; Fig. 3B) was observed for the half-lives calculated using this massively parallel kinetics compared to those determined for individual rPAI-1 variants in this study (Ile91Phe and Lys176Glu, SI Fig 1) or previously reported (Phe36Thr, Met45Lys, Thr49Ala, Thr93Ser, Lys141Gln, Lys154Thr, Glu212Val, Thr232Ser, Arg271Cys, Gln319Leu, Glu327Val, Ser332Gly, Met354Ile, Phe372Ile, and Phe372Leu) (45).

As the phPAI-1 mutational library used in these studies contains multiple amino acid substitutions per molecule, it is possible that epistatic interactions between mutations in cis with each other on different sequencing amplicons (SI Fig. 2) may confound the determination of quantitative half-lives in these studies. However, the excellent agreement between our massively parallel calculations and the values measured for individual variants, suggests that this factor does not significantly confound these data. Similar to the conclusions of Kretz *et al.* in their massively parallel kinetic studies of the rates for ADAMTS13 proteolysis of a phage displayed peptidyl library (46), we posit that any epistatic effects average out over multiple clones generating a neutral background.

DISCUSSION

We report the successful application of DMS to map PAI-1's mutational landscape with respect to functional stability, identifying a large number of variants that prolong PAI-1's lifetime in its metastable active or "stressed" state. Though DMS has been increasingly applied to catalog variants for a large number of proteins (47-49), only a few studies to date have attempted to extract kinetic parameters from such large data sets (46,50,51). The approach described here permits the massively parallel measurement of a kinetic parameter for a large set of amino acid substitutions. A similar approach should be applicable for the measurement of other parameters including inhibitory rate constants. The results of this and similar studies should also aid in the interpretation of variants of unknown significance as clinical use of whole human genome sequencing expands. For example, a PAI-1 variant with a prolonged functional half-life may present with a prothrombotic phenotype, while a PAI-1 variant with a decreased half-life may mimic PAI-1 deficiency (19,20).

Our DMS of PAI-1 functional stability identifies regions in which a large number of amino acid

substitutions that increase PAI-1 functional stability are present (Fig. 2, SI Fig. 3). The two

regions most enriched in variants that prolong PAI-1's functional stability are residues 73-98 and

319-347. When mapped to the crystal structure of PAI-1 (PDB: 3Q02 (52)), Fig. 4) residues 73-

98 correspond to helix D and β-strand 2A, while residues 319-347 correspond to β-strand 5A

and the N-terminus of the RCL (residues 331-350). Other regions enriched in amino acid

substitutions that prolong PAI-1's functional stability include residues 40-49 (helix B), and in the

central portion of PAI-1's primary structure, residues 145-174 (β-strand 2A and the Val^{157}-loop).

Helices D and F, along with the Val^{157}-loop and the loop connecting helix D and β-strand 2A, are

part of one of the most flexible regions of PAI-1, aptly referred to as the "flexible loop region"

(28). This flexibility may make it thermodynamically favorable for the F-loop, along with β-

strands 1A, 2A, and 3A, to "slide" out of the way as part of PAI-1's shutter region, permitting full

RCL insertion into the central β-sheet A from PAI-1's pre-latent state (27,53,54). We speculate

that the missense mutations in this region identified in our screen most likely decrease flexibility

leading to increased residence time of the RCL in the metastable active and pre-latent

conformations.

PAI-1 variants with increased functional stability are also clustered in the "breach region" of

PAI-1 between strands 3 and 5 of β-sheet A into which the RCL must insert to undergo either a

transition to latency or inhibit a target protease. We hypothesize that mutations in this region

may reduce that rate at which the RCL inserts into β-sheet A enough to impede the latency

transition but not enough to inhibit the efficient and effective inhibition of uPA (Fig. 4). In a

similar vein the enrichment of mutations in the N-terminal region of the RCL that prolong PAI-1's

functional stability suggests that latency is driven in part by high-flexibility in this region that

allows the RCL to exist in conformations that permit initial insertion into the breach region (Fig.

2).

Other variants of interest that may prolong functional stability of PAI-1 are clustered in the

regions of helices B (residues 40-49) and H (residues 261-276), as well as strand 2 of both β-

sheets B (residues 217-232) and C (residues 261-276) as illustrated in Figs. 2 and 4. Flexibility

in helix B, has previously been linked with increased probability of a latency transition (28,55).

Furthermore, helix H, along with β-strands B and C, comprise the sB/sC binding pocket of PAI-1

in which ligands can allosterically inhibit vitronectin binding to PAI-1 (56). We postulate that

mutations in this region that extend the functional half-life of PAI-1 may mimic the allosteric

changes in PAI-1 structure that are induced by vitronectin binding, potentially restricting the RCL

from transitioning to a latent state, while not rendering PAI-1 so rigid that it loses its inhibitory

capacity.

Of the 35 most enriched mutations listed in Table 1, two or more different amino acid

substitutions were observed at seven distinct sites (Ile^{91} (n = 3), Leu^{169} (n = 2), Asp^{222} (n = 4),

Tyr^{241} (n = 2), Gly^{332} (n = 3), Thr^{333} (n =2) and Phe^{372} (n = 4)). Among these sites, Gly^{332} and

Thr^{333} along with Ser^{331}, are located at the N-terminal hinge of the RCL. Mutation of Gly^{332} and

Thr^{333} to other small amino acids (Ser, Asp, and Ala for Gly^{332} , as well as Ala and Ser for

Thr^{333}), may not affect RCL insertion into β-sheet A, while moderately affecting the accessible

Ramachandran space, thereby increasing the time scale of RCL insertion—consistent with

limiting the ability of PAI-1 to enter a pre-latent state (27).

Residues Asp^{222} and Tyr^{241} are located on the periphery of β-sheet B. Sui and Wiman have

previously reported that mutations in this region are linked to prolonged functional stability of

PAI-1 (57). Although their mutational analysis suggests that substitution of a basic residue (Lys)

at position 222 is responsible for the change in functional stability, our more comprehensive

mutational mapping demonstrates that simple loss of an acidic amino acid at this position by

mutation to a neutral side chain (His, Tyr, Val, Asn) may be more critical. Sui and Wiman also

reported that Tyr241Ser resulted in a modest increase in PAI-1 functional activity, consistent

with our data for multiple substitutions at this site (Table 1; SI Data 1 and 2).

Three other amino acid positions Ile^{91}, Leu^{169}, and Phe^{372} with multiple enriched mutations are

clustered on strands 2 and 3 of β-sheet A (Table 1). Previously, the Ile91Leu mutation was

reported as the single amino acid substitution that most significantly prolongs the half-life of PAI-

1 (39). The identification of two additional amino acid substitutions at this position that increase

the functional stability of PAI-1 (Ile91Phe and Ile91Met) further highlights the importance of this

position and suggests that the presence of bulkier side chains stabilizes the shutter region in a

closed position. Likewise, two of the four mutations at Phe^{372} (Phe372Ile and Phe372Le) had

been previously identified to stabilize PAI-1 in its metastable state. The additional two variants

identified here (Phe372Val and Phe372Cys) further suggest that substitution of Phe with a

smaller hydrophobic residue promotes PAI-1's functional stability. Finally, though the

importance of Leu[169] in affecting PAI-1's functional stability has not been reported previously, our data identify both Leu169His and Leu169Ser as promoting enhanced stability in PAI-1's active stressed state—perhaps due to increased potential for hydrogen bond formation or hydrophobic interactions, as Ile[91] and His[169] make contact with each other on the interior of β-sheet A (Fig. 4).

The results of our screen further complement the molecular dynamics simulations of Kihn *et al.* (58), highlighting the utility of DMS in interpreting complex protein structure-function relationships. The models of Kihn *et al.* suggest that PAI-1's latency transition is facilitated by a transient salt bridge between Arg[30] and Asp[193] (43). While mutations at Asp[193] are not tolerated (44), substitutions at Arg[30] with Cys, His, Leu, and Ser lead to prolonged PAI-1 functional half-lives (4.3, 2.8, 3.4, and 4.5 h, respectively). Furthermore, given that regions of enriched stabilizing mutations overlap with the regions of PAI-1 perturbed in response to vitronectin binding, we hypothesize that these mutations may result in a rigidification of the gate-region that mimics that of vitronectin binding (58). Using dynamic network analysis, Kihn *et al.* have identified 100 trajectories of allosteric pathways that link vitronectin binding to enhanced functional stability. Residues that are degenerate between pathways are either intolerant of mutations (Thr[144], Tyr[170], and Met[202]) (44) or the majority of tolerated mutations result in a functionally stable PAI-1 (Leu[169], Asn[172], and Asn[329]). These findings are indicative that the former residues are not only critical for the identified allosteric networks but also proper protein folding and function, while the stabilizing mutations introduced in the latter may replicate or enhance side chain interactions that are induced by vitronectin binding.

In conclusion, we have applied DMS to map the mutational landscape of PAI-1 with respect to its latency transition. Overall, we confirm and identify structural features that affect the functional stability of PAI-1 that complement not only prior experimental work but also molecular dynamics simulations of PAI-1 behavior (58) highlighting the utility of high-throughput unbiased screens for protein variant function. We were also able to accurately and quantitatively determine the half-lives of PAI-1 variants across its mutational landscape, allowing biochemically meaningful value to be incorporated into our high-throughput screen. DMS combined with enzymatic and kinetic measurements has the potential to couple biochemistry, structural biology, and genomics in which the unbiased screens provide a unique understanding of the complex systems at play in both protein evolution and protein architecture.

EXPERIMENTAL PROCEDURES

Generation of phage display library

WT phPAI-1 and the phPAI-1 library were generated, characterized, and expressed as described previously, with the library exhibiting a depth of 8.04×10^6 independent clones with an average of 4.2 ± 1.8 amino acid substitutions per PAI-1 fusion protein (59).

phPAI-1 latency screen

phPAI-1 (100 uL at $\sim 10^{12}$ phage/mL, ~ 0.17 nM final concentration) were incubated in TBS containing 5% BSA (900 µL) at 37°C and aliquots assayed over a 48 h time course (0, 1, 2, 4, 8, 12, 24, 48 h). At each time point, uPA (1.7 nM) was incubated with the phage for 30 min at 37°C. With an estimated valency of less than one PAI-1 fusion protein per phage (60), this is equivalent to at least a 10-fold excess of uPA. Residual uPA activity was inhibited by the addition of 100 uL of a 10X stock of cOmplete EDTA-free protease inhibitor cocktail (Millipore Sigma) and incubated for 10 min at 37 °C. phPAI-1:uPA complexes were further immunoprecipitated, washed, eluted, and quantified as described previously (59). In parallel experiments, phPAI-1 was immunoprecipitated via its N-terminal myc-tag over the same time course (Figs. 1).

High-throughput sequencing (HTS)

Twelve overlapping amplicons (150 bp) were PCR amplified from pAY-FE PAI-1 (SI Table 1 and SI Fig. 2) and prepared for HTS as previously described (44). HTS data were analyzed using the DESeq2 software package (61). Statistical thresholds ($p_{adj} < 0.05$ and base mean score > 10) were used to classify a mutation as significant for its effect on functional stability, where p_{adj} is the minimum false discovery rate when calling a mutation significant and the base mean score is the mean of the normalized counts across all samples. Missense mutations in PAI-1 that were nonfunctional were removed from subsequent analyses (44), and a functional stability score was calculated for each mutation by subtracting the $\log_2$-fold change (uPA-selected over unselected) for WT PAI-1 from that of each amino acid substitution at each position.

Recombinant PAI-1 (rPAI-1)

Human *SERPINE1* cDNA was subcloned from pAY-FE-PAI-1 using ligation independent cloning into the pET Flag TEV LIC cloning vector (2L-T, a gift from Scott Gradia—Addgene plasmid #29175; http://n2t.net/addgene:29715; RRID:Addgene_29715) using primers pET PAI-1 LIC For and pET PAI-1 LIC Rev (SI Table 1) and transformed into BL21(DE3) chemically competent *E.*

coli (Invitrogen) (62). Point mutations (I91F and K176E) were introduced using the

QuickChange II XL Site-Directed Mutagenesis Kit (SI Table 1; Agilent Technologies). The full-

length sequence of the PAI-1 constructs was confirmed by Sanger sequencing (SI Table 1). For

recombinant PAI-1 (rPAI-1) expression and purification, *E. coli* containing pET-PAI-1 constructs

were grown at 37°C in LB media (100 mL; Invitrogen) supplemented with ampicillin (100 µg/mL)

to mid-log phase (OD_{600} ~0.40-0.45) prior to induction with IPTG (1 mM) and continued growth

for 2 h at 37 °C. Bacteria were pelleted by centrifugation (3500xg for 20 min at 4°C) and washed

twice with NaCl (0.85M). The bacterial pellet was resuspended in sodium phosphate (50 mM)

pH 7.0 containing NaCl (100 mM) and EDTA (4 mM) and subsequently incubated with T4

lysozyme (1 mg/mL; Millipore Sigma) for 30 min at ambient temperature. Following three freeze-

thaw cycles (1 min in liquid nitrogen, 5 min at 37°C), the bacterial genomic DNA was sheared by

passing 5-10 times through a 20-gauge needle, and crude cell extracts isolated following

centrifugation (16,000xg for 20 min at 4°C). rPAI-1 was further purified by binding to anti-FLAG

agarose beads and elution with 3X FLAG peptide (Millipore Sigma). rPAI-1 was quantified

spectrophotometrically (ε_{280nM} = 43,000 M^{-1} cm^{-1}). To measure its functional half-life, rPAI-1(20

nM) was incubated at 37°C over a 24 h time course and assayed for activity. At selected time

points, the rPAI-1 was diluted between 2- and 40-fold in HEPES (40 mM) pH 7.4 containing 100

mM NaCl and 0.005% Tween-20 and its inhibitory activity against uPA (2.5 nM) was measured

following incubation for 30 min at room temperature. Residual uPA activity was assayed by the

rate of proteolysis of the fluorescent substrate Z-Gly-Gly-Arg-AMC (Bachem) over a 10 min time

course (λ_{ex} = 370 nM, λ_{em} = 440 nM) (SI Fig. 1).

Transition to latency half-life calculation

The functional half-lives of both phPAI-1 and r PAI-1 were determined by plotting either the

fraction of phPAI-1 recovered or relative activity of rPAI-1 that remains active after fitting with a

one-phase exponential decay function (Eq.1)

$y = e^{(-kt)} + c$ Eq. 1

where y is the PAI-1 fractional activity, k is the rate constant for the first-order transition of PAI-1

from its active to latent form, t is time, and c is the asymptotic value. Half-life ($t_{1/2}$) was further

determined as described in Eq. 2.

$t_{1/2} = \frac{\ln(2)}{k}$ Eq. 2

Massively parallel half-life kinetics

To determine the half-lives of individual PAI-1 missense variants within the library, the fractional activity of the library at each time point over a 48 h time course (Fig. 1) was multiplied by the fraction of counts representing a given missense mutation that were present at each time point. Following normalization of the 0 h time point to 1, the data were fit to an exponential decay function (Eq. 1) and the apparent half-life extracted (Eq. 2). In order to account for variations in the mutational background arising from error prone PCR, the variants at a given position were normalized by setting the WT amino acid substitution to half-life of 2.1h.

DATA AVAILABILITY

All data is contained within the manuscript and supplementary files. Raw sequencing data will be made available upon request.

SUPPORTING INFORMATION

This article contains supporting information (3 Figures, 1 Table, and 2 Data Files).

FUNDING AND ADDITIONAL INFORMATION

This work was supported by the National Institutes of Health grants R35-HL135793T (DG),R01-HL055374 (DAL), R01-AG074552 (DAL) T32-GM007863 (ZMH), and T32-HL125242 (ZMH). LMH was supported by a National Hemophilia Foundation Judith Graham Pool Postdoctoral Research Fellowship. AY is supported by an American Society of Hematology Faculty Scholar Award, National Hemophilia Foundation Innovative Investigator Research Award, National Institutes of Health grant R01-HL154688, and the Mary R. Gibson Foundation. CAK is supported by an Early Career Award from Hamilton Health Sciences. DAL has additional support from the American Heart Association (19TPA34880040). DG is a Howard Hughes Medical Institute investigator. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest with the contents of this article.

1 **ACKNOWLEDGEMENTS**

2 We thank Kyle Kihn, Elisa Marchiori, Giovanni Spagnolli, Alberto Boldrini, Luca Terruzzi, Daniel

3 Lawrence, Anne Gershenson, Pietro Facciolo, and Patrick Wintrode for access to their

4 computational models prior to publication.

REFERENCES

1. Sanrattana, W., Maas, C., and de Maat, S. (2019) SERPINs—From Trap to Treatment. *Frontiers in Medicine* **6**

2. Spence, M. A., Mortimer, M. D., Buckle, A. M., Minh, B. Q., and Jackson, C. J. (2021) A Comprehensive Phylogenetic Analysis of the Serpin Superfamily. *Molecular Biology and Evolution* **38**, 2915-2929

3. Gettins, P. G., and Olson, S. T. (2009) Exosite determinants of serpin specificity. *Journal of Biological Chemistry* **284**, 20441-20445

4. Lawrence, D. A., Ginsburg, D., Day, D. E., Berkenpas, M. B., Verhamme, I. M., Kvassman, J. O., and Shore, J. D. (1995) Serpin-protease complexes are trapped as stable acyl-enzyme intermediates. *J. Biol. Chem.* **270**, 25309-25312

5. Lawrence, D. A., Olson, S. T., Muhammad, S., Day, D. E., Kvassman, J. O., Ginsburg, D., and Shore, J. D. (2000) Partitioning of serpin-proteinase reactions between stable inhibition and substrate cleavage is regulated by the rate of serpin reactive center loop insertion into beta-sheet A. *J. Biol. Chem.* **275**, 5839-5844

6. Sillen, M., and Declerck, P. J. (2021) A Narrative Review on Plasminogen Activator Inhibitor-1 and Its (Patho)Physiological Role: To Target or Not to Target? *International Journal of Molecular Sciences* **22**, 2721

7. Declerck, P. J., De Mol, M., Alessi, M. C., Baudner, S., Pâques, E. P., Preissner, K. T., Müller-Berghaus, G., and Collen, D. (1988) Purification and characterization of a plasminogen activator inhibitor 1 binding protein from human plasma. Identification as a multimeric form of S protein (vitronectin). *Journal of Biological Chemistry* **263**, 15454-15461

8. Keijer, J., Linders, M., Wegman, J. J., Ehrlich, H. J., Mertens, K., and Pannekoek, H. (1991) On the Target Specificity of Plasminogen Activator Inhibitor 1: The Role of Heparin, Vitronectin, and the Reactive Site. *Blood* **78**, 1254-1261

9. Rezaie, A. R. (2001) Vitronectin Functions as a Cofactor for Rapid Inhibition of Activated Protein C by Plasminogen Activator Inhibitor-1: Implications for the mechanism of profibrinolytic action of activated protein C. *J Biol Chem* **276**, 15567-15570

10. Berrettini, M., Schleef, R. R., España, F., Loskutoff, D. J., and Griffin, J. H. (1989) Interaction of type 1 plasminogen activator inhibitor with the enzymes of the contact activation system. *Journal of Biological Chemistry* **264**, 11738-11743

11. Bernot, D., Stalin, J., Stocker, P., Bonardo, B., Scroyen, I., Alessi, M. C., and Peiretti, F. (2011) Plasminogen activator inhibitor 1 is an intracellular inhibitor of furin proprotein convertase. *J Cell Sci* **124**, 1224-1230

12. Dittmann, M., Hoffmann, H. H., Scull, M. A., Gilmore, R. H., Bell, K. L., Ciancanelli, M., Wilson, S. J., Crotta, S., Yu, Y., Flatley, B., Xiao, J. W., Casanova, J. L., Wack, A., Bieniasz, P. D., and Rice, C. M. (2015) A serpin shapes the extracellular environment to prevent influenza A virus maturation. *Cell* **160**, 631-643

13. Böttcher-Friebertshäuser, E., Freuer, C., Sielaff, F., Schmidt, S., Eickmann, M., Uhlendorff, J., Steinmetzer, T., Klenk, H. D., and Garten, W. (2010) Cleavage of influenza virus hemagglutinin by airway proteases TMPRSS2 and HAT differs in subcellular localization and susceptibility to protease inhibitors. *J Virol* **84**, 5605-5614

14. Shen, L. W., Mao, H. J., Wu, Y. L., Tanaka, Y., and Zhang, W. (2017) TMPRSS2: A potential target for treatment of influenza virus and coronavirus infections. *Biochimie* **142**, 1-10

15. Hoffmann, M., Kleine-Weber, H., Schroeder, S., Krüger, N., Herrler, T., Erichsen, S., Schiergens, T. S., Herrler, G., Wu, N.-H., Nitsche, A., Müller, M. A., Drosten, C., and Pöhlmann, S. (2020) SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280.e278

16. Carmeliet, P., Kieckens, L., Schoonjans, L., Ream, B., van Nuffelen, A., Prendergast, G., Cole, M., Bronson, R., Collen, D., and Mulligan, R. C. (1993) Plasminogen activator inhibitor-1 gene-deficient mice. I. Generation by homologous recombination and characterization. *J. Clin. Invest.* **92**, 2746-2755

17. Carmeliet, P., Stassen, J. M., Schoonjans, L., Ream, B., van den Oord, J. J., De Mol, M., Mulligan, R. C., and Collen, D. (1993) Plasminogen activator inhibitor-1 gene-deficient mice. II. Effects on hemostasis, thrombosis, and thrombolysis. *The Journal of clinical investigation* **92**, 2756-2760

18. Cale, J. M., and Lawrence, D. A. (2007) Structure-function relationships of plasminogen activator inhibitor-1 and its potential as a therapeutic agent. *Curr Drug Targets* **8**, 971-981

19. Fay, W. P., Shapiro, A. D., Shih, J. L., Schleef, R. R., and Ginsburg, D. (1992) Brief report: complete deficiency of plasminogen-activator inhibitor type 1 due to a frame-shift mutation. *N. Engl. J. Med.* **327**, 1729-1733

20. Fay, W. P., Parker, A. C., Condrey, L. R., and Shapiro, A. D. (1997) Human plasminogen activator inhibitor-1 (PAI-1) deficiency: characterization of a large kindred with a null mutation in the PAI-1 gene. *Blood* **90**, 204-208

21. Khan, S. S., Shah, S. J., Klyachko, E., Baldridge, A. S., Eren, M., Place, A. T., Aviv, A., Puterman, E., Lloyd-Jones, D. M., Heiman, M., Miyata, T., Gupta, S., Shapiro, A. D., and Vaughan, D. E. (2017) A null mutation in SERPINE1 protects against biological aging in humans. *Sci. Adv.* **3**, eaao1617

22. Sillen, M., and Declerck, P. J. (2020) Targeting PAI-1 in Cardiovascular Disease: Structural Insights Into PAI-1 Functionality and Inhibition. *Frontiers in Cardiovascular Medicine* **7**

23. Morrow, G. B., Whyte, C. S., and Mutch, N. J. (2021) A Serpin With a Finger in Many PAIs: PAI-1's Central Function in Thromboinflammation and Cardiovascular Disease. *Front Cardiovasc Med* **8**, 653655

24. Declerck, P. J., De Mol, M., Alessi, M. C., Baudner, S., Paques, E. P., Preissner, K. T., Muller-Berghaus, G., and Collen, D. (1988) Purification and characterization of a plasminogen activator inhibitor 1 binding protein from human plasma. Identification as a multimeric form of S protein (vitronectin). *J Biol Chem* **263**, 15454-15461

25. Verhamme, I., Kvassman, J. O., Day, D., Debrock, S., Vleugels, N., Declerck, P. J., and Shore, J. D. (1999) Accelerated conversion of human plasminogen activator inhibitor-1 to its latent form by antibody binding. *J Biol Chem* **274**, 17511-17517

26. Hägglöf, P., Bergström, F., Wilczynska, M., Johansson, L. B. Å., and Ny, T. (2004) The Reactive-center Loop of Active PAI-1 is Folded Close to the Protein Core and can be Partially Inserted. *Journal of molecular biology* **335**, 823-832

27. Cazzolli, G., Wang, F., a Beccara, S., Gershenson, A., Faccioli, P., and Wintrode, P. L. (2014) Serpin latency transition at atomic resolution. *Proceedings of the National Academy of Sciences* **111**, 15414-15419

28. Petersen, M., Madsen, J. B., Jorgensen, T. J. D., and Trelle, M. B. (2017) Conformational preludes to the latency transition in PAI-1 as determined by atomistic computer simulations and hydrogen/deuterium-exchange mass spectrometry. *Sci Rep* **7**, 6636

29. Gorlatova, N. V., Elokdah, H., Fan, K., Crandall, D. L., and Lawrence, D. A. (2003) Mapping of a conformational epitope on plasminogen activator inhibitor-1 by random mutagenesis. Implications for serpin function. *J Biol Chem* **278**, 16329-16335

30. Li, S. H., Gorlatova, N. V., Lawrence, D. A., and Schwartz, B. S. (2008) Structural differences between active forms of plasminogen activator inhibitor type 1 revealed by conformationally sensitive ligands. *J Biol Chem* **283**, 18147-18157

31. Dewilde, M., Van De Craen, B., Compernolle, G., Madsen, J. B., Strelkov, S., Gils, A., and Declerck, P. J. (2010) Subtle structural differences between human and mouse PAI-1 reveal the basis for biochemical differences. *Journal of Structural Biology* **171**, 95-101

32. Jendroszek, A., Sønnichsen, M. S., Muñoz, A. C., Leyman, K., Christensen, A., Petersen, S. V., Wang, T., Bendixen, C., Panitz, F., Andreasen, P. A., and Jensen, J. K. (2017) Latency transition of plasminogen activator inhibitor type 1 is evolutionarily conserved. *Thrombosis and haemostasis* **117**, 1688-1699

33. Bager, R., Johansen, J. S., Jensen, J. K., Stensballe, A., Jendroszek, A., Buxbom, L., Sørensen, H. P., and Andreasen, P. A. (2013) Protein conformational change delayed by steric hindrance from an N-linked glycan. *Journal of molecular biology* **425**, 2867-2877

34. Kooistra, T., Sprengers, E. D., and van Hinsbergh, V. W. (1986) Rapid inactivation of the plasminogen-activator inhibitor upon secretion from cultured human endothelial cells. *Biochem J* **239**, 497-503

35. Levin, E. G., and Santell, L. (1987) Conversion of the active to latent plasminogen activator inhibitor from human endothelial cells. *Blood* **70**, 1090-1098

36. Mottonen, J., Strand, A., Symersky, J., Sweet, R. M., Danley, D. E., Geoghegan, K. F., Gerard, R. D., and Goldsmith, E. J. (1992) Structural basis of latency in plasminogen activator inhibitor-1. *Nature* **355**, 270-273

37. Lindahl, T. L., Sigurdardottir, O., and Wiman, B. (1989) Stability of plasminogen activator inhibitor 1 (PAI-1). *Thrombosis and haemostasis* **62**, 748-751

38. Mayer, E. J., Fujita, T., Gardell, S. J., Shebuski, R. J., and Reilly, C. F. (1990) The Pharmacokinetics of Plasminogen Activator Inhibitor-1 in the Rabbit. *Blood* **76**, 1514-1520

39. Berkenpas, M. B., Lawrence, D. A., and Ginsburg, D. (1995) Molecular evolution of plasminogen activator inhibitor-1 functional stability. *EMBO J* **14**, 2969-2977

40. Stoop, A. A., Eldering, E., Dafforn, T. R., Read, R. J., and Pannekoek, H. (2001) Different structural requirements for plasminogen activator inhibitor 1 (PAI-1) during latency transition and proteinase inhibition as evidenced by phage-displayed hypermutated PAI-1 libraries. *J. Mol. Biol.* **305**, 773-783

41. Jankun, J., Aleem, A. M., Selman, S. H., Skrzypczak-Jankun, E., Lysiak-Szydlowska, W., Grafos, N., Fryer, H. J., and Greenfield, R. S. (2007) Highly stable plasminogen activator inhibitor type one (VLHL PAI-1) protects fibrin clots from tissue plasminogen activator-mediated fibrinolysis. *International journal of molecular medicine* **20**, 683-687

42. Dewilde, M., Strelkov, S. V., Rabijns, A., and Declerck, P. J. (2009) High quality structure of cleaved PAI-1-stab. *J Struct Biol* **165**, 126-132

43. Lawrence, D. A., Olson, S. T., Palaniappan, S., and Ginsburg, D. (1994) Engineering plasminogen activator inhibitor 1 mutants with increased functional stability. *Biochemistry* **33**, 3643-3648

44. Huttinger, Z. M., Haynes, L. M., Yee, A., Kretz, C. A., Holding, M. L., Siemieniak, D. R., Lawrence, D. A., and Ginsburg, D. (2021) Deep mutational scanning of the plasminogen activator inhibitor-1 functional landscape. *Sci Rep* **11**, 18827

45. Sherman, P. M., Lawrence, D. A., Yang, A. Y., Vandenberg, E. T., Paielli, D., Olson, S. T., Shore, J. D., and Ginsburg, D. (1992) Saturation mutagenesis of the plasminogen activator inhibitor-1 reactive center. *J. Biol. Chem.* **267**, 7588-7595

46. Kretz, C. A., Dai, M., Soylemez, O., Yee, A., Desch, K. C., Siemieniak, D., Tomberg, K., Kondrashov, F. A., Meng, F., and Ginsburg, D. (2015) Massively parallel enzyme kinetics reveals the substrate recognition landscape of the metalloprotease ADAMTS13. *Proc Natl Acad Sci U S A* **112**, 9328-9333

47. Fowler, D. M., and Fields, S. (2014) Deep mutational scanning: a new style of protein science. *Nat. Methods* **11**, 801-807

48. Fowler, D. M., Stephany, J. J., and Fields, S. (2014) Measuring the activity of protein variants on a large scale using deep mutational scanning. *Nat. Protoc.* **9**, 2267-2284

49. Stein, A., Fowler, D. M., Hartmann-Petersen, R., and Lindorff-Larsen, K. (2019) Biophysical and Mechanistic Models for Disease-Causing Protein Variants. *Trends Biochem Sci* **44**, 575-588

50. McCormick, J. W., Russo, M. A. X., Thompson, S., Blevins, A., and Reynolds, K. A. (2021) Structurally distributed surface sites tune allosteric regulation. *eLife* **10**, e68346

51. Markin, C. J., Mokhtari, D. A., Sunden, F., Appel, M. J., Akiva, E., Longwell, S. A., Sabatti, C., Herschlag, D., and Fordyce, P. M. (2021) Revealing enzyme functional architecture via high-throughput microfluidic enzyme kinetics. *Science* **373**

52. Jensen, J. K., Thompson, L. C., Bucci, J. C., Nissen, P., Gettins, P. G. W., Peterson, C. B., Andreasen, P. A., and Morth, J. P. (2011) Crystal Structure of Plasminogen Activator Inhibitor-1 in an Active Conformation with Normal Thermodynamic Stability. *Journal of Biological Chemistry* **286**, 29709-29717

53. Hansen, M., Busse, M. N., and Andreasen, P. A. (2001) Importance of the amino-acid composition of the shutter region of plasminogen activator inhibitor-1 for its transitions to latent and substrate forms. *European Journal of Biochemistry* **268**, 6274-6283

54. Carrell, R. W., and Stein, P. E. (1996) The biostructural pathology of the serpins: critical function of sheet opening mechanism. *Biol Chem Hoppe Seyler* **377**, 1-17

55. Trelle, M. B., Dupont, D. M., Madsen, J. B., Andreasen, P. A., and Jørgensen, T. J. D. (2014) Dissecting the Effect of RNA Aptamer Binding on the Dynamics of Plasminogen Activator Inhibitor 1 Using Hydrogen/Deuterium Exchange Mass Spectrometry. *ACS Chemical Biology* **9**, 174-182

56. Li, S. H., Reinke, A. A., Sanders, K. L., Emal, C. D., Whisstock, J. C., Stuckey, J. A., and Lawrence, D. A. (2013) Mechanistic characterization and crystal structure of a small molecule inactivator bound to plasminogen activator inhibitor-1. *Proc Natl Acad Sci U S A* **110**, E4941-4949

57. Sui, G. C., and Wiman, B. (2000) The Bbeta-sheet in the PAI-1 molecule plays an important role for its stability. *Thrombosis and haemostasis* **83**, 896-901

58. Kihn, K., Marchiori, E., Spagnolli, G., Boldrini, A., Terruzzi, L., Lawrence, D., Gershenson, A., Faccioli, P., and Wintrode, P. L. (2022) Long range allostery mediates the regulation of plasminogen activator inhibitor-1 by vitronectin. *Unpublished*

59. Huttinger, Z. M., Haynes, L. M., Yee, A., Kretz, C. A., Siemieniak, D. R., Lawrence, D. A., and Ginsburg, D. (2021) Deep mutational scanning of the plasminogen activator inhibitor-1 functional landscape. *bioRxiv*, 2021.2004.2015.440003

60. Barbas, C. F., Burton, D. R., Scott, J. K., and Silverman, G. J. (2004) *Phage Display: A Laboratory Manual*, Cold Spring Harbor Laboratory Press

61. Love, M. I., Huber, W., and Anders, S. (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome biology* **15**, 550

62. Gradia, S. D., Ishida, J. P., Tsai, M.-S., Jeans, C., Tainer, J. A., and Fuss, J. O. (2017) Chapter One - MacroBac: New Technologies for Robust and Efficient Large-Scale Production of Recombinant Multiprotein Complexes. in *Methods in Enzymology* (Eichman, B. F. ed.), Academic Press. pp 1-26

**Figure 1. The phPAI-1 mutant library undergoes an extended latency transition compared
to WT phPAI-1.** Relative quantity of WT phPAI-1 (closed red circles) or mutant library phPAI-1
(open red circles) recovered by uPA complex formation and α-uPA immunoprecipitation at
selected time points (0, 2, 4, 8, 12, 24, 48 h) at 37° C . The data were fit to a one-phase
exponential decay function (Eq. 1) and the apparent half-lives of WT phPAI-1 and the mutant
library were determined (Eq. 2). *Myc*-tag immunoprecipitation (closed blue circles) preformed at
selected time points (0, 12, 24, 48 h) at 37° C shows no significant decrease in number of
phage recovered after 48 h incubation. Mean phage ± SD relative to amount
immunoprecipitated at time 0 is plotted. N=3 per group for all time points.

1
2
Figure 2. DMS identifies PAI-1 missense variants that exhibit inhibitory activity towards uPA with changes in functional
stability. Heatmap indicating functional stability relative to WT PAI-1 for phPAI-1 variants selected for uPA inhibitory activity at 48 h.
Amino acid position is indicated on the x-axis, while amino acid substitutions are indicated on the y-axis. Variants with increased
functional stability relative to WT are indicated in blue, while those with decreased functional stability are highlighted in red. WT
amino acid residues are shown in dark grey, while beige indicates missense mutations that were present in the mutational library but
did not meet statistical thresholds(44). Regions enriched in variants exhibiting enhanced functional stability include the RCL (residues
331-350, *black*) as well as residues 40-49 (*green*), residues 73-98 (*cyan*), residues 145-174 (*purple*), residues 217-232 *(purple)*.
residues 261-276 (*purple*), and residues 319-347 (*orange*).

1
2
Figure 3. Massively parallel kinetics can accurately determine the half-lives of PAI-1 variants. *(A)* Heatmap indicating average
half-lives of phPAI-1 variants. Missense variants with a prolonged half-life ($t_{1/2}$ > 2.1 h) are shown in blue, while those that are
decreased are shown in red ($t_{1/2}$ > 2.1 h). WT amino acid residues are indicated by dark grey, while beige indicates missense
mutations that were present in the mutational library but for which a half-life different from WT could not be distinguished. *(B)* Half-
lives for missense phPAI-1 variants determined using massively parallel kinetics were compared to those described for recombinant
PAI-1 both in this study and previously reported (45). A linear regression (slope = 0.5, R = 0.8) indicates a strong positive correlation
between our massively parallel method and direct half-life determination from individually expressed and purified proteins.

1
2
Figure 4. DMS of PAI-1 reveals key regions that regulate its functional stability. The
crystal structure of active PAI-1 (PBD: 3Q02 (52)) is shown *(grey)* with regions of increased
functional stability relative to WT PAI-1 highlighted: residues 40-49 map to α-helix B *(green)*,
residues 73-98 correspond to helix D and β-strand 2A *(cyan)*, residues 145-174 encompass the
Val[157]-loop and β-strand 3A *(purple)*, residues 217-232 correspond to β-strand 2B *(purple)*.
residues 261-276 encompass helix H and β-strand 2C *(purple)*, and residues 319-347 include β-
strand 5A as well as the N-terminal portion of the RCL *(orange)*. *Inset,* Ile[91] and Leu[169] are
located in proximity to each other on the interior of β-sheet A.
11
12
13
14

1 <u>Table 1. 35 most functionally stable PAI-1 variants relative to WT</u>

Mutation	Functional Stability Score	p_{adj}	Massively Parallel Half-life, h
I91L	3.73	2.27×10^{-188}	ND
L169S	3.67	6.25×10^{-29}	ND
I91F	3.47	$<10^{-294}$	7.7
F372V	3.39	$<10^{-294}$	6.1
D222H	3.22	5.14×10^{-136}	ND
I91M	3.20	5.51×10^{-165}	ND
S331G	2.97	$<10^{-294}$	6.8
D222Y	2.95	4.76×10^{-293}	ND
Y241R	2.94	2.66×10^{-26}	ND
M45K	2.82	$<10^{-294}$	3.9
G332S	2.81	4.63×10^{-242}	6.2
T333A	2.74	$<10^{-294}$	7.8
Y221C	2.72	5.47×10^{-240}	ND
F372I	2.72	$<10^{-294}$	4.0
G332D	2.70	9.24×10^{-225}	7.0
R271L	2.67	3.20×10^{-239}	3.7
K154I	2.62	1.21×10^{-266}	3.3
G332A	2.62	2.48×10^{-124}	ND
F372C	2.57	8.19×10^{-123}	ND
D222V	2.56	1.54×10^{-263}	ND
E378D	2.45	7.76×10^{-236}	4.9
G332R	2.44	1.60×10^{-55}	ND
T323I	2.40	$<10^{-294}$	ND
L272R	2.38	1.95×10^{-193}	3.4
K176E	2.38	1.26×10^{-210}	4.1
D222N	2.37	$<10^{-294}$	2.0
F372L	2.34	$<10^{-294}$	3.7
T333S	2.33	8.77×10^{-65}	8.4
Y241H	2.31	$<10^{-294}$	2.0
V334G	2.28	1.37×10^{-89}	ND
S198C	2.26	4.44×10^{-118}	ND
G332V	2.18	1.80×10^{-70}	7.6
W86L	2.17	9.72×10^{-274}	13.1
R271C	2.16	4.50×10^{-277}	3.9
L169H	2.14	1.68×10^{-88}	3.2

2   p_{adj} = adjusted p-value as defined by Love, *et al.* (61)

3   ND = not determined due to insufficient data

2
3
4 **SI Figure 1. Rates of latency transition for recombinant PAI-1 variants** *(A)* **I91F and** *(B)*
5 **K176E.** Activity for WT rPAI-1 is indicated in black and single amino acid substituted variants
6 are shown in red (n = 3). The half-life of I91L rPAI-1 was determined to be 5.4 h and while for
7 K176E it was 4.8 h. WT rPAI-1 had a half-life of 2.2 h.
8

1
2
3 **SI Figure 2. A hypothetical mutated PAI-1 cDNA is shown with 4 mutations.** Mutated PAI-1
4 cDNAs are sequenced using 12 overlapping 150 bp amplicons. Most amplicons will be WT with
5 no mutations. Most mutations will only appear in 1 amplicon (blue and brown) and phase will be
6 lost with mutations in *cis* in the original cDNA but occurring in different amplicons.
7

SI Figure 3. Identification of PAI-1 regions enriched in variants that stabilize the active
conformation. PAI-1 amino acid positions are plotted on the x-axis with black circles
representing the sum of functional stability scores divided by the number of variants scored at
each position The data were fit to a LOWESS regression (*right* axis, blue) with a 20 point
smoothing window determined with the GraphPad Prism software package (v. 9.0.2). Six
regions of increased functional stability were identified with this method (LOWESS regression >
0): residues 40-49, 73-98, 145-174, 217-232, 261-276, and 319-347.

1   SI Table 1. Primer Sequences
2

Primer	5' → 3' Sequence
Sanger Sequencing/Colony PCR Primers	
pAYE Sequencing For	TTCATGCTGCCGGCTTTCTCGG
pAYE Sequencing Rev	TCGTCATCGTCCTTGTAGTCACCACCA
HTS Sequencing Primers	
PAI-1 Amplicon 1 For	NNNNNNGAACAAAAACTCATCTCAGAAGAGGATCTG
PAI-1 Amplicon 1 Rev	NNNNNNGGGTGAGAAAACCACGTTGCG
PAI-1 Amplicon 2 For	NNNNNNTTTCAGCAGGTGGCGCAG
PAI-1 Amplicon 2 Rev	NNNNNNCTTGTCATCAATCTTGAATCCCATAGCTGC
PAI-1 Amplicon 3 For	NNNNNNATGCTCCAGCTGACAACAGGA
PAI-1 Amplicon 3 Rev	NNNNNNTGTGGTGCTGATCTCATCCTTGTT
PAI-1 Amplicon 4 For	NNNNNNGCCCTCCGGCATCTGTACAAG
PAI-1 Amplicon 4 Rev	NNNNNNTTGCTTGACCGTGCTCCGG
PAI-1 Amplicon 5 For	NNNNNNAAGCTGGTCCAGGGCTTCATG
PAI-1 Amplicon 5 Rev	NNNNNNCCCAAGCAAGTTGCTGATCATACC
PAI-1 Amplicon 6 For	NNNNNNATTCATCATCAATGACTGGGTGAAGAC
PAI-1 Amplicon 6 Rev	NNNNNNCTGGAGTCGGGGAAGGGAG
PAI-1 Amplicon 7 For	NNNNNNAATGCCCTCTACTTCAACGGC
PAI-1 Amplicon 7 Rev	NNNNNNGGGCGTGGTGAACTCAGTATAG
PAI-1 Amplicon 8 For	NNNNNNCCCATGATGGCTCAGACCAAC
PAI-1 Amplicon 8 Rev	NNNNNNGGCAGAGAGAGGCACCTCT
PAI-1 Amplicon 9 For	NNNNNNAGCATGTTCATTGCTGCCCC
PAI-1 Amplicon 9 Rev	NNNNNNTTCAGTCTCCAGGGAGAACTTGG
PAI-1 Amplicon 10 For	NNNNNNCAGGCTGCCCCGCCT
PAI-1 Amplicon 10 Rev	NNNNNNACGTGGAGAGGCTCTTGGTC
PAI-1 Amplicon 11 For	NNNNNNAGACAGTTTCAGGCTGACTTCAC
PAI-1 Amplicon 11 Rev	NNNNNNGGGGGCCATGCGGGC
PAI-1 Amplicon 12 For	NNNNNNTCATCCACAGCTGTCATA
PAI-1 Amplicon 12 Rev	NNNNNNTGCGGCCGCTGAACCACCTCC
pET-PAI-1 LIC Primers	
pET PAI-1 LIC For	TACTTCCAATCCAATGCAGTGCACCATCCCCCATCCTAC
pET PAI-1 LIC Rev	TTATCCACTTCCAATGTTATTATCAGGGTTCCATCACTTGGCCCA
pET PAI-1 Site Directed Mutagenesis Primers	
PAI-1 I91F SDM For	GCTCATGGGGCCATtGAACAAGGATGAGATCAGCACCACAGACGC
PAI-1 I91F SDM Rev	ATCGCGTCTGTGGTGCTGATCTCATCCTTGTTCaATGGCCCCATG
PAI-1 K176E SDM For	GCCCTCTACTTCAACGGCCAGTGGgAGACTCCCTTCCCCGACTCC
PAI-1 K176E SDM Rev	GGAGTCGGGGAAGGGAGTCTcCCACTGGCCGTTGAAGTAGAGGGC
PAI-1 Y221C SDM For	CCCGATGGCCATTACTgCGACATCCTGGAACTGCCCTACCACGGG
PAI-1 Y221C SDM Rev	GGTAGGGCAGTTCCAGGATGTCGcAGTAATGGCCATCGGGCGTGG

23