Full Title: *Rickettsia* infection rate along an altitudinal gradient as influenced by population

genetic structure of Ixodid ticks

Short Title: Tick population genetic structure and Rickettsia infection rate along an altitudinal

gradient

Author names and affiliations

Maria Angenica F. Regilme[a], Megumi Sato[b], Tsutomu Tamura[c], Reiko Arai[c], Marcello Otake Sato[d],

Sumire Ikeda[e], Kozo Watanabe[a]*

[a] Center for Marine Environmental Studies, Ehime University, Matsuyama, Ehime 790-8577, Japan

[b] Graduate School of Health Sciences, Niigata University, Niigata, 951-8518, Japan

[c] Niigata Prefectural Institute of Public Health and Environmental Sciences, Niigata, 950-2144,

Japan

[d] Department of Tropical Medicine and Parasitology, Dokkyo Medical University, 880

Kitakobayashi, Mibu-machi, Shimotsuga-gun, Tochigi 321-0293, Japan

[e] Research Laboratories, Research and Development Headquarters, Earth Corporation, Hyogo 678-

0192, Japan

Corresponding author*

watanabe.kozo.mj@ehime-u.ac.jp

Center for Marine Environmental Studies, Ehime University, Bunkyo-cho 3, Matsuyama, Ehime,

Japan 790-8577

Abstract

Ixodid ticks, such as *Ixodes ovatus* and *Haemaphysalis flava,* are important vectors of tick-borne diseases in Japan, such as Japanese spotted fever caused by *Rickettsia japonica*. This study describes the population genetic structure and gene flow of *I. ovatus* and *H. flava* and their *Rickettsia* infection rates along an altitudinal gradient. A total of 346 adult *I. ovatus* and 243 *H. flava* were analyzed for the presence of *Rickettsia* by nested PCR targeting the 17kDA, *gltA, rOmpA,* and *rOmpB* genes. Population genetic structure was performed using the mitochondrial cytochrome oxidase 1 (*cox1*) marker. The *Rickettsia* infection rates were 13.26% in *I. ovatus* and 6.17% in *H. flava.* For *I. ovatus,* the global F_{ST} value revealed significant genetic differentiation among the different populations, whereas *H. flava* showed non-significant genetic differentiation. The *cox1 I. ovatus* cluster dendrogram showed two cluster groups, while the haplotype network and phylogenetic tree showed three genetic groups. The two cluster groups and the three genetic groups found in *I. ovatus* showed a significant difference in their *Rickettsia* infection rates and mean altitude per group. No significant differences were found in the mean altitude or *Rickettsia* infection rates of *H. flava*. Our results suggest that low gene flow in the *I. ovatus* populations has caused spatially heterogenous *Rickettsia* infection rates along the altitudinal gradient. This information can be used in understanding the relationship between the tick vector, its pathogen, and environmental factors, such as altitude, and for the control of tick-borne diseases in Japan.

Keywords: environment, haplotype, hosts, pathogen, dispersal, vector

1. Introduction

Tick-borne diseases are a significant public health concern in Japan and are transmitted by a diverse range of tick species, such as *Ixodes ovatus* [1] that potentially transmit *Borrelia sp.* causing *Lyme* disease [2] and *Haemaphysalis flava* [3], which transmits *Rickettsia japonica* and is a suspected vector of severe fever with thrombocytopenia syndrome virus [4-6]. Since ticks are small, their dispersal is closely related to the movement of their hosts [7-8]. The dynamics of tick-borne pathogens are influenced by the habitat distribution and dispersal behaviors of vectors and hosts along environmental gradients [9]. Therefore, understanding the complex interaction between these factors is important in understanding the spread of tick-borne diseases in Japan.

Tick population genetic analysis provides data that help identify the dispersal pattern of ticks based on gene flow between local populations [10]. Tick dispersal of many three-host Ixodid ticks depends on the movements of their vertebrate hosts, which influences each tick's potential to spread its pathogen [7-8]. For example, contrasting patterns in the population genetic structures of *I. ovatus* and *H. flava* in the Niigata Prefecture of Japan suggest that host mobility during the immature stages of tick development may influence the genetic structure of adult ticks by affecting survivability into their adult stages [11-12]. *I. ovatus* populations had greater genetic divergence possibly due to the limited dispersal of their small mammalian hosts during the immature development stage; *H. flava* populations showed a more homogenized structure possibly due to the larger mobility of their large mammalian hosts and avian-mediated dispersal [11]. Other studies have also revealed low gene flow in ticks with low-mobility hosts (e.g., small mammals) and higher gene flow in ticks with highly mobile hosts (e.g., large mammals and birds) [10; 13-16].

The spatial distribution and movement of the vector (i.e., ticks) may determine the spatial

distribution of the pathogen's (i.e., *Rickettsia*) infection rate [17]. Previous studies have shown

that the pathogen infection rate can be influenced by many factors, such as the vector's genetic

diversity, gene flow, and spatial structure [5-6; 11; 18-25]. For example, previous studies have

shown that strong gene flow between local vector populations tends to reduce the spatial

heterogeneity of pathogen infection rates between populations [26-27]. Thus, the spatial

distribution and movement of the vector may affect the spatial distribution of the pathogen. To

our knowledge, no previous studies have examined the relationship between the spatial

heterogeneity of *Rickettsia* infection rates and population genetic structure in ticks.

Environmental factors may relate to the population genetic structure of ticks [28-29], with

limited gene flow increasing genetic variation between populations along an altitudinal gradient,

as reported in several studies on other species [30-32]. Thus, in this study, we expected to see a

highly divergent population genetic structure in *I. ovatus* along an altitudinal gradient due to the

limited movement of their mammalian hosts along that gradient; whereas *H. flava* should show a

less divergent structure along the altitudinal gradient due to the higher mobility of its hosts. To

our knowledge, no studies have focused on tick gene flow along an altitudinal gradient.

In the study by [11], no significant influence of environmental factors, including altitude,

was observed in the genetic structures of *I. ovatus* and *H. flava* based on the mantel test, but the

study did not use any other robust analytical methods to thoroughly examine the influence of

altitude on tick genetic structure. In another study, major spotted fever group *Rickettsia* (SFGR)

prevalence was analyzed in a total of 3,336 immature and adult ticks across the Niigata Prefecture,

Japan in the following tick species: *Dermacentor taiwanensis, Haemaphysalis flava,*

Haemaphysalis hystricis, Haemaphysalis longicornis, Haemaphysalis megaspinosa, Ixodes

columnae, Ixodes monospinosus, Ixodes nipponensis, Ixodes ovatus, and *Ixodes persulcatus* [6].

Three SFGR species namely *Rickettsia asiatica, R. helvetica* and *R. monacensis* were detected in *H.*

flava, Haemaphysalis longicornis, Ixodes monospinus, Ixodes nipponensis, and *Ixodes ovatus*, no

spatial distribution of *Rickettsia* infection rates was found among the local populations. To our

knowledge, no previous studies have considered the influence of environmental factors on tick

population genetic structure or the spatial distribution of *Rickettsia* infection rates along an

altitudinal gradient in local tick populations.

Here, we determine the relationship between the vector tick population genetic structure

and *Rickettsia* infection rates as influenced by an altitudinal gradient. Based on the isolation by

environment (IBE) theory, genetic differentiation increases with environmental variation,

regardless of geographic distance [33-35]. Thus, we hypothesized that limited gene flow among

tick populations along an altitudinal gradient might cause spatially heterogenous *Rickettsia*

infection rates in ticks. The present study used cytochrome oxidase 1 (*cox1*) mitochondrial gene

sequences and *Rickettsia*-infected and uninfected data for *I. ovatus* (n = 307) and *H. flava* (n = 220)

individuals collected in 2016–2017 as reported in [6; 11]. Additionally, we added new *cox1*

sequences and *Rickettsia*-infected/uninfected ticks from *I. ovatus* (n= 39)and *H. flava* (n=23)

individuals and five other species collected in 2018.

2. Materials and Method

Published data of [6;11]

In this study, we used *cox1* sequence data from [11] for *I. ovatus* (n = 307) and *H. flava*

(n = 220) ticks collected from April 2016 to November 2017 from 30 sites across the Niigata

Prefecture, Japan. *Rickettsia* infection rate data were obtained from [6] from *I. ovatus* (n = 29), *H.*

flava (n = 2), *I. monospinus* (n = 4), and *I. nipponensis* (n = 2) from 38 sites across Niigata

Prefecture. The 38 sites surveyed in the previous study by [6] include the 30 sites that were used

in this present study (S1 Table). Please refer to [6;11] for more information about the study sites,

collection, sampling identification, DNA extraction, PCR amplification, and sequencing methods

used in each respective study.

To strengthen our analysis, we also added data collected from April to October 2018 from

additional individuals (n=62;) sampled at 30 sites across the Niigata Prefecture, including two sites

not previously sampled by [11]. At these sites, ticks were collected 2–14 times from six core sites

among the 30 sites, while ticks were collected once at the remaining sites. The altitude at each site

ranged from 8 to 1402 meters above sea level (m.a.s.l.), with a mean altitude of 348 m.a.s.l.

Unpublished data from 2018 collection

Ticks collected were stored at 4 °C in microcentrifuge tubes with 70% ethanol. Each

collected tick was morphologically identified using a stereo microscope following the identification

keys of [36]. Genomic DNA was extracted using Isogenome DNA extraction kits (Nippon Gene Co.

Ltd. Tokyo, Japan) following the manufacturer's recommended protocol.

In this study, we combined previously published data from [6] with our newly collected

data to calculate the *Rickettsia* infection rate, which is the percentage of *Rickettsia*-infected ticks

from each obtained population. We analyzed the obtained tick DNA for spotted fever group

Rickettsia (SFGR) detection and host identification and amplified the mitochondrial gene *cox1* for

population genetic analysis. We performed nested PCR targeting the following genes for the

detection of *Rickettsia* sp.: 17-kDA antigen gene (17-kDA); citrate synthase gene (gltA); spotted

fever group (SFG)-specific outer membrane protein A gene (rOmpA); and outer membrane protein

B gene (rOmpB) as described and analyzed in [6]. Briefly, we first amplified the 17-kDa protein. If

the results were positive, then PCR was performed to target gltA. Samples that were positive with

both 17-kDA and gltA were regarded as positive for SFGR and a nested PCR was performed to

target the rOmpA and rOmpB gene, samples that are positive for 17-Kda, gltA, rOmpA, and rOmpB

genes were sequenced to identify the *Rickettsia* species. The amplified PCR products were purified

using AMPure XP (Beckman Coulter Co., Japan) and sequenced using the Big Dye Terminator Cycle

Sequence Kit (Thermo Fisher Scientific).

The *cox1* mitochondrial gene was amplified by PCR for *cox1* (658 base pairs) using the

primer pairs LCO-1490 (5⏘- AAACTTCAGGGTGACCAAAAAATCA-3') and HCO1–2198 (5⏘–

AAACTTCAGGGTGACCAAAAAATCA - 3) for phylogenetic analysis and tick species identification [37].

The PCR amplification profile included an initial denaturation of 94 °C for 2 min, followed by

denaturation at 94 °C for 30 s, then annealing at 38 °C for 30 s, followed by an extension of 72 °C

for 1 min for 30 cycles, and a final extension of 72 °C for 10 min. The obtained PCR products were

purified using the QIAquick 96 PCR Purification Kit (Qiagen, Germany) following the

manufacturer's instructions and were sequenced by Eurofin Genomics, Inc. (Tokyo, Japan).

Each forward and reverse read was assembled using CodonCode Aligner version 1.2.4

software (https://www.codoncode.com/aligner/). Low-quality bases were removed in the aligned

sequences, and no ambiguous bases were detected. We used the MAFFT alignment online

program (https://mafft.cbrc.jp/alignment/server/) to perform multiple alignments using the

default settings. The sequences were checked for similarities with the deposited reference

sequences from GenBank for sequence quality and tick species confirmation using BLAST

(https://blast.ncbi.nlm.nih.gov/Blast.cgi). All final aligned sequences were checked in Mesquite

version 3.5 [38]. The protein-coding genes were translated to amino acids to confirm the absence

of stop codons.

Population genetic analysis

Multiple sites that are in proximity to each other were combined for population genetic

analysis if less than eight individuals were obtained per site, which resulted in 8 populations

labeled A to H (S1 Table). Three sites were excluded from the population genetic analysis because

of the limited number of obtained individuals (<8) and the lack of a nearby site to combine into a

single population.

The final *cox1* sequences of the tick species: *I. ovatus* and *H. flava* were individually

analyzed using DNASp version 6.12.03 to determine the haplotype diversity per species [39]. The

level of genetic divergence between each population was quantified per species using global F_{ST}.

Significance was tested using Arlequin software version 3.5.2.2 [40] with 9999 permutations.

The genetic relationship between the *I. ovatus* populations was visualized using the

unweighted pair group with the arithmetic mean (UPGMA) cluster method using the APE package

[41] for the RStudio software (R Development Core Team, 2016). A cluster dendrogram was

created using pairwise F_{ST} values genetic distance matrix from GenAlEx.

Haplotype network and phylogenetic analyses

We constructed a haplotype network analysis using PopART program version 1.7

(http://popart.otago.ac.nz/index.shtml) on *cox1 I. ovatus* and *H. flava* sequences to assess

haplotype relationships using the median-joining network algorithm [42]. Briefly, we constructed a

Bayesian phylogenetic tree of *cox1* haplotypes for *I. ovatus* and *H. flava,* respectively, using BEAST

version 1.10.14 [43]. We used the Hasegawa-Kishino-Yano substitution model with estimated base

frequencies. We employed a strict clock model and used the coalescent prior as the tree prior. A

maximum clade credibility tree was acquired using TreeAnnotator version 1.10.14 using trees from

BEAUti version1.10.14 with 90% of the trees as the burn-in. We viewed the constructed maximum

clade credibility tree using FigTree version 1.4.4.

Statistical analysis

To determine whether there was a significant difference in the *Rickettsia* infection rate

between haplotype groups for *I. ovatus* and *H. flava,* we performed a z-score test at $p < 0.05$. To

determine whether there were differences in the mean altitude between the haplotype groups,

we used the Welch t-test at $p < 0.05$.

3. Results

The total number of positive (pos) and negative (neg) ticks for *Rickettsia* infections from

the ticks collected in 2018 were: *I. ovatus* (neg = 22, pos = 17); *H. flava* (neg = 10, pos = 13); *I.*

monospinus (neg = 11, pos = 6); *I. asanumai* (neg = 4, pos=5); *I. nipponensis* (pos = 4); *I. persulcatus*

(neg = 4); and *H. japonica* (pos = 2). In this study, the total number of samples from the previously

published data [6;11] and the unpublished data from 2018 were: *I. ovatus* (n = 346); *H. flava*

(n = 243); *I. monospinus* (n = 19); *I. asanumai* (n = 9); *I. nipponensis* (n = 4); *I. persulcatus* (n = 4);

and *H. japonica* (n = 2). The number of adult ticks whose *cox1* was successfully sequenced per

species were: *I. ovatus* (346); *H. flava* (243); *I. monospinus* (19); *I. asanumai* (9); *I. nipponensis* (4); *I.*

persulcatus (4); and *H. japonica* (2) (Table 1).

Table 1. Summary of the haplotype and *Rickettsia* infection rates among the 7 ixodid tick species

obtained in the Niigata Prefecture, Japan

Tick species ***	ns	n	nh	r	Global F_{ST}
1. *Ixodes ovatus*	30	346	59	46 (13.26%)	0.4154*
2. *Haemaphysalis flava*	18	243	66	15 (6.17%)	0.3597
3. *Ixodes monospinus*	6	19	9	6 (31.58%)	0.9034
4. *Ixodes asanumai*	7	9	4	5 (55.55%)	1.0000
5. *Ixodes nipponensis*	2	4	4	4 (100%)	0.3345
6. *Ixodes persulcatus*	3	4	3	0 (0%)	1.0000
7. *Haemaphysalis japonica*	2	2	2	2 (100%)	**
Total		628		78	

Abbreviations: ns no. of sampling sites; n sample size; nh no of haplotypes; r *Rickettsia* infection rate per species; *p < 0.05** not

enough samples for analysis***tick species identification is based on molecular identification using the *cox1* marker and BLAST results

We detected SFGR in 78 (12.44%) out of 627 ixodid ticks, with the highest detected in *I. ovatus* (46/346; 13.29%), *H. flava* (15/243; 6.17%), and *I. monospinus* (6/19; 31.58%) as summarized in Table 1. Out of the 46 *Rickettsia*-infected *I. ovatus* ticks, 25 displayed a 100% identity match with *Rickettsia asiatica* in the 17kDA, *gltA,* and *rOmpB* markers [6], while an additional 19 adult *I. ovatus* from the 2018 collection were also positive with *R. asiatica.* Two haplotypes were found in the rOmpB and 17kDA markers, respectively. One haplotype was found in only one individual (17369). Two out of the 15 *Rickettsia*-infected *H. flava* ticks were found to have the same haplotypes in the 17Kda, gltA,and rOmpA markers, and were identified as *Rickettsia sp.* (LC461063). The remaining 13 *Rickettsia*-infected *H. flava* ticks were identified as *Rickettsia sp.* Four out of the six *Rickettsia*-infected *I. monospinus* were identified as *Rickettsia helvetica* with the same haplotype found in the 17kDA, gltA, and rOmpB markers.

Based on the population genetic analysis of the *cox1* sequences, we found a significant global F_{ST} of 0.4154 at $p < 0.05$ for *I. ovatus* (Table 1). In contrast, no significant global F_{ST} values

were observed in the *H. flava, I. monospinus, I. asanumai, I. nipponensis,* or *I. persulcatus* samples

(Table 1). There were 59 and 66 *cox1* haplotypes found among the 346 *I. ovatus* and 243 *H. flava*

individuals, respectively. We found the following number of *cox1* haplotypes in the remaining

species: *I. monospinus* (n = 9), *I. asanumai* (n = 4), *I. nipponensis* (n = 4), *I. persulcatus* (n = 3), and

H. japonica (n = 2).

The *cox1* haplotype network of *I. ovatus* (Figure 1) revealed four genetic groups, wherein

three genetic groups (1, 2, and 3) were distributed along different altitudinal gradients, as shown

in Figure 2. These four genetic groups were concordant with the four clusters found in the *I.*

ovatus phylogenetic tree (S1 Figure). The habitat distribution of genetic group 3 was limited to

high altitude sites only (range = 16–912 m.a.s.l.), whereas genetic groups 1 (255–471 m.a.s.l.) and

2 (84–1354 m.a.s.l.) were distributed at lower altitudes (Figure 2). The Welch t-test revealed a

significant difference between the mean altitudes of genetic groups 1 and 3 at $p < 0.05$ (Table 2);

however, no significant difference was observed between genetic groups 1 and 2 or groups 2 and

3. We found a significant difference in the *Rickettsia* infection rates between *I. ovatus* genetic

groups 1 and 2 based on the z-score test, but no significant difference between groups 1 and 3 or

groups 2 and 3 (Table 2). The mean altitude between the *Rickettsia*-infected (=273.72 m.a.s.l.) and

non-infected *I. ovatus* (=369.61 m.a.s.l.) revealed a significant difference based on the Welch t-test

at $p < 0.05$ (Figure 3). The UPGMA dendrogram of *I. ovatus* revealed two genetic clusters, 1 and 2

using the genetic distance among the seven populations excluding one population due to the

limited number of samples (Figure 4).

Figure 1. Median-joining network of the 59 *cox1* haplotype sequences of *Rickettsia* positive and

negative *I. ovatus.* Haplotype groups are indicated by numbers (1 to 4).

Figure 2. The influence of altitude on the habitat distribution of *I. ovatus*. The points indicate the

sampling sites of the collected individuals per *I. ovatus* haplotype group (n = 3) in the haplotype

network (Figure 1). The map shows the elevation level across the sampling area, with Niigata

Prefecture Japan depicted as a color gradation. Group 4, not shown in this figure, with haplotypes

54 and 55 was found on Sado island, which is encircled in black in box 3.

Figure 3. The relationship between the mean altitude of *Rickettsia* positive (n = 46) and negative

(n = 300) *I. ovatus.* Welch t-tests revealed a significant difference in the mean altitude between

Rickettsia-infected and non-infected *I. ovatus* at $p < 0.05$. Populations with combined sites are

labeled A to H.

Figure 4. An unweighted pair group method with the arithmetic mean (UPGMA) dendrogram of *I.*

ovatus based on the pairwise genetic distance (F_{ST}) of *cox1* among the 7 populations across Niigata

Prefecture, Japan. We excluded one population due to the limited number of samples.

Table 2. The differences in *Rickettsia* infection rates and mean altitude in *I. ovatus* haplotype

groups and cluster dendrogram groups. The table shows the distribution of *Rickettsia*-infected and

uninfected *I. ovatus* and the mean altitude in each of the haplotype groups as shown in Figure 1 *(I.*

ovatus haplotype network) and the cluster dendrogram in Figure 4 (*I. ovatus* cluster dendrogram).

The z-score test showed a significant difference at $p < 0.05$ between the *Rickettsia* detection rates

in haplotype groups 1 and 2 (indicated by [ab]) and in cluster dendrogram groups 1 and 2. The Welch

t-test at $p < 0.05$ revealed a significant difference in the mean altitude of haplotype groups 1 and 3

and cluster dendrogram groups 1 and 2 (indicated by [ab]). Haplotype group 4 was not included in

the analysis due to its low sample size.

Haplotype group	Rickettsia infection rate			Mean altitude
	Positive	Negative	Detection rate	
1	1	92	1.07%[ab]	294[ab]
2	28	99	22.04%[ab]	341[a]
3	16	110	12.70%[a]	374[ab]
Dendrogram Group				
1	4	120	3.23%[ab]	30
2	40	227	14.98%[ab]	345

The *cox1* haplotype network of *H. flava* displayed two genetic groups (S2 Figure)

consistent with the *H. flava* phylogenetic tree (S3 Figure). Based on the Welch t-test, we found no

significant differences in mean altitude (205 and 165 m.a.s.l.) at $p < 0.05$ between the two *H. flava*

genetic groups (S3 Table). No significant difference between the *Rickettsia* infection rates of *H.*

flava genetic groups 1 and 2 was observed using the z-score test at $p < 0.05$ (S2 Table).

4. Discussion

Our findings support our hypothesis that a genetically structured tick population, such as *I.*

ovatus, can cause the *Rickettsia* infection rate to be spatially heterogenous due to limited gene

flow along an altitudinal gradient. The significant global F_{ST} estimate of 0.4154 among the *I. ovatus*

populations revealed genetic differentiation between the populations as supported by the

occurrence of two genetic clusters in the cluster dendrogram. Our results were consistent with our

previous study [11] which suggested that the low mobility of the host species for immature *I.*

ovatus contributed to low gene flow in the tick populations. The low *I. ovatus* gene flow along the

altitudinal gradient might have caused the spatial heterogeneity of *Rickettsia* infection rates

among these populations, which is supported by the significant difference found in *Rickettsia*

infection rates between genetic clusters 1 and 2. A similar pattern was observed in the studies of

[44-46] which found that the infection rate of *Borrelia burgdorferi*, the causative agent of Lyme

disease, decreased in ticks along the altitudinal gradient. Low gene flow can cause infected and

uninfected ticks to have limited opportunities to traverse a wider spatial area thus causing a

heterogeneous *Rickettsia* infection rate [47-48].

We found two genetic groups in the *H. flava* haplotype network, but no significant

difference in the *Rickettsia* infection rates between the two groups. These results might be due to

the high gene flow observed in the *H. flava* populations, which enable *Rickettsia*-infected and

uninfected *H. flava* individuals to traverse between the study sites. The high mobility of the large

mammalian hosts used by adult *H. flava* and avian-mediated dispersal during their immature stage

probably contributed to their homogenized population genetic structure [11], and the resulting

homogenized *Rickettsia* infection rates. Large mammalian hosts and birds have a wide dispersal

range that enables the broader movement of *Rickettsia*-infected ticks, as observed in previous

studies of *Amblyomma Americanum* [49-51], *H. flava* [11], and *I. ricinus* [52]. Birds are especially

good at dispersing over large areas since they can easily traverse landscape barriers such as

mountains, fences, glaciers, and oceans that would be difficult for mammals to cross [53].

In addition to host mobility, environmental factors can influence the population genetic

structure of ticks [54]. Each tick species has preferred environmental conditions that are conducive

to completing the tick life cycle, thus influencing the geographical distribution of ticks and the risk

areas for tick-borne diseases [55-56]. Altitude may influence the population genetic structure of

ticks through the effect of altitude on the distribution and abundance of ticks and/or their hosts

[57-59]. Altitudinal differences between populations can affect genetic divergence [60], through,

for example, ecological isolation, which causes natural selection against maladapted immigrants

and limits gene flow [61-62]. For example, organisms adapted to low altitudinal sites that cannot

tolerate the lower temperatures at higher altitudes would not survive if they dispersed to those

higher altitude sites, thus restricting gene flow across the altitudinal gradient [60].

The difference in mean altitude between *I. ovatus* cluster groups 1 and 2 might be due to

adaptive divergence along the altitudinal gradient. Individuals from cluster group 1 were

distributed in higher altitude areas of the northern mountainous area of Niigata Prefecture, while

cluster group 2 individuals were found in the lower altitude regions of northern Niigata Prefecture.

Populations along an altitudinal gradient are prone to differentiating selection pressures, which

result in local adaptation [63]. Altitudinal gradients may also cause the varying ambient

temperature, precipitation, and humidity levels essential to ticks' development and survival [64].

These varying environmental factors may cause ticks to have difficulty dispersing over a wide

habitat range. Thus, ticks may need to adapt to extreme habitats, such as extreme altitude or

precipitation levels, for survival. For example, *Rhipicephalus compositus* were found in altitudes of

1000–2500 m, but optimal conditions were between 1200–2500 m [64-66].

The different *Rickettsia* infection rates and altitudinal ranges between the *I. ovatus*

phylogenetic groups may be explained by adaptive evolutionary theory, which states that

organisms adjust to new or severe changes in their environment to become better suited to their

habitat[67-68]. Based on the relationship between the *I. ovatus* phylogenetic groups and their

mean attitudes, *I. ovatus* might be undergoing local adaptation along the altitudinal gradient due

to the higher genetic differentiation between populations as supported by the significant global F_{ST}

(0.4154) found in *I. ovatus*. Based on isolation by environment (IBE), genetic differentiation will

increase with increased environmental differences independent of geographic distances [33-

34;69]. When environmental conditions differ, the success of immigration in a new habitat is

reduced, which may increase the genetic fixation rate due to a lower chance of outcrossing;

thereby enhancing genetic isolation [70]. Thus, the lower gene flow along the altitudinal gradient

reduced the spatial homogeneity of *Rickettsia* infection rates among the *I. ovatus* tick populations,

thus causing the different *Rickettsia* infection rates obtained.

In this study, only a few (n < 20) *I. monospinus, I. asanumai, I. nipponensis* and *H. japonica*

were collected; however, a high *Rickettsia* infection rate (31.58%–100%) was found despite these

limited numbers. In previous studies, high *Rickettsia* infection rates despite a few individuals were

also observed in *I. monospinus* in several prefectures in Japan [71-73]. The extent of tick habitat

distribution may vary between species, which may have influenced the widely different *Rickettsia*

infection rates observed among the species in this study. The high *Rickettsia* infection rate in these

ticks is probably due to effective transovarial transmission [74]. Furthermore, *Rickettsial*

endosymbionts are inclined to have a high infection rate in some tick species populations because

they contribute to the tick microbiome which might imply nutritional symbiosis in the tick species

with high *Rickettsia* infection despite the low number of samples tested [75-77].   Additionally, the

few collected individuals of these species may be habitat specialists that thrive in a narrow habitat

range in contrast to *I. ovatus* and *H. flava*, which may be habitat generalists capable of thriving in a

wide habitat range [78]. The narrow habitat range of habitat specialist ticks might have enabled

increased interactions between infected and uninfected *Rickettsia* ticks within the small spatial

scale compared to the habitat generalist ticks, thus causing their high *Rickettsia* infection rates.

This conclusion should be considered with caution, since the limited number of individual ticks

collected across the Niigata Prefecture may have caused a bias in the estimated *Rickettsia*

infection rates determined in this study. We suggest conducting future research on *Rickettsia*

infection rates in a wider sampling range with an increased number of *I. monospinus, I. asanumai,*

I. nipponensis, and *H. japonica* individuals to further increase our understanding of the association

between *Rickettsiae* and their tick vectors.

The occurrence of local adaptation in tick populations could affect the future of the tick-

borne disease landscape. Environmental factors, such as precipitation, temperature, and altitude,

have been shown to drive population differentiation in insects, such as *Anopheles* mosquitoes and

Drosophila flies [79-81], but studies on the environmental adaptation of ixodid ticks, such as *I.*

ovatus, and its *Rickettsia* infection susceptibility, have not yet been performed. [7] suggested that

environmental conditions that affect bird hosts can also affect the local adaptation of ticks. Few

studies have assessed such local adaptation in multiple organisms with varying dispersal abilities

[82-86] and are an area in need of future research.

One of the limitations of this study is the use of one mitochondrial gene *cox1*, despite this,

we were able to determine the relationship between the tick population genetic structure and

Rickettsia infection rates as influenced by the altitudinal gradient. The mitochondrial *cox1* gene

has been widely used for population genetic analysis of many tick species and was proven to be

informative in determining the relationship from the subfamily to the population level [87-91].

Mitochondrial genes have a mutation rate that is useful in species-level phylogenetics and can be

used for wide geographic ranges however its resolution is not fine enough to study species

selection [10]. In future studies, we suggest including additional mitochondrial genes and or

nuclear genes.

Since ticks are blood-sucking ectoparasites, they directly influence their mammalian hosts

and the pathogens they transmit [94-96]. The interaction between the vector (tick), host, and

pathogen (*Rickettsia*) is essential in understanding and predicting the risk and transmission of tick-

borne diseases [97]. Understanding the genetic structure of ticks can serve as an alternative

indicator to infer the potential spread of its pathogen [98]. Our study found relationships between

(1) the population genetic structure of ticks and the corresponding *Rickettsia* infection rates, (2)

altitude and the population genetic structure of ticks, and (3) altitude and *Rickettsia* infection rates.

We found that host mobility may influence the genetic structure of ixodid ticks. This information

can be used to design more effective tick-borne disease control programs that focus on screening

and detecting pathogens found in ticks and their mammalian hosts. For example, patterns of

disease transmission from ticks with a high genetic divergence and less mobile hosts, such as *I.*

ovatus, are likely due to the movement of infected hosts rather than infected ticks. Thus,

screening prospective tick hosts for *Rickettsia* infection would be more suitable in this example.

We suggest screening the hosts of immature *I. ovatus*, such as small rodents, instead of screening

ticks. The *Rickettsia* infection rate in tick genetic groups can predict the spread of tick-borne

diseases caused by *Rickettsia,* such as the Japanese spotted fever. We also found that altitude may

influence the *Rickettsia* infection rate of *I. ovatus* genetic groups. This information can be used to

determine the high-risk areas (e.g., lowland, mountains, etc.) of tick-borne diseases along an

altitudinal gradient. Genetically structured arthropod vectors, such as ticks, can have different

vector competencies, and environmental factors, such as altitudinal gradients, that can influence

the vector's ability to acquire, transmit, and maintain the pathogen infection [99].

Acknowledgments

The authors are thankful for the assistance from the Niigata Prefectural Office during the

tick sampling collection. We would also like to thank the alumni of MECOH Lab Ehime University

for their help in molecular analyses: Masaya Doi, Kohki Tanaka, and Mizuki Ueda. We are also grateful to Micanaldo Francisco for the construction of the map for this manuscript. We would like to thank Enago (www.enago.jp) for the English language review and proofreading of the manuscript.

References

1. Neumann M. Eine hysterische Hausepidemie. European Neurology. 1899;5(6):405-10.

2. Miyamoto, K., Nakao, M., Fujita, H., Sato, F., 1993. The ixodid ticks on migratory birds in Japan and the isolation of Lyme disease spirochetes from bird-feeding ticks. Med. Entomol. and zool., 44(4), 315-326. https://doi.org/10.7601/mez.44.315

3. Neumann H. Zur Behandlung der Pleuritis exsudativa. DMW-Deutsche Medizinische Wochenschrift. 1897 Jan;23(05):73-5.

4. Yu XJ, Liang MF, Zhang SY, Liu Y, Li JD, Sun YL, Zhang L, Zhang QF, Popov VL, Li C, Qu J. Fever with thrombocytopenia associated with a novel bunyavirus in China. N. Engl. J. Med. 2011 Apr 21;364(16):1523-32. https://doi.org/10.1056/nejmoa1010095

5. Yamaji K, Aonuma H, Kanuka H. Distribution of tick-borne diseases in Japan: Past patterns and implications for the future. J. Infect. Chemother.2018 Jul 1;24(7):499-504. https://doi.org/10.1016/j.jiac.2018.03.012

6. Arai R, Sato M, Kato M, Aoki J, Nishida A, Watanabe K, Hirokawa C, Ikeda S, Watanabe K, Regilme MA, Sato MO. Spotted fever group rickettsiae (SFGR) detection in ticks following reported human case of Japanese spotted fever in Niigata Prefecture, Japan. Sci. Rep. 2021 Jan 28;11(1):2595. https://doi.org/10.1038/s41598-021-81587-9

7. McCoy KD, Boulinier T, Tirard C, Michalakis Y. Host specificity of a generalist parasite: genetic

evidence of sympatric host races in the seabird tick *Ixodes uriae*. J. Evol. Biol.. 2001 May

9;14(3):395-405. https://doi.org/10.1046/j.1420-9101.2001.00290.x

8. Oliver JD, Bennett SW, Beati L, Bartholomay LC. Range expansion and increasing *Borrelia*

burgdorferi infection of the tick *Ixodes scapularis* (Acari: Ixodidae) in Iowa, 1990–2013. J. Med.

Entomol. 2017 Nov 7;54(6):1727-34. https://doi.org/10.1093/jme/tjx121

9. Agany, D. D., Pietri, J. E., Gnimpieba, E. Z., 2020. Assessment of vector-host-pathogen

relationships using data mining and machine learning. Comput. Struct. Biotechnol. J., 18,

1704-1721. https://doi.org/10.1016/j.csbj.2020.06.031

10. Araya-Anchetta A, Busch JD, Scoles GA, Wagner DM. Thirty years of tick population genetics:

a comprehensive review. Infect. Genet. Evol. 2015 Jan 1;29:164-79.

https://doi.org/10.1016/j.meegid.2014.11.008

11. Regilme MA, Sato M, Tamura T, Arai R, Sato MO, Ikeda S, Gamboa M, Monaghan MT,

Watanabe K. Comparative population genetic structure of two ixodid tick species (Acari:

Ixodidae)(*Ixodes ovatus* and *Haemaphysalis flava*) in Niigata prefecture, Japan. Infect. Genet.

Evol. 2021 Oct 1;94:104999.

12. Estrada-Peña AJ. Ticks as vectors: taxonomy, biology and ecology. Rev. Sci. Tech. 2015 Apr

1;34(1):53-65. https://doi.org/10.20506/rst.34.1.2345.

13. Mixson TR, Lydy SL, Dasch GA, Real LA. Inferring the population structure and demographic

history of the tick, *Amblyomma americanum* Linnaeus. J. Vector Ecol. 2006 Jun;31(1):181-92.

https://doi.org/10.3376/1081-1710(2006)31[181:itpsad]2.0.co; 2.

14. Trout RT, Steelman CD, Szalanski AL. Population genetics of *Amblyomma americanum* (Acari:

Ixodidae) collected from Arkansas. J. Med. Entomol. 2014 Dec 1;47(2):152-61.

https://doi.org/10.1603/me09106

15. Guglielmone AA, Nava S, Mastropaolo M, Mangold AJ. Distribution and genetic variation of

Amblyomma triste (Acari: Ixodidae) in Argentina. Ticks Tick Borne Dis. 2013 Sep 1;4(5):386-90.

https://doi.org/10.1016/j.ttbdis.2013.01.009

16. Lampo M, Rangel Y, Matat A. Population genetic structure of a three-host population genetic

in *Amblyommma dissimile*. J. Parasitol. 2015;84:1137-42.

17. Dougherty ER, Seidel DP, Carlson CJ, Spiegel O, Getz WM. Going through the motions:

incorporating movement analyses into disease research. Ecol. Lett. 2018 Apr;21(4):588-604.

https://doi.org/10.1111/ele.12917

18. Regoes RR, Nowak MA, Bonhoeffer S. Evolution of virulence in a heterogeneous host

population. Evol. 2000 Feb;54(1):64-71. https://doi.org/10.1111/j.0014-3820.2000.tb00008.x

19. Boots M, Mealor M. Local interactions select for lower pathogen infectivity. Sci. 2007 Mar

2;315(5816):1284-6.https://doi.org/10.1126/science.1137126

20. Lion S, Gandon S. Evolution of spatially structured host–parasite interactions. Journal of Evol.

Biol. 2015 Jan;28(1):10-28. https://doi.org/10.1111/jeb.12551

21. Chin KM, Wolfe MS. Selection on *Erysiphe graminis* in pure and mixed stands of barley. Plant

Pathol. 1984;33:535-546. doi: 10.1111/j.1365-3059.1984.tb02878.x

22. Thrall PH, Burdon JJ. Evolution of virulence in a plant host-pathogen metapopulation. Science.

2003;299:1735-1737. doi: 10.1126/science.1080070.

23. White PS, Choi A, Pandey R, Menezes A, Penley M, Gibson AK, de Roode J, Morran L. Host

heterogeneity mitigates virulence evolution. Biol Lett. 2020;16:20190744. doi:

10.1098/rsbl.2019.0744.

24. Gibson AK, Baffoe‐Bonnie H, Penley MJ, Lin J, Owens R, Khalid A, Morran LT. The evolution

of parasite host range in heterogeneous host populations. J Evol Biol. 2020;33:773-782. doi:

10.1111/jeb.13608.

25. Bono LM, Gensel CL, Pfennig DW, Burch CL. Competition and the origins of novelty:

experimental evolution of niche-width expansion in a virus. Biol Lett. 2013;9:20120616. doi:

10.1098/rsbl.2012.0616.

26. Streicker DG, Winternitz JC, Satterfield DA, Condori-Condori RE, Broos A, Tello C, Recuenco S,

Velasco-Villa A, Altizer S, Valderrama W. Host–pathogen evolutionary signatures reveal

dynamics and future invasions of vampire bat rabies. Proc Natl Acad Sci U S A.

2016;113:10926-10931. doi: 10.1073/pnas.1606587113.

27. Fountain‐Jones NM, Craft ME, Funk WC, Kozakiewicz C, Trumbo DR, Boydston EE, Lyren LM,

Crooks K, Lee JS, VandeWoude S, Carver S. Urban landscapes can change virus gene flow and

evolution in a fragmentation‐sensitive carnivore. Mol Ecol. 2017;26:6487-6498. doi:

10.1111/mec.14375.

28. Daniel M, Malý M, Danielová V, Kříž B, Nuttall P. Abiotic predictors and annual seasonal

dynamics of *Ixodes ricinus*, the major disease vector of Central Europe. Parasit Vectors.

2015;8:1-12. doi: 10.1186/s13071-015-1092-y.

29. Eisen RJ, Eisen L, Beard CB. County-scale distribution of *Ixodes scapularis* and *Ixodes pacificus*

(Acari: Ixodidae) in the continental United States. J Med Entomol. 2016;53:349-386. doi:

10.1093/jme/tjv237.

30. Chapman MA, Hiscock SJ, Filatov DA. Genomic divergence during speciation driven by

adaptation to altitude. Mol Biol Evol. 2013;30:2553–2567. doi: 10.1093/molbev/mst168.

31. Osborne OG, Batstone TE, Hiscock SJ, Filatov DA. Rapid speciation with gene flow following

the formation of Mt. Etna. Genome Biol Evol. 2013;5:1704–1715. doi: 10.1093/gbe/evt127.

32. Polato NR, Gray MM, Gill BA, Becker CG, Casner KL, Flecker AS, Kondratieff BC, Encalada AC,

Poff NL, Funk WC, Zamudio KR. Genetic diversity and gene flow decline with elevation in

montane mayflies. Heredity. 2017 Aug;119(2):107-116. doi: 10.1038/hdy.2017.19. PMID:

28333156.

33. Wang ,J., Summers,K., 2010. Genetic structure is correlated with phenotypic divergence

rather than geographic isolation in the highly polymorphic strawberry poison-dart frog. Mol.

Ecol., 19, 447–458. https://doi.org/10.1111/j.1365-294x.2009.04465.x

34. Bradburd, G.S., Ralph, P.L., Coop, G.M., 2013. Disentangling the effects of geographic and

ecological isolation on genetic differentiation. Evolution, 67, 3258–3273.

https://doi.org/10.1111/evo.12193

35. Sexton,J.P., Hangartner,S.B., Hoffmann,A.A., 2014. Genetic isolation by environment or

distance: which pattern of gene flow is most common?. Evol., 68, 1–15.

https://doi.org/10.1111/evo.12258

36. Yamaguti, N., Tipton, V. J., Keegan, H. L., & Toshioka, S. (1971). Ticks of Japan, Korea, and the

Ryukyu islands. Brigham Young University Science Bulletin, Biological Series, 15, 1.

37. Vrijenhoek R. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I

from diverse metazoan invertebrates. Mol Mar Biol Biotechnol. 1994 Oct 1;3(5):294-9. PMID:

7881515.

38. Maddison, W.P., Maddison, D.R., 2011. Mesquite: A modular system for evolutionary analysis.

508   Available at: http://mesquiteproject.org.

509   39. Rozas J, Ferrer-Mata A, Sánchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-Onsins SE,

510   Sánchez-Gracia A. DnaSP 6: DNA sequence polymorphism analysis of large datasets. Mol Biol

511   Evol. 2017 Dec 1;34(12):3299-3302. doi: 10.1093/molbev/msx248. PMID: 29036515.

512   40. Excoffier, L., & Lischer, H. E. L. (2010). Arlequin suite ver 3.5: A new series of programs to

513   perform population genetics analyses under Linux and Windows. Molecular Ecology

514   Resources, 10(3), 564-567. https://doi.org/10.1111/j.1755-0998.2010.02847.x

515   41. Paradis, E., Schliep, K., 2018. APE 5.0: an environment for modern phylogenetics and

516   evolutionary analyses in R. Bioinformatics, 35, 526–528.

517   https://doi.org/10.1093/bioinformatics/bty633.

518   42. Bandelt, H.J., Forster, P., Röhl, A., 1999. Median-joining networks for inferring intraspecific

519   phylogenies. Mol. Biol. Evol. 16, 37–48.

520   https://doi.org/10.1093/oxfordjournals.molbev.a026036

521   43. Drummond AJ, Rambaut A. BEAST: Bayesian evolutionary analysis by sampling trees. BMC

522   Evol Biol. 2007 Nov 8;7:214. doi: 10.1186/1471-2148-7-214. PMID: 17996036; PMCID:

523   PMC2222611.

524   44. Burri, C., Cadenas, F.M., Douet, V., Moret, J., Gern, L., 2007. Ixodes ricinus density and

525   infection prevalence of *Borrelia burgdorferi* sensu lato along a north-facing altitudinal

526   gradient in the Rhône Valley (Switzerland). Vector-Borne and Zoonotic Diseases, 7, 50-58.

527   https://doi.org/10.1089/vbz.2006.0569.

45. Jouda, F., Perret, J.L., Gern, L., 2004. *Ixodes ricinus* density, and distribution and prevalence of

Borrelia burgdorferi sensu lato infection along an altitudinal gradient. J. Med. Entomol. 41,

162-169. doi: 10.1603/0022-2585-41.2.162.

46. Moran-Cadenas F, Schneider H, Lommano E, Burri C, Moret J, Gern L. A comparison of two

DNA extraction approaches in the detection of *Borrelia burgdorferi* sensu lato from live

Ixodes ricinus ticks by PCR and reverse line blotting. Vector Borne Zoonotic Dis. 2007 Dec

1;7(4):555-62. https://doi.org/10.1089/vbz.2006.0596

47. Gooding, R.H. Genetic variation in arthropod vectors of disease-causing organisms: Obstacles

and opportunities. Clin. Microbiol. Rev. 1996; 9: 301.https://doi.org/10.1128/cmr.9.3.301

48. Páez-Triana, L., Muñoz, M., Herrera, G., Moreno-Pérez, D.A., Tafur-Gómez, G.A., Montenegro,

D., Patarroyo, M.A., Paniz-Mondolfi, A., Ramírez, J.D., 2021. Genetic diversity and population

structure of *Rhipicephalus sanguineus* sensu lato across different regions of Colombia. Parasit

Vectors, 14, 1-11. https://doi.org/10.1186/s13071-021-04898-w

49. Reichard, M.V., Kocan, A.A., Van Den Bussche, A., Barker, R.W., Wyckoff, J., Ewing, S.A., 2005.

Sequence variation of the ribosomal DNA second internal transcribed spacer region in two

spatially distinct populations of *Amblyomma americanum* (L.)(Acari: Ixodidae). J. Parasitol. 91,

260–264. https://doi.org/10.1645/GE-292R1.

50. Trout, R.T., Steelman, C.D., Szalanski, A.L., 2010. Population genetics of *Amblyomma*

americanum (Acari: Ixodidae) collected from Arkansas. J. Med. Entomol. 47, 152–161.

https://doi.org/10.1603/me09106.

51. Mixson TR, Lydy SL, Dasch GA, Real LA. Inferring the population structure and demographic

history of the tick, *Amblyomma americanum* Linnaeus. J Vector Ecol. 2006 Jun;31(1):181-92.

doi: 10.3376/1081-1710(2006)31[181:itpsad]2.0.co; 2.

52. Casati, S., Bernasconi, M.V., Gern, L., Piffaretti, J.C., 2008. Assessment of intraspecific mtDNA

variability of European *Ixodes ricinus* sensu stricto (Acari: Ixodidae). Infect. Genet. Evol. 8,

152–158. https://doi.org/10.1016/j.meegid.2007.11.007

53. Hasle, G., 2013. Transport of ixodid ticks and tick-borne pathogens by migratory birds. Front.

Cell. Infect. Microbiol., 3, 48. https://doi.org/10.3389/fcimb.2013.00048

54. Eremeeva, M.E. and Dasch, G.A., 2015. Challenges posed by tick-borne rickettsiae: eco-

epidemiology and public health implications. Front. Public Health. 3, 55. doi:

10.3389/fpubh.2015.00055.

55. Gage, K.L., Burkot, T.R., Eisen, R.J., Hayes, E.B., 2008. Climate and vector-borne diseases. Am.

J. Prev. Med. 35, 436-50. doi: 10.1016/j.amepre.2008.08.030.

56. Parola, P., Raoult, D., 2001. Tick-borne bacterial diseases emerging in Europe. Clin. Microbiol.

Infect. 7,80-83. https://doi.org/10.1046/j.1469-0691.2001.00200.x

57. Van Treuren W, Ponnusamy L, Brinkerhoff RJ, Gonzalez A, Parobek CM, Juliano JJ, et al.

Variation in the microbiota of Ixodes ticks with regard to geography, species, and sex. Appl

Environ Microbiol. 2015;81(18):6200-9. https://doi.org/10.1128/AEM.01562-15.

58. Randolph SE, Miklisová D, Lysy J, Rogers DJ, Labuda M. Incidence from coincidence: patterns

of tick infestations on rodents facilitate transmission of tick-borne encephalitis virus.

Parasitology. 1999;118:177–186. https://doi.org/10.1017/s0031182098003643.

59. MacDonald AJ, Hyon DW, Iii JBB, Connor KEO, Swei A, Briggs CJ. Lyme disease risk in southern

California: abiotic and environmental drivers of *Ixodes pacificus* (Acari: Ixodidae) density and

infection prevalence with *Borrelia burgdorferi*. Parasit Vectors. 2017 Jan;10(1):7.

https://doi.org/10.1186/s13071-016-1938-y.

60. Funk WC, Murphy MA, Hoke KL, Muths E, Amburgey SM, Lemmon EM, Lemmon AR.

Elevational speciation in action? Restricted gene flow associated with adaptive divergence

across an altitudinal gradient. J Evol Biol. 2016 Feb;29(2):241-252.

https://doi.org/10.1111/jeb.12760.

61. Coyne JA, Orr HA. Speciation. Sunderland, MA: Sinauer Associates; 2004.

62. Nosil P. Ecological Speciation. Oxford University Press; 2012.

63. Lim MC, Bi K, Witt CC, Graham CH, Dávalos LM. Pervasive genomic signatures of local

adaptation to altitude across highland specialist Andean hummingbird populations. J Hered.

2021 Mar 1;112(2):229-240. https://doi.org/10.1093/jhered/esab008.

64. Juvenal N, Edward M. Seasonal dynamics and distribution of ticks in Rwanda: implications for

tick control strategy in Rwanda. J Anim Vet Adv. 2010;9(1):21-25.

65. Nzige T. Ecologie des tiques du bétail. Rapport de stage, Higher Institute of Agriculture and

Animal Husbandry (ISAE-Busogo) Publications, Musanze, Rwanda. 1978.

66. Morel P. Maladies à tiques du bétail en Afrique In: Précis de parasitologie vétérinaire

tropicale. Ministére de la Coopération et du Développement, République Française, Norval,

R.A.I., Donachie, pp: 471-717. 1981.

67. Kawecki TJ, Ebert D. Conceptual issues in local adaptation. Ecol Lett. 2004 Dec;7(12):1225-

1241. https://doi.org/10.1111/j.1461-0248.2004.00684.x.

68. Gregory TR. Understanding natural selection: essential concepts and common

misconceptions. Evol Educ Outreach. 2009 Jun;2(2):156-175.

https://doi.org/10.1007/s12052-009-0128-1.

69. Wang IJ, Bradburd GS. Isolation by environment. Mol. Ecol. 2014 Dec;23(23):5649-5662.

https://doi.org/10.1111/mec.12938.

70. Jiang S, Luo MX, Gao RH, Zhang W, Yang YZ, Li YJ, Liao PC. Isolation-by-environment as a

driver of genetic differentiation among populations of the only broad-leaved evergreen shrub

Ammopiptanthus mongolicus in Asian temperate deserts. Sci Rep. 2019 Jul 30;9(1):1-14.

https://doi.org/10.1038/s41598-019-48472-y.

71. Yamauchi T, Takeno K. Record of *Haemaphysalis flava* Neumann (Acari,Ixodidae) from

Zoothera dauma (Latham) (Aves) in Kyushu, Japan. Esakia. 2000;40:99-101.

https://doi.org/10.5109/2647

72. Ishiguro F, Takada N, Fujita H, Noji Y, Yano Y, Iwasaki H. Survey of the vectorial competence

of ticks in an endemic area of spotted fever group rickettsioses in Fukui Prefecture, Japan.

Microbiol Immunol. 2008;52:305-9. https://doi.org/10.1111/j.1348-0421.2008.00042.x

73. Sato M, Ikeda S, Arai R, Kato M, Aoki J, Nishida A, et al. Diversity and distribution of ticks in

Niigata prefecture, Japan (2016–2018): Changes since 1950. Ticks Tick Borne Dis.

2021;12:101683. https://doi.org/10.1016/j.ttbdis.2021.101683.

74. Parola, P., Paddock, C.D., Socolovschi, C., Labruna, M.B., Mediannikov, O., Kernif, T., Abdad,

M.Y., Stenos, J., Bitam, I., Fournier, P.E., Raoult, D., 2013. Update on tick-borne rickettsioses

around the world: a geographic approach. Clin. Microbiol. Rev., 26, 657-702. doi:

10.1128/cmr.00032-13.

75. Gillespie JJ, Joardar V, Williams KP, Driscoll T, Hostetler JB, Nordberg E, Shukla M, Walenz B,

Hill CA, Nene VM, Azad AF. A *Rickettsia* genome overrun by mobile genetic elements provides

insight into the acquisition of genes characteristic of an obligate intracellular lifestyle. J

Bacteriol. 2012 Jan 15;194(2):376-94. https://doi.org/10.1128/JB.06244-11

76. Clay K, Klyachko O, Grindle N, Civitello D, Oleske D, Fuqua C. Microbial communities and

interactions in the lone star tick, *Amblyomma americanum*. Mol Ecol. 2008 Oct;17(19):4371-

4381.

77. Budachetri K, Browning RE, Adamson SW, Dowd SE, Chao CC, Ching WM, Karim S. An insight

into the microbiome of the *Amblyomma maculatum* (Acari: Ixodidae). J Med Entomol. 2014

Jan;51(1):119-29. doi: 10.1603/me13139. PMID: 24605422.

78. Richmond, C.E., Breitburg, D.L., Rose, K.A., 2005. The role of environmental generalist species

in ecosystem function. Ecol. Model. 188, 279-295. doi: 10.1016/j.ecolmodel.2005.03.002.

79. Cheng, C., White, B.J., Kamdem, C., Mockaitis, K., Costantini, C., Hahn, M.W., Besansky, N.J.,

2012. Ecological genomics of *Anopheles gambiae* along a latitudinal cline: a population-

resequencing approach. Genetics, 190(4), 1417-1432.

https://doi.org/10.1534/genetics.112.139899

80. Kapun, M., Barrón, M.G., Staubach, F., Obbard, D.J., Wiberg, R.A.W., Vieira, J., Goubert, C.,

Rota-Stabelli, O., Kankare, M., Bogaerts-Márquez, M., Haudry, A. (2020). Genomic analysis of

European *Drosophila melanogaster* populations reveals longitudinal structure, continent-

wide selection, and previously unknown DNA viruses. Mol. Biol. Evol., 37, 2661-2678.

https://doi.org/10.1093/molbev/msaa120

81. Love RR, Steele AM, Coulibaly MB, Traore SF, Emrich SJ, Fontaine MC, Besansky NJ.

Chromosomal inversions and ecotypic differentiation in *Anopheles gambiae*: the perspective

from whole-genome sequencing. Mol Ecol. 2016 Dec;25(23):5889-906.

https://doi.org/10.1111/mec.13888

82. Exposito-Alonso M, Vasseur F, Ding W, Wang G, Burbano HA, Weigel D. Genomic basis and

evolutionary potential for extreme drought adaptation in *Arabidopsis thaliana*. Nat Ecol Evol.

2018 Feb;2(2):352-8., https://doi.org/10.1038/s41559-017-0423-0

83. Hancock AM, Brachi B, Faure N, Horton MW, Jarymowycz LB, Sperone FG, Toomajian C, Roux

F, Bergelson J. Adaptation to climate across the *Arabidopsis thaliana* genome. Science. 2011

Oct 7;334(6052):83-6.https://doi.org/10.1126/science.1209244

84. Zhen Y, Harrigan RJ, Ruegg KC, Anderson EC, Ng TC, Lao S, Lohmueller KE, Smith TB. Genomic

divergence across ecological gradients in the Central African rainforest songbird (*A ndropadus*

virens). Mol. Ecol. 2017 Oct;26(19):4966-77.https://doi.org/10.1111/mec.14270

85. Ahrens CW, Byrne M, Rymer PD. Standing genomic variation within coding and regulatory

regions contributes to the adaptive capacity to climate in a foundation tree species. Mol Ecol.

2019 May;28(10):2502-16. https://doi.org/10.1111/mec.15092

86. Miller AD, Hoffmann AA, Tan MH, Young M, Ahrens C, Cocomazzo M, Rattray A, Ierodiaconou

DA, Treml E, Sherman CD. Local and regional scale habitat heterogeneity contribute to

genetic adaptation in a commercially important marine mollusc (*Haliotis rubra*) from

southeastern Australia. Mol. Ecol. 2019 Jun;28(12):3053-72.

https://doi.org/10.1111/mec.15128

87. Crosbie PR, Boyce WM, Rodwell TC. DNA sequence variation in *Dermacentor hunteri* and

estimated phylogenies of *Dermacentor* spp.(Acari: Ixodidae) in the New World. J. Med.

Entomol. 1998 May 1;35(3):277-88. https://doi.org/10.1093/jmedent/35.3.277

88. Norris DE, Klompen JS, Keirans JE, Black IV WC. Population genetics of *Ixodes scapularis*

(Acari: Ixodidae) based on mitochondrial 16S and 12S genes. J. Med. Entomol.1996 Jan

1;33(1):78-89. https://doi.org/10.1093/jmedent/33.1.78

89. Qiu WG, Dykhuizen DE, Acosta MS, Luft BJ. Geographic uniformity of the Lyme disease
spirochete (*Borrelia burgdorferi*) and its shared history with tick vector (*Ixodes scapularis*) in
the northeastern United States. Genetics. 2002 Mar 1;160(3):833-49.
https://doi.org/10.1093/genetics/160.3.833

90. Rich SM, Caporale DA, Telford 3rd SR, Kocher TD, Hartl DL, Spielman A. Distribution of the
Ixodes ricinus-like ticks of eastern North America. Proceedings of the National Academy of
Sciences. 1995 Jul 3;92(14):6284-8. https://doi.org/10.1073/pnas.92.14.6284

91. de la Fuente J, Almazán C, Van Den Bussche RA, Bowman J, Yoshioka JH, Kocan KM.
Characterization of genetic diversity in *Dermacentor andersoni* (Acari: Ixodidae) with body
size and weight polymorphism. Exp. Parasitol. 2005 Jan 1;109(1):16-26.
https://doi.org/10.1016/j.exppara.2004.10.004

92. Caporale DA, Rich SM, Spielman A, Telford SR, Kocher TD. Discriminating between *Ixodes*
ticks by means of mitochondrial DNA sequences. Mol. Phylogenet. Evol. 1995 Dec 1;4(4):361-
5. https://doi.org/10.1006/mpev.1995.1033

93. Mitani H, Takahashi M, Masuyama M, Fukunaga M. *Ixodes philipi* (Acari: Ixodidae):
phylogenetic status inferred from mitochondrial cytochrome oxidase subunit I gene sequence
comparison. J. Parasitol. Res. 2007 Jun;93(3):719-22. https://doi.org/10.1645/ge-953r.1

94. Anderson JF, Magnarelli LA. Biology of ticks. Infectious disease clinics of North America. 2008
Jun 1;22(2):195-215.

95. Hill CA, Wikel SK. The *Ixodes scapularis* Genome Project: an opportunity for advancing tick
research. Trends Parasitol.. 2005 Apr 1;21(4):151-3.https://doi.org/10.1016/j.pt.2005.02.004

96. Van Zee JP, Geraci NS, Guerrero FD, Wikel SK, Stuart JJ, Nene VM, Hill CA. Tick genomics: the

Ixodes genome project and beyond. Int. J. Parasitol. 2007 Oct 1;37(12):1297-305.

https://doi.org/10.1016/j.ijpara.2007.05.011

97. Jongejan F, Nene V, de la Fuente J, Pain A, Willadsen P. Advances in the genomics of ticks and

tick-borne pathogens. Trends Parasitol. 2007 Sep 1;23(9):391-6.

https://doi.org/10.1016/j.pt.2007.07.004

98. Gray JS. Biology of Ixodes species ticks in relation to tick-borne zoonoses. Wiener Klinische

Wochenschrift. 2002 Jul 1;114(13-14):473-8.

99. Parola P, Paddock CD. Travel and tick-borne diseases: Lyme disease and beyond. Travel Med

Infect Dis. 2018;26:1-2. https://doi.org/10.1016/j.tmaid.2018.09.010

Supporting information

S1 Table. Summary of *Ixodes ovatus* and *Haemaphysalis flava* collected from the different

locations of Niigata Prefecture and its corresponding sample number and number of *Rickettsia*

infection per site

S2 Table. The difference in *Rickettsia* infection rates in *H. flava* haplotype groups and a table

showing the distribution of *Rickettsia*-infected and uninfected *H. flava* in each genetic groups as

shown in S2 Figure *(H. flava* median-joining network). The results of the z-score test for two

populations proportions at $p < 0.05$ showed no significant difference between the *Rickettsia*

detection rate in haplotype groups 1 and 2. The Welch t-test at $p < 0.05$ revealed no significant

difference in the mean altitude of the two groups.

S1 Fig. Phylogenetic tree from the BEAST analysis of 59 haplotype *cox1* sequences of *I. ovatus.* The

blue-labeled haplotypes indicate the presence of *Rickettsia* infection. The red parentheses provide

the number of *Rickettsia*-positive individuals per haplotype. The black labeled haplotypes are

negative for *Rickettsia* infection.

S2 Fig. Median-joining network of the 66 *cox1* haplotype sequences of *Rickettsia* positive and

negative *H. flava.* Haplotype groups are indicated as 1 and 2.

S3 Fig. Phylogenetic tree from the BEAST analysis of 66 haplotype *cox1* sequences of *H. flava.* The

blue-labeled haplotypes indicate *Rickettsia* infection in individual samples. The parentheses in red

provide the number of *Rickettsia*-infected ticks. The black-labeled haplotypes are negative for

Rickettsia infection.

Rickettsia (-)   Rickettsia (+)

Elevation
(meters)
High : 3039
Low : -10