Title

1   **Relationship of *Rickettsia* infection rates and population genetic structure**

2   **of ixodid ticks along an altitudinal gradient**

Author names and affiliations

Maria Angenica F. Regilme[a], Megumi Sato[b], Tsutomu Tamura[c], Reiko Arai[c], Marcello Otake Sato[d], Sumire Ikeda[e], Kozo Watanabe[a]*

[a] Center for Marine Environmental Studies, Ehime University, Matsuyama, Ehime 790-8577, Japan

[b] Graduate School of Health Sciences, Niigata University, Niigata, 951-8518, Japan

[c] Niigata Prefectural Institute of Public Health and Environmental Sciences, Niigata, 950-2144, Japan

[d] Department of Tropical Medicine and Parasitology, Dokkyo Medical University, 880 Kitakobayashi, Mibu-machi, Shimotsuga-gun, Tochigi 321-0293, Japan

[e] Research Laboratories, Research and Development Headquarters, Earth Corporation, Hyogo 678-0192, Japan

Corresponding author*

watanabe.kozo.mj@ehime-u.ac.jp

Center for Marine Environmental Studies, Ehime University, Bunkyo-cho 3, Matsuyama, Ehime, Japan 790-8577

Abstract

Ixodid ticks, such as *Ixodes ovatus* and *Haemaphysalis flava,* are important vectors of tick-

borne diseases in Japan, such as Japanese spotted fever caused by *Rickettsia sp* (i.e., the

pathogen). This study describes the population genetic structure and gene flow of *I. ovatus* and *H.*

flava and their *Rickettsia* infection rates along an altitudinal gradient. A total of 346 adult *I. ovatus*

and 243 *H. flava* were analyzed for the presence of *Rickettsia* using the primers 17kDA, *gltA,*

rOmpA, and *rOmpB.* Population genetic structure was performed using the mitochondrial

cytochrome oxidase 1 (*cox1*) marker. The *Rickettsia* infection rates were 13.26% in *I. ovatus* and

6.17% in *H. flava.* For *I. ovatus,* the global F_{ST} value revealed significant genetic differentiation

among the different populations, whereas *H. flava* showed non-significant genetic differentiation.

The *cox1 I. ovatus* cluster dendrogram showed two cluster groups, while the haplotype network

and phylogenetic tree showed three genetic groups. The two cluster groups and the three genetic

groups found in *I. ovatus* showed a significant difference in their *Rickettsia* infection rates and

mean altitude per group. No significant differences were found in the mean altitude or *Rickettsia*

infection rates of *H. flava.* Our results suggest that low gene flow in the *I. ovatus* populations have

caused spatially heterogenous *Rickettsia* infection rates along the altitudinal gradient. This

information is essential in understanding the relationship between the tick vector, its pathogen,

and environmental factors, such as altitude.

Keywords: environment, haplotype, hosts, pathogen, dispersal, vector

1. Introduction

Tick-borne diseases are a public health concern in Japan and are transmitted by a wide range of tick species, such as *Ixodes ovatus*, the primary vector of *Lyme borealis* and *Haemaphysalis flava*, a known vector of severe fever thrombocytopenia syndrome and Japanese spotted fever caused by the pathogen *Rickettsia japonica* (Mahara et al., 1997; Miyamoto et al., 1993; Yu et al., 2014, Yamaji et al., 2018, Arai et al., 2021). The dynamics of tick-borne pathogens are influenced by the habitat distribution and dispersal behaviors of vectors and hosts along environmental gradients (Agany et al., 2020; Jongejan et al., 2007).

Tick population genetic analysis provides dispersal pattern data based on gene flow between local populations (Arraya-Anchetta et al., 2015). Tick dispersal primarily depends on the movements of their vertebrate hosts, which influences each tick's potential to spread its pathogen (Falco and Fish, 1991; McCoy et al., 2001, Oliver et al., 2007). For example, contrasting patterns in the population genetic structures of *I. ovatus* and *H. flava* in the Niigata Prefecture of Japan indicate that host mobility during the immature stages of tick development may influence the genetic structure of adult ticks by affecting survivability into their adult stages (Regilme et al., 2021; Estrada-Pena et al., 2015). *I. ovatus* populations had greater genetic divergence due to the limited dispersal of their small mammalian hosts during the immature development stage; whereas *H. flava* populations showed a more homogenized structure due to the higher mobility of their large mammalian hosts and avian-mediated dispersal (Regilme et al., 2021). Other studies have also revealed low gene flow in ticks with low mobility hosts (e.g., small mammals) and higher gene flow in ticks with highly mobile hosts (e.g., large mammals and birds; Arraya-ancheta et al., 2015; Mixson et al., 2006; Trout et al., 2010; Guglielmone et al., 2013; Lampo et al., 2015).

The spatial distribution and movement of the vector (i.e., ticks) may determine the spatial

distribution of the pathogen's (i.e., *Rickettsia*) infection rate (Dougherty et al., 2018). Previous

studies have shown that pathogen infection rate can be influenced by many factors, such as vector

host genetic diversity, gene flow, and spatial structure (Regoes et al., 2000; Boots et al., 2007; Lion

et al., 2015; Chin et al., 1984, Thrall et al., 2003; White et al., 2020; Gibson et al., 2020; Bono et al.,

2012). For example, previous studies have shown that strong gene flow between local host

populations tend to reduce the spatial heterogeneity of pathogen infection rates between

populations (Streicker et al., 2016; Fountain-Jones et al., 2018). Thus, the exchange of infected and

uninfected individuals across a spatial scale is expected to spatially homogenize pathogen

infection rates. To our knowledge, no previous studies have examined the relationship between

the spatial heterogeneity of *Rickettsia* infection rates and population genetic structure in ticks.

Environmental factors, such as altitude, may relate to the population genetic structure of

ticks (Daniel et al., 2015; Eisen et al., 2016), with limited gene flow increasing genetic variation

between populations along an altitudinal gradient, as reported in several species (Hofreiter et al.,

2004; Chapman et al., 2013; Osborne et al., 2013, Polato et al., 2017). Thus, in this study, we

expected to see a highly divergent population genetic structure in *I. ovatus* along an altitudinal

gradient due to the limited movement of their mammalian hosts along that gradient; whereas *H.*

flava should show a less divergent structure along the altitudinal gradient due to the higher

mobility of its hosts. To our knowledge, no studies have focused on tick gene flow along an

altitudinal gradient.

In the study by Regilme et al. (2021), no significant influence of environmental factors,

including altitude, were observed in the genetic structures of *I. ovatus* and *H. flava* based on the

mantel test, but the study did not use any other robust analytical methods to more thoroughly

examine the influence of altitude on tick genetic structure. In another study, the relationship

between the tick vector and its pathogen (*Rickettsia* sp.) were analyzed in a total of 3,336

immature and adult ticks across the Niigata Prefecture, Japan by analyzing *Rickettsia* infection

rates per species (Arai et al., 2021). Although *Rickettsia* infections were found in *H. flava,*

Haemaphysalis longicornis, Ixodes monospinus, Ixodes nipponensis, and *Ixodes ovatus*, no spatial

distribution of *Rickettsia* infection rates were found among the local populations. To our

knowledge, no previous studies have considered the influence of environmental factors on tick

population genetic structure or the spatial distribution of *Rickettsia* infection rates along an

altitudinal gradient in local tick populations.

78   Here, we determine the relationship between the vector tick population genetic structure

and *Rickettsia* infection rates as influenced by an altitudinal gradient. Based on the isolation by

environment (IBE) theory, genetic differentiation increases with environmental variation,

regardless of geographic distance (Wang and Summers et al., 2010; Bradburd et al., 2013; Sexton

et al., 2014). Thus, we hypothesized that limited gene flow among tick populations along an

altitudinal gradient might cause spatially heterogenous *Rickettsia* infection rates in ticks. The

present study used cytochrome oxidase 1 (*cox1*) mitochondrial gene sequences and *Rickettsia*-

infected/uninfected data for *I. ovatus* (n = 307) and *H. flava* (n = 220) individuals collected in

2016–2017 as reported in Regilme et al. (2021) and Arai et al. (2021). Additionally, we added new

cox1 sequences and *Rickettsia* infection/noninfection data from *I. ovatus* and *H. flava* individuals

and five other species collected in 2018.

2. Materials and Method

Published data of Regilme et al. (2021) and Arai et al. (2021)

In this study, we used *cox1* sequence data from Regilme et al. (2021) for *I. ovatus* (n = 307) and *H. flava* (n = 220) ticks collected from April 2016 to November 2017 from 29 sites across the Niigata Prefecture, Japan. *Rickettsia* infection rate data were obtained from Arai et al. (2021) from *I. ovatus* (n = 29), *H. flava* (n = 2), *I. monospinus* (n = 4), and *I. nipponensis* (n = 2) from 38 sites. Please refer to Regilme et al. (2021) and Arai et al. (2021) for more information about the study sites, collection, sampling identification, DNA extraction, PCR amplification, and sequencing methods used in each study.

To strengthen our analysis, we also added data collected from April to October 2018 from additional individuals sampled at 31 sites across the Niigata Prefecture, including two sites not previously sampled by Regilme et al. (2021). At these sites, ticks were collected 2–14 times from six core sites among the 31 sites while the remaining sites, ticks were collected once. The altitude at each site ranged from 8 to 1402 m.a.s.l., with a mean altitude of 348 m.a.s.l. Multiple sites were combined for population genetic analysis if less than eight individuals were obtained per site, which resulted in 8 populations labeled A to H. Three sites were excluded from the population genetic analysis because of the limited number of obtained individuals (<8) and the lack of a nearby site to combine into a single population.

Unpublished data from 2018 collection

Ticks collected in 2018 were stored at 4 °C in microcentrifuge tubes with 70% ethanol. Each collected tick was morphologically identified using a stereo microscope following the

identification keys of Yamaguti et al. (1971). Genomic DNA was extracted using Isogenome DNA

extraction kits (Nippon Gene Co. Ltd. Tokyo, Japan) following the manufacturer's recommended

protocol.

In this study, we combined previously published data from Arai et al. (2021) with our

newly collected data to calculate the *Rickettsia* infection rate, which is the percentage of

Rickettsia-infected ticks from each obtained population. We analyzed the obtained tick DNA for

spotted fever group *Rickettsia* (SFGR) detection and host identification, and amplified the

mitochondrial gene *cox1* for population genetic analysis. We performed nested PCR targeting the

following genes for the detection of *Rickettsia* sp.: 17-kDA antigen gene (17-kDA); citrate synthase

gene (gltA); spotted fever group (SFG)-specific outer membrane protein A gene (rOmpA); and

outer membrane protein B gene (rOmpB) as described and analyzed in Arai et al. (2021). Briefly,

we first amplified the 17-kDa protein. If the results were positive, then PCR was performed to

target gltA. Samples that were positive with both 17-kDA and gltA were regarded as positive for

SFGR. PCR was also performed to target the rOmpA and rOmpB genes to identify *Rickettsia* species.

The amplified PCR products were purified using AMPure XP (Beckman Coulter Co., Japan) and

sequenced using the Big Dye Terminator Cycle Sequence Kit (Thermo Fisher Scientific).

The *cox1* mitochondrial gene was amplified by PCR for *cox1* (658 base pairs) using the

primer pairs LCO-1490 (5ꞌ- AAACTTCAGGGTGACCAAAAAATCA-3') and HCO1−2198 (5ꞌ–

AAACTTCAGGGTGACCAAAAAATCA - 3) for phylogenetic analysis and tick species identification

(Folmer et al., 1994). The PCR amplification profile included an initial denaturation of 94 °C for

2 min, followed by denaturation at 94 °C for 30 s, then annealing at 38 °C for 30 s, followed by an

extension of 72 °C for 1 min for 30 cycles, and a final extension of 72 °C for 10 min. The obtained

PCR products were purified using the QIAquick 96 PCR Purification Kit (Qiagen, Germany) following

the manufacturer's instructions and were sequenced by Eurofin Genomics, Inc. (Tokyo, Japan).

Each forward and reverse read was assembled using CodonCode Aligner version 1.2.4

software (https://www.codoncode.com/aligner/). Low-quality bases were removed in the aligned

sequences, and no ambiguous bases were detected. We used the MAFFT alignment online

program (https://mafft.cbrc.jp/alignment/server/) to perform multiple alignments using the

default settings. The sequences were checked for similarities with the deposited reference

sequences from GenBank for sequence quality and tick species confirmation using BLAST

(https://blast.ncbi.nlm.nih.gov/Blast.cgi). All final aligned sequences were checked in Mesquite

version 3.5 (Maddison et al., 2011). The protein-coding genes were translated to amino acids to

confirm the absence of stop codons.

Population genetic analysis

The final *cox1* sequences from all seven tick species were individually analyzed using

DNASp version 6.12.03 to determine the haplotype diversity per species (Rozas et al., 2017). The

level of genetic divergence between each population was quantified per species using global F_{ST}.

Significance was tested using Arlequin software version 3.5.2.2 (Excoffier and Lischer, 2010) with

9999 permutations.

The genetic relationship between the *I. ovatus* populations was visualized using the

unweighted pair group with the arithmetic mean (UPGMA) cluster method using the APE package

(Paradis and Schliep et al., 2018) for the RStudio software (R Development Core Team, 2016). A

cluster dendrogram was created using pairwise F_{ST} values genetic distance matrix from GenAlEx.

Haplotype network and phylogenetic analyses

We constructed a haplotype network analysis using PopART program version 1.7

(http://popart.otago.ac.nz/index.shtml) on *cox1 I. ovatus* and *H. flava* sequences to assess

haplotype relationships using the median-joining network algorithm (Bandeth et al., 1999). Briefly,

we constructed a Bayesian phylogenetic tree of *cox1* haplotypes for *I. ovatus* and *H. flava,*

respectively, using BEAST version 1.10.14 (Drummond and Rambaut, 2007). We used the

Hasegawa-Kishino-Yano substitution model with estimated base frequencies. We employed a

strict clock model and used the coalescent prior as the tree prior. A maximum clade credibility tree

was acquired using TreeAnnotator version 1.10.14 using trees from BEAUti version1.10.14 with

90% of the trees as the burn-in. We viewed the constructed maximum clade credibility tree using

FigTree version1.4.4.

Statistical analysis

To determine whether there was a significant difference in the *Rickettsia* infection rate

between haplotype groups for *I. ovatus* and *H. flava),* we performed a z-score test at $p < 0.05$. To

determine whether there were differences in the mean altitude between the haplotype groups,

we used the Welch t-test at $p < 0.05$.

3. Results

The total number of positive (pos) and negative (neg) ticks for *Rickettsia* infections from

the ticks collected in 2018 were: *I. ovatus* (neg = 22, pos = 17); *H. flava* (neg = 10, pos = 13); *I.*

monospinus (neg = 11, pos = 6); *I. asanumai* (neg = 4, pos=5); *I. nipponensis* (pos = 4); *I. persulcatus*

(neg = 4); and *H. japonica* (pos = 2). In this study, the total number of samples from the previously

published data (Regilme et al., 2021; Arai et al., 2021) and the unpublished data from 2018 were: *I.*

ovatus (n = 347); *H. flava* (n = 243); *I. monospinus* (n = 19); *I. asanumai* (n = 9); *I. nipponensis*

(n = 4); *I. persulcatus* (n = 4); and *H. japonica* (n = 2). The number of adult ticks whose *cox1* was

successfully sequenced per species were: *I. ovatus* (346); *H. flava* (243); *I. monospinus* (19); *I.*

asanumai (9); *I. nipponensis* (4); *I. persulcatus* (4); and *H. japonica* (2) (Table 1).

We detected SFGR in 78 (12.44%) out of 627 ixodid ticks, with the highest detected in *I.*

ovatus (46/346; 13.29%), *H. flava* (15/243; 6.17%), and *I. monospinus* (6/19; 31.58%) as

summarized in Table 1. Out of the 46 *Rickettsia*-infected *I. ovatus* ticks, 25 displayed a 100%

identity-match with *Rickettsia asiatica* in the 17kDA, *gltA,* and *rOmpB* markers (Arai et al., 2021),

while an additional 19 adult *I. ovatus* from the 2018 collection were also positive with *R. asiatica.*

Two haplotypes were found in the rOmpB and 17kDA markers, respectively. One haplotype was

found in only one individual (17369). Two out of the 15 *Rickettsia*-infected *H. flava* ticks were

found to have the same haplotypes in the 17Kda, gltA,and rOmpA markers, and were identified as

Rickettsia sp. (LC461063). The remaining 13 *Rickettsia*-infected *H. flava* ticks were identified as

Rickettsia sp. Four out of the six *Rickettsia*-infected *I. monospinus* were identified as *Rickettsia*

helvetica with the same haplotype found in the 17kDA, gltA, and rOmpB markers.

Based on the population genetic analysis of the *cox1* sequences, we found a significant

global F_{ST} of 0.4154 at $p < 0.05$ for *I. ovatus* (Table 1). In contrast, no significant global F_{ST} values

were observed in the *H. flava, I. monospinus, I. asanumai, I. nipponensis* or *I. persulcatus* samples

(Table 1). There were 59 and 66 *cox1* haplotypes found among the 346 *I. ovatus* and 243 *H. flava*

individuals, respectively. We found the following number of *cox1* haplotypes in the remaining

species: *I. monospinus* (n = 9), *I. asanumai* (n = 4), *I. nipponensis* (n = 4), *I. persulcatus* (n = 3), and

H. japonica (n = 2).

The *cox1* haplotype network of *I. ovatus* (Figure 1) revealed four genetic groups, wherein

three genetic groups (1, 2, and 3) were distributed along different altitudinal gradients, as shown

in Figure 2. These four genetic groups were concordant with the four clusters found in the *I.*

ovatus phylogenetic tree (Supplementary Figure 1). The habitat distribution of genetic group 3 was

limited to high altitude sites only (range = 16–912 m.a.s.l.), whereas genetic groups 1 (255–

471 m.a.s.l.) and 2 (84–1354 m.a.s.l.) were distributed at lower altitudes (Figure 2). The Welch t-

test revealed a significant difference between the mean altitudes of genetic groups 1 and 3 at

$p < 0.05$ (Table 2); however, no significant difference was observed between genetic groups 1 and

2 or groups 2 and 3. We found a significant difference in the *Rickettsia* infection rates between *I.*

ovatus genetic groups 1 and 2 based on the z-score test, but no significant difference between

groups 1 and 3 or groups 2 and 3 (Table 2). The mean altitude between the *Rickettsia*-infected

(=273.72 m.a.s.l.) and non-infected *I. ovatus* (=369.61 m.a.s.l.) revealed a significant difference

based on the Welch t-test at $p < 0.05$ (Figure 3). The UPGMA dendrogram of *I. ovatus* revealed two

genetic clusters, 1 and 2 using the genetic distance among the seven populations excluding one

population due to the limited number of samples (Figure 4).

The *cox1* haplotype network of *H. flava* displayed two genetic groups (Supplementary

Figure 2) consistent with the *H. flava* phylogenetic tree (Supplementary Figure 3). Based on the

Welch t-test, we found no significant differences in mean altitude (205 and 165 m.a.s.l.) at $p < 0.05$

between the two *H. flava* genetic groups (Supplementary Table 1). No significant difference

between the *Rickettsia* infection rates of *H. flava* genetic groups 1 and 2 were observed using the

z-score test at $p < 0.05$ (Supplementary Table 1).

4. Discussion

Our findings support our hypothesis that a genetically structured tick population, such as *I. ovatus*, can cause the *Rickettsia* infection rate to be spatially heterogenous due to limited gene flow along an altitudinal gradient. The significant global F_{ST} estimate of 0.4154 among the *I. ovatus* populations revealed genetic differentiation between the populations as supported by the occurrence of two genetic clusters in the cluster dendrogram. Our results were consistent with our previous study (Regilme et al., 2021) that suggested that the low mobility of the host species for immature *I. ovatus* contributed to low gene flow in the tick populations. The low *I. ovatus* gene flow along the altitudinal gradient might have caused the spatial heterogeneity of *Rickettsia* infection rates among these populations, which is supported by the significant difference found in *Rickettsia* infection rates between genetic clusters 1 and 2. A similar pattern was observed in the studies of Burri et al. (2007), Jouda et al. (2004), and Moran et al. (2007), which found that the infection rate of *Borrelia burgdorferi*, the causative agent of Lyme disease, decreased in ticks along the altitudinal gradient. Low gene flow can cause infected and uninfected ticks to have limited opportunities to traverse a wider spatial area thus causing heterogeneous *Rickettsia* infection rates (Gooding et al., 1996; Paez-Triana et al., 2021).

We found two genetic groups in the *H. flava* haplotype network, but no significant difference in the *Rickettsia* infection rates between the two groups. These results might be due to the high gene flow observed in the *H. flava* populations, which enable *Rickettsia*-infected and uninfected *H. flava* individuals to traverse between the study sites. The high mobility of the large mammalian hosts used by adult *H. flava* and avian-mediated dispersal during their immature stage probably contributed to their homogenized population genetic structure (Regilme et al., 2021), and the resulting homogenized *Rickettsia* infection rates. Large mammalian hosts and birds have a wide dispersal range that enable the broader movement of *Rickettsia*-infected ticks, as observed

in previous studies of *Amblyomma americanum* (Reichard et al., 2005; Trout et al., 2010; Mixson

et al., 2006), *H. flava* (Regilme et al., 2021), and *I. ricinus* (Casati et al., 2008). Birds are especially

good at dispersing over large areas, since they can easily traverse landscape barriers such as

mountains, fences, glaciers, and oceans that would be difficult for mammals to cross (Hasle et al.,

2013).

In addition to host mobility, environmental factors can influence the population genetic

structure of ticks (Eremeeva et al., 2015). Each tick species has preferred environmental conditions

that are conducive to completing the tick life cycle, thus influencing the geographical distribution

of ticks and the risk areas for tick-borne diseases (Gage et al., 2008; Parola et al., 2001). Altitude

may influence the population genetic structure of ticks through the effect of altitude on the

distribution and abundance of ticks and/or their hosts (Van Treuren et al., 2015; Randolph et al.,

1999; Macdonald et al., 2017). Altitudinal differences between populations can affect genetic

divergence (Funk et al., 2015), through, for example, ecological isolation, which causes natural

selection against maladapted immigrants and limits gene flow (Coyne and Orr, 2004; Nosil, 2012).

For example, organisms adapted to low altitudinal sites that cannot tolerate the lower

temperatures at higher altitudes would not survive if they dispersed to those higher altitude sites,

thus restricting gene flow across the altitudinal gradient (Funk et al., 2015)

The difference in mean altitude between *I. ovatus* cluster groups 1 and 2 might be due to

adaptive divergence along the altitudinal gradient. Individuals from cluster group 1 were

distributed in higher altitude areas of the northern mountainous area of Niigata Prefecture, while

cluster group 2 individuals were found in the lower altitude regions of northern Niigata Prefecture.

Populations along an altitudinal gradient are prone to differentiating selection pressures, which

result in local adaptation (Lim et al., 2021). Altitudinal gradients may also cause the varying

ambient temperature, precipitation and humidity levels that are essential to the development and

survival of ticks (Juvenal and Edward, 2010). These varying environmental factors may cause ticks

to have difficulty dispersing over a wide habitat range. Thus, ticks may need to adapt to extreme

habitats, such as extreme altitude or precipitation levels, for survival. For example, *Rhipicephalus*

compositus were found in altitudes of 1000–2500 m, but optimal conditions were between 1200–

2500 m (Juvenal and Edward, 2010; Nzige et al., 1978; Morel, 1981).

The different *Rickettsia* infection rates and altitudinal ranges between the *I. ovatus*

phylogenetic groups may be explained by adaptive evolutionary theory, which states that

organisms adjust to new or severe changes in their environment to become better suited to their

habitat (Kawecki and Ebert, 2004; Gregory, 2009). Based on the relationship between the *I. ovatus*

phylogenetic groups and their mean attitudes, *I. ovatus* might be undergoing local adaptation

along the altitudinal gradient due to the higher genetic differentiation between populations as

supported by the significant global F_{ST} (0.4154) found in *I. ovatus*. Based on isolation by

environment (IBE), genetic differentiation will increase with increased environmental differences

independent of geographic distances (Wang and Summers, 2010; Bradburd et al., 2013; Wang and

Bradburd, 2014). When environmental conditions differ, the success of immigrations in a new

habitat is reduced, which may increase the genetic fixation rate due to a lower chance of

outcrossing; thereby enhancing genetic isolation (Jiang et al., 2019). Thus, the lower gene flow

along the altitudinal gradient reduced the spatial homogeneity of *Rickettsia* infection rates among

the *I. ovatus* tick populations, thus causing the different *Rickettsia* infection rates obtained.

In this study, only a few (n < 20) *I. monospinus, I. asanumai, I. nipponensis* and *H. japonica*

were collected; however a high *Rickettsia* infection rate (31.58%–100%) was found despite these

limited numbers. In previous studies, high *Rickettsia* infection rates despite few individuals were

also observed in *I. monospinus* in several prefectures in Japan (Yamauchi et al., 2009; Ishiguro et

al., 2008; Sato et al., 2021). The extent of tick habitat distribution may vary between species,

which may have influenced the widely different *Rickettsia* infection rates observed among the

species in this study. The high Rickettsia infection rate in these ticks is probably due to effective

transovarial transmission (Parola et al., 2013). Additionally, the few collected individuals of these

species may be habitat specialists that thrive in a narrow habitat range in contrast to *I. ovatus* and

H. flava, which may be habitat generalists capable of thriving in a wide habitat range (Richmond et

al., 2005). The narrow habitat range of habitat specialist ticks might have enabled increased

interactions between infected and uninfected *Rickettsia* ticks within the small spatial scale

compared to the habitat generalist ticks, thus causing their high *Rickettsia* infection rates. This

conclusion should be considered with caution, since the limited number of individual ticks

collected across the Niigata Prefecture may have caused a bias in the estimated *Rickettsia*

infection rates determined in this study. We suggest conducting future research of *Rickettsia*

infection rates in a wider sampling range with an increased number of *I. monospinus, I. asanumai,*

I. nipponensis, and *H. japonica* individuals to further increase our understanding of the disease-

carrying potential of these species.

The occurrence of local adaptation in tick populations could affect the future of the tick-

borne disease landscape. Environmental factors, such as precipitation, temperature, and altitude,

have been shown to drive population differentiation in insects, such as *Anopheles* mosquitoes and

Drosophila flies (Cheng et al., 2012; Kapun et al., 2016; Love et al., 2016), but studies on the

environmental adaptation of ixodid ticks, such as *I. ovatus*, and its *Rickettsia* infection

susceptibility, have not yet been performed. McCoy et al. (2001) suggested that environmental

conditions that affect bird hosts can also affect the local adaptation of ticks. Few studies have

assessed such local adaptation in multiple organisms with varying dispersal abilities (Exposito et al.,

2018, Hancock et al., 2011, Oliviera et al., 2018, Zhen et al., 2017, Ahrens et al., 2019 Miller et al.,

2019) and is an area in need of future research.

Since ticks are obligate vectors, they directly influence their mammalian hosts and the

pathogens they transmit (Anderson and Magnarelli, 2008; Hill and Wikel, 2005; Van Zee et al.,

2007). The interaction between the vector (tick), host, and pathogen (*Rickettsia*) is essential in

understanding and predicting the risk and transmission of tick-borne diseases (Jongejan et al.,

2007). Understanding the genetic structure of ticks can serve as an alternative indicator to infer

the potential spread of pathogens, such as *Rickettsia* (Gray, 2002). Our study found relationships

between (1) the population genetic structure of ticks and the corresponding *Rickettsia* infection

rates, (2) altitude and the population genetic structure of ticks, and (3) altitude and *Rickettsia*

infection rates. We found that host mobility may influence the genetic structure of ixodid ticks.

This information can be used to design more effective tick-borne disease control programs that

focus on screening and detecting pathogens found in ticks and their mammalian hosts. For

example, patterns of disease transmission from ticks with a high genetic divergence and less

mobile hosts, such as *I. ovatus*, are likely due to the movement of infected hosts rather than

infected ticks. Thus, screening prospective tick hosts for *Rickettsia* infection would be more

suitable in this example. We suggest screening the hosts of immature *I. ovatus*, such as small

rodents, instead of screening ticks. The *Rickettsia* infection rate in tick genetic groups can predict

the spread of tick-borne diseases caused by *Rickettsia,* such as the Japanese spotted fever. We

also found that altitude may influence the *Rickettsia* infection rate of *I. ovatus* genetic groups. This

information can be used to determine the high-risk areas (e.g., lowland, mountains etc.) of tick-

borne diseases along an altitudinal gradient. Genetically structured arthropod vectors, such as

ticks, can have different vector competencies, and environmental factors, such as altitudinal

gradients, can influence the vector's ability to acquire, transmit, and maintain the pathogen

infection (Mclain et al., 1995).

Acknowledgements

The authors are thankful for the assistance from the Niigata Prefectural Office during the tick sampling collection. We would also like to thank the alumni of MECOH Lab Ehime University for their help in the molecular analyses: Masaya Doi, Kohki Tanaka and Mizuki Ueda. We are also grateful to Micanaldo Francisco for the construction of the map for this manuscript. We would like to thank Enago (www.enago.jp) for the English language review and proofreading of the manuscript.

Funding

References

Agany, D. D., Pietri, J. E., Gnimpieba, E. Z., 2020. Assessment of vector-host-pathogen relationships using data mining and machine learning. Comput. Struct. Biotechnol. J., 18, 1704-1721. https://doi.org/10.1016/j.csbj.2020.06.031

Ahrens, C.W., Byrne,M., Rymer,P.D., 2019.Standing genomic variation within coding and regulatory regions contributes to the adaptive capacity to climate in a foundation tree species. Mol Ecol 28, 2502–2516. https://doi.org/10.1111/mec.15092

Anderson, J.F., Magnarelli,L.A.,2008. Biology of ticks. Infect. Dis. Clin. North Am. 22, 195-215. https://doi.org/10.1016/j.idc.2007.12.006

Araya-Anchetta, A., Busch, J.D., Scoles, G.A., Wagner, D.M., 2015. Thirty years of tick population genetics: A comprehensive review. Infect. Genet. Evol. 29, 164–179. https://doi.org/10.1016/j.meegid.2014.11.008

Arai, R., Sato, M., Kato, M., Aoki, J., Nishida, A., Watanabe, K., Hirokawa, C., Ikeda, S., Watanabe, K., Regilme, M.A.F., Sato, M.O., Tamura, T., 2021. Spotted fever group rickettsiae (SFGR) detection in

ticks following reported human case of Japanese spotted fever in Niigata Prefecture, Japan. Sci.

Rep. 11, 1–9. https://doi.org/10.1038/s41598-021-81587-9

Bandelt, H.J., Forster, P., Röhl, A., 1999. Median-joining networks for inferringintraspecific

phylogenies. Mol. Biol. Evol. 16, 37–48. https://doi.org/10.1093/oxfordjournals.molbev.a026036

Bono, L. M., Gensel, C. L., Pfennig, D. W., Burch, C. L. 2013. Competition and the origins of novelty:

experimental evolution of niche-width expansion in a virus. Biol. Lett., 9, 20120616.

https://doi.org/10.1098/rsbl.2012.0616

Boots, M., and Mealor, M., 2007. Local interactions select for lower pathogen infectivity. Sci, 315,

1284-1286. https://doi.org/10.1126/science.1137126

Bradburd, G.S., Ralph, P.L., Coop, G.M., 2013. Disentangling the effects of geographic and

ecological isolation on genetic differentiation. Evol., 67, 3258–3273.

https://doi.org/10.1111/evo.12193

Burri, C., Cadenas, F.M., Douet, V., Moret, J., Gern, L., 2007. *Ixodes ricinus* density and infection

prevalence of *Borrelia burgdorferi* sensu lato along a north-facing altitudinal gradient in the Rhône

Valley (Switzerland). Vector-Borne Zoonotic Dis.7,50-58. https://doi.org/10.1089/vbz.2006.0569

Cadenas, F.M., Rais, O., Jouda, F., Douet, V., Humair, P.F., Moret, J., Gern, L., 2007. Phenology of

Ixodes ricinus and infection with *Borrelia burgdorferi* sensu lato along a north-and south-facing

altitudinal gradient on Chaumont Mountain, Switzerland. J. Med. Entomol.44, 683-693.

https://doi.org/10.1603/0022-2585(2007)44[683:poirai]2.0.co;2

Casati, S., Bernasconi, M.V., Gern, L., Piffaretti, J.C., 2008. Assessment of intraspecific mtDNA

variability of European *Ixodes ricinus* sensu stricto (Acari: Ixodidae). Infect.Genet. Evol. 8, 152–158.

https://doi.org/10.1016/j.meegid.2007.11.007`

Chapman, M.A., Hiscock, S.J., Filatov, D.A., 2013. Genomic divergence during speciation driven by

adaptation to altitude. *Mol. Biol. Evol.* 30, 2553–2567. https://doi.org/10.1093/molbev/mst168

Cheng, C., White, B.J., Kamdem, C., Mockaitis, K., Costantini, C., Hahn, M.W., Besansky, N.J.,

2012. Ecological genomics of *Anopheles gambiae* along a latitudinal cline: a population-

resequencing approach. *Genetics.* 190, 1417–1432. https://doi.org/10.1534/genetics.112.139899

Chin, K. M., Wolfe, M. S., 1984. Selection on *Erysiphe graminis* in pure and mixed stands of

barley. Plant Pathol., 33, 535-546. https://doi.org/10.1111/j.1365-3059.1984.tb02878.x

Coyne, J.A., Orr, H.A., 2004. *Speciation* (Vol. 37). Sunderland, MA: Sinauer associates.

Daniel, M., Malý, M., Danielová, V., Kříž, B., Nuttall, P., 2015. Abiotic predictors and annual

seasonal dynamics of *Ixodes ricinus*, the major disease vector of Central Europe. Parasit Vectors. 8,

1-12. https://doi.org/10.1186/s13071-015-1092-y

Dougherty, E. R., Seidel, D. P., Carlson, C. J., Spiegel, O., Getz, W. M., 2018. Going through the

motions: incorporating movement analyses into disease research. Ecol. Lett. 21, 588-604.

https://doi.org/10.1111/ele.12917

Drummond, A.J., Rambaut, A., 2007. BEAST: Bayesian evolutionary analysis by sampling trees.

BMC Evol. Biol.7, 1-8. https://doi.org/10.1186/1471-2148-7-214

Eisen, R. J., Eisen, L.,Beard, C. B., 2016. County-scale distribution of *Ixodes scapularis* and *Ixodes*

pacificus (Acari: Ixodidae) in the continental United States. J. Med. Entomol., 53, 349-386.

https://doi.org/10.1093/jme/tjv237

Eremeeva, M.E. and Dasch, G.A., 2015. Challenges posed by tick-borne rickettsiae: eco-

epidemiology and public health implications. Front. Public Health.*3*, 55.

https://doi.org/10.3389/fpubh.2015.00055

Estrada-Peña, A., 2015. Ticks as vectors: taxonomy, biology, and ecology. Rev. Sci. Tech. 34, 53–65.

https://doi.org/10.20506/rst.34.1.2345.

L. Excoffier, H.E.L. Lischer, 2010. Arlequin suite ver 3.5: a new series of programs to perform

population genetics analyses under linux and windows. Mol. Ecol. Resour., 10, 564-567.

https://doi.org/10.1111/j.1755-0998.2010.02847

Exposito-Alonso, M., Vasseur, F., Ding, W., Wang, G., Burbano, H.A., Weigel, D., 2018. Genomic

basis and evolutionary potential for extreme drought adaptation in Ara*bidopsis thaliana*. Nat Ecol

Evol, 2, 352-358. https://doi.org/10.1038/s41559-017-0423-0

Falco, R. C., Fish, D., 1991. Horizontal movement of adult *Ixodes dammini* (Acari: Ixodidae)

attracted to CO2-baited traps. J. Med. Entomol. 28, 726-729.

https://doi.org/10.1093/jmedent/28.5.726

Fountain-Jones, N.M., Craft, M.E., Funk, W.C., Kozakiewicz, C., Trumbo, D.R., Boydston, E.E., Lyren,

L.M., Crooks, K., Lee, J.S., VandeWoude, S., Carver, S., 2017. Urban landscapes can change virus

gene flow and evolution in a fragmentation-sensitive carnivore. Mol. Ecol., 26, 6487-6498.

https://doi.org/10.1111/mec.14375

Funk, W.C., Murphy, M.A., Hoke, K.L., Muths, E., Amburgey, S.M., Lemmon, E.M., Lemmon, A.R.,

2016. Elevational speciation in action? Restricted gene flow associated with adaptive divergence

across an altitudinal gradient. J. Evol. Biol. 29, 241-252. https://doi.org/10.1111/jeb.12760

Gage, K.L., Burkot, T.R., Eisen, R.J., Hayes, E.B., 2008. Climate and vectorborne diseases. Am. J.

Prev. Med. 35, 436-50. https://doi.org/10.1016/j.amepre.2008.08.030

Gibson, A. K., Baffoe-Bonnie, H., Penley, M. J., Lin, J., Owens, R., Khalid, A.,Morran, L. T., 2020. The

evolution of parasite host range in heterogeneous host populations. J. Evol. Biol., 33, 773-782.

https://doi.org/10.1111%2Fjeb.13608

Gooding, R.H., 1996. Genetic variation in arthropod vectors of diseasecausing organisms:

Obstacles and opportunities. Clin. Microbiol. Rev. 9, 301. https://doi.org/10.1128/cmr.9.3.301

Gray, J.S., 2002. Biology of *Ixodes* species ticks in relation to tick-borne zoonoses. Wien. Klin.

Wochenschr. 114, 473-8.

Gregory, T.R., 2009. Understanding natural selection: essential concepts and common

misconceptions. Evol.: Educ. Outreach, 2, 156-175. https://doi.org/10.1007/s12052-009-0128-1

Guglielmone, A.A., Nava, S., Mastropaolo, M., Mangold, A.J., 2013. Distribution and genetic

variation of *Amblyomma triste* (Acari: Ixodidae) in Argentina. Ticks Tick. Borne. Dis. 4, 386–390.

https://doi.org/10.1016/j.ttbdis.2013.01.009

Hancock, A.M., Brachi, B., Faure, N., Horton, M.W., Jarymowycz, L.B., Sperone, F.G., Toomajian, C.,

Roux, F., Bergelson, J., 2011. Adaptation to climate across the *Arabidopsis thaliana* genome.

Science, 334, 83-86. https://doi.org/10.1126/science.1209244

Hasle, G., 2013. Transport of ixodid ticks and tick-borne pathogens by migratory birds. Front. Cell.

Infect. Microbiol., 3, 48. https://doi.org/10.3389%2Ffcimb.2013.00048

Hill, C.A., Wikel, S.K., 2005. The *Ixodes scapularis* Genome Project: an opportunity for advancing

tick research. Trends Parasitol., 21, 151-153. https://doi.org/10.1016/j.pt.2005.02.004

Hofreiter, M., Rabeder, G., Jaenicke-Després, V., Withalm, G., Nagel, D., Paunovic, M., Jambrĕsić,

G., Pääbo, S., 2004. Evidence for reproductive isolation between cave bear populations. Curr. Biol.,

14, 40-43. https://doi.org/10.1016/j.cub.2003.12.035

453 Ishiguro, F., Takada, N., Fujita, H., Noji, Y., Yano, Y., Iwasaki, H., 2008. Survey of the vectorial

454 competence of ticks in an endemic area of spotted fever group rickettsioses in Fukui Prefecture,

455 Japan. Microbiol. Immunol. 52,305-309. https://doi.org/10.1111/j.1348-0421.2008.00042.x

456 Jiang, S., Luo, M.X., Gao, R.H., Zhang, W., Yang, Y.Z., Li, Y.J., Liao, P.C., 2019. Isolation-by-

457 environment as a driver of genetic differentiation among populations of the only broad-leaved

458 evergreen shrub *Ammopiptanthus mongolicus* in Asian temperate deserts. Sci. Rep.9, 1-14.

459 https://doi.org/10.1038/s41598-019-48472-y

460 Jongejan, F., Nene, V., de la Fuente, J., Pain, A., Willadsen, P., 2007. Advances in the genomics of

461 ticks and tick-borne pathogens. Trends Parasitol., 23, 391-396.

462 https://doi.org/10.1016/j.pt.2007.07.004

463 Jouda, F., Perret, J.L., Gern, L., 2004. *Ixodes ricinus* density, and distribution and prevalence of

464 *Borrelia burgdorferi* sensu lato infection along an altitudinal gradient. J. Med. Entomol. 41, 162-

465 169. https://doi.org/10.1603/0022-2585-41.2.162

466 Juvenal, N., Edward, M., 2010. Seasonal dynamics and distribution of ticks in Rwanda: implications

467 for tick control strategy in Rwanda. J Anim Vet Adv, 2,21-25.

468 Kapun, M., Barrón, M.G., Staubach, F., Obbard, D.J., Wiberg, R.A.W., Vieira, J., Goubert, C., Rota-

469 Stabelli, O., Kankare, M., Bogaerts-Márquez, M., Haudry, A., 2020. Genomic analysis of European

470 *Drosophila melanogaster* populations reveals longitudinal structure, continent-wide selection, and

471 previously unknown DNA viruses. Mol. Biol. Evol. 37, 2661-2678.

472 https://doi.org/10.1093/molbev/msaa120

473 Kawecki, T.J., Ebert, D., 2004. Conceptual issues in local adaptation. Ecol. Lett. 7, 1225-1241.

474 https://doi.org/10.1111/j.1461-0248.2004.00684.x

Lampo, M., Rangel, Y., Matat, A., 2015. Population genetic structure of a three-host population

genetic in *Amblyommma dissimile*. J. Parasitol. 84, 1137–1142. https:// doi.org/10.2307/3284662.

Lim, M.C., Bi, K., Witt, C.C., Graham, C.H. and Dávalos, L.M., 2021. Pervasive genomic signatures of

local adaptation to altitude across highland specialist Andean hummingbird populations. J.

Hered., 112, 229-240. https://doi.org/10.1093/jhered/esab008

Lion, S., Gandon, S., 2015. Evolution of spatially structured host–parasite interactions. *Journal of*

Evol. Biol., 28, 10-28. https://doi.org/10.1111/jeb.12551

Love,R.R., Steele, A.M., Coulibaly,M.B., Traore,S.F., Emrich,S.J., Fontaine,M.C., Besansky,N.J., 2016.

Chromosomal inversions and ecotypic differentiation in *Anopheles gambiae*: the perspective from

whole-genome sequencing. Mol. Ecol. 25, 5889-906. https://doi.org/10.1111/mec.13888

MacDonald, A.J., Hyon, D.W., Iii, J.B.B., Connor, K.E.O., Swei, A., Briggs, C.J., 2017. Lyme disease

risk in southern California: abiotic and environmental drivers of *Ixodes pacificus* (Acari: Ixodidae)

density and infection prevalence with *Borrelia burgdorferi.* Parasit Vectors 10:7 DOI

10.1186/s13071-016-1938-y.

Maddison, W.P., Maddison, D.R., 2011. Mesquite: A Modular System for Evolutionary Analysis, See.

http://mesquiteproject.org.

Mahara, F., 1997. Japanese spotted fever: report of 31 cases and review of the literature.Emerg.

Infect. Dis. 3, 105–111. https://doi.org/10.3201/eid0302.970203

Mclain, D.K., Wesson, D.M., Oliver Jr, J.H., Collins, F.H., 1995. Variation in ribosomal DNA internal

transcribed spacers 1 among eastern populations of *Ixodes scapularis* (Acari: Ixodidae).   J. Med.

Entomol, 32, 353-360. https://doi.org/10.1093/jmedent/32.3.353

McCoy, K. D., Boulinier, T., Tirard, C., Michalakis, Y., 2001. Host specificity of a generalist parasite:

genetic evidence of sympatric host races in the seabird tick *Ixodes uriae*. J. Evol. Biol., 14, 395-405.

https://doi.org/10.1046/j.1420-9101.2001.00290.x

Miller, A.D., Hoffmann, A.A., Tan, M.H., Young, M., Ahrens, C., Cocomazzo, M., Rattray, A.,

Ierodiaconou, D.A., Treml, E. and Sherman, C.D., 2019. Local and regional scale habitat

heterogeneity contribute to genetic adaptation in a commercially important marine mollusc

(Haliotis rubra) from southeastern Australia. Mol. Ecol., 28, 3053-3072.

https://doi.org/10.1111/mec.15128

Mixson, T.R., Lydy, S.L., Dasch, G.A., Real, L.A., 2006. Inferring the population structure and

demographic history of the tick, *Amblyomma americanum* Linnaeus. J. Vector Ecol. 31, 181–192.

https://doi.org/10.3376/1081-1710(2006)31[181:itpsad]2.0.co; 2.

Miyamoto, K., Nakao, M., Fujita, H., Sato, F., 1993. The ixodid ticks on migratory birds in Japan and

the isolation of Lyme disease spirochetes from bird-feeding ticks. *Med. Entomol. and zool.*, 44(4),

315-326. https://doi.org/10.7601/mez.44.315

Morel, P., 1981. Maladies à tiques du bétail en Afrique In: Précis de parasitologie vétérinaire

tropicale. Ministére de la Coopération et du Développement, République Française, Norval, R.A.I.,

Donachie, pp: 471-717.

Nosil, P., 2012. Ecological speciation. Oxford University Press.

Nzige, T., 1978. Ecologie des tiques du bétail. Rapport de stage, Higher Institute of Agriculture and

Animal Husbandry (ISAE-Busogo) Publications, Musanze, Rwanda.

Oliveira, E.F., Martinez, P.A., São-Pedro, V.A., Gehara, M., Burbrink, F.T., Mesquita, D.O., Garda,

A.A., Colli, G.R., Costa, G.C., 2018. Climatic suitability, isolation by distance and river resistance

explain genetic variation in a Brazilian whiptail lizard. Heredity, 120, 251-265.

https://doi.org/10.1038/s41437-017-0017-2

Oliver, J.D., Bennett, S.W., Beati, L., Bartholomay, L.C., 2017. Range expansion and increasing

Borrelia burgdorferi infection of the tick *Ixodes scapularis* (Acari: Ixodidae) in Iowa, 1990–2013. J.

Med. Entomol. 54, 1727–1734. https://doi.org/10.1093/jme/tjx121

Osborne, O.G., Batstone, T.E., Hiscock, S.J. & Filatov, D.A., 2013. Rapid speciation with gene flow

following the formation of Mt. Etna. *Genome Biol. Evol.* 5, 1704–1715.

https://doi.org/10.1093/gbe/evt127

Páez-Triana, L., Muñoz, M., Herrera, G., Moreno-Pérez, D.A., Tafur-Gómez, G.A., Montenegro, D.,

Patarroyo, M.A., Paniz-Mondolfi, A., Ramírez, J.D., 2021. Genetic diversity and population

structure of *Rhipicephalus sanguineus* sensu lato across different regions of Colombia. Parasit

Vectors, 14, 1-11. https://doi.org/10.1186/s13071-021-04898-w

Paradis, E., Schliep, K., 2018. APE 5.0: an environment for modern phylogenetics andevolutionary

analyses in R. Bioinformatics 35, 526–528. https://doi.org/10.1093/bioinformatics/bty633.P

Parola, P., Raoult, D., 2001. Tick-borne bacterial diseases emerging in Europe. Clin. Microbiol.

Infect. 7,80-83. https://doi.org/10.1046/j.1469-0691.2001.00200.x

Parola, P., Paddock, C.D., Socolovschi, C., Labruna, M.B., Mediannikov, O., Kernif, T., Abdad, M.Y.,

Stenos, J., Bitam, I., Fournier, P.E., Raoult, D., 2013. Update on tick-borne rickettsioses around the

world: a geographic approach. Clin. Microbiol. Rev., 26, 657-702.

https://doi.org/10.1128/cmr.00032-13

Randolph,S.E., Miklisová,D., Lysy,J., Rogers,D.J., Labuda, M., 1999. Incidence from coincidence:

patterns of tick infestations on rodents facilitate transmission of tick-borne encephalitis virus.

Parasitology. 118,177–186. https://doi.org/10.1017/s0031182098003643

Regilme, M.A.F., Sato, M., Tamura, T., Arai, R., Sato, M.O., Ikeda, S., Gamboa, M., Monaghan, M.T.

and Watanabe, K., 2021. Comparative population genetic structure of two ixodid tick species

(Acari: Ixodidae)(*Ixodes ovatus* and *Haemaphysalis flava*) in Niigata prefecture, Japan. Infect.

Genet. Evol.,, 94, 104999.

Reichard, M.V., Kocan, A.A., Van Den Bussche, A., Barker, R.W., Wyckoff, J., Ewing, S.A., 2005.

Sequence variation of the ribosomal DNA second internal transcribed spacer region in two

spatially distinct populations of *Amblyomma americanum* (L.)(Acari: Ixodidae). J. Parasitol. 91,

260–264. https://doi.org/10.1645/GE-292R1.

Regoes, R. R., Nowak, M. A., Bonhoeffer, S., 2000. Evolution of virulence in a heterogeneous host

population. Evol.54, 64-71. https://doi.org/10.1111/j.0014-3820.2000.tb00008.x

Richmond, C.E., Breitburg, D.L.,Rose, K.A., 2005. The role of environmental generalist species in

ecosystem function. Ecol. Model. 188, 279-295. https://doi.org/10.1016/j.ecolmodel.2005.03.002

Rozas, J., Ferrer-Mata, A., S̓anchez-DelBarrio, J.C., Guirao-Rico, S., Librado, P., Ramos-Onsins, S.E.,

S̓anchez-Gracia, A., 2017. DnaSP 6: DNA sequence polymorphism analysis of large data sets. Mol.

Biol. Evol. 34, 3299–3302. https://doi.org/10.1093/molbev/msx248.

Sato, M., Ikeda, S., Arai, R., Kato, M., Aoki, J., Nishida, A., Watanabe, K., Hirokawa, C., Watanabe, K.,

Regilme, M.A.F., Sato, M., 2021. Diversity and distribution of ticks in Niigata prefecture, Japan

(2016–2018): Changes since 1950. Ticks Tick Borne Dis, 12, 101683.

https://doi.org/10.1016/j.ttbdis.2021.101683

Sexton,J.P., Hangartner,S.B., Hoffmann,A.A., 2014. Genetic isolation by environment or distance:

which pattern of gene flow is most common?. Evol., 68, 1–15. https://doi.org/10.1111/evo.12258

Streicker, D.G., Winternitz, J.C., Satterfield, D.A., Condori-Condori, R.E., Broos, A., Tello, C.,

Recuenco, S., Velasco-Villa, A., Altizer, S. and Valderrama, W., 2016. Host–pathogen evolutionary

signatures reveal dynamics and future invasions of vampire bat rabies. Proc Natl Acad Sci U S

A, 113, 10926-10931. https://doi.org/10.1073/pnas.1606587113

Thrall, P. H., Burdon, J. J., 2003. Evolution of virulence in a plant host-pathogen

metapopulation. Science, 29, 1735-1737. https://doi.org/10.1126/science.1080070

Trout, R.T., Steelman, C.D., Szalanski, A.L., 2010. Population genetics of *Amblyomma americanum*

(Acari: Ixodidae) collected from Arkansas. J. Med. Entomol. 47, 152–161.

https://doi.org/10.1603/me09106.

Van Treuren,W., Ponnusamy,L., Brinkerhoff,R.J., Gonzalez,A., Parobek,C.M., Juliano,J.J., Andreadis

T.G., Falco,R.C., Ziegler,L.B., Hathaway,N., Keeler,C., Emch,M., Bailey,J.A., Roe,R.M., Apperson,C.S.,

Knight,R., Meshnick,S.R., 2015. Variation in the microbiota of Ixodes ticks with regard to

geography, species, and sex. Appl. Environ. Microbiol. 81, 6200–6209

https://doi.org/10.1128/aem.01562-15

Van Zee, J.P., Geraci, N.S., Guerrero, F.D., Wikel, S.K., Stuart, J.J., Nene, V.M., Hill, C.A., 2007. Tick

genomics: the Ixodes genome project and beyond. Int. J. Parasitol., 37, 1297-1305.

https://doi.org/10.1016/j.ijpara.2007.05.011

Wang, I.J., Bradburd, G.S., 2014. https://doi.org/10.1111/mec.12938. Mol. Ecol., 23, 5649-5662.

Wang,I.J., Glor,R.E., Losos,J.B., 2013. Quantifying the roles of ecology and geography in spatial

genetic divergence. Ecol. Lett., 16, 175–182. https://doi.org/10.1111/ele.12025

Wang ,J., Summers,K., 2010. Genetic structure is correlated with phenotypic divergence rather

than geographic isolation in the highly polymorphic strawberry poison-dart frog. Mol. Ecol., 19,

447–458. https://doi.org/10.1111/j.1365-294x.2009.04465.x

White, P.S., Choi, A., Pandey, R., Menezes, A., Penley, M., Gibson, A.K., de Roode, J. Morran, L.,

2020. Host heterogeneity mitigates virulence evolution. Biol. Lett., 16,20190744.

https://doi.org/10.1098/rsbl.2019.0744

Yamaguti, N., Tipton, V. J., Keegan, H. L., & Toshioka, S. 1971. Ticks of Japan, Korea, and the

Ryukyu islands. Brigham Young University Science Bulletin, Biological Series, 15, 1.

Yamaji, K., Aonuma, H., Kanuka, H., 2018. Distribution of tick-borne diseases in Japan:past patterns

and implications for the future. J. Infect. Chemother. 24,

499504.https://doi.org/10.1016/j.jiac.2018.03.012.

Yamauchi, T., Takeno, K., 2000. Record of Haemaphysalis flava Neumann (Acari,Ixodidae) from

Zoothera dauma (Latham) (Aves) in Kyushu, Japan. Esakia 40,99–101.

https://doi.org/10.5109/2647

Yu, X.J., Liang, M.F., Zhang, S.Y., Liu, Y., Li, J.D., Sun, Y.L., Zhang, L., Zhang, Q.F., Popov, V.L., Li, C.

and Qu, J., 2011. Fever with thrombocytopenia associated with a novel bunyavirus in

China. N. Engl. J. Med, 364, 1523-1532. DOI: 10.1056/NEJMoa1010095

Zhen, Y., Harrigan, R.J., Ruegg, K.C., Anderson, E.C., Ng, T.C., Lao, S., Lohmueller, K.E. and Smith,

T.B., 2017. Genomic divergence across ecological gradients in the Central African rainforest

songbird (A ndropadus virens). Mol. Ecol., 26, 4966-4977. https://doi.org/10.1111/mec.14270

Figure Captions

Figure 1. Median-joining network of the 59 *cox1* haplotype sequences of *Rickettsia* positive and

negative *I. ovatus.* Haplotype groups are indicated by numbers (1 to 4).

615 **Figure 2.** The influence of altitude on the habitat distribution of *I. ovatus*. The points indicate the

616 sampling sites of the collected individuals per *I. ovatus* haplotype group (n = 3) in the haplotype

617 network (Figure 1). The map shows the elevation level across the sampling area, with Niigata

618 Prefecture Japan depicted as a color gradation. Group 4, not shown in this figure, with haplotypes

619 54 and 55 were found on Sado island, which is encircled in black in box 3.

626   **Figure 3.** The relationship between the mean altitude of *Rickettsia-infected* (n = 46) and non-

627   infected (n = 300) *I. ovatus.* Welch t-tests revealed a significant difference in the mean altitude

628   between *Rickettsia-infected* and non-infected *I. ovatus* at $p < 0.05$. Populations with combined

629   sites are labeled A to H.

Figure 4. An unweighted pair group method with the arithmetic mean (UPGMA) dendrogram of *I.*

ovatus based on the pairwise genetic distance (FST) of *cox1* among the 7 populations across

Niigata Prefecture, Japan. We excluded one population due to limited number of samples.

Tables

Table 1. Summary of the haplotype and *Rickettsia* infection rates among the 7 ixodid tick species

obtained   in the

Niigata

Tick species ***	ns	n	nh	r	Global F$_{ST}$
1. *Ixodes ovatus*	29	346	59	46 (13.26%)	0.4154*
2. *Haemaphysalis flava*	18	243	66	15 (6.17%)	0.3597
3. *Ixodes monospinus*	6	19	9	6 (31.58%)	0.9034
4. *Ixodes asanumai*	7	9	4	5 (55.55%)	1.0000
5. *Ixodes nipponensis*	2	4	4	4 (100%)	0.3345
6. *Ixodes persulcatus*	3	4	3	0 (0%)	1.0000
7. *Haemaphysalis japonica*	2	2	2	2 (100%)	**
Total		628		78	

Prefecture, Japan

Abbreviations: ns no. of sampling sites; n sample size; nh no of haplotypes; r rickettsia infection rate per species; *p < 0.05** not

enough samples for analysis***tick species identification is based on molecular identification using the *cox1* marker and BLAST results

Table 2. The differences in *Rickettsia* infection rates and mean altitude in *I. ovatus* haplotype

groups and cluster dendrogram groups. The table shows the distribution of *Rickettsia*-infected and

uninfected *I. ovatus* and the mean altitude in each of the haplotype groups as shown in Figure 1 *(I.*

ovatus haplotype network) and the cluster dendrogram in Figure 4 (*I. ovatus* cluster dendrogram).

The z-score test showed a significant difference at $p < 0.05$ between the *Rickettsia* detection rates

in haplotype groups 1 and 2 (indicated by [ab)]) and in cluster dendrogram groups 1 and 2.he Welch t-

test at $p < 0.05$ revealed a significant difference in the mean altitude of haplotype groups 1 and 3

and cluster dendrogram groups 1 and 2 (indicated by [ab]). Haplotype group 4 was not included in

the analysis due to its low sample size.

Haplotype group	Rickettsia infection rate			Mean altitude
	Positive	Negative	Detection rate	
1	1	92	1.07%[ab]	294[ab]
2	28	99	22.04%[ab]	341[a]
3	16	110	12.70%[a]	374[ab]
Dendrogram Group				
1	4	120	3.23%[ab]	30
2	40	227	14.98%[ab]	345

Supplementary Tables and Figures

Supplementary Table 1. The difference in *Rickettsia* infection rates in *H. flava* haplotype groups

and a table showing the distribution of *Rickettsia*-infected and uninfected *H. flava* in each genetic

groups as shown in Supplementary Figure 2 *(H. flava* median-joining network). The results of the z-

score test for two populations proportions at $p < 0.05$ showed no significant difference between

the *Rickettsia* detection rate in haplotype groups 1 and 2. The Welch t-test at $p < 0.05$ revealed no

significant difference in the mean altitude of the two groups.

Haplotype group	*Rickettsia* infection			Mean altitude
	Positive	Negative	Detection rate	
1	1	45	2.17%[ab]	205[ab]
2	14	183	7.11%[ab]	165[ab]

684   **Supplementary Figure 3.** Phylogenetic tree from the BEAST analysis of 66 haplotype *cox1*

685   sequences of *H. flava.* The blue labeled haplotypes indicate *Rickettsia* infection in individual

686   samples. The parentheses in red provide the number of *Rickettsia*-infected ticks. The black labeled

687   haplotypes are negative for *Rickettsia* infection.