1

Stepwise recombination suppression around the mating-type locus in the fungus *Schizothecium tetrasporum* (Ascomycota, Sordariales)

Nina Vittorelli [1,2,3], Alodie Snirc [1], Emilie Levert [1,2], Valérie Gautier [2], Christophe Lalanne [2], Elsa De Filippo [1,2], Ricardo C. Rodríguez de la Vega [1], Pierre Gladieux [4], Sonia Guillou [4], Yu Zhang [5], Sravanthi Tejomurthula [5], Igor Grigoriev [5], Robert Debuchy [6], Philippe Silar [2]*, Tatiana Giraud [1]*, Fanny E. Hartmann [1]*

[1] Ecologie Systematique et Evolution, CNRS, Université Paris-Saclay, AgroParisTech, 91198 Gif-sur-Yvette, France

[2] Laboratoire Interdisciplinaire des Energies de Demain, Université Paris Cité, Paris, France

[3] Département de Biologie, École Normale Supérieure, PSL Université Paris, Paris, France

[4] PHIM Plant Health Institute, Univ Montpellier, INRAE, CIRAD, Institut Agro, IRD, Montpellier, France

[5] Joint Genome Institute, Lawrence Berkeley National Laboratory, University of California Berkeley

[6] Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC), 91198 Gif-sur-Yvette, France.

Running title: Recombination suppression around a mating-type locus

Corresponding author: Fanny E. Hartmann

E-mail: fanny.hartmann@universite-paris-saclay.fr

*co-senior authors

Keywords: fungi, *Neurospora tetrasperma,* sex chromosomes, pseudo-homothallism, evolutionary strata, mating system, interference, Sordariales*, Podospora tetraspora*

ABSTRACT

Recombination is often suppressed at sex-determining loci in plants and animals, and at self-

incompatibility or mating-type loci in plants and fungi. In fungal ascomycetes, recombination

suppression around the mating-type locus is associated with pseudo-homothallism, *i.e.,* the

production of self-fertile dikaryotic sexual spores carrying the two opposite mating types. This has

been well studied in two species complexes from different families of Sordariales: *Podospora anserina*

and *Neurospora tetrasperma.* However, it is unclear whether this intriguing convergent association

holds in other species. We show here that *Schizothecium tetrasporum,* a fungus from a third family in

the order Sordariales*,* also produces mostly self-fertile dikaryotic spores carrying the two opposite

mating types. This was due to a high frequency of second meiotic division segregation at the mating-

type locus, indicating the occurrence of a single and systematic crossing-over event between the

mating-type locus and the centromere, as in *P. anserina*. The mating-type locus has the typical

Sordariales organization*,* plus a *MAT1-1-1* pseudogene in the *MAT1-2* haplotype. High-quality

genome assemblies of opposite mating types and segregation analyses revealed a suppression of

recombination in a region of 1.3 Mb around the mating-type locus. We detected three evolutionary

strata, displaying a stepwise extension of recombination suppression, but no rearrangement or

transposable element accumulation in the non-recombining region. Our findings indicate a convergent

evolution of self-fertile dikaryotic sexual spores across multiple ascomycete fungi. The particular

pattern of meiotic segregation at the mating-type locus was associated with recombination

suppression around this locus, that had extended stepwise. This association is consistent with a

recently proposed mechanism of deleterious allele sheltering through recombination suppression

around a permanently heterozygous locus.

AUTHOR SUMMARY

Recombination allows faster adaptation and the purging of deleterious mutation but is often
paradoxically lacking in sex chromosomes. It has been recently recognized that recombination can
also be suppressed on fungal mating-type chromosomes, but the evolutionary explanation and the
proximal mechanism of this phenomenon remain unclear. By studying here the sexual biology of a
poorly studied mold living in rabbit dung, we reveal a striking convergence in three distant fungal
lineages of an independently evolved association between the production of self-fertile sexual spores
(carrying two nuclei with opposite mating types), a particular segregation of the mating-type locus and
the lack of recombination on mating-type chromosomes, having evolved stepwise. Such a convergent
association suggests causal relationships and will contribute to unveil the evolutionary causes of
recombination suppression.

Graphical summary

66
67

INTRODUCTION

In eukaryotes, recombination is often considered to be a major long-term evolutionary advantage. Indeed, it prevents the accumulation of deleterious mutations (1) and facilitates the combination of multiple advantageous alleles in the same genome (2). However, convergent recombination suppression events have been observed at sex-determining loci in plants and animals, and at self-incompatibility or mating-type loci in plants and fungi (3). It has been suggested that this recombination suppression links lock-and-key genes together, enabling self-incompatibility systems to function correctly (4–6). Surprisingly, recombination suppression has often been found to extend distally, in a stepwise manner, from sex and mating-type loci, generating evolutionary strata of differentiation between haplotypes (7–10). The evolutionary causes of this stepwise recombination suppression beyond mating-compatibility loci remain a matter of debate (9–12). Sexual antagonism was long thought to be the main driving force generating evolutionary strata on sex chromosomes, through selection for the linkage to the sex-determining locus of genes with alleles beneficial in only one sex (12). However, it has been difficult to obtain definitive evidence supporting this hypothesis, and evolutionary strata have also been reported in fungi without sexual antagonism or other types of antagonism between mating types (8–10,13). Alternative hypotheses have been developed (10,11), based on associative overdominance associated with the sheltering of deleterious mutations (14), dosage compensation (15) or the spread of transposable elements (16).

Fungi are useful models for studies of the evolution of recombination suppression: they have many assets for evolutionary genomics studies (17) and several convergent recombination suppression events around mating-type loci have been observed in this group (8,10,18). Fungi have highly diverse modes of reproduction, mating systems (selfing versus outcrossing) and breeding systems (mating compatibility), with considerable differences sometimes reported between closely related species (19). Among all breeding systems, heterothallism was the most investigated and seems to display the most conserved scheme in fungi (20). In heterothallic fungi, mating can occur only between haploid cells carrying different alleles at the mating-type locus. The term "heterothallism" was initially coined after observations of *in vitro* crosses, in which mating could occur only between different thalli (*i.e.,* mycelia or hyphal networks), each arising from a single spore. We now know that this heterothallism was due to the thalli mostly being haploid and carrying different alleles at the

mating-type locus. Other fungi were described as "homothallic" because mating within a single thallus

was possible. We now know that homothallism is possible because mating in these fungi does not

need to occur between cells of different mating types. However, some fungi capable of displaying

mating within a single thallus *in vitro* were later found to carry two types of nuclei, of different mating

types, in each cell. They were therefore called pseudo-homothallic.

The pseudo-homothallic breeding system has been studied in detail in two species from

different families of the order Sordariales (Ascomycota): *Neurospora tetrasperma* (Sordariaceae*),*

which is typically found as orange blooms on scorched vegetation at burnt sites after a fire, and

Podospora anserina (Podosporaceae), which is mostly found in herbivore dung. Sexual spore

formation is similar in these two species (21). Karyogamy occurs between two haploid nuclei of

different mating types, followed by meiosis, and then the post-meiotic mitosis of each meiosis product.

The ascus produced therefore contains eight haploid nuclei, which are packaged in pairs to form four

sexual spores (ascospores). The resulting ascospores are, thus, dikaryotic, *i.e.*, each contains two

haploid nuclei. A small fraction of asci (0.1% for *N. tetrasperma* and 1.0% in *P. anserina*) contains five

ascospores, with one dikaryotic spore replaced by a pair of monokaryotic ascospores (*i.e.,*

ascospores carrying a single haploid nucleus), recognizable in *P. anserina* by their smaller size. In

both species, the dikaryotic spores almost always contain nuclei of opposite mating types, and are

therefore referred to as heterokaryotic ascospores, each of which germinates into a self-fertile

mycelium (21). The particular pattern of segregation of the mating-type locus leads to the generation

of nuclei of different mating types in close physical proximity, which can be compacted into a single

ascospore. In *N. tetrasperma*, the mating-type locus segregates during the first meiotic division, due

to the linkage between the centromere and the mating-type locus. During meiosis II, the spindles are

parallel, and, following the post-meiotic mitosis, each ascospore is delineated around two non-sister

nuclei, resulting in the compaction of opposite mating types in the same cell. In *P. anserina*, the

mating-type locus segregates during the second meiotic division, due to the occurrence of a single

and systematic crossing-over (CO) event between the centromere and the mating-type locus. During

meiosis II, the spindles are aligned, leading to the same final nuclear distribution as in *N. tetrasperma*.

After the post-meiotic mitosis, the ascospores formed contain nuclei of opposite mating types.

In these two pseudo-homothallic species, there are large non-recombining regions around the

mating-type locus. In *N. tetrasperma*, the 7 Mb region without recombination encompasses the

mating-type locus and the centromere (22). In *P. anserina*, a 0.8 Mb region devoid of recombination

has been identified around the mating-type locus in the reference strain (23). Each of these two

species belongs to a complex of cryptic species (22,24,25), which also all harbor a region of

recombination suppression around their mating-type locus; however the size of the non-recombining

region differs between species within each species complex (22,26,27). In both *N. tetrasperma* and *P.*

pseudocomata (a cryptic species from the *P. anserina* species complex; (24)), the suppression of

recombination has extended stepwise, generating evolutionary strata (10,22,26,27). It is intriguing that

the presence of non-recombining regions around fungal mating-type loci and their stepwise extension

seem to be associated with pseudo-homothallism in ascomycetes (10). One hypothesis that could

account for this association is that most strains are heterokaryotic at the mating-type locus for most of

their life cycle in pseudo-homothallic ascomycetes. This could allow recombination suppression to be

selected as a means of sheltering recessive deleterious alleles, by linking them to a permanently

heterozygous locus (14). Indeed, non-recombinant haplotypes with fewer recessive deleterious

mutations than the population average should be selected for when rare, but if they increase in

frequency, they will start to occur as homozygotes, exposing their recessive load, unless they capture

a permanently heterozygous allele; in such cases, they continue to be selected and can rapidly reach

their maximal frequency, *i.e.* becoming fully associated with the heterozygous allele close to which

they appeared (14). The dikaryotic phase should also foster a stepwise extension of recombination

suppression around the mating-type locus by the same mechanism (14). However, studies of

additional pseudo-homothallic fungi are required, to assess the strength of the association between

pseudo-homothallism and stepwise recombination suppression around the mating-type locus.

We therefore studied *Schizothecium tetrasporum* (previously known as *Podospora tetraspora*

or *P. minuta var. tetraspora),* a third pseudo-homothallic species from order Sordariales (21,28).

Schizothecium   tetrasporum belongs to the family Schizotheciaceae (also known as

Neoschizotheciaceae), while *P. anserina* belongs to the Podosporaceae and *N. tetrasperma* to the

Sordariaceae (29,30). The phylogenetic relationships between these three families are not well

established, but they probably diverged over 150 million years ago (31). For example, the

phylogenetic distance between *P. anserina* and *N. crassa,* a member of the Sordariaceae (like *N.*

tetrasperma), corresponds roughly to that between humans and fishes (32). Phylogenetically,

Schizothecium tetrasporum is roughly equidistant from these two species (29). However, it displays

striking cytological similarities to *P. anserina*, particularly in terms of dikaryotic spore production: in the

FGSC7436 strain, 98% of asci were reported to produce four dikaryotic spores, with the spindles

aligned in the ascus during meiosis II (21). However, this previous study did not clearly investigate

whether the two nuclei of a dikaryotic spore contained opposite mating types and did not study the

mating-type locus or recombination suppression.

We identified media suitable for reproducible and fast growth, and media for sexual

reproduction in *S. tetrasporum.* We studied the dikaryotic strain CBS815.71, estimating the frequency

of second-division segregation of the mating-type locus at 99.5%. This indicates the existence of a

single and systematic CO event between the mating-type locus and the centromere. We also built

high-quality genome assemblies of homokaryons (*i.*e., with only one nucleus type) with opposite

mating types, making it possible: i) to elucidate the genetic organization of the mating-type locus, and

ii) to detect a 1.3 Mb region without recombination around the mating-type locus, through analysis of

differentiation between the mating-type chromosomes. We observed three evolutionary strata, with

significantly different levels of differentiation between mating-type chromosomes, suggesting a

stepwise extension of recombination suppression. We detected no increase in transposable element

content, non-synonymous substitutions or rearrangements in the non-recombining region.

Recombination suppression was confirmed by segregation analyses. Our findings indicate that the

convergent evolution of an extremely high frequency of self-fertile dikaryotic spores together with

recombination suppression around the mating-type locus has occurred in multiple pseudo-homothallic

fungal species.

RESULTS

***High frequency of self-fertile spores in* Schizothecium tetrasporum**

The study of a single strain of *Schizothecium tetrasporum* (also known as *P. tetraspora*; Figure 1)

suggested that this species was pseudo-homothallic and produced self-fertile heterokaryotic

ascospores similarly to *P. anserina* (21,28), *i.e.,* with most asci containing four dikaryotic spores,

probably carrying both mating types. We studied the breeding system of *S. tetrasporum* further, by

testing different media and identifying those optimal for reproducible and fast growth, and those

optimal for sexual reproduction (see Material & Method section). Raju & Perkins found that the

FGSC7436 strain produced 2% five-spore asci. In this study, we found that 10% of the asci produced by the CBS815.71 strain had five spores. We also observed, in rare frequencies, other kinds of asci, including six-spore asci, and some with "extra-large" ascospores, probably encompassing three nuclei, rather than the two usually encompassed during ascospore delimitation (Figure 1). The resulting mycelium produced numerous asexual single-cell propagules, for which we were unable to find appropriate conditions for germination and development. These cells probably, therefore, corresponded to spermatia rather than conidia, *i.e.* they are sexual cells acting as fertilizing elements and are unable to germinate, in contrast to asexual spores. *Podospora anserina* also produces spermatia, which function exclusively in the fertilization of the female structures, the protoperithecia (33). We generated an F1 progeny from the CBS815.71 dikaryotic strain, and isolated two homokaryotic offspring of opposite mating types, referred to hereafter as CBS815.71-sp3 and CBS815.71-sp6. The mating types of these strains were named according to the standard nomenclature (34): *MAT1-1* for CBS815.71-sp6 and *MAT1-2* for CBS815.71-sp3.

We assessed the frequency of second-division segregation (SDS) at the mating-type locus (Figure 2), by analyzing the progeny of a cross between CBS815.71-sp3 and CBS815.71-sp6. We collected 190 four-spore asci (each ascus containing the products of a given meiosis) and isolated large (dikaryotic) ascospores. Mating types were determined in the mycelium generated from germinating ascospores, by PCR with two primer pairs, each specific to one mating-type allele and yielding amplicons of different sizes. We initially genotyped only one ascospore per ascus, as this is usually sufficient to differentiate between two extreme cases: first-division segregation at meiosis (FDS) and second-division segregation (SDS). FDS yields four-spore asci containing two ascospores of a single mating type and another two ascospores of the opposite mating type, whereas SDS results in four ascospores, all containing both mating types (Figure 2A). We found that 87% (166/190) of the thalli resulting from ascospore germination carried both mating types, indicating that they resulted from SDS, and 13% (24/190) carried a single mating type, and were therefore either haploid or had two nuclei of the same mating type. Indeed, thalli carrying a single mating type may result not only from FDS, but also from the loss of a nucleus during growth in thalli resulting from SDS (Figure 2B). We differentiated between these two situations by determining the alleles at the mating-type locus in thalli resulting from the germination of the three remaining spores from the 24 four-spore asci in which we found a single mating type in the first genotyped ascospore. Only one of the 24 asci analyzed

proved to be a true FDS ascus (*i.e.*, with two *MAT1-1* and two *MAT1-2* ascospores). The other 23 asci

contained at least one ascospore carrying both mating types, indicating that the homokaryotic thalli

growing from their other ascospores resulted from the loss of nuclei during growth after SDS in asci

(Figure 2B). We estimated the overall frequency of SDS for mating type at 99.5% (189/190) in the

CBS815.71 strain.

We detected a substantial number of homokaryotic thalli originating from SDS asci (23 thalli

from 189 germinating heterokaryotic ascospores), suggesting that the loss of nuclei corresponding to

one mating type in heterokaryotic thalli may occur at a non-negligible frequency in the CBS815.71

strain. Raju and Perkins (21) showed that the two nuclei present in developing ascospores undergo

mitosis, which is rapidly followed by the migration of two of the four daughter nuclei into the future

pedicel cell, which are presumed to abort, or at least to fail to contribute to the nuclear composition of

the resulting culture. It is therefore possible that homokaryotic thalli result from the migration of two

nuclei carrying the same mating type into the future pedicel (leaving two nuclei of the same mating

type in the maturing large ascospore cell), rather than the migration of two nuclei carrying opposite

mating types (leaving two nuclei of opposite mating types in the maturing large ascospore cell; Figure

2B). It is difficult to test this hypothesis, because there is no easy way to follow the migration of nuclei

during ascospore maturation. Alternatively, *MAT1-1/MAT1-2* heterokaryons may sometimes lose one

type of nucleus during vegetative growth.

We tested this second hypothesis, by determining mating type in several sections of

heterokaryotic thalli. A heterokaryon was generated by vegetative fusion (*i.e.*, by mixing identical

amounts of ground mycelia of the two homokaryotic strains of opposite mating types, CBS815.71-sp3

and CBS815-sp6). This procedure of mixed homokaryons chopped in very fine grain leads to the

generation of a heterokaryotic mycelium through multiple vegetative fusions (35) and nuclear

movements in the mycelial interconnected network of cells (36); no sexual reproduction can occur

under these growth conditions. The resulting heterokaryotic mycelium was grown on V8 medium and

incubated until the colony had reached 4 cm in diameter. It was then sliced into 70 pieces, each with a

volume of about 1 mm^3. PCR analyses demonstrated that only 31 of these 70 pieces carried both

mating types, 29 contained only *MAT1-2* and 10 contained only *MAT1-1*. In a parallel experiment on

another thallus of the same heterokaryon, the 70 pieces of mycelium were used to inoculate fresh M2

medium, and regenerating thalli were allowed to grow to a diameter of 1 cm. A randomly selected 1

mm^3 piece of each of these thalli was sampled at the edge of growing colonies and genotyped by

PCR for mating type. We found that 34 of the 70 regenerated thalli carried both mating types,

whereas 25 carried only *MAT1-2* and 11 carried only *MAT1-1*. These results indicate that one of the

two nuclei can be lost during vegetative growth, and they confirm the high rate of SDS in *S.*

tetrasporum, yielding 99.5% of asci with all four ascospores initially containing both mating types, but

with the occasional loss of one nucleus during vegetative growth. The *MAT1-1* mating type was

preferentially lost in both experimental set-ups (chi^2=9.25, *p*<0.005 for the first experiment and

chi^2=4.44, *p*<0.025 in the second experiment).

We investigated whether this loss of nuclei could be due to the relative fitness of the two

types of nuclei, by investigating whether the two CBS815.71 homokaryons, carrying either the *MAT1-*

1 or *MAT1-2* mating type, displayed differences in growth rate or biomass production, and comparing

them with the *MAT1-1/MAT1-2* heterokaryon. We measured growth rates on three different media and

determined the dry weight obtained after growth on M2 medium for both homokaryons and the

heterokaryon (Table S1A). No significant differences in growth or biomass were found between the

homokaryons or between the homokaryons and the heterokaryon, the only significant effect being that

of culture medium on growth rate (ANOVA, Table S1B). The growth rate of the *MAT1-1* homokaryon,

corresponding to the mating type most often lost during growth, was slightly lower, although not

significantly so (Table S1A).

***Reference genome assembly for* Schizothecium tetrasporum**

Using both Illumina and Oxford Nanopore MinION technology, we sequenced the genomes of the two

homokaryotic strains of opposite mating types, CBS815.71-sp3 and CBS815.71-sp6. We built a *de*

novo long read-based genome assembly for each, the first such assemblies to have been generated

for *S. tetrasporum*, to our knowledge. For CBS815.71-sp3, we also had an Illumina-based assembly

available that had been obtained before the long-read assemblies. BLAST searches and analyses of

Illumina read coverage against the CBS815.71-sp3 long read-based assembly suggested the

presence of bacterial contamination in the Nanopore data, but not in the Illumina data. We discarded

scaffolds with high similarity scores to bacterial genomes in BLAST analyses for further analyses. The

resulting long read-based genome assemblies for CBS815.71-sp3 and CBS815.71-sp6 contained 16

and 23 contigs and were 32.88 Mb and 32.83 Mb long, respectively. We found 96% of the 3,817

highly conserved orthologs in Sordariomycota from BUSCO analyses on the assemblies (Table S2).

In the CBS815.71-sp3 assembly, seven contigs were more than 3.5 Mb long and contained all the

conserved BUSCO orthologs. One contig corresponded to the mitochondrial genome and the other 8

contigs were less than 70 kb long, which is consistent with the expected number of chromosomes in

Sordariales, an order in which several species have seven chromosomes (28,37). Telomeric repeats

were present at both ends of contigs 1, 2, 4 and 5 in the CBS815.71-sp3 assembly, indicating that

these contigs represent four entirely assembled chromosomes (Figure 3A). For CBS815.71-sp3, we

compared the long read-based assembly obtained with the more fragmented Illumina-based assembly

and found high levels of similarity and synteny (Figure S1). We used the long read-based assembly of

CBS815.71-sp3 for further analyses. We found a high overall level of collinearity between the two

homokaryotic long read-based genome assemblies for CBS815.71-sp3 and CBS815.71-sp6 (Figure

3B). The 2305 indels between assemblies that we detected were scattered throughout the genome

and all were less than 10 bp long. These small indels were probably due to an artifactual enrichment

of low-complexity regions in homopolymers in the Nanopore data and were not considered for

subsequent analyses.

Repetitive sequences accounted for about 7% of the assemblies. In the CBS815.71-sp3

assembly, we identified 182 transposable element (TE) family consensus sequences: 4.2% Gypsy

LTRs and 6.7% other LTRs, 7% non-LTR SINEs and LINEs, 7.5% TcMar-Fot1 DNA transposons,

3.4% other DNA transposons and the remaining 70.5% repetitive sequences could not be annotated

(Figure S2A). Both genomes contained very few (<0.1 %) signatures of RIP (repeat-induced point

mutations), a fungal genome defense against transposable elements inducing C to T mutations in

repeated sequences at target dinucleotides (38,39).

Structure of the mating-type locus in Schizothecium tetrasporum

The mating-type locus was located at about the 5.180 Mbp position on contig 1 of the CBS815.71-sp3

assembly, which was considered likely to correspond to a fully assembled chromosome (Figure 3).

The mating-type locus of *S. tetrasporum* had the typical structure of heterothallic Sordariales (20),

with two highly divergent "alleles", referred to as idiomorphs, flanked by the *APN2* and *SLA2* genes

(Figure 3C; Figure S3, (40,41)). The *MAT1-2* idiomorph, present in CBS815.71-sp6, harbored a single

functional gene (*MAT1-2-1*), orthologous to the *mat+ FPR1* (*Pa_1_20590*) gene of *P. anserina* and

the *mat_a-1* gene of *Neurospora crassa.* The other idiomorph, *MAT1-1*, carried three genes (*MAT1-1-*

1, *MAT1-1-2* and *MAT1-1-3*), orthologous and syntenic with the three genes present in the *mat-*

idiomorph of *P. anserina* (*Pa_1_20593*, *_20592*, *_20591*) and the *mat_A* idiomorph of *N. crassa*

(NCU01958, NCU01959, NCU01960). MAT1-1-1 contained the typical MATα_HMG domain

(pfam04769) and *MAT1-1-2* contained a HPG domain (pfam17043). *MAT1-1-3* and *MAT1-2-1* carried

a MATA_HMG domain (cd01389). The 3' end of the *MAT1-1-1* coding sequence was found to extend

outside the *MAT1-1* idiomorph in *S. tetrasporum*, as shown by its presence next to the *MAT1-2*

idiomorph*.* This truncated *MAT1-1-1* gene close to the *MAT1-2* idiomorph has a putative open reading

frame starting 68 residues downstreaming of the start codon of the typical *MAT1-1-1* gene structure

(Figure S3). However, this putative start codon is in a poor context for translation initiation, with a

pyrimidine in position -3 (40). The truncated *MAT1-1-1* gene encodes most of the MATα_HMG

domain, and the second conserved domain adjacent to the MATα_HMG domain (Figure S3; (42)).

This gene was therefore named *TαD1* (truncated MATα_HMG domain). This name does not follow

the standard mating-type gene nomenclature (34), as it is not included in the mating-type locus. TαD1

may be a pseudogene and its transcriptional status is unknown in *S. tetrasporum*. A similar truncated

MAT1-1-1 gene is present and has little or no expression in *MAT1-2* strains in the Chaetomiaceae

(Sordariales) *Myceliophthora heterothallica*, *M. fergusii*, *M. hinnulea* and *Humicola hyalothermophila*

(43). It is unknown whether *TαD1* has any role in sexual reproduction.

A region with stepwise recombination suppression around the mating-type locus in

Schizothecium tetrasporum, *located 1 Mb from the centromere*

We plotted the synonymous divergence (d_S) between the two genomes of opposite mating types for

each gene along the contigs of the CBS815.71-sp3 assembly (Figure 4; Figure S4). Most of the d_S

values obtained were zero, as expected in a selfing species with a high degree of homozygosity. Only

around the mating-type locus did we find a cluster of genes with high d_S values (Figure 4), indicating

recombination suppression. The high degree of homozygosity throughout the genome indicates that

many selfing events occurred in the ancestors of the CBS815.71 heterokaryotic strain; only the

regions fully linked to the mating-type locus have remained heterozygous under selfing (27). Indeed,

for any gene linked to the mating-type locus, alleles associated with the alternative mating types

gradually accumulate different mutations over the time since recombination suppression.

Synonymous divergence (d_S) can, thus, be used as a proxy for the age of the linkage between a gene
and the mating-type locus (7).

The cluster of genes with high d_S values around the mating-type locus spanned a 1.47 Mb
region located between 4.983 Mbp and 6.457 Mbp on scaffold 1 of the CBS815.71-sp3 assembly,
encompassing 406 genes in the CBS815.71-sp3 assembly and 413 genes in the CBS815.71-sp6
assembly. The d_S values were lowest in the center of the non-recombining region (Figure 4),
suggesting the occurrence of gene conversion. We found no predicted function potentially directly
related to mating compatibility for any of the 406 genes in the non-recombining region of the
CBS815.71-sp3 assembly other than the mating-type locus. A total of 15 genes in the non-
recombining region encoded proteins that were identified as putative transcription factors or involved
in signaling pathways (Table S3). The four orthologs of the *P. anserina* genes located in its non-
recombining region and known to be involved in post-mating sexual development (but not mating
compatibility) were also found in the non-recombining region in *S. tetrasporum*: *ami1*, controlling
nuclear migration events (44), *PaRID*, a DNA methyltransferase controlling sexual development in *P.*
anserina (45), *AS4,* encoding the eEF1A translation elongation factor and *su12,* encoding a ribosomal
protein, mutations of which often result in sterility in *P. anserina* (46,47).

The d_S values plotted along the mating-type chromosome seemed to indicate the existence of
three distinct evolutionary strata (Figure 4; Figure S6). We tested the existence of evolutionary strata
more formally, by splitting the heterozygous region with sliding limits in the CBS815.71-sp3 assembly
and calculating, for each limit, the difference in d_S values between the two segments; limits between
strata should correspond to a peak in d_S value differences according to stratum definition (27). We
first arbitrarily set the limit of the first stratum by eye at 5,350,000 bp and applied the automatic
procedure to the rest of the heterozygous region. The d_S values peaked at 5,768,335 bp, identifying
the limit between strata 1 and 2 (Figure S6). We then re-applied the procedure to the part of the
heterozygous region without stratum 2. The d_S values peaked at 5,388,708 bp, identifying the limit
between strata 1 and 3 (Figure S6). The median d_S values were significantly different between strata
1 and 2 (pairwise Wilcoxon test *p*-value = 0.00134 with the False Discovery Rate (FDR) correction)
and between strata 1 and 3 (pairwise Wilcoxon test *p*-value = 0.00015 with FDR correction),
supporting the existence of distinct evolutionary strata. Median d_S values did not differ significantly
between strata 2 and 3, located on opposite sides of stratum 1 (pairwise multiple-comparison

Wilcoxon test *p*-value = 0.27571 with FDR correction). Strata 1 and 3 each spanned 400 kb, and

stratum 2 spanned about 700 kb. The most ancient evolutionary stratum, stratum 1, with the highest

d_S values, does not encompass the mating-type locus. The mating-type locus was located in the

middle of stratum 3 (Figure S6).

We looked for the centromere on the mating-type chromosome because its position may

provide important insight into the segregation of the mating-type locus at the first meiotic division. We

found a single location with a drop in GC content along the mating-type chromosome of the

CBS815.71-sp3 assembly outside of the telomeric regions, with 53.5% GC between 3.990 Mbp and

4.025 Mbp, versus the genome-wide mean value of 58.76% (Figure S5). A drop in GC content can

pinpoint the centromere in some fungi, such as some *Neurospora spp.* in which GC content drops to

about 44%, whereas the genome-wide mean value is about 53% (48). The lower variation of GC

content observed in the CBS815.71-sp3 genome assembly may be due to a lack of RIP in *S.*

tetrasporum. For the other three fully assembled chromosomes of the CBS815.71-sp3 genome,

similar drops in GC content were also identified at a single location per chromosome, co-localizing

with clusters of repeats. We characterized the repeat content of these four putative centromere

regions. We found no specific repeats, but we did detect an enrichment in *TcMar-Fot1* DNA

transposons relative to the genome-wide mean (Figure S2A). The putative centromere of the mating-

type chromosome was located about 1 Mb from the edge of the non-recombining region around the

mating-type locus.

We found no enrichment in repetitive sequences in the non-recombining region around the

mating-type locus relative to autosomes and the recombining regions on the mating-type chromosome

in the CBS815.71-sp3 assembly. The mean repetitive sequence densities calculated in 10 kb

windows overlapping by 1 kb were 2.52%, 5.05% and 4.21% in the non-recombining region around

the mating-type locus, the autosomes and the recombining regions on the mating-type chromosome,

respectively; pairwise Wilcoxon tests revealed no significant differences in repetitive sequence

densities. We found no difference in TE family proportions between these regions either (Figure S2B).

Repetitive sequence density was significantly higher in stratum 2 than in stratum 3 (*p*-value for a

pairwise Wilcoxon test = 0.0053 with FDR correction). We found no genes with a d_N/d_S ratio >1 in the

non-recombining region, i.e. with higher non-synonymous than synonymous substitution rates (the

mean d_N/d_S was 0.11 for genes with positive d_S values). As noted above, we found no evidence for

genomic rearrangements between the two mating-type chromosomes in the non-recombining region

(Figure 3B).

Validation of the region without recombination by progeny analysis

We investigated the lack of recombination in the heterozygous region around the mating-type locus in

the CBS815.71 strain. We obtained offspring resulting from a selfing event in the CBS815.71 strain,

i.e., a cross between CBS815.71-sp3 and CBS815.71-sp6. We genotyped 107 offspring resulting

from haploid spores isolated from five-spore asci (i.e., asci with three large dikaryotic spores and two

small monokaryotic spores; Figure 1). We initially used two genetic markers, one at each end of the

heterozygous region, and then three additional markers in the three recombinant offspring to delimit

more finely the regions in which recombination events occurred (Figure S7). We detected no

recombination events at the edge of stratum 3 (*i.e*., between the *P_NRR2* marker and the mating-type

locus) but we found recombination events at the edge of stratum 2 (between the markers *P_NRR4*

and *P_NRR5*) in three of the 107 offspring. We found no recombination event between the mating-

type locus and the additional markers *P_NRR7*, *P_NRR6* and *P_NRR5* located 0.808 Mb, 1.054 Mb

and 1.088 Mb from the mating-type locus, respectively. Given the position of the MAT locus, this

suggests a complete lack of recombination over a 1.3 Mb region in strata 1 and 3, and in a large part

of stratum 2, as recombination occurred only at the edge of this stratum. Stratum 2 is, therefore, likely

slightly smaller than the heterozygous region in the CBS815.71 strain, the heterozygosity at its edge

probably resulting from an outcrossing event a few generations ago.

Orthologs in the non-recombining regions across Sordariales

We aimed at investigating gene orthology relationships and synteny between the non-recombining

regions on the mating-type chromosomes of the three pseudo-homothallic species, *S. tetrasporum*, *P.*

anserina and *N. tetrasperma.* However, several rearrangements occurred following recombination

suppression on the mating-type chromosomes in *N. tetrasperma.* We therefore used the genome of

the closely related heterothallic *N. crassa* species instead, as a proxy for the ancestral gene order in

N. tetrasperma before recombination suppression. No recombination suppression around the mating-

type locus has been reported in *N. crassa* (26,49). This analysis therefore also provided an

opportunity for an interesting contrast with species displaying recombination suppression. In an
analysis of single-copy orthologs, we found that the mating-type chromosomes, and all the other
chromosomes, displayed regions of synteny, but also breaks of synteny between species, and
mesosynteny, *i.e.,* a global conservation of gene content despite rearrangements in order and
orientation (Figure 5A; Figure S8), as expected given their phylogenetic distance (50). In particular, 69
orthologs common to the species studied and belonging to stratum 3 of *S. tetrasporum* had a well-
conserved gene order in a 300 kb region around the mating-type locus (Figure 5A), suggesting some
conservation of the ancestral gene order in the region around the mating-type locus in these distantly
related pseudo-homothallic species, and in other non-pseudo-homothallic Sordariales, as previously
reported (32). The region around the putative centromere identified on the *S. tetrasporum* mating-type
chromosome had orthologous genes and gene orders conserved with the *P. anserina* and *N. crassa*
centromeric region over a distance of 700 kb despite an inversion, providing support for the putative
location of the centromere on the *S. tetrasporum* mating-type chromosome identified above (Figure
5A).

We also found conserved gene content and synteny between parts of the non-recombining
regions in the three species despite rearrangements. In *S. tetrasporum*, 261 of the 406 genes located
in the identified region of suppressed recombination around the mating-type locus had orthologous
genes in both *P. anserina* and *N. crassa.* We found that 138 of these 261 genes were also located in
the region of suppressed recombination around the mating-type locus in the *P. anserina* S strain
(Figure 5B). Most of these shared genes (85%) were part of stratum 3, encompassing the mating-type
locus, and stratum 1, the oldest evolutionary stratum in *S. tetrasporum* (Figure 5A(2),5B). No
evolutionary strata have previously been identified in *P. anserina*, but the oldest evolutionary stratum
in *P. pseudocomata,* a cryptic species closely related to *P. anserina* (27), was orthologous to stratum
3 in *S. tetrasporum*. Most genes (82%) of stratum 2 in *S. tetrasporum* had orthologs in *P. anserina* in
the recombining region between the mating-type locus and the centromere (Figure 5A(ii) & 5B). The
region of recombination suppression was located from the mating-type locus in the direction of the
telomere in *S. tetrasporum*, whereas it was located from the mating-type locus in the direction of the
centromere in *P. anserina* and other species of the *P. anserina* species complex (Figure 5A(ii)&5B).
The distance between the centromere and the mating-type locus was 1.1 Mb in *S. tetrasporum* and
2.9 Mb in *P. anserina.*

We further found that 188 of the 406 genes in the non-recombining region of *S. tetrasporum,*

(mostly in strata 3 and 2) were also present in the non-recombining region in *N. tetrasperma* (Figure

5B), with the genes from *S. tetrasporum* stratum 1 present in the flanking region of the non-

recombining region in *N. tetrasperma.* In total, 90 genes were common to the non-recombining

regions of the three pseudo-homothallic species, including genes at the two edges of these non-

recombining regions (Figure 5A). The predicted functions of these 90 orthologs, when available, did

not appear to be related to mating or mating-type functions (Table S4).

DISCUSSION

Striking evolutionary convergence

This study of the breeding system and sexual biology of the little-studied pseudo-homothallic fungus

S. tetrasporum reveals a striking convergence with *P. anserina* and other cryptic species of the *P.*

anserina complex, for multiple ecological and reproductive traits. These two distantly related fungi

thrive in herbivore dung and produce predominantly dikaryotic spores carrying opposite mating types.

In both cases, this is due to mating-type segregation during the second meiotic division associated

with a single systematic CO event between the centromere and the mating-type locus. In addition,

both *S. tetrasporum* and *P. anserina* display recombination suppression around this locus ((23) and

this study). We found that recombination suppression had expanded in several successive steps in *S.*

tetrasporum, as in *P. peudocomata,* which belongs to the *P. anserina* species complex (27). The

association between pseudo-homothallism (self-fertile dikaryotic spores), recombination suppression

around the mating-type locus and evolutionary strata also evolved independently in the *N.*

tetrasperma complex, representing an additional case of convergence. However, unlike *S.*

tetrasporum and *P. anserina, N. tetrasperma* displays recombination suppression up to the

centromere and mating-type segregation during the first meiotic division (26,48).

Association between pseudo-homothallism and recombination suppression: evolutionary and

proximal causes

Such a convergent association between recombination suppression and pseudo-homothallism (with

the production of heterokaryotic spores carrying both mating types and growth as a heterokaryotic

mycelium rather than as a homokaryon in the predominant phase of the lifecycle, as in most

ascomycetes) suggests causal relationships. Suppression of recombination around the mating-type

locus may be selected for if it favors the production of heterokaryotic ascospores, thus facilitating the

compaction of nuclei of opposite mating types in dispersing spores under the particular pattern of

meiotic segregation. The resulting self-fertile heterokaryotic ascospores allow reproductive insurance

in short-lived habitats such as dung. However, the occurrence of a CO between the centromere and

the proximal end of the region of suppressed recombination in almost all meiosis I to cause mating-

type segregation during the second meiotic division still calls for an explanation. Indeed, if we

consider the 21 ± 3 COs occurring at each meiosis in one nucleus of *Sordaria macrospora* (51,52) as

a proxy for the number of COs occurring in *S. tetrasporum* displaying approximately the same

genome size, and if we take into account the number of obligate COs occurring on each of the seven

chromosomes, we can estimate that 11 to 17 COs remain to be randomly distributed across a

genome of 32 Mbp. It is therefore unlikely that one CO takes place between the centromere and the

mating-type locus in 99.5 % of the meioses in *P. anserina*, the distance between the centromere and

the mating-type locus being 2.9 Mbp, and in *S. tetrasporum,* the distance between the centromere

and the mating-type locus being 1.2 Mbp. The occurrence of a second CO between the centromere

and the mating-type locus should be prevented, as two COs in this region would result in FDS. This

second CO might be prevented by positive interference, but this mechanism alone may not be

sufficient to reach the very high frequency of SDS.

It therefore appears more likely that a yet unidentified mechanism has evolved that enforces

the occurrence of the single and systematic CO event between the centromere and the mating-type

locus, independently in both *P. anserina* and *S. tetrasporum*, together with a mechanism compacting

two nuclei per ascospore, for the production of self-fertile ascospores. Suppression of recombination

around the mating-type locus may be directly linked to the mechanism that prevents a second CO

around the mating type locus, which would yield FSD instead of SDS. The selective advantage of

producing self-fertile spores would then be the driver to develop this mechanism.

An alternative hypothesis may be proposed considering that the mating-type locus is always

heterozygous in heterokaryons. Indeed, a recent theoretical model showed that recombination

suppression capturing a permanently heterozygous locus may be selected due to its capacity to

shelter deleterious alleles in organisms with an extended diploid or dikaryotic phase (14). More

precisely, recombination suppression may be selected because it protects DNA fragments with fewer

deleterious mutations than average in the population, but the load of these mutations will still prove

problematic when they become frequent enough to become homozygous; the capture of a

permanently heterozygous allele prevents this effect (14). This mechanism can also explain the

evolution of successive evolutionary strata, as observed here in *S. tetrasporum* and, previously, in *N.*

tetrasperma and *P. pseudocomata* (22,27,27,48), the basidiomycete mushroom *Agaricus bisporus*

(53) and repeatedly in *Microbotryum* basidiomycete fungi, plant pathogens in natural ecosystems

(8,9,18,54). This mechanism may also account for the suppression of recombination in *N.*

tetrasperma.

In this study, we found no detectable differences in growth between homokaryons of the two

mating types and the heterokaryon in *S. tetrasporum*, but we found that one of the mating-type alleles

was more often lost during mycelial growth. In addition, fitness differences even smaller than those

detectable here can promote the evolution of recombination suppression (14), and genes with

deleterious alleles may be involved in specific parts of the life cycle not evaluated *in vitro,* and related

to the dung habitat or resistance to harsh conditions. We found that 13% of the thalli grown from

ascospores *in vitro* ended up homokaryotic, and the mechanism involved could function in

haplodiplontic lifecycles, depending on the mean length or frequency of the haploid phase and the

strength of selection (14). The mycelia in dung may not grow for long enough for the loss of a nucleus

to be occur very often, especially as *S. tetrasporum* is mostly found on small pellets of rabbit dung

(unpublished data).

The finding that the non-recombining regions in *P. anserina, N. tetrasperma* and *S.*

tetrasporum contain a number of shared orthologous genes, particularly at the two edges of the non-

recombining region, and that these regions display some level of synteny, raises the possibility of a

third hypothesis: the selection of recombination suppression to maintain beneficial allelic combinations

(55), as observed for supergenes in plants and animals (56,57). No function known to be directly

related to mating compatibility was found among these orthologous genes, except for the mating-type

locus, but several genes involved in sexual development in *P. anserina* are located in the non-

recombining regions in both *P. anserina* (23) and *S. tetrasporum* (this study): *ami1*, controlling nuclear

migration events (44) and *PaRID*, a RID-like putative cytosine methyltransferase that controls sexual

development (45). However, there is no evidence of antagonistic selection between mating types for

these functions (23) and we found no phenotypic differences between homokaryons of opposite

mating types. Nevertheless, we cannot rule out the possibility that there are differences in other parts

of the life cycle not investigated here *in vitro*.

These various evolutionary hypotheses for explaining recombination suppression are not

mutually exclusive, and the evolutionary causes of recombination suppression may differ between

evolutionary strata and species. The proximal (mechanistic) causes of recombination suppression

also remain unknown. We detected no chromosomal inversion between mating-type chromosomes in

S. tetrasporum, as in some other fungi in which recombination is also suppressed close to the mating-

type locus without inversion, such as *P. anserina* and other species of the *P. anserina* complex, *N.*

tetrasperma and *Microbotryum* fungi (8,23,26,27). Histone modification can also prevent

recombination (58,59) and this aspect warrants investigation in *S. tetrasporum*. There may also be

recombination modifiers in *cis*. The proximal cause for the single CO event between the centromere

and the mating-type locus is also unknown. The shorter genomic region between the centromere and

the mating-type locus in *S. tetrasporum* (1.2 Mb) than in *P. anserina* (2.9 Mb) makes this species an

excellent model for identifying the proximal factors responsible for the single CO event, if they are

located in *cis*. The convergent evolution of a single systematic CO event between the mating-type

locus and the centromere in these two species provides an opportunity to study the genetic basis of

the convergent evolution of this phenomenon.

Evolutionary strata

We detected three evolutionary strata in the CBS815.71 *S. tetrasporum* strain, indicating a stepwise

extension of recombination suppression. Our progeny analysis showed that stratum 2 did not cover

the entire corresponding heterozygous region, the extremity of the heterozygous region probably

being a remnant of an outcrossing event a few generations ago. Nevertheless, the progeny analysis

showed that strata 1 and 3, and much of stratum 2, probably display a full suppression of

recombination. This gradual progression of recombination suppression cannot be explained by sexual

antagonism, as male and female functions are not associated with mating types in fungi (19,60). In

addition, monokaryotic ascospores are rare in *S. tetrasporum* and traits and ecological features are

similar between mating types, so other types of antagonistic selection are also unlikely. Stepwise

recombination suppression in this dikaryotic fungus may have been selected to shelter deleterious

alleles (14). It may appear surprising that the most ancient evolutionary stratum, stratum 1, with the

highest d_S values, does not encompass the mating-type locus. However, this oldest stratum is very

close to the mating-type locus and the deleterious allele sheltering model predicts the possible

evolution of recombination suppression at sites tightly linked to but not encompassing the

permanently heterozygous locus (14). This situation has, in fact, been observed in the chestnut blight

fungus *Cryphonectria parasitica* (61). It is also possible that the most ancient evolutionary stratum

encompassed the mating-type locus but that the limits of the stratum were obscured by subsequent

gene conversion events. Alternatively, the different strata may correspond to gradual increases in the

SDS frequency, by stepwise increase of the efficiency of the mechanism preventing a second CO in

the region of mating-type locus and FDS. The selective pressure for this stepwise increase would be

to reach a higher number of dikaryotic ascospores growing into self-fertile mycelia.

No sign of degeneration in the non-recombining region

We found no enrichment in repeats, rearrangements or non-synonymous substitutions, whereas

transposable elements, non-synonymous substitutions and non-optimal codon usage have all been

found to increase with time since recombination suppression in the mating-type chromosomes of

Microbotryum anther-smut fungi (9,54,62). Gene disruptions due to small indels remain to be

investigated. The suppression of recombination may have occurred too recently in *S. tetrasporum*, or

recombination or gene conversion may occur occasionally, as suggested in *P. anserina* (27). The low

d_S values in the center of the non-recombining region, between strata 1 and 2, suggests that gene

conversion may occur, as observed in the non-recombining regions of mating-type chromosomes in

other fungi (8,9,23,63). We found no footprints of RIP, the fungal genome defense mechanism that

introduces cytosine-to-thymine transitions into repetitive sequences to control transposable elements

(38,39). Nevertheless, the TE load was not high in *S. tetrasporum*, suggesting that this fungus may

have evolved another mechanism of transposable element control, such as RNA silencing (64).

***Mating system in* Schizothecium tetrasporum**

We found a higher percentage of homokaryotic thalli resulting from ascospore germination in *S.*

tetrasporum than in *P. anserina* (13% versus 1%). Homokaryons may be recovered due to first-

division segregation, the loss of one nucleus after germination or a particular pattern of nucleus

migration following the post-meiosis mitosis. We found that nuclei could be lost during vegetative

growth *in vitro* in *S. tetrasporum* (here in 13% of the germinating ascospores), whereas this is not the

case in *P. anserina* (47), suggesting that *S. tetrasporum* may sometimes, albeit rarely, end up

homokaryotic in nature. Such nucleus losses may be due to active competition between nuclei

(65,66) or the presence of deleterious alleles associated with a mating type. The occurrence of

mycelia of a single mating type could promote some degree of outcrossing, which could be checked in

analyses of the levels of heterozygosity of dikaryons in natural populations. Nevertheless, the rate of

self-fertile (heterokaryotic) ascospores was high, and this would be expected to promote a high level

of selfing. Population genomics approaches detected widespread genome-wide homozygosity in

dikaryons of *P. anserina*, indicating high selfing rates, but with occasional stretches of heterozygosity,

also indicating the occurrence of outcrossing in nature (27,67,68). The optimal media and conditions

for vegetative growth and sexual reproduction identified here will facilitate further investigations of the

lifecycle and biology of *S. tetrasporum*. The reference strain was found to be highly homozygous, as

expected under high rates of selfing. If we assume that the non-recombining region evolved to prevent

a second CO, to maximize the frequency of SDS and thereby of self-fertile spores, it may appear

intriguing that no mechanism evolved to subsequently maintain a dikaryotic mycelium, as for instance

the crozier cells that are present in the perithecia and are analogous to the dikaryotic clamp cells of

Basidiomycetes (69). It may be that, in nature, the mycelium does not grow long enough to frequently

loose a nucleus or that the occurrence of mycelia carrying a single mating type is beneficial in

promoting outcrossing, as does the formation of a few asci with 5 ascospores including two

monokaryotic ascospores.

Conclusion

In conclusion, we found a striking convergence in lifecycle, sexual biology and stepwise

recombination suppression around the mating-type locus in phylogenetically distant fungi of the order

Sordariales. Documenting convergent evolution is important, as it shows that evolution can be

repeatable, and not only contingent (70). Studies of these traits in additional species would make it

possible to perform a more robust assessment of the degree of convergence in trait association and,

thus, of the causal relationship. Furthermore, the broad convergence towards an association between

dikaryotic life cycle and stepwise recombination suppression beyond the mating-type locus despite

the lack of sexual antagonism provides support for alternative hypotheses, such as deleterious allele

sheltering, as an explanation for the suppression of recombination beyond mating-type or sex-

determining alleles (14).

MATERIALS AND METHODS

Strain origins, experimental crosses and culture

The dikaryotic strain CBS815.71 of *S. tetrasporum* was obtained from the Centraalbureau voor

Schimmelcultures (CBS). No optimized methods have been described for culturing *S. tetrasporum*

and inducing sexual reproduction in this species. We therefore tested several media, to select those

optimal for growth and sexual reproduction (M2, M3, Oat Meal Agar and V8 agar; (71)), and for

ascospore germination (the same media plus *P. anserina* germination medium (G), (71)). Growth and

sexual reproduction were optimal on V8 medium (composition per liter: 15 g agar, 2 g $CaCO_3$, 200 mL

V8 tomato juice and 800 mL water) at 22°C with 12 h of illumination. Incubation at a higher

temperature, such as 26°C, often resulted in aborted fruiting bodies. Growth and ascospore recovery

were very slow at lower temperatures (*e.g.*, 18°C). We observed 100% germination on G medium

(composition per liter: 4.4 g ammonium acetate, 15 g bactopeptone and 13 g agar) with a 65°C heat

shock for 30 minutes just after the plating of ascospores. On all the other media tested, germination

rates were lower, with no germination in the absence of a heat shock. In optimal conditions, mature

perithecia ejecting ascospores were obtained three weeks after inoculation on V8 medium, and

germination occurred overnight on G medium after a heat shock. Ascospores could be collected and

plated with the same methods as used for *P. anserina* (71).

In these optimal conditions, we generated and isolated two F1 monokaryotic strains by

inducing a sexual cycle in the self-fertile heterokaryotic culture received from CBS815.71: the

CBS815.71-sp3 strain with a mating-type allele orthologous to the *mat+* mating type of *Podospora*

anserina, corresponding to the *MAT1-1* allele according to the official nomenclature (34), and the

CBS815.71-sp6 strain with a mating-type allele orthologous to the *mat-* mating type of *P. anserina*,

corresponding to the *MAT1-2* allele according to the official nomenclature (34). We assessed the

frequency of second-division segregation (SDS) at the mating-type locus and, therefore, the

frequency of heterokaryotic spore formation, by generating and isolating 190 four-spore asci from a

cross between CBS815.71-sp3 and CBS815.71-sp6. We studied the size of the region in which

recombination was fully abolished around the mating-type locus, by isolating 107 small ascospores

from five-spore asci (monokaryotic) from a cross between CBS815.71-sp3 and CBS815.71-sp6. After

isolation, the strains were maintained on M2 minimal medium to prevent contamination (71), and were

incubated at 22°C for growth. The growth of CBS815.71-sp3 and CBS815.71-sp6 homokaryons and

the CBS815.71 dikaryotic strain was assessed on OA, V8 and M2 media (71). The diameter of

mycelial colonies was measured after 3 and 10 days of growth at 22°C with 12 h illumination, with a

10 cm ruler. Mean growth rate per day was calculated over the seven-day period from day 3 to day

10. Dry weights were measured after 10 days of growth in liquid M2 medium (50 mL); the mycelium

was filtered and dried at 65°C overnight. Measurements were performed on five Petri dishes for each

strain and each medium. ANOVA was performed with R software v 4.2.0.

Genome sequencing and other genomic data

We generated long read-based genome assemblies by sequencing the genomes of the *S.*

tetrasporum strains CBS815.71-sp3 and CBS815.71-sp6 with both Illumina technology on the

HiSeq2500 platform (150 bp long paired-end reads) and Oxford Nanopore MinION technology with an

R9 flow cell. For each DNA extraction, we prepared 100–200 mg aliquots of mycelium, which were

stored in a 1.5 mL Eppendorf tube at -80°C for at least 24 hours before lyophilization. A sheet of

cellophane was placed over the M2 medium before inoculation with spores, such that the mycelium

grew on the cellophane for three to four days and could be scraped off easily. For Illumina

sequencing, DNA was extracted with the commercial Nucleospin Soil kit from Macherey Nagel.

Illumina library preparation and genome sequencing were outsourced to the sequencing core facility

of Genoscope (last accessed May 15, 2021). For Nanopore sequencing, mycelium DNA was

extracted with the NucleoBond High Molecular Weight DNA kit from Macherey-Nagel, with the

mechanical disruption of about 30 mg of lyophilized mycelium with beads for 5 min at 30 Hz. The

Nanopore library was prepared with the SQK-LSK109 ligation sequencing kit, and sequencing was

performed in house. The quality of the raw reads was analyzed with fastQC 0.11.9 (72) for the

Illumina raw reads and pycoQC 2.5.0.23 (73) for the Nanopore raw reads (Table S5).

In addition, the genome and transcriptome of *S. tetrasporum* strain CBS815.71-sp3 were

sequenced with Illumina technology by the Joint Genome Institute (JGI). DNA was extracted following

(74) from mycelium grown on cellophane-covered potato dextrose agar medium. Plate-based DNA

library was prepared for Illumina sequencing on the PerkinElmer Sciclone NGS robotic liquid handling

system with the Kapa Biosystems library preparation kit. Sample DNA (200 ng) was sheared into 600

bp fragments with a Covaris LE220 focused ultrasonicator. The sheared DNA fragments were size-

selected by double-SPRI and the selected fragments were end-repaired, A-tailed, and ligated with

Illumina-compatible sequencing adaptors from IDT containing a unique molecular index barcode. RNA

was extracted from the same mycelial colonies as DNA using TRIzol (Invitrogen) following

manufacturer's instructions (https://assets.thermofisher.com/TFS-

Assets/LSG/manuals/trizol_reagent.pdf). Plate-based RNA sample preparation was obtained with the

PerkinElmer Sciclone NGS robotic liquid handling system and the Illumina TruSeq Stranded mRNA

HT sample preparation kit, with the poly-A selection of mRNA according to the protocol in the Illumina

user guide (https://support.illumina.com/sequencing/sequencing_kits/truseq-stranded-mrna.html),

under the following conditions: 1 µg total RNA as the starting material, and eight PCR cycles for

library amplification. The prepared libraries were then quantified with the KAPA Biosystem next-

generation sequencing library qPCR kit (Roche) and a Roche LightCycler 480 real-time PCR

instrument. The quantified libraries were then then multiplexed with other libraries, and the pool of

libraries was prepared for sequencing on the Illumina NovaSeq 6000 sequencing platform with

NovaSeq XP v1.5 reagent kits, an S4 flow cell, following a 2x150 indexed run method. All raw Illumina

sequence data were filtered for artifacts/process contamination. DNA reads produced by the JGI were

then assembled with SPAdes v3.15.2 (75). The obtained genome assembly is referred to as the

Illumina read-based assembly of CBS815.71-sp3.

The genome of the *P. anserina* S *mat+* strain "Podan2" (32) was downloaded from the JGI

MycoCosm website (https://mycocosm.jgi.doe.gov/mycocosm/home, last accessed November 15,

2018) and improved annotations by (76) were downloaded from the GitHub repository,

https://github.com/johannessonlab/SpokBlockPaper (last accessed June 1, 2020). The genome of the

OR74A *N. crassa mat A* strain (77) was downloaded from GenBank under accession number

GCA_000182925.2 (last accessed May 10, 2022).

Long read-based genome assemblies for CBS815.71 strains

De novo assemblies of the genomes of the CBS815.71-sp3 and CBS815.71-sp6 strains were constructed from both Illumina reads produced at Genoscope and Nanopore reads. For each genome, the raw Nanopore reads were trimmed and assembled with Canu v1.8 (78) with the option genomeSize=35m. The assembly obtained with Canu was polished twice with Nanopore trimmed reads using first Racon v1.4.3 (79) with the options m=8 x=6 g=8 w=500 and then with medaka (https://github.com/nanoporetech/medaka) with the default settings. The assembly obtained was polished five times with the trimmed Illumina reads with Pilon v1.24 (80). For each round of polishing, BWA v0.7.17 (81) was used to align the Nanopore or Illumina trimmed reads with the assembly for polishing. The Illumina reads were first trimmed with Trimmomatic v0.36 (82) with the options: ILLUMINACLIP:TruSeq3-PE.fa:2:30:10 LEADING:10 TRAILING:10 SLIDINGWINDOW:5:10 MINLEN:50. The quality of the final assemblies was assessed with Quast v5.1 (83). We checked for contamination, by performing a BLAST analysis of the assembly scaffolds against the online NCBI nucleotide collection database (https://blast.ncbi.nlm.nih.gov/, last accessed March 5, 2022) and looking at the coverage of Illumina and Nanopore reads on the final assemblies. We also evaluated the completeness of the final assemblies with BUSCO v5.2.2 (84,85) and the Sordariomyceta ortholog set (sordariomycetes_odb10, 2020-08-05, *n*= 3817 orthologous groups searched). We also ran BUSCO analysis on the Podan2 *P. anserina* assembly for comparison.

Genome annotation

Before gene annotation, the final assemblies were formatted (contigs were renamed and ordered in descending order of size) and masked for repeats with RepeatMasker (86) and the funnannotate v1.8.9 pipeline (87). We performed gene annotation on the masked assemblies with BRAKER v2.1.6, which uses a combination of the *ab initio* gene predictors Augustus and GeneMark-ET (88). We ran BRAKER twice, first using the BUSCO dataset of proteins sordariomycetes_odb10 (84) as evidence, and then using the RNAseq dataset as evidence. We optimized the training steps, by concatenating the two homokaryotic genomes derived from the same heterokaryotic strain, CBS815.71, into a single fasta file and running BRAKER on the combined fasta file. For RNAseq mapping, we used STAR v2.7.10a (89) with the default settings. We combined the results of the two BRAKER runs with TSEBRA (the Transcript Selector for BRAKER; https://github.com/Gaius-Augustus/TSEBRA), with the

option 'intron_support' set to 0.2. We ran InterProScan5 v5.54-87.0 (90), TMHMM v2.0c (91), SignalP

v4.1 (92) and Phobius (93) and obtained predicted protein annotations from PFAM, InterPro, EggNog,

UniProtKB, MEROPS, CAZyme, and GO ontology. For repeat annotation, we performed a *de novo*

identification of transposable elements in the final assemblies with RepeatModeler v1.0.11 (94). We

merged the consensus sequences obtained wiereth the RepBase library available in RepeatMasker

v4.0.9 (86) and the custom library "PodoTE-1.00" built for the *Podospora anserina* species complex

by (95) and the *Neurospora* library of (96) "Gioti_neurospora_repeats.renamed" (both available from

the GitHub repository https://github.com/johannessonlab/SpokBlockPaper, last accessed June 1,

2020). We used RepeatMasker v4.0.9 (86) to annotate repeats. We parsed the RepeatMasker

outputs and removed low-complexity and simple repeats with the parseRM_merge_interrupted.pl

script from https://github.com/4ureliek/Parsing-RepeatMasker-Outputs (last accessed February 25,

2021). We retained only TEs longer than 150 bp, and overlapping TEs annotated as belonging to the

same family were merged into single elements for the final annotation. We used the TE annotation

suggested by (97) to sort TEs by family. We looked for signatures of RIP with the RIPper platform

http://theripper.hawk.rocks (default settings; (98)). The mating-type locus was identified by tBLASTn,

with the *mat+* and *mat-* idiomorphs of *P. anserina* as queries. The *SMR1*, *SMR2*, *FMR1*, *FPR1* gene

models were manually curated as previously described (99), in Artemis v18.0.2 (100). The sequences

of proteins containing a MATα_HMG domain were downloaded from the European Nucleotide Archive

(ENA) database with accession numbers CDN29981 and ESA43845 for *P. anserina* and *N. crassa*,

respectively. Proteins were aligned with Clustal Omega (101) and the alignments were color-coded

with JALVIEW v2 (102).

Comparative genomics and statistical analyses

For all comparative genomics analyses, we identified orthologous groups with OrthoFinder v2.3.7

(103). We calculated d_S values by studying one-to-one orthologs between the CBS815.71-sp3 and

CBS815.71-sp6 genome assemblies and retaining only those orthologs with the same number of

exons and the same transcript length as predicted. We aligned orthologous gene coding sequences,

using the codon-based approach implemented in translatorX v1.1 with default parameters (104). We

used the nucleotide alignment as input for the yn00 program in the PAML v.4.9 package (105,106).

Pairwise Wilcoxon tests were performed in R with false discovery rate correction. We excluded the

four genes located at the edges of strata to assess the differences in d_S between strata. We studied

synteny between assemblies, by investigating either collinearity between one-to-one orthologs or

running the genome sequence aligner nucmer from MUMmer v4.0.0rc1 (107) with default options. For

the identification of small indels between the long-read assemblies, we first aligned the CBS815.71-

sp6 genome assembly with the CBS815.71-sp3 genome assembly using Minimap2 (108) with the -cx

asm10 -t8 options. We then used samtools mpileup (109) with the following options: -d 250 -m 1 -E --

BCF --output-tags DP,DV,DP4,SP.

Genotyping of the mating-type locus alleles and surrounding regions

We devised a new method of DNA extraction for genotyping based on a published protocol (110).

Briefly, 1 mm^3 plugs were taken directly from cultures growing on agar plates and ground with 200 µL

extraction buffer, heated at 95°C for 15 minutes, placed on ice for one minute, briefly vortexed and

centrifuged at 15,900 x g for 30 seconds. We determined the mating-type allele of the offspring from a

cross between CBS815.71-sp3 and CBS815.71-sp6 by PCR with the primers and conditions

described in Table S6 and Table S7. Primers were designed with Primer3 software (111,112), based

on the mating-type locus region annotated in the two genome assemblies obtained for *S. tetrasporum.*

We visualized amplicons by agarose-gel electrophoresis.

Data availability

All genomic data produced by the JGI are available from the JGI MycoCosm website

(https://mycocosm.jgi.doe.gov/mycocosm/home, last accessed February 5, 2022). The raw Illumina

and Nanopore reads and the long read-based assemblies for the CBS815.71-sp3 and CBS815.71-

sp6 genomes were deposited in GenBank under the BioProject accession number PRJNAXXX

(BioSample ID: xxx, assembly accession number: xxx; available upon publication).

Acknowledgements

This work was supported by the Louis D. Foundation award and EvolSexChrom ERC advanced grant

#832352 to T.G. and DOE-JGI CSP grant #504394 to P.G. and P.S. We are grateful to J.P. Vernadet

for sharing the conda environment to run the BRAKER pipeline.

Author contributions

T.G., F.E.H. and P.S. conceptualized the study, acquired funding and supervised the study. N.V.,

A.S., V.G., E.D.F., E.L., F.E.H., and P.S. contributed to experiments. N.V., F.E.H., R.D., C.L., P.S.

and R.R.d.l.V. analyzed the genomes. N.V., V.G., E.D.F., E.L. and P.S. analyzed the progenies. P.G.,

S.G., Y. Z., S. T. and I.G. provided RNAseq data and the Illumina based-genome assembly of the

CBS815.71-sp3 strains. N.V., F.E.H., P.S., and T.G. wrote the manuscript draft. All authors edited the

manuscript.

REFERENCES

1.   Muller HJ. Some genetic aspects of sex. Am Nat. 1932 Mar 1;66(703):118–38.

2.   Hill WG, Robertson A. The effect of linkage on limits to artificial selection. Genet Res. 1966
Dec;8(3):269–94.

3.   Uyenoyama MK. Evolution under tight linkage to mating type. New Phytol. 2005;165(1):63–70.

4.   Hartmann FE, Duhamel M, Carpentier F, Hood ME, Foulongne-Oriol M, Silar P, et al.
Recombination suppression and evolutionary strata around mating-type loci in fungi:
documenting patterns and understanding evolutionary and mechanistic causes. New Phytol.
2021;229(5):2470–91.

5.   Goubet PM, Bergès H, Bellec A, Prat E, Helmstetter N, Mangenot S, et al. Contrasted patterns
of molecular evolution in dominant and recessive self-incompatibility haplotypes in *Arabidopsis*.
PLOS Genet. 2012 Mar 22;8(3):e1002495.

6.   Coelho MA, Bakkeren G, Sun S, Hood ME, Giraud T. Fungal sex: The Basidiomycota. Microbiol
Spectr. 2017 Jun;5(3):FUNK-0046-2016.

7.   Bergero R, Charlesworth D. The evolution of restricted recombination in sex chromosomes.
Trends Ecol Evol. 2009 Feb 1;24(2):94–102.

8.   Branco S, Carpentier F, Rodríguez de la Vega RC, Badouin H, Snirc A, Prieur SL, et al. Multiple
convergent supergene evolution events in mating-type chromosomes. Nat Commun. 2018 May
21;9(1):2000.

849 9. Branco S, Badouin H, Rodríguez de la Vega RC, Gouzy J, Carpentier F, Aguileta G, et al.
850   Evolutionary strata on young mating-type chromosomes despite the lack of sexual antagonism.
851   Proc Natl Acad Sci. 2017 Mar 7;114(27):7067–72.

852 10. Hartmann FE, Duhamel M, Carpentier F, Hood ME, Foulongne-Oriol M, Silar P, et al.
853   Recombination suppression and evolutionary strata around mating-type loci in fungi:
854   documenting patterns and understanding evolutionary and mechanistic causes. New Phytol.
855   2021;229(5):2470–91.

856 11. Ironside JE. No amicable divorce? Challenging the notion that sexual antagonism drives sex
857   chromosome evolution. BioEssays. 2010 Aug 1;32(8):718–26.

858 12. Charlesworth. The status of supergenes in the 21st century: recombination suppression in
859   Batesian mimicry and sex chromosomes and other complex adaptations. Evol Appl.
860   2016;9(1):74–90.

861 13. Bazzicalupo AL, Carpentier F, Otto SP, Giraud T. Little evidence of antagonistic selection in the
862   evolutionary strata of fungal mating-type chromosomes (*Microbotryum lychnidis-dioicae*). G3
863   Genes Genomes Genet. 2019 Jun 1;9(6):1987–98.

864 14. Jay P, Tezenas E, Véber A, Giraud T. Modeling the stepwise extension of recombination
865   suppression on sex chromosomes and other supergenes through deleterious mutation
866   sheltering. PLOS Biol. 2022 in press. available at
867   https://www.biorxiv.org/content/10.1101/2021.05.17.444504v2.full.pdf

868 15. Lenormand T, Roze D. Y recombination arrest and degeneration in the absence of sexual
869   dimorphism. Science. 2022 Feb 11;375(6581):663–6.

870 16. Kent TV, Uzunović J, Wright SI. Coevolution between transposable elements and
871   recombination. Philos Trans R Soc B Biol Sci. 2017 Dec 19;372(1736):20160458.

872 17. Gladieux P, Ropars J, Badouin H, Branca A, Aguileta G, de Vienne DM, et al. Fungal
873   evolutionary genomics provides insight into the mechanisms of adaptive divergence in
874   eukaryotes. Mol Ecol. 2014 Feb 1;23(4):753–73.

875 18. Duhamel M, Carpentier F, Begerow D, Hood ME, Rodríguez de la Vega RC, Giraud T. Onset
876   and stepwise extensions of recombination suppression are common in mating-type
877   chromosomes of *Microbotryum* anther-smut fungi. J Evol Biol. Available from:
878   https://onlinelibrary.wiley.com/doi/abs/10.1111/jeb.13991

879 19. Billiard, López-Villavicencio M, Hood ME, Giraud T. Sex, outcrossing and mating types:
880   unsolved questions in fungi and beyond. J Evol Biol. 2012;25(6):1020–38.

881 20. Dyer PS, Inderbitzin P, Debuchy R. 14 mating-type structure, function, regulation and evolution
882   in the Pezizomycotina. In: Wendland J, editor. Growth, Differentiation and Sexuality. Cham:
883   Springer International Publishing; 2016. p. 351–85. (The Mycota). Available from:
884   https://doi.org/10.1007/978-3-319-25844-7_14

885 21. Raju NB, Perkins DD. Diverse programs of ascus development in pseudohomothallic species of
886   *Neurospora*, *Gelasinospora*, and *Podospora*. Dev Genet. 1994;15(1):104–18.

887 22. Menkis A, Jacobson DJ, Gustafsson T, Johannesson H. The mating-type chromosome in the
888   filamentous ascomycete *Neurospora tetrasperma* represents a model for early evolution of sex
889   chromosomes. PLOS Genet. 2008 Mar 14;4(3):e1000030.

890 23. Grognet P, Bidard F, Kuchly C, Tong LCH, Coppin E, Benkhali JA, et al. Maintaining two mating
891   types: structure of the mating type locus and its role in heterokaryosis in *Podospora anserina*.
892   Genetics. 2014 May 1;197(1):421–32.

24.   Boucher C, Nguyen TS, Silar P. Species delimitation in the *Podospora anserina/p. pauciseta/p.*
comata species complex (Sordariales). Cryptogam Mycol. 2017 Dec;38(4):485–506.

25.   Corcoran P, Dettman JR, Sun Y, Luque EM, Corrochano LM, Taylor JW, et al. A global
multilocus analysis of the model fungus *Neurospora* reveals a single recent origin of a novel
genetic system. Mol Phylogenet Evol. 2014 Sep 1;78:136–47.

26.   Sun Y, Svedberg J, Hiltunen M, Corcoran P, Johannesson H. Large-scale suppression of
recombination predates genomic rearrangements in *Neurospora tetrasperma*. Nat Commun.
2017 Oct 26;8(1):1–8.

27.   Hartmann FE, Ament-Velásquez SL, Vogan AA, Gautier V, Le Prieur S, Berramdane M, et al.
Size variation of the nonrecombining region on the mating-type chromosomes in the fungal
Podospora anserina species complex. Mol Biol Evol. 2021 Jun 1;38(6):2475–92.

28.   Heslot H. Contribution à l'étude cytogénétique et génétique des Sordariacées. 1958.

29.   Marin-Felix Y, Miller AN, Cano-Lira JF, Guarro J, García D, Stadler M, et al. Re-evaluation of the
order Sordariales: Delimitation of Lasiosphaeriaceae s. str., and introduction of the new families
Diplogelasinosporaceae, Naviculisporaceae, and Schizotheciaceae. Microorganisms. 2020 Sep
17;8(9):E1430.

30.   Huang SK, Hyde KD, Mapook A, Maharachchikumbura SSN, Bhat JD, McKenzie EHC, et al.
Taxonomic studies of some often over-looked Diaporthomycetidae and Sordariomycetidae.
Fungal Divers. 2021 Nov 1;111(1):443–572.

31.   Hyde KD, Maharachchikumbura SSN, Hongsanan S, Samarakoon MC, Lücking R, Pem D, et al.
The ranking of fungi: a tribute to David L. Hawksworth on his 70th birthday. Fungal Divers. 2017
May 1;84(1):1–23.

32.   Espagne E, Lespinet O, Malagnac F, Da Silva C, Jaillon O, Porcel BM, et al. The genome
sequence of the model ascomycete fungus *Podospora anserina*. Genome Biol. 2008 May
6;9(5):R77.

33.   Esser K, Prillinger H. A new technique for using spermatia in the production of mutants in
Podospora. Mutat Res Mol Mech Mutagen. 1972 Dec 1;16(4):417–9.

34.   Turgeon BG, Yoder OC. Proposed nomenclature for mating type genes of filamentous
Ascomycetes. Fungal Genet Biol. 2000 Oct 1;31(1):1–5.

35.   Marcou D. Notion de longévité et nature cytoplasmique du déterminant de la sénescence chez
quelques champignons. Annales des Sciences Naturelles, Botaniques Paris. 1961;Sér. 12, 2:
653-764.

36.   Roper M, Simonin A, Hickey PC, Leeder A, Glass NL. Nuclear dynamics in a fungal chimera.
Proc Natl Acad Sci U S A. 2013 Aug 6;110(32):12875–80.

37.   Barth G. Genetic Maps 1987: a compilation of linkage and reconstruction maps of genetically
studied organisms, Volume 4. IX + 755 S. Cold Spring Harbor 1987. Cold Spring Harbor
Laboratory. ISBN: 0-87969-303-7. J Basic Microbiol. 1989;29(3):190–190.

38.   Cambareri EB, Jensen BC, Schabtach E, Selker EU. Repeat-induced G-C to A-T mutations in
Neurospora. Science. 1989 Jun 30;244(4912):1571–5.

39.   Galagan JE, Selker EU. RIP: the evolutionary cost of genome defense. Trends Genet. 2004
Sep;20(9):417–23.

40. Silar P, Barreau C, Debuchy R, Kicka S, Turcq B, Sainsard-Chanet A, et al. Characterization of
the genomic organization of the region bordering the centromere of chromosome V of
Podospora anserina by direct sequencing. Fungal Genet Biol. 2003 Aug 1;39(3):250–63.

41. Butler G. The Evolution of MAT: The Ascomycetes. Sex Fungi. 2007 Jan 1;3–18.

42. Martin T, Lu SW, Tilbeurgh H van, Ripoll DR, Dixelius C, Turgeon BG, et al. Tracing the origin of
the fungal α1 domain places its ancestor in the HMG-Box superfamily: Implication for fungal
mating-type evolution. PLOS ONE. 2010 Dec 8;5(12):e15199.

43. Hutchinson MI, Powell AJ, Tsang A, O'Toole N, Berka RM, Barry K, et al. Genetics of mating in
members of the Chaetomiaceae as revealed by experimental and genomic characterization of
reproduction in *Myceliophthora heterothallica*. Fungal Genet Biol. 2016 Jan 1;86:9–19.

44. Graïa F, Berteaux-Lecellier V, Zickler D, Picard M. *ami1*, an orthologue of the *Aspergillus
nidulans apsA* gene, is involved in nuclear migration events throughout the life cycle of
Podospora anserina. Genetics. 2000 Jun 1;155(2):633–46.

45. Grognet P, Timpano H, Carlier F, Aït-Benkhali J, Berteaux-Lecellier V, Debuchy R, et al. A RID-
like putative cytosine methyltransferase homologue controls sexual development in the fungus
Podospora anserina. PLOS Genet. 2019 Aug 14;15(8):e1008086.

46. Picard-Bennoun M. Does translational ambiguity increase during cell differentiation? FEBS Lett.
1982;149(2):167–70.

47. Silar P, Picard M. Increased longevity of EF-1 alpha high-fidelity mutants in *Podospora anserina*.
J Mol Biol. 1994 Jan 7;235(1):231–6.

48. Sun Y, Corcoran P, Menkis A, Whittle CA, Andersson SGE, Johannesson H. Large-scale
introgression shapes the evolution of the mating-type chromosomes of the filamentous
ascomycete *Neurospora tetrasperma*. PLOS Genet. 2012 Jul 26;8(7):e1002820.

49. Ellison CE, Stajich JE, Jacobson DJ, Natvig DO, Lapidus A, Foster B, et al. Massive changes in
genome architecture accompany the transition to self-fertility in the filamentous fungus
Neurospora tetrasperma. Genetics. 2011 Sep 1;189(1):55–69.

50. Hane JK, Rouxel T, Howlett BJ, Kema GH, Goodwin SB, Oliver RP. A novel mode of
chromosomal evolution peculiar to filamentous Ascomycete fungi. Genome Biol. 2011 May
24;12(5):R45.

51. Zickler D, Moreau PJ, Huynh AD, Slezec AM. Correlation between pairing initiation sites,
recombination nodules and meiotic recombination in *Sordaria macrospora*. Genetics. 1992 Sep
1;132(1):135–48.

52. Muyt AD, Zhang L, Piolot T, Kleckner N, Espagne E, Zickler D. E3 ligase Hei10: a multifaceted
structure-based signaling molecule with roles within and beyond meiosis. Genes Dev. 2014 May
15;28(10):1111–23.

53. Foulongne-Oriol M, Taskent O, Kües U, Sonnenberg ASM, van Peer AF, Giraud T. Mating-type
locus organization and mating-type chromosome differentiation in the bipolar edible button
mushroom *Agaricus bisporus*. Genes. 2021 Jul;12(7):1079.

54. Carpentier F, Rodríguez de la Vega RC, Jay P, Duhamel M, Shykoff JA, Perlin MH, et al. Tempo
of degeneration across independently evolved nonrecombining regions. Mol Biol Evol. 2022 Apr
1;39(4):msac060.

55. Idnurm A, Hood ME, Johannesson H, Giraud T. Contrasted patterns in mating-type
chromosomes in fungi: Hotspots versus coldspots of recombination. Fungal Biol Rev. 2015 Dec
1;29(3):220–9.

56.   Schwander T, Libbrecht R, Keller L. Supergenes and complex phenotypes. Curr Biol. 2014 Mar
31;24(7):R288–94.

57.   Berdan EL, Flatt T, Kozak GM, Lotterhos KE, Wielstra B. Genomic architecture of supergenes:
connecting form and function. Philos Trans R Soc B Biol Sci. 2022 Aug;377(1856):20210192.

58.   Maloisel L, Rossignol JL. Suppression of crossing-over by DNA methylation in *Ascobolus*.
Genes Dev. 1998 May 1;12(9):1381–9.

59.   Ben-Aroya S, Mieczkowski PA, Petes TD, Kupiec M. The compact chromatin structure of a Ty
repeated sequence suppresses recombination hotspot activity in *Saccharomyces cerevisiae*.
Mol Cell. 2004 Jul 23;15(2):221–31.

60.   Billiard S, López-Villavicencio M, Devier B, Hood ME, Fairhead C, Giraud T. Having sex, yes,
but with whom? Inferences from fungi on the evolution of anisogamy and mating types. Biol
Rev. 2011;86(2):421–42.

61.   Kubisiak TL, Milgroom MG. Markers linked to vegetative incompatibility (*vic*) genes and a region
of high heterogeneity and reduced recombination near the mating type locus (*MAT*) in
Cryphonectria parasitica. Fungal Genet Biol. 2006 Jun 1;43(6):453–63.

62.   Fontanillas E, Hood ME, Badouin H, Petit E, Barbe V, Gouzy J, et al. Degeneration of the non-
recombining regions in the mating-type chromosomes of the anther-smut fungi. Mol Biol Evol.
2015;32(4):928–43.

63.   Sun S, Hsueh YP, Heitman J. Gene conversion occurs within the mating-type locus of
Cryptococcus neoformans during sexual reproduction. PLoS Genet. 2012 Jul 5;8(7).

64.   Nakayashiki H. RNA silencing in fungi: Mechanisms and applications. FEBS Lett. 2005 Oct
31;579(26):5950–7.

65.   Meunier C, Hosseini S, Heidari N, Maryush Z, Johannesson H. Multilevel selection in the
filamentous ascomycete *Neurospora tetrasperma*. Am Nat. 2018 Mar 1;191(3):290–305.

66.   Nieuwenhuis BPS, Billiard S, Vuilleumier S, Petit E, Hood ME, Giraud T. Evolution of uni- and
bifactorial sexual compatibility systems in fungi. Heredity. 2013 Dec;111(6):445–55.

67.   Corcoran P, Anderson JL, Jacobson DJ, Sun Y, Ni P, Lascoux M, et al. Introgression maintains
the genetic integrity of the mating-type determining chromosome of the fungus *Neurospora
tetrasperma*. Genome Res. 2016 Jan 4;26(4):486–98.

68.   Ament-Velásquez SL, Vogan AA, Granger-Farbos A, Bastiaans E, Martinossi-Allibert I, Saupe
SJ, et al. Allorecognition genes drive reproductive isolation in *Podospora anserina*. Nat Ecol
Evol. 2022 Jul;6(7):910–23.

69.   Coppin E, Berteaux-Lecellier V, Bidard F, Brun S, Ruprich-Robert G, Espagne E, et al.
Systematic deletion of homeobox genes in *Podospora anserina* uncovers their roles in shaping
the fruiting body. PLOS ONE. 2012 May 25;7(5):e37488.

70.   Gould S. Wonderful life: the Burgess Shale and the nature of history. WW Norton&Company.
1990.

71.   Silar P. Podospora anserina. 2020.

72.   Andrews S. FastQC a quality control tool for high throughput sequence data. 2010; Available
from: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

73.   Leger A, Leonardi T. pycoQC, interactive quality control for Oxford Nanopore Sequencing. J
Open Source Softw. 2019 Feb 28;4(34):1236.

74. Thierry M, Charriat F, Milazzo J, Adreit H, Ravel S, Cros-Arteil S, et al. Maintenance of divergent
lineages of the rice blast fungus *Pyricularia oryzae* through niche separation, loss of sex and
post-mating genetic incompatibilities. PLOS Pathogens. 2022 in press. available at
https://www.biorxiv.org/content/10.1101/2020.06.02.129296v4.full.pdf

75. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new
genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol. 2012
Apr 16;19(5):455–77.

76. Vogan AA, Ament-Velásquez SL, Granger-Farbos A, Svedberg J, Bastiaans E, Debets AJ, et al.
Combinations of *Spok* genes create multiple meiotic drivers in *Podospora*. eLife. 2019;8.

77. Galagan JE, Calvo SE, Borkovich KA, Selker EU, Read ND, Jaffe D, et al. The genome
sequence of the filamentous fungus *Neurospora crassa*. Nature. 2003 Apr;422(6934):859–68.

78. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: scalable and
accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res.
2017;27(5):722–36.

79. Vaser R, Sović I, Nagarajan N, Šikić M. Fast and accurate de novo genome assembly from long
uncorrected reads. Genome Res. 2017 May;27(5):737–46.

80. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: An integrated
tool for comprehensive microbial variant detection and genome assembly improvement. PLOS
ONE. 2014 Nov 19;9(11):e112963.

81. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinforma Oxf Engl. 2009 Jul 15;25(14):1754–60.

82. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014 Apr 1;30:2114–20.

83. Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome
assemblies. Bioinformatics. 2013 Apr 15;29(8):1072–5.

84. Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO update: Novel and
streamlined workflows along with broader and deeper phylogenetic coverage for scoring of
eukaryotic, prokaryotic, and viral genomes. Mol Biol Evol. 2021 Oct 1;38(10):4647–54.

85. Manni M, Berkeley MR, Seppey M, Zdobnov EM. BUSCO: Assessing Genomic Data Quality and
Beyond. Curr Protoc. 2021;1(12):e323.

86. Smit AFA, Hubley R, Green P. RepeatMasker Open-4.0. Available at:
http://www.repeatmasker.org Accessed: November 30, 2017. 2013 2015;

87. Palmer JM, Stajich J. Funannotate v1.8.1: Eukaryotic genome annotation. Zenodo; 2020.
Available from: https://zenodo.org/record/4054262

88. Brůna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M. BRAKER2: automatic eukaryotic
genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein database.
NAR Genomics Bioinforma. 2021;3(1):lqaa108.

89. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal
RNA-seq aligner. Bioinformatics. 2013;29(1):15–21.

90. Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, et al. InterProScan 5: genome-scale
protein function classification. Bioinformatics. 2014 Jan 5;30(9):1236–40.

91.   Krogh A, Larsson B, von Heijne G, Sonnhammer ELL. Predicting transmembrane protein
1062   topology with a hidden markov model: application to complete genomes. J Mol Biol. 2001 Jan
1063   19;305(3):567–80.

1064   92.   Petersen TN, Brunak S, von Heijne G, Nielsen H. SignalP 4.0: discriminating signal peptides
1065   from transmembrane regions. Nat Methods. 2011 Oct;8(10):785–6.

1066   93.   Käll L, Krogh A, Sonnhammer ELL. Advantages of combined transmembrane topology and
1067   signal peptide prediction—the Phobius web server. Nucleic Acids Res. 2007 Jul;35(Web Server
1068   issue):W429-432.

1069   94.   Smit AFA, Hubley R. RepeatModeler Open-1.0.Available at: http://www.repeatmasker.org.
1070   Accessed: November 30, 2017. 2008 2015;

1071   95.   Vogan AA, Ament-Velásquez SL, Bastiaans E, Wallerman O, Saupe SJ, Suh A, et al. The
1072   *Enterprise*, a massive transposon carrying *Spok* meiotic drive genes. Genome Res. 2021
1073   May;31(5):789–98.

1074   96.   Gioti A, Mushegian AA, Strandberg R, Stajich JE, Johannesson H. Unidirectional evolutionary
1075   transitions in fungal mating systems and the role of transposable elements. Mol Biol Evol. 2012
1076   Oct 1;29(10):3215–26.

1077   97.   Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, et al. A unified classification
1078   system for eukaryotic transposable elements. Nat Rev Genet. 2007 Dec;8(12):973–82.

1079   98.   Wyk S van, Harrison CH, Wingfield BD, Vos LD, Merwe NA van der, Steenkamp ET. The
1080   RIPper, a web-based tool for genome-wide quantification of Repeat-Induced Point (RIP)
1081   mutations. PeerJ. 2019 Aug 26;7:e7447.

1082   99.   Debuchy R, Berteaux-Lecellier V, Silar P. Mating systems and sexual morphogenesis in
1083   Ascomycetes. In: Cellular and Molecular Biology of Filamentous Fungi [Internet]. John Wiley &
1084   Sons, Ltd; 2010. p. 499–535. Available from:
1085   https://onlinelibrary.wiley.com/doi/abs/10.1128/9781555816636.ch33

1086   100. Carver T, Harris SR, Berriman M, Parkhill J, McQuillan JA. Artemis: an integrated platform for
1087   visualization and analysis of high-throughput sequence-based experimental data.
1088   Bioinformatics. 2012 Feb 15;28(4):464–9.

1089   101. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of
1090   high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol. 2011 Oct
1091   11;7:539.

1092   102. Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. Jalview Version 2--a multiple
1093   sequence alignment editor and analysis workbench. Bioinforma Oxf Engl. 2009 May
1094   1;25(9):1189–91.

1095   103. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative genomics.
1096   Genome Biol. 2019 Nov 14;20(1):238.

1097   104. Abascal F, Zardoya R, Telford MJ. TranslatorX: multiple alignment of nucleotide sequences
1098   guided by amino acid translations. Nucleic Acids Res. 2010 Jul;38(Web Server issue):W7-13.

1099   105. Yang Z. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol. 2007 Aug
1100   1;24(8):1586–91.

1101   106. Yang Z, Nielsen R. Estimating synonymous and nonsynonymous substitution rates under
1102   realistic evolutionary models. Mol Biol Evol. 2000 Jan;17(1):32–43.

107. Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. MUMmer4: A fast and
versatile genome alignment system. PLOS Comput Biol. 2018 Jan 26;14(1):e1005944.

108. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018 Sep
15;34(18):3094–100.

109. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009 Aug 15;25(16):2078–9.

110. Liu KH, Yeh YL, Shen WC. Fast preparation of fungal DNA for PCR screening. J Microbiol
Methods. 2011 May 1;85(2):170–2.

111. Rozen S, Skaletsky H. Primer3 on the WWW for general users and for biologist programmers.
Methods Mol Biol Clifton NJ. 2000;132:365–86.

112. Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et al. Primer3--new
capabilities and interfaces. Nucleic Acids Res. 2012 Aug;40(15):e115.

1115

1116

1117

Supporting information captions

1119

Supplementary note S1: Frequency of second-division segregation of the mating-type locus under random segregation of the mating-type locus.

Supplementary Table S1: Tests of differences in growth speed and dry weight between *Schizothecium tetrasporum MAT1-1 and MAT1-2* homokaryons and heterokaryons on different media.

Supplementary Table S2: Genome assembly of the *Schizothecium tetrasporum* monokaryotic strains CBS815.71-sp3 and CBS815.71-sp6.

Supplementary Table S3: List of 15 genes in the recombination suppression region around the mating-type locus in *Schizothecium tetraporum* predicted to encode proteins that are putative transcription factors or involved in signalling pathways.

Supplementary Table S4: List of genes in the recombination suppression region around the mating-type locus in *Schizothecium tetraporum* having orthologues also located in the recombination suppression region around the mating-type locus in *Podospora anserina* and *Neurospora tetrasperma*.

Supplementary Table S5: Quality of the raw reads produced and used in this study.

Supplementary Table S6: PCR primers used in this study.

Supplementary Table S7: PCR conditions used in this study.

Supplementary Figure S1: Genome collinearity between the long read-based assembly and the Illumina-based assembly of *Schizothecium tetrasporum* CBS815.71-sp3 strain.

Supplementary Figure S2: Transposable element annotation in different genomic regions of the *Schizothecium tetrasporum* CBS815.71 strain.

Supplementary Figure S3: Amino-acid alignment of proteins containing a MATα_HMG domain.

Supplementary Figure S4: Genome-wide distribution of synonymous divergence (dS) in the *Schizothecium tetrasporum* CBS815.71 strain. dS was computed per-gene between the CBS815.71-sp3 and CBS815.71-sp6 assemblies and plotted according to the CBS815.71-sp3 assembly gene position.

Supplementary Figure S5: GC content variation and determination of centromeres the *Schizothecium tetrasporum* CBS815.71-sp3 assembly.

Supplementary Figure S6: Delimitation of three evolutionary strata in the S*chizothecium tetrasporum* CBS815.71 strain.

Supplementary Figure S7: Probability of recombination in centimorgans (cM) between markers and the mating-type locus in the heterokaryotic strain CBS815.71 after one round of experimental selfing measured in 107 of the offspring.

Supplementary figure S8: Genome-wide collinearity based on single orthologous genes between *Neurospora crassa*, *Podospora anserina* and *Schizothecium tetrasporum* genomes.

1156

1157

1160   **FIGURES**

1161

Figure 1: Sexual reproductive structures and ascospores of *Schizothecium tetrasporum*

CBS815.71. A. Perithecia on *Miscanthus* (1) and squashed perithecia showing the rosette of asci (2);

bars = 200 µm. **B.** Enlargements of asci within the rosettes (1) and a five-spore ascus with three

dikaryotic ascospores and two monokaryotic ascospores (2); bars= 20 µm. **C.** Other abnormal asci

observed in CBS815.71: (1) a six-spore ascus with four monokaryotic ascospores (small) and two

dikaryotic ones (large); (2) a five-spore ascus with three monokaryotic ascospores (the three small

spores in the middle), one dikaryotic (the large ascospore at the bottom) and another, probably

trikaryotic, ascospore (arrowhead showing the very large ascospore); bars= 20 µm.

1170

1171

1172

1173

1174

Figure 2: Segregation of the mating-type locus during meiosis, ascus composition and

hypothetical scenarios for the loss of one mating type in thalli of *Schizothecium tetrasporum*

A. Theoretical composition of an ascus according to meiotic segregation pattern at the mating-type

locus: if alleles at the mating-type locus (represented as a red or blue vertical bar) segregate during

the first meiotic division (FDS, top), two *MAT-1-1* spores and two *MAT-1-2* spores are formed;

conversely, if alleles at the mating-type locus segregate during the second meiotic division (SDS,

bottom), four heterokaryotic spores carrying both mating types are formed. Only the chromosomes

carrying the mating-type locus are represented, with their centromeres. The *MAT-1-1* and *MAT1-2*

alleles at the mating-type locus are represented by red and blue vertical bars, respectively.

Centromeres are represented by yellow circles at the diploid and meiosis I stages. Chromosomal

arms are colored in red or blue according to the parental origin of the chromosomal fragment. **B.** Two

hypothetical scenarios for the loss of one mating type despite SDS at the mating-type locus. SDS at

the mating-type locus generates a spore carrying two nuclei of opposite mating types, but one nucleus

may be lost during spore development (top) when half the nuclei gather in a region of the spore that

subsequently undergoes apoptosis, or during mycelium growth (bottom).

1190

1191   **A**

1192

1193

1194   **B**

1195

1196

Figure 3: Genome features and annotation of the mating-type locus in the *Schizothecium*
***tetrasporum* CBS815.71 strain. A.** Ideograms representing the seven largest contigs of the
CBS815.71-sp3 genome assembly. The mating-type locus is indicated with a green triangle, telomeric
repeats with red boxes and putative centromeres with yellow circles. Gene density in 10 kb non-
overlapping windows is represented by vertical bars on the contigs and the heatmap colors indicate
low to high density. The non-recombining region around the mating-type locus is indicated by a black
rectangle. **B.** Genome collinearity between opposite mating types of the *Schizothecium tetrasporum*
CBS815.71 strain. Circular plot showing contigs from the CBS815.71-sp3 genome (green, right) and
the CBS815.71-sp6 genome (purple, left). Only the largest contigs (>500 kb) were plotted. Blue links
indicate the locations of orthologous genes (Mb). The mating-type locus is indicated with a green
triangle. Putative centromeres and the non-recombining region around the mating-type locus identified
on the CBS815.71-sp3 genome assembly are indicated by yellow circles and a black rectangle,
respectively. **C.** Organization of the mating-type locus in CBS815.71-sp3 (top) and CBS815.71-sp6
(bottom) assemblies viewed in Artemis v18.0.2 (100). The *APN2* and *SLA2* genes flanking the
mating-type locus are indicated. The green boxes represent the nucleoide sequences differing
between the two idiomorphs that were used to design markers for assessing mating type in the
progeny. The *MAT1-1-1* pseudogene (*TαD1*) present in the CBS815.71-sp3 (*MAT1-2*) genome is
boxed in red. The three forward and three reverse frames of the DNA sequence are presented;
vertical bars indicate putative STOP codons.

1216

1217

1218

Figure 4: Synonymous divergence (d$_S$) along the mating-type chromosome in the

Schizothecium tetrasporum **CBS815.71 strain.** d$_S$ was calculated per gene, between the orthologs

common to the CBS815.71-sp3 and CBS815.71-sp6 assemblies and was plotted, according to the

CBS815.71-sp3 assembly gene position (contig 1), along the *x*-axis. Evolutionary strata 1, 2, and 3

are indicated by red, green and blue rectangles, respectively. The mating-type locus is indicated with

a black dotted vertical line and a green triangle. The putative centromere is indicated with a yellow

circle. The *P_NRR2* and *P_NRR5* markers, which displayed no recombination event with the mating-

type locus, are indicated with gray diamonds.

1227

1228

1229

 Figure 5: Collinearity and shared gene content between the mating-type chromosomes of
Schizothecium tetrasporum, Podospora anserina and Neurospora tetrasperma. A. Collinearity
between (i) *N. crassa* (left) and *P. anserina* (right); (ii) *P. anserina* (left) and *S. tetrasporum* (right);
(iii) *N. crassa* (left) and *S. tetrasporum* (right). Collinearity is based on single-copy orthologous genes.
Centromere is indicated with a yellow circle and the region of well-conserved gene order around the
centromere is highlighted with a yellow rectangle. The mating-type locus location is shown with a
green triangle and the non-recombining region around the mating-type locus in *P. anserina* (23) and
S. tetrasporum with a black track. Along the recombining mating-type chromosome of *N. crassa* a
black rectangle indicates the orthologous region that is non recombining in multiple *N. tetrasperma*
lineages (26). Red, green and blue links highlight orthologous genes located in the evolutionary strata
1, 2, and 3 of *S. tetrasporum,* respectively. Grey links indicate other orthologous genes. On panel (ii),
the evolutionary strata identified previously in the *P. pseudocomata* CBS415.72 strain, that is closely
related to the *P. anserina* S strain, are indicated by purple (for the oldest stratum) and dark green (for
the youngest stratum) tracks (27), at left. B. Number of single-copy orthologous genes present in the
non-recombining region of *S. tetrasporum* that are also in the non-recombining regions of *P. anserina*
and/or *N. tetrasperma*, as indicated by the squares at the bottom (a filled square indicates the
presence in the non-recombining region of the focal species).

1248

A

1249
1250

B

1251