Male Guinea baboon tracking of female whereabouts

Dominique Treschnak[a], Dietmar Zinner[a, b, c] Julia Fischer[a, b, c]

[a] Cognitive Ethology Laboratory, German Primate Center, Göttingen, Germany

[b] Department of Primate Cognition, Georg-August-University of Göttingen, Göttingen, Germany

[c] Leibniz ScienceCampus Primate Cognition, Göttingen, Germany

Orcid: Treschnak: https://orcid.org/0000-0002-8249-191X

Zinner: http://orcid.org/0000-0003-3967-8014

Fischer: http://orcid.org/0000-0002-5807-0074

Abstract

In group-living species, evolution puts a premium on the ability of individuals to track the state, whereabouts, and interactions of others. The value of social information might vary with the degree of competition within and between groups, however. We investigated male monitoring of female location in wild Guinea baboons (*Papio papio*). Guinea baboons live in socially tolerant multi-level societies with one-male-units comprising 1-6 females and young at the core. Using field playback experiments, we tested whether males (N=22 males, N=62 trials) keep track of the whereabouts of associated females by playing back unit females' calls from locations that were either consistent or inconsistent with the actual position of the female. Contrary to predictions, males responded equally strongly in both conditions. In a preparatory experiment, males (N=14) responded more strongly to playbacks of unit vs. non-unit females. While males seem to recognize their females by voice, they were not able or not motivated to track their females' movements. These results reinforce the view that the value of social information may vary substantially with the distribution of power in a society. While highly competitive regimes necessitate high attention to deviations from expected patterns, egalitarian societies allow for a certain degree of obliviousness.

Introduction

Knowledge about conspecifics and their relationships guides social decision-making in many group-living animals. The use of such social knowledge is documented for a large number of species, ranging from simple and more complex forms of individual recognition [1] to the assessment and monitoring of stable or transient social attributes of group members, like kinship, rank, or bond strengths. Such knowledge extends not only to an individual's direct associations but also to third-party relationships [2]. When navigating the social environment, knowledge about previous interactions with group members, the capabilities of potential partners or competitors, and the nature and quality of relationships between others, aids in

predicting the outcomes of future interactions and allows to act strategically. For example,

spotted hyenas (*Crocuta crocuta*) joining into dyadic fights mainly support the dominant

individual and are subsequently also more likely to attack relatives of the subordinate [3].

Pinyon jays (*Gymnorhinus cyanocephalus*) assess their relative rank difference to strangers

by observing them in encounters with known individuals [4]. Tonkean macaques (*Macaca*

tonkeana) respond more strongly to conflicts between strongly bonded individuals ('friends')

compared to non-friends [5].

Besides kin and allies, mating partners are of particular value to an individual. Males

compete not only for access to females [6,7]; they are also under selection to monitor the state

and behaviour of females. Males may increase their reproductive success by assessing suited

mating partners [8] or mating opportunities [9,10]. In many species, females become the centre

of male attention when they approach the fertile phase of their reproductive cycle. In contrast,

in species where males and females form long-lasting bonds as in monogamous [11] or

polygynandrous species (e.g., plains zebras (*Equus burchellii*) [12], hamadryas baboons

(*Papio hamadryas*) [13]), males are permanently incentivised to monitor and control associated

females' whereabouts and interactions with other group members.

We tested male knowledge of female whereabouts in wild Guinea baboons (*Papio*

papio). The species lives in multi-level societies. At the core are one-male units consisting of

one primary male, one to six associated females, and their offspring. Bachelor males may be

associated with several such units [14]. Several units form a party, which in turn aggregate into

gangs [15]. Females associate with one primary male and show mate fidelity [16], but in

contrast to hamadryas baboons, they also enjoy spatial freedom, i.e., they may spend

considerable time away from their male [16].

We hypothesized that males keep track of the movement patterns of associated

females. To test this hypothesis, we conducted a playback experiment [17], in which we

presented female grunts from a location that was either consistent or inconsistent with the

actual position of the female. We made use of the violation-of-expectation paradigm and

presented the animals with a physically impossible scenario, similar to Townsend et al. (2012).

We tested a male immediately after the female had left him and assumed that he would have

noticed the direction in which she disappeared. We predicted that males would show 'signs of

surprise', meaning a stronger response, when they were confronted with information that the

female was in an unexpected - indeed physically impossible – location compared to their

response when the female's vocalisation came from the direction into which she had recently

disappeared. In a preparatory experiment, we tested the prerequisite that males can recognise

their associated females by voice. We tested if males respond more strongly to the

vocalizations of females from their unit compared to the vocalizations from females of another

unit, but the same party. We predicted that males would show stronger responses when

presented with vocalisation from unit-females.

Methods

The experiments took place between January 2019 and August 2021 at the Centre de

Recherche de Primatologie Simenti in the Niokolo-Koba National Park in Senegal, a field

station maintained by the German Primate Center (see Fischer et al., 2017 for details). The

study population comprised ~ 200 individually identified Guinea baboons that belonged to three

parties, with a varying number of reproductive units (between 15 and 25 per year across three

parties) suitable for the experiments. For the experimental stimuli, we recorded 'grunt'

vocalisations of sub-adult and adult females during their non-receptive phase (electronic

supplementary material, appendix S1, S2). Grunts are the most frequently occurring

vocalisation in Guinea baboons and are mainly produced in affiliative contexts [20].

In Experiment 1 (individual recognition), we presented males with calls from a female

from their unit (*unit-female* condition) and a female from another unit (*non-unit-female*

condition). Trials were separated by at least five days and conducted only when females were

non-receptive. Once the female whose call was to be played back was not visible to the

subject, a loudspeaker was positioned at a 90° angle to the left or right of the male depending

on the actual position of the female, and the stimulus presented. Male responses were video

recorded for three minutes after the onset of the stimulus. We conducted 28 playback trials

testing 14 primary males.

In Experiment 2 (spatial monitoring), we tested males in a within-subject design and

presented grunts from a unit-female on two occasions separated by at least seven days. As

above, trials were conducted only when females were non-receptive. In the *consistent*

condition, the speaker was hidden in a location matching the actual direction of the departed

female, whereas in the *inconsistent* condition in the opposite direction, presenting an

impossible scenario (figure 1). A male was tested after he had been near a unit-female, she

had then walked away and was no longer in sight (median time out-of-sight: 70s, range 8 s –

273 s). A loudspeaker was then hidden in vegetation, at a 90° angle to the left or right of the

male and a distance of approximately 10 m. Male responses were video recorded for 10 min.

after the onset of the stimulus. We conducted 62 playback trials with 22 primary males. Nine

of these males were tested twice with the call of a different female (average time between first

and second run: 43 weeks (min: 3, max: 100)) (electronic supplementary material, appendix

S3).

Video recordings were coded using Solomon coder beta (András Péter,

solomoncoder.com) on a frame-by-frame basis (25 frames/s). We examined male responses

by coding changes in their head orientation; i.e., changes between the neutral position: male

faces camera, and subsequent looks exceeding an angle of 45° towards the direction of the

speaker or away from it. We measured the duration of the first look and the latency to respond.

Trials where the latency exceeded the cut-off were coded as "no response". As the first look in

the inconsistent condition could be truncated because the male may turn his attention to look

into the direction where the female was last seen, we additionally measured the total time

vigilant (all looks toward the speaker or actual position of the female) within 30 s after stimulus

onset in the social monitoring experiment, (electronic supplementary material, appendix S4

(observer reliability); appendix S5, figure S1, S2 (classification of responses)).

Analyses were carried out in R (version 4.1.1; R Core Team, 2021). GLMMs, LMMs,

and Cox proportional hazard model were fitted using the R package *lme4*, *survival* and *coxme*

(electronic supplementary material, appendix S6, S7). We used a Linear Mixed model [22] for

first look duration (experiment 1) and vigilance time, a Generalized Linear Mixed Model with

gamma error structure and log link function [22] for the duration of the first look (experiment

2), and a survival analysis [23] for latencies. In addition to the main predictor 'experimental

condition', we included unit size as a fixed effect to control for the influence of the number of

unit-females and male identity as random intercept. As in experiment 1, the same call could

be used in the unit-female and non-unit-female conditions for different males, and some calls

(stimuli) stemmed from the same female, we included female ID and stimulus ID as additional

random intercept effects. To investigate the effect of the main predictor we compared the full

model to a null model lacking experimental condition using a likelihood ratio test (Dobson &

Barnett, 2008). Confidence intervals of estimates and fitted values were determined using a

parametric (LMM & GLMM) and non-parametric (survival analysis) bootstrap (N=1000

bootstraps).

Figure 1. Set-up experiment 2 (spatial monitoring). In the consistent condition, a speaker is positioned close to the actual position of the female, in the inconsistent condition, the speaker is placed in the opposite direction.

Results

In experiment 1 (individual recognition), males responded to the playback of calls in 24 out of 28 trials. The average duration of the first response was 3.18 s ± 2.53 s (median ± IQR). Males looked longer when presented with calls from unit-females (3.38 s ± 4.60 s) compared to non-unit females (2.32 s ± 3.80 s) (full-null model comparison: χ^2_1=8110, p=0.004, table S1a). The average latency of responses was 0.96 s ± 0.86 s for the unit-female and 1.26 s ± 3.21 s for

non-unit-females (median ± IQR). Unit size had no obvious effect on response duration or

latency (Duration: p=0.48; Latency: p=0.37, table S1a, S2).

In experiment 2 (spatial monitoring), males responded to the playback in 49 out of 62

trials (consistent condition: N=22, inconsistent: N=27). There was no obvious difference in the

duration of first look in the consistent (2.80 s ± 4.36 s, median ± IQR) compared to the

inconsistent (2.92 s ± 2.62 s) condition (full-null model comparison: χ^2_1=0.0002, p=0.99, table

S3) (figure 2b). There were no obvious differences in response latencies between the two

conditions (consistent: 0.72 s ± 0.48 s; inconsistent: 0.80 ± 0.64 (median ± IQR); full-null

model comparison: χ^2_1 =1.10, p=0.29, table S4). There were also no obvious differences in the

overall time vigilant (consistent: 7.84 s ± 7.19 s (median ± IQR); inconsistent: 8.08 s ± 6.39 s;

full-null model comparison: χ^2_1 =0.04, p =0.84, table S5). We found no evidence that unit size

influenced any of the response variables (Duration: p=0.38; Latency: p=0.63, Vigilance:

p=0.15, table S3, S4, S5).

Figure 2. First look duration for males in the a) individual recognition experiment and b) spatial

monitoring experiment. Connected points represent data from the same individual (a: N=14; b:

N=22). Thick black lines depict bootstrapped mean and 95% confidence intervals for males

with average unit size.

Discussion

Male Guinea baboons showed no signs of surprise when calls from associated females were

played back from an impossible location. Instead, they responded equally strongly to

playbacks of calls from an impossible or a possible location. Further, males responded more

strongly to the playback of vocalizations from unit-females compared to non-unit-females.

While males seemed to be able to recognise their unit's females by voice, they lacked either

the ability or the motivation to track their females' positions.

These findings were not in line with our initial prediction that primary males monitor the

whereabouts of their females. Guinea baboons form one-male units similar to hamadryas

baboons or mountain gorillas (*Gorilla b. beringei*). In both of these species, sexual coercion

[25] is used by leader males to control female movement and interactions and to prevent

transfers to other males [26,27]. In Guinea baboons, we did not observe such overt aggression

towards females, except for some occasional chasing of females. Indeed, female Guinea

baboons can roam relatively unimpeded and interact socially with other group members,

including other adult males [16].

The lack of differentiated response fits with the relatively laid-back stance of Guinea

baboon males. Males form strong bonds with other males [14,15,28]. They also show low

levels of overt aggression, preventing us from discerning a clear dominance hierarchy [14]. At

the same time, female Guinea baboons have considerable leverage in mate choice and

intersexual bond maintenance [16]. Male strategies mainly seem to consist of investing their

social time into female grooming and support. Interestingly, males appear to face a trade-off

in the allocation of social time, as male investment into socio-positive interactions with other

male declines with increasing unit size [28]. Social investment into females thus might be

important for intersexual bond maintenance and potentially female mate choice in the first

place.

Since we tested males when the female whose calls were played was not receptive,

we do not know whether males would be more attentive if the female would be able to conceive.

We conducted the trials only while females were non-receptive because, during females'

oestrus, primary males and females are less likely to separate [29], leaving very few

opportunities for conducting the experimental trials. Thus, we cannot exclude the possibility

that males would respond differently in conditions where they should be more motivated to

track their female's whereabouts.

Our study adds to the accumulating evidence that the need to monitor the social

environment varies between species with the degree of competition among individuals. For

instance, the highly competitive chacma baboons (*Papio ursinus*), which live in female

philopatric groups show strong responses to the playback of vocalisation from unfamiliar males

[30,31], while Guinea baboons showed greater attention to vocalisations from familiar males

compared to neighbours or strangers [20]. In geladas (*Theropithecus gelada*), which live in a

multi-level society in aggregations of up to several hundred individuals, vocal recognition

seems to be limited to individuals with a high degree of social overlap [32]. Additionally, when

presenting individuals with information about changes in association patterns, chacma

baboons responded strongly to simulated separations of consortships [10], while Guinea

baboons paid more attention to information consistent with current male-female association

patterns [33]. Similarly, Geladas did not differentiate between consistent or inconsistent

information about male-female relationships at all [34].

While the link between group-living and sophisticated social knowledge is well

documented [2], it is still unclear whether life in a socially complex environment per se [35] or

rather the degree of competition within and between groups selects for advanced socio-

cognitive skills ("Machiavellian intelligence") [36]. Bergman (2010, p. 2050) argued that

"missing social knowledge" might be a consequence of the absence of a competitive

environment that offers no benefits for the ability to assess and use of specific social

information of conspecifics. Our results as well as results of previous from the same population

[20,33] suggest that a reduced competitive environment affects the value of social information,

and as a consequence, the motivation or ability of an individual to attend to them. At the same

time, both Guinea baboons and geladas live in highly structured multi-level groups, suggesting

that a complex social organisation does not per se select for a high motivation to monitor the

social environment. We contend that a skewed distribution of power influences the value of

social information and therefore the motivation to attend to events in the social environment.

Acknowledgements

We thank the Direction des Parcs Nationaux and Ministère de l'Environnement et de la Protection de la Nature du Sénégal for permission to work in the Niokolo-Koba National Park. We are grateful to all research assistants for their help and thank Roger Mundry for valuable statistical advice and Robert Seyfarth for helpful comments on the manuscript.

Funding

This project was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) project number 254142454.

References

1. Wiley RH. 2013 Specificity and multiplicity in the recognition of individuals: implications for the evolution of social behaviour. *Biol. Rev.* **88**, 179–195. (doi:https://doi.org/10.1111/j.1469-185X.2012.00246.x)

2. Seyfarth RM, Cheney DL. 2015 Social cognition. *Anim. Behav.* **103**, 191–202. (doi:10.1016/j.anbehav.2015.01.030)

3. Engh AL, Siebert ER, Greenberg DA, Holekamp KE. 2005 Patterns of alliance formation and postconflict aggression indicate spotted hyaenas recognize third-party relationships. *Anim. Behav.* **69**, 209–217. (doi:https://doi.org/10.1016/j.anbehav.2004.04.013)

4. Paz-y-Miño C G, Bond AB, Kamil AC, Balda RP. 2004 Pinyon jays use transitive inference to predict social dominance. *Nature* **430**, 778–781. (doi:10.1038/nature02723)

5. Whitehouse J, Meunier H. 2020 An understanding of third-party friendships in a tolerant macaque. *Sci. Rep.* **10**, 9777. (doi:10.1038/s41598-020-66407-w)

6. Clutton-Brock TH, Parker GA. 1992 Potential Reproductive Rates and the Operation of Sexual Selection. *Q. Rev. Biol.* **67**, 437–456. (doi:10.1086/417793)

7. Clutton-Brock TH, Vincent ACJ. 1991 Sexual selection and the potential reproductive rates of males and females. *Nature* **351**, 58–60. (doi:10.1038/351058a0)

8. Davies AD, Lewis Z, Dougherty LR. 2020 A meta-analysis of factors influencing the strength of mate-choice copying in animals. *Behav. Ecol.* **31**, 1279–1290. (doi:10.1093/beheco/araa064)

9. Balsby TJ, Dabelsteen T. 2005 Simulated courtship interactions elicit neighbour intrusions in the whitethroat, Sylvia communis. *Anim. Behav.* **69**, 161–168. (doi:https://doi.org/10.1016/j.anbehav.2004.01.021)

10. Crockford C, Wittig RM, Seyfarth RM, Cheney DL. 2007 Baboons eavesdrop to deduce
mating opportunities. *Anim. Behav.* **73**, 885–890.
(doi:https://doi.org/10.1016/j.anbehav.2006.10.016)

11. Birkhead TR, Møller AP. 1995 Extra-pair copulation and extra-pair paternity in birds.
Anim. Behav. **49**, 843–848.

12. Rubenstein DI, Hack M. 2004 Natural and sexual selection and the evolution of multi-
level societies: insights from zebras with comparisons to primates. In *Sexual Selection in*
Primates (eds PM Kappeler, CP van Schaik), pp. 266–279. Cambridge University Press.
(doi:10.1017/CBO9780511542459.017)

13. Swedell L, Plummer T. 2012 A papionin multilevel society as a model for hominin social
evolution. *Int. J. Primatol.* **33**, 1165–1193. (doi:https://doi.org/10.1007/s10764-012-9600-
9)

14. Dal Pesco F, Trede F, Zinner D, Fischer J. 2021 Kin bias and male pair-bond status
shape male-male relationships in a multilevel primate society. *Behav. Ecol. Sociobiol.* **75**,
1–14. (doi:https://doi.org/10.1007/s00265-020-02960-8)

15. Patzelt A, Kopp GH, Ndao I, Kalbitzer U, Zinner D, Fischer J. 2014 Male tolerance and
male–male bonds in a multilevel primate society. *Proc. Natl. Acad. Sci.* **111**, 14740–
14745. (doi:https://doi.org/10.1073/pnas.1405811111)

16. Goffe AS, Zinner D, Fischer J. 2016 Sex and friendship in a multilevel society:
behavioural patterns and associations between female and male Guinea baboons.
Behav. Ecol. Sociobiol. **70**, 323–336. (doi:https://doi.org/10.1007/s00265-015-2050-6)

17. Fischer J, Noser R, Hammerschmidt K. 2013 Bioacoustic Field Research: A Primer to
Acoustic Analyses and Playback Experiments With Primates: Bioacoustic Field Methods.
Am. J. Primatol. **75**, 643–663. (doi:10.1002/ajp.22153)

18. Townsend SW, Allen C, Manser MB. 2012 A simple test of vocal individual recognition in
wild meerkats. *Biol. Lett.* **8**, 179–182. (doi:https://doi.org/10.1098/rsbl.2011.0844)

19. Fischer J *et al.* 2017 Charting the neglected West: The social system of Guinea baboons.
Am. J. Phys. Anthropol. **162**, 15–31. (doi:10.1002/ajpa.23144)

20. Maciej P, Patzelt A, Ndao I, Hammerschmidt K, Fischer J. 2013 Social monitoring in a
multilevel society: a playback study with male Guinea baboons. *Behav. Ecol. Sociobiol.*
67, 61–68. (doi:https://doi.org/10.1007/s00265-012-1425-1)

21. R Core Team. 2021 *R: A Language and Environment for Statistical Computing*. Vienna,
Austria: R   Foundation for Statistical Computing. See URL https://www.R-project.org/.

22. Baayen RH, Davidson DJ, Bates DM. 2008 Mixed-effects modeling with crossed random
effects for subjects and items. *J. Mem. Lang.* **59**, 390–412.
(doi:https://doi.org/10.1016/j.jml.2007.12.005)

23. Jahn-Eimermacher A, Lasarzik I, Raber J. 2011 Statistical analysis of latency outcomes
in behavioral experiments. *Behav. Brain Res.* **221**, 271–275.
(doi:https://doi.org/10.1016/j.bbr.2011.03.007)

24. Dobson AJ, Barnett AG. 2008 *An introduction to generalized linear models*. Chapman
and Hall/CRC.

25. Smuts BB, Smuts RW. 1993 Male aggression and sexual coercion of females in
nonhuman primates and other mammals: evidence and theoretical implications. In
Advances in the Study of Behavior, pp. 1–63.

26. Schreier AL, Swedell L. 2009 The fourth level of social structure in a multi-level society:
ecological and social functions of clans in hamadryas baboons. *Am. J. Primatol.* **71**, 948–
955. (doi:10.1002/ajp.20736)

27. Sicotte P. 1993 Inter-group encounters and female transfer in mountain gorillas:
Influence of group composition on male behavior. *Am. J. Primatol.* **30**, 21–36.
(doi:10.1002/ajp.1350300103)

28. Dal Pesco F, Trede F, Zinner D, Fischer J. 2022 Male–male social bonding, coalitionary
support and reproductive success in wild Guinea baboons. *Proc. R. Soc. B* **289**,
20220347. (doi:https://doi.org/10.1098/rspb.2022.0347)

29. Goffe AS. 2016 Social relationships of female Guinea baboons (Papio papio) in Senegal.
Dissertation, Göttingen, Georg-August Universität, 2016.

30. Kitchen DM, Cheney DL, Seyfarth RM. 2005 Male chacma baboons (Papio hamadryas
ursinus) discriminate loud call contests between rivals of different relative ranks. *Anim.
Cogn.* **8**, 1–6. (doi:https://doi.org/10.1007/s10071-004-0222-2)

31. Kitchen DM, Cheney DL, Engh AL, Fischer J, Moscovice LR, Seyfarth RM. 2013 Male
baboon responses to experimental manipulations of loud "wahoo calls": testing an honest
signal of fighting ability. *Behav. Ecol. Sociobiol.* **67**, 1825–1835.
(doi:https://doi.org/10.1007/s00265-013-1592-8)

32. Bergman TJ. 2010 Experimental evidence for limited vocal recognition in a wild primate:
implications for the social complexity hypothesis. *Proc. R. Soc. B Biol. Sci.* **277**, 3045–
3053. (doi:10.1098/rspb.2010.0580)

33. Faraut L, Fischer J. 2019 How life in a tolerant society affects the attention to social
information in baboons. *Anim. Behav.* **152**, 11–17.
(doi:https://doi.org/10.1016/j.anbehav.2019.04.004)

34. le Roux A, Bergman TJ. 2012 Indirect rival assessment in a social primate,
Theropithecus gelada. *Anim. Behav.* **83**, 249–255.
(doi:https://doi.org/10.1016/j.anbehav.2011.10.034)

35. Holekamp KE. 2007 Questioning the social intelligence hypothesis. *Trends Cogn. Sci.*
11, 65–69. (doi:10.1016/j.tics.2006.11.003)

36. Whiten A, Byrne RW. 1988 The Machiavellian intelligence hypotheses. Clarendon
Press/Oxford University Press.

335