Predictors of genomic diversity within North American squamates

Ivy E. Larkin[1] (larkin.132@buckeyemail.osu.edu)

Edward A. Myers[2] (eddie.a.myers@gmail.com; https://orcid.org/0000-0001-5369-6329)

Bryan C. Carstens[1] (carstens.12@osu.edu; https://orcid.org/0000-0002-1552-227X)

Lisa N. Barrow[1,3*] (lnbarrow@unm.edu; https://orcid.org/0000-0001-7081-2432)

[1]Department of Evolution, Ecology and Organismal Biology, The Ohio State University, 318 W.

12th Ave, Columbus, OH 43210

[2]Department of Biological Sciences, Clemson University, Clemson, SC 29634

[3]Current Address: Museum of Southwestern Biology and Department of Biology, University of

New Mexico, Albuquerque NM 87131

*Corresponding author: lnbarrow@unm.edu

Running Title: GENOMIC DIVERSITY IN SNAKES

Abstract

Comparisons of intraspecific genetic diversity across species can reveal the roles of geography, ecology, and life history in shaping biodiversity. The wide availability of mitochondrial DNA (mtDNA) sequences in open-access databases makes this marker practical for conducting analyses across several species in a common framework, but patterns may not be representative of overall species diversity. Here, we gather new and existing mtDNA sequences and genome-wide nuclear data (genotyping-by-sequencing; GBS) for 30 North American squamate species sampled in the Southeastern and Southwestern United States. We estimated mtDNA nucleotide diversity for two mtDNA genes, COI (22 species alignments; average 16 sequences) and cytb (22 species; average 58 sequences), as well as nuclear heterozygosity and nucleotide diversity from GBS data for 118 individuals (30 species; four individuals and 6,820–44,309 loci per species). We showed that nuclear genomic diversity estimates were highly consistent across individuals for some species, while other species showed large differences depending on the locality sampled. Range size was positively correlated with both cytb diversity (Phylogenetically Independent Contrasts: R^2 = 0.31, p = 0.007) and GBS diversity (R^2 = 0.21; p = 0.006), while other predictors differed across the top models for each dataset. Mitochondrial and nuclear diversity estimates were not correlated within species, although sampling differences in the data available made these datasets difficult to compare. Further study of mtDNA and nuclear diversity sampled across species' ranges is needed to evaluate the roles of geography and life history in structuring diversity across a variety of taxonomic groups.

42   **Keywords**: (3-6) genotyping-by-sequencing, mitochondrial DNA, nucleotide diversity,

43   Serpentes, Florida, American Southwest

Introduction

Genetic information is key for investigating the evolutionary histories and demographic trajectories of populations and species. Since the advent of DNA sequencing in the late 1970s (Sanger et al. 1977), analysis of sequence variation has enabled biologists to discover species, determine relationships among lineages, reveal demographic histories, and provide assessments of population structure and diversity (Avise 2000; Hebert et al. 2003; Brito and Edwards 2009). Genetic diversity within species is especially critical for persistence in changing environments and is clearly of interest for global conservation (Laikre 2010; Laikre et al. 2020), but several challenges remain for gathering and comparing genetic information from large numbers of species. The extent to which genetic diversity is predictable given species characteristics such as range size or life history traits is an area of active interest (e.g., Leffler et al. 2012; Romiguier et al. 2014; Singhal et al. 2017; Pelletier and Carstens 2018; Barrow et al. 2021). These studies demonstrate that predictors often differ across different taxonomic groups and scales, prompting the need for additional investigations into understudied clades.

The type of genetic data chosen for comparative studies is important to consider. Most work to date has used widely available markers from organellar genomes, but the extent to which these data are representative of broader genomic patterns remains unclear. Mitochondrial DNA (mtDNA) is easily sequenced and has been used for decades, making it readily accessible from public databases for comparisons across large numbers of species (Zink and Barrowclough 2008; Miraldo et al. 2016). While certain characteristics of mtDNA, such as a high mutation rate and lack of recombination, make it desirable for population-level studies and the inference of phylogenetic relationships at shallow scales (Avise et al. 1987), mtDNA

may be poorly suited as a representation of genomic diversity in a given species (Galtier et al.

2009). As a nonrecombinant unit that is matrilineally inherited, mtDNA represents a single

realization of many possible coalescent histories of sampled taxa, and it is well appreciated that

any single genealogy may not accurately reflect population or species history (Hudson and

Turelli 2003) particularly in cases where sex-biased dispersal or multiple divergences in a short

timespan have occurred (Morin et al. 2004; Galtier et al. 2009). Notably, several investigations

comparing genetic diversity from mitochondrial and nuclear genomes within species have

found conflicting results across different taxonomic groups (Bazin 2006; Mulligan et al. 2006;

Singhal et al. 2017b). Bazin et al. (2006) showed that mtDNA did not correspond with

expectations of population abundance when comparing across broad animal groups, e.g.,

invertebrate groups did not have higher mtDNA diversity than vertebrate groups, while nuclear

sequence and allozyme datasets did meet this expectation. Comparisons within vertebrate

groups such as mammals (Mulligan et al. 2006) and lizards (Singhal et al. 2017) have

demonstrated a positive correlation between estimates of mtDNA and nuclear diversity,

however, suggesting that mtDNA may be a useful marker for understanding patterns of genetic

diversity in animals with smaller population sizes. Taken together, the contrasting results of

these studies also illustrate how the taxonomic scale of a given study can lead to different

findings.

Sampling many nuclear loci enables more robust estimates of species relationships,

population history, diversity estimates, and demographic parameters of interest (Edwards and

Beerli 2000; Carling and Brumfield 2007). The growing availability of genome-scale nuclear

datasets is promising for comparative studies (Garrick et al. 2015), but sampling many

individuals and populations for multiple species in a single study remains cost prohibitive. One

solution implemented in previous comparative studies is to obtain genome-wide estimates of

diversity from a single or few individuals as a representative of each species (e.g., Romiguier et

al. 2014; Singhal et al. 2017; Grundler et al. 2019). It is not yet clear how consistent genomic

diversity estimates are across populations of wide-ranging species that have experienced

different histories across their ranges (but see Nazareno et al. (2017) for an example in plant

populations separated by 20 km). We begin to address this question in squamates by evaluating

genomic diversity from multiple individuals and localities within species in the present study.

In addition to considering appropriate sampling strategies, identifying potential

predictors of genomic diversity within species is challenging because many species-level

characteristics may correspond with diversity within populations, diversity between

populations (i.e., genetic structure), or both. Species with larger census population sizes are

expected to have larger effective population sizes and therefore higher levels of neutral genetic

diversity within populations (Kimura 1979). This assumed relationship may also extend to total

range size, which has been considered as a proxy for census population size (e.g., Leffler et al.

2012; Singhal et al. 2017) because species that occupy large areas are presumed to be more

locally abundant. The abundance-range size relationship may be explained by a variety of

mechanisms (Gaston et al. 1997). In analyses of some taxonomic groups, population density

appears to be unrelated to range size (Novosolov et al. 2017), while empirical examples of

island and mainland bird species have demonstrated the expected positive relationship

between genomic diversity and range size (Brüniche-Olsen et al. 2019; Leroy et al. 2021).

Furthermore, the size, shape, and characteristics of a species' range can impact how individuals

move across the landscape, thus influencing rates of dispersal, gene flow, and the maintenance

of genetic structure and overall diversity within species (Wright 1943; Sexton et al. 2014; Myers

et al. 2019).

Life history and ecological traits also correspond with genetic diversity across broad

taxonomic groups (Romiguier et al. 2014; Chen et al. 2017). Body size is negatively correlated

with genetic diversity within and between populations for some groups (e.g., mammals:

Brüniche-Olsen et al. 2018; bees: López-Uribe et al. 2019; butterflies: Mackintosh et al. 2019),

which could relate to limits on population size (leading to lower within-population diversity) or

to higher dispersal (leading to lower genetic structure) in larger-bodied species (White et al.

2007; Paz et al. 2015). Fecundity, or clutch size, is also expected to relate to abundance, where

species that have more offspring will have larger and more stable population sizes through time

and therefore higher neutral diversity (Kimura 1979). Indeed, clutch size is positively correlated

with genetic diversity across animals (Romiguier et al. 2014), where species that have many,

small offspring have higher within-population genetic diversity than long-lived species with few,

large offspring. Aspects of mating system or reproductive mode have also been associated with

genetic variation in different groups. In plants, within-population genomic diversity is higher in

outcrossing species compared to selfing species (Chen et al. 2017), which is expected because

inbreeding reduces effective population size.

Different reproductive strategies may also lead to differences in dispersal distance and

resulting genetic structure. For example, in Panamanian frogs, species with direct development

exhibit greater genetic structure compared to larval developing species (Paz et al. 2015); and in

marine invertebrates, benthic direct-developing species have higher genetic structure than

those with a pelagic larval stage (Collin 2001; Lee and Boulding 2009). Another aspect of

reproduction that has not been thoroughly addressed is parity mode, whether species lay eggs

(oviparous) or have live young (viviparous). Oviparity could be associated with higher dispersal,

and less genetic structure, if females must travel long distances to locate suitable nest sites,

while viviparous species have reduced movements because of increased energetic costs (Shine

2015). In the common lizard (*Zootoca vivipara*), greater genetic structure was found in an

oviparous lineage compared to a viviparous lineage, but in this case, the pattern could be

explained by differences in demographic history (Cornetti et al. 2015). Comparisons of species-

wide genetic diversity across multiple oviparous and viviparous species are lacking thus far.

North American Squamates in Regions of Contrasting Topographic Complexity

This study focuses on the genetic diversity of squamates from North America in two

regions that were climatically suitable during the Last Glacial Maximum (LGM; Waltari et al.

2007), but differ in their topographic complexity (Badgley et al. 2017). We investigated species

from the North American Coastal Plain and the Desert Southwest (Fig. 1), both of which harbor

high reptile diversity (Jenkins et al. 2015) and are currently under threat from habitat

fragmentation, climate change, and change in wildfire regimes (Archer and Predick 2008; Noss

et al. 2015; Briggs et al. 2020). The North American Coastal Plain includes the Gulf and Atlantic

coasts in the southeastern United States and has been described as a biodiversity hotspot (Noss

et al. 2015). Although the region was never glaciated, when climatic and habitat shifts during

Pleistocene glacial cycles occurred (Williams et al. 2004), the region likely provided stable

refugial areas for many of the species in this study (Soltis et al. 2006; Weinell and Austin 2017).

Within the Desert Southwest, located in the southwestern United States and north-central

Mexico, the Chihuahuan Desert is the most biologically diverse desert and is considered among

the 200 most biologically valuable ecoregions globally (Olson and Dinerstein 1998; Briggs et al.

2020). This region is also predicted to have provided refugial areas during the LGM (Waltari et

al. 2007), but is more topographically complex with high environmental heterogeneity that

plays an important role in population genetic structure and lineage divergence (Badgley et al.

2017; Myers et al. 2019). Both regions should retain much of the ancestral genetic diversity of

the sampled species since each is suspected to have harbored refugial populations during the

Pleistocene. Furthermore, the high squamate diversity in these regions provides the

opportunity to compare species with a variety of range sizes, body sizes, and life history traits.

In this study, we combined new and existing mtDNA and genome-scale nuclear

sequence data for 30 North American squamate species to address three main questions. First,

how consistent are nuclear genomic estimates across individuals and localities within species?

Second, are measures of mtDNA and nuclear diversity within species correlated, suggesting that

mtDNA is a useful proxy for within-species diversity? Third, are species geographic range size,

body size, or life history traits (number of offspring and reproductive mode) associated with

either mtDNA or nuclear diversity within squamates?

Methods

Sample Collection

We gathered and analyzed sequence data from 30 species representing four snake

families (Viperidae, Colubridae, Natricidae, Dipsadidae) and two lizard families (Anolidae and

Anguidae) (Fig. 1). We followed the most recent taxonomic revisions in the literature, ensuring

mtDNA sequences could be assigned to a single species based on their specific localities, and

excluding sequences without locality information (Table S1). Tissue samples for 12 Coastal Plain

species (4–7 individuals each) were collected in northern Florida from 2009 to 2018, preserved

in either 95% ethanol or DMSO tissue buffer, frozen at or below -20 °C, and subsequently

archived at the Museum of Southwestern Biology (Fig. 1; Table S2). We also included previously

published data for 18 snake species distributed across the Desert Southwest (Myers et al. 2019,

2020; Myers 2021). For the nuclear dataset of these 18 species, we chose four individuals with

a similar sampling strategy as the Coastal Plain species. Briefly, we chose two individuals each

from two localities per species when possible, focusing primarily on the Chihuahuan Desert and

a similar distance (~200 km) between localities. This standardized distance enabled us to make

comparisons among species without the potentially confounding effects of geographic distance

on genetic structure.

Range Maps and Trait Data

Species range maps were downloaded as shapefiles from the IUCN Red List of

Threatened Species Version 6.1 (www.iucnredlist.org/). Maps were edited in QGIS 3.12 as

needed to exclude non-native parts of the range where species have been introduced and to

edit species ranges according to recent taxonomic changes (Table S1). The R packages 'rgdal'

(Bivand et al. 2021) and 'geosphere' (Karney 2013; Hijmans 2021) were used to visualize ranges

and calculate total range area in kilometers squared. Species trait information was obtained

primarily from Burbrink and Myers (2015), including maximum body size (log-transformed),

average clutch size (log-transformed), and parity (viviparous or oviparous). Trait data for the

two lizard species were obtained from field guides and online natural history accounts (Powell

et al. 2016; https://www.virginiaherpetologicalsociety.com; https://animaldiversity.org).

Mitochondrial Sequences

For the 12 Coastal Plain species, genomic DNA was extracted from tail or liver tissue

with the E.Z.N.A. DNA Tissue Kit (Omega Bio-Tek) according to the manufacturer's instructions.

We amplified a 658-base pair portion of the mtDNA cytochrome c oxidase 1 gene (COI) using

primers ReptBCF_COI and ReptBCR_COI (Castañeda and de Queiroz 2011). The thermal profile

included an initial denaturation step at 94° C for 3 minutes, 35 cycles of denaturing at 94° for 45

seconds, annealing at 53° for 30 seconds, elongating at 72° for 60 seconds, and a final extension

step at 72° for 10 minutes. PCR products were visualized on agarose gels to ensure successful

amplification, cleaned with EXOSAP-IT (Affymetrix Inc.), and sequenced at The Ohio State

University (OSU) Comprehensive Cancer Center Genomics Shared Resource. Species-specific

sequencing primers were designed when the original PCR primers did not perform adequately

(Table S3). Additional mtDNA sequences for two genes, COI and cytochrome b (cytb), were

obtained from NCBI GenBank for species with data available. Sequences for each species were

aligned in Geneious 2020.1.1 (https://www.geneious.com) using MAFFT 7.450 (Katoh and

Standley 2013).

Nuclear Data Collection

Genotyping-by-sequencing (GBS) libraries were prepared following a modified version of

the protocol described in Elshire et al. (2011). For four samples per Coastal Plain species (48

individuals), we digested 100 ng of input DNA with the enzyme Pst1, ligated 8 µL of adapter mix

including unique barcodes, pooled libraries, and performed a bead cleanup with Sera-Mag

Speedbeads (Rohland and Reich 2012). Final PCR amplification was conducted with 16 cycles in

8 replicate reactions, followed by pooling, bead cleanup, and quantification via a Qubit

fluorometer (Life Technologies). Libraries were size selected to 200-500 bp using a Blue Pippen

(Sage Science Inc.), quantified with a Bioanalyzer (Agilent Technologies), and sequenced at the

OSU Comprehensive Cancer Center Genomics Shared Resource on an Illumina HiSeq 4000 with

150 bp paired-end sequencing. For four samples per Desert Southwest species (72 individuals),

we downloaded GBS reads from the NCBI Sequence Read Archive (Table S2).

Sequenced GBS reads from all 30 species were assembled through the ipyrad 0.7.28

pipeline (Eaton and Overcast 2020) with the following settings. After demultiplexing (no

mismatches allowed in barcodes), the R1 reads for the four individuals per species were

assembled to generate within-species datasets. We trimmed reads for the Southeastern U.S.

species to 125 bp prior to assembly using the built-in ipyrad option which uses the software

tool 'cutadapt' (trim_reads = 0, -25). We used the de novo assembly method with a clustering

similarity threshold of 85% (clust_threshold = 0.85), maximum clustering depth of 10000 reads

(maxdepth = 10000), maximum of five low quality base calls per read (max_low_qual_bases = 5;

with Q<20), minimum depth for base calling of six reads (mindepth_statistical = 6 and

mindepth_majrule = 6, strict filtering for adapters (filter_adapters = 2), and we retained reads

longer than 35 bp after trimming adapters (filter_min_trim_len = 35). We allowed a maximum

of eight heterozygous sites in consensus sequences (max_Hs_consens = 5), a maximum of 20

SNPs per locus (max_SNPs_locus = 20) and required all four individuals (no missing data) to be

included in a locus to retain that locus in output files (min_samples_locus = 4).

Genetic Diversity Metrics

Measures of mtDNA diversity were calculated for each species using the R package

'pegas' (Paradis 2010). We calculated nucleotide diversity (pi or π, Nei 1987) for COI and cytb

alignments of each species with the nuc.div function. Nuclear diversity within species was

estimated from each GBS dataset using the R packages 'adegenet' (Jombart 2008; Jombart and

Ahmed 2011) and 'PopGenome' (Pfeifer et al. 2014). Expected heterozygosity (Hs) was

computed from the unique SNPs outfile from ipyrad for each species using 'adegenet'. To

calculate π, 'PopGenome' reads in a directory of FASTA files including an alignment for each

locus; we used a custom Python script (available on Dryad) to generate this directory for each

species using the alleles outfiles (full sequences) from ipyrad. We calculated π within each

sample (two alleles per individual) and then determined the mean and standard deviation

across individuals, hereafter referred to as "mean within-individual π". For comparison, we also

calculated π within each species from the alignments including four individuals (eight alleles)

per species, hereafter "within-species π". Species-level and within-population nuclear diversity

can be estimated from a single individual when sufficient loci are sampled (e.g., Chen et al.

2017; Grundler et al. 2019). Where possible, we included two individuals per species from each

sampled locality to investigate how consistent genomic diversity estimates are across different

individuals and localities within species.

Statistical Analyses

We compared mtDNA (COI π, cytb π) and nuclear (mean within-individual π, within-

species π) diversity estimates within species using pairwise correlations in R. We then

constructed phylogenetic generalized least square (PGLS) models to assess the importance of

different predictors on genetic diversity, while accounting for the possibility that closely related

species may share similar genomic diversity and species traits. We downloaded 100 phylogeny

subsets for the 30 species in our study from VertLife.org, which includes the squamate

relationships from Tonini et al. (2016), and generated a least-squares consensus tree using

phytools (Revell 2012) for our analyses. We constructed a model set for each of three

responses: mtDNA COI π, mtDNA cytb π, and nuclear within-species π. For each response,

predictors of interest included geographic range size, maximum body size, average clutch size,

and parity. To assess effects of sample size on diversity metrics, we included the number of

individuals sampled as a covariate in mtDNA models, and the number of GBS loci as a covariate

for nuclear models. We included the region (Coastal Plain or Desert Southwest) in our initial

PGLS models for nuclear within-species π and used Welch's two-sample t-tests to check for

differences between the GBS datasets generated in this study and those generated previously.

Analyses were conducted using the R packages 'ape' (Paradis and Schliep 2019), 'geiger'

(Harmon et al. 2008), and 'nlme' (Pinheiro et al. 2022). We constructed PGLS models using two

models of evolution, Brownian Motion (BM) and Ornstein-Uhlenbeck (OU). The BM model

assumes traits evolve according to a random walk, while the OU model incorporates a trait

optimum towards which traits are pulled (Felsenstein 1985; Martins and Hansen 1997). For

each model set, all combinations of the predictors of interest were included and models were

ranked based on the corrected Akaike Information Criterion (AICc). We ran model selection

analyses using the R package 'MuMIn' (Bartón 2022) to determine which model of evolution

had a better fit. We then constructed models incorporating all combinations of the predictors

of interest and ran model selection analyses to assess which sets of predictors were included in

the best-fit models for each diversity measure. We further assessed relationships between

continuous variables using Spearman's rank correlation tests, phylogenetically independent

contrasts implemented in 'ape' (Felsenstein 1985), and linear models.

Results

Data Summary

We generated 62 new COI sequences (GenBank accession numbers ON911378–

ON911439) and combined these with available sequences on GenBank. Of the 30 squamates in

our study, we were able to analyze COI alignments for 22 species (4–69 sequences per

alignment; average 16) and cytb alignments for 22 species (8–143 sequences; average 58).

Three species (*Crotalus oreganus*, *Hypsiglena jani*, *Trimorphodon vilkinsonii*) did not have

sufficient mtDNA data available for either COI or cytb and 17 species had alignments for both

genes (Table 1). We sequenced new GBS data for 47 individuals from 12 species (NCBI SRA

accession numbers SAMN31800422–SAMN31800468; PRJNA903383) and calculated genomic

diversity metrics for 118 individuals from 30 species. One *Thamnophis sirtalis* sample had poor

sequencing success and was removed from subsequent analyses. The assembled GBS datasets

included 2,923–25,086 loci for the unique SNP datasets and 6,820–44,309 loci for the full

sequence datasets depending on the species (Table 1).

Mitochondrial and Nuclear Diversity

Mitochondrial diversity within species ranged from 0.0015 to 0.0812 for COI π and from

0.0005 to 0.0717 for cytb π. Diversity estimates for the two mtDNA genes were correlated

within species (Fig. S1; Spearman's rho (ρ) = 0.897, p < 2.2e-16). For the nuclear GBS data,

within-species heterozygosity ranged from 0.3 to 0.346 and within-species π ranged from

0.0005 to 0.0083. Mean within-individual π ranged from 0.0004 to 0.0056 and there was

substantial variation among individuals and localities for some species (Fig. 1). For species such

as *Thamnophis marcianus* and *Masticophis flagellum*, the within-individual π estimates were

consistent between the two individuals within a locality, but values for one locality were nearly

double the values for the second locality. For other species such as *Crotalus adamanteus* and

Sonora episcopa, the within-individual π estimates were highly consistent for all four

individuals. For still other species such as *Thamnophis sauritus* and *Lampropeltis splendida*,

there was large variation in π estimates even for individuals collected from the same locality.

The within-species π estimates were strongly correlated with the mean within-individual

π estimates (Fig. S1; Spearman's ρ = 0.986, p < 2.2e-16), and we used the within-species

estimate to represent overall species nuclear diversity for subsequent analyses. Metrics of

diversity for mtDNA and nuclear GBS data were not correlated for the species represented in

both datasets (Fig. 2). Nuclear within-species π was not correlated with mtDNA COI π

(Spearman's ρ = 0.16, p = 0.476) or mtDNA cytb π (Spearman's ρ = 0.293, p = 0.185). We did not

detect any clear differences in nuclear diversity between datasets sampled from the Desert

Southwest compared to the Coastal Plain (Fig. S2).

Predictors of Squamate Genetic Diversity

333   The OU model of evolution was the better fit for all three response variables (model

weights > 0.986; Table 2). Different sets of predictors were included in the top models for the

three response variables (Table 3). For mtDNA COI, parity was the only variable consistently

included in the top models. Diversity between viviparous and oviparous species was not

significantly different according to the top PGLS model (Table 4), although oviparous species did

have higher COI diversity than viviparous species without phylogenetic correction (Fig. 3a; t-

test: t = 2.34, p = 0.03). For mtDNA cytb, range size and parity were always included in the top

models. Within-species cytb diversity had a positive relationship with range size (Fig. 3),

suggesting that species with larger ranges tend to have higher mtDNA diversity. This

relationship was consistent for both uncorrected (Spearman's ρ = 0.64; p = 0.002; Fig. 3c) and

phylogenetically corrected datasets (phylogenetically independent contrasts: R^2 = 0.31; p =

0.007; Fig. 3d). Oviparous species had higher cytb diversity compared to viviparous species (Fig.

3b; t-test: t = 3.11, p = 0.008).

346   Range size and clutch size were consistently included in the top models for nuclear

within-species π (Table 2). Nuclear within-species π had a positive relationship with both range

size and average clutch size (Table 4; Fig. 3). The relationship between nuclear π and range size

was consistent for both uncorrected (Spearman's ρ = 0.56; p = 0.001; Fig. 3e) and

phylogenetically corrected datasets (PIC: R^2 = 0.21; p = 0.006; Fig. 3f). The relationship

between nuclear π and average clutch size was significant for the uncorrected data (Spearman's

ρ = 0.53; p = 0.003; Fig. 3g), but not based on phylogenetically independent contrasts (R^2 =

0.04; p = 0.288; Fig. 3h).

Discussion

We compared genetic diversity estimates from mitochondrial and nuclear genomes for

30 North American squamate species. Our analyses demonstrated that range size is positively

correlated with mtDNA and nuclear diversity in squamates and that life history traits are

inconsistent predictors of diversity for these datasets. We found that mtDNA and nuclear

diversity estimates were not correlated, but we note that several aspects of sampling made

these datasets difficult to compare. Here, we discuss sampling considerations for this and

future studies, including the number of individuals, localities, species, and loci available or

feasible to sample. Despite these challenges, we provide empirical evidence that geography is a

key element for understanding patterns of genomic diversity within and across species.

Sampling Considerations Across Geography, Taxonomy, and Genomes

Comparative population genomics is a developing field that aims to uncover the forces

underlying genomic diversity by comparing across multiple species (Ellegren and Galtier 2016;

Edwards et al. 2022). Studies typically focus on the roles of selection, neutrality, and

recombination in structuring genetic variation in a primarily non-geographic context, thus

sampling usually consists of a single or few individuals to represent each species (e.g. Romiguier

et al. 2014; Chen et al. 2017). In contrast, comparative phylogeography takes a geography-

centric view to understanding historical processes underlying current genetic variation within

species and lineages. Ideally, species have been sampled across their entire range and as

densely as possible, often revealing the presence of multiple lineages within nominal species

(e.g., Schield et al. 2018). A third emerging field, macrogenetics, focuses on an even broader

geographic and taxonomic scale, typically using repurposed data from relatively few genetic

markers to investigate large-scale patterns and predictors of genetic variation within species

(Leigh et al. 2021). Macrogenetic studies rely on sampling choices and data from previous

studies, but until relatively recently, genome-wide datasets were not common in

phylogeographic studies (Garrick et al. 2015). The gradual accumulation of these data over time

and the introduction of new data aggregators (e.g. Pelletier et al. 2022) potentially increases

the scale of comparative studies that use opportunistic sampling, but the novel data collected

here offer a direct comparison between two ecoregions.

Thoroughly sampling many individuals from many species across geographic space

remains costly, thus a pressing question relevant to the design of comparative studies is: How

representative are genomic diversity estimates from a few individuals or from a narrow part of

a species' range? Our GBS data indicate that this answer depends on several factors – certain

species (e.g., *Crotalus adamanteus*) had highly consistent genomic diversity estimates from

localities sampled ~200 km apart, while others (e.g., *Thamnophis marcianus*) had two-fold

differences in genomic diversity between localities (Fig. 1). This result indicates that using a

single individual or locality to represent diversity for an entire species does not always provide a

complete picture of diversity within species. We expected to find similar levels of genetic

diversity across individuals because we sampled them from historically stable regions. Thus, it

may seem surprising that such differences are already emerging. We predict even greater

differences when populations are sampled from topographically complex regions or in areas

that have been influenced by historical climate change (e.g., Howes and Lougheed 2008; Nali et

al. 2020), a consideration that should continue to be addressed in future studies.

Our study provides an initial comparison of squamate genomic diversity from five

families sampled in two species-rich regions, the North American Desert Southwest and North

American Coastal Plain. We found considerable variation in diversity among species within

some families (e.g., Colubridae and Natricidae; Fig. 1), but further taxon sampling is needed for

robust comparisons of genomic diversity across families. In general, the Desert Southwest has

more heterogenous environments and landscapes compared to the Coastal Plain, and Florida in

particular, where populations were sampled (Noss et al. 2015; Badgley et al. 2017). Thus far, we

did not detect any clear differences in average genomic diversity levels between the two

regions (Fig. S2), although several Desert Southwest species had high variation in genomic

diversity among individuals and localities (Fig. 1). This result is expected given the

heterogeneous nature of the region, where large differences in climate, elevation, and habitat

on relatively small spatial scales can conceivably lead to differences in population sizes and

standing genomic diversity among nearby localities (Myers et al. 2019).

We found that mtDNA and GBS diversity were not correlated, suggesting that mtDNA

may not be a reliable proxy for within-species diversity in North American squamates. Several

differences in our sampling for mtDNA and nuclear data made these datasets difficult to

compare, however. The number of mtDNA sequences available for some species was limited,

producing COI and cytb alignments for only 22 species each. These mtDNA sequences most

likely represent a much broader geographic area within each species compared to our GBS

data, though it is not possible to quantify the influence of geographic distance because sample

localities were not associated with the majority of GenBank sequences (Marques et al. 2013).

On the other hand, the GBS data consist of several thousand loci and provide a better

representation of genome-wide diversity within species compared to a single mtDNA gene.

These differences in power and geographic representation, along with relatively few data

points for comparison (22 species), may explain the lack of correlation between mtDNA and

nuclear diversity metrics in our study. Previous studies have found a relationship between

mtDNA and nuclear diversity in some taxonomic groups including mammals (47 species;

Mulligan et al. 2006) and Australian lizards (60 species; Singhal et al. 2017). Further study of this

topic is important for comparative studies because mtDNA is readily available for many more

species, but reduced representation nuclear genome datasets are increasing and have the

potential to provide new insights on overall species diversity.

Additional Challenges for Repurposed Data

One challenge for comparative studies such as ours is the difficulty of gathering

different data types from species complexes that have undergone recent taxonomic revisions.

When possible, we manually edited mtDNA alignments and IUCN range maps to reflect data

from a single described lineage (Table S1). In some cases, however, recent taxonomic revisions

and limited geographic information about existing sequences and traits made it impossible to

assign data to newly-described species. For those taxa, we excluded mtDNA data and assumed

trait information from former species designations would be representative for the whole

complex. We provide the alignments and GenBank accession numbers for the data we analyzed

(https://doi.org/10.5061/dryad.qfttdz0ks) such that they can be reassigned and reanalyzed in

follow-up studies. Future efforts to link geographic coordinates with genetic sequences in open

access databases will continue to improve the prospects of comparative studies (Pelletier et al.

2022). Frequent updates to geographic distribution and trait databases will also be needed to

facilitate large-scale comparative studies.

The approaches used to obtain and analyze nuclear genome-wide datasets are

important to consider for comparative studies of genetic diversity. One appealing characteristic

of mtDNA data for data repurposing and comparing species sequenced across different studies

is the ease of aligning known genes. Restriction site associated methods such as GBS provide

many more loci for analysis, but without an annotated reference genome for most species of

interest, these loci are anonymous. Furthermore, the locus assemblies and resulting

parameters are highly dependent on the parameter settings of the pipeline used for analysis

(Harvey et al. 2016; Shafer et al. 2017). Our study used existing squamate datasets from the

Desert Southwest that we reassembled to ensure the same ipyrad version and parameters were

used for comparison with new GBS data we collected for the Coastal Plain species. We found

that heterozygosity values were highly sensitive to the parameter settings used. Prior to

reassembly, heterozygosity estimates for the Desert Southwest species were an order of

magnitude smaller than those of the Coastal Plain species (Table S4). These contrasting results

demonstrate the additional challenges of comparing GBS and similar datasets collected from

different studies. Targeted sequence capture approaches may be preferable to increase

comparability across studies, though they remain more expensive and require more initial

effort to design probe kits (Harvey et al. 2016; Singhal et al. 2017a; Hutter et al. 2022).

Predictors of Squamate Genomic Diversity

Range size was positively correlated with nucleotide diversity within squamate species

for both mtDNA (cytb) and nuclear data, despite differences in geographic and locus sampling.

Previous studies have found mixed results in a variety of taxonomic groups and with different

sampling strategies. For example, a global study including single-gene sequences from more

than 8,000 eukaryotic species found that total range size was one of the most important

predictors of genetic structure within species (Pelletier and Carstens 2018). In contrast,

(Romiguier et al. 2014) analyzed transcriptome data from 2-10 individuals of 76 animal species

and found no correlation between genomic diversity and range size. Detailed focus within

taxonomic groups demonstrated correlations between range size and genetic diversity in

Drosophila (Leffler et al. 2012), Australian lizards (Singhal et al. 2017b), and cetaceans (Vachon

et al. 2018), but not in butterflies (Mackintosh et al. 2019). To our knowledge, we provide the

first evidence that total range size may be a useful predictor of squamate genomic diversity.

Future studies including additional species, biogeographic regions, and nuclear sampling across

species' ranges would provide important insights about the generality of these findings and the

potential influence of fluctuations in population size on genomic variation.

The life history and ecological traits we included were less consistent predictors of

mitochondrial and nuclear diversity in squamates. Body size has negative associations with

genetic diversity in various taxonomic groups including mammals (Brüniche-Olsen et al. 2018),

frogs (Paz et al. 2015), and insects (López-Uribe et al. 2019; Mackintosh et al. 2019). One

possible explanation is that larger-bodied animals can disperse over longer distances, leading to

less genetic structure and overall diversity across a species range. Our results do not provide

compelling evidence for squamates, perhaps indicating that body size is not a good indicator of

dispersal for this group. Our results for clutch size are consistent with the prediction that

species with larger clutch sizes harbor higher genomic diversity (e.g., Romiguier et al. 2014),

though this relationship did not remain after phylogenetic correction (Fig. 3h). These patterns

will require further scrutiny with increased species sampling in future studies.

The potential association of parity with mitochondrial diversity, where oviparous (egg-

laying) species had higher diversity on average than viviparous (live-bearing) species, is

intriguing and should also be investigated further with more comprehensive sampling. Since the

mitochondrial sequences were sampled across species ranges, our results may reflect greater

genetic structure for oviparous species, in contrast with the prediction that oviparous species

must disperse to find suitable nest sites (Shine 2015). The evolution of viviparity is associated

with colder climates and faster development times in cool temperatures (Ma et al. 2018), but

our study lacked sampling across broad latitudinal or climatic gradients. Furthermore, most of

the oviparous species included in this study belong to the family Colubridae, but viviparity in

squamates has evolved from oviparity at least 34 and perhaps more than 100 times (Pyron and

Burbrink 2014; Blackburn 2015). Squamates therefore present an excellent opportunity to

continue investigating whether life history traits such as clutch size or parity are associated with

intraspecific genomic diversity while taking phylogeny into account.

Overall, we provide new insights into the predictors of diversity within squamates

considering both mitochondrial and nuclear genome patterns. We showed that geographic

range size corresponds with intraspecific genomic diversity even when a relatively small part of

the range is sampled. We also demonstrate that genomic diversity can vary widely between

populations on small spatial scales, suggesting that sampling a few individuals may not

adequately represent diversity levels within species. As genome-wide datasets continue being

generated, comparative studies of intraspecific diversity across species ranges will increase

understanding of the roles of geography and life history in structuring diversity across a variety

of taxonomic groups.

Funding

This work was supported by The Ohio State University (OSU) Undergraduate Research

Scholarship awarded to IEL and by the OSU President's Postdoctoral Scholars Program through

an award to LNB. Computational resources were provided by the Ohio Supercomputer Center

via a grant to BCC (PAA0202). LNB and BCC were supported via grants from the National

Science Foundation (LNB: DEB-2112946 and BCC: DBI-1910623).

Acknowledgements

We thank Moses Michelsohn for providing tissues from Florida and Megan Smith for laboratory

and analytical support. Samples were obtained in accordance with collecting permits from the

Florida Fish and Wildlife Conservation Commission and animal care protocols approved by the

Florida State University Animal Care and Use Committee.

Data Availability

We have deposited the primary data underlying these analyses as follows:

- Sampling locations, field numbers, museum catalogue numbers, R/Python scripts, and

datasets used for analyses and visualization: Dryad link -

https://doi.org/10.5061/dryad.qfttdz0ks

- DNA sequences: GenBank accession numbers ON911378–ON911439; NCBI SRA

accession numbers: SAMN31800422–SAMN31800468 (PRJNA903383)

References

References

Archer S.R., Predick K.I. 2008. Climate Change and Ecosystems of the Southwestern United States.
Rangelands. 30:23–28.

Avise J.C. 2000. Phylogeography: the history and formation of species. Cambridge, Massachussetts:
Harvard University Press.

Avise J.C., Arnold J., Ball R.M., Bermingham E., Lamb T., Neigel J.E., Reeb C.A., Saunders N.C. 1987.
Intraspecific Phylogeography: The mitochondrial DNA bridge between population genetics and
systematics. Annual Review of Ecology and Systematics. 18:489–522.

Badgley C., Smiley T.M., Terry R., Davis E.B., DeSantis L.R.G., Fox D.L., Hopkins S.S.B., Jezkova T., Matocq
M.D., Matzke N., McGuire J.L., Mulch A., Riddle B.R., Roth V.L., Samuels J.X., Strömberg C.A.E.,
Yanites B.J. 2017. Biodiversity and Topographic Complexity: Modern and Geohistorical
Perspectives. Trends in Ecology & Evolution. 32:211–226.

Barrow L.N., Masiero da Fonseca E., Thompson C.E.P., Carstens B.C. 2021. Predicting amphibian
intraspecific diversity with machine learning: Challenges and prospects for integrating traits,
geography, and genetic data. Molecular Ecology Resources. 21:2818–2831.

Bartón K. 2022. MuMIn: Multi-Model Inference. R package version 1.46.0. https://CRAN.R-
project.org/package=MuMIn. .

Bazin E. 2006. Population Size Does Not Influence Mitochondrial Genetic Diversity in Animals. Science.
312:570–572.

Bivand R., Keitt T., Rowlingson B. 2021. rgdal: Bindings for the "Geospatial" Data Abstraction Library. R
package version 1.5-23. https://CRAN.R-project.org/package=rgdal. .

Blackburn D.G. 2015. Evolution of viviparity in squamate reptiles: Reversibility reconsidered. Journal of
Experimental Zoology Part B: Molecular and Developmental Evolution. 324:473–486.

Briggs M.K., Lozano-Cavazos E.A., Mills Poulos H., Ochoa-Espinoza J., Rodriguez-Pineda J.A. 2020. The
Chihuahuan Desert: A Binational Conservation Response to Protect a Global Treasure.
Encyclopedia of the World's Biomes. Elsevier. p. 126–138.

Brito P.H., Edwards S.V. 2009. Multilocus phylogeography and phylogenetics using sequence-based
markers. Genetica. 135:439–455.

Brüniche-Olsen A., Kellner K.F., Anderson C.J., DeWoody J.A. 2018. Runs of homozygosity have utility in
mammalian conservation and evolutionary studies. Conservation Genetics. 19:1295–1307.

Brüniche-Olsen A., Kellner K.F., DeWoody J.A. 2019. Island area, body size and demographic history
shape genomic diversity in Darwin's finches and related tanagers. Molecular Ecology. 28:4914–
4925.

Burbrink F.T., Myers E.A. 2015. Both traits and phylogenetic history influence community structure in
snakes over steep environmental gradients. Ecography. 38:1036–1048.

Carling M.D., Brumfield R.T. 2007. Gene sampling strategies for multi-locus population estimates of
genetic diversity. PLoS ONE. 2:e160–e160.

Castañeda M. del R., de Queiroz K. 2011. Phylogenetic relationships of the Dactyloa clade of Anolis
lizards based on nuclear and mitochondrial DNA sequence data. Molecular Phylogenetics and
Evolution. 61:784–800.

Chen J., Glémin S., Lascoux M. 2017. Genetic diversity and the efficacy of purifying selection across plant
and animal species. Molecular Biology and Evolution. 34:1417–1428.

Collin R. 2001. The effects of mode of development on phylogeography and population structure of
North Atlantic Crepidula (Gastropoda: Calyptraeidae). Molecular Ecology. 10:2249–2262.

Cornetti L., Ficetola G.F., Hoban S., Vernesi C. 2015. Genetic and ecological data reveal species
boundaries between viviparous and oviparous lizard lineages. Heredity. 115:517–526.

Eaton D.A.R., Overcast I. 2020. ipyrad: Interactive assembly and analysis of RADseq datasets.
Bioinformatics. 36:2592–2594.

Edwards S.V., Beerli P. 2000. Perspective: Gene divergence, population divergence, and the variance in
coalescence time in phylogeographic studies. Evolution. 54:1839–1854.

Edwards S.V., Robin V.V., Ferrand N., Moritz C. 2022. The Evolution of Comparative Phylogeography:
Putting the Geography (and More) into Comparative Population Genomics. Genome Biology and
Evolution. 14:evab176.

Ellegren H., Galtier N. 2016. Determinants of genetic diversity. Nat Rev Genet. 17:13.

Elshire R.J., Glaubitz J.C., Sun Q., Poland J.A., Kawamoto K., Buckler E.S., Mitchell S.E. 2011. A robust,
simple genotyping-by-sequencing (GBS) approach for high diversity species. PLoS ONE. 6:1–10.

Felsenstein J. 1985. Phylogenies and the comparative method. American Naturalist. 125:1–15.

Galtier N., Nabholz B., Glémin S., Hurst G.D.D. 2009. Mitochondrial DNA as a marker of molecular
diversity: a reappraisal. Molecular Ecology. 18:4541–4550.

Garrick R.C., Bonatelli I.A.S., Hyseni C., Morales A., Pelletier T.A., Perez M.F., Rice E., Satler J.D., Symula
R.E., Thomé M.T.C., Carstens B.C. 2015. The evolution of phylogeographic datasets. Molecular
Ecology. 24:1164–1171.

Gaston K.J., Blackburn T.M., Lawton J.H. 1997. Interspecific Abundance-Range Size Relationships: An
Appraisal of Mechanisms. Journal of Animal Ecology. 66:579–601.

Grundler M.R., Singhal S., Cowan M.A., Rabosky D.L. 2019. Is genomic diversity a useful proxy for census
population size? Evidence from a species-rich community of desert lizards. Molecular
Ecology.:1664–1674.

Harmon L.J., Weir J.T., Brock C.D., Glor R.E., Challenger W. 2008. GEIGER: investigating evolutionary
radiations. Bioinformatics. 24:129–131.

Harvey M.G., Smith B.T., Glenn T.C., Faircloth B.C., Brumfield R.T. 2016. Sequence capture versus
restriction site associated DNA sequencing for shallow systematics. Systematic Biology. 65:910–
924.

Hebert P.D.N., Cywinska A., Ball S.L., deWaard J.R. 2003. Biological identifications through DNA
barcodes. Proc Biol Sci. 270:313–321.

Hijmans R.J. 2021. geosphere: Spherical Trigonometry. R package version 1.5-14. https://CRAN.R-
project.org/package=geosphere. .

Howes B.J., Lougheed S.C. 2008. Genetic diversity across the range of a temperate lizard. Journal of
Biogeography. 35:1269–1278.

Hudson R.R., Turelli M. 2003. Stochasticity Overrules the "Three-Times Rule": Genetic Drift, Genetic
Draft, and Coalescence Times for Nuclear Loci Versus Mitochondrial Dna. Evolution. 57:182–190.

Hutter C.R., Cobb K.A., Portik D.M., Travers S.L., Wood Jr. P.L., Brown R.M. 2022. FrogCap: A modular
sequence capture probe-set for phylogenomics and population genetics for all frogs, assessed
across multiple phylogenetic scales. Molecular Ecology Resources. 22:1100–1119.

Jenkins C.N., Houtan K.S.V., Pimm S.L., Sexton J.O. 2015. US protected lands mismatch biodiversity
priorities. PNAS. 112:5081–5086.

Jombart T. 2008. adegenet: a R package for the multivariate analysis of genetic markers. Bioinformatics.
24:1403–1405.

Jombart T., Ahmed I. 2011. adegenet 1.3-1: new tools for the analysis of genome-wide SNP data.
Bioinformatics. 27:3070–3071.

Karney C.F.F. 2013. Algorithms for geodesics. J Geod. 87:43–55.

Katoh K., Standley D.M. 2013. MAFFT Multiple Sequence Alignment Software Version 7: Improvements
in Performance and Usability. Molecular Biology and Evolution. 30:772–780.

Kimura M. 1979. The Neutral Theory of Molecular Evolution. Scientific American. 241:98–129.

Laikre L. 2010. Genetic diversity is overlooked in international conservation policy implementation.
Conserv Genet. 11:349–354.

Laikre L., Hoban S., Bruford M.W., Segelbacher G., Allendorf F.W., Gajardo G., Rodríguez A.G., Hedrick
P.W., Heuertz M., Hohenlohe P.A., Jaffé R., Johannesson K., Liggins L., MacDonald A.J.,
OrozcoterWengel P., Reusch T.B.H., Rodríguez-Correa H., Russo I.-R.M., Ryman N., Vernesi C.
2020. Post-2020 goals overlook genetic diversity. Science. 367:1083–1085.

Lee H.J., Boulding E.G. 2009. Spatial and temporal population genetic structure of four northeastern
Pacific littorinid gastropods: the effect of mode of larval development on variation at one
mitochondrial and two nuclear DNA markers. Molecular Ecology. 18:2165–2184.

Leffler E.M., Bullaughey K., Matute D.R., Meyer W.K., Ségurel L., Venkat A., Andolfatto P., Przeworski M.
2012. Revisiting an Old Riddle: What Determines Genetic Diversity Levels within Species? PLoS
Biology. 10.

Leigh D.M., van Rees C.B., Millette K.L., Breed M.F., Schmidt C., Bertola L.D., Hand B.K., Hunter M.E.,
Jensen E.L., Kershaw F., Liggins L., Luikart G., Manel S., Mergeay J., Miller J.M., Segelbacher G.,
Hoban S., Paz-Vinas I. 2021. Opportunities and challenges of macrogenetic studies. Nat Rev
Genet. 22:791–807.

Leroy T., Rousselle M., Tilak M.-K., Caizergues A.E., Scornavacca C., Recuerda M., Fuchs J., Illera J.C., De
Swardt D.H., Blanco G., Thébaud C., Milá B., Nabholz B. 2021. Island songbirds as windows into
evolution in small populations. Current Biology. 31:1303-1310.e4.

López-Uribe M.M., Jha S., Soro A. 2019. A trait-based approach to predict population genetic structure in
bees. Mol Ecol. 28:1919–1929.

Ma L., Buckley L.B., Huey R.B., Du W.-G. 2018. A global test of the cold-climate hypothesis for the
evolution of viviparity of squamate reptiles. Global Ecology and Biogeography. 27:679–689.

Mackintosh A., Laetsch D.R., Hayward A., Charlesworth B., Waterfall M., Vila R., Lohse K. 2019. The
determinants of genetic diversity in butterflies. Nature Communications. 10:1–9.

Marques A.C., Maronna M.M., Collins A.G. 2013. Putting GenBank Data on the Map. Science. 341:1341–
1341.

Martins E.P., Hansen T.F. 1997. Phylogenies and the Comparative Method: A General Approach to
Incorporating Phylogenetic Information into the Analysis of Interspecific Data. The American
Naturalist. 149:646–667.

Miraldo A., Li S., Borregaard M.K., Flórez-Rodríguez A., Gopalakrishnan S., Rizvanovic M., Wang Z.,
Rahbek C., Marske K.A., Nogués-Bravo D. 2016. An Anthropocene map of genetic diversity.
Science. 353:1532–1535.

Morin P.A., Luikart G., Wayne R.K., the SNP workshop group. 2004. SNPs in ecology, evolution and
conservation. Trends in Ecology & Evolution. 19:208–216.

Mulligan C.J., Kitchen A., Miyamoto M.M. 2006. Comment on "Population Size Does Not Influence
Mitochondrial Genetic Diversity in Animals." Science. 314:1390–1390.

Myers E.A. 2021. Genome-wide data reveal extensive gene flow during the diversification of the western
rattlesnakes (Viperidae: Crotalinae: Crotalus). Molecular Phylogenetics and Evolution.
165:107313.

Myers E.A., McKelvy A.D., Burbrink F.T. 2020. Biogeographic barriers, Pleistocene refugia, and climatic
gradients in the southeastern Nearctic drive diversification in cornsnakes (*Pantherophis*
guttatus complex). Mol Ecol. 29:797–811.

Myers E.A., Xue A.T., Gehara M., Cox C.L., Davis Rabosky A.R., Lemos-Espinal J., Martínez-Gómez J.E.,
Burbrink F.T. 2019. Environmental heterogeneity and not vicariant biogeographic barriers
generate community-wide population structure in desert-adapted snakes. Mol Ecol. 28:4535–
4548.

Nali R.C., Becker C.G., Zamudio K.R., Prado C.P.A. 2020. Topography, more than land cover, explains
genetic diversity in a Neotropical savanna tree frog. Diversity and Distributions. 26:1798–1812.

Nazareno A.G., Bemmels J.B., Dick C.W., Lohmann L.G. 2017. Minimum sample sizes for population
genomics: an empirical study from an Amazonian plant species. Molecular Ecology Resources.
17:1136–1147.

Nei M. 1987. Molecular evolutionary genetics. New York: Columbia University Press.

Noss R.F., Platt W.J., Sorrie B.A., Weakley A.S., Means D.B., Costanza J., Peet R.K. 2015. How global
biodiversity hotspots may go unrecognized: lessons from the North American Coastal Plain.
Diversity Distrib. 21:236–244.

Novosolov M., Rodda G.H., North A.C., Butchart S.H.M., Tallowin O.J.S., Gainsbury A.M., Meiri S. 2017.
Population density–range size relationship revisited. Global Ecology and Biogeography.
26:1088–1097.

Olson D.M., Dinerstein E. 1998. The Global 200: A Representation Approach to Conserving the Earth's
Most Biologically Valuable Ecoregions. Conservation Biology. 12:502–515.

Paradis E. 2010. pegas: an R package for population genetics with an integrated–modular approach.
Bioinformatics. 26:419–420.

Paradis E., Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary analyses
in R. Bioinformatics. 35:526–528.

Paz A., Ibáñez R., Lips K.R., Crawford A.J. 2015. Testing the role of ecology and life history in structuring
genetic variation across a landscape: A trait-based phylogeographic approach. Molecular
Ecology. 24:3723–3737.

Pelletier T.A., Carstens B.C. 2018. Geographical range size and latitude predict population genetic
structure in a global survey. Biol. Lett. 14:20170566.

Pelletier T.A., Parsons D.J., Decker S.K., Crouch S., Franz E., Ohrstrom J., Carstens B.C. 2022. phylogatR:
Phylogeographic data aggregation and repurposing. Molecular Ecology Resources. 22:2830–
2843.

Pfeifer B., Wittelsbuerger U., Ramos-Onsins S.E., Lercher M.J. 2014. PopGenome: An Efficient Swiss
Army Knife for Population Genomic Analyses in R. Molecular Biology and Evolution. 31:1929–
1936.

Pinheiro J., Bates D., DebRoy S., Sarkar D., R Core Team. 2022. nlme: Linear and Nonlinear Mixed Effects
Models. R package version 3.1-155. https://CRAN.R-project.org/package=nlme. .

Powell R., Conant R., Collins J.T. 2016. Peterson Field Guide to Reptiles and Amphibians of Eastern and
Central North America 4th Edition. Boston, MA: Houghton Mifflin Harcourt.

Pyron R.A., Burbrink F.T. 2014. Early origin of viviparity and multiple reversions to oviparity in squamate
reptiles. Ecol Lett. 17:13–21.

Revell L.J. 2012. phytools: an R package for phylogenetic comparative biology (and other things).
Methods in Ecology and Evolution. 3:217–223.

Rohland N., Reich D. 2012. Cost-effective, high-throughput DNA sequencing libraries for multiplexed
target capture. Genome Res. 22:939–946.

Romiguier J., Gayral P., Ballenghien M., Bernard A., Cahais V., Chenuil A., Chiari Y., Dernat R., Duret L.,
Faivre N., Loire E., Lourenco J.M., Nabholz B., Roux C., Tsagkogeorga G., Weber A.A.T., Weinert
L.A., Belkhir K., Bierne N., Glémin S., Galtier N. 2014. Comparative population genomics in
animals uncovers the determinants of genetic diversity. Nature. 515:261–263.

Sanger F., Nicklen S., Coulson A.R. 1977. DNA sequencing with chain-terminating inhibitors. Proceedings
of the National Academy of Sciences. 74:5463–5467.

Schield D.R., Adams R.H., Card D.C., Corbin A.B., Jezkova T., Hales N.R., Meik J.M., Perry B.W., Spencer
C.L., Smith L.L., García G.C., Bouzid N.M., Strickland J.L., Parkinson C.L., Borja M., Castañeda-
Gaytán G., Bryson R.W., Flores-Villela O.A., Mackessy S.P., Castoe T.A. 2018. Cryptic genetic
diversity, population structure, and gene flow in the Mojave rattlesnake (Crotalus scutulatus).
Molecular Phylogenetics and Evolution. 127:669–681.

Sexton J.P., Hangartner S.B., Hoffmann A.A. 2014. Genetic isolation by environment or distance: Which
pattern of gene flow is most common? Evolution. 68:1–15.

Shafer A.B.A., Peart C.R., Tusso S., Maayan I., Brelsford A., Wheat C.W., Wolf J.B.W. 2017. Bioinformatic
processing of RAD-seq data dramatically impacts downstream population genetic inference.
Methods in Ecology and Evolution. 8:907–917.

Shine R. 2015. The evolution of oviparity in squamate reptiles: An adaptationist perspective. Journal of
Experimental Zoology Part B: Molecular and Developmental Evolution. 324:487–492.

Singhal S., Grundler M., Colli G., Rabosky D.L. 2017a. Squamate Conserved Loci (SqCL): A unified set of
conserved loci for phylogenomics and population genetics of squamate reptiles. Molecular
Ecology Resources. 17:e12–e24.

Singhal S., Huang H., Title P.O., Donnellan S.C., Holmes I., Rabosky D.L. 2017b. Genetic diversity is largely
unpredictable but scales with museum occurrences in a species-rich clade of Australian lizards.
Proceedings of the Royal Society B: Biological Sciences. 284.

Soltis D.E., Morris A.B., McLachlan J.S., Manos P.S., Soltis P.S. 2006. Comparative phylogeography of
unglaciated eastern North America. Molecular Ecology. 15:4261–4293.

Tonini J.F.R., Beard K.H., Ferreira R.B., Jetz W., Pyron R.A. 2016. Fully-sampled phylogenies of squamates
reveal evolutionary patterns in threat status. Biological Conservation. 204:23–31.

Vachon F., Whitehead H., Frasier T.R. 2018. What factors shape genetic diversity in cetaceans? Ecology
and Evolution. 8:1554–1572.

Waltari E., Hijmans R.J., Peterson a T., Nyári A.S., Perkins S.L., Guralnick R.P. 2007. Locating pleistocene
refugia: comparing phylogeographic and ecological niche model predictions. PloS ONE. 2:e563–
e563.

Weinell J.L., Austin C.C. 2017. Refugia and Speciation in North American Scarlet Snakes (*Cemophora*).
Journal of Herpetology. 51:161–171.

White E.P., Ernest S.K.M., Kerkhoff A.J., Enquist B.J. 2007. Relationships between body size and
abundance in ecology. Trends in Ecology and Evolution. 22:323–330.

Williams J.W., Shuman B.N., Webb T., Bartlein P.J., Leduc P.L. 2004. Late-Quaternary vegetation
dynamics in north America: Scaling from taxa to biomes. Ecological Monographs. 74:309–334.

Wright S. 1943. Isolation by Distance. Genetics. 28:114–138.

Zink R.M., Barrowclough G.F. 2008. Mitochondrial DNA under siege in avian phylogeography: nuDNA VS.
mtDNA in phylogeography. Molecular Ecology. 17:2107–2121.

 Table and Figure Captions

Table 1. Sample sizes for mtDNA and GBS datasets for each species.

Table 2. Model comparison of PGLS models under the Brownian Motion (BM) or Ornstein-

Uhlenbeck (OU) models of evolution.

Table 3. Model comparison of PGLS models with different sets of predictors. The top three

models for each response are shown, which all had a delta AIC value of <= 2. Note that none of

the models for mtDNA COI π were substantially improved over others (delta AIC all < 4.4). The

plus ("+") sign in the predictor columns indicates that predictor was included in that model.

Table 4. Model coefficients, standard errors (SE), t, and p-values for the top model listed in

Table 3 for each response.

Figure 1. Individual sample localities and nucleotide diversity estimates for species included in

the study. (a) Map of all samples for which GBS data were analyzed. Desert Southwest samples

are shown in turquoise and Coastal Plain species in brown. Dotted black boxes indicate the

inset maps to the right, which depict localities sampled within two species as examples. Shapes

indicate individuals were sampled from the same locality within a species, but do not

necessarily indicate the same locality across different species. (b) Phylogenetic relationships for

the 30 species included in the study based on Tonini et al. (2016). Within-individual nucleotide

estimates are shown to the right with colors and shapes corresponding to the maps in (a).

Figure 2. Lack of correlation between mtDNA and nuclear diversity estimates within species.

Spearman's rank correlation sample estimates and p-values are shown.

Figure 3. Predictors of mtDNA (a-d) and nuclear diversity (e-h) within species. Scatter plots and

dotted lines depict simple linear relationships without phylogenetic correction. a, b) Welch's

two-sample t-test statistics and p-values are shown. c, e, g) Spearman's rank correlation sample

estimates and p-values are shown. d, f, h) Phylogenetically-corrected R^2 values and p-values are

shown.

Table 1. Sample sizes for mtDNA and GBS datasets for each species.

Species	Family	# COI seqs	# cytb seqs	# GBS samples	# GBS loci (unique SNPs)	# GBS loci (seqs)
Ophisaurus ventralis	Anguidae	6	NA	4	22951	42875
Anolis carolinensis	Anolidae	16	NA	4	15708	28909
Agkistrodon conanti	Viperidae	5	46	4	5961	15540
Sistrurus miliarius	Viperidae	10	31	4	14356	23527
Crotalus ornatus	Viperidae	NA	27	4	9047	29251
Crotalus atrox	Viperidae	5	66	4	6926	19307
Crotalus scutulatus	Viperidae	4	74	4	14696	29502
Crotalus oreganus	Viperidae	NA	NA	4	11032	27615
Crotalus viridis	Viperidae	NA	8	4	15565	29627
Crotalus tigris	Viperidae	5	NA	4	9003	37903
Crotalus pyrrhus	Viperidae	6	NA	4	3274	6820
Crotalus adamanteus	Viperidae	10	126	4	8424	27387
Nerodia fasciata	Natricidae	30	143	4	15130	25148
Thamnophis marcianus	Natricidae	NA	22	4	7467	17648
Thamnophis sauritus	Natricidae	7	NA	4	14540	17945
Thamnophis sirtalis	Natricidae	7	37	3	15664	25409
Hypsiglena jani	Dipsadidae	NA	NA	4	19304	29950
Sonora episcopa	Colubridae	7	20	4	2923	12181
Trimorphodon vilkinsonii	Colubridae	NA	NA	3	5566	26644
Salvadora hexalepis	Colubridae	8	31	4	3513	20475
Masticophis flagellum	Colubridae	55	136	4	7512	18350
Coluber constrictor	Colubridae	51	22	4	18737	22267
Lampropeltis splendida	Colubridae	NA	25	4	6485	19095
Cemophora coccinea	Colubridae	69	9	4	15437	25907
Rhinocheilus lecontei	Colubridae	9	105	4	4713	15022
Arizona elegans	Colubridae	8	92	4	16879	37596
Pantherophis quadrivittatus	Colubridae	12	20	4	14352	26136
Pituophis catenifer	Colubridae	9	98	4	18205	32161
Pantherophis emoryi	Colubridae	NA	129	4	25086	44309
Pantherophis guttatus	Colubridae	11	11	4	14377	25517

789 Table 2. Model comparison of PGLS models under the Brownian Motion (BM) or Ornstein-

790 Uhlenbeck (OU) models of evolution.

Response	Model	log Likelihood	AIC	ΔAIC	Weight
mtDNA COI π	OU	57.452	-98.9	0.00	0.998
	BM	50.320	-86.6	12.26	0.002
mtDNA cytb π	OU	66.965	-117.9	0.00	0.997
	BM	60.257	-106.5	11.42	0.003
Within-species π	OU	157.485	-297.0	0.00	0.994
	BM	151.435	-286.9	10.1	0.006

793 Table 3. Model comparison of PGLS models with different sets of predictors. The top three

794 models for each response are shown, which all had a delta AIC value of < 2. Note that none of

795 the models for mtDNA COI π were substantially improved over others (delta AIC all < 4.4). The

796 plus ("+") sign in the predictor columns indicates that predictor was included in that model.

Response	df	log Like	AIC	ΔAIC	Weight	Range size	Body size	Clutch size	Parity
mtDNA COI π (OU models)	6	55.745	-99.5	0.00	0.129	+			+
	5	54.731	-99.5	0.03	0.127				+
	6	55.729	-99.5	0.03	0.127		+		+
mtDNA cytb π (OU models)	7	66.402	-118.8	0.00	0.379	+	+		+
	8	66.964	-117.9	0.88	0.245	+	+	+	+
	6	64.615	-117.2	1.58	0.172	+			+
Within-species π (OU models)	7	156.994	-300.0	0.00	0.342	+	+	+	
	6	155.531	-299.1	0.92	0.215	+		+	
	8	157.310	-298.6	1.37	0.172	+	+	+	+

799   Table 4. Model coefficients, standard errors (SE), t-, and p-values for the top model listed in

800   Table 3 for each response.

Response	Coefficient	Value	SE	t	p-value
mtDNA COI π (OU model)	Intercept	-0.0565	0.0662	-0.8540	0.4043
	log(Range size)	0.0062	0.0047	1.3184	0.2039
	Parity_vivip	-0.0165	0.0100	-1.6530	0.1157
	N sequences	-0.0001	0.0003	-0.2777	0.7844
mtDNA cytb π (OU model)	Intercept	-0.1522	0.0539	-2.8252	0.0117
	log(Range size)	0.0078	0.0033	2.3328	0.0322*
	log(Body size)	0.0315	0.0182	1.7320	0.1014
	Parity_vivip	-0.0179	0.0061	-2.9503	0.0090*
	N sequences	0.0001	0.0001	0.8222	0.4223
Within-species π (OU model)	Intercept	-0.0099	0.0041	-2.4112	0.0236
	log(Range size)	0.0010	0.0003	3.8158	0.0008*
	log(Body size)	-0.0022	0.0014	-1.6001	0.1221
	log(Clutch size)	0.0043	0.0017	2.5383	0.0177*
	N loci	0.0000	0.0000	0.0665	0.9475

Figure 1. Individual sample localities and nucleotide diversity estimates for species included in
the study. (a) Map of all samples for which GBS data were analyzed. Desert Southwest samples
are shown in turquoise and Coastal Plain samples in brown. Dotted black boxes indicate the
inset maps to the right, which depict localities sampled within two species as examples. Shapes
indicate individuals were sampled from the same locality within a species. (b) Phylogenetic
relationships for the 30 species included in the study based on Tonini et al. (2016). Within-
individual nucleotide estimates are shown to the right with colors and shapes corresponding to
the maps in (a).

Figure 2. Lack of correlation between mtDNA and nuclear diversity estimates within species.
Spearman's rank correlation sample estimates and p-values are shown.

820 **Figure 3.** Predictors of mtDNA (a-d) and nuclear diversity (e-h) within species. a, b) Welch's two-
821 sample t-test statistics and p-values are shown. c, e, g) Scatter plots and dotted lines depict
822 simple linear relationships without phylogenetic correction. Spearman's rank correlation
823 sample estimates and p-values are shown. d, f, h) Phylogenetically-corrected R^2 values and p-
824 values are shown.

Supplementary Materials

Table S1. Taxonomy used for this study and details about how different datasets were handled

to reflect recent taxonomic changes. References that were used to manually edit mitochondrial

alignments and range maps are provided.

[provided as Excel sheet – tab 1]

Table S2. List of all samples for which mtDNA COI was sequenced and GBS data were analyzed.

Sample names, museum catalogue numbers, geographic coordinates, locality information, and

total number of GBS reads sequenced are included.

[provided as Excel sheet – tab 2]

Table S3. Additional species-specific sequencing primers designed in this study to improve COI

sequencing results.

[provided as Excel sheet – tab 3]

Table S4. Data for 30 squamate species used as input for analyses.

[provided as Excel sheet – tab 4 and with PGLS R script in Dryad package]

Table S5. Data for 118 squamate individuals used to make figures and summaries.

[provided as Excel sheet – tab 5 and with R script in Dryad package]

Additional files on Dryad (https://doi.org/10.5061/dryad.qfttdz0ks): Mitochondrial (COI and

cytb) alignments including previous GenBank samples; GBS files; Range maps; Python/R scripts

Figure S1. (a) Diversity estimates for mtDNA genes COI and cytb are correlated. (b) Mean

within-individual pi estimates are strongly correlated with within-species pi estimates.

Figure S2. There are no clear differences in nuclear diversity estimates or the GBS datasets

collected from the two different regions.