Experimental determination and mathematical modeling of standard shapes of forming autophagosomes

Determination of standard shapes of phagophores

Yuji Sakai[1,2,3]*, Satoru Takahashi[1,4], Ikuko Koyama-Honda[1], Chieko Saito[1], and Noboru Mizushima[1]*

[1]Department of Biochemistry and Molecular Biology, Graduate School of Medicine, The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan

[2]Department of Biosystems Science, Institute for Life and Medical Sciences, Kyoto University, Sakyo-ku, Kyoto 606-8507, Japan

[3]Interdisciplinary Theoretical and Mathematical Sciences (iTHEMS) Program, RIKEN, Wako, Saitama 351-0198, Japan

[4]Department of Neurosurgery, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University, Bunkyo-ku, Tokyo 113-8510, Japan

*Correspondence to Yuji Sakai: yuji-sakai@m.u-tokyo.ac.jp; Noboru Mizushima: nmizu@m.u-tokyo.ac.jp.

Summary

The formation of autophagosomes involves dynamic morphological changes of membrane cisternae. Sakai et al. determined the average shapes of forming autophagosomes by statistically investigating three-dimensional electron micrographs and established a theoretical model that quantitatively reproduces the phagophore shapes.

Abstract

The formation of autophagosomes involves dynamic morphological changes of a phagophore from a disk-shaped membrane cisterna into a cup-shaped intermediate and a spherical autophagosome. However, the physical mechanism behind these morphological changes remains elusive. Here, we determined the average shapes of phagophores by statistically investigating three-dimensional electron micrographs of more than 100 phagophores. The results showed that the cup-shaped structures adopted a characteristic morphology; they were longitudinally elongated, and the rim was catenoidal with an outwardly recurved shape. To understand these characteristic shapes, we established a theoretical model of the shape of entire phagophores. The model quantitatively reproduced the average morphology and revealed that the characteristic shape of phagophores (i.e., an elongated shape with a catenoidal rim) was primarily determined by the relative size of the open rim to the total surface area. These results suggest that autophagosomal membranes are highly flexible and that the morphological changes during autophagosome formation follow a stable path determined by elastic bending energy minimization.

Introduction

Eukaryotic cells have membranous organelles with a variety of characteristic shapes. The autophagosome, which mediates macroautophagy (hereafter simply referred to as autophagy), is unique among the various organelles in that its generation involves characteristic membrane deformation. Autophagy is a bulk degradation process, through which a portion of the cytoplasm is sequestered into autophagosomes and is degraded upon fusion with lysosomes (Fig. 1) (Mizushima et al., 2011). In this process, a flat cisterna called a phagophore (also called an isolation membrane) grows from an initial disk through a cup-shaped intermediate and into a spherical stomatocyte. After the closure of the edge, it becomes a double-membrane autophagosome. Despite these characteristic morphological changes, there have been no systematic and quantitative studies to date on the standard morphology of phagophores during autophagosome formation. The morphology of phagophores is often approximated as a simple geometric shape, such as a part of a sphere or an ellipsoid, but the complexity and diversity of their morphology remain largely unexplored.

Quantitative and theoretical analyses can elucidate the mechanisms that control organelle morphology. Quantitative comparisons between experimentally observed and theoretically calculated shapes provide a mesoscopic, molecular-level explanation of the mechanisms that control organelle morphology (Lipowsky, 1991; Bassereau et al., 2018; Ramakrishnan et al., 2018). In theoretical analyses, organelles are often considered to be simple geometric shapes, such as spheres or tubes, and their complex and diverse morphologies, and thus the mechanisms behind them, are often ignored. Although theoretical research on autophagy is only in its early phase, the morphology of phagophores has been analyzed by approximating parts of a sphere or an ellipse (Knorr et al., 2012; Sakai et al., 2020). However, the actual morphology of phagophores observed in cells is more complex, and previous studies have oversimplified their morphology.

In the present study, we statistically and quantitatively investigated the morphology of phagophores. First, we determined the average morphology of more than 100 phagophores obtained from three-dimensional (3D) electron micrographs. The results showed that phagophores were elongated vertically, with a rim curved outward to form a catenoid-like shape. To understand the morphological characteristics, we developed a theoretical model based on the elastic bending energy. The morphology of the phagophore was considered to be in equilibrium at each step (Sakai et al., 2020). The physical parameter, Gaussian modulus, which determines the elastic properties of the phagophore membrane, was estimated from the experimental shape of the rim. The resulting theoretical model quantitatively reproduced the morphology during autophagosome formation observed by electron microscopy. These results suggest that autophagosomal membranes are highly

78 flexible and that the characteristic shapes of phagophores are primarily determined by
79 elastic bending energy minimization.

Results

Quantification of phagophore morphology from 3D-electron micrographs

To understand the standard shapes of phagophores, we determined the average morphology
of phagophores in mouse embryonic fibroblasts by array tomography, a technique that can
be used in 3D electron microscopy (Micheva and Smith, 2007). Images of more than 100–
200 serial sections (25-nm thickness) of a region containing several cells were captured
using a scanning electron microscope(Takahashi et al., 2022). We collected 117 entire 3D
structures of phagophores from 23 chemically fixed cells (Fig. 2A). Next, each phagophore
membrane was segmented and approximated as a point cloud $\{\vec{X}_{i,j}\}$ of 300-600 points,
where each point $\vec{X}_{i,j}$ represented the location of the j-th point on the membrane of the i-
th phagophore structure (Fig. 2B). In this electron microscopy experiment, cells were fixed
with reduced osmium tetroxide to keep the inner and outer membranes attached to each
other, which is their natural physiological state; if conventional osmium tetroxide had been
used, the space between the outer and inner membranes would have been artificially
expanded (Uemura et al., 2014). Thus, the inner and outer membranes were treated as a
single membrane in each point cloud representation. Each 3D morphology was constructed
by stacking a set of points obtained from a series of images (Fig. 2C).

To obtain an average shape of structures with different sizes, we normalized the size of
each 3D phagophore using standard z-score normalization since the morphologies were
almost size-independent (Fig. S1). In z-score normalization, each point $\vec{X}_{i,j}$ was converted
to a standardized point $\vec{Y}_{i,j}$ as

104
$$\vec{Y}_{i,j} = \frac{\vec{X}_{i,j} - \vec{\mu}_X}{\sigma_{X_i}}, \qquad \vec{\mu}_{X_i} = \frac{1}{N_i}\sum_{j=1}^{N_i}\vec{X}_{i,j}, \qquad \sigma_{X_i} = \sqrt{\frac{1}{N_i}\sum_{j=1}^{N_i}|\vec{X}_{i,j} - \vec{\mu}_{X_i}|^2}, \qquad (1)$$

where $\vec{\mu}_{X_i}$, σ_{X_i}, and N_i represent the mean, deviation, and number of the points belonging
to the i-th phagophore, respectively. σ_{X_i} means the structure size. Then, the normalized
point cloud $\{\vec{Y}_{i,j}\}$ was obtained.

Because phagophores in the electron microscopy images were oriented in various
directions, we rotated each structure to be oriented in a uniform direction (along the z-axis).

The average edge vector of a normalized point cloud, $\vec{\mu}_{e_i}$, was used as the orientation of
the i-th phagophore and was obtained as

$$\vec{\mu}_{e_i} = \frac{1}{M_i} \sum_{\substack{j \in edge}}^{M_i} \vec{Y}_{i,j}, \tag{2}$$

110 where the edge points belonging to the cloud were summed and M_i was the number of edge
111 points. From the average edge vector, the polar angle, θ_{e_i}, and azimuth angle, φ_{e_i}, of the
112 structure were determined. Each point $\vec{Y}_{i,j}$ was rotated according to

$$\vec{Z}_{i,j} = R_y(-\theta_{e_i})R_z(-\varphi_{e_i})\vec{Y}_{i,j}, \tag{3}$$

where R_y and R_z are the matrices that rotate $\vec{Y}_{i,j}$ by $-\theta_{e_i}$ about the y-axis and $-\varphi_{e_i}$ about
the z-axis, respectively, and then, a point cloud with uniform size and orientation, $\{\vec{Z}_{i,j}\}$,
was obtained. The surface was constructed from the normalized point cloud by Delaunay
triangulation (Fig. 2D and Video 1). Thus, we reconstructed the normalized shapes of
phagophores obtained from the 3D electron micrographs.

Because the obtained shapes of phagophores were diverse (reflecting the morphological
changes during autophagosome formation), we selected four representative categories
based on the rim size. The rim size σ_{e_i} was expressed as the magnitude of the deviation of
the normalized point cloud belonging to the rim as

$$\sigma_{e_i} = \sqrt{\frac{1}{M_i} \sum_{\substack{j \in edge}}^{M_i} \left| \vec{Y}_{i,j} - \vec{\mu}_{e_i} \right|^2}. \tag{4}$$

The four categories of observed shapes were as follows: disk ($\sigma_{e_i} > 1.1$), early-cup ($0.8 <$
$\sigma_{e_i} < 1.0$), middle-cup ($0.5 < \sigma_{e_i} < 0.7$), and late-cup ($\sigma_{e_i} < 0.4$) (examples of 3D
electron micrographs of each category are shown in Fig. S1 and S2). The point clouds in
each category were superposed (Fig. 3A–D, Video 2–5). The spread of the point clouds
indicated the variation within each morphology, while the regions with a high density of
points represented the modal morphology. Although individual phagophores were not
axisymmetric, probably owing to deformation caused by intracellular fluctuations, the
superimposed point clouds appeared to be axisymmetric around the z-axis.

Thus, these point clouds were projected onto the radial and axial planes (Fig. 3E-H). The
average morphology obtained suggested that the cup-shaped phagophores had two
characteristic features. First, they were elongated vertically along the rotation axis. This

result was consistent with our previous observations by fluorescence microscopy (Tsuboyama et al., 2016). Second, as the phagophore membrane approached the rim, it curved inward and then outward, forming a catenoid shape. The outward curvature was higher in late-cups. However, the rim appeared to be straight and not curved outward in disks. These characteristic features were also observed in individual phagophores in electron micrographs (Fig. 3I-L).

One methodological concern was that these characteristic shapes might have resulted from the chemical fixation of cells during sample preparation for electron microscopy. To rule out this possibility, we prepared samples by the high-pressure freezing and freeze-substitution method and subjected them to electron microscopy. The results showed similar morphological features of phagophores, such as elongated cups with catenoid-like rims (Fig. S3). Furthermore, we directly observed phagophores in live cells expressing GFP-LC3, a marker for phagophores and autophagosomes. Typical shapes of elongated phagophores with a catenoidal rim, which were consistent with the superimposed electron microscopy data, were frequently observed (Fig. 3M-P, Fig. S4). Thus, the structures observed by electron microscopy should represent the actual morphology of those in live cells.

To determine the average shape and quantify the curvature of the membrane, the projected point clouds were fitted with the z-axis symmetric polynomial function

$$x^2 = az^3 + bz^2 + cz + d, \tag{5}$$

where the fitting parameters are summarized in Table S1 (Fig. 4A-D). The standard error of each parameter in the three cup shapes was within a few percentage points, while the error of the disk shape was larger because of the small number of statistics and the large classification range ($\sigma_e > 1.1$). Each fitting curve well reproduced the point clouds that were experimentally obtained by electron microscopy. The fitted curves of middle- and late-cups clearly had a catenoidal rim (curved outward). The 3D surface in the axisymmetric approximation was represented using the fitting function

$$(f(v)\cos u, f(v)\sin u, v), \qquad f(v) = \sqrt{av^3 + bv^2 + cv + d}, \tag{6}$$

where u represents the azimuth angle. The total curvature J (indicated by the line color in Fig. 4A-D) on the surface at $z = v$ was obtained by

$$J(v) = J_m + J_p, \qquad J_m = -\frac{f''(v)}{[(f'(v))^2 + 1]^{3/2}}, \qquad J_p = \frac{1}{f(v)[(f'(v))^2 + 1]^{1/2}}, \tag{7}$$

where J_m and J_p represent the meridional (i.e., perpendicular) and parallel curvature relative to the xy-plane, respectively (Fig. 4E, line colors in Fig. 4F-I). In middle- and late-

cups, the parallel curvature, J_p, increased, with a positive sign as the surface approached

the rim from the bottom, while the meridional curvature, J_m, decreased, with a negative

sign. Furthermore, the meridional curvature decreased with the rim size. This trend was

also observed in individual phagophores (Fig. 4J), where the rim curvature was obtained

by fitting the point cloud of each structure with Eq. (6). Thus, the two orthogonal curvatures

J_m and J_p had opposite signs and seemed to cancel each other out, thus reducing the total

curvature J.

Modeling of phagophore morphology

Next, to investigate whether the morphological features of the phagophores obtained by

electron microscopy were reasonable, we conducted a mathematical model analysis. Each

shape of the phagophore membrane shown in Fig. 3 (i.e., disc, early-, middle-, and late-

cups) is stable for several minutes during autophagosome formation (Tsuboyama et al.,

2016; Sakai et al., 2020). While the rim length changes on this time scale, $\tau_{grow} \sim 1min$,

the mechanical relaxation time of the membrane is $\tau_{relax} \sim 1ms$ (Sakai et al., 2020).

Because $\tau_{relax} \ll \tau_{grow}$, it was considered that the rim length remained constant during

the mechanical relaxation and each morphology observed in electron microscopy was in

equilibrium with a fixed rim length. The membrane shapes at equilibrium were determined

from the elastic bending energy model. In the elastic bending energy model (Helfrich,

1973), the free energy F for a fixed rim length was given by

203
$$F = \int \left(\frac{\kappa_b}{2} J^2 + \kappa_G K \right) dA + \gamma_A A + \gamma_L L. \tag{8}$$

Here, the first term is the elastic bending energy with the total curvature, $J = J_m + J_p$, and

Gaussian curvature, $K = J_m J_p$, of the inner and outer membranes, which is scale-invariant.

The spontaneous curvature of the inner and outer membranes was assumed to be zero

because the inner and outer phagophore membranes contain relatively few proteins

(Fengsrud et al., 2000). It is considered that the phagophore membranes expand by

acquiring lipids from the endoplasmic reticulum through the lipid transfer protein ATG2

(Maeda et al., 2019; Osawa et al., 2019; Valverde et al., 2019), thereby keeping the lumen

(i.e., intermembrane space) volume small and the inner and outer membranes almost

parallel (Fig. 3I–L). Thus, the phagophore was assumed to have closely juxtaposed inner

and outer membranes with the same curvature and area, and the rim was approximated as

a closed line (Fig. 5A). The bending modulus is $\kappa_b = 20 k_B T$ with the Boltzmann constant

k_B and temperature T. The Gaussian modulus of elasticity, which was estimated from the

experimental morphology using the force balancing conditions acting on the rim at

equilibrium, $\kappa_G = -0.2\kappa_b$, was used (Fig. 5B). The smaller the Gauss modulus, the more

the rim curves outward (Fig. 5C-F). The second and third terms are the constraints for the

membrane area A and the rim length L, respectively, while γ_A and γ_L are the Lagrange

multipliers for fixing A and L, respectively. The second term is the rim line energy, which corresponds to the elastic bending energy of the rim. The rim length is in principle controlled by the abundance of curvature-inducing proteins that stabilize a highly curved rim (Sakai et al., 2020). The line energy includes the effects of the spontaneous curvature induced by these proteins (see Materials and methods). For an axisymmetric membrane, the free energy was rewritten as

$$F = 2\pi \int_0^{s_1} \left[\frac{\kappa_b}{2} x (J_m + J_p)^2 + \kappa_G x J_m J_p + \gamma_A x + \gamma_L \dot{x} \right] ds , \tag{9}$$

with

$$J_m = \dot{\theta}, \qquad J_p = \frac{\sin \theta}{x} , \tag{10}$$

where s, s_1 x, and θ are the arc length, location of the rim, radial coordinate, and tilt angle, respectively (Fig. 5A). The dots represented the derivative with respect to s. The membrane morphology was obtained by the variational method (Tu and Ou-Yang, 2003). A complete method for obtaining the morphology from the bending energy is described in the Materials and methods.

First, we determined the morphology of a stable cup-shaped membrane with several different rim radiuses (Fig. 6A-D). Because the free energy and the stable membrane morphology are scale-invariant, the rim radius was rescaled as

$$l = x_r / \sqrt{A/2\pi} , \tag{11}$$

with membrane area $A = 4\pi \int_0^{s_1} x ds$ and rim radius $x_r = \int_0^{s_1} \cos \theta \, ds$, where $l = 1$ represents a flat disk shape and $l = 0$ a closed sphere shape. Cups for which $0.2 < l < 0.6$ were not spherical but elongated vertically along the rotational axis. This trend was also observed in our previous models (Sakai et al., 2020). In these structures, the curvature in the elongated direction was reduced, thus lowering the bending energy (a cylinder [radius r, height h] had less bending energy than a spherical cap [radius r] of the same area when $r > h$). Furthermore, these structures had a catenoid-like rim. The catenoidal rims were attributed to the fact that the Gaussian modulus values of the phagophores were not particularly small. As the Gaussian modulus became even lower, the catenoid-like rim disappeared, and the theoretical model could no longer reproduce the experimental results (Fig. 5C-F). As the rim radius became smaller, the warping became stronger. These morphological features obtained by the theoretical model were consistent with those obtained by electron microscopy (Fig. 3). Note that γ_A was always negative (Fig. 6A-D), which indicates that membrane area growth decreased the energy of cup shapes [see Eq. (10)].

Next, we calculated the total curvature to quantitatively evaluate the shapes obtained by
the theoretical model (Fig. 6A-D). In all structures, from a disc to a cup, the total curvature
was reduced near the rim. The meridional curvature, J_m, and parallel curvature, J_p, were
also calculated (Fig. 6E-H). The meridional curvature was negative near the rim, reflecting
the outward curvature of the membrane, while the parallel curvature was positive. As a
result of the two competing curvatures, negative and positive, the total curvature near the
rim seemed to be smaller. Except for the rim, which was isotropic, the two curvatures were
of approximately the same magnitude.

We further investigated the change of curvatures during the morphological transition from
a disc to a cup (Fig. 6I). As the rim radius decreased, the parallel curvature J_p at the rim
increased monotonically because the curvature given by $J_p = \sin\theta / x$ was approximately
inversely proportional to the radial distance x. In the middle- and late-cup shapes, the
surface near the rim got closer to the rotational axis, and therefore the parallel curvature,
J_p, increased. On the other hand, as the rim radius decreased, the meridional curvature, J_m,
decreased (became more negative) in opposition to the parallel curvature, J_p, keeping the
total curvature ($J = J_m + J_p$) small. Thus, the two orthogonal curvatures canceled each
other out, making the rim catenoidal.

Comparison between theoretical and experimental morphologies

We compared the results of our theoretical calculations with our experimental data. The
point clouds obtained from electron micrographs (Fig. 3E-H) were superimposed on the
theoretically calculated morphologies with the same rim sizes (Fig. 7). At each step, the
experimental and theoretical results overlapped well. In particular, the longitudinally
elongated shape and the catenoidal rim were well captured by the theoretical calculations.
The individual structures of various sizes at each step were also compared with the
theoretical morphology (Fig. S1). Over a wide range of size scales, the theoretical model
reproduced the experimentally observed morphology well. The relative error of the
theoretical curves to the experimental fitting curves was within 30% across all shapes (Fig.
S5). The magnitude of this error was $\sim k_B T$ in energy, which was within the range of
thermal fluctuations. These results suggest that the morphological transition of
phagophores during autophagosome formation follows a stable path determined by the
elastic bending energy.

However, there was a slight discrepancy between the theoretical expectation and
experimental data, especially at the late-cup stage (Fig. 7). This discrepancy may be
partially explained by non-equilibrium fluctuations. The outer and rim membranes of the
phagophore often attach to the endoplasmic reticulum (Hayashi-Nishino et al., 2009; Ylä-

Anttila et al., 2009; Uemura et al., 2014), and the phagophore membrane may actively fluctuate like the endoplasmic reticulum membrane. Cytoskeletons may also contribute to autophagosome shaping (Mi et al., 2015; Kast et al., 2015). Another possibility is the influence of lipid asymmetry. During autophagosome formation, lipids are provided to the outer leaflet by the lipid transfer protein ATG2 (Maeda et al., 2019; Osawa et al., 2019; Valverde et al., 2019). Although it is hypothesized that the autophagosomal scramblase ATG9 resolves lipid asymmetry (Matoba et al., 2020), the remaining asymmetry may contribute to the elongation of phagophores (Seifert et al., 1991). However, these effects on the shape of the phagophore would be within the range of thermal fluctuation.

Discussion

The shape of phagophores has been empirically assumed, but their standard shapes have not yet been determined experimentally. In this study, we quantitatively analyzed the average morphology of phagophores using 3D electron microscopy data and found that phagophores are vertically elongated and that their rims have a catenoidal shape. These morphological features are also observed in cryo-electron tomography images of yeast phagophores (Bieber et al., 2022). We also showed that these characteristic features could be well explained by an elastic bending energy model. These results suggest that phagophore membranes are highly flexible and the morphological changes during autophagosome formation follow a stable path determined by elastic bending energy minimization. Our results also suggest that the shape of phagophores is primarily determined by the relative size of the open rim to the total surface area without any external influences. We do not rule out the possibility that some internal forces (e.g., lipid asymmetry) or external forces (e.g., exerted by the endoplasmic reticulum or cytoskeletons) influence the phagophore morphology, but these effects would be relatively minor.

It should be noted that the shape of phagophores is scale-independent (Fig. S1). Over a wide range of size scales, the theoretical model reproduced the experimentally observed morphology well. Generally, the bending energy of membranes [e.g., the outer and inner membranes of phagophores, the first term of Eq. (8)] is scale-independent (Helfrich, 1973), whereas that of the rim [i.e., the line energy, the second term of Eq. (8)] is scale (length)-dependent. Our result that the shape of phagophores is scale-independent suggests that the length of the rim is determined by a mechanism other than the simple bending energy model, probably by stabilization by curvature generators accumulating on the rim (Sakai et al., 2020).

The rim length is considered to be constant on the time scale of mechanical relaxation as we assumed in this study, but it changes with phagophore growth on larger time scales.

The change in the rim length could be explained by a change in the abundance of curvature
generators at the rim that stabilizes the highly curved rim (Sakai et al., 2020).

A recent cryo-electron tomography study in yeast shows that the intermembrane space
between the inner and outer membranes is not constant, with slight swelling at the rim
(Bieber et al., 2022). Although the rim swelling would be important in determining the
bending elasticity at the rim, the qualitative results of this study for overall morphology are
expected to remain the same. The model also indicates that the surface energy [$\gamma_A A$ in Eq.
(8)] is negative ($\gamma_A < 0$ in Fig. 6A-D), predicting that a larger membrane area is
energetically more stable for the cup shapes. This stability could be one of the mechanisms
underlying the directional lipid transfer from the endoplasmic reticulum to phagophores
during phagophore growth.

Materials and methods

Cell culture

Mouse embryonic fibroblasts (MEFs) were cultured in Dulbecco's modified Eagle's
medium (DMEM: D6546, Sigma-Aldrich) supplemented with 10% fetal bovine serum
(172012, Sigma-Aldrich), and 2 mM L-glutamine (25030-081, Gibco) in a 5% CO_2
incubator. For the starvation treatment, cells were washed twice and incubated in amino
acid-free DMEM (048-33575, Wako Pure Chemical Industries) without serum. MEFs that
stably express GFP-LC3 were generated previously (Itakura and Mizushima, 2010).

Three-dimensional electron microscopy of phagophores using array tomography

For chemical fixation, a glass base dish was coated with carbon using a vacuum evaporator
(IB-29510VET, JEOL) and then coated with 0.1% gelatin and dried under UV irradiation,
as described previously (Maeda et al., 2019). Cells were cultured in the pretreated glass
base dish for 2 days and starved for 30 min just before fixation. The cells were first fixed
with 2.5% glutaraldehyde (G018/1, TAAB) in 0.1 M sodium cacodylate buffer, pH 7.4
(37237-35, Nacalai Tesque), overnight at 4ºC and then with 1% osmium tetroxide (3020-
4, Nisshin EM) and 1.5% potassium ferrocyanide (161-03742, Wako) in 0.065 M sodium
cacodylate buffer for 2 h at 4ºC. The fixed cells were washed five times with Milli-Q water,
block-stained with 3% uranium acetate for 1 h, and dehydrated with an ethanol series. After
a 1-h dehydration in 100% ethanol, the cells were embedded in Epon (EPON812, TAAB),
polymerized at 40ºC for 12 h, and then polymerized at 60ºC for 48 h. Glass bases were
removed from blocks by soaking them in liquid nitrogen, and then, the top surface of the
blocks was trimmed to 100 × 100 mm using razor blades.

For rapid freeze fixation, we used the high-pressure freezing and freeze substitution method.
A sapphire glass (EM ICE accessory, cat. number 10702766; Leica Microsystems, Wetzlar,
Germany) was cleaned with ethanol, coated with carbon by using a vacuum evaporator
(IB-29510VET), heat treated (60ºC for >48 h), coated with poly-L-lysin, and sterilized.
Cells were cultured on the pretreated sapphire glass for 2 days and starved for 30–45 min
before freeze fixation. The sapphire glass was then mounted between two aluminum
carriers (EM ICE accessory, type A, cat. number 10660141). The space between the
sapphire glass and carrier was filled with 3% percoll solution to prevent the formation of
ice crystals. After freezing in a high-pressure freezer (HPM010; Bal-Tec, Balzers,
Liechtenstein), each sapphire disk was detached from an aluminum carrier in liquid
nitrogen and quickly transferred to a tube for freeze substitution. In the tube, acetone was
prepared containing 0.1% uranium acetate, 2% osmium tetroxide, and 3% water and then
frozen with liquid nitrogen. Freeze substitution was carried out for 5 days at −80ºC, and
then, the temperature was gradually raised, to −30ºC for 1 h and 40 min, to 4ºC for 1 h and
10 min, and to room temperature for 1 h. After being washed with dry acetone three times,
the sapphire disks were collected and embedded in EPON812, with a gradually increasing
EPON812 concentration, as follows: a 2:1 ratio of acetone:EPON812 overnight, a 1:1 ratio
of acetone:EPON812 for 1 day, and 100% EPON 812 for 6 h. Then, the sapphire disks
were placed in a silicon mold and polymerized. After polymerization, the sapphire disks
were removed with a razor and by soaking in liquid nitrogen repeatedly.

To cut ultra-thin serial sections, we used a diamond knife with an ultra-jumbo boat (Ultra
Jumbo 45 degree; Diatome, Hatfield, PA) mounted on an ultramicrotome (UC7; Leica).
For chemical fixation, 25-nm-thick sections were cut, creating a ribbon of 130 serial
sections, and for HPF fixation, 50-nm-thick sections were cut, creating a ribbon of 55 serial
sections. The ribbons were collected onto a silicon wafer that was held and manipulated by
an MM33 micromanipulator (Märzhäuser Wetzlar, Wetzlar, Germany). Electron
microscopic images were acquired using a scanning electron microscope (JSM7900F;
JEOL, Tokyo, Japan) as described previously (Maeda et al. 2020).

In the 3D reconstruction, the scale factors in the stack direction were adjusted assuming
axisymmetry of phagophores as described previously (Hayashi-Nishino et al. 2009, Ylä-
Anttila et al. 2009) in order to compensate for the compression of the structures during
sample preparation (i.e., fixation and sectioning) for electron microscopy. For 3D
reconstruction, the z-axis was roughly aligned in a large area (~10 × 10 mm) around
autophagosomes (including other subcellular organelles) by StackerNEO (System in
Frontier Inc., Tokyo, Japan), and serial images of each autophagosome were extracted.
Next, the serial images were aligned by StackerNEO. For the segmentation of
autophagosomes, their shapes were traced manually by Plot Digitizer.

Fluorescence microscopy

MEFs stably expressing GFP-LC3B were subjected to live-cell fluorescence imaging using a SpinSR10 spinning-disk confocal microscope (Olympus) equipped with an ORCA-Flash 4.0 camera (Hamamatsu), a UPLAPO OHR 60× lens (NA 1.50, Olympus), a 3.2× optical zoom lens, and a SORA disk in place (Yokogawa). The microscope was operated with cellSens Dimension v2.3 (Olympus). During live-cell imaging, a dish was mounted in a chamber (INUB-ONI-F2, TOKAI HIT) to maintain the incubation conditions at 37ºC and 5% CO_2. Five slices with a thickness of 250 nm were acquired. The images were processed with cellSens Dimension software using the following settings: deconvolution with advanced maximum likelihood algorithm, 5× interaction, and adaptive PSF filter settings.

Elastic bending energy and shape equations of membranes

For an axisymmetric geometry, the free energy of the membrane was given by

$$F = 2\pi \int_0^{s_1} \left[\kappa_b x \left(\dot{\theta} + \frac{\sin\theta}{x} \right)^2 + 2\kappa_G \dot{\theta} \sin\theta + 2\gamma_A x + \gamma_L \dot{x} \right] ds, \tag{12}$$

where s, x, and θ are the arc length, radial coordinate, and tilt angle, respectively (Fig. 5A) (Tu and Ou-Yang, 2003); s_1 represents the location of the rim; κ_b and κ_G are the bending elastic moduli; and γ_A and γ_L are the Lagrange multipliers for the area A and the rim length L, respectively. The free energy should be minimized with the constraints $\dot{x} = \cos\theta$, and thus the action $S = 2\pi \int_0^{s_1} \mathcal{L} ds$ was constructed with the Lagrangian

$$\mathcal{L} = \kappa_b x \left(\dot{\theta} + \frac{\sin\theta}{x} \right)^2 + 2\kappa_G \dot{\theta} \sin\theta + 2\gamma_A x + \gamma_L \dot{x} + \gamma_x (\dot{x} - \cos\theta), \tag{13}$$

where γ_x is the Lagrange multiplier for $\dot{x}$. Based on variational theory (Tu and Ou-Yang, 2003), the shape equation to determine the membrane shape with the minimum free energy is given by

$$\ddot{\theta} = -\frac{\tan\theta}{2} \dot{\theta}^2 - \frac{\dot{\theta}\cos\theta}{x} + \frac{\sin 2\theta}{2x^2} + \gamma_L \tan\theta + \frac{\tan\theta}{2} \left(\frac{\sin\theta}{x} \right)^2, \tag{14}$$

with the boundary conditions at the rim ($s = s_1$)

$$\left[\dot{\theta} + \left(1 + \frac{\kappa_G}{\kappa_b} \right) \frac{\sin\theta}{x} \right]_{s=s_1} = 0, \tag{15}$$

$$\left[-2 \left(1 + \frac{\kappa_G}{2\kappa_b} \right) \kappa_G \left(\frac{\sin\theta}{x} \right)^2 + 2\gamma_L + \gamma_A \frac{\cos\theta}{x} \right]_{s=s_1} = 0. \tag{16}$$

These boundary conditions at the rim indicate that Gaussian curvature, line tension, and
surface tension are obtained from the curvature at the rim. The initial conditions at $s = 0$
for x and θ are $x(0) = \theta(0) = 0$.

By solving the shape equation with the boundary conditions, a stable shape is obtained for
a membrane area $A = 4\pi \int_0^{s_1} x ds$ and rim radius $x_r = \int_0^{s_1} \cos \theta \, ds$, with the
corresponding Lagrange multipliers γ_A and γ_L, as shown in in Fig. 6A-D.

To compare theoretical expectations with the empirical data shown in Fig. 7, the computed
shapes were normalized in the same way as in the experiment. That is, we constructed a
point cloud of axisymmetric 3D shapes from the 2D shapes obtained in the calculation and
normalized them using z-score normalization [see Eq. (1)-(4)].

Relation between line energy and bending energy at the rim

The elastic bending energy of the rim was determined by

$$F_r = \frac{\kappa_b}{2} \int (J_r - \bar{J}_r)^2 dA_r \tag{17}$$

417 where κ_b is the bending rigidity and $J_r, \bar{J}_r$, and A_r are the total curvature, spontaneous
418 curvature, and area, respectively, of the rim. The rim geometry was modeled as a part of a
419 torus with an intermembrane radius r (Fig. 5A). In the torus approximation, the total
420 curvature and the surface element of the rim were respectively given by

$$J_r = \frac{1}{r} - \frac{\cos \varphi}{l - r \cos \varphi}, \qquad dA_r = 2\pi r(l - r \cos \varphi) d\varphi, \tag{18}$$

422 where l is the rim radius (Sakai et al., 2020). The integration variable φ spans the interval
423 $[\theta - \pi/2, \theta + \pi/2]$, with the tilt angle θ at the rim (Fig. 5A). The intermembrane distance
424 of phagophores was very small, and $l \gg r$ was assumed. In this case, the total curvature
425 and the surface element became $J_r = \frac{1}{r}$ and $dA_r = 2\pi r l d\varphi$, respectively, and then the
426 bending energy became

$$F_r = \frac{\kappa_b}{2} \int_{\alpha-\pi/2}^{\alpha+\pi/2} \left(\frac{1}{r} - \bar{J}_r\right)^2 2\pi r l d\varphi = \kappa_b \pi^2 r \left(\frac{1}{r} - \bar{J}_r\right)^2 l = \gamma_L L, \tag{19}$$

428 with the rim length $L = 2\pi l$ and the line tension $\gamma_L = \frac{\kappa_b}{2} \pi r \left(\frac{1}{r} - \bar{J}_r\right)^2$. Therefore, the
429 bending energy of the rim was converted into the line energy, which included the effects
430 of the spontaneous curvature. The spontaneous curvature is induced by the curvature

generating proteins, and the rim length is controlled by the abundance of curvature-generating proteins (Sakai et al., 2020).

Estimation of the Gaussian modulus with phagophore growth

According to the elastic bending energy model interpreted in terms of stresses (Evans and Skalak, 2018), the membrane can develop a meridional bending moment, a force that resists bending, $M_m = \kappa_b J + \kappa_G J_p$; the first and second terms in this equation are associated with the total curvature, $J = J_m + J_p$, and the Gaussian curvature, $K = J_m J_p$, respectively (Fig. 5A), while κ_b and κ_G represent the bending and the Gaussian moduli, respectively, which characterize the elastic properties of the membrane. As shown in the electron micrographs (Fig. 3 and S1, S2), the phagophore had closely juxtaposed inner and outer membranes (*a* and *b* in Fig. 5A) with the same curvatures, and the rim can be approximated as a line (*c* in Fig. 5A). The spontaneous curvature was assumed to be zero because the outer and inner membranes contain relatively few proteins (Fengsrud et al., 2000). Because the phagophore rim was not exposed to any external rotational moment, the total moment should be zero, that is, $M_m = 0$. From this moment balancing condition, the Gaussian modulus can be obtained as

$$\frac{\kappa_G}{\kappa_b} = -\frac{J}{J_p}. \tag{20}$$

Therefore, the Gaussian modulus, κ_G, is determined by fitting the experimental phagophore membrane shapes (Fig. 4), where the curvatures are obtained from Eq. (7).

The Gaussian modulus obtained by fitting the experimental phagophore membrane shapes with a polynomial function was $-(0.21 \pm 0.05)\kappa_b$, with no significant differences among the disc, early-, middle-, and late-cups (Fig. 5B). The deviation σ_X defined in Eq. (1), which represents the spatial extent of the point clouds of each morphology (Fig. 3), was used as a unit of length (Fig. S1). The obtained values of the phagophore membrane were within the reported range of biological membranes, namely, $-\kappa_b < \kappa_G < 0$ (Siegel and Kozlov, 2004; Hu et al., 2012).

Data Availability

The numerical code in the mathematical model is publicly available on GitHub at https://github.com/YSakaiGH/autophagosome_shapes.

Online supplemental material

Fig. S1 shows the comparison of individual shapes of phagophores of different sizes obtained experimentally with the theoretical model. Fig. S2 shows examples of serial electron microscopy images of phagophores. Fig. S3 shows high-pressure freezing and freeze substitution electron microscope images of phagophores. Fig. S4 shows examples of in vivo fluorescent images of phagophores. Fig. S5 shows quantitative comparison of the shapes between experimental and theoretical results. Video 1 shows an example of a rotating phagophore structure, corresponding to Fig. 2*D*. Video 2–5 show the rotating view of the superposed morphology of phagophores at the disk, early-cup, middle-cup, and late-cup, respectively, corresponding to Fig. 3*A-D*.

Acknowledgments

We would like to thank Michael M. Kozlov and Masashi Tachikawa for useful discussions. This work was supported by a Grant-in-Aid for Scientific Research (KAKENHI) from JSPS (20K06482 to Y.S., 22H04635 to I. K-.H., and 22H04919 to N.M.) and Exploratory Research for Advanced Technology (ERATO; grant number JPMJER1702) from the Japan Science and Technology Agency (JST) (to N.M.). The numerical computations have been performed with the RIKEN supercomputer system (HOKUSAI).

Author Contributions:

Conceptualization, Y.S. and N.M.; Methodology, Y.S.; Investigation, Y.S., S.T., C.S., and I.K-.H.; Formal analysis, Y.S.; Writing, Y.S., I.K-.H., and N.M.; Visualization, Y.S., S.T., C.S., and I.K-.H.; Funding acquisition, Y.S., I.K-.H. and N.M.; Resources, Y.S., S.T., C.S., and I.K-.H.; Supervision, Y.S. and N.M.

Competing Interest Statement: The authors declare no competing interests.

References

Bassereau, P., R. Jin, T. Baumgart, M. Deserno, R. Dimova, V.A. Frolov, P. v. Bashkirov, H. Grubmüller, R. Jahn, H.J. Risselada, L. Johannes, M.M. Kozlov, R. Lipowsky, T.J. Pucadyil, W.F. Zeno, J.C. Stachowiak, D. Stamou, A. Breuer, L. Lauritsen, C. Simon, C. Sykes, G.A. Voth, and T.R. Weikl. 2018. The 2018 biomembrane curvature and remodeling roadmap. *J Phys D Appl Phys*. 51:343001. doi:10.1088/1361-6463/aacb98.

Bieber, A., C. Capitanio, P.S. Erdmann, F. Fiedler, F. Beck, C.-W. Lee, D. Li, G. Hummer, B.A. Schulman, W. Baumeister, and F. Wilfling. 2022. In situ structural analysis reveals membrane shape transitions during autophagosome formation. *Proceedings of the National Academy of Sciences*. 119. doi:10.1073/pnas.2209823119.

Evans, E.A., and R. Skalak. 2018. Mechanics and Thermodynamics of Biomembranes. CRC Press. 1–254 pp.

Fengsrud, M., E.S. Erichsen, T.O. Berg, C. Raiborg, and P.O. Seglen. 2000. Ultrastructural
characterization of the delimiting membranes of isolated autophagosomes and amphisomes by
freeze-fracture electron microscopy. *Eur J Cell Biol*. 79. doi:10.1078/0171-9335-00125.

Hayashi-Nishino, M., N. Fujita, T. Noda, A. Yamaguchi, T. Yoshimori, and A. Yamamoto. 2009.
A subdomain of the endoplasmic reticulum forms a cradle for autophagosome formation. *Nat
Cell Biol*. 11:1433–1437. doi:10.1038/ncb1991.

Helfrich, W. 1973. Elastic properties of lipid bilayers: theory and possible experiments. *Z
Naturforsch C*. 28:693–703. doi:10.1515/znc-1973-11-1209.

Hu, M., J.J. Briguglio, and M. Deserno. 2012. Determining the Gaussian curvature modulus of
lipid membranes in simulations. *Biophys J*. 102:1403–10. doi:10.1016/j.bpj.2012.02.013.

Itakura, E., and N. Mizushima. 2010. Characterization of autophagosome formation site by a
hierarchical analysis of mammalian Atg proteins. *Autophagy*. 6:764–776.
doi:10.4161/auto.6.6.12709.

Kast, D.J., A.L. Zajac, E.L.F. Holzbaur, E.M. Ostap, and R. Dominguez. 2015. WHAMM Directs
the Arp2/3 Complex to the ER for Autophagosome Biogenesis through an Actin Comet Tail
Mechanism. *Current Biology*. 25:1791–1797. doi:10.1016/j.cub.2015.05.042.

Knorr, R.L., R. Dimova, and R. Lipowsky. 2012. Curvature of Double-Membrane Organelles
Generated by Changes in Membrane Size and Composition. *PLoS One*. 7:e32753.
doi:10.1371/journal.pone.0032753.

Lipowsky, R. 1991. The conformation of membranes. *Nature*. 349:475–481.
doi:10.1038/349475a0.

Maeda, S., C. Otomo, and T. Otomo. 2019. The autophagic membrane tether ATG2A transfers
lipids between membranes. *Elife*. 8. doi:10.7554/eLife.45777.

Matoba, K., T. Kotani, A. Tsutsumi, T. Tsuji, T. Mori, D. Noshiro, Y. Sugita, N. Nomura, S.
Iwata, Y. Ohsumi, T. Fujimoto, H. Nakatogawa, M. Kikkawa, and N.N. Noda. 2020. Atg9 is a
lipid scramblase that mediates autophagosomal membrane expansion. *Nat Struct Mol Biol*.
27:1185–1193. doi:10.1038/s41594-020-00518-w.

Mi, N., Y. Chen, S. Wang, M. Chen, M. Zhao, G. Yang, M. Ma, Q. Su, S. Luo, J. Shi, J. Xu, Q.
Guo, N. Gao, Y. Sun, Z. Chen, and L. Yu. 2015. CapZ regulates autophagosomal membrane
shaping by promoting actin assembly inside the isolation membrane. *Nat Cell Biol*. 17:1112–
1123. doi:10.1038/ncb3215.

Micheva, K.D., and S.J. Smith. 2007. Array Tomography: A New Tool for Imaging the Molecular
Architecture and Ultrastructure of Neural Circuits. *Neuron*. 55:25–36.
doi:10.1016/j.neuron.2007.06.014.

Mizushima, N., T. Yoshimori, and Y. Ohsumi. 2011. The Role of Atg Proteins in Autophagosome
Formation. *Annu Rev Cell Dev Biol*. 27:107–132. doi:10.1146/annurev-cellbio-092910-154005.

Osawa, T., T. Kotani, T. Kawaoka, E. Hirata, K. Suzuki, H. Nakatogawa, Y. Ohsumi, and N.N.
Noda. 2019. Atg2 mediates direct lipid transfer between membranes for autophagosome
formation. *Nat Struct Mol Biol*. 26:281–288. doi:10.1038/s41594-019-0203-4.

Ramakrishnan, N., R.P. Bradley, R.W. Tourdot, and R. Radhakrishnan. 2018. Biophysics of
membrane curvature remodeling at molecular and mesoscopic lengthscales. *Journal of Physics:*
Condensed Matter. 30:273001. doi:10.1088/1361-648X/aac702.

Sakai, Y., I. Koyama-Honda, M. Tachikawa, R.L. Knorr, and N. Mizushima. 2020. Modeling
Membrane Morphological Change during Autophagosome Formation. *iScience*. 23:101466.
doi:10.1016/j.isci.2020.101466.

Seifert, U., K. Berndl, and R. Lipowsky. 1991. Shape transformations of vesicles: Phase diagram
for spontaneous- curvature and bilayer-coupling models. *Phys Rev A (Coll Park)*. 44:1182–
1202. doi:10.1103/PhysRevA.44.1182.

Siegel, D.P., and M.M. Kozlov. 2004. The Gaussian Curvature Elastic Modulus of N-
Monomethylated Dioleoylphosphatidylethanolamine: Relevance to Membrane Fusion and
Lipid Phase Behavior. *Biophys J*. 87:366–374. doi:10.1529/biophysj.104.040782.

Takahashi, S., C. Saito, I. Koyama-Honda, and N. Mizushima. 2022. Quantitative 3D correlative
light and electron microscopy of organelle association during autophagy. *Cell Struct Funct*.
47:22071. doi:10.1247/csf.22071.

Tsuboyama, K., I. Koyama-Honda, Y. Sakamaki, M. Koike, H. Morishita, and N. Mizushima.
2016. The ATG conjugation systems are important for degradation of the inner autophagosomal
membrane. *Science (1979)*. 354:1036–1041. doi:10.1126/science.aaf6136.

Tu, Z.C., and Z.C. Ou-Yang. 2003. Lipid membranes with free edges. *Phys Rev E*. 68:061915.
doi:10.1103/PhysRevE.68.061915.

Uemura, T., M. Yamamoto, A. Kametaka, Y. Sou, A. Yabashi, A. Yamada, H. Annoh, S.
Kametaka, M. Komatsu, and S. Waguri. 2014. A Cluster of Thin Tubular Structures Mediates
Transformation of the Endoplasmic Reticulum to Autophagic Isolation Membrane. *Mol Cell*
Biol. 34:1695–1706. doi:10.1128/MCB.01327-13.

Valverde, D.P., S. Yu, V. Boggavarapu, N. Kumar, J.A. Lees, T. Walz, K.M. Reinisch, and T.J.
Melia. 2019. ATG2 transports lipids to promote autophagosome biogenesis. *Journal of Cell*
Biology. 218:1787–1798. doi:10.1083/jcb.201811139.

Ylä-Anttila, P., H. Vihinen, E. Jokitalo, and E.-L. Eskelinen. 2009. 3D tomography reveals
connections between the phagophore and endoplasmic reticulum. *Autophagy*. 5:1180–1185.
doi:10.4161/auto.5.8.10274.

Figures

Figure 1. Schematic representation of autophagosome formation.

Each shape represents an axially symmetric membrane structure. As surface area increases,
the shape of a forming autophagosome (phagophore) changes from an initial disk, through
a cup-shaped intermediate, to a spherical closed autophagosome. The regions labeled a, b,
and c correspond to the inner membrane, the outer membrane, and the rim, respectively.
Eventually, autophagosomes and lysosomes fuse to degrade the contents of the former.

Figure 2. Three-dimensional electron microscopy of phagophores.

(**A**) Serial electron microscopy images of a phagophore. (**B**) Left, an electron microscopy section image; the red dots represent the contour of the phagophore. Right, one segment of a point cloud of the extracted contour; the two red dots represent the edges of the phagophore. (**C**) 3D reconstruction of a point cloud. (**D**) A normalized shape of a phagophore, in which the deviation of the point cloud was normalized to one and rotated so that the direction of the edge aligns with the z-axis. Each structure is shown rotated 90 degrees about the z-axis. Rotating views (the z-axis is the rotation axis) are shown in Video 1. The surface was constructed by Delaunay triangulation, in which the open edge of the cup (the area surrounded by the red line) was also considered as part of the surface.

Figure 3. Average shape of phagophores.

(**A–D**) Three-dimensional point clouds of disks (**A**), early-cups (**B**), middle-cups (**C**), and late-cups (**D**) are shown. Point clouds were obtained from electron microscopy images of 15 disks (3,491 points), 16 early-cups (4,585 points), 24 middle-cups (8,578 points), and 45 late-cups (17,692 points) and then superposed. Rotating views (the z-axis is the rotation axis) are shown in Video 2–5. (**E–H**) The projected point clouds of disks (**E**), early-cups (**F**), middle-cups (**G**), and late-cups (**H**) onto the radial (x) and axial (y) plane. The unit of length was normalized so that the variance of the point set was one. (**I–L**) Examples of electron microscopy images of phagophores at the disk (**I**), early-cup (**J**), middle-cup (**K**), and late-cup (**L**) stages. The red arrows indicate phagophores; scale bar, 1 μm. (**M–P**) Examples of live-cell imaging of GFP-LC3 showing phagophores at the disk (**M**), early-cup (**N**), middle-cup (**O**), and late-cup (**P**) stages; scale bar, 1 μm.

Figure 4. Shapes and curvatures fitted with approximate curves.

(**A–D**) The shapes fitted with polynomial functions of a disk (**A**), an early-cup (**B**), a middle-cup (**C**), and a late-cup (**D**), in which the line color represents the total curvature. In each panel, the black dots represent the same point set as in Fig. 3*E–H*, respectively. The unit of length was normalized so that the variance of the point set was one. (**E**) A schematic description of the meridional and parallel curvatures relative to the *xy*-plane, J_m and J_p, respectively. (**F–I**) The meridional (J_m, $x \geq 0$, right) and parallel (J_p, $x \leq 0$, left) curvatures of a disk (**F**), an early-cup (**G**), a middle-cup (**H**), and a late-cup (**I**) are shown by the various line colors; the shape of the fitting curve in each panel is the same as those in panels **A–D**. (**J**) A scatter plot of the meridional outward curvature near the rim obtained by polynomial fitting of individual phagophores.

Figure 5. Model description of phagophore and Gaussian modulus.

(**A**) A schematic description of the phagophore. a and b represent the inner and outer membranes, respectively, which are approximated by two paralleled axisymmetric membranes. c represents the rim which is approximated as a line. z is the axis of symmetry, s is the length along the contour measured from the origin, and θ is the angle between the tangent to the contour and the x-axis. M_m is the meridional moments. J_m and J_p are the two principal curvatures. J and K are the total and Gaussian curvatures, respectively, with $J > 0, K > 0$ for convex surfaces and $J \approx 0, K < 0$ for catenoidal surfaces. (**B**) Gaussian modulus obtained by fitting the experimental shapes, where $\kappa_b = 20k_BT$. The membrane area, $A = 4\pi \int f\sqrt{1 + (f')^2}dx$, is also obtained from the polynomial fitting. (**C**-**F**) Membrane shapes obtained from the bending energy with different values of the Gaussian modulus, $\kappa_G = 0$ (red), $-0.5\kappa_b$ (green), and $-\kappa_b$ (blue), for different values of the rim radius, $l = 0.8$ (**C**), $l = 0.6$ (**D**), $l = 0.4$ (**E**), and $l = 0.2$ (**F**). The unit of length was non-dimensionalized by the length, $\sqrt{A/2\pi}$.

Figure 6. Shapes and curvatures obtained from theoretical calculations.

(**A–D**) Membrane shapes obtained from the bending energy with Gaussian modulus $\kappa_G = -0.2\kappa_b$ for several values of the rim radius $l = 0.8$ (**A**), $l = 0.6$ (**B**), $l = 0.4$ (**C**), and $l = 0.2$ (**D**), where the line color represents the total curvature. (**E–H**) The meridional and parallel curvatures J_m and J_p are shown in the right ($x \geq 0$) and left ($x \leq 0$) regions, respectively. The same shapes as Panel **A–D** were used. (**I**) The meridional, parallel, and total curvatures, J_m (red), J_p (blue), and J (green), respectively, at the rim as a function of the rim radius, l. The rim radii corresponding to Panel **A–D** are indicated. The unit of length was non-dimensionalized by the length $\sqrt{A/2\pi}$.

Figure 7. Comparison of theoretical and experimental shapes.

The theoretically calculated shapes (black lines) were superimposed over the experimentally obtained point clouds (red) of disks (**A**), early-cups (**B**), middle-cups (**C**), and late-cups (**D**) [using the same data as shown in Fig. 3 (**E**–**H**)]. The theoretical results were normalized so that the mean of the point cloud representing each shape was zero and the variance was one, in the same way as the experimental results were normalized. The rim size σ_{e_i} was fixed at $\sigma_{e_i} = 1.45$ (**A**), 0.75 (**B**), 0.48 (**C**), and 0.28 (**D**), corresponding to $l = 0.8$, 0.39, 0.24, and 0.13, respectively.