Temporal progression along discrete coding states during decision-making in the mouse gustatory cortex

Liam Lang[1,2,3], Giancarlo La Camera[1,2,3,*], and Alfredo Fontanini[1,2,3,*]

[1]Graduate Program in Neuroscience, [2]Center for Neural Circuit Dynamics, and [3]Department of Neurobiology and Behavior

Stony Brook University

*Equal contribution and co-corresponding authors:

giancarlo.lacamera@stonybrook.edu, alfredo.fontanini@stonybrook.edu

The mouse gustatory cortex (GC) is involved in taste-guided decision-making in addition to sensory processing. Rodent GC exhibits metastable neural dynamics during ongoing and stimulus-evoked activity, but how these dynamics evolve in the context of a taste-based decision-making task remains unclear. Here we employ analytical and modeling approaches to i) extract metastable dynamics in ensemble spiking activity recorded from the GC of mice performing a perceptual decision-making task; ii) investigate the computational mechanisms underlying GC metastability in this task; and iii) establish a relationship between GC dynamics and behavioral performance. Our results show that activity in GC during perceptual decision-making is metastable and that this metastability may serve as a substrate for sequentially encoding sensory, abstract cue, and decision information over time. Perturbations of the model's metastable dynamics indicate that boosting inhibition in different coding epochs differentially impacts network performance, explaining a counterintuitive effect of GC optogenetic silencing on mouse behavior.

Introduction

Over the past decade, the gustatory cortex (GC) has emerged as a model for studying cortical dynamics. Analysis of neural activity in alert rats has revealed that ensembles of GC neurons consistently hop between semi-stable patterns of activity (Jones et al., 2007; Moran and Katz, 2014; Mazzucato et al., 2016; Mazzucato et al., 2015; Sadacca et al., 2016) termed "metastable dynamics" (for reviews, see La Camera et al., 2019; Brinkman et al., 2022). While these metastable dynamics were originally found in the context of taste processing, recent work has reported that metastability can also be associated with cognitive processes such as taste expectation (Samuelsen et al., 2012; Mazzucato et al., 2019) and sensorimotor transformations (Sadacca et al., 2016; Mukherjee et al., 2019). The latter was shown in the case of rats producing gapes—orofacial reactions signaling aversion and rejection (Grill and Norgren, 1978)—in response to bitter stimulation. During the course of such a consummatory experience, GC neural ensembles switch from taste-encoding to gape-anticipating metastable states (Sadacca et al., 2016; Mukherjee et al., 2019). Optogenetic manipulations of activity preceding this switch lead to a delayed onset of gapes (Mukherjee et al., 2019), demonstrating a causal role of GC's metastable dynamics in naturalistic consummatory behavior.

Recent studies indicated that, in addition to representing taste and naturalistic sensorimotor transformations, GC can also contribute to taste-based decisions in mice engaged in a decision-making paradigm (Vincis et al., 2020). Mice were given one of four tastants—two sweets and two bitters—and, after a brief delay period, had to register a decision by licking one of two lateral spouts. Electrophysiological recordings in mice performing this task uncovered that GC neurons encode sensory and decision-making information during the sampling and delay periods. Interestingly, and counterintuitively for a sensory region, optogenetic perturbation of GC activity during stimulus sampling had no impact on behavioral performance, whereas silencing during the delay period significantly impaired task performance (Vincis et al., 2020). While important in highlighting the role of GC in taste-based decision-making, this research raised two fundamental questions: 1) Is GC's activity metastable during decision-making and, if so, what role does GC metastability play in perceptual decisions? 2) Why is optogenetic perturbation of GC neural activity during stimulus sampling less disruptive than during the delay period?

Here, we investigate these questions using a Hidden Markov Model (HMM) analysis of the data (Radons et al., 1994; Abeles et al., 1995; Jones et al., 2007; Ponce-Alvarez et al., 2012; Mazzucato et al., 2015; Sadacca et al., 2016) and a biophysically grounded model of GC neural activity. Using HMM, we demonstrate the existence of a temporal progression of metastable states emerging, sequentially, for coding sensory information (taste quality), internal deliberations, and finally forthcoming action. To understand the computational mechanisms underlying the genesis of metastable activity and the emergence of coding states in the context of decision-making, we developed a metastable spiking network model of GC. The model provides a mechanistic understanding of the results of our HMM analysis—in particular, the temporal progression of hidden coding states for different sensory and cognitive variables. The model is then used to explain the experimental effects of the optogenetic perturbation—in particular, why performance is more sensitive to perturbations during the delay period than during the stimulus-evoked period (Vincis et al., 2020).

Altogether, our results show that GC supports complex taste-related decisions via sequences of metastable states that follow a precise temporal progression. The mechanism for this progression is explained by a biologically plausible model of GC, which also clarifies the counterintuitive effects of optogenetic silencing on performance, and further provides testable predictions on the role of intensity and

timing of optogenetic perturbations.

Results

Decision-making neural activity in mouse GC is metastable

Mice implanted with drivable multi-tetrode arrays in the gustatory cortex were trained on a taste-based perceptual decision-making task (Vincis et al., 2020). In this task, head-fixed, water-restricted animals sampled a tastant—sucrose, quinine, maltose, or sucrose octaacetate—from a central spout and learned to lick one of two lateral spouts based on the following rule: sucrose or quinine means lick left, and maltose or octaacetate means lick right. A correct directional lick resulted in a water reward from the lateral spout. The intentional counterbalancing of taste quality between cued directions—sucrose and maltose are sweet, while quinine and octaacetate are bitter—ensured that the animal could not rely simply on the generalization of sweet or bitter to perform well on the task, but rather had to recategorize the four stimuli according to the rule. A variable delay between sampling and availability of the lateral spouts was implemented such that the average time between the sampling/taste event (first lick to the central spout) and the decision event (first lick to the lateral spout) was 2.53 s (range: $1.40 - 6.66$ s) (**Figure 1a**). Once mice learned the task (i.e., they reached criterion performance of at least 75% for 3 consecutive days), recording sessions began. We recorded from a total of 16 sessions in four mice, with an average of 198 trials per session (range: $75 - 292$) and 5 simultaneously recorded neurons per session (range: $3 - 9$).

We hypothesized that metastable dynamics support the ability of GC to code for different task-related variables, informed by recent findings revealing a crucial role for GC in sensory and decision-making aspects of this task (Vincis et al., 2020), as well as work demonstrating metastability in GC (Jones et al., 2007; Mazzucato et al., 2015). To test this hypothesis, we fitted a Hidden Markov Model (HMM) to our ensembles of simultaneously recorded spike trains. This method allowed us to segment the neural activity into recurrent sequences of discrete states that are not directly observable—a signature of metastable dynamics previously described in this area (Jones et al., 2007; La Camera et al., 2019; Brinkman et al., 2022). The hidden states were defined as vectors of firing rates across the simultaneously recorded neurons (Gat and Tishby, 1993; Seidemann et al., 1996; Jones et al., 2007; Bollimunta et al., 2012; Ponce-Alvarez et al., 2012; Mazzucato et al., 2015).

HMM was fitted to the spike trains of each session separately, and then used to decode, trial-by-trial, the sequence of hidden states (see **Methods** for details). The neural activity in all sessions and trial types could be characterized by sequences of metastable states, with a median of 6 decoded states per session (range: $3 - 9$). Examples of state sequences are shown in **Figure 1b-d (left panel)**. Note how state transitions occur at random times in different trials. The distribution of state durations resembled an exponential distribution with mean 539 ms (standard deviation: 585 ms; median: 304 ms; range: $50 - 5,546$ ms), in keeping with previous findings in the rat GC (Jones et al., 2007; Mazzucato et al., 2015). These results demonstrate that ensemble activity of mouse GC during decision-making is metastable. Remarkably, some states occurred more often in specific trial types, with their onset times obeying distinct probability distributions, as we show next.

Figure 1 Decision-making activity in GC is metastable and distinct metastable states encode different task-relevant variables. **a**: Perceptual decision-making task schematic (Vincis et al., 2020). S: sucrose, Q: quinine, M: maltose, and O: sucrose octaacetate. **b-d (left)**: Raster plots and HMM-decoded metastable states for in vivo spiking activity during decision-making behavior. "Taste" indicates the animal's first lick to the central spout; "Decision" indicates the animal's first lick to a lateral spout. **b-d (right)**: Classification of hidden states. Quality-coding (**b**, red), Cue-coding (**c**, cyan), and Action-coding (**d**, blue) states. * indicates significant difference by Chi-squared test ($p < 0.05$, corrected for the number of decoded states in the session).

Distinct discrete states encode task-related variables

To identify whether hidden states encoded task-related variables, state occurrences were compared across different trial types (i.e., characterized by different stimuli or actions; see **Figure 1b-d, right** and **Supplementary Figure 1**). For a state to be considered a coding state, occurrence frequencies had to differ across correct trials of the four different stimuli (Chi-squared test, $p < 0.05$). An additional comparison was performed to determine whether the probability of observing the state was significantly different for one stimulus out of the four (assessed by running pairwise Marascuilo post-hoc tests; see **Methods**). If this was the case, the state was considered "Taste ID-coding." If not, we grouped correct trials by cued direction (cue left: sucrose or quinine; cue right: maltose or octaacetate) and by taste quality (sweet: sucrose or maltose; bitter: quinine or octaacetate). If the state appeared with significantly different frequency between correct cue left and correct cue right trials, but not between correct sweet

and correct bitter trials, it was considered a "Decision-coding" state. Similarly, if a state appeared with significantly different frequency between correct sweet and correct bitter trials, but not between correct cue left and correct cue right trials, it was considered a "Quality-coding" state. In the case that both of these comparisons were significantly different, the state was considered "Dual-coding" (see **Methods** for details). Over all the 16 recording sessions we found 10 Taste ID-coding states, 4 Dual-coding states, 18 Decision-coding states, and 5 Quality-coding states. We chose to focus on Quality-coding and Decision-coding states because they have intuitive interpretations in the context of our behavioral task: Quality-coding states are associated with either sweet or bitter taste qualities, which reflect processing of sensory information. Decision-coding states are associated with either the left or right directions, which reflect decision-making activity. These states were distributed sparsely across sessions: 56.3% of sessions contained a Decision-coding state and 25% of sessions contained a Quality-coding state, with only 12.5% of sessions containing both (**Supplementary Table 1, Column 1**).

Decision-coding states were additionally divided into two categories: "Cue-coding" and "Action-coding" states, the former being associated with the perceptual categorization of the stimulus based on the direction it cues (i.e., sucrose or quinine → left; maltose or octaacetate → right), and the latter being associated with the execution of the action (i.e., go left; go right). This separation was based on the frequency of occurrence of states in correct and incorrect trials. Decision-coding states associated with the same cued direction in correct and incorrect trials were deemed Cue-coding since they were consistently associated with the cued direction, whereas Decision-coding states associated with opposite cued directions in correct and incorrect trials were deemed Action-coding since they were consistently associated with the chosen direction (see **Methods** for details). We found 10 Action-coding states and 7 Cue-coding states (**Supplementary Table 1, Column 1**). Imposing stricter criteria at this step of the classification (i.e., requiring that sessions contain at least 10 incorrectly performed trials of each cued direction) did not qualitatively alter any of the results shown here, but decreased the number of Action- and Cue-coding states to 8 and 4, respectively.

Temporal sequences of coding states reflect task structure

Previous literature highlights the importance of temporal dynamics in GC (Katz et al., 2001; Fontanini and Katz, 2006; Jones et al., 2007; Sadacca et al., 2012; Sadacca et al., 2016; Mazzucato et al., 2019), with different variables being encoded at different times following stimulus delivery. To determine whether the variables associated with a taste-based, 2-alternative choice (2-AC) task are encoded sequentially, we analyzed the within-trial onset times of coding states. Consistent with the literature, we hypothesized that Quality-coding states should appear early in the trial, followed by Cue-coding and Action-coding states. To test this hypothesis, all within-trial onset times of the classified coding states were pooled across all sessions. Trial durations were "warped" so that taste delivery occurred at time 0 and decision occurred at time 1; this way timings could be compared across different trials in common time units. The distributions of coding states' onset times in these common time units are plotted in **Figure 2a**. The distributions are clearly separated in time and arranged in a sequence where Quality-coding states appear before Cue-coding states, which, in turn, appear before Action-coding states. Mean onset times for Quality-coding, Cue-coding, and Action-coding states were 0.10, 0.42, and 0.68 (common time units), with inferred peaks occurring at 0.05, 0.39, and 0.88, respectively. This result is consistent with sequential encoding in each trial unfolding from sensory information processing to perceptual categorization, and finally to decision-making and action execution.

Figure 2 Coding state onset times exhibit an intuitive temporal sequence. **a**: Distribution of onset times of all states classified as either Quality- (red), Cue- (cyan), or Action-coding (blue) for experimental data showing a progression from coding for sensory information to abstract cue information to action information. **b**: Distribution of onset times for spiking data that are shuffled circularly. **c**: Distribution of onset times for spiking data that are swapped across time.

To ensure that coding states were meaningful, and not a by-product of HMM analysis, a series of control analyses were performed. Randomly permuting trial type (i.e., stimulus) labels within correct and incorrect trials and repeating the state coding classification analysis severely reduced the probability of finding coding states in the data, as expected (5 Quality-coding states in unshuffled data vs. an average of 0.4 in shuffled datasets; 18 Decision-coding states in unshuffled data vs. none in shuffled datasets). In addition, two surrogate datasets were generated from raw data by shuffling the spike trains circularly or by swapping segments of simultaneous activity across time (see **Methods**). The

circular shuffle preserves individual spike train autocorrelations but disrupts co-activation (hence, pairwise correlations) between neurons; the swap shuffle does the opposite, i.e., it preserves neuron co-activations but disrupts autocorrelations (Maboudi et al., 2018). Both types of shuffling preserve the individual neurons' firing rates. These shuffled datasets were fitted with new HMMs and the resulting states were re-classified for their coding properties. Overall, we found only a modest reduction in the numbers of different coding states (**Supplementary Table 1**), which may indicate that neural autocorrelations or pairwise correlations in the data alone contain enough information to identify activity associated with task variables. However, onset time distribution plots constructed for the shuffled datasets showed no clear separation among coding states of different types (**Figure 2b-c**). Thus, even though there was enough information in the shuffled datasets for the model to find coding states, the temporal structure of these states was completely disrupted.

Modeling metastable decision-making activity in GC

Results from HMM analysis demonstrate that GC relies on metastability to represent variables associated with the performance of a taste-based 2-AC task. To investigate the potential circuit-level mechanisms driving metastable activity in the context of the 2-AC task, we developed a recurrent spiking network model of GC that could replicate the key experimental findings.

Metastability in spiking network models can be achieved by a clustered architecture where synaptic connections among neurons within the same cluster are stronger than those among neurons of different clusters (Deco and Hugues, 2012; Litwin-Kumar and Doiron, 2012; Mazzucato et al., 2015; Rostami et al., 2020). Our network contained both excitatory (E) and inhibitory (I) neurons, each organized into 14 uniformly-sized clusters with stronger synaptic connections than among neurons of different clusters. Each excitatory cluster was paired with a partner inhibitory cluster via stronger synaptic connections than among neurons of non-partner E-I pairs (Rostami et al., 2020) (see **Figure 3a**, **Supplementary Figure 2a**, and **Methods** for details).

The model was built to encode the crucial variables associated with the decision-making task, namely stimulus (taste), cued direction (cue), and selected direction (action). This was achieved by wiring the network such that specific properties could be assigned to 8 randomly selected excitatory clusters, coding respectively for: sucrose, quinine, maltose, sucrose octaacetate, cue left, cue right, action left, and action right. Taste-selective clusters were characterized by having half of the neurons receive additional external input when either sucrose, quinine, maltose, or sucrose octaacetate were presented. The increased input varied in time as an alpha function (difference of two exponentials) that began at the start of the trial (**Figure 3b, top left**).

To replicate the experimental finding of coding states associated with a particular taste quality (i.e., sweet or bitter), clusters of neurons coding for sucrose and maltose were more strongly connected than clusters coding for, e.g., sucrose and quinine. Similarly, clusters coding for quinine and octaacetate were more strongly connected than clusters coding for, e.g., quinine and maltose. The same structure was imposed on the corresponding inhibitory partners. This structure was obtained by randomly selecting one quarter of the neurons in each partner E-I pair and connecting them with strong synaptic weights $J_{++}J_{\alpha\beta}$, with $J_{\alpha\beta}$ being a common reference synaptic weight from neurons in cluster β to neurons in cluster α ($\alpha, \beta \in \{E, I\}$). Neurons in the same taste cluster, but outside this more strongly connected subpopulation, had weaker synaptic connections (mean $J_+J_{\alpha\beta}$, with $J_{++} > J_+ > 1$; see **Methods**, **Supplementary Figure 2b**, and **Supplementary Table 3** for model parameters). This structure

Figure 3 Clustered spiking network model for decision-making activity. **a**: General structure of the network. Neurons in the network are organized into fourteen excitatory (black) and partner inhibitory (red) clusters and connected in a highly recurrent manner. The zoomed inset indicates the generic structure of all of the connections that exist between any two excitatory clusters and their inhibitory cluster partners (pointed arrowheads are excitatory connections, flat arrowheads are inhibitory connections). **b**: Details of synaptic connections between taste, cue and action clusters, with inputs simulating the sensory stimulus (**left, top**) and the gating GO signal (**left, bottom**). **c**: Raster plot of metastable network activity in the absence of a stimulus (**top**) and after stimulation (**bottom**). Neurons are grouped by clusters (black: excitatory, red: inhibitory). S: sucrose, M: maltose, Q: quinine, O: sucrose octaacetate, CL: cue left, CR: cue right, AL: action left, and AR: action right.

mimics overlapping populations of neurons with similar response properties, and reflects the fact that many neurons in the GC (Jezzini et al., 2013) and elsewhere in the cortex (Curti et al., 2004) respond to multiple stimuli.

The 2 cue-selective clusters received stronger input from the appropriate pair of taste-selective clusters (i.e., connections from sucrose and quinine clusters to the "cue left" cluster and from maltose and sucrose octaacetate clusters to the "cue right" cluster were stronger compared to the cross-connections; see **Figure 3b** and **Supplementary Figure 3** for additional details); similarly the 2 action-selective clusters received stronger input from the appropriate cue-selective cluster (i.e., stronger connections from the "cue left" cluster to the "action left" cluster, and from the "cue right" cluster to the "action right"

cluster). Cue- and action-selective clusters also had stronger intracluster connections than other clusters (**Supplementary Table 3**) to promote more "persistent" activity intended to reflect stable mental constructs.

To represent the approach of the lateral spouts in the behavior task, which prevents mice from acting on a decision until the end of the delay period, we incorporated a preparatory action gating signal in the model (**Figure 3b, bottom left**). The gate prevents the action-selective clusters from receiving input until a randomly chosen point in time in each trial (uniformly distributed over [1.6 s, 2.4 s]), representing availability of the spouts.

Altogether, the anatomy and topology of the model network gave rise to a random-looking, metastable dynamics in the absence of a stimulus (**Figure 3c, top**), and to a cascade of processes—from taste to cue to action—in the presence of a stimulus (Miller and Katz, 2010), due to preferential connections embedded within the highly recurrent network model (**Figure 3c, bottom**). The network typically responded to stimuli with high firing rates initially in the stimulus-responsive cluster and its overlapping partner, then in a cue cluster, and then in an action cluster (**Figure 3c, bottom**).

To infer the model's choice on each trial, we computed firing rates for action clusters in 50 ms bins and considered a cluster to be "on" when it fired at $\geq 40\%$ of its maximum firing rate across the whole session. Behavioral scoring of trials was then evaluated following this simple rule: if the correct action cluster came on between 0.5 s and 3.0 s after stimulus onset and the incorrect action cluster did not, the trial was scored as correct; if the incorrect action cluster came on during the same window and the correct action cluster did not, the trial was scored as incorrect; if neither action cluster or both action clusters came on during this window, the decision was assigned randomly.

To test the model, we randomly generated 10 different synaptic weight matrices, which may be thought of as 10 independent network instances. For each instance of the network, we ran 100 trial simulations (25 simulations per tastant); in each simulation, a stimulus was applied and the network was allowed to evolve over 3 seconds. Over all 10 realizations, the network had an average accuracy rate of 80.1%, which is comparable to the experimental performance of mice performing the real task (82.8%; Vincis et al., 2020). To assess whether the network model could reproduce the metastable dynamics observed experimentally, we randomly selected one neuron from each excitatory cluster to form a 14-neuron sample. The spike trains of these sampled 14-neuron ensembles were fitted with an HMM over the 100 trials (**Figure 4a**). Metastable states were decoded on a trial-by-trial basis as was done for the experimental data, and state coding properties were classified according to the same definitions used in **Figure 1b-d** (see **Figure 4c-e**).

Analysis of state onset times (**Figure 5a**) revealed that these distributions corresponded quite well with those observed in the experimental data. Most notably, the model captured the temporal sequence of distribution peaks observed in the data, showing a progression through time from taste-coding to abstract cue-coding and finally to action-coding. The mean onset times for Quality-coding, Cue-coding, and Action-coding states were 0.05, 0.50, and 0.77 common time units, with inferred peaks occurring at 0.03, 0.41, and 0.82, respectively. Similarly to the experimental data, no coding states with any meaningful temporal dynamics were found when shuffling the simulated data (circular or swap shuffle, see **Methods** and **Figure 5b-c**). This finding confirms that coding the temporal structure of the task via a sequence of metastable states requires a particular network structure, with our model providing a specific link between network structure and function.

Figure 4 HMM analysis of network-simulated data. **a**: Example network activity showing sucrose-triggered response. **b**: Raster plot with overlaid HMM-decoded hidden states corresponding to the trial shown in **a**. Arrows indicate onsets of different coding states. **c-e (left)**: Raster plots and HMM-decoded metastable states for simulated spiking. **c-e (right)**: Classification of hidden states. Quality-coding (**b**, red), Cue-coding (**c**, cyan), and Action-coding (**d**, blue) states. * indicates significant difference by Chi-squared test ($p < 0.05$, corrected for the number of decoded states in the session).

Relationship between GC dynamics and behavioral performance

To further probe the relationship between network structure and function, we subjected our model network to a series of external perturbations. Inspired by the optogenetically-induced silencing of GC used in Vincis et al. (2020), we inhibited network activity at different time points during simulated trials, and measured the effect of the manipulation on task accuracy. This allowed us to identify temporal windows of "weakness" and "robustness" to perturbation in the model. Network activity was modulated by applying a square pulse stimulus—parameterized by strength, duration, and center point—to all inhibitory neurons in the network (**Figure 6a**). When applied for 250 ms and centered on the middle of the trial, a square pulse increasing input current by 100% relative to baseline typically produced an 11.4% increase in inhibitory neurons' mean firing rates. On average, mean firing rates increased in 12 out of 14 inhibitory clusters and decreased in the remaining 2. The latter decrease in firing rate occurred when the cluster was already active (i.e., firing rates $>\sim$ 6 Hz) prior to the square pulse. The same

Figure 5 Coding state onset times in the simulated data qualitatively reflect those in experimental data. **a**: Distribution of onset times of all states classified as either Quality- (red), Cue- (cyan), or Action-coding (blue) for simulated data showing a progression from coding for sensory information to abstract cue information to action information. **b**: Distribution of onset times for spiking data that are shuffled circularly. **c**: Distribution of onset times for spiking data that are swapped across time.

square pulse reduced mean firing rates of excitatory clusters by 99.4%. **Figure 6b** shows the effects of the square pulse strength on cluster firing rates.

We tested various silencing strengths (square pulse amplitudes) and calculated the mean network per-

Figure 6 Simulated optogenetic silencing disrupts network accuracy and reveals temporally-specific points of sensitivity and robustness in network structure. **a**: Example network simulation with 250 ms simulated optogenetic perturbation (100% increase in external input current to inhibitory neurons). **Bottom**: Inputs to the network are sucrose stimulus (solid black), silencing (gold), and action gate (dotted black). **Middle**: Raster plot of network activity. **Top**: Changes in firing rates of excitatory (black) and inhibitory neurons (red) resulting from simulated optogenetic perturbation. **b**: Effect of variable-strength silencing (centered at 1.5 s) on cluster firing rates (each dot represents a cluster averaged across 10 networks). Percent changes are expressed relative to the mean firing rates prior to silencing (black: excitatory, red: inhibitory). The outlier points ($> 50\%$ increase) are inhibitory action clusters, which have very low mean firing rates prior to silencing due to the action gate mechanism (see the main text). Open circles represent medians over all clusters. **c**: Effect of variable-strength silencing on network accuracy. Solid curves are mean accuracies as functions of the silencing center for a particular silencing strength and fixed 250 ms duration. Gray band represents mean ± 1 standard deviation of network performance without silencing. Shaded red, cyan, and blue curves are coding state onset time distributions (same as in **Figure 5**, but scaled to the same height) to provide within-trial temporal context. **d**: Silencing-induced accuracy impairment is mitigated by temporally extending the stimulus input. **Top**: Stimuli with different decay time constants (τ_D). **Middle**: Curves representing mean accuracies as functions of the silencing center (as in **c**) averaged across 10 networks, 100 trials each. **Bottom**: Correlation between average slope of each mean accuracy curve (over the first 0.5 s) versus corresponding stimulus τ_D.

formance accuracy (10 networks, 100 trials each) as a function of the silencing center point (**Figure 6c**). Albeit limited in its effect, weak silencing (25% and 50%) was capable of reducing performance accuracy below one standard deviation of the baseline mean (gray band), but only when it was centered on particular points during the simulation (**Figure 6c**). These points of sensitivity occurred during the network's transition from taste-coding to cue-coding and during the onset of action-coding. On the contrary, strong silencing (100% and 125%) reduced network performance accuracy below baseline for

a wide range of center point values. The only points at which performance was resistant to silencing were at the very beginning of the trial (0 ms and 50 ms centers, corresponding to silencing from 0 ms to 175 ms after stimulus onset) and at the very end (3 s center, corresponding to silencing from 565 ms to 690 ms after the average onset of Action-coding states; **Figure 6c**). The relative ineffectiveness of perturbation during sensory processing is consistent with the counterintuitive observations from the optogenetic silencing reported in Vincis et al. (2020). Those experiments showed that task performance is unaffected by silencing GC during the sampling period (taste-coding stage) but impaired by silencing GC during the delay period (cue-coding stage).

Ineffectiveness of silencing during the sampling period is presumably due to lingering of the taste stimulus past the perturbation window. To test this hypothesis, we systematically varied the time course of the simulated gustatory input (decay time varied from 160 ms to 705 ms). For a 250 ms long pulse with 100% strength relative to baseline, increasing the stimulus' decay time constant resulted in increased resistance to silencing over the first half of the trial—average performance increased from 57.5% to 75.6%, and the average initial slope of the performance curve increased from -41.2%/s to -2.5%/s (**Figure 6d**). In summary, the results of perturbing the model network are consistent with the experimental findings showing that task performance is resilient to GC inhibition during the sampling period, and further imply that this robustness is the result of stimulus information persisting past the silencing period.

Discussion

In this study, we demonstrate that neural activity associated with taste-based decisions is metastable and temporally organized to sequentially encode information related to sensory stimuli, decision variables, and upcoming actions. We further provide a computational model that accounts for metastable dynamics in GC and its link with behavior. This model allows us to explain why silencing GC during sensory stimulation has a smaller impact on performance compared to perturbation during the deliberation period. These results significantly advance our understanding of GC and its role in decision-making, and confirm metastability as a dynamic regime relevant for GC's function.

Temporal dynamics of GC activity and functional roles of metastability in GC

GC neurons produce time-varying responses following gustatory stimulation (Katz et al., 2001; Katz et al., 2002; Bouaichi and Vincis, 2020). Firing rates transition through three successive epochs, each having a different onset and encoding a distinct aspect of taste. The earliest epoch lasts for ~ 250 ms and encodes the contact of taste with the tongue. The second epoch (from ~ 250 ms to ~ 1 s) represents chemosensory information—that is, taste quality (i.e., bitter, sweet, salty, sour, and savory). Finally, the third epoch (after ~ 1 s) encodes the palatability of the stimulus and correlates with ingestive or aversive behaviors. This sequence is believed to result from recurrent interactions among GC, thalamic, and limbic areas such as the basolateral nucleus of the amygdala (Piette et al., 2012; Samuelsen et al., 2012; Samuelsen et al., 2013; Livneh et al., 2017; Lin et al., 2021). These dynamics were initially uncovered through a careful analysis of single neuron firing rate modulations averaged across trials (Katz et al., 2001). Additional studies of the ensemble activity from simultaneously recorded neurons further confirmed the existence of these dynamics at the single trial level, revealed trial-by-trial consistency of epochs, and identified epochs as metastable states with rapid onsets (Jones et al., 2007; Mazzucato et al., 2015; Mazzucato et al., 2016).

Recent experiments in rodents engaged in behavioral tasks suggest that additional epochs containing task-dependent variables could exist. For instance, recordings from GC of rodents intentionally licking a spout show preparatory activity that precedes the somatosensory epoch (Stapleton et al., 2007; Dikecligil et al., 2020). Studies using taste-anticipating cues prove the existence in GC of anticipatory firing rate modulations, which encode general and specific expectations (Samuelsen et al., 2012; Gardner and Fontanini, 2014; Livneh et al., 2017). Similarly, experiments in rodents performing a decision-making task show neural activity consistent with sequences of epochs encoding decision-related variables (Fonseca et al., 2018; Vincis et al., 2020). While this more recent work has been relevant in showing that GC does not just multiplex sensory variables, it has largely relied on single neuron analyses and across-trial averages.

The work presented here is the first that shows that variables associated with reward-driven decision-making are encoded by GC as metastable states. Our HMM analysis of GC ensemble activity during the four-taste, perceptual decision-making task has demonstrated that GC activity traverses sequences of metastable states encoding relevant task variables. These results are in line with prior evidence from higher order areas (Durstewitz et al., 2010; Ponce-Alvarez et al., 2012; Stokes et al., 2013; Rich and Wallis, 2016; Benozzo et al., 2021) and are the first to show such a phenomenon in a sensory area. Our analysis further shows that, in the rodent GC, metastable states undergo a temporal progression with taste-coding states occurring first, followed by cue-coding states, and finally by action-coding states. This progression is consistent with a model in which GC first encodes sensory information (e.g., sweet vs. bitter), then decisional variables (e.g., predictive value of taste or preparatory activity associated with actions), and then actions (e.g., lick left vs. lick right). Importantly, coding state transitions occur at variable times in different trials, allowing for pinpointing of the internal occurrence time of sensory perception or decision. This also allows for disentangling of the neural activity related to internal processes from the neural activity immediately following external triggers.

Perturbing GC dynamics

GC dynamics and their behavioral significance have been extensively tested via temporally precise optogenetic perturbations. Studies in rats receiving tastants through intraoral cannulae demonstrated that the effects of perturbations on gaping depend on the specific timing of silencing (Mukherjee et al., 2019). The differential impact of perturbations during specific temporal epochs was also present in mice performing a four-taste, perceptual decision-making task (Vincis et al., 2020). Optogenetic activation of parvalbumin-positive interneurons—an established method for silencing cortical activity (Guo et al., 2014)—was effective in reducing behavioral accuracy only if performed during the delay period, when cue-related, preparatory activity occurs, and not during the period of taste sampling. While these results were interpreted as evidence that GC's cognitive signals play a role in guiding taste-based decision-making, they also raised questions about the behavioral role of GC taste-coding states. In this work, we provide a possible explanation for these results by analyzing the behavior of our model under a silencing perturbation. Strong perturbations dramatically affected performance, with a stronger effect during the cue-coding state and a limited effect during taste-coding and action-coding states. The limited effect of silencing during taste sampling on network performance bears a remarkable qualitative resemblance to the experimental results of Vincis et al. (2020); furthermore, the model predicts that the effect depends on the duration of the thalamic input relaying the gustatory stimulus to GC. The limited impact during the action-coding state results from the metastable nature of the dynamics and is also a testable prediction of the model: if the correct action cluster is activated prior to the perturbation, the correct decision will

be taken despite the perturbation. These predictions can be tested in additional experiments.

Modeling GC: Metastability and decision-making

The observation of reliable and reproducible metastable dynamics in GC has inspired modeling efforts aimed at explaining the network mechanisms underlying this phenomenon and its functional significance. Miller and Katz (2010) showed how a spiking metastable network can account for state sequences observed during consummatory behavior (associated with ingestion or rejection of tastants); later, Mazzucato et al. (2015) demonstrated the existence of metastable dynamics in the absence of overt stimulation, and built a spiking network model of GC able to spontaneously produce metastable dynamics. This network has formed the basis of subsequent models that have captured a wealth of data pertaining to neural dynamics in GC (Mazzucato et al., 2016; Mazzucato et al., 2019). The model described here represents another significant advancement, in that the model can further process stimuli based on the abstract decision they instruct. The four tastants instructing decisions are categorized according to their taste quality first and their predictive value after, reproducing the temporal sequence of metastable states representing taste quality, predictive value of the taste, and action. The model also produces decisions with a similar performance to our experimental subjects.

In order to reproduce the experimental results observed in mice performing the four-taste, 2-AC task, the network had to include the following features: i) partial overlapping of clusters representing stimuli that share the same taste quality; ii) biased intercluster connections creating a flow of information from sensory clusters to cue clusters to action clusters; iii) an action gate signal priming action. These ingredients were absent in previous models and suggest the existence of possible biological counterparts in GC. Partially-overlapping clusters take into account that GC neurons may respond to multiple tastes (Jezzini et al., 2013) and may be the basis of stimulus generalization based on predominant taste quality. Biased intercluster connectivity may be the result of synaptic plasticity leading to the formation of neuron ensembles producing preparatory and action-related activity. The action gate was inspired by the presence of the lateral spouts moving toward the mouse, hence providing an implicit, ramping "GO" signal that may correspond either to an anticipatory input from areas like the amygdala, or to a disinhibitory signal triggered by the cue. Additional experiments are needed to test these predicted mechanisms.

The model also explains the effects of optogenetic perturbations and makes further predictions based on the strength of the perturbation and the period in which it is applied, as discussed in the previous section. We emphasize that, once the network model is structured to obtain metastable dynamics coding sequentially for taste, cue, and action, these additional model predictions emerge without further parameter tuning.

In summary, the present work shows that metastable dynamics occurs in GC during decision-making and provides a computational model explaining the genesis of metastability and its role in taste-based decisions. The ability of our model to reproduce the experimental results, together with our past body of work (Mazzucato et al., 2015; Mazzucato et al., 2016; Mazzucato et al., 2019), suggest that the clustered inhibitory-excitatory network model may truly underlie the anatomical and functional organization of the rodent GC.

Acknowledgements

We thank Dr. Luca Mazzucato, Xiaoyu Yang, and Tianshu Li for sharing their computer code, and Dr. Arianna Maffei for helpful discussions and feedback on the manuscript. This work was partially supported by a U01 grant from the NIH/NINDS Brain Initiative (1UF1NS115779) to G.L.C. and A.F and by grants R01DC015234 and R01DC018227 to A.F. The content of this article is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Methods

Summary of behavioral procedure and electrophysiology

Methods for behavior and electrophysiology are available in Vincis et al. (2020), which presents the dataset analyzed in this study. Briefly, mice were unilaterally implanted with 8 tetrodes in the left gustatory cortex for chronic electrophysiological recording. After recovering from surgery, mice were water-restricted and trained to perform a four-taste, perceptual decision-making task to obtain water rewards. For the task, head-fixed mice sample a tastant—either sucrose, quinine, maltose, or sucrose octaacetate—from a central spout. The sampling period (time between first and last lick to the central spout) was ~ 0.5 s. After a delay period (~ 2 s) two lateral spouts advance in front of the animal's mouth. To receive the water reward, the mouse must lick the left spout if the tastant was sucrose or quinine and must lick the right one if the tastant was maltose or sucrose octaacetate. Testing commenced once mice achieved a criterion performance of at least 75% for 3 consecutive days. Electrophysiological recordings were obtained while mice performed the task.

Raw data and preprocessing

Raw data consisted of spike trains from simultaneously recorded neurons over 21 recording sessions involving 5 mice. Within each recording session, trained mice performed an average of 190 trials (range: $75 - 292$) of the behavioral task at an average accuracy of 82.8% (range: $71.0\% - 93.3\%$). Only spikes within 100 ms before the taste event (first lick to central spout) to 100 ms after the decision event (first lick to lateral spout) were considered for each trial (we refer to these 100 ms segments before and after events as "padding"). These windows of analysis for each trial were concatenated and, prior to model fitting, neurons whose firing rates averaged over this total window were < 2 Hz were excluded from analysis. Sessions with < 3 simultaneously recorded neurons were then excluded from analysis. This resulted in 16 recording sessions involving 4 mice, with an average of 5.4 neurons per session (median: 5.5; range: $3 - 9$).

Model fitting and hidden state decoding

Hidden Markov Models (HMMs) were fit to spiking data from Vincis et al. (2020) as described in Mazzucato et al. (2019). Briefly, spike sequences in a single session were binned into 2 ms bins, and each bin was assigned a symbol from 1 to $N + 1$, where N is the number of neurons and symbol s means that only the sth neuron fired in the current bin (multiple firings were rare and were replaced with a random

selection of one of the firing neurons). The HMM is completely determined by the number of hidden states, M, the transition probability matrix, T, and the emission probability matrix, E. The entry T_{ij} of the $M \times M$ transition probability matrix is the probability of making a transition from state i to state j; the entry E_{is} of the $M \times (M+1)$ emission probability matrix is the probability of producing symbol s while in state i. This model can be considered an approximation to a Poisson-HMM in the case of very short time bins, with the exception that firing of multiple neurons in the same bin is not permitted (see Mazzucato et al., 2015 for details). The HMMs were fitted to the data by maximum likelihood, using the standard Baum-Welch expectation-maximization algorithm (Rabiner, 1989). For each dataset, we fitted HMMs with values of M ranging from 2 to 50 and then chose the model with the lowest BIC score as the best model (see below). The maximal number of hidden states selected in the best model never exceeded 29. For each value of M, 10 different random initial conditions were used for T and E. For initial values of T, off-diagonal entries were close to zero and diagonal terms were close to 1. For initial values of E, entries were drawn independently from a uniform distribution between 0 and 1.

From each initial condition, the Baum-Welch algorithm was performed for 50 iterations (or until convergence at $< 10^{-10}$ tolerance on the log-likelihood, transition, and emission probabilities). Out of the 10 models, the model with the highest log-likelihood was selected as the best model for that given value of M. The best models for different M values were then compared using the Bayesian Information Criterion (BIC) to obtain the optimal model with the optimal number of states (Ponce-Alvarez et al., 2012; Mazzucato et al., 2019; Benozzo et al., 2021). Specifically, the optimal model was the one that minimized the BIC score: $-2LL + [M(M-1) + MN]\ln B$, where LL is the log-likelihood of the model given the data, M is the number of hidden states, N is the number of neurons, and B is the total number of observations in each session (i.e., the total number of time bins across all trials).

After fitting the models, we computed the posterior probabilities of each hidden state given the model and the data in each bin in single trials. A state admissibility criterion was then imposed: a state was considered detected with enough confidence only if its posterior probability was ≥ 0.8 for ≥ 50 consecutive ms (Mazzucato et al., 2019; see also Seidemann et al., 1996; Jones et al., 2007; Miller and Katz, 2010; Bollimunta et al., 2012; Ponce-Alvarez et al., 2012; Mazzucato et al., 2015; Benozzo et al., 2021 for similar methods).

Classification of decoded states

For each session, all trials were associated with two labels: a stimulus label (sucrose, quinine, maltose, or sucrose octaacetate) and an outcome label (correct or incorrect). This allowed us to compute occurrence probabilities of states given the trial labels, $P(S \mid \text{Stimulus, Outcome})$, as the fraction of trials with the given labels in which state S was detected at least once. Each decoded state's coding properties were determined by first looking only at correct trials and comparing the state occurrence probability $P(S \mid \text{Stimulus, Correct})$ across all 4 stimulus labels through a Chi-squared test. If significant, all 6 pairwise post-hoc Marascuilo tests (Prins, 2013) were examined. If there was exactly one stimulus label for which all post-hoc tests involving it were significant, S was classified as a Taste ID-Coding state. Otherwise, coding properties were investigated by grouping all trials into cue left vs. cue right trials (sucrose and quinine vs. maltose and sucrose octaacetate trials) and sweet vs. bitter trials (sucrose and maltose vs. quinine and sucrose octaacetate trials). The fraction of cue left, correct trials featuring S, $P(S \mid \text{Cue left, Correct})$, was compared to the fraction of cue right, correct trials featuring S, $P(S \mid \text{Cue right, Correct})$ through a second Chi-squared test, and the fraction of sweet, correct

trials featuring S, $P(S \mid \text{Sweet, Correct})$, was compared to the fraction of bitter, correct trials featuring S, $P(S \mid \text{Bitter, Correct})$, through a third Chi-squared test (see **Supplementary Figure 1**). In each case, significance level was taken to be 0.05, corrected for the number of decoded states, N_d (i.e., $p < 1 - (1 - 0.05)^{1/N_\text{d}}$ is considered significant). Results from these tests were used to classify states as Decision-coding (significant result on second test but not on third test), Quality-coding (significant result on third test but not on second test), Dual-coding (significant results on both tests), and Non-coding (non-significant results on both tests). Decision-coding states were further divided into Cue-coding and Action-coding, as explained below. To control for chance occurrences of coding states, stimulus labels on correct trials were randomly permuted 10 times and, following each permutation, proportions of trials containing each decoded state were re-computed and the same Chi-squared tests were re-run to re-classify the states. We then compared the numbers of coding states found with this shuffling procedure to the number found from unshuffled data.

Decision-coding states were further subclassified by whether their "preferred direction" switched or remained the same between correct and incorrect trials. A coding state's preferred direction was defined, separately for correct and incorrect trials, as the cued direction that maximized its occurrence frequency (assuming this frequency differed between trials of opposite cued directions): if $P(S \mid \text{Cue left}) > P(S \mid \text{Cue right})$ in correct and incorrect trials, Decision-coding state S codes for the left cued direction, and analogously for states coding for the right cued direction. Vice versa, if $P(S \mid \text{Cue left}) > P(S \mid \text{Cue right})$ in correct trials, but $P(S \mid \text{Cue left}) < P(S \mid \text{Cue right})$ in incorrect trials, Decision-coding state S codes for the action taken; the same conclusion is reached if $P(S \mid \text{Cue left}) < P(S \mid \text{Cue right})$ in correct trials, but $P(S \mid \text{Cue left}) > P(S \mid \text{Cue right})$ in incorrect trials. Finally, if the state S occurred equally often in cue left and cue right trials (either for correct or incorrect trials), we did not subclassify the Decision-coding state.

Randomization controls

Prior to model fitting, raw data were manipulated to form two additional surrogate data sets that served as additional controls (Maboudi et al., 2018; Benozzo et al., 2021; Recanatesi et al., 2022). For the first control, each spike train was shuffled circularly in each trial of a given session: a random number Δt from the uniform distribution on $[0, t_\text{trial}]$, where t_trial is the time length of the trial (including the padding), was added to each neuron's spike train in each trial independently such that the portion of the spike train that now exceeded the end of the trial would "wrap back around" to the beginning (i.e., if a trial window starts at 0 and ends at t_trial, and the spike train $\{t_i\}$ is shifted by Δt, the new spike train is $\{t_j\} = \{t_i + \Delta t, \text{ if } t_i + \Delta t < t_\text{trial}; \ t_i + \Delta t - t_\text{trial}, \text{ otherwise}\}$). This procedure preserves the autocorrelations of individual neurons but alters the co-activation/pairwise correlations between neurons. For the second control, each session had its raw spiking data swap-shuffled: each trial's neuron-by-time spike raster was segmented into 5 ms bins and then these bins/columns were randomly permuted for each trial independently. This randomization procedure preserves the co-activation/pairwise correlations between neurons but removes the autocorrelations of individual neurons. Both procedures preserve the trial-averaged firing rates of individual neurons.

Onset time analysis

For each session, lists of decoded states of each classified type (Quality-coding, Cue-coding, and Action-coding) were obtained. For each state in each list, all onset times of the state in each trial were obtained. Onset times were expressed relative to their trial's taste event ($t = 0$), with any negative times (i.e., states beginning during the left padding period) taken to be 0. To account for the fact that the inter-event interval (IEI) between the taste and decision event varied across trials (average: 2.52 s; range: $2.01 - 6.66$ s), time was rescaled (or "warped") in each trial by dividing onset times by the value of IEI in that trial. As a result, warped onset times were between 0 and 1 (where 0 is the time of the taste event and 1 is the time of the decision event) regardless of trial, neuron, session, and classification of the hidden state, and could be directly compared. From this, we obtained a list of (warped) onset times for coding states of each classified type. These lists were pooled across all sessions to obtain histograms of the onset times for coding states of each classified type (bin size: 0.05 warped time units). From the histograms, probability distributions of the onset times were estimated from linear interpolation on the peaks (100 points between any two adjacent bins). Distributions were then smoothed with a Gaussian filter (400 point width).

Spiking network model

The core of the model was a clustered excitatory/inhibitory spiking network with $4,000$ excitatory (E) neurons and 994 inhibitory (I) leaky integrate-and-fire (LIF) neurons. Each neuron i had subthreshold dynamics that obeyed the differential equation in time t:

$$\frac{dV_i(t)}{dt} = -\frac{V_i(t) - V_L}{\tau_{\mathrm{m}}} + \frac{I_{i,\mathrm{ext}}(t) + I_{i,\mathrm{syn}}(t)}{C},$$

where $V_i(t)$ is neuron i's membrane potential, V_L its equilibrium/leak potential, τ_{m} its membrane time constant, C its capacitance, and its total current $I_i(t)$ is the sum of its external input current $I_{i,\mathrm{ext}}(t)$ and its synaptic current $I_{i,\mathrm{syn}}(t)$. The synaptic current to neuron i in turn obeyed the differential equation in time t:

$$\frac{dI_{i,\mathrm{syn}}(t)}{dt} = -\frac{I_{i,\mathrm{syn}}(t)}{\tau_{\mathrm{syn}}} + \frac{1}{\tau_{\mathrm{syn}}} \sum_j J_{ij} \sum_n \delta(t - t_j^{(n)}),$$

where τ_{syn} is the synaptic time constant (different for excitatory and inhibitory presynaptic neurons), J_{ij} is the synaptic connection strength from presynaptic neuron j, δ is the Dirac delta function, and $t_j^{(n)}$ is a presynaptic spike time with n indexing over all spikes from a particular presynaptic neuron j. This system of differential equations was integrated numerically using the forward Euler algorithm with time step $dt = 0.05$ ms. Whenever $V_i(t)$ hit a threshold V_{th}, a spike was emitted and the membrane potential was immediately reset to a value V_{r} for a refractory period τ_{r} (see **Supplementary Table 3** for parameter values).

The E neurons were arranged into 14 clusters of 250 neurons each, plus 500 neurons not arranged in clusters (the "background population"). The I neurons were arranged into 14 clusters of 71 neurons each. Neurons were randomly connected with probability $P_{\alpha\beta}$, with $\alpha, \beta \in \{\mathrm{E,I}\}$. In general, neurons in the same cluster had stronger synaptic connections than neurons belonging to separate clusters, given by $J_{\mathrm{E+}+}J_{\mathrm{EE}}$ inside E clusters and $J_{\mathrm{I+}+}J_{\mathrm{II}}$ inside I clusters. Each E cluster was paired with a partner

I cluster. Synaptic weights between E and I neurons belonging to E-I partner clusters ($J_{I+}J_{\alpha\beta}$, with $\alpha, \beta \in \{E, I\}$) were stronger than between E and I neurons in non-partner E-I clusters ($J_{I-}J_{\alpha\beta}$, with $\alpha, \beta \in \{E, I\}$) (see **Supplementary Figure 2a** and **Supplementary Table 3** for network parameters).

Eight out of the 14 excitatory clusters were randomly selected and assigned "roles" to represent crucial task information: taste clusters (sucrose, quinine, maltose, and sucrose octaacetate), cue clusters (cue left and cue right), and action clusters (action left and action right). For each taste cluster, half of the neurons inside the cluster responded to additional external input when that particular stimulus was presented to the network. All presented stimuli had the functional form $f(t) = (G/h)(e^{-t/\tau_D} - e^{-t/\tau_R})$, where G is the maximum gain, τ_D and τ_R are the decay and rise time constants, respectively, of the function $f(t)$, $h = (\tau_R/\tau_D)^{\tau_R/(\tau_D-\tau_R)} - (\tau_R/\tau_D)^{\tau_D/(\tau_D-\tau_R)}$ is a normalization constant, and the value of the function determined the percentage increase in external current at time t for selective neurons in the appropriate cluster (i.e., $I_{i,\text{ext}}(t) = I_{i,\text{ext}}^* \times (1 + f(t)/100)$ for the constant baseline external current $I_{i,\text{ext}}^*$).

Unlike generic clusters (i.e., clusters not ascribed a role), only 25% of neurons in a taste cluster were strongly connected (via synapses $J_{E++}J_{EE}$ or $J_{I++}J_{II}$, see above), while the remaining 75% were less strongly connected ($J_{E+}J_{EE}$ or $J_{I+}J_{II}$, respectively for E and I clusters, with $J_{++} > J_+ > 1$; see **Supplementary Figure 2b**). The subcluster of taste neurons with strong interconnections was also strongly connected with the subcluster of neurons representing the same taste quality (sweet or bitter), i.e., the sucrose and the maltose subclusters were connected with synapses equal to $J_{E++}J_{EE}$ or $J_{I++}J_{II}$ (for E or I neurons, respectively), and the same was true for the quinine and sucrose octaacetate subclusters. The rest of the connections were as for generic clusters shown in **Supplementary Figure 2a**. As shown in **Supplementary Figure 2b (top)**, it may help to think of this structure as introducing overlaps between taste clusters that share the same taste quality (i.e., sweet or bitter).

Excitatory cue clusters had stronger synaptic connections from the appropriate E taste clusters. For example, 60% of the weights from sucrose and quinine clusters to the cue left cluster and 60% of the weights from maltose and sucrose octaacetate clusters to the cue right cluster were increased by 85% ($J_{CT,E}J_{E-}J_{EE}$, see **Supplementary Figure 3** and **Supplementary Table 3** for parameter values). To facilitate longer lasting representations of cue information in the model, cue clusters had 60% of their excitatory intracluster weights scaled up by 5% ($J_{CC,E}J_{E++}J_{EE}$), and their synaptic time constants ($\tau_{\text{syn,CueE}}$, $\tau_{\text{syn,CueI}}$) increased (2.3 times greater for excitatory clusters, 1.3 times greater for their inhibitory partners). To discourage simultaneous activation of both cue clusters, 50% of the connections from inhibitory partner clusters of one cue direction to excitatory clusters of the opposite cue direction ($J_{CC,I}J_{I-}J_{EI}$) were increased by 40%.

Excitatory action clusters had potentiated connections from E neurons in cue clusters, with a preference for the appropriate direction: 50% of the weights from, e.g., cue right to action left were multiplied by 2.60 ($J_{AC,Inc}J_{E-}J_{EE}$), while 50% of the weights from, e.g., cue left to action left were multiplied by 2.75 ($J_{AC,Cor}J_{E-}J_{EE}$). We encouraged action clusters, similar to cue clusters, to represent information over longer time scales by scaling up 50% of their intracluster weights by 7% ($J_{AA,E}J_{E++}J_{EE}$) and increasing their synaptic time constants ($\tau_{\text{syn,ActE}}$ and $\tau_{\text{syn,ActI}}$) 3.30 times for excitatory clusters and 3.25 times for their inhibitory partners. As done with cue clusters, we discouraged co-activation of action clusters by increasing the strength of 50% of the connections from inhibitory partner clusters of one action direction to excitatory clusters of the opposite action direction three-fold ($J_{AA,I}J_{I-}J_{EI}$). Finally, to promote a transfer of information from cue to action via negative feedback, we increased the strength of 50% of the connections from inhibitory partners of action clusters to excitatory neurons in cue clusters ($J_{CA,I}J_{I-}J_{EI}$) by 40%. The full synaptic matrix for the whole model network is shown in **Supplementary Figure 4**.

HMM analysis of the model simulations

For HMM analysis of model data, 10 different networks were used to mimic variability across sessions and/or subjects. The networks were obtained each time by sampling a new synaptic matrix (all having the same basic structure described earlier). Although each network had random connectivity, the in-degree connectivity from each presynaptic population was fixed across the 10 networks. This enforced some degree of uniformity in firing rates across clusters and minimized the presence of bias in the performance of each network.

A simulation of each network was run for 100 trials (25 trials per taste stimulus). For each trial, each neuron started with a random initial membrane potential (normally distributed with mean 0 and standard deviation 4) and received a constant external input current. At the start of the trial, the network was presented with one of the stimuli (which modified the external input current for select neurons) and evolved over 3 seconds. We then sampled one neuron per excitatory cluster, excluding those that fired at < 2 Hz on average whenever possible. An HMM was fitted to the spike trains of the sampled neurons following the same procedure used for the HMM analysis of experimental data. Trials were scored by inferring the model's selected direction from the firing rates of its action clusters. Cluster firing rates were calculated by binning spikes in 50 ms windows, and a cluster was considered "on" when its firing rate was at least 40% of its maximum observed over the 100-trial simulation (setting a constant threshold at 34 Hz yielded exactly the same results). A trial was deemed correct if the appropriate action cluster came on after 0.5 s after stimulus onset while the opposite action cluster did not. A trial was deemed incorrect if the opposite action cluster came on during this window and the correct one did not. For an ambiguous trial, in which neither action cluster came on or both action clusters came on, we assigned a decision randomly such that the expected accuracy was 0.5, but also excluded the trial from further analysis. We then used the scored trials to classify the coding properties of the hidden states from each HMM, following the same procedure used for the experimental data. Onset times of all coding states were pooled over the 10 models, and distributions of these onset times were compared for Quality-coding states, Cue-coding states, and Action-coding states.

Simulated optogenetic silencing experiments

The same 10 networks (characterized by the same 10 synaptic weight matrices used above) were subjected to various perturbations in a series of simulations meant to mimic experimental optogenetic silencing of cortical activity (Vincis et al., 2020). Silencing was implemented as a simple, square stimulus pulse to all inhibitory neurons in the network. The height of the square pulse quantified the percent increase in baseline external current to inhibitory neurons. As a function of the square pulse's center c, its duration Δ_{sil}, and its strength κ_{sil}, the formula for the square pulse in time t read:

$$\varphi(t; c, \Delta_{\text{sil}}, \kappa_{\text{sil}}) = \kappa_{\text{sil}} \times [\Theta(t - (c - \Delta_{\text{sil}}/2)) - \Theta(t - (c + \Delta_{\text{sil}}/2))],$$

where Θ is the Heaviside step function. We studied the effect of silencing on the networks' performance accuracy as a function of parameter values. We tested perturbation strengths of 25%, 50%, 75%, 100%, and 125% for a constant duration of 250 ms. We also tested how the effect of the 100%, 250 ms silencing depends on the dynamics of the stimulus, $f(t) = (G/h)(e^{-t/\tau_{\text{D}}} - e^{-t/\tau_{\text{R}}})$, by increasing τ_{D} from a baseline value of 160 ms to 280 ms, 400 ms, 455 ms, 520 ms, and 705 ms.

For each configuration of parameters $\{\kappa_{\text{sil}}, \Delta_{\text{sil}}, \tau_{\text{D}}\}$, we computed mean network accuracy as a function of the silencing center c by "sliding" the square pulse center along the trial window from 0 to 3 s in

increments of $\Delta c = 50$ ms. For each value of c, we simulated 100 trials for each of the 10 networks under the given parameter set, and then averaged the accuracy over the collated $1,000$ trials. Accuracy-versus-c curves were smoothed by upsampling via linear interpolation (point density between adjacent points increased from 2 to 100) and then convolving with a Gaussian filter (250 point width). Average initial slopes of accuracy-versus-c curves were computed over the first 0.5 s (where all curves could be well-approximated by lines) for the scatter plot in **Figure 6d**. For silencing centered at $c = 1.5$ s and with parameter set $\{\kappa_{\mathrm{sil}} = 100\%, \Delta_{\mathrm{sil}} = 250$ ms, $\tau_{\mathrm{D}} = 160$ ms$\}$, we computed the average firing rate of each cluster in the 250 ms baseline window before silencing (from $t = 1.125$ s to $t = 1.375$ s) and in the 250 ms silencing window (from $t = 1.375$ s to $t = 1.625$ s) in each trial. We computed the percent change in averaged firing rates for each cluster to obtain the points shown in **Figure 6b**. Because excitatory action clusters are prevented from receiving input due to the action gate during this baseline period, their baseline rates are 0 and their relative changes are, therefore, undefined and excluded.

Supplementary Information

Temporal progression along discrete coding states during decision-making in the mouse gustatory cortex

Liam Lang[1,2,3], Giancarlo La Camera[1,2,3,*], and Alfredo Fontanini[1,2,3,*]

[1]Graduate Program in Neuroscience, [2]Center for Neural Circuit Dynamics, and [3]Department of Neurobiology and Behavior

Stony Brook University

[*]Equal contribution and co-corresponding authors:

giancarlo.lacamera@stonybrook.edu, alfredo.fontanini@stonybrook.edu

HIDDEN STATES IN DATA

EXPERIMENT	Model fit to experimental data	Model fit to circularly shuffled data	Model fit to swap-shuffled data
Hidden states	Mean: 7 Median: 7 Range: $3-12$	Mean: 5.7 Median: 6 Range: $3-9$	Mean: 4.8 Median: 4 Range: $3-10$
Decoded states	Total: 90 Mean: 5.6 Median: 6 Range: $3-9$	Total: 67 Mean: 4.2 Median: 4 Range: $2-7$	Total: 48 Mean: 3 Median: 3 Range: $2-6$
Decision-coding states	18 (over 9 sessions)	13 (over 8 sessions)	12 (over 8 sessions)
Cue-coding states	7 (over 5 sessions)	8 (over 6 sessions)	5 (over 5 sessions)
Action-coding states	10 (over 6 sessions)	5 (over 4 sessions)	6 (over 4 sessions)
Quality-coding states	5 (over 4 sessions)	5 (over 5 sessions)	3 (over 3 sessions)
Taste ID-coding states	10 (over 6 sessions)	11 (over 7 sessions)	11 (over 8 sessions)
Dual-coding states	4 (over 3 sessions)	1	2 (over 2 sessions)
Non-coding states	53 (over 15 sessions)	37 (over 12 sessions)	20 (over 10 sessions)
Sessions with Quality- and Decision-coding states	2	1	2
Sessions with Cue- and Action-coding states	3	2	1
Sessions with Quality-, Cue-, and Action-coding states	1	1	0

Supplementary Table 1 Summary of the numbers of states found by HMM models fit to unshuffled, circularly shuffled, and swap-shuffled experimental data. Also indicated is the distribution of these states over sessions (16 sessions total). The same coding state classification procedure (see **Supplementary Figure 1**) was applied to the trial-by-trial decoding results from each model. "Hidden states" is the theoretical best number determined from model fitting; "Decoded states" is how many were actually found after decoding trial-by-trial.

Hidden states in model

SIMULATION	Model fit to simulation data	Model fit to circularly shuffled data	Model fit to swap-shuffled data
Hidden states	Mean: 24.3 Median: 25.5 Range: 17 − 29	Mean: 21.8 Median: 23 Range: 14 − 27	Mean: 12.4 Median: 12 Range: 8 − 18
Decoded states	Total: 194 Mean: 19.4 Median: 19.5 Range: 16 − 23	Total: 185 Mean: 18.5 Median: 19 Range: 13 − 22	Total: 95 Mean: 9.5 Median: 9 Range: 6 − 15
Decision-coding states	85 (over 10 sessions)	78 (over 10 sessions)	4 (over 3 sessions)
Cue-coding states	57 (over 10 sessions)	52 (over 10 sessions)	1
Action-coding states	16 (over 9 sessions)	18 (over 8 sessions)	1
Quality-coding states	21 (over 10 sessions)	4 (over 4 sessions)	1
Taste ID-coding states	43 (over 10 sessions)	50 (over 10 sessions)	47 (over 10 sessions)
Dual-coding states	0	0	1
Non-coding states	45 (over 10 sessions)	53 (over 10 sessions)	42 (over 10 sessions)
Sessions with Quality- and Decision-coding states	10	4	0
Sessions with Cue- and Action-coding states	9	8	0
Sessions with Quality-, Cue-, and Action-coding states	9	3	0

Supplementary Table 2   Summary of numbers of states found by HMM models fit to unshuffled, circularly shuffled, and swap-shuffled simulation data. Also indicated is the distribution of these states over sessions (10 sessions total). The same coding state classification procedure (see **Supplementary Figure 1**) was applied to the trial-by-trial decoding results from each model. "Hidden states" is the theoretical best number determined from model fitting; "Decoded states" is how many were actually found after decoding trial-by-trial.

Model Parameters

Symbol	Value	Name/Description
NEURONS AND NETWORK MODEL		
V_{L}	0 mV	Resting membrane potential
V_{th}	20 mV	Spiking threshold
V_{r}	0 mV	Reset membrane potential
C	1 pF	Membrane capacitance
$\tau_{\mathrm{m,E}}$	20 ms	E membrane time constant
$\tau_{\mathrm{m,I}}$	10 ms	I membrane time constant
$\tau_{\mathrm{syn,E}}$	3 ms	E synaptic time constant
$\tau_{\mathrm{syn,CueE}}$	6.9 ms	E synaptic time constant for neurons in cue clusters
$\tau_{\mathrm{syn,ActE}}$	9.9 ms	E synaptic time constant for neurons in action clusters
$\tau_{\mathrm{syn,I}}$	2 ms	I synaptic time constant
$\tau_{\mathrm{syn,CueI}}$	2.6 ms	I synaptic time constant for neurons in cue clusters
$\tau_{\mathrm{syn,ActI}}$	6.5 ms	I synaptic time constant for neurons in action clusters
τ_{r}	5 ms	Absolute refractory period
N_{e}	4,000	Number of E neurons
N_{i}	994	Number of I neurons
$I_{\mathrm{E,ext}}^{*}$	2.05 pA	Baseline external input current for E neurons
$I_{\mathrm{I,ext}}^{*}$	2.16 pA	Baseline external input current for I neurons
Q	14	Number of clusters
f_{bg}	0.125	Fraction of E neurons in "background" population
f_{ov}	0.25	Fraction of neurons in E and I taste clusters assigned to "overlapping" subpopulations
t_{Total}	3 s	Trial simulation length
dt	0.05 ms	Simulation time step
P_{ee}	0.2	Probability of E $\rightarrow$ E connection
P_{ei}	0.5	Probability of I $\rightarrow$ E connection
P_{ie}	0.5	Probability of E $\rightarrow$ I connection
P_{ii}	0.5	Probability of I $\rightarrow$ I connection
J_{EE}	0.16 pA×ms	E→E synaptic weight (reference value)
J_{EI}	−0.63 pA×ms	I→E synaptic weight (reference value)
J_{IE}	0.23 pA×ms	E→I synaptic weight (reference value)
J_{II}	1.2 pA×ms	I→I synaptic weight (reference value)
$J_{\mathrm{E++}}$	12	E→E maximum intracluster potentiation factor
$J_{\mathrm{E+}}$	$0.25 \times J_{\mathrm{E-}} + 0.75 \times J_{\mathrm{E++}}$	E→E intermediate intracluster potentiation factor
$J_{\mathrm{E-}}$	0.267	E→E generic depression factor for synaptic weights between neurons in different clusters
$J_{\mathrm{I++}}$	5.4	I→E, E→I, I→I maximum intracluster potentiation factor
$J_{\mathrm{I+}}$	$0.25 \times J_{\mathrm{I-}} + 0.75 \times J_{\mathrm{E++}}$	I→E, E→I, I→I intermediate potentiation factor
$J_{\mathrm{I-}}$	0.685	I→E, E→I, I→I generic depression factor for synaptic weights between neurons in different clusters
$J_{\mathrm{CT,E}}$; $P_{\mathrm{CT,E}}$	1.85 ; 0.60	Weight modifier for connections from E taste cluster to appropriate E cue cluster; Probability of applying modifier
$J_{\mathrm{CC,E}}$; $P_{\mathrm{CC,E}}$	1.04 ; 0.60	Weight modifier for connections from E cue cluster to itself; Probability of applying modifier
$J_{\mathrm{CC,I}}$; $P_{\mathrm{CC,I}}$	1.40 ; 0.50	Weight modifier for connections from I cue cluster to E opposite cue cluster; Probability of applying modifier

Symbol	Value	Name/Description
\multicolumn{3}{center}{*Continuation of Supplementary Table 3*}		
$J_{\text{AC,Corr}}$; $P_{\text{AC,Corr}}$	2.75 ; 0.50	Weight modifier for connections from E cue cluster to appropriate ("correct") E action cluster; Probability of applying modifier
$J_{\text{AC,Inc}}$; $P_{\text{AC,Inc}}$	2.60 ; 0.50	Weight modifier for connections from E cue cluster to inappropriate ("incorrect") E action cluster; Probability of applying modifier
$J_{\text{CA,I}}$; $P_{\text{CA,I}}$	1.40 ; 0.50	Weight modifier for connections from I action clusters to E cue clusters; Probability of applying modifier
$J_{\text{AA,E}}$; $P_{\text{AA,E}}$	1.07 ; 0.50	Weight modifier for connections from E action cluster to itself; Probability of applying modifier
$J_{\text{AA,I}}$; $P_{\text{AA,I}}$	3.00 ; 0.50	Weight modifier for connections from I action cluster to E opposite action cluster; Probability of applying modifier
STIMULUS		
f_{stim}	0.5	Fraction of neurons in taste-selective clusters that respond to external stimulus
G	60%	Maximum stimulus gain (external current modifier)
τ_{R}	150 ms	Stimulus rise time constant
τ_{D}	160 ms (but varied up to 705 ms for silencing experiment)	Stimulus decay time constant
ACTION GATE		
μ_{ag}	1.5 s	Action gate mean start time
η_{ag}	0.4 s	Halfwidth of uniformly random action gate start time
Δ_{ag}	0.5 s	Time for action gate to increase from 0 to 1
SILENCING PERTURBATION		
κ_{sil}	25% − 125%	Silencing strength (external current modifier)
Δ_{sil}	250 ms	Silencing duration
Δc	50 ms	Increment for sliding center of silencing window

Supplementary Table 3 Model parameters. Symbols, values, and brief descriptions for all parameters used in the simulations of the spiking network model presented in the main text.

Supplementary Figure 1 Pipeline for classification of HMM states. Given a state S, the probability of observing S in a correctly performed trial was compared across the four trial types using a Chi-squared test. If the result was not significant (N), S was labeled "Non-coding." Otherwise (Y), all six pairwise Marascuilo post-hoc tests were performed. If there was exactly one trial type for which all post-hoc tests involving it were significant (Y), S was labeled "Taste ID-coding." Otherwise (N), the four trial types were grouped in two orthogonal ways: sweet (sucrose and maltose) and bitter (quinine and sucrose octaacetate), and cue left (sucrose and quinine) and cue right (maltose and sucrose octaacetate). Two additional Chi-squared tests were run: one to compare the probability of observing S in correctly performed trials between the sweet and bitter categories, and another to compare the probability of observing S in correctly performed trials between the cue left and cue right categories. These two tests are run independently, leading to four possible significance outcomes. If neither test was significant, S was labeled "Non-coding." If both tests were significant, S was labeled "Dual-coding." If the sweet vs. bitter test was significant and the cue left vs. cue right test was not, S was labeled "Quality-coding." If the sweet vs. bitter test was not significant and the cue left vs. cue right test was, S was labeled "Decision-coding," but could be further classified by examining its frequency of occurrence in correctly vs. incorrectly performed trials. If the cued direction for which S was more likely to be observed in correctly performed trials was the same for incorrectly performed trials (referred to as the "preferred direction" for S in correct and incorrect trials), S was labeled "Cue-coding." If the preferred directions in correct and incorrect trials were opposite, S was labeled "Action-coding."

Supplementary Figure 2 Synaptic structure for taste clusters. **a**: Generic synaptic structure between clusters without task roles. **Top**: Cartoon representing a pair of E-I partner clusters with their synaptic connection weights (red, flathead arrows: inhibitory connections; black, pointed arrows: excitatory connections). **Bottom**: synaptic matrix for the generic pair of E-I partner clusters (colormap corresponding to units of pA×ms). Given any two excitatory clusters, E1 and E2, and their inhibitory partners, I1 and I2, synaptic connections were strong between neurons within the same cluster or belonging to partner clusters ($J_{E++}J_{EE}$ or $J_{I++}J_{\alpha\beta}$, with $\alpha, \beta \in \{E, I\}$), and weak between neurons in different clusters ($J_{E-}J_{EE}$ or $J_{I-}J_{\alpha\beta}$, with $\alpha, \beta \in \{E, I\}$). **b**: **Top**: Venn diagram depicting the "overlapping" structure of two E-I taste cluster pairs with the same quality (sweet or bitter). **Bottom**: synaptic matrix of "overlapping" taste clusters depicted at top (colormap corresponding to units of pA×ms). Taste neurons in different clusters with the same taste quality had strongly connected overlapping subclusters (E1∩E2 and I1∩I2 in the Venn diagram), with synaptic strength $J_{E++}J_{EE}$ or $J_{I++}J_{\alpha\beta}$, with $\alpha, \beta \in \{E, I\}$. These overlapping subclusters are shown in the bottom right corner of each taste cluster in the synaptic matrix. Taste neurons in the same cluster but outside the overlapping subclusters were connected with synaptic strength $J_{E+}J_{EE}$ or $J_{I+}J_{\alpha\beta}$, with $\alpha, \beta \in \{E, I\}$. All other neurons were connected as in the generic clusters of panel **a**, with $J_{E-} < 1 < J_{E+} < J_{E++}$ and $J_{I-} < 1 < J_{I+} < J_{I++}$.

Supplementary Figure 3 Details of synaptic connections among taste, cue, and action E-I cluster pairs (compare with main **Figure 3b**). The main synaptic connections are depicted by curved lines (pointed arrowheads are excitatory connections, flat arrowheads are inhibitory connections). The strength of each connection is indicated next to the corresponding curved line (see **Supplementary Table 3** for parameter values). Synaptic weights from excitatory taste clusters to appropriate excitatory cue clusters (i.e., S → CL, M → CR, Q → CL, and O → CR) were larger by a factor $J_{CT,E}$ compared to generic weights between different clusters (e.g., S → CR, M → CL, Q → CR, and O → CL). Excitatory cue clusters had stronger intracluster connections (by a factor $J_{CC,E}$) compared to generic clusters; the synaptic weights from an inhibitory cue cluster to the opposite excitatory cue cluster were larger by a factor $J_{CC,I}$. Connections from excitatory cue clusters to correct excitatory action clusters (i.e., CL → AL and CR → AR) were larger by a factor $J_{AC,Cor}$, while the connections to incorrect excitatory action clusters (i.e., CL → AR and CR → AL) were larger by a factor $J_{AC,Inc}$, with $J_{AC,Cor} > J_{AC,Inc}$. Excitatory action clusters had stronger intracluster connections (scaled up by $J_{AA,E}$) and the weights from their inhibitory partners to the opposite excitatory action cluster were scaled up by $J_{AA,I}$. Connections from inhibitory action clusters to excitatory cue clusters were magnified by $J_{CA,I}$. All synaptic weight modifiers were applied with a given probability (reported in **Supplementary Table 3**). Connections among clusters not explicitly depicted here were as in **Supplementary Figure 2** (also see **Supplementary Figure 4** for the full synaptic weight matrix). Key: S: sucrose, M: maltose, Q: quinine, O: octaacetate, CL: cue left, CR: cue right, AL: action left, AR: action right.

Full synaptic matrix

Presynaptic

Post \ Pre	S	Q	M	O	CL	CR	AL	AR	9	10	11	12	13	14	B	S	Q	M	O	CL	CR	AL	AR	9	10	11	12	13	14
S	0.296	0.009	0.033	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	-1.370	-0.211	-0.307	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211
Q	0.009	0.296	0.009	0.033	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	-0.211	-1.370	-0.211	-0.307	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211
M	0.033	0.009	0.296	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	-0.307	-0.211	-1.370	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211
O	0.009	0.033	0.009	0.296	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	-0.211	-0.307	-0.211	-1.370	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211
CL	0.013	0.013	0.009	0.009	0.396	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	-0.211	-0.211	-0.211	-0.211	-1.725	-0.254	-0.253	-0.254	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211
CR	0.009	0.009	0.013	0.013	0.009	0.396	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	-0.211	-0.211	-0.211	-0.211	-0.254	-1.725	-0.254	-0.254	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211
AL	0.009	0.009	0.009	0.009	0.016	0.015	0.397	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-1.725	-0.423	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211
AR	0.009	0.009	0.009	0.009	0.015	0.016	0.009	0.397	0.009	0.009	0.009	0.009	0.009	0.009	0.009	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.423	-1.725	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211
9	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.384	0.009	0.009	0.009	0.009	0.009	0.009	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-1.725	-0.211	-0.211	-0.211	-0.211	-0.211
10	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.384	0.009	0.009	0.009	0.009	0.009	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-1.725	-0.211	-0.211	-0.211	-0.211
11	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.384	0.009	0.009	0.009	0.009	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-1.725	-0.211	-0.211	-0.211
12	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.384	0.009	0.009	0.009	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-1.725	-0.211	-0.211
13	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.384	0.009	0.009	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-1.725	-0.211
14	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.384	0.009	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-1.725
B	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211	-0.211
S (I)	0.494	0.076	0.111	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	-2.611	-0.403	-0.585	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403
Q (I)	0.076	0.494	0.076	0.111	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	-0.403	-2.611	-0.403	-0.585	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403
M (I)	0.111	0.076	0.494	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	-0.585	-0.403	-2.611	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403
O (I)	0.076	0.111	0.076	0.494	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	-0.403	-0.585	-0.403	-2.611	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403
CL (I)	0.076	0.076	0.076	0.076	0.621	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	-0.403	-0.403	-0.403	-0.403	-3.286	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403
CR (I)	0.076	0.076	0.076	0.076	0.076	0.621	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	-0.403	-0.403	-0.403	-0.403	-0.403	-3.286	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403
AL (I)	0.076	0.076	0.076	0.076	0.076	0.076	0.621	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-3.286	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403
AR (I)	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.621	0.076	0.076	0.076	0.076	0.076	0.076	0.076	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-3.286	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403
9 (I)	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.621	0.076	0.076	0.076	0.076	0.076	0.076	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-3.286	-0.403	-0.403	-0.403	-0.403	-0.403
10 (I)	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.621	0.076	0.076	0.076	0.076	0.076	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-3.286	-0.403	-0.403	-0.403	-0.403
11 (I)	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.621	0.076	0.076	0.076	0.076	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-3.286	-0.403	-0.403	-0.403
12 (I)	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.621	0.076	0.076	0.076	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-3.286	-0.403	-0.403
13 (I)	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.621	0.076	0.076	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-3.286	-0.403
14 (I)	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.076	0.621	0.076	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-0.403	-3.286

Postsynaptic

Supplementary Figure 4 Full synaptic weight matrix of the network model. The value in a block occupying row i and column j is the *average* synaptic weight (in units of pA×ms) from a presynaptic neuron belonging to cluster j to a postsynaptic neuron belonging to cluster i. The average synaptic values shown in the table were obtained (up to rounding) as: $P_{\alpha\beta}[f_{\mathrm{ov}}^2 J_{\gamma++} + (1-f_{\mathrm{ov}}^2) J_{\gamma X}] J_{\alpha\beta}$ for connections involving taste clusters with the same taste quality (sweet or bitter), and $P_{\alpha\beta}(P_M J_M + 1 - P_M) J_{\gamma Y} J_{\alpha\beta}$ for all other connections. Here $\alpha, \beta, \gamma \in \{E, I\}$, $\gamma = E$ if and only if $\alpha = \beta = E$, $X \in \{-, +\}$, $Y \in \{-, ++\}$, $J_M \in \{J_{\mathrm{CT,E}}, J_{\mathrm{CC,E}}, J_{\mathrm{CC,I}}, J_{\mathrm{AC,Cor}}, J_{\mathrm{AC,Inc}}, J_{\mathrm{AA,E}}, J_{\mathrm{AA,I}}, J_{\mathrm{CA,I}}\}$ is a weight modifier, and $P_M \in \{0, P_{\mathrm{CT,E}}, P_{\mathrm{CC,E}}, P_{\mathrm{CC,I}}, P_{\mathrm{AC,Cor}}, P_{\mathrm{AC,Inc}}, P_{\mathrm{AA,E}}, P_{\mathrm{AA,I}}, P_{\mathrm{CA,I}}\}$ is the corresponding probability of applying the modifier ($P_M = 0$ when modifier was not applied). Key: S: sucrose cluster, M: maltose cluster, Q: quinine cluster, O: octaacetate cluster, CL: cue left cluster, CR: cue right cluster, AL: action left cluster, AR: action right cluster, 9-14: clusters without task roles, B: background (excitatory) population.

References

Abeles M, Bergman H, Gat I, Meilijson I, Seidemann E, Tishby N, Vaadia E (1995) Cortical activity flips among quasi-stationary states. *Proc. Natl. Acad. Sci. U. S. A.* 92: 8616–8620.

Benozzo D, La Camera G, Genovesio A (2021) Slower prefrontal metastable dynamics during deliberation predicts error trials in a distance discrimination task. *Cell Rep.* 35: 108934.

Bollimunta A, Totten D, Ditterich J (2012) Neural Dynamics of Choice: Single-Trial Analysis of Decision-Related Activity in Parietal Cortex. *Journal of Neuroscience* 32: 12684–12701.

Bouaichi CG, Vincis R (2020) Cortical processing of chemosensory and hedonic features of taste in active licking mice. *J. Neurophysiol.* 123: 1995–2009.

Brinkman BAW, Yan H, Maffei A, Park IM, Fontanini A, Wang J, La Camera G (2022) Metastable dynamics of neural circuits and networks. *Appl Phys Rev* 9: 011313.

Curti E, Mongillo G, La Camera G, Amit DJ (2004) Mean field and capacity in realistic networks of spiking neurons storing sparsely coded random memories. *Neural Comput.* 16: 2597–2637.

Deco G, Hugues E (2012) Neural network mechanisms underlying stimulus driven variability reduction. *PLoS Comput. Biol.* 8: e1002395.

Dikecligil GN, Graham DM, Park IM, Fontanini A (2020) Layer- and Cell Type-Specific Response Properties of Gustatory Cortex Neurons in Awake Mice. *J. Neurosci.* 40: 9676–9691.

Durstewitz D, Vittoz NM, Floresco SB, Seamans JK (2010) Abrupt transitions between prefrontal neural ensemble states accompany behavioral transitions during rule learning. *Neuron* 66: 438–448.

Fonseca E, de Lafuente V, Simon SA, Gutierrez R (2018) Sucrose intensity coding and decision-making in rat gustatory cortices. *eLife* 7: e41152.

Fontanini A, Katz DB (2006) State-dependent modulation of time-varying gustatory responses. *J. Neurophysiol.* 96: 3183–3193.

Gardner MPH, Fontanini A (2014) Encoding and tracking of outcome-specific expectancy in the gustatory cortex of alert rats. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 34: 13000–13017.

Gat I, Tishby N (1993) Statistical modeling of cell-assemblies activities in associative cortex of behaving monkeys In *Neural Information Process. Syst.*, Vol. 5, pp. 945–53.

Grill HJ, Norgren R (1978) The taste reactivity test. I. Mimetic responses to gustatory stimuli in neurologically normal rats. *Brain Res.* 143: 263–279.

Guo ZV, Li N, Huber D, Ophir E, Gutnisky D, Ting JT, Feng G, Svoboda K (2014) Flow of cortical activity underlying a tactile decision in mice. *Neuron* 81: 179–194.

Jezzini A, Mazzucato L, La Camera G, Fontanini A (2013) Processing of hedonic and chemosensory features of taste in medial prefrontal and insular networks. *J. Neurosci.* 33: 18966–18978.

Jones LM, Fontanini A, Sadacca BF, Miller P, Katz DB (2007) Natural stimuli evoke dynamic sequences of states in sensory cortical ensembles. *Proc. Natl. Acad. Sci. U. S. A.* 104: 18772–18777.

Katz DB, Simon SA, Nicolelis MA (2001) Dynamic and multimodal responses of gustatory cortical neurons in awake rats. *J. Neurosci.* 21: 4478–4489.

Katz DB, Nicolelis MAL, Simon SA (2002) Gustatory processing is dynamic and distributed. *Curr. Opin. Neurobiol.* 12: 448–454.

La Camera G, Fontanini A, Mazzucato L (2019) Cortical computations via metastable activity. *Current Opinion in Neurobiology* 58: 37–45.

Lin JY, Mukherjee N, Bernstein MJ, Katz DB (2021) Perturbation of amygdala-cortical projections reduces ensemble coherence of palatability coding in gustatory cortex. *Elife* 10.

Litwin-Kumar A, Doiron B (2012) Slow dynamics and high variability in balanced cortical networks with clustered connections. *Nat. Neurosci.* 15: 1498–1505.

Livneh Y, Ramesh RN, Burgess CR, Levandowski KM, Madara JC, Fenselau H, Goldey GJ, Diaz VE, Jikomes N, Resch JM, Lowell BB, Andermann ML (2017) Homeostatic circuits selectively gate food cue responses in insular cortex. *Nature* 546: 611–616.

Maboudi K, Ackermann E, de Jong LW, Pfeiffer BE, Foster D, Diba K, Kemere C (2018) Uncovering temporal structure in hippocampal output patterns. *Elife* 7.

Mazzucato L, Fontanini A, La Camera G (2015) Dynamics of Multistable States during Ongoing and Evoked Cortical Activity. *Journal of Neuroscience* 35: 8214–8231.

Mazzucato L, La Camera G, Fontanini A (2019) Expectation-induced modulation of metastable activity underlies faster coding of sensory stimuli. *Nat. Neurosci.* 22: 787–796.

Mazzucato L, Fontanini A, La Camera G (2016) Stimuli Reduce the Dimensionality of Cortical Activity. *Front. Syst. Neurosci.* 10: 11.

Miller P, Katz DB (2010) Stochastic transitions between neural states in taste processing and decision-making. *J. Neurosci.* 30: 2559–2570.

Moran A, Katz DB (2014) Sensory cortical population dynamics uniquely track behavior across learning and extinction. *J. Neurosci.* 34: 1248–1257.

Mukherjee N, Wachutka J, Katz DB (2019) Impact of precisely-timed inhibition of gustatory cortex on taste behavior depends on single-trial ensemble dynamics. *Elife* 8.

Piette CE, Baez-Santiago MA, Reid EE, Katz DB, Moran A (2012) Inactivation of basolateral amygdala specifically eliminates palatability-related information in cortical sensory responses. *J. Neurosci.* 32: 9981–9991.

Ponce-Alvarez A, Nácher V, Luna R, Riehle A, Romo R (2012) Dynamics of cortical neuronal ensembles transit from decision making to storage for later report. *J. Neurosci.* 32: 11956–11969.

Prins J (2013) *NIST/SEMATECH e-Handbook of Statistical Methods.* https://www.itl.nist.gov/div898/handbook/ [Accessed June 29, 2022].

Rabiner LR (1989) A tutorial on hidden Markov models and selected applications in speech recognition. *Proc. IEEE* 77: 257–286.

Radons G, Becker JD, Dülfer B, Krüger J (1994) Analysis, classification, and coding of multielectrode spike trains with hidden Markov models. *Biol. Cybern.* 71: 359–373.

Recanatesi S, Pereira-Obilinovic U, Murakami M, Mainen Z, Mazzucato L (2022) Metastable attractors explain the variable timing of stable behavioral action sequences. *Neuron* 110: 139–153.E9.

Rich EL, Wallis JD (2016) Decoding subjective decisions from orbitofrontal cortex. *Nat. Neurosci.* 19: 973–980.

Rostami V, Rost T, Riehle A, van Albada SJ, Nawrot MP (2020) Spiking neural network model of motor cortex with joint excitatory and inhibitory clusters reflects task uncertainty, reaction times, and variability dynamics. *bioRxiv* 2020.02.27.968339.

Sadacca BF, Mukherjee N, Vladusich T, Li JX, Katz DB, Miller P (2016) The Behavioral Relevance of Cortical Neural Ensemble Responses Emerges Suddenly. *J. Neurosci.* 36: 655–669.

Sadacca BF, Rothwax JT, Katz DB (2012) Sodium concentration coding gives way to evaluative coding in cortex and amygdala. *J. Neurosci.* 32: 9999–10011.

Samuelsen CL, Gardner MPH, Fontanini A (2012) Effects of cue-triggered expectation on cortical processing of taste. *Neuron* 74: 410–422.

Samuelsen CL, Gardner MPH, Fontanini A (2013) Thalamic contribution to cortical processing of taste and expectation. *J. Neurosci.* 33: 1815–1827.

Seidemann E, Meilijson I, Abeles M, Bergman H, Vaadia E (1996) Simultaneously recorded single units in the frontal cortex go through sequences of discrete and stable states in monkeys performing a delayed localization task. *J. Neurosci.* 16: 752–768.

Stapleton JR, Lavine ML, Nicolelis MAL, Simon SA (2007) Ensembles of gustatory cortical neurons anticipate and discriminate between tastants in a single lick. *Front. Neurosci.* 1: 161–174.

Stokes MG, Kusunoki M, Sigala N, Nili H, Gaffan D, Duncan J (2013) Dynamic coding for cognitive control in prefrontal cortex. *Neuron* 78: 364–375.

Vincis R, Chen K, Czarnecki L, Chen J, Fontanini A (2020) Dynamic Representation of Taste-Related Decisions in the Gustatory Insular Cortex of Mice. *Curr. Biol.* 30: 1834–1844.e5.