Method article

Pitfalls in methods to study colocalization of nanoparticles in mouse macrophage
lysosomes

Aura Maria Moreno-Echeverri[1,†], Eva Susnik[1,†], Dimitri Vanhecke[1], Patricia Taladriz-Blanco[3],

Sandor Balog[1], Alke Petri-Fink[1,2], Barbara Rothen-Rutishauser[1*]

* Correspondence: barbara.rothen@unifr.ch; Tel.: +41 26 300 95 02

[†] **Equal contributor**

Graphical Abstract

Abstract

Background: In the field of nanoscience there is an increasing interest to follow dynamics of
nanoparticles (NP) in cells with an emphasis on endo-lysosomal pathways and long-term NP fate.
The accuracy of fluorescence microscopy methods has an important role in obtaining insights into
NP interactions with lysosomes at the single cell level including quantification of NP uptake in a
specific cell type. During our research on this topic, we encountered several pitfalls, which can
bias the experimental outcome. We address some of these pitfalls and suggest possible solutions.

Methods: Here we use J774A.1 cells as a model for professional phagocytes. We expose them
to fluorescently-labelled amorphous silica NP with different sizes and quantify the colocalization
of fluorescently-labelled NP with lysosomes over time. We focus on confocal laser scanning
microscopy (CLSM), due to its ability to obtain 3D spatial information and follow live cell imaging
to study NP colocalization with lysosomes.

Results: We evaluate different experimental parameters that can bias the colocalization
coefficients (i.e, Pearson's and Manders'), such as the interference of phenol red in the cell culture
medium with the fluorescence intensity and image post-processing (effect of spatial resolution,
optical slice thickness, pixel saturation and bit depth). Additionally, we determine the correlation
coefficients for NP entering the lysosomes under four different experimental set-ups. First, we
found out that not only Pearsons', but also Manders' correlation coefficient should be considered
in lysosome-NP colocalization studies; second, there is a difference in NP colocalization when
using NP of different sizes and fluorescence dyes and last, the correlation coefficients might
change depending on live-cell and fixed-cell imaging set-up.

Conclusions: The results summarize detailed steps and recommendations for the experimental
design, staining, sample preparation and imaging to improve the reproducibility of colocalization
studies between the NP and lysosomes.

Keywords: nanoparticles; lysosomes; LysoTracker probes; colocalization; macrophages; pitfalls

1. Background

The use of engineered nanoparticles (NP) for biomedical applications requires a detailed
understanding of the NP-cell interaction, NP uptake and intracellular fate. Upon deposition at the
outer cell membrane, the NP are internalized via a variety of endocytic mechanisms[1–5] and pass
through an endosomal sorting process. Finally, depending on their properties, NP localize in the
subcellular compartments, including lysosomes, endoplasmic reticulum, Golgi apparatus,
mitochondria and nucleus[6,7] or bypass these compartments and end up in the cytoplasm.[8]
Lysosomes are dynamic cellular catabolic organelles responsible for the degradation of
macromolecules[9] and play a key role in the accumulation of internalized materials such as NP.[10,11]
Lysosome dysfunction is associated with multiple diseases, such as cancer, inflammatory and
neurodegenerative diseases, and Tay Sachs disease.[12–14] Therefore, lysosomes are a target for
a number of therapeutics[15,16] and there are several methods available to study the intracellular
fate and localization of NP in lysosomes.[17] Most recognized methods are transmission electron
microscopy,[18] fluorescence microscopy i.e., conventional bright field,[19] confocal,[20] and super-
resolution microscopy,[21] correlative microscopy,[22] focused ion beam-scanning electron
microscopy (FIB-SEM)[23] and lysosomal isolation.[24] The choice of each method strongly depends
on NP physicochemical properties (i.e., core material, size, shape, surface charge, porosity,
surface roughness, surface area, crystalline structure, etc.)[25] and complexity of the intracellular
environment.[26] In this study, we focused on confocal laser scanning microscopy (CLSM), due to
the possibility of imaging fluorescently labelled NP, fast sample preparation, obtaining 3D spatial
information without the need for physical sectioning of the sample, live cell imaging, and the
possibility to obtain information about NP colocalization with cellular compartments.

Different approaches are available to visualize lysosomes within live and fixed cells.[27] The most
used are the LysoTracker probes, Lysosomal-Associated Membrane Protein (LAMP) antibodies[28]
and cells transfected with fluorescent constructs targeting sequence for LAMP.[29] Here, the focus
is on the LysoTracker probes due to the possibility of dynamic lysosome imaging in unfixed cells,
low phototoxicity, and ease of use for long term imaging. Considering its many advantages, this
approach is widely used staining in research.[28,30–33] LysoTracker probes contain a conjugated
multi-pyrrole ring structure with a weakly basic amine, which gets protonated and accumulates in
the acidic environment of the lysosomes and late endosomes.[34] LysoTracker probes has some

drawbacks: **i)** Low pH is not unique as late endosomes can have a similar pH as lysosomes (~ 4.5

- 6.5) end, **ii)** Fluorophore instability[26], e.g. LysoTracker Red can undergo photoconversion to a

green-emitting fluorophore,[34,35] **iii)** Variability in intracellular NP trafficking between individual cells

(the time for NP to internalize into lysosomes can range from 15 min to several hours),[20] **iv)**

LysoTracker probes and LAMP signal miscorrelation may occur in some cell lines.[28]

Given these challenges, further experimental and observation-based biases must be avoided to

provide more reliable data interpretation. Moreover, there are other experimental factors that can

lead to misinterpretation of biological results, particularly those that rely on fluorescence and

image analysis of live or fixed cells.

Colocalization analysis is a common approach to evaluate NP trafficking and accumulation in

lysosomes.[36] Based on optical sections recorded by CLSM, colocalization analysis requires

statistical methods. The data consists of two channels, such as i) the lysosome visualized by

LysoTracker and ii) the fluorescently labelled NP. Fluorescence intensity data is obtained from

each recorded pixel between the LysoTracker channel and fluorescently labelled NPs. Such

analysis provides information about co-occurrence (i.e., the extent of spatial overlap between the

two fluorophores), as assessed by Manders' coefficients[37] and cross-correlation (the functional

relationship between the signal intensities of the two fluorophores), as described by Pearson's

correlation coefficient (PCC).[38] Additional measures and tests may include the Spearmann's Rank

Coefficient[39], Costes' randomization approach[40], Intensity Correlation Quotient[41] or various object-

based measures.[42] For accurate data interpretation, a combination of multiple metrics is

recommended. However, such sound interpretations are only achievable if biases in data

gathering are avoided. In our microscopic investigations on intracellular trafficking of NP, in

particular to the lysosomes, we encountered several methodical and technical pitfalls when using

probes to label lysosomes, each leading to misinterpretation and bias of the lysosomal labelling.

Here, we describe the most common pitfalls and provide recommendations on how to circumvent

them to yield accurate conclusions.

2. Results and discussion

The purpose of the present study is to determine the necessary steps for reliable colocalization of

NP within lysosomes with a focus on CLSM. J774A.1 mouse macrophages were used as a cell

model for professional phagocytes.[43] Cells were exposed to amorphous silica NP because they

can be synthesized in different sizes and labelled with a variety of different fluorophores, allowing
their tracking in living cells.

2.1. Fluorescent lysosomal probe characteristics

When designing an experiment, the key experimental parameters (i.e. LysoTracker probe
selection and the stability of fluorophores as well as equipment available) should be taken into
consideration. Since lysosomes were visualized by labeling the cells with LysoTracker probes, we
first extracted the main characteristics of LysoTracker probes (Table 1). Experiments must be
designed in a way to minimize fluorescence cross-talk (i.e., overlap of fluorochromes' excitation
spectra) and bleed-through (i.e overlap of fluorochromes' emission spectra), either by using a free
online spectral analyzer (available by Thermo Fisher Scientific) or experimentally.[44]

Table 1. Main LysoTracker probes characteristics and experimental parameters.

LysoTracker*	Excitation (nm)[1]	Emission (nm)[1]	Recommended concentration (nM)	Incubation time (37°C)	Ref.
Green (DND-26)	504	511	50	15-30 min	[45–47]
Blue (DND-22)	373	422	50-100	30 min – 1.5h	[48–50]
Red (DND-99)	577	590	10-50	15 min – 4h	[20,31,51]

112   1   Wavelength. The information in this table refers to LysoTracker probes purchased from ThermoFisher Scientific.

2.2. Influence of phenol red in cell culture medium on fluorescence intensity of LysoTracker probes

Our first attempt was to investigate whether the presence of phenol red, a routinely added pH
indicator, in the culture medium influences the fluorescence intensity of the red, green and blue
LysoTracker probes. Cells were grown either in complete Roswell Park Memorial
Institute medium, supplemented with 10% fetal bovine serum (FBS), 1 % (v/v) L-glutamine and 1
% (v/v) penicillin/streptomycin (cRPMI) with or without phenol red. After 24 h of incubation, the
cells were exposed to LysoTracker probes and the effect was monitored under continuous live cell
imaging. A representative image of an individual cell for each LysoTracker probe can be found in
(Figure SI1). Our results show that the presence of phenol red decreases the relative signal
intensity and biases the fluorescent signal as demonstrated by the significant differences in the
raw integrated density (i.e., sum of pixel values) of LysoTracker probes **(Figure 1)**. This distorts
the correlation coefficients and the interpretation of the colocalization results. The phenol red emits
light in the blue and red region of the spectrum, causing mainly interference with the blue and red

LysoTracker probes spectra **(Figure SI2**). On the other hand, no significant impact of phenol red
was detected on the intensity of LysoTracker Green. Based on these observations, we suggest
using phenol red-free cell culture medium for experiments involving LysoTracker probes. In case
the use of phenol red-containing medium cannot be avoided recommended alternative is to use
LysoTracker Green instead.

Figure 1. Comparison of the normalized raw integrated density of three different LysoTracker probes (red, blue and green) obtained
by cell imaging in cRPMI with phenol red (white bars) and without phenol red (gray bars). Error bars are standard deviation, n=10 cells
per treatment. * $p<0.05$ by a one-way ANOVA.

137 **2.3.   Stability of LysoTracker Red probe under continuous and staggered**
138 **imaging set-up**

The ability to trace the behaviour of lysosomes over longer periods opens a possibility to study
cellular processes upon prolonged and repeated exposure to NP. The suitability of LysoTracker
probes for prolonged imaging was evaluated by monitoring the LysoTracker Red fluorescent signal
over a period of 24 h in a continuous experimental set-up **(Figure 2B)**. J774A.1 cells were first
stained with LysoTracker Red for 1 hour, the medium was replaced with phenol red-free medium,
and cells were imaged live for 24 hours at 20 minutes intervals. Randomized datasets[17] were
recorded.

In a continuous imaging set-up, the fluorescent signal of LysoTracker Red did not significantly
alter over the first 8 h **(Figure 2C).** Longer periods were accompanied by a decrease in the raw
integrated intensity of the LysoTracker Red and increased cell death over time **(Video SI1)**. This
observation is supported by Chen et al., who demonstrated that prolonged intracellular
accumulation of LysoTracker probes causes an increase in intracellular pH and enhances
quenching of the fluorescent dye.[52] However, as assessed by lactate dehydrogenase (LDH) assay

(Figure 2A) we did not observe any significant changes in LDH release in the medium from
LysoTracker treated cells compared to the control cells. Hence, adding LysoTracker does not
cause cytotoxicity within the time range investigated. Therefore, we conclude that the cytotoxicity
is rather influenced by phototoxicity, an effect caused by extensive imaging.

To test whether there is a difference in LysoTracker Red signal intensity depending on the imaging
conditions an additional staggered experimental set-up was conducted as shown in **(Figure 2B)**.
We observed that the raw integrated density does not differ significantly among the staggered
imaging set-up of 1-, 4- and 24-hours after LysoTracker incubation **(Figure 2C)**. However, a
significant decrease in raw integrated density of LysoTracker Red signal was found in continuous
live-cell imaging set-up **(Figure 2C)** and the effect accumulated with increasing imaging time. This
suggests that the fluorescent intensity of the LysoTracker Red drops mostly because of
photobleaching effect due to continuous imaging for a prolonged time.[53] The effect might be cell
specific and is affected by the imaging settings.

Figure 2. Stability of LysoTracker Red probe under different imaging conditions. **a.** Cell viability after incubation with LysoTracker Red probe for 1 h, 4 h and 24 h assessed via membrane rupture assay (lactate dehydrogenase; LDH) and presented as fold change over untreated cells (control). Data shows the mean of three replicates. Cells exposed to 0.2% Triton X-100 (v/v) served as positive control for LDH assay. **b.** Schematic representation of staggered experimental set-up (cells were stained for 1 h with LysoTracker Red probe. The supernatant containing LysoTracker Red was removed and cells were further incubated and imaged at 1 h, 4 h and 24 h as hyperstacks) and continuous set-up (cells were stained for 1 h with LysoTracker Red probe and imaged continuously for 24 h as hyperstacks). **c.** Comparison of raw integrated densities of LysoTracker Red probe versus time for continuous and staggered imaging. Each time-point (hour) shows the raw integrated density measured for each cell individually. The whiskers are the standard deviations. **** ($p < 0.05$) is the significant difference analyzed by One-way ANOVA.

Once the LysoTracker probe stability and imaging set-up is optimized, the selection of the optimal NP to assess the specific research question is important. For our experiments, we selected synthetic amorphous silica NP of different sizes and labelled with different fluorophores.

2.4. Physicochemical characteristics of silica nanoparticles

Table 2 represents the main physicochemical characteristics of different sizes of silica (SiO_2) NP (59- and 920-nm) used in this study; NP of 59 nm size range are internalized into cells mainly via pinocytosis while 920 nm SiO_2 particles are taken up by phagocytosis.[43] A concentration of 20 µg/mL of the NP suspension was chosen as this concentration does not impair cell viability [32] and all the particles used in our studies were negative for the endotoxin test (data not shown).

The core particle diameter was measured by transmission electron microscopy (**Figure SI3**). The hydrodynamic diameter was assessed both in Milli-Q water and complete medium via dynamic light scattering (DLS). All SiO_2 particles were colloidally stable in Milli-Q and in cRPMI over 24 h of incubation time.

Table 2. Representation of the main physicochemical characteristics of SiO_2 particles.

Particles	TEM[5] d_c [nm] [1]	DLS[6] Milli-Q d_h [nm] [2] t = 0 h	d_h [nm] [2] t = 24 h	PDI[3]	ζ [mV] [4]	cRPMI d_h [nm] [2] t = 0 h	d_h [nm] [2] t = 24 h
SiO_2-Bodipy	59 ± 6	77 ± 0	76 ± 1	0.2	- 52 ± 2	91 ± 2	87 ± 3
SiO_2-RhoB	59 ± 5	71 ± 1	72 ± 1	0.04	- 34 ± 1	82 ± 4	69 ± 3
SiO_2-Cy5	920 ± 88	820 ± 3	847 ± 6	0.03	- 44 ± 1	825 ± 11	941 ± 10

[1] core diameter; [2] hydrodynamic diameter; [3] polydispersity index; [4] zeta potential; [5] transmission electron microscopy; [6] dynamic light scattering. Particles: SiO_2-Bodipy, SiO_2-Rhodamine B, SiO_2-Cyanine-5

2.5. Confocal microscopy imaging and post-processing

Even optimally designed experiments – no cross talk, no bleed through – and sample preparation can be undone by errors during sample imaging.[54] Small variations from the optimal settings can significantly alter the correlation coefficients. Besides resolution (spatial and axial), suboptimal illumination and even the way of digital storage of the data can affect the outcome. To demonstrate these biases effectively, suboptimal imaging conditions were simulated starting from an optimally imaged dataset, considering the spatial resolution, slice thickness, pixel saturation, or bit depth **(Figure 3)**. By simulating different imaging conditions, we analyzed the changes in the colocalization parameters from a confocal data set. Briefly, J774A.1 macrophages were exposed to 59 nm SiO_2-Bodipy NP for 13 h. Cells were washed to remove the remaining NP, followed by resting phase fixation and immunostaining for LAMP-2.

Lowering the resolution by decreasing XY-pixel size (i.e., the distance between points on a specimen, also known as "downsampling") can cause a decrease in Pearson's correlation coefficient (PCC) and especially in overlap (Manders M1). Lower resolutions cause the integration of fluorescence signals thereby affecting the interactions between the two channels. This is most clear in the overlap (M1), shown in the grey boxplots of **Figure 3.** Therefore, the optimal resolution to image the objects must be taken into consideration when setting up the experiment.

Besides spatial resolution, slice thickness of the optical slices will also affect the colocalization parameters. Simulation of the opening of the pinhole causing lower axial resolution (Z-axis thickness) yielded an increase in colocalization measures as objects separated along the Z-axis are increasingly projected on top of each other and end up falsely overlapping **(Figure 3).**

A brighter illumination will improve the signal to noise ratio of the fluorescence signal, which may yield a lower variance of the correlation coefficients. However, a saturation of pixels will bias the result. Saturation occurs when the fluorescence intensity exceeds the acquisition or storage limit of the detector. Such clipping of data has especially a significant impact on correlation routines, such as PCC, which try to explain the intensity of the first channel with the information of the second. Our simulations show that even a fraction of saturated pixels as low as 0.3 % of the total pixel number will significantly alter the colocalization **(Figure 3).** The pitfall of saturation can be completely circumvented by using photon-counting detectors,[55] which do not convert the signal to binary output (see below) but store the actual number of photons that have arrived at the detector.

CLSM detectors convert the analogue fluorescent intensities to a digital signal and store it on a computer. Usually, you will get the choice to save the data with an information depth ranging from 8 bit (2^8, i.e., 256 different intensity values) up to 16 bit (2^{16}, i.e., 65536 different intensity values). Recording data in 16 bit but saving the data at 8 bit, i.e., compressing the information, may cause significant shifts in PCC **(Figure 3).**

Figure 3. Effect of spatial resolution, optical slice thickness, pixel saturation and bit depth on the Pearson correlation coefficients (PCC, white boxes) and Manders' M1 coefficient (gray boxes). The data correspond to the experimental set-up where cells were exposed to 59 nm SiO$_2$-Bodipy NP for 13 h, fixed and stained with Alexa Fluor 647-conjugated LAMP-2 antibody. The error bars represent the standard deviations. All the images were analysed using ImageJ with the JACoP plugin comparing the correlation coefficients after each specific simulation.

Before proceeding with further experiments, all microscopy settings, such as resolution, optical slice thickness, saturation and bit depth should be taken into consideration for the reproducibility of the data.

2.6. Analysis of nanoparticles within the lysosomes

In this section, we describe four different experiments illustrating the colocalization of silica NP with lysosomes and the corresponding interpretation. The experimental design is matched to the specific aim of each study.

Experiment I: The effect of continuous live cell imaging on colocalization

Understanding NP localization in lysosomes after their cellular uptake is crucial for their use in biomedical applications.[56] The advantage of live cell imaging is to get a better insight into intracellular NP trafficking [20], and accumulation. The variability in NP uptake dynamics depends on the cell type[57] and cell division.[20,30,58,59] The following experimental approach enabled us to investigate whether NP colocalization with lysosomes varies over time under a continuous imaging set-up. J774A.1 cells were initially stained with LysoTracker Red for 1 h and then sequentially exposed to 59 nm SiO_2-Bodipy NP for a duration of 4-8 h **(Figure 4A)**. The colocalization was assessed using PCC as an indicator of functional relationship and Manders' coefficients (M1 and M2) as quantifiers of spatial overlap between fluorophores**.** PCC values indicate that approximately 50% (PCC: 0.5) of intracellular NP colocalized with lysosomes and there were no significant differences in PCC between 4- to 8-h **(Figure 4B)**. However, the PCC does not provide detailed information about the distribution of NP arriving into lysosomes and the fraction of remaining empty lysosomes **(Figure 4B)**. In contrast, M1 and M2 coefficients vary between the different time points **(Figure 4C)** and provide information on the fraction of the NP co-occurring with lysosomes within the resolution of a single voxel. We observed that less than 10% (<0.1 of fraction; M1) of NP reached the lysosomes within the first 4 h of exposure (**Figure 4C**). The fraction of NP that accumulated in the lysosomes increased gradually over time as confirmed by increasing M1 fraction up to 0.5 after 8 h of incubation. However, at 8 h post-incubation, less than 20% of the lysosomes colocalized with NP (M2), which is a significantly lower amount compared to the 4 h time-point. This suggests a possible formation of new lysosomes in macrophages over time[60] (**Figure 4C**).

Figure 4. The effect of continuous live cell imaging on NP colocalization with lysosomes. **a.** Experimental workflow: Cells were first stained with LysoTracker Red probe for 1 h, and then exposed to 20 µg/mL of 59 nm SiO$_2$-Bodipy NP in phenol free cRPMI. Cells were imaged continuously from 4 h to 8 h. All the images were analyzed using ImageJ with a script (Figure SI1). Each cell was analyzed individually. **b.** Pearson correlation coefficient (PCC) demonstrating correlation between NP and LysoTracker Red probe (lysosomes) at different time points. **c.** Manders' correlation coefficient M1 corresponds to the colocalization of 59 nm SiO$_2$-Bodipy NP and M2 corresponds to the colocalization of LysoTracker Red (lysosomes) filled with NP.

Based on different NP physicochemical properties such as core material, size, shape, surface charge, porosity, surface roughness, surface area, crystalline structure, NP uptake and their intracellular fate can vary over time. In order to investigate these phenomena, we monitored the changes in colocalization of NP of different sizes and incorporated fluorophores.

Experiment II: The effect of different NP sizes and on colocalization with lysosomes

A combination of the PCC and Manders' colocalization coefficients can provide different messages of NP accumulation in lysosomes.[40] Since particles of different sizes are widely used in the research, we chose two different-sized silica NPs (59 nm SiO$_2$-Bodipy NP and 920 nm SiO$_2$-Cyanine-5 particles) and applied them to J774A.1 macrophages. After exposure to a single NP type for 4 h or 24 h, cells were washed and stained with LysoTracker Red for 1 h followed by live cell imaging **(Figure 5A).**

The accumulation of 59 nm SiO$_2$-Bodipy NP within lysosomes was observed 4 h after exposure, as demonstrated by the PCC of 0.4, which was significantly different from 24h. The PCC increased to 0.7 over the next 20 hours further confirming the NP uptake and intracellular sorting to the

lysosomes. Such information cannot be extracted by the naked eye from the CLSM images but
can be visualized by the cytofluorograms **(Figure 5 B):** the point cloud expanded towards the top
right corner, and the increase in PCC reflected this process. Four hours post-exposure, Manders'
coefficient showed that 70% of the pixels associated with the 59 nm SiO_2-Bodipy NP were co-
occurring with pixels associated with LysoTracker Red, i.e. a lysosome (M1). However, less than
10% of the lysosomes were filled with NPs (M2). After 20 h, almost all internalized NP were
associated with the lysosomes (M1 > 80%) and most lysosomes were filled with NPs - we
observed about 20 % of lysosomes without NPs (M2, **Figure 5C**).

A different outcome was observed with the 920 nm SiO_2-Cyanine5 particles **(Figure 5 D, E)**. Four
hours post-exposure, these particles arrived at the lysosomes (PCC: 0.3, significantly different
from 0), but no significant increase in accumulation was observed over the next 20 h. Manders'
coefficients show that only 40% of particles reached lysosomes within 24 h (M1) and the majority
(about 80%) of lysosomes were still without particles (M2, **Figure 5E**). Thus, micron-sized particles
were less likely to enter the lysosomes within 24 h and possibly sedimented in the cells'
surroundings or accumulated in other intracellular compartments.[61–63] A possible explanation for
lower colocalization of bigger particles with lysosomes could be as well the differences in particle
number added to the cells ($7x10^{10}$ of SiO_2-Bodipy NP vs $2x10^7$ of SiO_2-Cy5 particles) or,
alternatively, different mobility of NP within the cells.[64]

Figure 5. Colocalization of different-sized SiO$_2$ particles with lysosomes in J774A.1 macrophages. **a.** Experimental workflow: Cells were initially exposed to NP for 4 h or 24 h followed by LysoTracker Red staining for 1 h and imaged live. **b.** Representative images of single cells with the corresponding cytofluorograms show colocalization of 59 nm SiO$_2$-Bodipy NP (green) and **d.** 920 nm SiO$_2$-Cyanine-5 particles (green) with lysosomes (red) 4 hour and 24 h post-exposure. Co-localized pixels appear in yellow. **c.** Quantification of colocalization of 59 nm SiO$_2$-Bodipy NP and **e.** 920 nm SiO$_2$-Cyanine 5 particles with LysoTracker Red probe was performed using the ImageJ software with JACoP plugin (n= 10 cells). Data is presented as Pearson correlation coefficient (PCC) and Manders' overlap coefficients, where M1 represents the fraction of NPs-associated pixels overlapping with lysosomes and M2 the fraction of lysosome-

 associated pixels overlapping with NPs. Statisitcal analysis was performed using unpaired t-test in GraphPad Prism: **** p < 0.0001,
312   ns – not significant.

Experiment III: Colocalization between NP with different fluorophores and lysosomes

A potential colocalization pitfall is the dependence of fluorescent dye encapsulation on the physicochemical properties of NP. The dye encapsulation might change the physicochemical properties of NP and the fluorescence properties of a dye, for example, by inducing fluorescence quenching or enhancing emission.[26,65,66] An experimental design was implemented to compare the colocalization of NP with similar sizes but different fluorophores. J774.A1 cells were initially exposed to 59 nm SiO_2-Bodipy NP and 59 nm SiO_2-Rhodamine B NP for a period of 13 h. The medium containing NP was subsequently removed and cells were washed. Next, a fresh medium without NP was added and a resting phase of 6 hours was performed to investigate whether all the NP that have just reached the cell membrane continued being internalized and accumulate in the lysosomes. The resting time was chosen based on known literature, as it has been shown that different NP or macromolecules reach the lysosomes in a period from 3 h to 6 h.[20,67–71] Finally, the cells were stained with LysoTracker probes, which were selected due to their compatibility with NP-conjugated fluorophores **(Figure 6 A)**. 59 nm SiO_2-Bodipy NP were combined with LysoTracker Red probe and 59 nm SiO_2-RhodamineB NP with LysoTracker Green probe **(Figure 6 B and D)**.

Based on PCC values, the results showed no significant differences in the colocalization of lysosomes and NP with different fluorophores (PCC: 0.9). Nevertheless, Manders' correlation coefficients (M1, fraction of NP inside the lysosomes) confirmed that 90% of SiO_2-Rhodamine B NP are localized in the lysosomes **(Figure 6 C)** compared to 70% of SiO_2-Bodipy NP **(Figure 6 E)**. This suggests that most likely the dye molecules incorporated in the silica NP change the surface properties of NP, thus affecting their uptake and their intracellular fate.[18,72,73] On the other hand, similar fraction of lysosomes filled with NP was observed when comparing NP with different fluorophores (M2: 0.7).

Figure 6. Colocalization between NP with different fluorophores and lysosomes. **a.** Experimental workflow. Representative images of single cells exposed to two different NP (green), **b.** SiO_2-Rhodamine-NP and **d.** SiO_2-Bodipy NP. Lysosomes were stained with LysoTracker Green and LysoTracker Red probes, respectively (shown in red). The plots represent main parameters (PCC, M1 and M2) to estimate colocalization between different NP and lysosomes. **c.** The colocalization between the 59 nm SiO_2-Rhodamine NP stained with LysoTracker Green probe and **e.** the colocalization of the 59 nm SiO_2-Bodipy NP stained with LysoTracker Red probe. PCC: Pearson correlation coefficient, Manders' correlation coefficients M1 and M2. The whiskers represent the standard deviations. Analysis was performed in individual cells for each experiment (n=10 to 13 cells). All the images were analysed using ImageJ, with the JACoP plugin comparing the correlation coefficients.

Experiment IV: Comparison of NP colocalization with lysosomes between fixed-and live-cell imaging

While performing lysosomal staining, both LysoTracker probes and LAMP antibodies are available. However, there are differences both in their specificity as well as in staining protocols, which can significantly impact the experimental outcome.[28] With this experimental approach, the changes in colocalization of NP with lysosomes between live- and fixed-cell imaging were evaluated. The same exposure time and resting phase was kept as described in section III **(Figure 7A).** In addition to the LysoTracker staining, cells were immunostained with LAMP-2 antibody **(Figure 7A, B and D)** and the results from fixed-cells were compared to the data collected in section III**.**

No significant changes in PCC values were observed among live and fixed cells for both NP types. Nevertheless, we observe differences for Manders' correlation coefficients (M1 and M2) **(Figure 7C and E)**. In live cells, exposed to 59 nm SiO_2-Bodipy NP, there were roughly 70 % of NP colocalized with the lysosomes (M1), but in fixed-cell the colocalization decreased to 40 % **(Figure 7C).** This difference is evident in the cells exposed to 59 nm SiO_2-Rhodamine B where 86 % of the NP are colocalized with the lysosomes of living cells stained with LysoTracker in contrast to only 24 % of colocalized NP inside the lysosomes of fixed-cells **(Figure 7E)**. Likewise, a decrease in the fraction of lysosomes filled with NP (M2) between live and fixed cells was noted. These differences are attributed to image processing, and selectivity of LysoTracker probe (stains lysosomes and late endosomes) vs LAMP-2 (specific to LAMP-2 protein present on lysosomal membrane).[74]

Correlative studies between live and fixed cells are challenging because the shape of the fixed cells varies from its natural state.[75] Moreover, a quantitative analysis that compares and correlates the intensities of fluorophores such as the LysoTracker probes after cell fixation is expected to be less precise due to the decrease of LysoTracker intensity and further changes in cellular shape **(Figure SI4**). We have already described that each experimental factor that alters the intensity of the fluorophores will have repercussions, especially on the Manders' coefficients, as shown in **Figure SI5** and **Figure SI6**. In conclusion, many parameters influence the NP-lysosome colocalization outcome such as the fluorophore dye to label the NP, the staining procedure for lysosomes in cells, the lysosome marker specificity, and the imaging set-up. Therefore, it is recommended to take these parameters into consideration when planning and conducting the experiments as well as to adjust each experiment individually based on the research question to minimize the variability.

Figure 7. NP colocalization with lysosomes between fixed-and live-cell imaging. **a.** Experimental workflow. Representative images of a single cell exposed to two different NP types, **b.** SiO$_2$-Rhodamine NP (green) and **d.** SiO$_2$-Bodipy NP (green). Lysosomes were stained with Lamp-2 antibody (red). The graphs depict the main parameters (Pearson correlation coefficient - PCC, Manders' correlation coefficients - M1 and M2) to estimate the colocalization between **c.** SiO$_2$-Rhodamine NP and lysosomes (LysoTracker Green) and **e.** SiO$_2$-Bodipy NP and lysosomes (LysoTracker Red) in live cells (white box) and fixed cells stained with and LAMP-2 (grey box). The whiskers are the standard deviations. The analysis was done in individual cells for each experiment (n=8-13 cells). All the images were analysed using ImageJ with the JACoP plugin comparing the correlation coefficients.

3. *Conclusions*

We visualized the lysosomes by labeling cells with LysoTracker probes and calculated the colocalization between fluorescently labelled NP and lysosomes using different correlation

coefficients. Herein, we described detailed steps and recommendations from the experimental
design, staining, sample preparation and imaging on the colocalization parameters.

Key considerations for conducting NP-lysosome imaging and colocalization analysis are:

i) Fluorescent lysosomal probe properties (see Table 1), such as concentration to be
used, incubation time and fluorophore, [18]
ii) Physicochemical characteristics of the NP (see Table 2) such as exposure time,
concentration, and administration method,[76]
iii) CLSM imaging setup including resolution, illumination, and storage considerations,[77,78]
iv) Colocalization analysis of NP in the lysosomes.[36]

We strongly advise to fully report the parameters of the image acquisition to allow correct
comparisons and reproducibility between experiments reported from different studies. The human
eye is not capable of extracting the correct colocalization interpretation from CLSM images alone:
a cytofluorogram must be added. Similarly, a single PCC often does not provide information about
the fractional distribution of the fluorophores co-occurring within the resolution of a single voxel,
for this an additional measure such as Manders' co-occurrence coefficients should be taken into
analysis. Additional recommendations are summarized in the **Table 3**.

Table 3. List of recommendations based on the different parameters.

Parameter	Recommendation
LysoTracker probe	- Check for intensity overlap of LysoTracker probe with the NP fluorescence properties - Do not exceed maximal incubation time - Perform imaging of LysoTracker probes in cell culture medium without phenol red - Perform a pilot study and cell viability test to acquire the time range wherein your cell's viability remains acceptable. - Do not exceed this time scale for continuous live-cell imaging (e.g., for J774.A1: 8 hours)

	• For experiments exceeding this time range, conduct a staggered set-up and stain with LysoTrackers before imaging
CLSM Imaging	• Apply optimal imaging parameters considering optimal resolution of sample to proceed with the colocalization analysis, as bias will increase with decreasing resolution • Optimize the pinhole setting • Never use (maximum) intensity projections • Avoid noise and saturation events • If available, use photon-counting detectors
Nanoparticle-lysosome colocalization analysis	• Describe data with adequate colocalization coefficients to provide the precise information about NP intracellular fate • Include the cytofluorograms • Analyse enough cells e.g. eight to ten single cells in the entire z-stack for each experimental condition, in case of high variability the number has to be increased • Perform analysis on single cells selected from region of interest (ROI) to avoid noise contribution • For colocalization analysis 3D data is recommended to be used. Keep in mind the different NP uptake kinetics and variability among cell lines. • Adjust experimental set-up specifically to the research question
Data analysis	• PCC does not provide detailed information about the distribution of NP arriving into lysosomes and the fraction of remaining or newly formed empty lysosomes • Manders' correlation coefficients M1 and M2 provide information on the fraction of the NP co-occurring with lysosomes within the resolution of a single voxel.

412   *4. Methods*
413

4.1. Synthesis of silica nanoparticles

The synthesis of fluorescently labelled silica (SiO_2) NP was performed following a Stöber method.[79] For a detailed description of the 59 nm SiO_2-Bodipy and 920 nm SiO_2-Cyanine-5 NP synthesis and characterization, refer to the publication from Susnik et al.[32] Similar synthesis protocol was adapted for 59 nm SiO_2-Rhodamine B NP using a Rhodamine B isothiocyanate (RBITC) dye (CAS 36877-69-7, Sigma- Aldrich, USA). **Table 2** summarizes the physicochemical characteristics of three different SiO_2 particles, such as core diameter measured by transmission electron microscopy (TEM; FEI Technai G2Spirit, Thermo Fisher Scientific, Waltham, MA, USA) and hydrodynamic diameter in water and complete RPMI by dynamic light scattering (DLS; LS instruments AG, Fribourg, Switzerland).

4.2. Cell culture and seeding

Mouse monocyte/macrophages J774A.1, obtained from the American Type Culture Collection (ATCC, Rockville, MD, USA) were subcultured Roswell Park Memorial Institute 1640 medium (RPMI; Cat.No. 11835030, Gibco, Life Technologies, Zug, Switzerland) supplemented with 10 % v/v FBS, 1 % v/v penicillin/streptomycin and 1 % v/v L-glutamine. Cells were maintained in T75 polystyrene cell culture flasks (TRP, Trasadingen, Switzerland) in cRPMI until reaching 80 % confluence. Cell seeding procedure has been performed as described by Susnik et al.[32]

4.3. The interference of phenol red with the LysoTracker probes

LysoTracker™ Red DND-99 (Cat. No. L7528), Green DND-26 (Cat. No. L7526) and Blue DND-22 (Cat. No. L7525) (Thermo Fisher Scientific, MA, USA) probes at final concentration 75 nM were prepared in cRPMI medium containing phenol red or in phenol red-free medium. J774A.1 macrophages were incubated with LysoTracker probes for 1 h at 37 °C, 5 % pCO_2 and 95 % relative humidity, following providers' protocol (Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA). After incubation, the medium containing LysoTracker probes was removed and replaced with fresh phenol red-containing or phenol red-free medium. Live cell imaging was done in cRPMI containing phenol red or in phenol free medium (refer to chapter Image acquisition). The images were analyzed using ImageJ software with a JACoP plugin and a custom-written script **(see Figure SI1)**. Briefly, each cell contour was manually defined based on the LysoTracker channel to define the region of interest (ROI) for the analysis. The detected signals from the LysoTracker

channel were masked, followed by an automatically adjusted threshold. An inverted mask was
used to remove the background noise and to measure the raw integrated density by excluding
background signals. The raw integrated densities for ten individual cells were analyzed for each
LysoTracker probe. The following parameters have been used: 16-bit image (1024x1024 pixels),
pixel dwell of 25.5,1 Airy unit and the gain for each individual fluorophore as it follows: LT blue
850, pinhole 43, channel range 410-431 nm; LT green 583 pinhole 52, channel range 493-606
nm; LT Red 446, pinhole 60, channel range 566-690 nm.

453 4.4. Stability of LysoTracker Red probe
454

For continuous imaging set-up, LysoTracker™ Red DND-99 (Cat. No. L7528), (Thermo Fisher
Scientific, MA, USA) probe at final concentration 75 nM was prepared in cRPMI medium
containing phenol red or in phenol red-free medium. J774A.1 macrophages were incubated with
LysoTracker probes for 1 h at 37 °C, 5 % pCO_2 and 95 % relative humidity, following providers'
protocol (Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA). After incubation, the medium
containing LysoTracker probes was replaced with fresh phenol red-free medium without
LysoTracker probe. Cells were immediately imaged live (refer to chapter Image acquisition) in a
continuous (24h) or staggered set-up (1h, 4h, 24h). The images were analyzed using ImageJ
software with a JACoP plugin and a custom-written script as in described previously **(see Figure
SI1)**. The raw integrated densities for individual cells were analyzed. The following parameters
have been used: 16-bit image (1024x1024 pixels) and pixel dwell of 25.5,1 Airy unit.

466 4.5. Cell viability
467

Cells were incubated with LysoTracker Red probe for 1 h, 4 h and 24 h under standard cell culture
conditions. Supernatants were collected and cell viability was assessed based on the lactate
dehydrogenase (LDH) enzyme released from the damaged cells. Data was analyzed in triplicate
using LDH cytotoxicity detection kit (Roche Applied Science, Mannheim, Germany) according to
the manufacturer's protocol. The absorbance of colorimetric product formazan was recorded using
a microplate reader (Bio-Rad, Switzerland) at 490 nm with a reference wavelength of 630 nm. The
values were expressed as fold increase of the treated samples relative to the untreated (control)
samples. Triton X-100 (0.2% v/v; in phosphate buffered saline (PBS)) (Merck, Switzerland) was
added to the positive control 15 min prior to collecting the supernatant of each well.

477 4.6. Cell exposure to silica NP and LysoTracker probes
478

A suspension of 1 mg/mL silica NP was first prepared in Milli-Q water. NP were dispersed in
cRPMI to the final concentration of 20 μg/mL and administered to the cells via a pre-mixed method
i.e., NP were added to cRPMI immediately prior to the cell exposure. Three experimental setups
were designed based on the addition of the LysoTracker probes:

i. Addition of LysoTracker probe before NP exposure for live cell imaging: Cells were
seeded in uncoated glass-bottom MatTek culture dishes (35 mm, Life Sciences,
Massachusetts, USA) for 24 h and incubated with fresh cRPMI supplemented with 75
nM LysoTracker Red probe for 1 h at 37 °C. Afterwards, the supernatant containing
LysoTracker Red probe was removed, and cells were exposed to 59 nm SiO_2-Bodipy NP
in cRPMI for a period of 8 h. The image acquisition was performed from 4 h to 8 h in a
chamber at 37 °C and 5 % pCO_2, adapted for the Zeiss LSM710 confocal microscope.
The cells were imaged as hyperstacks over time. Time lapse and different z-stacks were
set for each sample. The time point was defined upon NP exposure and the analysis was
done from 4 - 8 h. For each time point, 13 individual cells were analyzed to evaluate the
raw integrated intensity of LysoTracker probe over time. The raw integrated density was
processed following the same procedure as described in section 4.3. The correlation
coefficients were calculated using ImageJ software with JACoP plugin and a custom-
written script (see **Script SI2**).

ii. Addition of the LysoTracker probe after the NP exposure: Cells were initially
exposed to a single particle type for different time-points (4 h and 24 h). The cells were
washed 3-times with 1X PBS and incubated with fresh cRPMI supplemented with 75 nM
LysoTracker probe for 1 h (according to the manufacturers' recommendations).
Afterwards, the supernatant containing LysoTracker probe was removed, and fresh
cRPMI was added to the cells for immediate live-cell imaging.

**iii. Addition of the LysoTracker probe after NP incubation and followed by resting
period:** Cells were exposed to 59 nm SiO_2-Rhodamine B NP or 59 nm SiO_2- Bodipy NP
for 13 hours, washed 3-times with 1X PBS and further incubated with fresh cRPMI. Cells
were incubated for an additional 6 h to ensure that most of the NP reached the
lysosomes. After, cells were stained with 75 nM LysoTracker Green or 75 nM
LysoTracker Red for 60 minutes. The LysoTracker probes were replaced with fresh
cRPMI and cells underwent immediate live-cell imaging.

4.7. Image acquisition
511

During live-cell imaging, cells were kept in phenol-red containing cRPMI or alternatively in cRPMI
without phenol red. Image acquisition was performed with Zeiss LSM 710 Meta confocal
microscope Axio Observer.Z1 (Zeiss GmbH, Jena, Germany), using a Plan-Apochromat objective
63×/NA 1.4-immersion oil objective lens fitted with diode-pumped solid-state laser (DPSS) 561-
10, argon 458, 488, 514 and HeNe 633 nm lasers. An immersion oil (Carl Zeiss Immersol™ 518F)
with a refractive index 1.518 was used. The excitation/emission spectra peaks for LysoTracker
probes: Blue (λ_{ex}: 373 nm; λ_{em}: 422 nm), Green (λ_{ex}: 504 nm; λ_{em}: 511 nm) and Red (λ_{ex}: 577 nm;
λ_{em}: 590 nm), using the respective lasers (405 nm, argon 458 nm and DPSS 561 nm) with adjusted
master gain and pinhole for each of the experiments. The excitation/emission spectra peaks for
NP are the following: SiO_2- Bodipy NP (laser λ_{ex}: 488 nm; λ_{em}: 520/30 nm), SiO_2-RhoB NP (λ_{ex}:
561 nm; λ_{em}: 620/30 nm), and SiO_2-Cyanine-5 NP (λ_{ex}: 633 nm; λ_{em}: 700/60 nm). After acquisition,
the images were analyzed using FIJI software (ImageJ, National Institutes of Health, Bethesda,
MD, USA).

4.8.   Colocalization analysis

To quantify the degree of colocalization between fluorescently labeled silica NP and LysoTracker
probes, the Pearson correlation coefficient (PCC) and the Manders' correlation coefficient (M1
and M2) were determined from the corresponding confocal images using FIJI software (ImageJ,
National Institutes of Health, Bethesda, MD, USA) with JACoP plugin and a custom-written script
(Script SI2). We used an approach without Costes randomizations, taking the automatic adjusted
threshold for the two different channels. The cell contours were manually defined based on the
LysoTracker channel to select the region of interest (ROI) for the analysis and avoid background
signals. In the script run for each channel the automatic threshold was adjusted, where channel 1
represented nanoparticles and channel 2 corresponded to a LysoTracker probe. All the images
were analyzed with the same script and the threshold of each individual channel was adjusted
each time. From the JaCoP analysis we extracted Pearson and Manders' (M1 and M2) correlation
coefficients for data representation.

4.9.   Data post-processing and simulation

The two channels of the dataset were treated to mimic different imaging setups. The simulations
were conducted in Fiji software and colocalization was performed in JaCoP plugin. **i)** To mimic
loss of spatial resolution, the dataset was binned in XY using the build-in scaling function of Fiji.
ii) Loss of axial resolution (increase in optical slice thickness) was created by the sum of slices of

the grouped Z-project routine. **iii)** Pixel saturation was simulated by iteratively multiplying the
intensity of each pixel by a factor 1.1 of an oversaturation-free stack and calculating the
colocalization measures after each step. **iv)** The loss of bit depth was made by converting the
original 16-bit stack to 8-bit and repeating the colocalization analysis. This part is divided in two
segments: **i)** Immunostaining: Upon completion of the NP exposures, cells were washed with PBS
and fixed with 4 % paraformaldehyde (in PBS, v/v) for 15 minutes. After washing, cells were
blocked with 4 % bovine serum albumin (BSA) (in PBS, v/v) for 1 hour, washed with PBS and
stained overnight at 4 °C with 20 µg/mL of LAMP-2 primary antibody (Thermo Fisher Scientific,
MA, USA) in 1 % BSA (in PBS, v/v). After washing, Alexa 647-conjugated secondary antibody was
applied (1:500 dilution) for 1 hour. Cells were washed and mounted (Fluoromount™ Aqueous
Mounting Medium (F4680-25mL, Sigma-Aldrich) before proceeding with imaging with Zeiss LSM
710 confocal microscope. LAMP-2 corresponding excitations were for Alexa 647 varying the
master gain and pinhole for each of the experiments. This data was used for the analysis of the
colocalization coefficients for fixed-cells described in section VI-experiment III. **ii)** The simulations
using scripts and macros (see SI): cLSM images were manipulated to evaluate the effect of spatial
resolution, optical slice thickness, pixel saturation and bit depth on the colocalization parameters.
The colocalization was analyzed in ImageJ software using the custom-written script and JACoP
plugin. Representative data was selected to show the effect of different imaging conditions on the
correlation coefficients.

4.10 Statistical Analysis
566

The statistical analyses were performed in GraphPad Prism 9 software, Origin 2016, 64 bit
adjusting each of the treatments to a corresponding statistical analysis. One-way ANOVA, t-test
or Dunnett's test analysis were performed. The results are reported as the mean with standard
deviation of the mean (SD).

Abbreviations

BSA: Bovine serum albumin

CLSM: Confocal laser scanning microscopy

cRPMI: Complete Roswell Park Memorial Institute medium, supplemented with 10% fetal bovine serum, 1
% (v/v) L-glutamine and 1 % (v/v) penicillin/streptomycin

DLS: Dynamic light scattering

FBS: Fetal bovine serum

FIB-SEM: Focused ion beam-scanning electron microscopy

LAMP: Lysosomal-Associated Membrane Protein

LDH: Lactate dehydrogenase

M1 and M2: Manders' coefficients

NP: Nanoparticles

PBS: Phosphate buffered saline

PCC: Pearson's correlation coefficient

RBITC: Rhodamine B isothiocyanate

ROI: Region of interest

SiO$_2$: Silica

Supplementary information

Figure SI1. Comparison of the absorbance and emission spectra of the different LysoTracker probes to
compare the interference of the phenol red. **Figure SI2.** Confocal microscopy images of different
LysoTracker probes. **Figure SI3.** Transmission electron micrographs (TEM) of the different silica NP. **Figure**
SI4. Representative images of LysoTracker and Lamp-2 stainings in a single cell. **Figure SI5.** Comparison
of Pearson's and Manders' coefficients between live and fixed cells. **Figure SI6.** Comparison of raw
integrated intensity between live and fixed cells. **Video SI1.** Continuous live cell imaging of J774A.1 cells.
Script SI1. Raw integrated densities. **Script SI2.** Colocalization analysis.

Acknowledgments

The authors would like to thank Liliane Ackermann Hirschi for supporting the synthesis of the nanoparticles.

Author contributions

The influence of phenol red in cell culture medium on fluorescence intensity of LysoTracker probes and the
stability of LysoTracker Red probe were tested by AMME. The analysis of nanoparticles within the
lysosomes for the experiment I, III, IV was done by AMME and experiment II was done by ES. the
simulations and post-processing imaging and the design of figure panels were done by DV. The support for
the synthesis of the nanoparticles was done by PTB. The dynamic light scattering analysis was supported
by SB. AMME and ES conducted the literature search, designed, and prepared a manuscript. AMME, ES,

DV, BR, AK, DV participated in the writing, discussing and preparing the final manuscript. The supervision
by BR and AK. All authors have read and agreed to the published version of the manuscript.

Founding

610 The authors acknowledge funding from the Swiss National Science Foundation (SNSF) (project
611 310030_159847/1) and the Adolphe Merkle Foundation. Additionally, this work benefitted from support
612 from the Swiss National Science Foundation through the National Center of Competence in Research Bio-
613 Inspired Materials.

Declarations

Consent for publication

616 All authors have approved the submitted manuscript.

Competing interests

618 The authors declare no competing interests.

Author details

620 [1] Adolphe Merkle Institute, University of Fribourg, Switzerland, Chemin des Verdiers 4, 1700 Fribourg,
621 Switzerland. [2] Department of Chemistry, University of Fribourg, Chemin du Musée 9, 1700 Fribourg,
622 Switzerland. [3] International Iberian Nanotechnology Laboratory, Avenida Mestre José Veiga, s/n 4715-330
623 Braga, Portugal.

References

1. Zhao J, Stenzel MH. Entry of nanoparticles into cells: The importance of nanoparticle
properties. *Polym Chem*. 2018;9(3):259-272. doi:10.1039/c7py01603d

2. Park JH, Oh N. Endocytosis and exocytosis of nanoparticles in mammalian cells. *Int J
Nanomedicine*. Published online 2014:51. doi:10.2147/ijn.s26592

3. Iversen TG, Skotland T, Sandvig K. Endocytosis and intracellular transport of
nanoparticles: Present knowledge and need for future studies. *Nano Today*.
2011;6(2):176-185. doi:10.1016/j.nantod.2011.02.003

4. Almeida MS De, Susnik E, Drasler B, Rothen-rutishauser B. Understanding nanoparticle
endocytosis to improve targeting strategies in nanomedicine. Published online 2021.
doi:10.1039/d0cs01127d

5. Rennick JJ, Johnston APR, Parton RG. Key principles and methods for studying the
endocytosis of biological and nanoparticle therapeutics. *Nat Nanotechnol*.
2021;16(3):266-276. doi:10.1038/s41565-021-00858-8

6. Brandenberger C, Mühlfeld C, Ali Z, et al. Quantitative Evaluation of Cellular Uptake and
Trafficking of Plain and Polyethylene Glycol-Coated Gold Nanoparticles. *Small*.
2010;6(15):1669-1678. doi:10.1002/smll.201000528

7.   Treuel L, Jiang X, Nienhaus GU. New views on cellular uptake and trafficking of
manufactured nanoparticles. *J R Soc Interface*. 2013;10(82):20120939.
doi:10.1098/rsif.2012.0939

8.   Cupic KI, Rennick JJ, Johnston AP, Such GK. Controlling endosomal escape using
nanoparticle composition: current progress and future perspectives. *Nanomedicine*.
2019;14(2):215-223. doi:10.2217/nnm-2018-0326

9.   Saftig P, Klumperman J. Lysosome biogenesis and lysosomal membrane proteins:
trafficking meets function. *Nat Rev Mol Cell Biol*. 2009;10(9):623-635.
doi:10.1038/nrm2745

10. Ding L, Zhu X, Wang Y, et al. Intracellular Fate of Nanoparticles with Polydopamine
Surface Engineering and a Novel Strategy for Exocytosis-Inhibiting, Lysosome
Impairment-Based Cancer Therapy. *Nano Lett*. 2017;17(11):6790-6801.
doi:10.1021/acs.nanolett.7b03021

11. Wong C-O, Gregory S, Hu H, et al. Lysosomal Degradation Is Required for Sustained
Phagocytosis of Bacteria by Macrophages. *Cell Host Microbe*. 2017;21(6):719-730.e6.
doi:10.1016/j.chom.2017.05.002

12. Platt FM. Sphingolipid lysosomal storage disorders. *Nature*. 2014;510(7503):68-75.
doi:10.1038/nature13476

13. Kim SE, Overholtzer M. Autophagy proteins regulate cell engulfment mechanisms that
participate in cancer. *Semin Cancer Biol*. 2013;23(5):329-336.
doi:10.1016/j.semcancer.2013.05.004

14. Jiang P, Mizushima N. Autophagy and human diseases. *Cell Res*. 2014;24(1):69-79.
doi:10.1038/cr.2013.161

15. Appelqvist H, Wäster P, Kågedal K, Öllinger K. The lysosome: from waste bag to potential
therapeutic target. *J Mol Cell Biol*. 2013;5(4):214-226. doi:10.1093/jmcb/mjt022

16. Padh H, Niraj Sakhrani. Organelle targeting: third level of drug targeting. *Drug Des Devel
Ther*. Published online July 2013:585. doi:10.2147/DDDT.S45614

17. Drasler B, Vanhecke D, Rodriguez-Lorenzo L, Petri-Fink A, Rothen-Rutishauser B.
Quantifying nanoparticle cellular uptake: which method is best? *Nanomedicine*.
2017;12(10):1095-1099. doi:10.2217/nnm-2017-0071

18. Brandenberger C, Mühlfeld C, Ali Z, et al. Quantitative Evaluation of Cellular Uptake and
Trafficking of Plain and Polyethylene Glycol-Coated Gold Nanoparticles. *Small*.
2010;6(15):1669-1678. doi:10.1002/smll.201000528

19. Calero M, Chiappi M, Lazaro-Carrillo A, et al. Characterization of interaction of magnetic
nanoparticles with breast cancer cells. *J Nanobiotechnology*. 2015;13(1):16.
doi:10.1186/s12951-015-0073-9

20. Vtyurina N, Åberg C, Salvati A. Imaging of nanoparticle uptake and kinetics of intracellular
trafficking in individual cells. *Nanoscale*. 2021;13(23):10436-10446.
doi:10.1039/d1nr00901j

21. Qiu K, Du Y, Liu J, Guan J-L, Chao H, Diao J. Super-resolution observation of lysosomal
dynamics with fluorescent gold nanoparticles. *Theranostics*. 2020;10(13):6072-6081.

doi:10.7150/thno.42134

22.   Liu M, Li Q, Liang L, et al. Real-time visualization of clustering and intracellular transport
of gold nanoparticles by correlative imaging. *Nat Commun.* 2017;8(1):15646.
doi:10.1038/ncomms15646

23.   Guehrs E, Schneider M, Günther CM, et al. Quantification of silver nanoparticle uptake
and distribution within individual human macrophages by FIB/SEM slice and view. *J*
Nanobiotechnology. 2017;15(1):21. doi:10.1186/s12951-017-0255-8

24.   Tharkeshwar AK, Demedts D, Annaert W. Superparamagnetic Nanoparticles for
Lysosome Isolation to Identify Spatial Alterations in Lysosomal Protein and Lipid
Composition. *STAR Protoc.* 2020;1(3):100122. doi:10.1016/j.xpro.2020.100122

25.   Zhu M, Nie G, Meng H, Xia T, Nel A, Zhao Y. Physicochemical Properties Determine
Nanomaterial Cellular Uptake, Transport, and Fate. *Acc Chem Res.* 2013;46(3):622-631.
doi:10.1021/ar300031y

26.   Milosevic AM, Rodriguez-Lorenzo L, Balog S, Monnier CA, Petri-Fink A, Rothen-
Rutishauser B. Assessing the Stability of Fluorescently Encoded Nanoparticles in
Lysosomes by Using Complementary Methods. *Angew Chemie Int Ed.*
2017;56(43):13382-13386. doi:10.1002/anie.201705422

27.   Chen X, Bi Y, Wang T, et al. Lysosomal Targeting with Stable and Sensitive Fluorescent
Probes (Superior LysoProbes): Applications for Lysosome Labeling and Tracking during
Apoptosis. *Sci Rep.* 2015;5:1-10. doi:10.1038/srep09004

28.   Baba K, Kuwada S, Nakao A, et al. Different localization of lysosomal-associated
membrane protein 1 (LAMP1) in mammalian cultured cell lines. *Histochem Cell Biol.*
2020;153(4):199-213. doi:10.1007/s00418-019-01842-z

29.   Arruda LB, Sim D, Chikhlikar PR, et al. Dendritic Cell-Lysosomal-Associated Membrane
Protein (LAMP) and LAMP-1-HIV-1 Gag Chimeras Have Distinct Cellular Trafficking
Pathways and Prime T and B Cell Responses to a Diverse Repertoire of Epitopes. *J*
Immunol. 2006;177(4):2265-2275. doi:10.4049/jimmunol.177.4.2265

30.   Bourquin J, Septiadi D, Vanhecke D, et al. Reduction of Nanoparticle Load in Cells by
Mitosis but Not Exocytosis. *ACS Nano.* 2019;13(7):7759-7770.
doi:10.1021/acsnano.9b01604

31.   Schütz I, Lopez-Hernandez T, Gao Q, et al. Lysosomal dysfunction caused by cellular
accumulation of silica nanoparticles. *J Biol Chem.* 2016;291(27):14170-14184.
doi:10.1074/jbc.M115.710947

32.   Susnik E, Taladriz-Blanco P, Drasler B, Balog S, Petri-Fink A, Rothen-Rutishauser B.
Increased Uptake of Silica Nanoparticles in Inflamed Macrophages but Not upon Co-
Exposure to Micron-Sized Particles. *Cells.* 2020;9(9). doi:10.3390/cells9092099

33.   Sugimoto Y, Ninomiya H, Ohsaki Y, et al. Accumulation of cholera toxin and GM1
ganglioside in the early endosome of niemann-pick C1-deficient cells. *Proc Natl Acad Sci*
U S A. 2001;98(22):12391-12396. doi:10.1073/pnas.221181998

34.   Freundt EC, Czapiga M, Lenardo MJ. Photoconversion of Lysotracker Red to a green
fluorescent molecule. *Cell Res.* 2007;17(11):956-958. doi:10.1038/cr.2007.80

35. Kim D-H, Chang Y, Park S, et al. Blue-conversion of organic dyes produces artifacts in multicolor fluorescence imaging. *Chem Sci*. 2021;12(25):8660-8667. doi:10.1039/d1sc00612f

36. Aaron JS, Taylor AB, Chew T-L. Image co-localization – co-occurrence versus correlation. *J Cell Sci*. 2018;131(3):jcs211847. doi:10.1242/jcs.211847

37. Manders EMM, Verbeek FJ, Aten JA. Measurement of co-localization of objects in dual-colour confocal images. *J Microsc*. 1993;169(3):375-382. doi:10.1111/j.1365-2818.1993.tb03313.x

38. Pearson K. VII. Mathematical contributions to the theory of evolution.—III. Regression, heredity, and panmixia. *Philos Trans R Soc London Ser A, Contain Pap a Math or Phys Character*. 1896;187:253-318. doi:10.1098/rsta.1896.0007

39. Adler J, Pagakis SN, Parmryd I. Replicate-based noise corrected correlation for accurate measurements of colocalization. *J Microsc*. 2008;230(1):121-133. doi:10.1111/j.1365-2818.2008.01967.x

40. Costes S V., Daelemans D, Cho EH, Dobbin Z, Pavlakis G, Lockett S. Automatic and Quantitative Measurement of Protein-Protein Colocalization in Live Cells. *Biophys J*. 2004;86(6):3993-4003. doi:10.1529/biophysj.103.038422

41. Li Q. A Syntaxin 1, G o, and N-Type Calcium Channel Complex at a Presynaptic Nerve Terminal: Analysis by Quantitative Immunocolocalization. *J Neurosci*. 2004;24(16):4070-4081. doi:10.1523/JNEUROSCI.0346-04.2004

42. Woodcroft BJ, Hammond L, Stow JL, Hamilton NA. Automated organelle-based colocalization in whole-cell imaging. *Cytom Part A*. 2009;75A(11):941-950. doi:10.1002/cyto.a.20786

43. Kinnear C, Moore TL, Rodriguez-Lorenzo L, Rothen-Rutishauser B, Petri-Fink A. Form Follows Function: Nanoparticle Shape and Its Implications for Nanomedicine. *Chem Rev*. 2017;117(17):11476-11521. doi:10.1021/acs.chemrev.7b00194

44. Scientific TF. Fluorescence SpectraViewer. Published 2022. https://www.thermofisher.com/order/fluorescence-spectraviewer#!/

45. Ashino Y, Ying X, Dobbs LG, Bhattacharya J. [Ca 2+] i oscillations regulate type II cell exocytosis in the pulmonary alveolus. *Am J Physiol Cell Mol Physiol*. 2000;279(1):L5-L13. doi:10.1152/ajplung.2000.279.1.L5

46. Wemhöner A, Frick M, Dietl P, Jennings P, Haller T. A Fluorescent Microplate Assay for Exocytosis in Alveolar Type II Cells. *J Biomol Screen*. 2006;11(3):286-295. doi:10.1177/1087057105285284

47. Li Y, Wang S, Zhao Y, et al. A Model In Vitro Study Using Hypericin: Tumor-Versus Necrosis-Targeting Property and Possible Mechanisms. *Biology (Basel)*. 2020;9(1):13. doi:10.3390/biology9010013

48. Viswanathan G, Hsu Y-H, Voon SH, et al. A Comparative Study of Cellular Uptake and Subcellular Localization of Doxorubicin Loaded in Self-Assemblies of Amphiphilic Copolymers with Pendant Dendron by MDA-MB-231 Human Breast Cancer Cells. *Macromol Biosci*. 2016;16(6):882-895. doi:10.1002/mabi.201500435

49. Bussi C, Peralta Ramos JM, Arroyo DS, et al. Alpha-synuclein fibrils recruit TBK1 and
OPTN to lysosomal damage sites and induce autophagy in microglial cells. *J Cell Sci*.
Published online January 1, 2018. doi:10.1242/jcs.226241

50. Xu H, Lee S-J, Suzuki E, et al. A lysosomal tetraspanin associated with retinal
degeneration identified via a genome-wide screen. *EMBO J*. 2004;23(4):811-822.
doi:10.1038/sj.emboj.7600112

51. Pol A, Luetterforst R, Lindsay M, Heino S, Ikonen E, Parton RG. A Caveolin Dominant
Negative Mutant Associates with Lipid Bodies and Induces Intracellular Cholesterol
Imbalance. *J Cell Biol*. 2001;152(5):1057-1070. doi:10.1083/jcb.152.5.1057

52. Chen X, Bi Y, Wang T, et al. Lysosomal Targeting with Stable and Sensitive Fluorescent
Probes (Superior LysoProbes): Applications for Lysosome Labeling and Tracking during
Apoptosis. *Sci Rep*. 2015;5(1):9004. doi:10.1038/srep09004

53. Lou X, Zhang M, Zhao Z, et al. A photostable AIE fluorogen for lysosome-targetable
imaging of living cells. *J Mater Chem B*. 2016;4(32):5412-5417. doi:10.1039/c6tb01293k

54. Jonkman J, Brown CM, Wright GD, Anderson KI, North AJ. Tutorial: guidance for
quantitative confocal microscopy. *Nat Protoc*. 2020;15(5):1585-1611.
doi:10.1038/s41596-020-0313-9

55. Leng S, Bruesewitz M, Tao S, et al. Photon-counting Detector CT: System Design and
Clinical Applications of an Emerging Technology. *RadioGraphics*. 2019;39(3):729-743.
doi:10.1148/rg.2019180115

56. Chikte S, Panchal N, Warnes G. Use of LysoTracker dyes: A flow cytometric study of
autophagy. *Cytom Part A*. 2014;85(2):169-178. doi:10.1002/cyto.a.22312

57. Kuhn DA, Vanhecke D, Michen B, et al. Different endocytotic uptake mechanisms for
nanoparticles in epithelial cells and macrophages. *Beilstein J Nanotechnol*.
2014;5(1):1625-1636. doi:10.3762/bjnano.5.174

58. Kim JA, Aberg C, Salvati A, Dawson KA. Role of cell cycle on the cellular uptake and
dilution of nanoparticles in a cell population. *Nat Nanotechnol*. 2012;7(1):62-68.
doi:10.1038/nnano.2011.191

59. Åberg C. Kinetics of nanoparticle uptake into and distribution in human cells. *Nanoscale
Adv*. 2021;3(8):2196-2212. doi:10.1039/d0na00716a

60. Yin Q, Jian Y, Xu M, et al. CDK4/6 regulate lysosome biogenesis through TFEB/TFE3. *J
Cell Biol*. 2020;219(8). doi:10.1083/JCB.201911036

61. Cheng W, Kim S, Zivkovic S, Chung H, Ren Y, Guan J. Specific Labelling of Phagosome-
Derived Vesicles in Macrophages with a Membrane Dye Delivered with Microfabricated
Microparticles. *SSRN Electron J*. Published online 2021:1-30. doi:10.2139/ssrn.3937057

62. Li Y, Hu Q, Miao G, et al. Size-dependent mechanism of intracellular localization and
cytotoxicity of mono-disperse spherical mesoporous nano-and micron-bioactive glass
particles. *J Biomed Nanotechnol*. 2016;12(5):863-877. doi:10.1166/jbn.2016.2235

63. Seydoux E, Rothen-Ruthishauser B, Nita I, et al. Size-dependent accumulation of
particles in lysosomes modulates dendritic cell function through impaired antigen
degradation. *Int J Nanomedicine*. Published online August 2014:3885.

doi:10.2147/IJN.S64353

64.   Oh YK, Swanson JA. Different fates of phagocytosed particles after delivery into
macrophage lysosomes. *J Cell Biol*. 1996;132(4):585-593. doi:10.1083/jcb.132.4.585

65.   Zhukova V, Osipova N, Semyonkin A, et al. Fluorescently Labeled PLGA Nanoparticles
for Visualization In Vitro and In Vivo: The Importance of Dye Properties. *Pharmaceutics*.
2021;13(8):1145. doi:10.3390/pharmaceutics13081145

66.   Rodriguez-Lorenzo L, Fytianos K, Blank F, Von Garnier C, Rothen-Rutishauser B, Petri-
Fink A. Fluorescence-encoded gold nanoparticles: Library design and modulation of
cellular uptake into dendritic cells. *Small*. 2014;10(7):1341-1350.
doi:10.1002/smll.201302889

67.   Hirota Y, Masuyama N, Kuronita T, Fujita H, Himeno M, Tanaka Y. Analysis of post-
lysosomal compartments. *Biochem Biophys Res Commun*. 2004;314(2):306-312.
doi:10.1016/j.bbrc.2003.12.092

68.   Wang F, Salvati A, Boya P. Lysosome-dependent cell death and deregulated autophagy
induced by amine-modified polystyrene nanoparticles. *Open Biol*. 2018;4(8).
doi:10.1098/rsob.170271

69.   Solorio-Rodríguez A, Escamilla-Rivera V, Uribe-Ramírez M, et al. In vitro cytotoxicity
study of superparamagnetic iron oxide and silica nanoparticles on pneumocyte
organelles. *Toxicol Vitr*. 2021;72(November 2020). doi:10.1016/j.tiv.2020.105071

70.   van Meel E, Bos E, van der Lienden MJC, et al. Localization of active endogenous and
exogenous β-glucocerebrosidase by correlative light-electron microscopy in human
fibroblasts. *Traffic*. 2019;20(5):346-356. doi:10.1111/tra.12641

71.   Banquy X, Suarez F, Argaw A, et al. Effect of mechanical properties of hydrogel
nanoparticles on macrophage cell uptake. *Soft Matter*. 2009;5(20):3984-3991.
doi:10.1039/b821583a

72.   Yu T, Malugin A, Ghandehari H. Impact of silica nanoparticle design on cellular toxicity
and hemolytic activity. *ACS Nano*. 2011;5(7):5717-5728. doi:10.1021/nn2013904

73.   Abdelkhaliq A, Van Der Zande M, Punt A, et al. Impact of nanoparticle surface
functionalization on the protein corona and cellular adhesion, uptake and transport 03
Chemical Sciences 0306 Physical Chemistry (incl. Structural). *J Nanobiotechnology*.
2018;16(1):1-13. doi:10.1186/s12951-018-0394-6

74.   Ramsey N. Majzoub, Emily Wonder, Kai K. Ewert, Venkata Ramana Kotamraju, Tambet
Teesalu CRS. Rab11 and Lysotracker Markers Reveal Correlation between Endosomal
Pathways and Transfection Efficiency of Surface- Functionalized Cationic Liposome–DNA
Nanoparticles. *Physiol Behav*. 2016;120(26):6439–6453.
doi:10.1021/acs.jpcb.6b04441.Rab11

75.   De Boer P, Hoogenboom JP, Giepmans BNG. Correlated light and electron microscopy:
Ultrastructure lights up! *Nat Methods*. 2015;12(6):503-513. doi:10.1038/nmeth.3400

76.   Moore TL, Urban DA, Rodriguez-Lorenzo L, et al. Nanoparticle administration method in
cell culture alters particle-cell interaction. *Sci Rep*. 2019;9(1):1-9. doi:10.1038/s41598-
018-36954-4

77.   Bolte S, Cordelières FP. A guided tour into subcellular colocalization analysis in light
microscopy. *J Microsc*. 2006;224(3):213-232. doi:10.1111/j.1365-2818.2006.01706.x

78.   Ettinger A (2014). Fluorescence live cell imaging. *Methods Cell Biol*. 2014;123:77-94.
doi:10.1016/B978-0-12-420138-5.00005-7

79.   Stöber W, Fink A, Bohn E. Controlled growth of monodisperse silica spheres in the micron
size range. *J Colloid Interface Sci*. 1968;26(1):62-69. doi:10.1016/0021-9797(68)90272-5

876 **Figure SI1.** Comparison of the absorbance and emission spectra of the different LysoTracker probes to
877 compare the interference with the phenol red.

880 **Figure SI2.** Confocal microscopy images of different LysoTracker probes.

Figure SI3. Transmission electron micrographs (TEM) of the different silica NP. From left to right: 59 nm SiO$_2$-Bodipy NP, 59 nm SiO$_2$-Rhodamine NP and 920 nm SiO$_2$-Cyanine5 particles.

Figure SI4. Representative images of LysoTracker and Lamp-2 stainings in a single cell. Left: Lysosomes stained with LysoTracker Red probe (green), middle: Lysosomes stained with Lamp-2 antibody (red color) and right: merged channels LysoTracker and Lamp-2 of the same cell.

Figure SI5. Comparison of Pearson's and Manders' coefficients between live and fixed cells. The cells were exposed to SiO_2 Bodipy NP for 1h, stained with LysoTracker probe and imaged by CLSM. The same sample was fixed, stained for LAMP-2 and imaged. A total of 4 cells were analyzed for comparision, each cell has a representative color as it is shown in the coefficients plot. Each cell varies its colocalization coefficient for the NP with the lysosomes. This data demonstrate that the coefficients from live and fixed cells cannot be compared. Additionally, Pearson correlation coefficients is not enough to confirm the distribution of the fraction of the NP between the lysosomes.

Figure SI6. Comparison of raw integrated intensity between live and fixed cells. Top two images represent CLSM data from live cells (left) and fixed cells (right). The cells were stained with LysoTracker Red probe for 1h and imaged. The same cells were fixed and washed following a fixation protocol and imaged using the same settings. At the bottom the boxplot show the raw integrated densities of live and fixed cells. The whiskers represent the standard deviations. Analysis was performed in 8 individual cells for each experiment. All the images were analyzed using ImageJ. There is a significant decrease in LysoTracker Red intensity between live and fixed cells.

Video 1-SI-2.
J774A.1-LysoTracker

919　**Video SI1**. Continuous live cell imaging of J774A.1 cells, stained with LysoTracker Red for 24 h, showing
920　increased cell death over time.

922　**Script SI1.** Raw integrated densities

924　It was done comparing 10 cells for each lysotracker using image J comparing the raw intensity densities and the mean gray values,
925　selecting each cell with a mask using the next script and manual threshold:

```
926    //setTool("rectangle");
927    rename("source");
928    run("In [+]");
929    run("In [+]");
930    run("Duplicate...", "title=mask");
931    run("In [+]");
932    run("In [+]");
933    run("Median...", "radius=5");
934    setAutoThreshold("Default dark");
935    //run("Threshold...");
936    setOption("BlackBackground", true);
937    run("Convert to Mask");
938    run("Dilate");
939    run("Create Selection");
940    run("Make Inverse");
941    run("Make Inverse");
942    roiManager("Add");
943    roiManager("Select", 4);
944    roiManager("Select", 0);
945    roiManager("Select", 2);
946    selectWindow("source");
947    roiManager("Select", 0);
948    roiManager("Select", 2);
949    run("Measure");
```

950
951
952

953　Masked, automatically adjusted the threshold., selected the interest area removing background noise. Measure the intensities and
954　compare the raw integrated intensity for each treatment vs with phenol free

955
956

957　**Script SI2.**　Colocalization analysis

958
path1 = " __________________ **";**
timepoint= _________________ **;**
//channel 1 - NPS
ThresholdA = _________________ **;**
//channel 2 - LT
ThresholdB = _________________ **;**

```
965    Path2 = "                        /JACOP/";
966    filename= "                    .tif";
967
968    // IMPORTANT: DO THE NEXT BEFORE EACH MEASURMENT
969
970    //Check the channels order>
971    //channel 1 - NPS
972    //channel 2 - LT
973    // ADJUSTED IN CASE IT IS NEEDED THE ORDER AND CHECK EACH CHANNEL THE THRESHOLD
974
975    // #########
976    // OPEN DATA
977    // #########
978    open(filename);
979    selectWindow(filename);
980    rename("raw");
981    run("Select None");
982    roiManager("reset")
983    roiManager("Open", path1 +"/Roi.zip");
984    run("Colors...", "foreground=black background=black selection=yellow");
985    setBatchMode(true);
986    print("\\Clear");
987
988
989
990    // #######################
991    // CYCLE THROUGH EACH CELL
992    // #######################
993
994    for (cell = 0; cell < roiManager("count"); cell++){
995
996    //-------------------------------------------
997    // Cut out the cell and delete the background
998    //-------------------------------------------
999    selectWindow("raw");
1000    run("Enhance Contrast", "saturated=0.35");
1001    run("Select None");
1002    roiManager("Select", cell);
1003    run("Duplicate...", "duplicate title=[cell 1] frames="+timepoint);
1004    run("Make Inverse");
1005    run("Fill", "stack");
1006    run("Select None");
1007    rename("data");
1008    run("Split Channels");
1009
1010
1011
1012    //----------------------
1013    // Run the colocalization
1014    //----------------------
1015
1016    //without Costes randomizations
1017    run("JACoP ", "imga=C1-data imgb=C2-data thra="+ThresholdA+" thrb="+ThresholdB+" pearson overlap mm ccf=20 cytofluo ica
1018    costesrand=2-1-1000-0.001-0-false-true-true");
1019
1020
1021    // with Costes randomizations (takes long)
1022    //run("JACoP ", "imga=C1-data imgb=C2-data thra="+ThresholdA+" thrb="+ThresholdB+"  pearson overlap mm costesthr ccf=20 cytofluo ica
1023    costesrand=2-1-1000-0.001-0-false-true-true");
1024
1025
1026    //----------------------------------------------------------
1027    // Check the resulting windows, save them and then close them
1028    //----------------------------------------------------------
1029
```

```
1030          if (isOpen("Costes' threshold C1-data and C2-data")){
1031              selectWindow("Costes' threshold C1-data and C2-data");
1032          saveAs("Tiff", Path2 +cell+"_Costes' threshold C1-data and C2-data.tif");
1033                          close();
1034                          }
1035
1036              if (isOpen("Costes' mask")){
1037              selectWindow("Costes' mask");
1038          saveAs("Tiff", Path2 +cell+"_Costes' mask.tif");
1039                          close();
1040                          }
1041
1042          if (isOpen("Van Steensel's CCF between C1-data and C2-data")){
1043          selectWindow("Van Steensel's CCF between C1-data and C2-data");
1044                  Plot.setStyle(3, "blue,none,2.0,Line");
1045          saveAs("Tiff", Path2 +cell+"_Van Steensel's CCF between C1-data and C2-data.tif");
1046                          close();
1047                          }
1048
1049          if (isOpen("Cytofluorogram between C1-data and C2-data")){
1050          selectWindow("Cytofluorogram between C1-data and C2-data");
1051                  Plot.setStyle(0, "black,none,2.0,Dot");
1052          saveAs("Tiff", Path2 +cell+"_Cytofluorogram between C1-data and C2-data.tif");
1053                          close();
1054                          }
1055
1056              if (isOpen("ICA A (C1-data)")){
1057              selectWindow("ICA A (C1-data)");
1058              Plot.setStyle(1, "black,none,2.0,Dot");
1059          saveAs("Tiff", Path2 +cell+"_ICA A (C1-data).tif");
1060                          close();
1061                          }
1062
1063              if (isOpen("ICA B (C2-data)")){
1064              selectWindow("ICA B (C2-data)");
1065              Plot.setStyle(1, "black,none,2.0,Dot");
1066          saveAs("Tiff", Path2 +cell+"_ICA B (C2-data).tif");
1067                          close();
1068                          }
1069
1070          if (isOpen("Randomized images of C2-data")){
1071              selectWindow("Randomized images of C2-data");
1072          saveAs("Tiff", Path2 +cell+"_Randomized images of C2-data.tif");
1073                          close();
1074                          }
1075
1076          if (isOpen("Costes' method (C1-data & C2-data)")){
1077              selectWindow("Costes' method (C1-data & C2-data)");
1078                  Plot.setStyle(3, "blue,none,3.0,Line");
1079              Plot.setStyle(0, "white,none,1.0,Linehidden");
1080              Plot.setStyle(1, "black,none,2.0,Circle");
1081          saveAs("Tiff", Path2 +cell+"_Costes' method (C1-data & C2-data).tif");
1082                          close();
1083                          }
1084
1085              if (isOpen("C2-data")) {
1086              selectWindow("C2-data");
1087          saveAs("Tiff", Path2 +cell+"_C2-data.tif");
1088                          close();
1089                          }
1090              if (isOpen("C1-data")) {
1091              selectWindow("C1-data");
1092          saveAs("Tiff", Path2 +cell+"_C1-data.tif");
1093                          close();
1094                          }
```

```
1095
1096
1097
1098                              //---------------------------------------------------------------------
1099                              // Extract the coloc data from the Log window an post it in a result table
1100                              //---------------------------------------------------------------------
1101
1102                                      setResult("Cell", cell, "cell "+ cell);
1103                                      logString = getInfo("Log");
1104                                      selectWindow("Log");
1105                                  saveAs("Text", Path2 +cell+"_Log.txt");
1106                                      logString = getInfo("Log");
1107                                      selectWindow("Log");
1108          setResult("Pearsons correlation coefficient", cell, substring(logString, indexOf(logString, "Pearson's Coefficient:")+25,
1109                              indexOf(logString, "Overlap Coefficient:")-2));
1110          setResult("Overlap coefficient (no threshold)", cell, substring(logString, indexOf(logString, "Overlap Coefficient:")+23,
1111                              indexOf(logString, "r^2=k1xk2:")-2));
1112  setResult("k1 (no threshold)", cell, substring(logString, indexOf(logString, "r^2=k1xk2:")+14, indexOf(logString, "r^2=k1xk2:")+19));
1113          setResult("k2 (no threshold)", cell,  substring(logString, indexOf(logString, "r^2=k1xk2:")+23, indexOf(logString,
1114                              "r^2=k1xk2:")+28));
1115          setResult("Threshold A", cell, substring(logString, indexOf(logString, "Using thresholds (thrA=")+23, indexOf(logString, " and
1116                              thrB=")));
1117                      logString = substring(logString,  indexOf(logString, "r^2=k1xk2:")+28, lengthOf(logString));
1118          setResult("Threshold B", cell,  substring(logString, indexOf(logString, "thrB=")+5,indexOf(logString, "Overlap Coefficient:")-3));
1119                      logString = substring(logString,  indexOf(logString, "Using thresholds ")+28, lengthOf(logString));
1120          setResult("Overlap coefficient (with threshold)", cell, substring(logString, indexOf(logString, "Overlap Coefficient:")+23,
1121                              indexOf(logString, "r^2=k1xk2:")-2));
1122          setResult("k1 (with threshold)", cell, parseFloat(substring(logString, indexOf(logString, "r^2=k1xk2:")+14, indexOf(logString,
1123                              "r^2=k1xk2:")+19)));
1124          setResult("k2 (with threshold)", cell, parseFloat(substring(logString, indexOf(logString, "r^2=k1xk2:")+23, indexOf(logString,
1125                              "r^2=k1xk2:")+28)));
1126          setResult("Manders coefficient (without threshold) M1", cell, substring(logString, indexOf(logString, "Manders' Coefficients
1127                              (original")+37, indexOf(logString, " (fraction of A overlapping B)")));
1128          setResult("Manders coefficient (without threshold) M2", cell, substring(logString, indexOf(logString, " (fraction of A overlapping
1129                              B)")+34, indexOf(logString, " (fraction of B overlapping A)")));
1130          logString = substring(logString,  indexOf(logString, "Manders' Coefficients (using threshold value of"), lengthOf(logString));
1131          setResult("Manders coefficient (with threshold) M1", cell,  substring(logString, indexOf(logString, "M1=")+3, indexOf(logString, "
1132                              (fraction of A overlapping B)")));
1133          setResult("Manders coefficient (without threshold) M2", cell,  substring(logString, indexOf(logString, "M2=")+3, indexOf(logString,
1134                              " (fraction of B overlapping A)")));
1135          setResult("Van Steensel CCFmin", cell,  substring(logString, indexOf(logString, "CCF min.:")+10, indexOf(logString, "(obtained")-1));
1136          setResult("Van Steensel CCFmin peak value", cell, substring(logString, indexOf(logString, "dx=")+3, indexOf(logString, "CCF max.")-
1137                              2));
1138                      logString = substring(logString,  indexOf(logString, "CCF max.")-2, lengthOf(logString));
1139          setResult("Van Steensel CCFmax", cell, substring(logString, indexOf(logString, "CCF max.:")+10, indexOf(logString, "(obtained")-1));
1140          setResult("Van Steensel CCFmax peak value", cell, substring(logString, indexOf(logString, "dx=")+3, indexOf(logString, "Results for
1141                              fitting")-3));
1142          setResult("Cytofluorogram a", cell, substring(logString, indexOf(logString, "Cytofluorogram's parameters:")+32, indexOf(logString,
1143                              "b: ")-3));
1144          setResult("Cytofluorogram a", cell, substring(logString, indexOf(logString, "b: ")+3, indexOf(logString, "Correlation coefficient:")-
1145                              3));
1146          setResult("Cytofluorogram Correlation Coeffcient", cell, substring(logString, indexOf(logString, "Correlation coefficient: ")+25,
1147                              indexOf(logString, "Li's")-2));
1148          setResult("Li's ICQ", cell, substring(logString, indexOf(logString, "ICQ")+4, indexOf(logString, "Costes' randomization based
1149                              colocalization:")-2));
1150  setResult("Costes's R original ", cell, substring(logString, indexOf(logString, "r (original)")+13, indexOf(logString, "r (randomized)")-
1151                              1));
1152          setResult("Costes's R randomized ", cell, substring(logString, indexOf(logString, "r (randomized)")+15, indexOf(logString, "±")));
1153          setResult("Costes's R original SD", cell, substring(logString, indexOf(logString, "±")+1, indexOf(logString, "(calculated from the
1154                              fitted data")));
1155          setResult("Costes's R P-value in %", cell,substring(logString, indexOf(logString, "(calculated from the fitted data")+42,
1156                              indexOf(logString, "%")));
1157
1158
1159                              //----------------------
```

```
1160                              // Reset the log window
1161                               //----------------------
1162
1163                               print("\\Clear");
1164                            selectWindow("Log");
1165                               run("Close" );
1166
1167          //-----------------------------------------
1168          // End of cycle, goto next cell, if available
1169          //-----------------------------------------
1170
1171                                    }
1172
1173
1174                            // #########
1175                            // CLOSE DATA
1176                            // #########
1177
1178                          selectWindow("raw");
1179                                 close();
1180

1181
```