Evolutionary consequences of nascent multicellular life cycles

Jennifer T. Pentz[1*], Kathryn MacGillivray[2,3], James G. DuBose[2], Peter L. Conlin[2], Emma

Reinhardt[4], Eric Libby[5] and William C. Ratcliff[2*]

[1]Los Alamos National Laboratory, Los Alamos, NM, 87544, USA

[2]School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA 30332, USA

[3]Interdisciplinary Graduate Program in Quantitative Biosciences, Georgia Institute of

Technology, Atlanta, GA 30332, USA

[4]Department of Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599,

USA

[5]Department of Mathematics and Mathematical Statistics, Umeå University, Umeå 90187,

Sweden

* For correspondence: jenn.pentz@gmail.com and ratcliff@gatech.edu

Abstract

A key step in the evolutionary transition to multicellularity is the origin of multicellular groups as biological individuals capable of adaptation. Comparative work, supported by theory, suggests clonal development should facilitate this transition, though this hypothesis has never been tested in a single model system. We evolved 20 replicate populations of otherwise isogenic clonally-reproducing 'snowflake' yeast (Δ*ace2*/Δ*ace2*) and aggregative 'floc' yeast (*GAL1*p::*FLO1/ GAL1*p::*FLO1*) with daily selection for rapid growth in liquid media, which favors faster cell division, followed by selection for rapid sedimentation, which favors larger multicellular groups. While both genotypes adapted to this regime, growing faster and having higher survival during the group-selection phase, there was a stark difference in evolutionary dynamics. Aggregative floc yeast obtained nearly all of their increased fitness from faster growth, not improved group survival; indicating that selection acted primarily at the level of cells. In contrast, clonal snowflake yeast mainly benefited from higher group-dependent fitness, indicating a shift in the level of biological individuality from cells to groups. Through genome sequencing and mathematical modeling, we show that the genetic bottlenecks in a clonal life cycle also drive much higher rates of genetic drift—a result with complex implications for this evolutionary transition. Our results highlight the central role that early multicellular life cycles play in the process of multicellular adaptation.

Introduction

Multicellularity has evolved more than 50 times across the tree of life [1,2]. Multicellular organisms vary widely in their life cycles, ecological niches, and traits, and there are few features shared by all members of this diverse group [3]. However, it is generally accepted that nascent multicellular lineages start out relatively simple (*e.g.*, as undifferentiated groups of cells), gradually evolving increased organismal integration and functionality [2,4-7]. In some cases, multicellular groups that gain the capacity for group reproduction undergo a shift in evolutionary mode: the origin of multicellular Darwinian individuality [8-13]. This shift, from cell-level traits being the focal point of adaptation in the unicellular ancestor, to group-level traits becoming the focal point of adaptation in a nascent multicellular lineage, represents a crucial tipping point in evolutionary dynamics. In this paper, we examine the impact of early multicellular life cycles on the capacity for groups to serve as evolutionary units capable of adaptation.

There are two basic routes to forming a multicellular body. Nascent multicellular organisms employing either of these routes to group formation possess distinct multicellular life cycles. Individual cells can 'stay together' by forming permanent cell-cell bonds, forming clonal clusters that exhibit little within-group genetic variation. Alternatively, free-living single cells can 'come together', or aggregate, and often evolve to do so in response to some stimuli, such as starvation [14-18]. Multicellularity has evolved multiple times via both staying together and coming together [2,14,19], but 'complex multicellularity' [6] (*e.g.*, plants, animals, fungi, red algae and brown algae) has only evolved in lineages that develop clonally [4,6,20]). Evolutionary theory explains this observation as a consequence of social evolution: clonal life cycles are a simple and powerful mechanism that 'de-Darwinizes' cells [9,10,21] while Darwinizing multicellular groups. By limiting within-group genetic diversity, clonal development prevents intra-organism genetic conflict, as

there is little standing within-group genetic variation for selection to act on [11,22,23]. Any genetic

variation that arises due to mutation gets partitioned among multicellular offspring, allowing

selection to act on the group-level effects of *de novo* mutations [24]. Clonal groups align the fitness

interests of cells and groups, allowing cells to evolve altruistic social traits necessary for cellular

differentiation [9,25]. Finally, organisms that aggregate from a free-living state may experience

contrasting selection on the fitness of free-living single cells and the fitness of these cells in a

multicellular group [26].

While the numerous origins of multicellularity provide a uniquely rich set of natural

experiments to examine correlations between developmental mode and organismal complexity

[1,4,27,28], there is no direct evidence that clonal development has played a causal role in the transition

to multicellular individuality, and subsequent evolution of organismal complexity. Indeed, a

number of alternative explanations exist. For example, clonal development has evolved more

frequently in aquatic environments, while aggregation has evolved more often in terrestrial

environments [14,20]. The lower complexity of aggregative organisms may be due simply to different

life history selection in these fundamentally different environments, rather than evolutionary

constraints due to within-organism genetic diversity. Alternatively, the potential for cellular

differentiation appears to be highly contingent on the cell biology and behavioral repertoire of the

unicellular ancestor [4,29-32]. The fact that complex multicellularity evolved in just five lineages may

reflect historical contingency in developmental plasticity in these specific lineages, rather than

evolutionary consequences of early developmental mode. Progress has been limited by the fact

that all known transitions to multicellularity occurred in the deep past (>200 MYA [2,33]), obscuring

the evolutionary dynamics of early multicellular evolution in extant lineages. Experiments directly

comparing the evolutionary consequences of developmental mode have not yet been conducted,

in part due to a lack of experimentally-tractable model systems that can be induced to undergo

either clonal or aggregative development.

Here we circumvent these constraints by synthetically generating an isogenic yeast model

system capable of either clonal or aggregative development. We created clonal 'snowflake yeast'

by knocking out *ACE2* in a unicellular ancestor [34,35], and aggregative floc yeast by placing the

dominant *FLO1* gene under transcriptional control of the *GAL1* promoter [35,36]. While these strains

differ only in these two genes, these differences produce two fundamentally different life cycles:

obligately-multicellular snowflake yeast undergo a unicellular genetic bottleneck during ontogeny,

making them clonal [34,37,38], while floc yeast can form genetically diverse aggregates [36]. We evolved

20 populations of each genotype for 24 weeks with galactose as the main carbon source, ensuring

robust flocculation, selecting daily for both faster growth and increased multicellular size. We

chose this selective regime because it is a simple and powerful way to examine the evolutionary

consequences of selection acting simultaneously on both cell-level traits (such as growth rate) and

group-level traits (such as settling rate). While buoyancy regulation may be an important driver of

multicellularity in some lineages [39], we chose this selective regime because it is an efficient way

to select on multicellular size, a fundamentally important multicellular trait [40].

In our experiment, both floc and snowflake yeast adapted to this fluctuating environment,

settling faster and growing faster. However, competition experiments with their ancestors reveal

fundamentally different modes of adaptation. In clonal snowflake yeast, groups of cells served as

the primary evolutionary units, gaining nearly all their increased fitness during the settling

selection phase of the experiment. In contrast, aggregative floc gained most of their fitness as the

cell-level, with evolved floc growing faster but possessing no measurable advantage during settling

selection when competing against their ancestor. Snowflake yeast, but not floc, are evolving as

primarily multicellular Darwinian individuals. Clonal development, however, resulted in far more

genetic drift than aggregation. Mathematical modeling suggests that this is the result of a key

difference in their life cycles. In floc, a rare beneficial mutant can disperse into many groups, while

in clonal snowflake yeast it is constrained to a few, exposing them to greater sampling error during

group selection. This is a general property of clonal multicellular life cycles, scaling with the

number of cells within the organism prior to reproduction. Together, our results show a simple

difference in the mechanism of group formation, whether cells adhere with reformable bonds or

adhere permanently, can fundamentally change their subsequent evolutionary dynamics and

impact the level of biological individuality [41].

Results

Experimental evolution

Our selection regime involved 24h of batch culture followed by daily selection for rapid

sedimentation in liquid media [37]. This selective scheme has previously been shown to promote

multicellular adaption in snowflake yeast [37,38,42,43] and has led to increases in cluster size of up to

20,000-fold over 600 consecutive rounds of selection [43]. We have previously quantified the effect

of settling selection on snowflake yeast by using a variety of tools (*i.e.*, microscopy and flow

cytometry [37,38,42,43]) that cannot be used for floc, because floc aggregates form dynamically as the

clusters are settling. Thus, we developed a method to measure the settling rate of both floc and

snowflake yeast populations during sedimentation [35], calculating the displacement of biomass as

they settle via high-resolution video (see Methods). We measured the settling speed of each of the

40 populations weekly over the course of the 24-week experiment (Figure 1A,B). Both snowflake

yeast and floc evolved to settle significantly faster (one-way ANOVA; floc: $F_{23,456}=16.57$,

$p<0.0001$; snowflake: $F_{23,456}=13.65$, $p<0.0001$, pairwise differences assessed with Tukey's post-

doc HSD with $\alpha=0.05$). Floc, however, exhibited a much larger increase in settling rate than

snowflake yeast (12-fold increase vs 1.3-fold increase after 24 weeks, respectively, Figure 1C).

Figure 1 - Dynamics of settling rate evolution in snowflake and floc populations. The settling
rate (as determined by time-lapse imaging, see Methods) of 40 independently evolving populations
of snowflake yeast (A) and floc yeast (b) over 24 weeks of evolution. Settling rate was measured
for each population weekly. Snowflake yeast evolved to settle 30% faster over 24 weeks of
evolution, while floc evolved to settle an average of 12-fold faster (C).

We next examined the traits underlying increased settling speed. Using flow cytometry, we

measured the size of 24-week evolved snowflake yeast. Biomass weighted mean forward scatter

(a proxy for group size) increased by an average of 33% (Figure 2A,B; one-sided t-test, $t=7.48$,

$n=24$, $p<0.0001$; non-overlayed histograms for individual snowflake isolates shown in

Supplementary Figure 2). Evolved snowflake isolates retain the same basic growth form (Figure

2D,E). Examining one lineage through time, we found that size appeared to plateau after 8-10

weeks of settling selection (Figure 2C), which is consistent with previous work in this model

system where aerobic metabolism, and the corresponding reliance on diffused oxygen for growth,

strongly inhibits the evolution of increased size [43,44]. By 24 weeks, floc yeast appeared to be

aggregating far more efficiently than their ancestors. To quantify this, we measured the coefficient

of variation in pixel opacity in a well-mixed population just prior to settling selection (Figure 2F).

This standardized variance measurement is as a proxy for flocculation efficiency, as when more

cells are in flocs, the biomass will be more heterogeneously distributed within the cuvette. Thirteen

populations showed significantly increased flocculation efficiency relative to the ancestor (one-

way ANOVA, $F_{(20,46)}$=45.53, P<0.0001, pairwise differences assessed with Tukey's honestly

significant difference [HSD] with α = 0.05). This is also noticeable as a reduction in the density of

free un-flocculated unicells, see examples of the ancestor (G) and a representative 24-week isolate

(H). Thus, floc yeast evolved to settle more rapidly by forming larger aggregative groups and

reducing the proportion of non-aggregated cells in the population (Figure 2G,H).

Figure 2 – Both floc and snowflake yeast evolved to form larger groups. A,B) Snowflake yeast
increased in settling rate by evolving larger cluster size, but snowflake yeast reach their maximum
cluster size after 8-10 weeks of evolution. Shown in (C) are the size distribution of a representative
snowflake population (S8) every three weeks for the duration of the experiment. Relative to their
ancestor (D), evolved snowflake yeast (E) are larger but retain a similar pattern of cellular
attachment (insets). F) We estimated the flocculation efficiency of ancestral and evolved floc by
measuring the coefficient of variation in pixel intensity within cuvettes of a well-mixed population
taken from shaking incubation (data on top, images from each population on the bottom).
Flocculation efficiency was significantly higher in 13/20 evolved populations (significance at the
overall $\alpha= 0.05$ level denoted by asterisks). Shown are the mean and standard deviation of 4
biological replicates normalized to the mean of the ancestor. Representative images of ancestral
(G) and evolved (H) floc yeast (genotype F4), showing fewer planktonic cells in the evolved isolate
with higher flocculation efficiency.

Partitioning fitness between growth and multicellular-dependent survival

A common way to analyze ETIs is to use the Price equation to partition fitness arising from

selection acting at distinct levels, i.e., cells and groups [41,45]. Our system is not amenable to this

kind of decomposition given the dynamic nature of flocs: groups rapidly form and fuse during

settling selection, changing in size and composition until they either succeed at joining the pellet

at the bottom of the tube, where they rapidly adhere to these cells, or fail to do so and are discarded.

The fluid nature of flocs, and corresponding difficulty of measuring their traits without changing

the traits we seek to measure, prevents us from quantifying the genetic composition and fitness of

flocs during settling selection- data that is necessary for a Price equation fitness decomposition.

Fortunately, we can still gain insight into how selection is acting on cell and group-level traits by

leveraging the biphasic nature of our experiment.

Fitness in our experiment depends on growth rate during the 24h of batch culture when the

population is competing for resources, and survival during the subsequent settling selection step

[35,37,46]. Growth rate during batch culture is a classic trait under strong selection during experimental

evolution [47]. The main way that a lineage can increase growth rates is by increasing the rate at

which cells divide. In contrast, the main way in which a lineage can increase survival during

settling selection is through changes in the phenotype of multicellular groups (*i.e.*, increasing their

size, packing density, or hydrodynamic profile [42]). Thus, if populations evolve higher fitness

during the growth phase, this can be taken as evidence that selection is acting effectively at the

cell-level. Correspondingly, fitness improvement during the settling phase can be taken as

evidence that selection is acting at the group-level.

We measured the fitness of evolved isolates (one isolate per 24-week population, or 40

isolates in total) in competition against their unevolved ancestor across one 24h period of growth,

and one round of settling selection. Floc yeast showed a significant fitness advantage during

growth (Figure 3A; mean relative fitness=1.24, one-sample *t*-test, $n=20$ $t=10.55$, $p<0.0001$), but

despite settling an average of 12-fold faster than their ancestor when grown in monoculture, were

only slightly significantly more fit than their ancestor during settling selection (Figure 3A; mean

relative fitness=1.03, one-sample *t*-test, n=20, $t=1.633$, $p=0.12$). Snowflake yeast displayed the

opposite behavior, possessing a 5.5% fitness benefit during growth (Figure 3A; one-sample *t*-test,

n=20, $t=8.374$, $p<0.0001$) and a 22% fitness advantage over their ancestor during settling selection

(Figure 3A; mean relative fitness=1.28, one-sample *t*-test, $n=20$, $t=10.29$, $p<0.0001$).

Figure 3 – Examining fitness during growth and group-dependent competition. A) When competing against their ancestors, snowflake and floc yeast evolved distinct differences in their competitiveness during the growth and competition for survival during settling selection. Floc yeast obtained a 30% fitness advantage during growth, but showed no detectable increase in fitness during settling selection. Alternatively, snowflake yeast had a 22% fitness advantage during settling but only a 6% advantage during growth. B) When competing against their ancestor, representative floc isolates formed groups with low genetic assortment. This may explain the decoupling between dramatically increased group sedimentation rate (Figure 1B), and the negligible increase in fitness during settling selection. Representative flocs are shown in C&D, with the ancestor competing against itself, or a RFP-labeled ancestor competing against an GFP-labeled 24-week evolved isolate (genotype F10).

Despite floc as a treatment not having higher fitness during settling selection, isolates from

2/20 replicate populations did have detectably higher fitness than their ancestor when competing

during settling selection ($F_{20,63}$ = 4.528, $p<0.0001$, multiple comparisons controlled by a

Bonferroni correction with overall α = 0.05, the same two populations were identified using a

Dunnett's test against an ancestor:ancestor control). In contrast, 12/20 snowflake populations

significantly increased fitness during settling selection ($F_{20,63}$ = 4.89, $p<0.0001$, multiple

comparisons controlled using a Dunnett test against ancestor:ancestor control). To determine why

floc obtained such marginal fitness benefit during settling selection despite evolving such a large

increase in their rate of flocculation, we measured the genetic structure of flocs formed by five 24-

week isolates from across the range of settling speeds by calculating their assortment [48], a scaled

statistic of genotypic enrichment relative to what would be expected by chance, which ranges from

-1 to 1 (representative images of flocs, compressed to a single-cell thickness, shown in Figure

3C,D). Multicellular adaptation requires a positive correlation between group phenotype and

underlying cell-level genotype; without this, selection acting on groups cannot drive changes in

allele frequency [49,50]. Genotypes that achieve high assortment should thus have greater potential

for multicellular adaptation.

Overall assortment in floc was relatively low (mean of five randomly-selected strains from

Figure 3B was 0.06), though there was significant among-strain variation (one-way ANOVA,

$F_{4,10}$=4.159, p=0.008). Low assortment impedes the potential for selection acting on groups to

drive changes in allele frequencies, explaining why evolved floc yeast obtained little fitness benefit

from their remarkably improved sedimentation rates. In contrast, snowflake yeast canonically have

an assortment of 1 (when competing two strains, every group is entirely clonal (Supplementary

Figure 1A; [35]), allowing for selection acting on emergent group-level traits (*i.e.*, settling speed) to

act on underlying genetic mechanisms.

Genomic analysis provides insight into evolutionary dynamics

We sequenced the genome of one isolate per 24-week population (40 isolates in total), representing

about 700 generations of evolution (Supplementary Figure 3). Floc yeast accumulated more

mutations than snowflake yeast, an average of 5 vs 3 mutations per genotype, respectively (Figure

4A; two-tailed *t*-test, *t*=2.49, *n*=40, *p*=0.017). All mutations are listed in Supplemental Table 1

(Snowflake) and Supplemental Table 2 (Floc). To develop insight into how each life cycle affected

the potential for selection to act on mutations, we compared the frequency of different types of

point mutations (which constituted the large majority of total mutations; Figure 4B) to those

predicted by a null model of evolution lacking selection. Specifically, we generated a distribution

of the expected frequency of different types of mutation (i.e., missense, nonsense, synonymous, or

intergenic) in yeast by simulating 100,000 mutations via Mutation-Simulator [51], then ran 1,000

bootstrap simulations in which we sampled the actual number of mutations identified in both floc

(104 mutations total) and snowflake (69 mutations total) populations. Then we compared the

observed frequency of these four types of mutation in both floc and snowflake populations to this

null distribution, which is the distribution of mutations expected without selection (i.e., under

genetic drift).

The overall pattern of mutations observed in snowflake yeast were not distinguishable from

drift, with the number of mutations in each of the four categories being near the mean of the

expectation under selective neutrality (Figure 4C). In contrast, the mutations in floc yeast show a

strong signature of selection. We observed more missense mutations than expected based on drift

(there was a 3% chance of observing at least this many missense mutations in our null model).

Similarly, we observed fewer synonymous mutations and fewer mutations in intergenic regions

than expected based on our null model (in both cases, there was a 3% chance of observing this

many mutations or fewer in our null model). Nonsense mutations occur infrequently, and do not

provide a sufficiently large sample size for analysis. Because we did not have a prior expectation

about the deviation from the null, each of the above tests should be treated as a two-sided test,

meaning there is a 6% chance of observing a deviation in either direction at least as large as the

one observed for the three categories of mutation described above, even when the null hypothesis

of no selection is true. Taken together, our results show that selection was acting strongly on the

mutations found in floc, but not snowflake yeast.

Figure 4 – Examining mutations for signatures of selection. After 24 weeks, floc yeast
accumulated more mutations on average than snowflake yeast (5 versus 3, respectively, A). We
categorized these mutations into four broad classes (B), then compared the number of each type of
mutation to expectations based on a simulation of neutral evolution (C). We cannot distinguish the
overall pattern of mutations in snowflake yeast from neutral expectations. In contrast, floc yeast

showed strong evidence of selection, with more missense mutations and fewer synonymous and
intergenic mutations than expected. The number in upper left-hand corner of each subplot reports
the shaded are of each distribution, which is the proportion of simulated runs in that observed a
result at least as extreme as the observed number of mutations of that type we identified, pooling
across all 20 floc and snowflake genotypes.

The analysis above only examined the overall statistical properties of mutations, and should

not be interpreted to mean that snowflake yeast did not undergo adaptive evolution. Indeed, we

see that all evolved isolates taken from all 20 populations have evolved to form larger groups

(Figure 2A-E, Supplementary Figure 2) that settle faster (Figure 1C) and are much more fit (Figure

3A). Some of the mutations identified within snowflake yeast are putatively adaptive, such as

missense mutations in the cell cycle (*i.e.*, *ULP1*, *SLK19*; Supplemental Table 1) or mutations affect

cellular morphology (*i.e.*, *AYR1, ACM1, GSC2, CHC1, ARP7, HKR1*), which prior work in

snowflake yeast has shown are a common mode of evolving larger group size [24,43,44,52]. We saw

only a single *GAL2* mutant (in contrast to seven arising independently in floc, Supplementary Table

2), which is consistent with less growth rate adaptation in snowflake yeast. Indeed, it is plausible

that selection acting on multicellular traits also contributed to genetic drift through the hitchhiking

of otherwise non-adaptive alleles (an outcome that has been seen previously in yeast selection

experiments [53]).

Our results (Figure 3) suggest that faster settling in floc, in contrast to snowflake yeast, is

only marginally adaptive. Yet if that is the case, why would floc convergently evolve to settle an

average of 12 times faster than their ancestor (Figure 1)? We see two potential explanations, which

are not mutually exclusive. The first is that low genetic assortment during flocculation reduces,

but does not eliminate, the benefit of faster settling. Indeed, we found that 24-week evolved floc

were an average of 10% more fit than their ancestor during settling selection, though this difference

was not statistically significant for the treatment as a whole. Consistent with flocculation itself

being under selection, we saw five parallel missense mutations in *FLO1*, and four in *MSN1,* a

transcriptional activator of another major flocculin gene, *FLO11* [54-56]; Supplemental Table 2).

Greater statistical power during competitions may be required to detect the benefit of increased

flocculation if it exists, however.

Alternatively, it may be that faster settling has evolved as a pleiotropic consequence of

increased growth rate. In our experiments, mutations that increase growth rate by altering carbon

metabolism may also increase flocculation efficiency as a side-effect because *FLO1* expression is

under the control of a *GAL1* promoter. We saw seven parallel missense mutations in *GAL2,* a

transmembrane galactose permease, which may increase the amount of galactose entering the cell.

As the main carbon source in the medium and promoter of *FLO1*, this may both increase growth

rate and flocculation rate simultaneously. We conducted an additional experiment to control for

this possibility, evolving five floc genotypes for eight weeks with daily selection for growth rate,

but not settling selection, in the same growth medium containing galactose. These controls evolved

a 20.7% increase in fitness during the growth phase, but evolved a 40% decrease in flocculation

efficiency (Supplementary Figure 4). We conclude that cell-level adaptation to growth on

galactose does not drive increased flocculation as a pleiotropic side-effect. It is thus likely that the

dramatically increased flocculation efficiency seen in our main experiment (Figure 1B) is an

adaptation, albeit one that provides only a relatively small advantage (Figure 3A).

Bottlenecks in the life cycle: modeling adaptation in aggregative and clonally-developing life
cycles

To reconcile the results above, in which clonal multicellularity both facilitates multicellular

adaptation but results in more genetic drift, we developed a theoretical model. Here we consider

the fate of a rare mutant that arises during the population growth phase. In order for a beneficial

mutant to have a chance of fixing in the population, it must be in a group that gets selected. If we

ignore that cells are partitioned across groups, then the probability that an entire mutant lineage

goes extinct (call it p_e) is described by a hypergeometric distribution: $p_e = \prod_{i=0}^{fN} \frac{N-k-i}{N-i}$, where N

is the total population size, k is the total number of mutants, and f is the fraction of the population

selected. If N is large, we can simplify this to $p_e \approx e^{-kf/(1-f)}$. Here the units of the key parameter

k are determined solely by the units of selection: so if groups of cells are selected, then k is the

number of groups containing at least one mutant. The way in which mutant cells are partitioned

across groups heavily impacts the value of k. In the absence of any preferential assortment, (i.e.,

an idealized form of aggregative development), the distribution of rare mutants across groups

follows the size distribution of groups, so groups with more cells are expected to contain more

mutants. For simplicity, suppose groups are all roughly the same size. When the mutant is rare and

the number of groups greatly exceeds the number of mutant cells, then the number of groups

containing a rare mutant is approximately the same as the number of mutant cells. In contrast,

when groups are highly assorted (e.g., clonal development), the initial rare mutant and all of its

descendants will be represented in a small number of groups, reducing k by a factor that

corresponds to mean group size. To get a sense of this effect, suppose a mutant arises at the

beginning of the growth phase. The population then grows 100-fold, resulting in 100 mutant cells.

Assuming the average size of a group is 50 cells, then if 1% of the population survives settling

selection, the probability that the mutant lineage goes extinct in a population that develops clonally

is 98%, while it is just 36.4% for a population developing aggregatively.

To assess the robustness of our theoretical predictions, we used experimental data

measuring the distribution of group sizes in floc and snowflake yeast and simulated different

mutations arising in these populations. We considered mutations that either: 1. Alter cell growth

by a factor $(1+s_c)$ where positive s_c corresponds to a beneficial mutation, or 2. Improve the settling

rate of that group by a factor of $s_g \cdot m$, where m is the number of mutant cells. We assume the best

scenario for a mutation to fix, *i.e.,* it arises at the very start of an exponential growth phase. We

then used experimentally-observed size distributions of groups to allocate the offspring of this

original mutant cell. For simplicity, in the well-mixed aggregative life cycle, we randomly selected

cells to be mutant, thereby causing larger groups to have more mutant cells (though keeping the

proportion of mutants similar). For the clonal life cycle, we randomly selected groups, allocating

mutant cells to it until we either ran out of cells, or had to choose a new group (biologically, these

would be propagules of the first cluster, as they all descend from a common mutant ancestor).

After allocating mutants, we simulated settling selection by probabilistically selecting groups, with

survival weighted linearly by size. We iterated over five rounds of growth and size-dependent

selection.

We recapitulate key dynamics from our experiments in this simple model. In the

aggregative life cycle, selection readily acts on mutations that affect cellular growth rate (either

favoring beneficial mutants or purging those that are deleterious; Figure 5A). In contrast, the clonal

life cycle struggles to capture growth beneficial mutations- even mutants that doubled cellular

growth rates went extinct ~40% of the time. When we examine mutations that increase group

survival, we find that while they rarely go extinct in aggregative groups (Figure 5B), neither can

they be efficiently selected upon (Figure 5C&D). In the clonal life cycle, even strongly group-

beneficial mutants (*e.g.*, increasing the probability of group survival 10-fold) go extinct far more

often than not, as they are extremely susceptible to being lost in the first several rounds of the

simulation when they are found in only a small number of groups. If they escape being lost by

drift, however, they rapidly fix in the population (Figure 5C&D). The clonal life cycle thus allows

selection to act, albeit inefficiently, on group-survival beneficial mutations, while the aggregative

life cycle is strongly biased in favor of capturing growth-enhancing mutations.

Figure 5. Examining clonal vs. aggregative development in a simple model. We examined the
ability of organisms that develop clonally or aggregatively to fix mutations that increase growth
rate (A) or group survival (B-D). Aggregative organisms readily fixed growth beneficial mutations
(A), but were unable to act upon group-beneficial mutations (B&C). In contrast, clonally-
developing were capable of fixing mutations that improve both growth and the survival of their
group, though they faced far more stochastic loss of each type of beneficial mutation than floc. B-
D show that clonal development stochastically loses even strongly group beneficial mutants a
majority of the time due to sampling error (in D, $s_g=10$), but when these mutations persist beyond
the first two rounds of selection, they rapidly fix. Note that 88% of the beneficial mutants within
the clonal lineages went extinct in D.

Discussion

      During the evolutionary transition to multicellularity, groups of cells become Darwinian

individuals, capable of reproducing and possessing heritable variation in multicellular traits that

affect fitness [11,21,23,57-59]. Using a simple yeast model system, in which we engineered either an

aggregative life cycle or one in which groups develop clonally, we used experimental evolution to

examine how developmental mode impacts this evolutionary transition in individuality. Our results

highlight how simple differences in the nature of cellular bonds required for group formation can

instantiate distinct proto-multicellular life cycles with divergent evolutionary outcomes. The

clonal multicellular life cycle, but not aggregation, allowed for efficient multicellular adaptation

in response to selection on group-level traits, suggesting a shift in the level of biological

individuality.

While both clonally-developing snowflake yeast and aggregative floc evolved to settle

faster in response to settling selection (forming larger, faster-settling clusters or flocs,

respectively), only snowflake yeast showed a shift in the balance of selection from cells to groups.

When competing evolved isolates against their ancestor, we found that snowflake yeast improved

their fitness mainly during settling selection (a phase of the culture cycle that primarily reflects

group-level properties), not the growth phase of the experiment (a period of the culture cycle that

mainly reflects changes in cellular growth rate). Floc yeast adapted in the opposite manner, gaining

a majority of their fitness increase during the growth phase, with little detectable increase in fitness

during settling selection. When floc cells aggregate, they form groups with a genetic composition

resembling the population as a whole. This low genetic assortment diminishes the ability for group-

level selection to drive evolutionary change, impeding the transition to multicellular individuality.

The differential pattern of mutations in floc and snowflake yeast, informed by a simple

mathematical model, highlights another significant impact of the mechanism of group formation:

the rate of genetic drift. In floc yeast, natural selection efficiently acted on novel mutations, with

fewer synonymous and intergenic mutations than expected by chance, and more missense

mutations than expected by chance. As our method detects a divergence in the proportion of each

mutation type from neutrality, these results could either be the result of selection against

synonymous and intergenic mutations, selection for missense mutations, or both. While

synonymous mutations may be non-neutral [60,61], the vast majority of mutations that fix during

microbial experimental evolution are nonsynonymous [53,60], as these are often subject to positive

selection [62,63]. When we query all putative loss of function mutations that arose in our experiment

in a genome-wide deletion collection grown on galactose [64], we find that 46% of the mutations in

floc are predicted to increase growth rates on this carbon source, in contrast to just 35% in

snowflake yeast. This is consistent with selection on mutations affecting cellular growth rate being

more efficacious in floc than snowflake yeast.

Snowflake yeast, on the other hand, showed no statistical deviation from the expectations

of neutrality. While initially surprising, this finding is consistent with group formation and

selection playing a more central role in their evolution. Our model demonstrates that clonal

development allows selection on group-level traits to be more efficacious, but simultaneously

drives a much a higher overall rate of genetic drift, due to the higher probability that novel mutants

will be lost during the group selection phase of the experiment. This result echoes prior theoretical

work showing that the effective population size in a metapopulation is greatly reduced by strong

among-deme selection, limited migration, and a small number of colonizing cells relative to

carrying capacity [65].

Our model highlights how the severity of genetic bottlenecking due to clonal development

scales with organismal size, with larger organisms exacerbating drift. Relative to the standards of

most extant multicellular organisms [66], snowflake yeast are small. As a result, the realized disparity

between clonal and aggregative development in many organisms may be considerably greater than

what was observed in this paper. It is not yet clear what impact this has on the evolution of complex

multicellularity. Reducing the ability for selection to act efficiently on cell-level fitness (*i.e.*,

growth rate) may relax constraints on the evolution of novel group-level traits that come at a cost

to growth [33]. In addition, relaxed selection can facilitate a broader search of potential trait space,

which may open up novel adaptive routes that would have been constrained by strong selection.

This may be especially important during an evolutionary transition in individuality, as the nature

of the organism is being fundamentally reshaped and genotypes are presumably far from potential

fitness optima. Indeed, relaxed selection appears to have played a central role in the evolution of

larger and more complex genomes within eukaryotes [67], freeing these populations from the ruthless

efficiency of strong purifying selection.

While floc and snowflake yeast differ starkly in evolutionary mode, our experiment was

too short to examine long-term evolutionary consequences. Large scale change is possible in our

system, but the evolution of macroscopic size appears to be much more challenging for snowflake

yeast than in floc, requiring more innovation. Snowflake yeast must resolve fundamental

biophysical challenges before they can form large groups, evolving larger size via three distinct

biophysical mechanisms: reducing cellular density within clusters to limit cell-cell jamming,

increasing the size and strength of the bonds connecting mother and daughter cells, and evolving

branch entanglement so that group fracture requires breaking many cellular bonds, not just one [43].

These innovations took 400-600 serial transfers to evolve, and resulted in snowflake yeast that

were 20,000 times larger than their ancestor, and 10,000 times more biomechanically tough. In

contrast, evolving macroscopic size (Figure 2F) appears somewhat trivial for floc, likely due the

relative ease and efficacy of increasing the strength of cell-cell adhesion.

There is an emerging consensus that clonal multicellular development provides the

foundation upon which sustained multicellular adaptation may ultimately drive the evolution of

dramatically increased multicellular complexity. This is supported by phylogenetic inference

[20,28,30], first-principles theory [17,26,34,57,68,69], and now, experimental evolution. In this paper, we

show that only clonally-developing snowflake yeast exhibited a shift in biological individuality,

such that multicellular groups were the primary unit of selection, with increased fitness arising

mainly via the group-selection phase of the life cycle. Aggregative floc, in contrast, behaved as a

social unicellular organism, adapting in both phases of the experiment, but low assortment limited

the potential fitness benefits of selection acting on groups. Persistent low genetic assortment

during group formation presents a challenge to this evolutionary transition, blunting the impact of

group-level selection, enabling within-group genetic conflict, and ultimately inhibiting a shift in

the level of Darwinian individuality from cells to groups.

And yet, aggregative multicellularity is clearly a successful life history strategy, evolving

repeatedly in diverse lineages despite the issues described above. Our results thus raise a number

of key questions for future research: how do lineages evolving aggregative multicellularity

overcome the constraint of limited assortment? Is active kin recognition a pre-requisite to the

evolution of multicellular individuality for aggregative life cycle, or can spatial structure generated

by a patchy environment or viscous media like soil [48] provide sufficiently high assortment for

efficient, sustained multicellular adaptation? To this latter point, most lineages of aggregative

multicellularity are terrestrial [14,20] - does this reflect the potential of for a highly structured

environment to scaffold the origin of aggregative multicellularity [70], or are there simply more

ecological opportunities for a biphasic life cycle in terrestrial habitats? We still have only a

rudimentary understanding of how key components of early multicellular life cycles (e.g.,

developmental mode, level of genetic assortment within groups, type and strength of selection on

cellular and group-level traits, etc.) influence the evolution of multicellular complexity.

Developing a robust, bottom-up theory of multicellular evolution from first-principles theory will

require the integration of multiple approaches- including mathematical theory, fieldwork, and

experiments incorporating both synthetic and naturally-evolved multicellular organisms.

Methods

Strains and media

Table 1 – Strains used in this study.

Strain	Relevant Genotype	Reference
Snowflake	Δ*ace2*::HYGMX / Δ*ace2*::HYGMX	Pentz *et al*. (2020)
Floc	Δ*ura3*::*KAN-GAL1*p*::FLO1* / Δ*ura3*::*KAN-GAL1*p*::FLO1*	Pentz *et al*. (2020)
Snowflake-GFP	Δ*lys2*::*TEF2*p-yeGFP / Δ*lys2*::*TEF2*p-yeGFP	Pentz *et al*. (2020)
Floc-GFP	Δ*lys2*::*TEF2*p-yeGFP / Δ*lys2*::*TEF2*p-yeGFP	Pentz *et al*. (2020)

Strains used in this study are listed in Table 1 and the construction of these is described in

Pentz *et al*. (2020). Briefly, flocculant yeast were created by replacing the *URA3* open reading

frame (ORF) with the *KAN-GAL1p::FLO1* cassette [36] and snowflake yeast were created by

replacing the *ACE2* ORF with the drug marker, *KANMX*. These genotypes were created from the

same homozygous diploid unicellular ancestor (strain Y55), so these strains differ only in their

mode of cluster formation. All experiments were performed in rich medium composed of a mix of

glucose and galactose (YPGal+Dex; per liter; 18 g galactose, 2 g glucose, 20 g peptone, 10 g yeast

extract), shaking at 250 rpm at 30°C. These growth conditions yield clusters of similar size after

24 h of growth.

Experimental evolution

Twenty replicate populations of both snowflake and floc yeast were initiated into 10 mL

of YPGal+Dex from a single clone and grown overnight. Every 24 h, each population (40

populations total) was subjected to daily selection for settling for 5 minutes on the bench as

described in Ratcliff *et al*. (2012). Briefly, 1.5 mL of each overnight culture was placed into a 2

mL microcentrifuge tube, left to settle on the bench for 5 minutes, after which the top 1.4 mL of

culture was discarded. The remaining 100 µL of the pellet was transferred to a fresh tube

containing 10 mL of culture media for the next round of growth and selection. Our selection regime

yielded ~4-5 generations per day for both snowflake and floc yeast (Supplementary Figure 3).

Every 7 days, whole populations were cryogenically stored at -80°C. As a control, five populations

of snowflake and floc yeast were evolved without settling selection for eight weeks. Specifically,

five replicates of both snowflake and floc yeast were initiated as described above. Every 24 h, each

population (10 populations total) was briefly vortexed, then 100 µL was transferred to 10 mL of

fresh medium. In the absence of settling selection, this would result in 6.6 generations of growth

per day. The number of generations realized per transfer was lower than this upper ceiling

(Supplementary Figure 3), because groups of cells surviving settling selection allow more biomass

to be transferred. Whole populations were cryogenically stored at -80 °C every 7 days.

Measuring settling rate

To explore the dynamics of multicellular adaptation in floc and snowflake yeast

populations, we measured the settling rate of each population every 7 days over the 24-week

experiment. We thawed cryogenically-stored whole populations and subsequently inoculated

100μL into 10 mL of YPGal+Dex and grew them for 24 h. Then, 100 μL of overnight cultures

was inoculated into fresh YPGal+Dex media and grown for an additional 24 h. We measured the

settling rate of populations as described in Pentz *et al*. (2020). Briefly, high-speed high-resolution

videos of yeast populations settling in back-illuminated cuvettes were recorded using a Sony a7RII

and 90 mm macro lens (24 fps, 3840 x 2160 pixels). Then, custom scripts were used to determine

the rate of yeast biomass displacement, or settling rate, based on changes in pixel densities over

settling time [35].

Phenotypic assessment of evolved populations.

Size distributions of evolved snowflake yeast were obtained via flow cytometry on a

CyFlow® Space flow cytometer using the forward scatter (FSC) channel as a proxy for cluster

size. To account for the fact that larger cluster contain more biomass, we calculated a biomass-

weighted mean size for each population. To do so, we developed a Python script that divides the

distribution of cluster size into 100 bins, then determines the proportion of the population's total

biomass that is found in each bin (sum of the FSC values for each cluster within that size range).

This gives us a new distribution of the population's biomass across the range of cluster sizes. In

Figure 2A, we report the means of this biomass-scaled distribution.

Flocculation efficiency was measured using the first frame from the timelapse videos used

to measure settling rate in the week-24 evolved floc populations, as well as the ancestor (Pentz *et*

al. 2020). The population is well-mixed in this frame, so floc aggregates will result in optically

dense regions while the planktonic culture (single cells that are not in flocs) will be less optically

dense. Thus, higher flocculation efficiency will result in a higher variance in the pixel density

between floc aggregates and planktonic cells. A custom Python script was used to calculate the

maximum variance in pixel density on four biological replicates for each population.

Fitness competitions

To determine the fitness of evolved populations, a representative genotype was isolated

from each population at the end of the experiment (24 weeks) by three rounds of single-colony

selection on YPGal+Dex agarose plates (YPGal+Dex with 15 g/L agarose). Our selection regime

is characterized by fluctuating periods of selection for growth and selection for rapid settling, or

large size [35,37]. Thus, it is important to measure fitness in both of these important life history traits.

To do so, we quantified the fitness of evolved isolates relative to their ancestor over one round of

growth and one round of settling selection. Specifically, to initiate competitions, we inoculated 10

mL cultures of YPGal+Dex with isolates from each population as well as a GFP-marked ancestor

and grown for 24 h. Then, we mixed each of the evolved isolates in equal volumes with its marked

ancestor (floc or snowflake), and 100 µL of this mixture was inoculated into 10 mL of YPGal+Dex

to start competitions. Counts of the GFP-tagged ancestor and evolved isolate were obtained via

flow cytometry using a CyFlow® Space flow cytometer where GFP and non-GFP can be

distinguished using the FL1 fluorescence channel. Prior to running on the flow cytometer, floc

competitions were deflocculating using 50 mM EDTA (pH 7). Counts were obtained at time 0 and

after 24 h of growth to determine the fitness of the evolved isolate over one period of growth. To

measure fitness over one round of settling selection, 2 mL of the overnight mixed culture was

aliquoted into a microcentrifuge tube, and 500 µL was used to determine pre-selection counts. The

remaining 1.5 mL was used to perform one round of settling selection (5 min on the bench), after

which the top 1.4 mL was discarded. The remaining bottom 100 µL was used to determine post-

selection counts. In all competitions, relative fitness was calculated using the ratio of Malthusian

growth parameters [47]. Relative fitness was normalized to the fitness of the ancestral strain for each

environment.

Fitness was also measured for the five control populations of floc yeast relative to their

ancestor over three transfers. Competitions were initiated as described above. Every 24 h for three

days, 100 µL was transferred to fresh medium. Counts at the beginning and end of the competition

were obtained using flow cytometry as described above, and the relative fitness was calculated

using the ratio of Malthusian growth parameters [46].

Floc assortment measurements

To calculate assortment, we co-cultured GFP-tagged evolved strains with an RFP-tagged

ancestor. Strains were grown overnight at 30°C in 10 mL of YP + 1.8% galactose + 0.2% dextrose.

The next day, cultures were mixed by vortexing and a 1 mL sample was deflocculated by

centrifuging and resuspending in 100 mM EDTA. To remove EDTA, which inhibits growth,

strains were again centrifuged and resuspended in YP + 1.8% galactose + 0.2% dextrose. From

this, 100 uL of each strain to be co-cultured were added to 10 mL of the same media to grow

overnight at 30°C. For imaging the next day, co-cultures were vortexed for 10 seconds, and 1.5

mL of culture was added to each of two tubes. One tube was for measuring the baseline population

frequency of the two strains, and EDTA was added to a final concentration of 100 mM. The other

tube for each co-culture was for settling selection, and was left on the bench for 5 minutes, then

all but the bottom 100 uL was removed. The remaining 100 uL were deflocculated by adding

EDTA to a final concentration of 200 mM. Both samples were concentrated by centrifugation, and

a small sample was imaged with a 20X microscope objective. Three images were taken for each

sample. Cells in the red and green channels were counted using Otsu thresholding and

watershedding in FIJI. Assortment was calculated using the following equation that controls for

population frequency, where f_{set} and f_{pop} are the frequency of the evolved strain after settling

selection and in the general population, respectively:

585
$$Assortment = \frac{f_{set} - f_{pop}}{1 - f_{pop}}$$

Genomic DNA preparation

To determine the genetic basis of observed fitness differences, we performed whole-

genome sequencing of 24-week evolved isolates and the starting ancestral genotypes. Yeast strains

were streaked out for single colonies from -80°C glycerol stocks. Single colonies were grown

overnight in 10 mL YPGal+Dex and genomic DNA was isolated from 1 mL aliquots using the

VWR® Life Science Yeast Genomic DNA Purification Kit (VWR 89492-616, us.vwr.com).

Whole-genome sequencing

DNA libraries were prepared using the NEBNext® Ultra™ II FS DNA Library Prep Kit

for Illumina (www.neb.com) and were sequenced on an Illumina HiSeq 2500. Paired-end 150bp

reads were used for all samples. Mean coverage across the genome was 200X for evolved isolate

DNA and 50X for ancestor DNA.

Sequencing analysis

DNA sequences were quality trimmed using Trimmomatic [71] and then aligned to the S288C

reference genome R64-2-1 using the Burrows-Wheeler Aligner [72]. Duplicates were marked using

SAMBLASTER [73] then converted to a BAM file, then sorted and indexed. Variants were called

using the Genome Analysis Tool Kit (GATK) HaplotypeCaller [74]. SNPs and INDELs were first

filtered based on read depth and quality using vcffilter (https://github.com/vcflib/vcflib). Variants

were removed with a read depth less than 10 and a quality score less than 20. Then, bcftools isec

(https://github.com/samtools/bcftools) was used to filter out variants shared between the ancestor

and evolved isolates, accounting for variants called due to aligning *S. cerevisiae* strain Y55 used

in our experiments to the S288C reference genome. Finally, bcftools isec was used again to identify

unique variants for each evolved isolate. Variants were manually validated using the Integrated

Genomics Viewer [75]. Final validated variants were pooled and annotated using SnpEff [76].

Next, we performed a bootstrap analysis to compare the classes of variants called

experimentally to a randomly generated sample of SNPs to determine if different mutational

classes are over- or underrepresented in our experimental populations. First, we used Mutation-

Simulator [51] to generate a null distribution of 100,000 random SNPs from across the *S. cerevisiae*

S288C genome and annotated using SnpEff [76]. Then, a custom Python script was used to perform

a bootstrap analysis by first generating a random subsample of SNPs from the null distribution.

The quantity of SNPs subsampled was equal to the pooled number of mutants seen experimentally

for either snowflake or floc yeast (69 or 104 mutations, respectively). Next, we compared the

number of SNPs in four mutational classes (missense, nonsense, synonymous/silent, and upstream

gene variant) generated experimentally or simulated. We performed the bootstrap analysis 1000

times each for snowflake and floc yeast. Histograms for the number of simulated SNPs generated

for different mutational classes in the bootstrap analysis can be seen in Figure 4C (experimental

number shown as vertical blue and orange lines for snowflake and floc yeast, respectively). Finally,

we determined the proportion (P) of runs where the # observed mutations > # simulated mutations

and the P # observed mutations < simulated mutations (see Figure 4C). The code for the Python

script   is   available   at   GitHub   (https://github.com/Ratcliff-Lab/genome-

analysis/tree/main/comparing-mutation-frequencies).

Mathematical modeling

We consider the survival and fixation of mutant lineages by distinguishing between two

phases of the experiment between transfers: the population growth phase and the settling selection

phase.

During the population growth phase, we assume that both the mutant and ancestral lineages

reproduce exponentially until the total population increases by a factor of 100, i.e. they reach the

carrying capacity. So if the initial population is I and there are m mutants, we assume the

population grows via $(I - m)e^{\lambda t} + m\, e^{\lambda(1+s_c)t}$, where the sign of s_c determines whether the

mutation is beneficial ($s_c > 0$) or deleterious ($s_c < 0$). We compute the number of mutants at carrying

capacity and then place them in groups, according to whether we are simulating clonal or

aggregative development. In the case of clonal development, we note that because of the branching

pattern of snowflake yeast there can only be a maximum of one group that is mixed with mutant

and ancestral lineages [38,77]; all other groups with mutant cells are homogeneously mutant. We can

simulate the distribution of mutants in groups of varying sizes by either tuning computational

models of populations of snowflake yeast to fit experimental data or, instead, by directly using

experimental data of group size distributions. We use the latter because it can easily be modified

to consider aggregative development. Thus, for clonal development, we randomly select groups to

place mutants. If there will be more mutant cells generated during growth (in this scenario, we are

generating 100 mutant cells during the growth phase) than the size of the group, then the entire

group is filled with mutant cells and a new group is selected to receive mutants, and so on until all

mutant cells are allocated. In the case of aggregative development, we also use an experimentally

derived distributions of group sizes but we select groups randomly weighted by their size to place

individual mutant cells.

We simulate the settling selection phase of the experiment by randomly selecting groups

weighted linearly by their size. For mutations that alter the survival of groups we assume that each

mutant contributes an additional factor s_g to the size of the group, i.e. the weight of a group is its

size plus s_g*m, or equivalently if there are n ancestral cells the weight is $n+(1+ s_g)*m$. We then

select groups randomly without replacement according to this weight until we have selected

1/100th of the population. If the number of cells exceeds 1/100th of the population we simply

rescale the selected number of cells to fit. Following selection, we compute the number of

surviving mutants and, if they exceed 0, we return to the growth phase. We iterate this process five

times.

Acknowledgements

This work was supported by NSF grant DEB-1845363 and a Packard Fellowship for Science and

Engineering to W.C.R. K.M. was supported by NIH T32 grant T32GM142616. We thank Jordi

van Gestel and the Ratcliff lab for constructive comments.

Competing Interests

The authors have no competing interests to declare.

664   **References**

1   Umen, J. & Herron, M. D. Green algal models for multicellularity. *Annual Review of*
Genetics **55**, 603-632 (2021).
2   Grosberg, R. K. & Strathmann, R. R. The evolution of multicellularity: a minor major
transition? *Annu. Rev. Ecol. Evol. Syst.* **38**, 621-654 (2007).
3   O'Malley, M. A. in *The Evolution of Multicellularity*   9-24 (CRC Press, 2022).
4   Brunet, T. & King, N. The origin of animal multicellularity and cell differentiation.
Developmental cell **43**, 124-140 (2017).
5   Pfeiffer, T. & Bonhoeffer, S. An evolutionary scenario for the transition to
undifferentiated multicellularity. *Proceedings of the National Academy of Sciences* **100**,
1095-1098, doi:doi:10.1073/pnas.0335420100 (2003).
6   Knoll, A. H. The multiple origins of complex multicellularity. *Annual Review of Earth*
and Planetary Sciences **39**, 217-239 (2011).
7   Herron, M. D., Conlin, P. L. & Ratcliff, W. C. *The Evolution of Multicellularity*. (CRC
Press, 2022).
8   Rose, C. & Hammerschmidt, K. What do we mean by multicellularity? The Evolutionary
Transitions Framework provides answers. *Frontiers in Ecology and Evolution* **9** (2021).
9   Michod, R. E. Evolution of individuality during the transition from unicellular to
multicellular life. *Proceedings of the National Academy of Sciences* **104**, 8613-8618
(2007).
10   Godfrey-Smith, P. *Darwinian populations and natural selection*. (Oxford University
Press, 2009).
11   Buss, L. W. *The evolution of individuality*. (Princeton University Press, 2014).
12   West, S. A., Fisher, R. M., Gardner, A. & Kiers, E. T. Major evolutionary transitions in
individuality. *Proceedings of the National Academy of Sciences* **112**, 10112-10119
(2015).
13   Smith, J. M. & Szathmary, E. *The major transitions in evolution*. (OUP Oxford, 1997).
14   Bonner, J. T. The origins of multicellularity. *Integrative Biology Issues News and*
Reviews **1**, 27-36 (1998).
15   Bonner, J. T. *First Signals: The Evolution of Multicellular Development*. (Princeton
University Press, 2000).
16   Crespi, B. J. The evolution of social behavior in microorganisms. *Trends in ecology &*
evolution **16**, 178-183 (2001).
17   Tarnita, C. E., Taubes, C. H. & Nowak, M. A. Evolutionary construction by staying
together and coming together. *Journal of theoretical biology* **320**, 10-22 (2013).
18   Velicer, G. J. & Vos, M. Sociobiology of the myxobacteria. *Annual review of*
microbiology **63**, 599-623 (2009).
19   Du, Q., Kawabe, Y., Schilde, C., Chen, Z.-h. & Schaap, P. The evolution of aggregative
multicellularity and cell–cell communication in the Dictyostelia. *Journal of molecular*
biology **427**, 3722-3733 (2015).
20   Fisher, R., Shik, J. & Boomsma, J. The evolution of multicellular complexity: the role of
relatedness and environmental constraints. *Proceedings of the Royal Society B* **287**,
20192963 (2020).
21   Godfrey-Smith, P., Bouchard, F. & Huneman, P. Darwinian individuals. *From groups to*
individuals: evolution and emerging individuality **16**, 17 (2013).
22   Szathmáry, E. & Smith, J. M. *Major Transitions in Evolution*. (Oxford University Press
Oxford, 1997).
23   Clarke, E. Origins of evolutionary transitions. *Journal of biosciences* **39**, 303-317 (2014).

24 Zamani-Dahaj, S. A. *et al.* Spontaneous emergence of multicellular heritability. *bioRxiv*
(2021).
25 Michod, R. E. On the transfer of fitness from the cell to the multicellular organism.
Biology and Philosophy **20**, 967-987 (2005).
26 Márquez-Zacarías, P., Conlin, P. L., Tong, K., Pentz, J. T. & Ratcliff, W. C. Why have
aggregative multicellular organisms stayed simple? *Current Genetics* **67**, 871-876 (2021).
27 Bonner, J. T. Perspective: the size-complexity rule. *Evolution* **58**, 1883-1890 (2004).
28 Fisher, R. M., Cornwallis, C. K. & West, S. A. Group formation, relatedness, and the
evolution of multicellularity. *Current Biology* **23**, 1120-1125 (2013).
29 Hanschen, E. R. *et al.* The Gonium pectorale genome demonstrates co-option of cell
cycle regulation during the evolution of multicellularity. *Nature Communications* **7**, 1-10
(2016).
30 King, N. The unicellular ancestry of animal development. *Developmental cell* **7**, 313-325
(2004).
31 King, N. *et al.* The genome of the choanoflagellate Monosiga brevicollis and the origin of
metazoans. *Nature* **451**, 783-788 (2008).
32 Suga, H. *et al.* The Capsaspora genome reveals a complex unicellular prehistory of
animals. *Nature communications* **4**, 1-9 (2013).
33 Herron, M. D., Hackett, J. D., Aylward, F. O. & Michod, R. E. Triassic origin and early
radiation of multicellular volvocine algae. *Proceedings of the National Academy of*
Sciences **106**, 3254-3258 (2009).
34 Ratcliff, W. C., Herron, M., Conlin, P. L. & Libby, E. Nascent life cycles and the
emergence of higher-level individuality. *Phil. Trans. R. Soc. B* **372**, 20160420 (2017).
35 Pentz, J. T., Márquez-Zacarías, P., Yunker, P. J., Libby, E. & Ratcliff, W. C. Ecological
advantages and evolutionary limitations of aggregative multicellular development.
Current Biology **30**, 4155-4164 (2020).
36 Smukalla, S. *et al.* FLO1 is a variable green beard gene that drives biofilm-like
cooperation in budding yeast. *Cell* **135**, 726-737 (2008).
37 Ratcliff, W. C., Denison, R. F., Borrello, M. & Travisano, M. Experimental evolution of
multicellularity. *Proceedings of the National Academy of Sciences* **109**, 1595-1600
(2012).
38 Ratcliff, W. C., Fankhauser, J. D., Rogers, D. W., Greig, D. & Travisano, M. Origins of
multicellular evolvability in snowflake yeast. *Nature communications* **6**, 6102 (2015).
39 Dudin, O., Wielgoss, S., New, A. M. & Ruiz-Trillo, I. Regulation of sedimentation rate
shapes the evolution of multicellularity in a unicellular relative of animals. *bioRxiv*
(2021).
40 Tong, K., Bozdag, G. O. & Ratcliff, W. C. Selective drivers of simple multicellularity.
Current Opinion in Microbiology **67**, 102141 (2022).
41 Hammerschmidt, K. & Rose, C. J. What do we mean by multicellularity? The
Evolutionary Transitions Framework provides answers. *Frontiers in Ecology and*
Evolution, 784.
42 Ratcliff, W. C., Pentz, J. T. & Travisano, M. Tempo and mode of multicellular adaptation
in experimentally evolved Saccharomyces cerevisiae. *Evolution* **67**, 1573-1581 (2013).
43 Bozdag, G. O. *et al.* De novo evolution of macroscopic multicellularity. *bioRxiv* (2021).
44 Bozdag, G. O., Libby, E., Pineau, R., Reinhard, C. T. & Ratcliff, W. C. Oxygen
suppression of macroscopic multicellularity. *Nature communications* **12**, 1-10 (2021).

45   Shelton, D. E. & Michod, R. E. Group and individual selection during evolutionary
760   transitions in individuality: meanings and partitions. *Philosophical Transactions of the*
761   *Royal Society B* **375**, 20190364 (2020).
762   46   Conlin, P. L. & Ratcliff, W. C. Trade-offs drive the evolution of increased complexity in
763   nascent multicellular digital organisms. *Multicellularity: origins and evolution* **131**
764   (2016).
765   47   Lenski, R. E., Rose, M. R., Simpson, S. C. & Tadler, S. C. Long-term experimental
766   evolution in Escherichia coli. I. Adaptation and divergence during 2,000 generations. *The*
767   *American Naturalist* **138**, 1315-1341 (1991).
768   48   Yanni, D., Márquez-Zacarías, P., Yunker, P. J. & Ratcliff, W. C. Drivers of spatial
769   structure in social microbial communities. *Current biology* **29**, R545-R550 (2019).
770   49   Fletcher, J. A. & Doebeli, M. A simple and general explanation for the evolution of
771   altruism. *Proceedings of the Royal Society B: Biological Sciences* **276**, 13-19 (2009).
772   50   Pepper, J. W. & Smuts, B. B. A mechanism for the evolution of altruism among nonkin:
773   positive assortment through environmental feedback. *The American Naturalist* **160**, 205-
774   213 (2002).
775   51   Kühl, M., Stich, B. & Ries, D. Mutation-Simulator: fine-grained simulation of random
776   mutations in any genome. *Bioinformatics* **37**, 568-569 (2021).
777   52   Jacobeen, S. *et al.* Cellular packing, mechanical stress and the evolution of
778   multicellularity. *Nature physics* **14**, 286-290 (2018).
779   53   Lang, G. I. *et al.* Pervasive genetic hitchhiking and clonal interference in forty evolving
780   yeast populations. *Nature* **500**, 571-574 (2013).
781   54   Fichtner, L., Schulze, F. & Braus, G. H. Differential Flo8p-dependent regulation of FLO1
782   and FLO11 for cell–cell and cell–substrate adherence of S. cerevisiae S288c. *Molecular*
783   *microbiology* **66**, 1276-1289 (2007).
784   55   Van Dyk, D., Pretorius, I. S. & Bauer, F. F. Mss11p is a central element of the regulatory
785   network that controls FLO11 expression and invasive growth in Saccharomyces
786   cerevisiae. *Genetics* **169**, 91-106 (2005).
787   56   Bayly, J. C., Douglas, L. M., Pretorius, I. S., Bauer, F. F. & Dranginis, A. M.
788   Characteristics of Flo11-dependent flocculation in Saccharomyces cerevisiae. *FEMS*
789   *Yeast Research* **5**, 1151-1156, doi:10.1016/j.femsyr.2005.05.004 (2005).
790   57   Van Gestel, J. & Tarnita, C. E. On the origin of biological construction, with a focus on
791   multicellularity. *Proceedings of the National Academy of Sciences* **114**, 11018-11026
792   (2017).
793   58   Libby, E. & Rainey, P. B. A conceptual framework for the evolutionary origins of
794   multicellularity. *Physical biology* **10**, 035001 (2013).
795   59   Rainey, P. B. & De Monte, S. Resolving conflicts during the evolutionary transition to
796   multicellular life. *Annual Review of Ecology, Evolution, and Systematics* **45**, 599-620
797   (2014).
798   60   Bailey, S. F., Alonso Morales, L. A. & Kassen, R. Effects of synonymous mutations
799   beyond codon bias: the evidence for adaptive synonymous substitutions from microbial
800   evolution experiments. *Genome biology and evolution* **13**, evab141 (2021).
801   61   Shen, X., Song, S., Li, C. & Zhang, J. Synonymous mutations in representative yeast
802   genes are mostly strongly non-neutral. *Nature*, 1-7 (2022).
803   62   Good, B. H., McDonald, M. J., Barrick, J. E., Lenski, R. E. & Desai, M. M. The
804   dynamics of molecular evolution over 60,000 generations. *Nature* **551**, 45-50 (2017).

63   Johnson, M. S. *et al.* Phenotypic and molecular evolution across 10,000 generations in
laboratory budding yeast populations. *Elife* **10**, e63910 (2021).
64   Giaever, G. *et al.* Functional profiling of the Saccharomyces cerevisiae genome. *nature*
418, 387-391 (2002).
65   Whitlock, M. C. & Barton, N. H. The effective size of a subdivided population. *Genetics*
146, 427-441 (1997).
66   Bell, G. & Mooers, A. O. Size and complexity among multicellular organisms. *Biological*
Journal of the Linnean Society **60**, 345-363 (1997).
67   Lynch, M. & Conery, J. S. The origins of genome complexity. *science* **302**, 1401-1404
(2003).
68   Grosberg, R. K. & Strathmann, R. R. One cell, two cell, red cell, blue cell: the persistence
of a unicellular stage in multicellular life histories. *Trends in ecology & evolution* **13**,
112-116 (1998).
69   Queller, D. C. Relatedness and the fraternal major transitions. *Philosophical Transactions*
of the Royal Society of London. Series B: Biological Sciences **355**, 1647-1655 (2000).
70   Black, A. J., Bourrat, P. & Rainey, P. B. Ecological scaffolding and the evolution of
individuality. *Nature Ecology & Evolution* **4**, 426-436 (2020).
71   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120 (2014).
72   Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows–Wheeler
transform. *Bioinformatics* **26**, 589-595 (2010).
73   Faust, G. G. & Hall, I. M. SAMBLASTER: fast duplicate marking and structural variant
read extraction. *Bioinformatics* **30**, 2503-2505 (2014).
74   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome research* **20**, 1297-1303
(2010).
75   Robinson, J. T. *et al.* Integrative genomics viewer. *Nature biotechnology* **29**, 24 (2011).
76   Cingolani, P. *et al.* A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster
strain w1118; iso-2; iso-3. *Fly* **6**, 80-92 (2012).
77   Libby, E., Ratcliff, W., Travisano, M. & Kerr, B. Geometry shapes evolution of early
multicellularity. *PLoS computational biology* **10**, e1003803 (2014).

Supplementary Figures

Supplementary Figure 1 - Clonal and aggregative yeast model system. A) Snowflake yeast
were constructed by deleting a gene responsible for mother-daughter cell separation following
division, *ACE2*, resulting in clonal clusters, shown by fluorescently tagged clusters that are remain
all green or all red during co-culture. Images adapted from Pentz *et. al*, 2020. B) Floc yeast were
constructed by inserting a gene that confers strong flocculation, *FLO1*, resulting in chimeric
clusters where fluorescently tagged cells will randomly aggregate, creating chimeric green and red
clusters. C,D) Simple schematics representing snowflake and flocculating yeast life cycles.

Supplementary Figure 2 - Size distributions for evolved snowflake isolates. Size distributions
of evolved isolates of individual snowflake populations. Overlay shown in Figure 2B.

Supplementary Figure 3 - Number of doublings per day in snowflake and floc yeast over the course of our 24-week selection experiment. To assay the number of generations floc and snowflake yeast experienced over the duration of our selection experiment, we measured the number of doublings per day, quantified as the log2(density after 24h/density post-selection) every three weeks. On average, both snowflake and floc yeast experienced 4-5 doublings per day. Thus, over the course of 24-weeks, both snowflake and floc yeast underwent ~700 generations. Error bars represent standard deviation of three biological replicates.

Supplementary Figure 4. Growth adaptation control for floc. To determine if adaptation to growth on galactose increases flocculation as a pleiotropic side effect, we evolved five populations of floc for eight weeks with daily growth rate selection on YPG. These evolved a ~20% increase in fitness during growth (one-way ANOVA; $F_{5,12}$=59.45, p<0.0001, pairwise differences assessed with Tukey's post-doc HSD with α=0.05), but a ~40% reduction in flocculation efficiency (one-way ANOVA; $F_{5,18}$=27.26, p<0.0001, pairwise differences assessed with Tukey's post-doc HSD with α=0.05), demonstrating that adaptation to faster growth alone does not drive increased flocculation. Shown is the mean and standard deviation of 3 and 4 biological replicates for fitness during growth and flocculation efficiency, respectively, normalized to the mean of the ancestor.