Molecular insight into the enzymatic macrocyclization of multiply backbone N-methylated peptides

Emmanuel Matabaro3#, Haigang Song1,2#, Lukas Sonderegger3, Fabio Gherlone3, Andrew Giltrap1, Sam Liver1, Alvar Gossert4, Markus Künzler3*, James H Naismith1,2*

[1] Structural Biology, The Rosalind Franklin Institute, Harwell Science & Innovation Campus, Didcot, UK. OX11 0QS
[2] Division of Structural Biology, University of Oxford, The Wellcome Centre for Human Genetics, Headington, Oxford, UK. OX3 7BN
[3] Institute of Microbiology, Department of Biology, ETH Zürich, Vladimir-Prelog-Weg 4, CH-8093 Zürich, Switzerland
[4] Biomolecular NMR Spectroscopy Platform, Department of Biology, ETH Zürich, Otto-Stern-Weg 5, CH-8093 Zürich, Switzerland

#equal contribution
*correspondence to (MK for protein and peptide production and function, JHN for protein structure and molecular mechanism)

Abstract

The enzyme OphP is essential for the biosynthesis of the macrocyclic peptide omphalotin A, a dodecamer with 9 backbone N-methylations produced by the wood-degrading fungus *Omphalotus olearius*. Heterologous expression of OphP and the peptide-precursor protein OphMA in yeast, yields omphalotin A. Thus, Oph P was hypothesized to have a dual function; catalyzing both endoproteolytic release of a peptide intermediate from OphMA, and macrocyclization of the multiply α-N-methylated core peptide with concomitant release of a C-terminal follower peptide. In our *in vitro* activity assays, OphP showed robust endoproteolytic and macrocyclase activity on α-N-methylated peptides but was unable to cleave OphMA. The enzyme had a strong preference for hydrophobic, highly α-N-methylated peptides and an α-N-methylated glycine residue at the P1 site. OphP adopts a canonical prolyl oligopeptidase (POP) fold with a predominantly hydrophobic substrate binding cleft, and a small and hydrophobic P1 binding pocket. We demonstrate that OphP is a POP-type macrocyclase with a specificity and a substrate route to the active site different from other members of the family. These results could be exploited for the biotechnological production of macrocyclic peptides with multiple backbone N-methylations, which are interesting due to their favorable pharmacological properties.

Introduction

Despite violating Lipinski's 'rule of five' (Lipinski et al., 2001), some peptide macrocycles show potent biological activity. This is because the macrocyclization restricts the bond rotations of the peptide backbone and, thus, the favourable conformation of the peptide has a much lower entropic penalty for target engagement (Marsault & Peterson, 2011; Marti-Centelles et al., 2015). In addition, peptide macrocycles are metabolically more stable as they are intrinsically resistant to proteases, and generally show an improved membrane permeability and oral availability when compared to linear peptides (Giordanetto & Kihlberg, 2014). Macrocycles are in clinical use against G-protein coupled receptors and protein-protein interfaces, latter being traditionally challenging targets for small (Lipinski compliant) molecules (Driggers et al., 2008; Surade & Blundell, 2012). Peptide macrocycles are particularly attractive for drug development due to their modular structure and potential chemical diversity arising from the flexible choice of building blocks (natural and unnatural amino acids, α-N-methylated amino acids, D-amino acids, β-amino acids, heterocyclised amino acids and thioamides). Macrocyclization of peptides can be accomplished either by forming a peptide bond between the N- and the C-terminus or by linking two or more functional groups forming amide, lactone, ether, thioether, disulfide or even C-C bonds (Arnison et al., 2013; Koehnke et al., 2017; Montalbán-López et al., 2021). There are powerful routes to synthetic peptide macrocycles with simple amino acids, but syntheses with more complex building blocks, that can significantly improve the potency and pharmacokinetics, are challenging. As a consequence, natural products remain the main sources of marketed peptide macrocycle drugs (Lee & van der Donk, 2022).

Naturally occurring peptide macrocycles are produced by two broad biosynthetic routes, resulting in non-ribosomal peptides (NRPs) and ribosomally synthesized and post-translationally modified peptides (RiPPs). In the case of NRPs, multidomain megaenzymes, non-ribosomal peptide synthetases (NRPS), catalyse the synthesis of the linear peptides from proteinogenic and non-proteinogenic amino acids using linearly arranged, amino acid specific modules in an assembly line fashion. The macrocylization of NRPs is mediated by a final thioesterase domain of the NRPS in bacteria or a condensation-like domain in fungi (Gao et al., 2012; Kohli et al., 2001; Zhang et al., 2016). Several enzyme classes catalyse macrocyclization of RiPPs, including PatG in patellamide biosynthesis (Donia et al., 2008; McIntosh et al., 2010), GmPOPB in fungal amatoxin (Czekster et al., 2017; Luo et al., 2014), and PCY1 in plant orbitide (Barber et al., 2013; Chekan et al., 2017; Ludewig et al., 2018). Mechanistically, these enzymes first cleave off a C-terminal sequence (the follower peptide) from the ribosomally synthesized precursor peptide, forming an acyl enzyme intermediate which is then resolved by the intramolecunucleophilic attack of the free amino terminus of the peptide substrate to form the macrocycle. The macrocyclase domain of PatG belongs to the S8 family of subtilisin-type serine peptidases (MEROPS database of known proteases and their inhibitors, https://www.ebi.ac.uk/merops/) and recognizes a four-residue follower peptide and a preceding heterocyclic ring residue (Koehnke et al., 2012; Koehnke et al., 2014; McIntosh et al., 2010). PatG tolerates a wide range of peptide substrates but is slow (Oueis et al., 2016; Emilia Oueis et al., 2017; E Oueis et al., 2017). PCY1 and GmPOPB belong to post-proline cleaving prolyl oligopeptidases (POP) of the S9A family of serine peptidases. GmPOPB is a dual function macrocyclase as it sequentially catalyses the removal of a leader and follower sequence from the same peptide substrate (precursor peptide) (Czekster et al., 2017; Czekster & Naismith, 2017; Luo et al., 2014). PCY1 relies on an additional protease to first remove the leader sequence to create the substrate for macrocyclization (Barber et al., 2013). Structural and biophysical analysis shows that both PCY1 and GmPOPB recognise the residues in the follower region for substrate binding and catalysis (Chekan et al., 2017;

Ludewig et al., 2018). Butelase 1, a fast and promiscuous cyclase from the asparaginyl endopeptidase (AEP) family, has been reported to recognize the C-terminal Asn/Asp-His-Val motif of its peptide substrate, to cleave off C-terminus and ligate the exposed Asn/Asp to the N-terminus of substrate (Nguyen et al., 2015; Nguyen et al., 2014).

A considerable fraction of naturally occurring macrocyclic peptides is backbone N-methylated at one or multiple amino acids. The function of backbone N-methylation of peptides is to enhance their conformational rigidity, membrane permeability, protease resistance and – most relevant in their use as drugs – oral availability (Chatterjee et al., 2008; Ovadia et al., 2011). GmPOPB and PCY1 have been shown to catalyse the macrocyclization of singly backbone N-methylated linear peptides (Ludewig et al., 2018; Sgambelluri et al., 2018) but no enzyme has been characterised, which can efficiently macrocyclize multiply backbone N-methylated peptides. Synthesis and macrocyclization of peptides with multiple backbone N-methylations by entirely chemical means (Marsault & Peterson, 2011; Marti-Centelles et al., 2015; Yu & Sun, 2013) is challenging. Yet, they have enormous potential, as demonstrated by the most well-known example, cyclosporin A, an orally available natural product of the insect pathogenic fungus *Tolypocladium inflatum* (Olarte et al., 2019) that is used as an immunosuppressant during organ transplantation (Cohen et al., 1984). The macrocycle consists of 11 amino acids, 7 of which are backbone N-methylated and 3 are non-proteinogenic (Dreyfuss et al., 1976). The oral availability is a consequence of its chameleon-like property, where the molecule reversibly switches between a non-polar arrangement for membrane transit and a polar arrangement for water solubility (Alex et al., 2011; El Tayar et al., 1993). The conformational flipping depends on the methylated backbone. Cyclosporin A is synthesised by an NRPS pathway with methylations preinstalled on the building blocks prior to peptide bond formation and ring closure by condensation (Dittmann et al., 1994; Lawen & Zocher, 1990).

The peptide natural product at the center of this study is omphalotin A, a 12-residue macrocyclic RiPP with 9 backbone N-methylations (**Figure 1a**). The molecule is produced by the wood-degrading fungus *Omphalotus olearius* and has anti-nematode activity (Mayer et al., 1997). Two posttranslationally modifying enzymes, a peptide α-N-methyltransferase, OphMA, and a serine peptidase of the S9A family, OphP, contribute to its biosynthesis (Ramm et al., 2017; van der Velden et al., 2017). OphMA catalyses the iterative backbone N-methylation of the amide bonds within the core peptide located at its own C-terminus. The enzyme uses SAM as cofactor and has promiscuity towards proteinogenic hydrophobic, aromatic and small hydrophilic amino acid residues, and even accepts non-proteinogenic residues (Song et al., 2021; Song et al., 2020; Song & Naismith, 2020). Coexpression of OphMA and OphP in the yeast *Pichia pastoris* yielded omphalotin A and swapping of the core peptide in OphMA with sequences for lentinulin A and dendrothelin A led to the production of the respective backbone N-methylated macrocycles (**Figure 1b**), consistent with a promiscuous and functional biosynthetic pathway of the two enzymes (Matabaro, Kaspar, et al., 2021; Ramm et al., 2017) (**Figure 1a**). Based on these data and the sequence similarity between the two enzymes, it was hypothesized that OphP, like GmPOPB (Czekster et al., 2017), possesses two catalytic activities: (1) proteolytic cleavage of the OphMA protein at the N-terminus of the backbone N-methylated core peptide, and (2) proteolytic cleavage of the released precursor peptide at the C-terminus of the core peptide and formation of an acyl-enzyme intermediate between the core peptide and OphP. This intermediate is then resolved by the nucleophilic attack of a water molecule or the N-terminus of the core peptide, leading to the release of the linear or macrocyclized (omphalotin A) core peptide, respectively (**Figure 1a**) (Matabaro, Kaspar, et al., 2021; Matabaro, Song, et al., 2021; Ramm et al., 2017).

In this study, we purified OphP from *P. pastoris* and assessed its *in vitro* activity towards a
variety of substrates. We found no evidence for OphP being able to process intact OphMA
but detected a robust proteolytic and macrocyclase activity of the enzyme towards multiply
backbone N-methylated peptides derived from OphMA. In contrast to other macrocyclase
systems, the enzyme appears to be insensitive to the sequence of the follower peptide but
have a preference for hydrophobic, multiply backbone N-methylated peptides and recognize a
backbone N-methylated glycine residue within the OphMA core peptide as P1 site for
proteolytic cleavage. These findings are supported by the structural analysis of enzyme-
substrate complexes. This knowledge may enable the exploitation of the enzyme for the
biotechnological production of new-to-nature multiply backbone N-methylated peptide
macrocycles for applications in agriculture or medicine.
150

Results

OphP proteolytically cleaves and macrocyclizes OphMA-derived, multiply backbone N-methylated peptides in vitro

OphP was purified from *P. pastoris* as an N-terminal His$_8$SUMO* fusion protein (**Figure 1-figure supplement 1**) and confirmed to be catalytically active using a standard chromogenic substrate for POPs, benzyloxycarbonyl-Gly-Pro-p-nitroanilide (Z-Gly-Pro-4-pNA) (Nagatsu et al., 1976) (**Figure 1-figure supplement 2**). OphP showed optimal activity at 30°C and pH 6.0. The reaction was dependent on enzyme and substrate concentration. OphP activity was reduced by Z-Pro-prolinal (ZPP), a common covalent inhibitor for POPs, but the enzyme was not inhibited as strongly as other members of the family (**Figure 1-figure supplement 2**) (Yoshimoto et al., 1985). Mutation of the putative active site Ser residue at position 580 to Ala (S580A) abolished the detectable enzymatic activity of the fusion protein *in vivo*. Co-incubation of catalytically active OphP with intact OphMA protein showed no detectable peptide product (LC-MS) nor did we observe any mass loss of the protein (ESI-MALDI-TOF) (**Figure 1-figure supplement 3**). Intact OphMA was expressed as an N-terminal His$_8$-fusion protein in *E. coli* for 72 hours and purified as previously described (Song et al., 2020; Song et al., 2018; van der Velden et al., 2017). The purified protein is a mixture of different species regarding the degree of backbone N-methylation with the predominant protein species containing 10 methylations (includes one residue in the follower peptide) (**Figure 1a**). The experiment was repeated by purifying OphMA with a lower degree of methylation (predominantly zero, one or two methylations) and again we detected no turnover by OphP. Addition of SAM *in vitro*, which led to conversion of OphMA to the completely methylated state over time, to the reaction, did not change this negative result. Based on these data, we conclude that OphP is not able to process the intact OphMA protein.

To characterize the presumed peptidase and/or macrocyclization activity of OphP, we tested a variety of backbone N-methylated, OphMA-derived C-terminal peptides, consisting of clasp domain, core peptide and follower peptide sequences (**Figure 1a**), for *in vitro* processing by OphP. Since, in our hands, the chemical synthesis of methylated or unmethylated OphMA-derived C-terminal peptides failed (an issue we attributed to their highly hydrophobic character), the peptides were generated by cleaving purified N-terminally His$_8$-tagged recombinant wildtype OphMA using trypsin (Oph-30mer), and variants thereof containing cleavage sites for the tobacco etch virus (TEV) protease in the unmethylated clasp domain at various distances from the core peptide, using TEV protease. Some TEV-cleavable OphMA variants also contained deletions and mutations in the C-terminal core and follower peptide sequences. The OphMA variants were expressed for a sufficient period in *E. coli* to allow backbone N-methylation to proceed, purified using metal affinity chromatography and processed by trypsin or TEV protease. The proteolytically released peptides were purified by preparative HPLC (**Figure 2a**). Peptides produced in this way (Oph-30mer, Oph-24mer, Oph-21mer, Oph-18mer, Oph-15mer, Led-21mer and Dbi-21mer), included between three and 12 residues of the unmethylated clasp domain preceding the core peptide. Whereas all other peptides contained the complete 12-residue core and 6-residue follower peptide, Oph-18mer[Tr] and Oph-15mer contained a truncated core peptide lacking the three C-terminal residues and a truncation of the entire follower peptide, respectively. Led-21mer and Dbi-21mer, contained the 12-residue cores and the follower peptides (SVVSSA) of the lentinulin A and dendrothelin A precursor proteins, LedMA and DbiMA1 (Matabaro, Kaspar, et al., 2021; Matabaro, Song, et al., 2021), respectively (**Figure 2a, Figure 2-figure supplement 1**). MS analysis indicated that the Oph-30mer, Oph-24mer, Oph-21mer, Dbi-21mer, and Led-21mer peptides contained up to ten methylations, both the Oph-18mer[Tr] had up to 8 methylations and the Oph-15mer carried 7 methylations (**Figure 2-figure supplements 1, 2**).

201   All our attempts to produce and purify non-methylated peptide substrates, using either the
202   catalytically inactive R72A mutant of OphMA or maltose-binding protein fusions of C-
203   terminal parts of OphMA, failed due to the insolubility of the peptides.
204
With the exception of Oph-30mer, where no product could be detected, incubation of all
these peptides with OphP led to the formation of peptide macrocycles where the
unmethylated residues of the clasp domain that precede the methylated core peptide in
OphMA, were incorporated into the product (**Figure 2a, 2b**). The Oph-24mer peptide thereby
yielded a mixture of the 15-residue macrocycle cyclo(SQNGFPW^{me}VI^{me}V^{me}V^{me}G^{me}V^{me}I^{me}G)
([M+H]$^+$=1651.958)   and   the   18-residue   macrocycle
cyclo(SQNGFPW^{me}VI^{me}V^{me}V^{me}G^{me}V^{me}I^{me}GV^{me}I^{me}G)   ([M+H]$^+$=1949.163)   (^{me} denotes
backbone N-methylation (**Figure 2b, Figure 2-figure supplement 3**). Similarly, the Oph-
21mer peptide yielded the 12-residue macrocycle cyclo(GFPW^{me}VI^{me}V^{me}V^{me}G^{me}V^{me}I^{me}G)
([M+H]$^+$=1322.824)   and   the   15-residue   macrocycle
cyclo(GFPW^{me}VI^{me}V^{me}V^{me}G^{me}V^{me}I^{me}GV^{me}I^{me}G) ([M+H]$^+$=1620.0293) (**Figure 2c, Figure
2-figure supplement 3**). The Oph-18mer^{Tr} and -15mer peptides both produced the 12-residue
macrocycle cyclo(GFPW^{me}VI^{me}V^{me}V^{me}G^{me}V^{me}I^{me}G) (**Figure 2d**). These results suggest that
OphP is able to *in vitro* macrocyclize OphMA-derived, multiply backbone N-methylated
peptides at two different sites within the methylated core but unable to remove the
unmethylated clasp domain residues resulting in peptide macrocycles including these residues.
221
In order to test whether OphP is also able to produce macrocycles consisting only of the
methylated core peptides, we exploited the fact that the clasp domain ends with a proline
residue in OphMA, LedMA and DbiMA1 and treated derived peptides with the proline-
specific endopeptidases recombinant GmPOPB (Czekster et al., 2017) or ProAlanase
(Promega, Madison WI, USA). In detail, these were Oph-21mer (leading to 18mer^{NL}; ^{NL}
denotes 'no leader'), Oph-18mer^{Tr} (leading to Oph-15mer^{Tr}), Oph-15mer (leading to Oph-
12mer), Led-21mer (leading to Led-18mer) and Dbi-21mer (leading to Dbi-18mer).
Treatment of the purified peptides with these enzymes removed the unmethylated residues
preceding the methylated core peptides (**Figure 2a, Figure 2-figure supplement 4**).
Subsequent incubation of Oph-18mer^{NL}, Led-18mer and Dbi-18mer with OphP led to the
formation of omphalotin A, lentinulin A and dendrothelin A, respectively (**Figure 2e**).
Noteworthy, in all reactions where peptide macrocycles were detected, also linear peptide
products were found (e.g. **Figure 2-figure supplement 2, 4 and 6**). This is typical for
peptide macrocyclases and a consequence of the inherent endopeptidase activity of these
enzymes (Czekster et al., 2017). Incubation of Oph-15mer^{Tr} and Oph-12mer with OphP only
produced linear peptides (W^{me}VI^{me}V^{me}V^{me}G^{me}V^{me}I^{me}G), suggesting OphP cannot form a 9-
residue macrocycle. These results are in accordance with the detection of linear 9-residue
core peptides in a recent *in vivo* study (Matabaro, Kaspar, et al., 2021).
240
To test whether OphP could process known peptide substrates for POPs, GmPOPB peptide
substrates, namely the 25mer amatoxin precursor (AMA1) and the 17mer phalloidin
precursor peptide (PHA1) were chemically synthesized. Consistent with the previous report
(Czekster et al., 2017; Luo et al., 2014), GmPOPB converted both substrates and produced
cyclic octapeptide from AMA1 and linear peptides in the case of PHA1 (**Figure 2-figure
supplement 5**). In contrast, neither OphP nor its homolog LedP was able to process these
non-native peptides. These results are in agreement with the specificity of OphP (and LedP)
for α-N-methylated peptide substrates.
249

Taken together, our results show that OphP is able to proteolytically cleave and macrocyclize
OphMA-derived peptides of 12 to 24 residues containing a multiply backbone N-methylated
core of at least nine residues. The OphP-mediated macrocyclization reaction involves the
proteolytic clipping of three to six unmethylated or partially methylated residues following
the methylated core peptide and produces peptide macrocycles of 12 to 18 residues that can
include stretches of up to six unmethylated residues preceding the methylated core peptide.
On the other hand, OphP appears to be unable to remove unmethylated residues preceding the
methylated core peptide or process intact OphMA.
258

OphP peptide cleavage and macrocyclization activity has a strong preference for an α-N-
methylated glycine residue at the P1 site
A closer analysis of the data revealed that peptide processing by OphP depended on the
presence of an α-N-methylated glycine (^{me}Gly) residue at the P1 site and an unmethylated
residue at the P1' site. This preference is particularly evident on the processing of Oph-
18mer^{NL} and Oph-15mer by OphP (**Figure 3a**). MS analysis of purified Oph-18mer^{NL}
showed predominantly 10 methylations with 3- to 11-fold methylated species being present in
lower amounts (11[th] methylation of the Met414 residue in the follower peptide) whereby the
methylation pattern of the various species can easily be inferred since methylation of the core
peptide by OphMA occurs sequentially from the N- to the C-terminus (van der Velden et al.,
2017). Incubation of Oph-18mer^{NL} with OphP showed that OphP consumed all species but
the 3-, 5- and 6-fold methylated ones. The peptide species consumed by OphP, all have a
^{me}Gly followed by an non-methylated residue while the three non-processed ones do not.
These results were confirmed by the *in vitro* processing of a 1:1 mixture of 6- and 7-fold
Oph-15mer. It should be noted that 7-fold methylation is the maximal degree of methylation
that is achievable for a OphMA lacking the 6-residue follower peptide (Song et al., 2020).
While the 7-fold methylated peptide species was readily consumed by OphP, the 6-fold
methylated one was hardly diminished (**Figure 3a and Figure 3-figure supplement 1**).
These results were confirmed by analyzing the processing of another batch of Oph-15mer
where purified OphMA was incubated with SAM *in vitro* before cleavage by TEV protease
and HPLC-purification of the peptide (**Figure 3-figure supplement 2**). This procedure is
known to increase the degree of core peptide methylation (Song et al., 2018) and, accordingly,
yielded Oph-15mer with a 1:9 ratio of 6-fold:7-fold methylated peptide. MS analysis of
consumption of this batch of Oph-15mer by OphP and the formation of the respective
macrocyclic products, cyclo(GFPW^{me}VI^{me}V^{me}V^{me}G^{me}V^{me}I^{me}G) and
cyclo(GFPW^{me}VI^{me}V^{me}V^{me}G^{me}V^{me}IG), demonstrated that the 7-fold methylated species was
strongly preferred over the 6-fold methylated one (**Figure 3b,c and Figure 3-figure**
supplement 2). The only difference between the two peptide substrates is the α-N-
methylation of the glycine residue at the P1 site of the proteolytic cleavage.
288

OphP adopts a typical POP-fold with an unusual hydrophobic tunnel
OphP shares 35% and 38% sequence identity to PCY1 from *Saponaria vaccaria* and
GmPOPB from *Galerina marginata*, respectively, including conservation of the putative
catalytic triad residues Ser580, Asp665 and His701 (**Figure 4-figure supplement 1**). Despite
these similarities, OphP has two remarkable features that distinguish it from these and other
members of the POP family: its preference for multiply backbone N-methylated substrates
including a ^{me}Gly residue (rather than a proline residue) at the P1 site, and the apparent lack
of a site for the recognition of the substrate C-terminus (follower peptide) (**Figures 2 and 3;**
Figure 4-figure supplement 2). In order to elucidate the structural basis of these differences,
we determined the crystal structures of OphP alone (apo-structure) and in complex with
substrates and inhibitors.

300
We first determined the structure of the putatively catalytically inactive OphP variant S580A.
The structure of the apo-form was determined in space group P1 (eight monomers in the
asymmetric unit) to 1.9 Å by molecular replacement using GmPOPB structure (PDB entry
5N4C) as the search model. Visual analysis and the PISA server suggest OphP is a monomer
in the crystal and other POP enzymes. The eight monomers differ only in flexible loops some
of which are involved in crystal contacts. Similar to other POPs, the monomer has two
domains, an α/β hydrolase domain (residues 1–82, 453–738) and the seven-bladed β-
propeller domain (residues 83–452) (**Figure 4a**). All the monomers in the asymmetric unit
adopt a "closed" conformation with the two domains packed against each other which closes
off the direct route to the active site. OphP is most similar to GmPOPB (5N4B) with a root-
mean-square-deviation (RMSD) of 1.5 Å over 686 residues and closely related to PCY1
(5O3W, RMSD of 1.7 Å over 650 residues) with a more distant relation to porcine muscle
POP (1QFS, RMSD of 1.7 Å over 650 residues). Examination of the surface of OphP
revealed a large and neutral, hydrophobic tunnel in the middle of the β-propeller domain that
connects to the putative active site with its key residue S580 (**Figure 4b**). His701 of OphP
sits in a flexible loop region (Leu697–Thr707) which is only resolved in two chains (chain A
and chain H) where it adopts different conformations (**Figure 4-figure supplement 3**). In
both subunits His701 is around 15 Å away from Ser580 and, thus, a structural rearrangement
is required to form the canonical catalytic triad.
320
The structure of the covalent complex between OphP and ZPP (**Figure 4-figure supplement
3)** was obtained by soaking wild-type OphP crystals with 5 mM ZPP overnight. The structure
of the complex was determined at 2.0 Å resolution. Additional electron density, which we
fitted as ZPP covalently linked to Ser580 was seen in two subunits (**Figure 4-figure
supplement 3**). Crystal packing would appear to limit the ability of the domains to have
separated sufficiently to allow ZPP to access the active site. The calculated solvent content of
the crystals is below 40 % consistent with visual observation that there are no large solvent
channels that would allow large domain movements that do not disrupt crystal packing. The
benzene ring of ZPP is located in an aromatic hydrophobic pocket. The other two subunits
had weak electron density suggesting low occupancy. The overall structure of the OphP:ZPP
complex is largely unchanged from the apo-structure of OphP(S580A) (0.5 Å over 716 Cα
atoms. There are differences in loops Leu139–Ala146, Ser164–Met171, Ser179–Met195,
Pro222–Gly230, Leu696–Ser706 and Ala622–Tyr664. As a result, His701 has moved closer
(7.6 Å to serine) to both Asp665 and Ser580 to form the catalytic triad (**Figure 4-figure
supplement 3**). These results confirm Ser580 as the key catalytic residue and support our
suggestion of structural rearrangement of loops is required to form the catalytic triad.
337

^{me}Gly at the P1 site of the peptide substrate is coordinated by a specific binding pocket
To characterize the interactions of OphP with its presumed natural substrate, crystals of
OphP(S580A) were soaked with mainly 10-fold methylated Oph-18mer^{NL}
($\underline{W_1^{me}V_2I_3^{me}V_4^{me}V_5^{me}G_6}$-$^{me}V_7^{me}I_8^{me}G_9V_{10}^{me}I_{11}^{me}G_{12}S_{13}^{me}V_{14}M_{15}S_{16}T_{17}E_{18}$) and mainly 7-fold
methylated Oph-15mer peptide substrate
($\underline{G_1F_2P_3W_4^{me}V_5I_6^{me}V_7^{me}V_8^{me}G_9^{me}V_{10}^{me}I_{11}^{me}G_{12}}V_{13}I_{14}G_{15}$) (**Figure 5a-c and Figure 5-figure
supplement 1, respectively**) (both substrates yield dodecameric peptide macrocycles with
the underlined residues; the numbering of the residues eases the visualization of the structural
arrangement of the peptide in the active site of the protein displayed below). The Oph-
18mer^{NL} complex was determined to 2.0 Å and the Oph-15mer complex at 2.5 Å resolution.
The protein structures superimpose well with the apo structure with an RMSD of 0.4 Å over
715 Cα atoms (Oph-18mer^{NL}) and 0.8 Å over 714 Cα atoms (Oph-15mer). The peptide

substrates are located at the interface between the two domains with its N-terminus extending into the β-propeller domain and its C-terminus contacting the hydrolase domain (**Figure 5a; Figure 5-figure supplement 1**). Substrate binding displaced both glycerol and water molecules found in the apo-structure.

For Oph-18mer[NL], the clearest density for the substrate was found in subunit A. There is some weak electron density for the loop (698–704) which contains the catalytic histidine His701, but we did not model it. Only residues ^{me}I8^{me}G9V10^{me}I11^{me}G12S13^{me}V14 were fitted to the electron density whilst the remainder were presumed to be disordered (**Figure 5b**). The side chain of ^{me}I8 packs against Arg316 whilst ^{me}G9 makes almost no contact; the side chain of V10 interacts strongly with the aromatic ring of Phe617, ^{me}I11 makes a similar interaction with Phe502 whilst its N-methyl group points to solvent (**Figure 5c**). The P1 binding pocket that accommodates ^{me}G12 can be decomposed into two components (amide and side chain) for ease of discussion. The N-methyl group sits in a pocket formed by Phe502, Ile606 and Trp621 (indole makes van der Waal contact) (**Figure 5d**). The Cα of ^{me}G12 is surrounded by Ile606, Val668 and Ser580 (< 5 Å). Further away from the Cα of ^{me}G12 are the indole of Trp621 (6 Å) and the hydroxyl of Tyr625 (7 Å) which complete the pocket. The peptide bond between ^{me}G12 and S13 is positioned for nucleophilic attack by Ser580. The hydroxyl of Tyr499 and the backbone amide of Asn581 are well-positioned to function as the canonical oxyanion hole pair to stabilize the tetrahedral intermediate formed by the nucleophilic attack of Ser580. A different conformer of the side chain of Arg667 would contribute to the stabilization of the negatively charged intermediate. Both Tyr499 and Arg667 are found in many other members of the POP family where they play a similar role[52].

In the Oph-15mer complex, residues G1F2P3W4^{me}V5I6^{me}V7^{me}V8^{me}G9^{me}V10^{me}I11 were located in the electron density in subunit D (**Figure 5-figure supplement 1**). The peptide bond of ^{me}V8 is closest to Ser580 but its orientation is not consistent with catalysis (**Figure 5-figure supplement 1**). We conclude this complex is not directly relevant to understanding catalysis but rather is evidence for substrate binding plasticity.

To further investigate the role of Ile606 and Trp621 in substrate recognition and catalysis, we produced OphP variants I606A and W621A. Incubation with Oph-15mer (in almost exclusively 7-fold methylated form) showed that I606A is only slightly less active than the native enzyme while W621A had no activity (**Figure 5e**). We hypothesised that Ile606 might play a role in limiting the volume of the P1 pocket thus introducing selectivity. To test this, we explored whether bulkier side chains at P1 could be processed by the I606A variant of OphP. Unfortunately, we were unable to purify G12V and G12L substrate variants of Oph-15mer and the variant G12A resulted in a complex mixture that prevented further analysis.

Discussion

The multiply backbone N-methylated peptide macrocycle omphalotin A from *Omphalotus olearius* originates from a RiPP pathway involving the peptide auto-α-N-methyltransferase OphMA and the serine peptidase OphP (Matabaro, Kaspar, et al., 2021; Ramm et al., 2017; van der Velden et al., 2017). Similar macrocyclic peptides, lentinulin A and dendrothelin A, both with nine-fold backbone N-methylations have been isolated from *Lentinula edodes* and *Dendrothele bispora* harboring biosynthetic gene clusters containing OphMA and OphP homologs (Matabaro, Kaspar, et al., 2021) (**Figure 1a**). The simplest model of the biosynthetic pathway is that OphP represents a dual function macrocyclase, that releases a peptide intermediate (core plus follower peptide) from OphMA (function 1) and cleaves off the follower peptide and concomitantly macrocyclizes the core peptide to yield the final product omphalotin A (function 2).

OphP is not bifunctional

Our *in vitro* experiments with various backbone N-methylated peptide substrates showed oligopeptidase and macrocyclase activity of purified OphP (**Figures 2 and 3 including respective figure supplements**). In contrast, OphP showed no proteolytic activity towards full-length OphMA, irrespective of its methylation state (**Figure 1-figure supplement 3**). OphP was unable to remove unmethylated residues preceding the methylated core peptide, rather in some cases these additional residues were incorporated into the resulting macrocycles (**Figure 2**). We therefore conclude that OphP is not bifunctional and that other enzyme(s) are required to cleave OphMA into the peptide intermediate that then can be macrocyclized by OphP.

Since no candidate for such an endoproteinase has been identified in the gene clusters reported for the biosynthesis of omphalotin A in *O. olearius* (or homologs in *L. edodes* and *D. bispora* (Quijano et al., 2019; Ramm et al., 2017; van der Velden et al., 2017)), the most likely explanation for these results is the existence of a cellular endoproteinase that cleaves OphMA in the clasp region close to the core peptide in order to create the OphP substrate. It has previously been shown that the presumed cleavage site in OphMA is flexible and accessible (Song et al., 2018). In the pathway for the peptide macrocycle segetalin A, an oligopeptidase (OLP1) cleaves off a leader peptide from the precursor peptide to generate the substrate for the POP macrocyclase PCY1(Barber et al., 2013). Thus there is clear precedent for the requirement of separate enzymes. If the cellular endoproteinase responsible for the cleavage of OphMA does not cut after Pro399, additional aminopeptidases will be needed to remove eventual unmethylated residues preceding the omphalotin core peptide and expose Trp400 for macrocyclization of the core peptide by OphP. Based on the sensitivity of the *in vivo* but not *in vitro* production of omphalotin A to proteasomal inhibitors (**Figure 2-figure supplement 6**), the proteasome is a possible candidate for both activities. In agreement with this hypothesis, omphalotin A could be produced by heterologous expression of OphMA and OphP in the eukaryotic host *P. pastoris* (Matabaro, Kaspar, et al., 2021; Matabaro, Song, et al., 2021; Ramm et al., 2017) but not the prokaryotic host *E. coli* (**Figure 2-figure supplement 7**). Further experiments are needed to verify this hypothesis. Since no partially methylated omphalotin A macrocycles have been identified in nature or in expression hosts (Matabaro, Kaspar, et al., 2021; Mayer et al., 1997; Ramm et al., 2017), a critical question arises as to how it is ensured that only the completely methylated OphMA is processed.

Substrate is routed to the active site most likely through a central hydrophobic channel

The entry of the substrate into active site of POP family enzymes has been proposed to occur by different mechanisms. In one, a hinge between the two domains gives rise to an open

(substrate binding/product release) and closed (catalysis) conformation of the enzyme
(Canning et al., 2013; Czekster et al., 2017; Kaushik et al., 2014; Li et al., 2010; Shan et al.,
2005). Apo OphP(S580A) crystallized in a closed conformation in the absence of ligand
(**Figure 4a**). We obtained ZPP, Oph-18mer[NL] and Oph-15mer co-complexes by soaking the
peptide substrates into crystals, with only some small changes in active site loops but no large
domain movements. Crystal packing would seem to prevent the large closed to open to closed
conformational changes required by the "hinge" mechanism. We cannot exclude the
possibility of such gross movements and annealing in the crystal, but it would be unusual.
Thus, we rather considered two alternative routes for large peptides to reach the active site.
The first is an entrance between the two α/β and β-propeller domains (the side cavity)
(**Figure 4a)**. Such side entry has been observed previously, in dipeptidyl peptidase IV, a
prolyl-specific exopeptidase (Aertgeerts et al., 2004; Shan et al., 2005; Thoma et al., 2003).
However, this side cavity was much larger than that seen in apo OphP. In OphP, the
disordered loops Pro222–Gly230 and Leu697–Gly704 would be expected to further constrict
the side entry. Furthermore, the protein surface for this route is polar which does not match
the nature of the substrate (**Figure 4b**). We cannot exclude this means of entry but do not
consider it as very likely. A second and more plausible route is via the central hydrophobic
tunnel (**Figure 4b**) that leads to the cleft. Compared to the other POP enzymes (1NU6, 5O3U
and 5N4F), the tunnel in OphP is wider (**Figure 4-figure supplement 4)**. The apparent lack
of conformational change, the tunnel width match to substrate size, and the low polarity of
this route make the tunnel an attractive route of substrate entry (**Figure 4b**). Entry by this
route that has not been observed in any member of the POP family. Interestingly, in a study
of porcine muscle prolyl oligopeptidase (Fülöp et al., 1998; Fülöp et al., 2000), the authors
proposed the opening of the central tunnel of the β-propeller domain with blades 1 and 7
acting as the gate to allow substrate to gain access to the active site.
464

The tight P1 pocket controls selectivity
Detailed analysis of the products derived from the various substrates revealed a [me]Gly residue
as the point of macrocyclization (P1 site) (**Figures 2 and 3 including respective figure**
supplements). The requirement for [me]Gly contrasts the requirement for Pro at P1 which
defines the POP family ([me]Gly resembles Pro in the sense that both are tertiary amides).
Structural analysis shows no specific interaction of OphP with residues C-terminal to the core
peptide (**Figure 5 including respective figure supplements**). Accordingly, the six C-
terminal (follower) residues show no binding to OphP (**Figure 4-figure supplement 2**) and
there is no conservation in length or amino acid sequence of the follower residue of peptides
processed by OphP (**Figure 2a**). We conclude that OphP does not utilize the C-terminal
follower residues as a recognition sequence, in contrast to GmPOPB and PCY1 (Chekan et al.,
2017; Czekster & Naismith, 2017; Ludewig et al., 2018; Luo et al., 2014).
477

OphP tolerated some sequence variations within the core peptide to produce omphalotin A,
lentinulin A and dendrothelin A (Matabaro, Kaspar, et al., 2021; Ramm et al., 2017) (**Figure**
2e). The enzyme also makes macrocycles of different lengths, incorporating additional
residues at the N-terminus (**Figure 2a**). This plasticity is consistent with the observation that
the interactions between OphP and the peptide substrate side chains (**Figure 5c)** are
dominated by unspecific hydrophobic contacts. Indeed, comparison of the structures of OphP
complexed with Oph-15mer and -18mer reveals different side chains of the peptide substrates
in the same protein location consistent with recognition plasticity. Furthermore, the different
conformations of the catalytic triad residues observed in different structures, indicate
considerable flexibility of the pocket and thus capability to adapt to different substrates.
488

In the Oph-18mer^{NL} complex, ^{me}G12 is located at the P1 site consistent with the expected
product. The active site configuration creates an appropriately positioned oxy-anion hole for
catalysis. ^{me}Gly12 binds to a hydrophobic pocket formed by Trp621, Tyr625, Val668,
Asn581, Phe502 and Ile606 (**Figure 5d**). The N-methyl group interacts with Trp621 and the
mutant W621A is inactive (**Figure 5e**). These favourable hydrophobic interactions are not
possible for an unmethylated amide. N-methylation profoundly reduces (Simon et al., 1992)
the conformational flexibility of glycine (Lovell et al., 2003; Ramachandran et al., 1963) and,
thus, the entropic penalty for binding (^{me}Gly12 has Φ and Ψ dihedral angles that are similar to
the prolines (P1 site) in structures of PCY1 (PDB 5O3U) and GmPOPB (5N4C)). We suggest
this combination of enthalpy and entropic factors drives the preference of OphP (and its
relatives) for tertiary amide substrates.
500
Except for Ile606, the other residues that could interact with a side chain at P1 (Phe502,
Val668, Trp621, Tyr625) are strictly conserved across all POP enzymes to date (**Figure 4-**
figure supplement 1, Figure 5-figure supplement 2). Isoleucine is almost exclusively found
in OphP and closely related methylated peptide macrocyclases (**Figure 4-figure supplement**
1, Figure 5-Figure supplement 2), POP enzymes have the smaller Val in this position. A
simple model based on the Oph-18mer^{NL} complex which places a proline residue at P1 result
in a clash (distance < 2.5 Å) with Ile606 (**Figure 5-figure supplement 2**), We propose that
Ile606 selects against recognition of Pro. In support of this argument, the covalent OphP:ZPP
complex, which has a proline group at the P1 site, results in Ile606 adopting an unfavorable
conformation (**Figure 5-figure supplement 2**). This correlates with the lower-than-expected
inhibitory potency of ZPP against OphP (**Figure 2-figure supplement 6**). The ZPP result
does show the selection is not absolute and the pocket can undergo conformational change.
Gymnopeptides are members of the borosin family that includes the omphalotins (Quijano et
al., 2019) and they possess ^{me}Val at the P1 position. The valine side chain would clash not
only with Ile606 but also with Val668. The sequences of the OphP homologues of the
producing fungus *Gymnopus fusipes* are not known but we would predict their P1 pocket has
to be altered from OphP to to accomodate the larger side chain at the P1 position.
518
In summary, we show that OphP is not a dual function proteinase/macrocyclase and that *in*
vivo a third enzyme is almost certainly required for the production of ribosomally synthesized,
backbone N-methylated peptide macrocycles from auto-α-N-methylating precursor proteins
like OphMA. OphP recognises an N-methylated glycine rather than a proline residue at the
P1 site of macrocyclization using a hydrophobic pocket that selects against residues larger
than alanine whilst selecting for methylated amides. The hydrophobic and spacious nature of
the substrate-binding pocket results in few specific contacts endowing OphP with some
substrate promiscuity and the unique ability to cyclize multiply methylated substrates. These
unique properties of OphP may turn this enzyme into a valuable tool for producing highly α-
N-methylated cyclic peptides for pharmaceutical applications.
529
530

Materials and methods
Cells, antibodies, and reagents
ProAlanase (mass spec grade) was ordered from Promega. Expression plasmids for recombinant GmPOPB (plasmid PMA 1237: pJ414-GmPOPB) and TEV protease (pRK793_MBP-TEV site-His6-TEV-C4) were previously described (Czekster et al., 2017) and provided by Alvar Gossert (ETH Zürich, Switzerland), respectively. Phusion high-fidelity DNA polymerase was ordered from ThermoFisher Scientific (USA). Peptides AMA1 and PHA1 were chemically synthesized by GenScript Biotech (USA). The chromogenic substrate Gly-Pro-pNA and the POP inhibitor Z-Pro-prolinal (ZPP) were ordered from Sigma-Aldrich. Follower and leader (clasp domain) peptides were synthesized by Genscript.

Plasmid construction
The plasmids expressing OphP or LedP were constructed from pPIC3.5K-strepII-SUMO*-TEVcs-OphP (PMA1382) and pPIC3.5K-StrepII-SUMO*-TEVcs-LedP (PMA1379) that were previously used in omphalotin A production in *P. pastoris* (Matabaro, Kaspar, et al., 2021). The His$_8$-tag was fused to SUMO* by PCR amplification using PMA1379 as a template and His-SUMOSTAR-FW and His-SUMOSTAR-RV as primers. The obtained fragment was cloned into the *Spe*I and *BamH*I restriction sites of plasmids PMA1382 and PMA1379 to generate pPIC3.5K-His$_8$-SUMO*-TEVcs-OphP (PMA1596) and pPIC3.5K-His$_8$-SUMO*-TEVcs-LedP (PMA1595), respectively. The plasmid coding for His$_8$-SUMO*-TEVcs-OphP(S580A) was constructed by exchanging the *Not*I-*Spe*I restriction fragment of PMA1596 by the respective fragment of PMA1374 (pCDFDuet-sMBP-OphP(S580A)). For most of the His$_8$-OphMA-TEVcs constructs, one common primer OphMA-FW was used. For plasmids expressing OphMA-TEVcs-ΔVIG, OphMA-TEVcs-DbiCterm and OphMA-TEVcs-LedCterm, primers OphMA-ΔVIG-RV, OphMA-DbiCterm-RV, OphMA-LedCterm-RV were used together with OphMA-FW on PMA1488, respectively. The obtained PCR fragments were digested with *Not*I and *Nde*I and inserted in the respective sites on the pET24b vector. For the amplification of the coding region of His$_8$-OphMA-TEVcs-24mer, primer OphMA-24mer-RV was used on PMA1004. The PCR fragment was inserted into *Nde*I and *Nco*I of PMA1488. In all cases, the plasmids were amplified in *E. coli* DH5α. Plasmid sequences were confirmed by Sanger sequencing. Plasmids encoding His$_8$-SUMO*-TEVcs-OphP(I606A), His$_8$-SUMO*-TEVcs-OphP(W721A), His$_8$-OphMA-TEVcs-ΔC6(G408A), His$_8$-OphMA-TEVcs-ΔC6(G408V) and His$_8$-OphMA-TEVcs-ΔC6(G408L) were created using published protocols (Liu & Naismith, 2008). Plasmids PMA1488 (His$_8$-OphMA-TEVcs-21mer) and PMA1304 (His$_8$-OphMA-TEVcs-15mer) were part of our plasmid library. They were used to generate Oph-21mer and -15mer peptides, respectively. PMA1304 and PMA1488 were derived from plasmid PMA1004 (pET24-His$_8$-OphMA-cDNA) described earlier (van der Velden et al., 2017). Used primers and plasmids are listed in Supplementary File 1-Tables S1 and S2, respectively.

Production of recombinant OphP in *P. pastoris*
*Pichia pastori*s GS115 cells were transformed by electroporation using the plasmids pPIC3.5K-His$_8$-SUMO*-TEVcs-OphP or pPIC3.5K-His$_8$-SUMO*-TEVcs-LedP, linearized by *Bsp*EI for the integration into the *HIS4* locus. Transformants were selected first on a standard SD-HIS minimal medium and then on 250 µg/ml geneticin for high-copy number screening. To produce OphP, cells were cultured at 20 °C in a standard methanol-containing complex medium (BMMY) for three days following the recommendations for *Pichia* protein expression(Matabaro, Song, et al., 2021). For harvesting, cells were separated from the culture medium by centrifugation, and cell pellets were washed with cold lysing buffer (2xPBS) or double-distilled water (dd H$_2$O) and centrifuged again at $8000 \times g$ for 10 min at

4 °C. Cell pellets were resuspended in cell lysis buffer [2xPBS, 10% glycerol, 1 mM DTT, 20
mM imidazole, and protease inhibitor cocktail (PIC) (Roche)]. Cells lysis was performed on a
planetary mill (Pulverisette 7) as previously described (Matabaro, Song, et al., 2021). Cell
debris were removed by spinning down the cell lysate at 16, 000 $x\ g$ for 30 min at 4 °C. The
cell lysate supernatant was collected and subjected to purification via a Ni^{2+}-based
immobilized metal affinity chromatography (Ni-IMAC) system using the washing buffer (50
mM HEPES pH 8.0, 10% glycerol and 20 mM imidazole), and elution buffer (50mM HEPES
pH 8.0, 10% glycerol and 400 mM imidazole). The purified proteins were stored in storage
buffer (50 mM HEPES pH 8.0 + 10% glycerol + 5 mM DTT) at -80 °C. The His_8SUMO* tag
was cleaved off by incubation with TEV protease (1:10, w/w) in assay buffer [300 mM NaCl,
50 mM TRIS pH 8.0, 1mM DTT and 10% Glycerol] overnight, at 25 °C. The reaction
mixture was reloaded on a Ni-IMAC system to remove the His_8-SUMO* tag. The fractions
containing OphP were collected and concentrated using an Amicon filter column (50 kDa
cut-off). OphP was further polished across an Äkta FPLC system for size exclusion
chromatography (SEC) equipped with a Superdex 200 Increase column. For the this final
purification step, HEPES buffer [50 mM HEPES pH 8, 1 mM DTT, 10% glycerol (v/v)] was
used, as previously described (Matabaro, Song, et al., 2021).
598
Protein expression in *E. coli*
OphMA, OphMA-TEV, GmPOPB, and TEV protease were produced in *E. coli* BL21 as
previously described (Czekster et al., 2017; Matabaro, Song, et al., 2021). Briefly, chemically
competent BL21 cells were transformed with respective expression plasmids and selected on
appropriate antibiotics. A pre-culture was incubated overnight in 10 ml at 37 ºC, 180 rpm. 5
ml of the preculture was used to inoculate 1 L of terrific broth (TB) medium. The culture was
maintained at 37ºC, 160 rpm until OD_{600} of 1.5 to 2.0. The culture was chilled on ice for 30
min before the addition of 200 µM isopropyl β-D-thiogalactopyranoside (IPTG) and
maintained at 16 ºC, 160 rpm for 2 days (GmPOPB and TEV) and 3-5 days for OphMA and
OphMA-TEV mutants. Cells were harvested by centrifugation at $8000 \times g$ for 20min, at 4ºC.
Cells pellets were resuspended in 100 mL ice-cold lysis buffer [300 mM NaCl, 50 mM Tris
pH 8.0, 10% glycerol, 1 mM DTT, DNase I, 20 mM Imidazole and protease inhibitor cocktail
(Roche)]. For OphMA and OphMA-TEV mutants, NaCl and Tris in the buffer above, were
replaced by 50mM HEPES pH 8.0, 0.1% Triton x-100. Cells were lysed by a cold French
press, two passages, and clearing of cell debris by centrifugation at 12 000 rpm, 30 min at 4
ºC. The cell lysate supernatant was subjected to protein purification either by Ni-IMAC
systems (for TEV and GmPOPB) or by Ni-NTA beads (Macherey Nagel). Elution and wash
buffers and the SEC method were the same as for OphP purification above. The protein
concentration was determined by BCA assay (Thermo Scientific). To confirm the protein
quality, 10 µg of each protein was loaded on a 12% SDS gel and run by electrophoresis and
stained by coomassie brilliant blue. The purified proteins were shock-frozen in liquid
nitrogen and then stored at -80 ºC.
621
Peptide production and purification
To produce OphMA-C-terminal fragments, purified and TEV-cleavable OphMA proteins
were mixed with TEV protease (1:10) to a concentration of 10 mg/ml in assay buffer [300
mM NaCl, 50 mM TRIS pH 8.0, 1 mM DTT and 10 % glycerol]. The reaction was mixed
well by pipetting and kept overnight at 25 °C. The reaction mixture was loaded on a pre-
equilibrated C8 or C18-SepPak cartridge following the user instruction and
recommendations. The peptides were eluted in 1-3 ml 100% methanol depending on the
concentration. 5 µl of the eluate was used for HPLC-MS/MS analysis before further
purification on an Agilent 1260 infinity preparative HPLC harboring a C18 column (Luna 5

μm C18(2) 100Å, 250X10 mm. The mobile phase consisted of a gradient between water (A)
and acetonitrile (B), both supplemented with 0.1% formic acid. The flow rate was set at 1.2
ml/min, and one fraction each minute. The method started with a system equilibration for 5
min at 5% B, followed by a linear gradient up to 40 % B in 10 min, then increased to 98% B
in 30 min. The column was finally washed with 98% B for 5 min before gradually dropping
to 5% B in a 5 min span. Fractions corresponding to the expected products were collected,
and 5 µl of each fraction are tested by HPLC-MS/MS for the presence of peptides. To
determine the concentration, samples were analysed by an Agilent 1100 series UV-HPLC
calibrated to the bovine albumin serum (BSA) as an internal standard, at λ = 210 nm. To
produce Oph-30mer, recombinant His_8-OphMA lacking a TEV protease cleavage site was
digested with trypsin. For this purpose, 100 µg of the purified protein was incubated with
1.25 mg trypsin in 25 µl HEPES buffer pH 8.0, 37 °C, 650 rpm, overnight. 3 µl of the
reaction was used for MS analysis. To purify the peptide, C18 cartridges (Sep-Pak C18 1cc
(100mg), Waters) were used. The cartridge was first flushed with 8 ml 100% methanol and
then equilibrated with 8 ml water. The trypsin reaction mixture was then loaded on the
cartridge, followed by 3 washes with 1 mL water supplemented with 1% methanol. The
peptides were eluted in 1 mL 100% methanol. The sample was concentrated by evaporating
the solvent using a speed vac, and the resulting peptide pellets were resuspended in the assay
buffer.
650
651 **Protein sequences (the portion used after TEV protease cleavage site (cs) is underlined,**
652 **the TEV protease cleavage site is in bold)**
653
His_8OphMA-
TEVcs(21mer)MEHHHHHHHHTSTQTKAGSLTIVGTGIESIGQMTLQALSYIEAAAKVF
YCVIDPATEAFILTKNKNCVDLYQYYDNGKSRLNTYTQMSELMVREVRKGLDVVG
VFYGHPGVFVNPSHRALAIAKSEGYRARMLPGVSAEDCLFADLCIDPSNPGCLTYEA
SDFLIRDRPVSIHSHLVLFQVGCVGIADFNFTGFDNNKFGVLVDRLEQEYGAEHPVV
HYIAAMMPHQDPVTDKYTVAQLREPEIAKRVGGVSTFYIPPKARKASNLDIIRRLELL
PAGQVPDKKARIYPANQWEPDVPEVEPYRPSDQAAIAQLADHAPPEQYQPLATSKA
MSDVMTKLALDPKALADYKADHRAFAQSVPDLTPQERAALELGDSWAIRCAMKN
MPSSLLDAARESG**ENLYFQG**FPWVIVVGVIGVIGSVMSTE
663
His_8OphMA-TEVcs-ΔVIG(18mer)
MEHHHHHHHHTSTQTKAGSLTIVGTGIESIGQMTLQALSYIEAAAKVFYCVIDPATE
AFILTKNKNCVDLYQYYDNGKSRLNTYTQMSELMVREVRKGLDVVGVFYGHPGVF
VNPSHRALAIAKSEGYRARMLPGVSAEDCLFADLCIDPSNPGCLTYEASDFLIRDRPV
SIHSHLVLFQVGCVGIADFNFTGFDNNKFGVLVDRLEQEYGAEHPVVHYIAAMMPH
QDPVTDKYTVAQLREPEIAKRVGGVSTFYIPPKARKASNLDIIRRLELLPAGQVPDKK
ARIYPANQWEPDVPEVEPYRPSDQAAIAQLADHAPPEQYQPLATSKAMSDVMTKLA
LDPKALADYKADHRAFAQSVPDLTPQERAALELGDSWAIRCAMKNMPSSLLDAAR
ESG**ENLYFQG**FPWVIVVGVIGVIGSVMSTE
673
His_8OphMA-TEVcs-ΔC6(15mer)
MEHHHHHHHHTSTQTKAGSLTIVGTGIESIGQMTLQALSYIEAAAKVFYCVIDPATE
AFILTKNKNCVDLYQYYDNGKSRLNTYTQMSELMVREVRKGLDVVGVFYGHPGVF
VNPSHRALAIAKSEGYRARMLPGVSAEDCLFADLCIDPSNPGCLTYEASDFLIRDRPV
SIHSHLVLFQVGCVGIADFNFTGFDNNKFGVLVDRLEQEYGAEHPVVHYIAAMMPH
QDPVTDKYTVAQLREPEIAKRVGGVSTFYIPPKARKASNLDIIRRLELLPAGQVPDKK
ARIYPANQWEPDVPEVEPYRPSDQAAIAQLADHAPPEQYQPLATSKAMSDVMTKLA

LDPKALADYKADHRAFAQSVPDLTPQERAALELGDSWAIRCAMKNMPSSLLDAAR
ESG**ENLYFQG**FPWVIVVGVIGVIG

His₈OphMA-TEVcs-24mer.
MEHHHHHHHHTSTQTKAGSLTIVGTGIESIGQMTLQALSYIEAAAKVFYCVIDPATE
AFILTKNKNCVDLYQYYDNGKSRLNTYTQMSELMVREVRKGLDVVGVFYGHPGVF
VNPSHRALAIAKSEGYRARMLPGVSAEDCLFADLCIDPSNPGCLTYEASDFLIRDRPV
SIHSHLVLFQVGCVGIADFNFTGFDNNKFGVLVDRLEQEYGAEHPVVHYIAAMMPH
QDPVTDKYTVAQLREPEIAKRVGGVSTFYIPPKARKASNLDIIRRLELLPAGQVPDKK
ARIYPANQWEPDVPEVEPYRPSDQAAIAQLADHAPPEQYQPLATSKAMSDVMTKLA
LDPKALADYKADHRAFAQSVPDLTPQERAALELGDSWAIRCAMKNMPSSLLDAA**E**
NLYFQGSQNGFPWVIVVGVIGVIGSVMSTE

His8-SUMO*-TEVcs-OphP
MGHHHHHHHHGGSDSEVNQEAKPEVKPEVKPETHINLKVSDGSSEIFFKIKKTTPLR
RLMEAFAKRQGKEMDSLTFLYDGIEIQADQTPEDLDMEDNDIIEAHREQIGG**ENLYF**
QGTSMSFPGWGPYPPVERDETSAITYSSKLHGSVTVRDPYSQLEVPFEDSEETKAFV
HSQRKFARTYLDENPDREAWLETLKKSWNYRRFSALKPESDGHYYFEYNDGLQSQL
SLYRVRMGEEDTVLTESGPGGELFFNPNLLSLDGNAALTGFVMSPCGNYWAYGVSE
HGSDWMSIYVRKTSSPHLPSQERGKDPGRMNDKIRHVRFFIVSWTSDSKGFFYSRYP
PEDDEGKGNAPAMNCMVYYHRIGEDQESDVLVHEDPEHPFWISSVQLTPSGRYILFA
ASRDASHTQLVKIADLHENDIGTNMKWKNLHDPWEARFTIVGDEGSKIYFMTNLKA
KNYKVATFDANHPDEGLTTLIAEDPNAFLVSASIHAQDKLLLVYLRNASHEIHIRDLT
TGKPLGRIFEDLLGQFMVSGRRQDNDIFVLFSSFLSPGTVYRYTFGEEKGYRSLFRAIS
IPGLNLDDFMTESVFYPSKDGTSVHMFITRPKDVLLDGTSPVLQYGYGGFSLAMLPT
FSLSTLLFCKIYRAIYAIPNIRGGSEYGESWHREGMLDKKQNVFDDFNAATEWLIAN
KYASKDRIAIRGGSNGGVLTTACANQAPGLYRCVITIEGIIDMLRFPKFTFGASWRSE
YGDPEDPEDFDFIFKYSPYHNIPPPGDTVMPAMLFFTAAYDDRVSPLHTFKHVAALQ
HNFPKGPNPCLMRIDLNSGHFAGKSTQEMLEETADEYSFIGKSMGLTMQTQGSVDSS
RWSCVTV

Enzyme activity assays
For the chromogenic substrate and the POP inhibitor, stocks of 50 mM and 10 mM,
respectively, were prepared in methanol. Before the reaction, the substrate was solubilized at
40 °C and then the required amounts were mixed with the standard assay buffer [50 mM
HEPES pH 6.0 + 10 mM DTT]. The enzyme was added last, to make the total reaction
volume 50 µL. The reaction mixture was kept at 30°C while shaking at 600 rpm. The reaction
was quenched by the addition of 50 µl methanol. The enzymatic activity was monitored by
absorbance measurement at 410 nm (for *p*-nitroanilide) on an Infinite 200Pro M Plex
spectrophotometer (Tecan). Continuous reaction monitoring was ensured by incubating the
96-well plates at 30°C, with absorbance measurements each 30 min.

For peptide substrates, reactions were also carried out in 50 µl with a standard buffer [50 mM
HEPES pH 7.0, 10 mM DTT], at 30°C, 600 rpm unless stated otherwise. The peptide was
first mixed with the buffer before adding the enzymes. Both substrate and enzyme
concentrations varied up to 100 µM. The reaction was stopped by the addition of 50 µl
methanol, then spun down at the highest speed (20,000 *x g*), 4ºC. For HPLC-MS/MS analysis,
5 µl supernatant was used. For the synthetic peptides, PHA1 and AMA1 were stored in 1
mg/ml. For the reaction, 20 µM substrate was used with 1 µM of protease. For ProAlanase, a
1:50 enzyme/substrate (w/w) was used in a 25 µl reaction volume, in either 50 mM HEPES

(pH 6.0) or 50 mM sodium acetate (pH 4.0). These reactions were carried out at 37°C for 4 h
and stopped by adding 25 µl methanol. After centrifugation at the highest speed at 4 °C, 5 µl
supernatant was used for HPLC-MS/MS analysis.
734
For the time course experiment with OphP and Oph-15mer, the reactions were performed in
buffer containing 20 mM HEPES (pH 7.0), 100 mM KOAc, 2mM Mg(OAc)2, 10% glycerol
and 10 mM DTT at a concentration of 100 μM total Oph-15mer (90 μM 7-fold methylation,
10 μM 6-fold methylation) and 10 μM OphP. The reactions were performed in 70 μl reaction
volume at 30°C and at each time point, 5 μl were sampled and added to 20 μl of MeOH to
stop the reaction. The samples were centrifuged for 5 minutes at 15000 rcf and 10 μl were
injected for HPLC-MS/MS analysis.
742
For the experiment of OphP with Oph-15mer (mixture of 7-fold and 6-fold methylations) and
protease inhibitors (**Figure 2-figure supplement 6**), the reactions were performed in buffer
containing 20 mM HEPES (pH 7.0), 100 mM KOAc, 2mM Mg(OAc)2, 10% glycerol and 10
mM DTT at a concentration of 100 μM total Oph-15mer (90 μM 7-fold methylation, 10 μM
6-fold methylation) and 10 μM OphP. DMSO stocks of the different inhibitors were diluted
in the HEPES buffer and added to the reaction to the indicated final concentration. The
highest used DMSO concentration was used in a control reaction without inhibitors The
reactions were performed in 5 μl reaction volume at 30°C and 20 μl of MeOH were added
after 30 minutes to stop the reaction. The samples were centrifuged for 5 minutes at 15000 xg
and 10 μl were injected for HPLC-MS/MS analysis.
For the experiment testing for preferred production of 7-fold over 6-fold methylated species
(**Figure 3-figure supplement 2**), the reactions were performed as described above, but with
varying Oph-15mer substrate concentrations of 100 μM and 1000 μM total Oph-15mer
substrate.
757
For assays of OphP variants with Oph-15mer (almost pure 7-fold methylated peptide), the
reactions were performed in buffer containing 50 mM HEPES (pH 7.0), 100 mM NaCl, and 5
mM DTT at a concentration of 200 μM Oph-15mer and 20 μM OphP. The reaction mixtures
in Eppendorf tubes were incubated at 25 $^{\circ}$C and 1000 rpm using a ThermoMixer C for 18
hours. Reactions were quenched by adding an equal amount of methanol and protein
precipitation was removed by centrifugation. The supernatant was analysed using Waters
Acquity H-Class plus UPLC equipped with SQ, PDA and ELS detectors. Buffer A (water
with 0.1% FA) and buffer B (acetonitrile with 0.1% FA) are used as mobile phase. Typically,
10 µL of supernatant was loaded and separated on a ACQUITY UPLC BEH C18 column (1.7
µM, 2.1 x 100 mm) at a flow rate of 0.6 ml/min with following gradient: 0-0.25 min for 5%
buffer B, 0.25-0.4 min from 5% to 40% buffer B, 0.4-3.6 min from 40% to 55% buffer B,
3.6-3.7 min from 55% to 60% buffer B, 3.7-6.0 min from 60% to 70% buffer B, 6.0-6.15 min
from 70% to 95% buffer B, 6.15-7.6 min for 95% buffer B.
771
LC-ESI-MS/MS analysis
Peptide analysis was performed by following previous protocols(Matabaro, Song, et al., 2021)
for HPLC-MS/MS analysis of borosins. Briefly, data were generated from Dionex Ultimate
3000 UHPLC HPLC coupled to an MS/MS system (Thermo Scientific Q Exactive classic). A
list of expected masses (inclusion list) was always included in the HPLC-MS/MS method.
The peptide was applied on a C18 column (Phenomenex Kinetex ®µm XB-C18 100 Å (150 x
2.1 mm)) pre-heated at 50 °C. For UV-vis spectra, the signals were recorded by a diode array
detector (DAD) set at 210, 230, 250, and 280 nm. Water (solvent A) and acetonitrile (solvent
B) both containing 0.1% FA served as a mobile phase, with a 0.5 ml/min flow rate. The

HPLC method started by equilibrating the system for 2 min at 5% B, followed by a linear
gradient up to 70 % B in 13 min, and then increased to 100% B in 3 min. The column was
rinsed first with 100% B for 1 min and then with 5% B for 5 min. The MS/MS system used
the heated electrospray ionization in a positive ion mode. The full MS was set as follows: an
automatic gain control (AGC) target of 70,000 [AGC target 1e6, maximum ion trap (IT), 120
ms, and scan range of 200-2500m/z]. Data-dependent (dd) MS/MS was performed at a
resolution of 17,500 [AGC target at 1e5, maximum IT 120 ms, and isolation window of 3.0
m/z]. The normalized collision energy (NCE) was set at 30% for macrocycles, and 18% or
stepped NCE (16, 18, 22%) for linear peptides.
790
ESI-TOF MS and intact mass measurement
Intact protein mass analysis was performed using a Waters Xevo G2-QS QToF MS coupled
to an ACQUITY UPLC I-Class LC system (Waters Corporation). The spectrometer was auto-
calibrated throughout the experiment using an internal lock-spray mass calibrant (at 556.2771
m/z of leucine enkephalin, 1 s scan every 120 s interval). The data was acquired in
Continuum format, in a 300 – 2500 m/z spectral window with a scan time of 1 s and an
interscan time of 0.014 s. The mass spectrometer and ESI ionization source were operated
under the following parameters: capillary voltage at 3 kV, cone voltage at 20 V, source offset
voltage at 40 V, source temperature at 100 °C, desolvation temperature at 400 °C and
desolvation gas (nitrogen) flow at 650 L/h, cone gas flow at 30 L/h. In a typical analysis, 1
µL (30 ng) of the protein sample was injected and desalted on a ProSwift™ RP-1S HPLC
(250 x 4.6 mm x 5 µm, Thermo Scientific™) column maintained at 60 ^{o}C. The elution was
performed using solvent A (water with 0.1% formic acid) and solvent B (acetonitrile with 0.1%
formic acid) with a gradient elution using isocratic 95% solvent A (0 - 1 min), linear gradient
to 95% solvent B (1.1 – 7 min), and isocratic 95% solvent A (7.1 – 10 min) at a flowrate of
0.2 ml/min. Combined mass spectra from the total chromatographic protein peak were used
for intact mass reconstruction using MassLynx (Waters) Max(imum) Ent(ropy) 1
deconvolution algorithm (Resolution: 0.5 Da/channel, Width at half-height: ion
series/protein-dependent, Minimum intensity ratios: 33% Left and Right). Spectra were
deconvoluted between 40000 and 55000 for OphMA.
811
ITC studies of OphP with peptides
Isothermal titration calorimetry (ITC) experiments were performed using a MicroCal PEAQ-
ITC (Malvern). Leader peptides and follower peptides (SVMSTE and VIGSVMSTE) were
dissolved in DMSO at sufficient concentration as a stock. A typical ITC experiment was
performed by repeated injections 2 µL of peptide solution eighteen times into a cell solution
containing 0.4 ml of S580A. The measurements were done at 25 ^{o}C at a stirring rate of 750
rpm with a reference power (DP) of 5 mcal/s or 10 mcal/s. Control titrations were performed
similarly except for the absence of protein in the cell solution. Data processing and fitting
were performed using MicroCal PEAQ-ITC analysis software (Malvern).
821
Crystallization, data collection, reduction, and refinement
Freshly purified OphP and S580A proteins were screened for apo crystals using the sitting
drop method using commercial screening kits from Hampton Research or Molecular
Dimensions. OphP and S580A were eventually crystallized at 40-50 mg/ml by mixing 0.3 µL
of proteins with 0.3 µL of the reservoir (0.2 M sodium acetate, 0.1 M Bis-Tris propane (pH
6.0-6.5) and 28% PEG 3350). Crystals appeared within eight days and grew to full size
within three weeks at room temperature. Crystals were fished, transferred to cryo-protectant
(reservoir solution supplemented with 40% PEG 3350) and flash-frozen in liquid nitrogen.
The structure of the OphP-ZPP complex was determined by soaking OphP crystals with 5

mM Z-Pro-prolinal overnight at room temperature. Crystals were in similar cryoprotectant
with 1 mM ZPP. For the S580A:Oph-15mer complex, apo S580A crystals were soaked with
1 mM purified Oph-15mer overnight at room temperature. Due to the poor solubility of Oph-
18mer$^{\mathrm{NL}}$, apo OphP(S580A) crystals were soaked with a cloudy reservoir solution containing
18mer overnight to obtain the OphP(S580A)-18mer$^{\mathrm{CF}}$ complex. More than 50 crystals dataset
were collected for each complex and screened based on the density.
837
All the X-ray diffraction data were recorded at Diamond Light Source beamlines (I03, I04,
I04-1, and I24). The diffraction images were reduced, integrated, and scaled using
xia2(Winter, 2010), DIALS(Winter et al., 2018), Apo structures of S580A were determined
by molecular replacement using a Phaser(McCoy et al., 2007) with GmPOPB structure (PDB
entry 5N4C) as the initial model. Autobuilding was performed using Buccanner from the
molecular replacement solutions and further refined using Refmac5(Murshudov et al., 1997)
and manually built with Coot(Emsley et al., 2010), and improved using PDB-redo(Joosten et
al., 2012). Simulated omit maps for the ligands were calculated using Phenix(Liebschner et
al., 2019). All the other structures were solved by molecular replacement using apo S580A
structure as initial models, refined, manually built and improved similarly. All
crystallographic figures were generated using Pymol (Schrödinger LLC), CCP4mg
(McNicholas et al., 2011) or Chimera(Pettersen et al., 2004). Structure-based sequence
alignment was created using MUSCLE (Edgar, 2004) and ESPript 3.0 (Robert & Gouet,
2014). Full crystallographic details are given in Table 3.
852

Acknowledgements
We would like to thank Prof. Jörn Piel and his staff, in particular, Dr. Clara Chepkirui and Dr.
Silke Probst for their support in the HPLC purification, and Philip Rößler for supplying TEV
protease and helping with protein purification. We are grateful to Dr. Chris Arter from
Rosalind Franklin Institute for the assistance with the MS experiments and Dr. Xiaping Fu
for the assistance with the purification of peptides. We would also like to thank Diamond
Light source for the access to the synchrotron. This work was funded by the Swiss National
Science Foundation (Grant no. 31003A-173097), ETH Zürich and UK BBSRC
(BB/R018189/1).
862

863 **Competing interests**
864 MK is an inventor on a patent application filed by ETH Zurich (no. WO2017174760A1,
865 priority date: 7 April 2016). The authors declare no other competing interests.
866

867 **Data availability**
868 Crystallographic and structural data reported in the paper are available from the RCSB; 7ZB2
869 (apo-structure), 7ZAZ (ZPP complex). 7ZB0 (Oph-15mer complex) 7ZB1 (Oph-18mer
870 complex). Raw data for the Figures are shown in Supporting Information.
871

 Figure legends
Figure 1. Biosynthesis of omphalotin A inferred from the co-expression of OphMA and
OphP in *Pichia pastoris*. (**a**). Proposed biosynthetic pathway of omphalotin A, where
scissors indicate proteolytic cleavage (black) and macrocyclization sites (green). (**b**).
Sequences and chemical structures of omphalotin A and homologous peptide natural products
dendrothelin A and lentinulin A shown with methylation sites in orange and the cyclization
site in green. These peptide macrocycles were produced in *Pichia pastoris* by expressing
OphP together with OphMA variants with respective C-termini.
880
Figure 2. Macrocyclization of OphMA-derived peptide substrates by OphP *in vitro*. (**a**)
Sequences of peptides produced in this work, with α-N-methylated residues highlighted in
orange. Note that methylation by OphMA proceeds from N- to C-terminus and that only the
completely methylated forms of the peptides are displayed. The P1 residues recognized by
OphP are highlighted in purple and the P1' residues in grey. Leader peptide removal was
achieved by incubating TEV protease-cleavable OphMA variants with TEV protease and
GmPOPB or ProAlanase. Macrocyclization of the peptides by OphP is indicated by green
brackets above the peptide sequence. Note that only the macrocyclic products are shown. (**b**
and c) Extracted ion chromatograms (EICs) of macrocyclic peptides produced by incubation
of Oph-24mer (**b**) or Oph-21mer (**c**)with OphP. (**d**) RP-HPLC UV absorption profiles at 280
nm for Oph-15mer or Oph-18mer^{Tr} in the absence and presence of OphP showing the
production of the same macrocyclic peptide. (**e**) MS-analysis of omphalotin A and its
orthologues lentinulin A and dendrothelin A produced from Oph-18mer^{NL}, Led-18mer^{NL} and
Dbi-18mer^{NL} using OphP. The MS chromatograms show masses of both protonated cyclic
peptides and corresponding sodium and ammonium adducts.
896
Figure 3. Substrate preference of OphP. (**a**) Mass spectrometric analysis showing ion
chromatograms of peptide substrates Oph-15mer and Oph-18mer^{NL} before and after
processing by OphP. Red dashed lines indicate the various methylation states of the
individual peptides. For comparison, the peak height of the most abundant non-reactive
species was drawn at the same level (6-fold methylated species of Oph-15mer and Oph-
18mer^{NL}). These species were marked with blue asterisks. The sequence of the various
peptide species and their processing by OphP is indicated on the right. Green ticks stand for
processing and red crosses for lack of processing by OphP. (**b,c**) Time course of *in vitro*
reaction between 10µM OphP and 100µM total Oph-15mer Substrate (mixture of 6Me and
7Me species) in HEPES buffer at pH 7.0. EIC peak areas of 6- and 7-fold methylated peptide
substrate (**b**) and product (**c**) species were measured at 7 time points over 4 hours. Left panels
show the absolute EIC peak areas and the right panels the ratio between the 7- to 6-fold
methylated peptide species over the 7 time points.
910
Figure 4. Crystal structures of OphP and possible substrate tunnel. (**a**) In all structures in
this study OphP is found with the two domains close together (so called closed state). The β-
propellor domain is in blue and the N-terminal domain in green. The active site is located at
the interface of the domains. A molecule of the inhibitor ZPP (carbon atoms yellow, oxygen
red, and nitrogen blue) shown in sphere form has been placed at the active site circled in red
to aid visualization. Experimental details of the ZPP complex structure are shown **Figure 4-**
figure supplement 3. On the right is the electrostatic surface calculated APBS-PDB2PQR
software suite(Jurrus et al., 2018) under the default setting and drawn in PyMol. The
electrostatic potential is set at ± 5 kT/e. There is no access to the active site (black circle)
from the side of the structure in the closed state. (**b**) Rotation of 90° around the X-axis shows
there is a tunnel through the β-propellor domain (circled in red in cartoon). On the right is the

is electrostatic surface calculated as above, the tunnel allows access to the active site (black
circle). The yellow carbon atoms of ZPP can be seen through the pore in both representations.
924
Figure 5. ^{me}Gly12 recognition site of OphP. (**a**) OphP S580A complex to peptide substrate
Oph-18mer^{NL} is shown in spheres. (The color schemes is as **Figure 4a**) (**b**) The σ_A-weighted
2mFo-DFc map (contoured at 1σ) for Oph-18mer^{NL} (**c**) LigPlot diagram (Laskowski &
Swindells, 2011) showing interactions of the 18mer^{NL} with OphP. (**d**) Surfaces of ^{me}Gly12
and its surrounding hydrophobic pocket in Oph-18mer^{NL} complex. The pocket is composed
of Tyr625, Trp621, Val668, Phe502, Asn581 and Ile606. (**e**) HPLC analysis of assays using
almost pure 7-methylated Oph-15mer as substrate for OphP and its variants. The control
reaction without OphP shows the absorption profile of the starting material. While the
W621A mutation abolished macrocyclization, I606A showed an activity comparable to wild
type OphP. The reactions were performed at 25 °C in buffer containing 50 mM HEPES pH
7.0, 100 mM NaCl and 5 mM DTT at a concentration of 20 µM enzymes and 200 µM Oph-
15mer.
937

References

Aertgeerts, K., Ye, S., Tennant, M. G., Kraus, M. L., Rogers, J., Sang, B. C., Skene, R. J., Webb, D. R., & Prasad, G. S. (2004). Crystal structure of human dipeptidyl peptidase IV in complex with a decapeptide reveals details on substrate specificity and tetrahedral intermediate formation. *Protein Science*, *13*(2), 412-421.

Alex, A., Millan, D. S., Perez, M., Wakenhut, F., & Whitlock, G. A. (2011). Intramolecular hydrogen bonding to improve membrane permeability and absorption in beyond rule of five chemical space. *MedChemComm*, *2*(7), 669-674.

Arnison, P. G., Bibb, M. J., Bierbaum, G., Bowers, A. A., Bugni, T. S., Bulaj, G., Camarero, J. A., Campopiano, D. J., Challis, G. L., & Clardy, J. (2013). Ribosomally synthesized and post-translationally modified peptide natural products: overview and recommendations for a universal nomenclature. *Natural product reports*, *30*(1), 108-160.

Barber, C. J., Pujara, P. T., Reed, D. W., Chiwocha, S., Zhang, H., & Covello, P. S. (2013). The two-step biosynthesis of cyclic peptides from linear precursors in a member of the plant family Caryophyllaceae involves cyclization by a serine protease-like enzyme. *Journal of Biological Chemistry*, *288*(18), 12500-12510.

Canning, P., Rea, D., Morty, R. E., & Fülöp, V. (2013). Crystal structures of Trypanosoma brucei oligopeptidase B broaden the paradigm of catalytic regulation in prolyl oligopeptidase family enzymes. *PloS one*, *8*(11), e79349.

Chatterjee, J., Gilon, C., Hoffman, A., & Kessler, H. (2008). N-methylation of peptides: a new perspective in medicinal chemistry. *Accounts of chemical research*, *41*(10), 1331-1342.

Chekan, J. R., Estrada, P., Covello, P. S., & Nair, S. K. (2017). Characterization of the macrocyclase involved in the biosynthesis of RiPP cyclic peptides in plants. *Proceedings of the National Academy of Sciences*, *114*(25), 6551-6556.

Cohen, D. J., Loertscher, R., Rubin, M. F., Tilney, N. L., Carpenter, C. B., & Strom, T. B. (1984). Cyclosporine: a new immunosuppressive agent for organ transplantation. *Annals of internal medicine*, *101*(5), 667-682.

Czekster, C. M., Ludewig, H., McMahon, S. A., & Naismith, J. H. (2017). Characterization of a dual function macrocyclase enables design and use of efficient macrocyclization substrates. *Nature communications*, *8*(1), 1-10.

Czekster, C. M., & Naismith, J. H. (2017). Kinetic landscape of a peptide bond-forming prolyl oligopeptidase. *Biochemistry*, *56*(15), 2086-2095.

Dittmann, J., Wenger, R. M., Kleinkauf, H., & Lawen, A. (1994). Mechanism of cyclosporin A biosynthesis. Evidence for synthesis via a single linear undecapeptide precursor. *Journal of Biological Chemistry*, *269*(4), 2841-2846.

Donia, M. S., Ravel, J., & Schmidt, E. W. (2008). A global assembly line for cyanobactins. *Nature chemical biology*, *4*(6), 341-343.

Dreyfuss, M., Härri, E., Hofmann, H. e. a., Kobel, H., Pache, W., & Tscherter, H. (1976). Cyclosporin a and C. *European journal of applied microbiology and biotechnology*, *3*(2), 125-133.

Driggers, E. M., Hale, S. P., Lee, J., & Terrett, N. K. (2008). The exploration of macrocycles for drug discovery—an underexploited structural class. *Nature Reviews Drug Discovery*, *7*(7), 608-624.

Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic acids research*, *32*(5), 1792-1797.

El Tayar, N., Mark, A. E., Vallat, P., Brunne, R. M., Testa, B., & van Gunsteren, W. F. (1993).
Solvent-dependent conformation and hydrogen-bonding capacity of cyclosporin A:
evidence from partition coefficients and molecular dynamics simulations. *Journal of*
medicinal chemistry, *36*(24), 3757-3764.
Emsley, P., Lohkamp, B., Scott, W. G., & Cowtan, K. (2010). Features and development of
Coot. *Acta Crystallographica Section D: Biological Crystallography*, *66*(4), 486-501.
Fülöp, V., Böcskei, Z., & Polgár, L. (1998). Prolyl oligopeptidase: an unusual β-propeller
domain regulates proteolysis. *Cell*, *94*(2), 161-170.
Fülöp, V., Szeltner, Z., & Polgár, L. (2000). Catalysis of serine oligopeptidases is controlled by
a gating filter mechanism. *EMBO reports*, *1*(3), 277-281.
Gao, X., Haynes, S. W., Ames, B. D., Wang, P., Vien, L. P., Walsh, C. T., & Tang, Y. (2012).
Cyclization of fungal nonribosomal peptides by a terminal condensation-like domain.
Nature chemical biology, *8*(10), 823-830.
Giordanetto, F., & Kihlberg, J. (2014). Macrocyclic drugs and clinical candidates: what can
medicinal chemists learn from their properties? *Journal of medicinal chemistry*, *57*(2),
278-295.
Joosten, R. P., Joosten, K., Murshudov, G. N., & Perrakis, A. (2012). PDB_REDO: constructive
validation, more than just looking for errors. *Acta Crystallographica Section D:*
Biological Crystallography, *68*(4), 484-496.
Jurrus, E., Engel, D., Star, K., Monson, K., Brandi, J., Felberg, L. E., Brookes, D. H., Wilson, L.,
Chen, J., & Liles, K. (2018). Improvements to the APBS biomolecular solvation
software suite. *Protein Science*, *27*(1), 112-128.
Kaushik, S., Etchebest, C., & Sowdhamini, R. (2014). Decoding the structural events in
substrate-gating mechanism of eukaryotic prolyl oligopeptidase using normal mode
analysis and molecular dynamics simulations. *Proteins: Structure, Function, and*
Bioinformatics, *82*(7), 1428-1443.
Koehnke, J., Bent, A., Houssen, W. E., Zollman, D., Morawitz, F., Shirran, S., Vendome, J.,
Nneoyiegbe, A. F., Trembleau, L., & Botting, C. H. (2012). The mechanism of
patellamide macrocyclization revealed by the characterization of the PatG
macrocyclase domain. *Nature structural & molecular biology*, *19*(8), 767-772.
Koehnke, J., Bent, A. F., Houssen, W. E., Mann, G., Jaspars, M., & Naismith, J. H. (2014). The
structural biology of patellamide biosynthesis. *Current opinion in structural biology*,
29, 112-121.
Koehnke, J., Naismith, J., & Van der Donk, W. A. (2017). *Cyclic Peptides: From Bioorganic*
Synthesis to Applications (Vol. 6). Royal Society of Chemistry.
Kohli, R. M., Trauger, J. W., Schwarzer, D., Marahiel, M. A., & Walsh, C. T. (2001). Generality
of peptide cyclization catalyzed by isolated thioesterase domains of nonribosomal
peptide synthetases. *Biochemistry*, *40*(24), 7099-7108.
Laskowski, R. A., & Swindells, M. B. (2011). LigPlot+: multiple ligand-protein interaction
diagrams for drug discovery. *J Chem Inf Model*, *51*(10), 2778-2786.
https://doi.org/10.1021/ci200227u
Lawen, A., & Zocher, R. (1990). Cyclosporin synthetase. The most complex peptide
synthesizing multienzyme polypeptide so far described. *Journal of Biological*
Chemistry, *265*(19), 11355-11360.
Lee, H., & van der Donk, W. A. (2022). Macrocyclization and Backbone Modification in RiPP
Biosynthesis. *Annual Review of Biochemistry*, *91*.

Li, M., Chen, C., Davies, D. R., & Chiu, T. K. (2010). Induced-fit mechanism for prolyl
endopeptidase. *Journal of Biological Chemistry*, *285*(28), 21487-21495.

Liebschner, D., Afonine, P. V., Baker, M. L., Bunkóczi, G., Chen, V. B., Croll, T. I., Hintze, B.,
Hung, L.-W., Jain, S., & McCoy, A. J. (2019). Macromolecular structure determination
using X-rays, neutrons and electrons: recent developments in Phenix. *Acta*
Crystallographica Section D: Structural Biology, *75*(10), 861-877.

Lipinski, C. A., Lombardo, F., Dominy, B. W., & Feeney, P. J. (2001). Experimental and
computational approaches to estimate solubility and permeability in drug discovery
and development. *Advanced drug delivery reviews*, *46*, 3-26.

Liu, H., & Naismith, J. H. (2008). An efficient one-step site-directed deletion, insertion, single
and multiple-site plasmid mutagenesis protocol. *BMC biotechnology*, *8*(1), 1-10.

Lovell, S. C., Davis, I. W., Arendall III, W. B., De Bakker, P. I., Word, J. M., Prisant, M. G.,
Richardson, J. S., & Richardson, D. C. (2003). Structure validation by Cα geometry: φ,
ψ and Cβ deviation. *Proteins: Structure, Function, and Bioinformatics*, *50*(3), 437-450.

Ludewig, H., Czekster, C. M., Oueis, E., Munday, E. S., Arshad, M., Synowsky, S. A., Bent, A. F.,
& Naismith, J. H. (2018). Characterization of the fast and promiscuous macrocyclase
from plant PCY1 enables the use of simple substrates. *ACS chemical biology*, *13*(3),
801-811.

Luo, H., Hong, S.-Y., Sgambelluri, R. M., Angelos, E., Li, X., & Walton, J. D. (2014). Peptide
macrocyclization catalyzed by a prolyl oligopeptidase involved in α-amanitin
biosynthesis. *Chemistry & biology*, *21*(12), 1610-1617.

Marsault, E., & Peterson, M. L. (2011). Macrocycles are great cycles: applications,
opportunities, and challenges of synthetic macrocycles in drug discovery. *Journal of*
medicinal chemistry, *54*(7), 1961-2004.

Marti-Centelles, V., Pandey, M. D., Burguete, M. I., & Luis, S. V. (2015). Macrocyclization
reactions: the importance of conformational, configurational, and template-induced
preorganization. *Chemical reviews*, *115*(16), 8736-8834.

Matabaro, E., Kaspar, H., Dahlin, P., Bader, D. L., Murar, C. E., Staubli, F., Field, C. M., Bode, J.
W., & Künzler, M. (2021). Identification, heterologous production and bioactivity of
lentinulin A and dendrothelin A, two natural variants of backbone N-methylated
peptide macrocycle omphalotin A. *Scientific reports*, *11*(1), 1-12.

Matabaro, E., Song, H., Chepkirui, C., Kaspar, H., Witte, L., Naismith, J. H., Freeman, M. F., &
Künzler, M. (2021). Enzyme-mediated backbone N-methylation in ribosomally
encoded peptides. In *Methods in Enzymology* (Vol. 656, pp. 429-458). Elsevier.

Mayer, A., Anke, H., & Sterner, O. (1997). Omphalotin, a new cyclic peptide with potent
nematicidal activity from Omphalotus olearius I. Fermentation and biological activity.
Natural Product Letters, *10*(1), 25-32.

McCoy, A. J., Grosse-Kunstleve, R. W., Adams, P. D., Winn, M. D., Storoni, L. C., & Read, R. J.
(2007). Phaser crystallographic software. *Journal of applied crystallography*, *40*(4),
658-674.

McIntosh, J. A., Robertson, C. R., Agarwal, V., Nair, S. K., Bulaj, G. W., & Schmidt, E. W.
(2010). Circular logic: nonribosomal peptide-like macrocyclization with a ribosomal
peptide catalyst. *Journal of the American Chemical Society*, *132*(44), 15499-15501.

McNicholas, S., Potterton, E., Wilson, K., & Noble, M. (2011). Presenting your structures: the
CCP4mg molecular-graphics software. *Acta Crystallographica Section D: Biological*
Crystallography, *67*(4), 386-394.

Montalbán-López, M., Scott, T. A., Ramesh, S., Rahman, I. R., Van Heel, A. J., Viel, J. H.,
Bandarian, V., Dittmann, E., Genilloud, O., & Goto, Y. (2021). New developments in
RiPP discovery, enzymology and engineering. *Natural product reports*, *38*(1), 130-239.
Murshudov, G. N., Vagin, A. A., & Dodson, E. J. (1997). Refinement of macromolecular
structures by the maximum-likelihood method. *Acta Crystallographica Section D:*
Biological Crystallography, *53*(3), 240-255.
Nagatsu, T., Hino, M., Fuyamada, H., Hayakawa, T., Sakakibara, S., Nakagawa, Y., &
Takemoto, T. (1976). New chromogenic substrates for X-prolyl dipeptidyl-
aminopeptidase. *Analytical biochemistry*, *74*(2), 466-476.
Nguyen, G. K., Kam, A., Loo, S., Jansson, A. E., Pan, L. X., & Tam, J. P. (2015). Butelase 1: a
versatile ligase for peptide and protein macrocyclization. *Journal of the American*
Chemical Society, *137*(49), 15398-15401.
Nguyen, G. K., Wang, S., Qiu, Y., Hemu, X., Lian, Y., & Tam, J. P. (2014). Butelase 1 is an Asx-
specific ligase enabling peptide macrocyclization and synthesis. *Nature chemical*
biology, *10*(9), 732-738.
Olarte, R. A., Menke, J., Zhang, Y., Sullivan, S., Slot, J. C., Huang, Y., Badalamenti, J. P.,
Quandt, A. C., Spatafora, J. W., & Bushley, K. E. (2019). Chromosome rearrangements
shape the diversification of secondary metabolism in the cyclosporin producing
fungus Tolypocladium inflatum. *BMC Genomics*, *20*(1), 120.
https://doi.org/10.1186/s12864-018-5399-x
Oueis, E., Jaspars, M., Westwood, N. J., & Naismith, J. H. (2016). Enzymatic Macrocyclization
of 1, 2, 3-Triazole Peptide Mimetics. *Angewandte Chemie International Edition*,
55(19), 5842-5845.
Oueis, E., Nardone, B., Jaspars, M., Westwood, N. J., & Naismith, J. H. (2017). Synthesis of
hybrid cyclopeptides through enzymatic macrocyclization. *ChemistryOpen*, *6*(1), 11-
14.
Oueis, E., Stevenson, H., Jaspars, M., Westwood, N., & Naismith, J. (2017). Bypassing the
proline/thiazoline requirement of the macrocyclase PatG. *Chemical Communications*,
53(91), 12274-12277.
Ovadia, O., Greenberg, S., Chatterjee, J., Laufer, B., Opperer, F., Kessler, H., Gilon, C., &
Hoffman, A. (2011). The effect of multiple N-methylation on intestinal permeability
of cyclic hexapeptides. *Molecular pharmaceutics*, *8*(2), 479-487.
Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E. C., &
Ferrin, T. E. (2004). UCSF Chimera—a visualization system for exploratory research
and analysis. *Journal of computational chemistry*, *25*(13), 1605-1612.
Quijano, M. R., Zach, C., Miller, F. S., Lee, A. R., Imani, A. S., Künzler, M., & Freeman, M. F.
(2019). Distinct autocatalytic α-N-methylating precursors expand the borosin RiPP
family of peptide natural products. *Journal of the American Chemical Society*,
141(24), 9637-9644.
Ramachandran, G. N., Ramakrishnan, C., & Sasisekharan, V. (1963). Stereochemistry of
polypeptide chain configurations. *Journal of Molecular Biology 7*, 95-99.
Ramm, S., Krawczyk, B., Mühlenweg, A., Poch, A., Mösker, E., & Süssmuth, R. D. (2017). A
self-sacrificing N-methyltransferase is the precursor of the fungal natural product
omphalotin. *Angewandte Chemie International Edition*, *56*(33), 9994-9997.
Robert, X., & Gouet, P. (2014). Deciphering key features in protein structures with the new
ENDscript server. *Nucleic acids research*, *42*(W1), W320-W324.

Sgambelluri, R. M., Smith, M. O., & Walton, J. D. (2018). Versatility of prolyl oligopeptidase B
in peptide macrocyclization. *ACS synthetic biology*, *7*(1), 145-152.
Shan, L., Mathews, I. I., & Khosla, C. (2005). Structural and mechanistic analysis of two prolyl
endopeptidases: role of interdomain dynamics in catalysis and specificity.
Proceedings of the National Academy of Sciences, *102*(10), 3599-3604.
Simon, R. J., Kania, R. S., Zuckermann, R. N., Huebner, V. D., Jewell, D. A., Banville, S., Ng, S.,
Wang, L., Rosenberg, S., & Marlowe, C. K. (1992). Peptoids: a modular approach to
drug discovery. *Proceedings of the National Academy of Sciences*, *89*(20), 9367-9371.
Song, H., Burton, A. J., Shirran, S. L., Fahrig-Kamarauskaitė, J., Kaspar, H., Muir, T. W.,
Künzler, M., & Naismith, J. H. (2021). Engineering of a Peptide α‐N‐
Methyltransferase to Methylate Non‐Proteinogenic Amino Acids. *Angewandte*
Chemie International Edition, *60*(26), 14319-14323.
Song, H., Fahrig-Kamarauskaitè, J. r., Matabaro, E., Kaspar, H., Shirran, S. L., Zach, C., Pace,
A., Stefanov, B.-A., Naismith, J. H., & Künzler, M. (2020). Substrate plasticity of a
fungal peptide α-N-methyltransferase. *ACS chemical biology*, *15*(7), 1901-1912.
Song, H., & Naismith, J. H. (2020). Enzymatic methylation of the amide bond. *Current*
opinion in structural biology, *65*, 79-88.
Song, H., Van Der Velden, N. S., Shiran, S. L., Bleiziffer, P., Zach, C., Sieber, R., Imani, A. S.,
Krausbeck, F., Aebi, M., & Freeman, M. F. (2018). A molecular mechanism for the
enzymatic methylation of nitrogen atoms within peptide bonds. *Science advances*,
4(8), eaat2720.
Surade, S., & Blundell, T. L. (2012). Structural biology and drug discovery of difficult targets:
the limits of ligandability. *Chemistry & biology*, *19*(1), 42-50.
Thoma, R., Löffler, B., Stihle, M., Huber, W., Ruf, A., & Hennig, M. (2003). Structural basis of
proline-specific exopeptidase activity as observed in human dipeptidyl peptidase-IV.
Structure, *11*(8), 947-959.
van der Velden, N. S., Kälin, N., Helf, M. J., Piel, J., Freeman, M. F., & Künzler, M. (2017).
Autocatalytic backbone N-methylation in a family of ribosomal peptide natural
products. *Nature chemical biology*, *13*(8), 833-835.
Winter, G. (2010). xia2: an expert system for macromolecular crystallography data reduction.
Journal of applied crystallography, *43*(1), 186-190.
Winter, G., Waterman, D. G., Parkhurst, J. M., Brewster, A. S., Gildea, R. J., Gerstel, M.,
Fuentes-Montero, L., Vollmar, M., Michels-Clark, T., & Young, I. D. (2018). DIALS:
implementation and evaluation of a new integration package. *Acta Crystallographica*
Section D, *74*(2), 85-97.
Yoshimoto, T., Kawahara, K., Matsubara, F., Kado, K., & Tsuru, D. (1985). Comparison of
inhibitory effects of prolinal-containing peptide derivatives on prolyl endopeptidases
from bovine brain and Flavobacterium. *Journal of biochemistry*, *98*(4), 975-979.
Yu, X., & Sun, D. (2013). Macrocyclic drugs and synthetic methodologies toward macrocycles.
Molecules, *18*(6), 6230-6268.
Zhang, J., Liu, N., Cacho, R. A., Gong, Z., Liu, Z., Qin, W., Tang, C., Tang, Y., & Zhou, J. (2016).
Structural basis of nonribosomal peptide macrocyclization in fungi. *Nature chemical*
biology, *12*(12), 1001-1003.
1166

omphalotin A

dendrothelin A

lentinulin A

Abbreviation	OphP peptide substrate	Produced peptide macrocycle(s)
Oph-30mer	ESGEEASQNGFPWVIVVGVIGVIGSVMSTE	None
Oph-24mer	SQNGFPWVIVVGVIGVIGSVMSTE	SQNGFPWVIVVGVIGVIG
Oph-21mer	GFPWVIVVGVIGVIGSVMSTE	GFPWVIVVGVIGVIG
Oph-18mer^Tr	GFPWVIVVGVIGSVMSTE	GFPWVIVVGVIG
Oph-15mer	GFPWVIVVGVIGVIG	GFPWVIVVGVIG
Oph-18mer^NL	WVIVVGVIGVIGSVMSTE	WVIVVGVIGVIG (omphalotin A)
Oph-15mer^Tr	WVIVVGVIGSVMSTE	None
Oph-12mer	WVIVVGVIGVIG	None
Led-21mer	GFPWIIVVGVVGVIGSVVSSA	GFPWIIVVGVVGVIG
Dbi-21mer	GFPWVIVTGIVGVIGSVVSSA	GFPWVIVTGIVGVIG
Led-18mer	WIIVVGVVGVVGSVVSSA	WIIVVGVVGVVG (lentinulin A)
Dbi-18mer	WVIVTGIVGVIGSVVSSA	WVIVTGIVGVIG (dendrothelin A)

b

SQNGFPWVIVVGVIG
[M+H]²⁺ 826.4793
Oph-24mer + OphP

SQNGFPWVIVVGVIGVIG
[M+H]²⁺ 975.5828
Oph-24mer + OphP

Time (min)

c

GFPWVIVVGVIG
[M+H]⁺ 1322.8244
Oph-21mer + OphP

GFPWVIVVGVIGVIG
[M+H]⁺ 1620.0292
Oph-21mer + OphP

Time (min)

d

GFPWVIVVGVIG

UV absorbance at 280 nm

Oph-15mer
Oph-18mer^Tr

Oph-15mer
Oph-15mer + OphP
Oph-18mer^Tr
Oph-18mer^Tr + OphP

Time (min)

e

[M+H]⁺ 1318.8887 **omphalotin A**
[M+NH₄]⁺ 1335.9136
[M+Na]⁺ 1340.8629

[M+H]⁺ 1304.8751 **lentinulin A**
[M+NH₄]⁺ 1321.9021
[M+Na]⁺ 1326.8561

[M+H]⁺ 1320.8694 **dendrothelin A**
[M+NH4]⁺ 1337.8964
[M+Na]⁺ 1342.8508

m/z

a

Oph-18mer^{NL}

Oph-18mer^{NL}
+ OphP

3Me: WVIVVGVIGVIGSVMSTE ❌
4Me: WVIVVGVIGVIGSVMSTE ✅
5Me: WVIVVGIGVIGSVMSTE ❌
6Me: WVIVVGVIGVIGSVMSTE ❌
7Me: WVIVVGVIGVIGSVMSTE ✅
8Me: WVIVVGVIGVIGSVMSTE ✅
9Me: WVIVVGVIGVIGSVMSTE ✅
10Me: WVIVVGVIGVIGSVMSTE ✅
11Me: WVIVVGVIGVIGSVMSTE ✅

Oph-15mer

Oph-15mer
+ OphP

6Me: GFPWVIVVGVIGVIG ❌
7Me: GFPWVIVVGVIGVIG ✅

m/z

b

Oph-15mer substrates

GFPWVIVVGVIGVIG (7Me)
GFPWVIVVGVIGVIG (6Me)

Ratio of Oph-15mer substrates

c

Oph-15mer cyclic products

GFPWVIVVGVIG (7Me)
GFPWVIVVGVIG (6Me)

Ratio of Oph-15mer cyclic products

a

Oph-18mer^{NL}

b

$^{me}I8$
$^{me}G9$
$V10$
$^{me}I11$
$^{me}G12$
$S13$
$^{me}V14$

Oph-18mer^{NL}

c

Trp621
Phe502
$^{me}I11$
Ala284
$^{me}G12$
$S13$
His701
Ser281
$^{me}G9$
$V10$
$\underline{S580A}$
$^{me}V14$
$^{me}I8$
Phe617
Gly579
Y497
Arg316
His286
Arg667
Tyr449

d

Tyr625
Ile606
Asn581
Trp621
Phe502
Val668
$^{me}G12$

e

GFPWVIVVGVIG
Oph-15mer

GFPWVIVVGVIG

Oph-15mer + S580A

Oph-15mer + I606A

Oph-15mer + W621A

Oph-15mer + OphP

Oph-15mer

mAU at 280 nm

2.0   3.0   4.0   5.0   6.0

Time (h)