Discovering non-additive heritability using additive GWAS summary statistics

Samuel Pattillo Smith[1-4,*], Gregory Darnell[1,5,*], Dana Udwin[6], Julian Stamp[1], Arbel Harpak[3,4], Sohini Ramachandran[1,2,7,8,§], and Lorin Crawford[1,6,8,§,†]

1 Center for Computational Molecular Biology, Brown University, Providence, RI, USA

2 Department of Ecology and Evolutionary Biology, Brown University, Providence, RI, USA

3 Department of Integrative Biology, University of Texas at Austin, Austin, TX, USA

4 Department of Population Health, University of Texas at Austin, Austin, TX, USA

5 Institute for Computational and Experimental Research in Mathematics, Brown University, Providence, RI, USA

6 Department of Biostatistics, Brown University, Providence, RI, USA

7 Data Science Institute, Brown University, Providence, RI, USA

8 Microsoft Research, Cambridge, MA, USA

* Authors Contributed Equally

§ Authors Contributed Equally

† Corresponding E-mail: lcrawford@microsoft.com

Abstract

LD score regression (`LDSC`) is a method to estimate narrow-sense heritability from genome-wide association study (GWAS) summary statistics alone, making it a fast and popular approach. In this work, we present interaction-LD score (`i-LDSC`) regression: an extension of the original `LDSC` framework that accounts for interactions between genetic variants. By studying a wide range of generative models in simulations, and by re-analyzing 25 well-studied quantitative phenotypes from 349,468 individuals in the UK Biobank and up to 159,095 individuals in BioBank Japan, we show that the inclusion of a *cis*-interaction score (i.e., interactions between a focal variant and proximal variants) recovers genetic variance that is not captured by `LDSC`. For each of the 25 traits analyzed in the UK Biobank and BioBank Japan, `i-LDSC` detects

additional variation contributed by genetic interactions. The `i-LDSC` software and its application to these biobanks represent a step towards resolving further genetic contributions of sources of non-additive genetic effects to complex trait variation.

Introduction

Heritability is defined as the proportion of phenotypic trait variation that can be explained by genetic effects [1–3]. Until recently, studies of heritability in humans have been reliant on typically small sized family studies with known relatedness structures among individuals [4,5]. Due to advances in genomic sequencing and the steady development of statistical tools, it is now possible to obtain reliable heritability estimates from biobank-scale data sets of unrelated individuals [1,3,6,7]. Computational and privacy considerations with genome-wide association studies (GWAS) in these larger cohorts have motivated a recent trend to estimate heritability using summary statistics (i.e., estimated effect sizes and their corresponding standard errors). In the GWAS framework, additive effect sizes and standard errors for individual single nucleotide polymorphisms (SNPs) are estimated by regressing phenotype measurements onto the allele counts of each SNP independently. Through the application of this approach over the last two decades, it has become clear that many traits have a complex and polygenic basis—that is, hundreds to thousands of individual genetic loci across the genome often contribute to the genetic basis of variation in a single trait [8].

Many statistical methods have been developed to improve the estimation of heritability from GWAS summary statistics [1,3,9,10]. The most widely used of these approaches is linkage disequilibrium (LD) score regression and the corresponding `LDSC` software [1], which corrects for inflation in GWAS summary statistics by modeling the relationship between the variance of SNP-level effect sizes and the sum of correlation coefficients between focal SNPs and their genomic neighbors (i.e., the LD score of each variant). The formulation of the `LDSC` framework relies on the fact that the expected relationship between chi-square test statistics (i.e., the squared magnitude of GWAS allelic effect estimates) and LD scores holds when complex traits are generated under the infinitesimal (or polygenic) model which assumes: (i) all causal variants have the same expected contribution to phenotypic variation and (ii) causal variants are uniformly distributed along the genome. Initial simulations in Bulik-Sullivan et al. [1] showed that violations of these assumptions can be tolerated to a point, but begin to affect the estimation of narrow-sense heritability

once a certain proportion of variants have nonzero effects. Importantly, the estimand of the `LDSC` model is the proportion of phenotypic variance attributable to additive effects of genotyped SNPs. The main motivation behind the `LDSC` model is that, for polygenic traits, many marker SNPs tag nonzero effects. This may simply arise because some of these SNPS are in LD with causal variants[1] or because their statistical association is the product of a confounding factor such as population stratification.

As of late, there have been many efforts to build upon and improve the `LDSC` framework. For example, recent work has shown that it is possible to estimate the proportion of phenotypic variation explained by dominance effects[11] and local ancestry[12] using extensions of the `LDSC` model. One limitation of `LDSC` is that, in practice, it only uses the diagonal elements of the squared LD matrix in its formulation which, while computationally efficient, does not account for information about trait architecture that is captured by the off-diagonal elements. This tradeoff helps `LDSC` to scale genome-wide, but it has also been shown to lead to heritability estimates with large standard error[10,13,14]. Recently, newer approaches have attempted to reformulate the `LDSC` model by using the eigenvalues of the LD matrix to leverage more of the information present in the correlation structure between SNPs[3,10].

In this paper, we show that the LDSC framework can be extended to estimate greater proportions of genetic variance in complex traits (i.e., beyond the variance that is attributable to additive effects) when a subset of causal variants is involved in a gene-by-gene (G×G) interaction. Indeed, recent association mapping studies have shown that G×G interactions can drive heterogeneity of causal variant effect sizes[15]. Importantly, non-additive genetic effects have been proposed as one of the main factors that explains "missing" heritability—the proportion of heritability not explained by the additive effects of variants[16].

The key insight we highlight in this manuscript is that SNP-level GWAS summary statistics can provide evidence of non-additive genetic effects contributing to trait architecture if there is a nonzero correlation between individual-level genotypes and their statistical interactions. We present the "interaction-LD score" regression model or `i-LDSC`: an extension of the `LDSC` framework which recovers "missing" heritability by leveraging this "tagged" relationship between linear and nonlinear genetic effects. To validate the performance of `i-LDSC` in simulation studies, we focus on synthetic trait architectures that have been generated with contributions stemming from second-order and *cis*-acting statistical SNP-by-SNP interaction effects; however, note that the general concept underlying `i-LDSC` can easily be extended to other sources of non-additive genetic effects (e.g., gene-by-environment interactions). The main difference

⁸⁶ between `i-LDSC` and `LDSC` is that the `i-LDSC` model includes an additional set of "*cis*-interaction" LD

⁸⁷ scores in its regression model. These scores measure the amount of phenoytpic variation contributed by

⁸⁸ genetic interactions that can be explained by additive effects. In practice, these additional scores are

⁸⁹ efficient to compute and require nothing more than access to a representative pairwise LD map, same as

⁹⁰ the input required for LD score regression.

⁹¹ Through extensive simulations, we show that `i-LDSC` recovers substantial non-additive heritability

⁹² that is not captured by `LDSC` when genetic interactions are indeed present in the generative model for a

⁹³ given complex trait. More importantly, `i-LDSC` has a calibrated type I error rate and does not overesti-

⁹⁴ mate contributions of genetic interactions to trait variation in simulated data when only additive effects

⁹⁵ are present. While analyzing 25 complex traits in the UK Biobank and BioBank Japan, we illustrate

⁹⁶ that pairwise interactions are a source of "missing" heritability captured by additive GWAS summary

⁹⁷ statistics—suggesting that phenotypic variation due to non-additive genetic effects is more pervasive in

⁹⁸ human phenotypes than previously reported. Specifically, we find evidence of tagged genetic interac-

⁹⁹ tion effects contributing to heritability estimates in all of the 25 traits in the UK Biobank, and 23 of

¹⁰⁰ the 25 traits we analyzed in the BioBank Japan. We believe that `i-LDSC`, with our development of a

¹⁰¹ new *cis*-interaction score, represents a significant step towards resolving the true contribution of genetic

¹⁰² interactions.

Results

Overview of the interaction-LD score regression model

¹⁰⁵ Interaction-LD score regression (`i-LDSC`) is a statistical framework for estimating heritability (i.e., the

¹⁰⁶ proportion of trait variance attributable to genetic variance). Here, we will give an overview of the

¹⁰⁷ `i-LDSC` method and its corresponding software, as well as detail how its underlying model differs from

¹⁰⁸ that of `LDSC`[1]. We will assume that we are analyzing a GWAS dats set $\mathcal{D} = \{\mathbf{X}, \mathbf{y}\}$ where $\mathbf{X}$ is an $N \times J$

¹⁰⁹ matrix of genotypes with J denoting the number of SNPs (each of which is encoded as $\{0, 1, 2\}$ copies of

¹¹⁰ a reference allele at each locus j) and $\mathbf{y}$ is an N-dimensional vector of measurements of a quantitative

¹¹¹ trait. The `i-LDSC` framework only requires summary statistics of individual-level data: namely, marginal

¹¹² effect size estimates for each SNP $\widehat{\boldsymbol{\beta}}$ and a sample LD matrix $\mathbf{R}$ (which can be provided via reference

¹¹³ panel data).

114   We begin by considering the following generative linear model for complex traits

$$\mathbf{y} = b_0 + \mathbf{X}\boldsymbol{\beta} + \mathbf{W}\boldsymbol{\theta} + \boldsymbol{\varepsilon}, \qquad \boldsymbol{\varepsilon} \sim \mathcal{N}(\mathbf{0}, (1 - H^2)\mathbf{I}), \tag{1}$$

where b_0 is an intercept term; $\boldsymbol{\beta} = (\beta_1, \ldots, \beta_J)$ is a J-dimensional vector containing the true additive
effect sizes for an additional copy of the reference allele at each locus on $\mathbf{y}$; $\mathbf{W}$ is an $N \times M$ matrix
of (pairwise) *cis*-acting SNP-by-SNP statistical interactions between some subset of causal SNPs, where
columns of this matrix are assumed to be the Hadamard (element-wise) product between genotypic vectors
of the form $\mathbf{x}_j \circ \mathbf{x}_k$ for the j-th and k-th variants; $\boldsymbol{\theta} = (\theta_1, \ldots, \theta_M)$ is an M-dimensional vector containing
the interaction effect sizes; $\boldsymbol{\varepsilon}$ is a normally distributed error term with mean zero and variance scaled
according to the proportion of phenotypic variation not explained by genetic effects [17], which we will refer
to as the broad-sense heritability of the trait denoted by H^2; and $\mathbf{I}$ denotes an $N \times N$ identity matrix.
For convenience, we will assume that the genotype matrix (column-wise) and the trait of interest have
been mean-centered and standardized [18–20]. Lastly, we will let the intercept term b_0 be a fixed parameter
and we will assume that the effect sizes are each normally distributed with variances proportional to their
individual contributions to trait heritability [17,20–23]

$$\beta_j \sim \mathcal{N}(0, \varphi_\beta^2/J), \qquad \theta_m \sim \mathcal{N}(0, \varphi_\theta^2/M). \tag{2}$$

Effectively, we say that $\mathbb{V}[\mathbf{X}\boldsymbol{\beta}] = \varphi_\beta^2$ is the proportion of phenotypic variation contributed by additive SNP
effects under the generative model, while $\mathbb{V}[\mathbf{W}\boldsymbol{\theta}] = \varphi_\theta^2$ makes up the proportion of phenotypic variation
contributed by genetic interactions. While the appropriateness of treating genetic effects as random
variables in analytical derivations has been questioned [24], later, we will justify the theory presented here
with simulation results showing that `i-LDSC` accurately recovers non-additive genetic variance in Eq. (1)
under a broad range of conditions.

There are two key takeaways from the generative model specified above. First, Eq. (2) implies that
the additive and non-additive components in Eq. (1) are orthogonal to each other. In other words,
$\mathbb{E}[\boldsymbol{\beta}^\intercal \mathbf{X}^\intercal \mathbf{W}\boldsymbol{\theta}] = \mathbb{E}[\boldsymbol{\beta}^\intercal]\mathbf{X}^\intercal \mathbf{W}\mathbb{E}[\boldsymbol{\theta}] = \mathbf{0}$. This is important because it means that there is a unique par-
titioning of genetic variance when studying a trait of interest. The second key takeaway is that the
genotype matrix $\mathbf{X}$ and the matrix of genetic interactions $\mathbf{W}$ themselves are correlated despite being
linearly independent (see Material and Methods). This property stems from the fact that the pairwise

interaction between two SNPs is encoded as the Hadamard product of two genotypic vectors in the form $\mathbf{w}_m = \mathbf{x}_j \circ \mathbf{x}_k$ (which is a nonlinear function of the genotypes).

A central objective in GWAS studies is to infer how much phenotypic variation can be explained by genetic effects. To achieve that objective, a key consideration involves incorporating the possibility of non-additive sources of genetic variation to be explained by additive effect size estimates obtained from GWAS analyses[25]. If we assume that the genotype and interaction matrices are correlated, then $\mathbf{X}$ and $\mathbf{W}$ are not completely orthogonal (i.e., such that $\mathbf{X}^\mathsf{T}\mathbf{W} \neq \mathbf{0}$) and the following relationship between the moment matrix $\mathbf{X}^\mathsf{T}\mathbf{y}$, the observed marginal GWAS summary statistics $\widehat{\boldsymbol{\beta}}$, and the true coefficient values $\boldsymbol{\beta}$ from the generative model in Eq. (1) holds in expectation (see Materials and Methods)

$$\mathbb{E}[\mathbf{X}^\mathsf{T}\mathbf{y}] = (\mathbf{X}^\mathsf{T}\mathbf{X})\boldsymbol{\beta} + (\mathbf{X}^\mathsf{T}\mathbf{W})\boldsymbol{\theta} \qquad \overset{\approx}{\Longleftrightarrow} \qquad \mathbb{E}[\widehat{\boldsymbol{\beta}}] = \mathbf{R}\boldsymbol{\beta} + \mathbf{V}\boldsymbol{\theta} \tag{3}$$

where $\mathbf{R}$ is a sample estimate of the LD matrix, and $\mathbf{V}$ represents a sample estimate of the correlation between the individual-level genotypes $\mathbf{X}$ and the span of genetic interactions between causal SNPs in $\mathbf{W}$. Intuitively, the term $\mathbf{V}\boldsymbol{\theta}$ can be interpreted as the subset of pairwise interaction effects that are tagged by the additive effect estimates from the GWAS study. Note that, when (i) non-additive genetic effects do not contribute to the overall architecture of a trait (i.e., such that $\boldsymbol{\theta} = \mathbf{0}$) or ($ii$) the genotype and interaction matrices $\mathbf{X}$ and $\mathbf{W}$ are uncorrelated, the equation above simplifies to a relationship between LD and summary statistics that is assumed in many GWAS studies and methods[26–32].

The goal of `i-LDSC` is to increase estimates of genetic variance by accounting for sources of non-additive genetic effects that can be explained by additive GWAS summary statistics. To do this, we extend the LD score regression framework and the corresponding `LDSC` software[17]. Here, according to Eq. (3), we note that $\widehat{\boldsymbol{\beta}} \sim \mathcal{N}(\mathbf{R}\boldsymbol{\beta} + \mathbf{V}\boldsymbol{\theta}, \lambda\mathbf{R})$ where λ is a scale variance term due to uncontrolled confounding effects[10,33]. Next, we condition on $\boldsymbol{\Theta} = (\boldsymbol{\beta}, \boldsymbol{\theta})$ and take the expectation of chi-square statistics $\boldsymbol{\chi}^2 = N\widehat{\boldsymbol{\beta}}\widehat{\boldsymbol{\beta}}^\mathsf{T}$ to yield

$$\begin{aligned}
\mathbb{E}[\widehat{\boldsymbol{\beta}}\widehat{\boldsymbol{\beta}}^\mathsf{T}] &= \mathbb{E}\left[\mathbb{E}\left[\widehat{\boldsymbol{\beta}}\widehat{\boldsymbol{\beta}}^\mathsf{T} \,|\, \boldsymbol{\Theta}\right]\right] = \mathbb{E}\left[\mathbb{V}\left[\widehat{\boldsymbol{\beta}} \,|\, \boldsymbol{\Theta}\right] + \mathbb{E}\left[\widehat{\boldsymbol{\beta}} \,|\, \boldsymbol{\Theta}\right]\mathbb{E}\left[\widehat{\boldsymbol{\beta}} \,|\, \boldsymbol{\Theta}\right]^\mathsf{T}\right] \\
&= \mathbb{E}\left[\lambda\mathbf{R} + (\mathbf{R}\boldsymbol{\beta} + \mathbf{V}\boldsymbol{\theta})(\mathbf{R}\boldsymbol{\beta} + \mathbf{V}\boldsymbol{\theta})^\mathsf{T}\right] \\
&= \mathbb{E}\left[\lambda\mathbf{R} + \mathbf{R}\boldsymbol{\beta}\boldsymbol{\beta}^\mathsf{T}\mathbf{R} + 2\mathbf{R}\boldsymbol{\beta}\boldsymbol{\theta}^\mathsf{T}\mathbf{V}^\mathsf{T} + \mathbf{V}\boldsymbol{\theta}\boldsymbol{\theta}^\mathsf{T}\mathbf{V}^\mathsf{T}\right] \\
&= \lambda\mathbf{R} + \left(\frac{\varphi_\beta^2}{J}\right)\mathbf{R}^2 + \left(\frac{\varphi_\theta^2}{M}\right)\mathbf{V}^2.
\end{aligned} \tag{4}$$

164 We define $\ell_j = \sum_k r_{jk}^2$ as the LD score for the additive effect of the j-th variant[17], and $f_j = \sum_m v_{jm}^2$

165 represents the "*cis*-interaction" LD score which encodes the pairwise interaction between the j-th variant

166 and all other variants within a genomic window that is a pre-specified number of SNPs wide[23], respec-

167 tively. By considering only the diagonal elements of LD matrix in the first term, similar to the original

168 LDSC approach[10,17], we get the following simplified regression model

$$\mathbb{E}[\boldsymbol{\chi}^2] \propto \mathbf{1} + \boldsymbol{\ell}\tau + \boldsymbol{f}\vartheta \tag{5}$$

where $\boldsymbol{\chi}^2 = (\chi_1^2, \ldots, \chi_J^2)$ is a J-dimensional vector of chi-square summary statistics, and $\boldsymbol{\ell} = (\ell_1, \ldots, \ell_J)$

and $\boldsymbol{f} = (f_1, \ldots, f_J)$ are J-dimensional vectors of additive and *cis*-interaction LD scores, respectively.

Furthermore, we define the variance components $\tau = N\varphi_\beta^2/J$ and $\vartheta = N\varphi_\theta^2/M$ as the additive and

non-additive regression coefficients of the model, and $\mathbf{1}$ is the intercept meant to model the bias factor

due to uncontrolled confounding effects (e.g., cryptic relatedness structure). In practice, we efficiently

compute the *cis*-interaction LD scores by considering only a subset of interactions between each j-th

focal SNP and SNPs within a *cis*-proximal window around the j-th SNP. In our validation studies and

applications, we base the width of this window on the observation that LD decays outside of a window

of 1 centimorgan (cM); therefore, SNPs outside the 1 cM window centered on the j-th SNP will not

significantly contribute to its LD scores. Note that the width of this window can be relaxed in the

i-LDSC software when appropriate. We fit the i-LDSC model using weighted least squares to estimate

regression parameters and derive P-values for identifying traits that have significant statistical evidence

of tagged *cis*-interaction effects by testing the null hypothesis $H_0 : \vartheta = 0$. Importantly, under the null

model of a trait being generated by only additive effects, the i-LDSC model in Eq. (5) reduces to an

infinitesimal model[34] or, in the case some variants have no effect on the trait, a polygenic model.

Lastly, we want to note the empirical observation that the additive ($\boldsymbol{\ell}$) and interaction ($\boldsymbol{f}$) LD scores

are lowly correlated. This is important because it indicates that the presence of *cis*-interaction LD scores

in the model specified in Eq. (5) has little-to-no influence over the estimate for the additive coefficient

τ. Instead, the inclusion of $\boldsymbol{f}$ creates a multivariate model that can identify the proportion of variance

explained by both additive and non-additive effects in summary statistics. In other words, we can

interpret $\widehat{\vartheta}$ as an estimate of the phenotypic variation explained by tagged *cis*-acting interaction effects.

The concept of additive genetic effects partially explaining non-additive variation has also described in

various studies from quantitative genetics [25,35,36]. Under Hardy-Weinberg equilibrium, it can be shown

that the additive variance explained by J SNPs takes on the following form (Materials and Methods) [37]

$$\sigma_A^2 = \sum_{j=1}^{J} 2p_j(1-p_j) \left[\beta_j + 2\sum_{k \neq j}^{J} p_k \theta_{jk} \right]^2 . \tag{6}$$

The expression for the additive variance σ_A^2 in Eq. (6) is important because it represents the theoretical

upper bound on the proportion of total phenotypic variance that can be recovered from GWAS summary

statistics using the i-LDSC framework. As a result, we use the sum of coefficient estimates $\widehat{\tau} + \widehat{\vartheta} \leq \sigma_A^2$

to construct i-LDSC heritability estimates. A full derivation of the *cis*-interaction regression framework

and details about its corresponding implementation in our software i-LDSC can be found in Materials

and Methods.

Detection of tagged pairwise interaction effects using i-LDSC in simulations

We illustrate the power of i-LDSC across different genetic trait architectures via extensive simulation

studies (Materials and Methods). We generate synthetic phenotypes using real genome-wide genotype

data from individuals of self-identified European ancestry in the UK Biobank. To do so, we first assume

that traits have a polygenic architecture where all SNPs have a nonzero additive effect. Next, we randomly

select a set of causal *cis*-interaction variants and divide them into two interacting groups (Materials and

Methods). One may interpret the SNPs in group #1 as being the "hubs" in an interaction map [23]; while,

SNPs in group #2 are selected to be variants within some kilobase (kb) window around each SNP in

group #1. We assume a wide range of simulation scenarios by varying the following parameters:

• heritability: $H^2 = 0.3$ and 0.6;

• proportion of phenotypic variation that is generated by additive effects: $\rho = 0.5$, 0.8, and 1;

• percentage of SNPs selected to be in group #1: 1%, 5%, and 10%;

• genomic window used to assign SNPs to group #2: ± 10 and ± 100 kb.

We also varied the correlation between SNP effect size and minor allele frequency (MAF) (as discussed

in Schoech et al. [38]). All results presented in this section are based on 100 different simulated phenotypes

for each parameter combination.

₂₁₈   Figure 1 demonstrates that `i-LDSC` robustly detects significant tagged non-additive genetic variance,

₂₁₉ regardless of the total number of causal interactions genome-wide. Instead, the power of `i-LDSC` depends

₂₂₀ on the proportion of phenotypic variation that is generated by additive versus interaction effects (ρ),

₂₂₁ and its power tends to scale with the window size used to compute the *cis*-interaction LD scores (see

₂₂₂ Materials and Methods). `i-LDSC` shows a similar performance for detecting tagged *cis*-interaction effects

₂₂₃ when the effect sizes of causal SNPs depend on their minor allele frequency and when we varied the

₂₂₄ number of SNPs assigned to be in group #2 within 10 kb and 100kb windows, respectively (Figure 1 –

₂₂₅ figure supplement 1-5).

₂₂₆   Importantly, `i-LDSC` does not falsely identify putative non-additive genetic effects in GWAS summary

₂₂₇ statistics when the synthetic phenotype was generated by only additive effects ($\rho = 1$). Figure 2 illustrates

₂₂₈ the performance of `i-LDSC` under the null hypothesis $H_0 : \vartheta = 0$, with the type I error rates for different

₂₂₉ estimation window sizes of the *cis*-interaction LD scores highlighted in panel A. Here, we also show

₂₃₀ that, when no genetic interaction effects are present, `i-LDSC` unbiasedly estimates the *cis*-interaction

₂₃₁ coefficient in the regression model to be $\widehat{\vartheta} = 0$ (Figure 2B), robustly estimates the heritability (Figure 2C),

₂₃₂ and provides well-calibrated P-values when assessed over many traits (Figure 2D). This behavior is

₂₃₃ consistent across different MAF-dependent effect size distributions, and P-value calibration is not sensitive

₂₃₄ to misspecification of the estimation windows used to generate the *cis*-interaction LD scores (Figure 2 –

₂₃₅ figure supplement 1-2).

₂₃₆   One of the innovations that `i-LDSC` offers over the traditional `LDSC` framework is increased heritability

₂₃₇ estimates after the identification of non-additive genetic effects that are tagged by GWAS summary

₂₃₈ statistics. Here, we applied both methods to the same set of simulations in order to understand how

₂₃₉ `LDSC` behaves for traits generated with *cis*-interaction effects. Figure 3 depicts boxplots of the heritability

₂₄₀ estimates for each approach and shows that, across an array of different synthetic phenotype architectures,

₂₄₁ `LDSC` captures less of phenotypic variance explained by all genetic effects. It is important to note that

₂₄₂ `i-LDSC` can yield upwardly biased heritability estimates when the *cis*-interaction scores are computed

₂₄₃ over genomic window sizes that are too small; however, these estimates become more accurate for larger

₂₄₄ window size choices (Figure 3 – figure supplement 1). In contrast to `LDSC`, which aims to capture

₂₄₅ phenotypic variance attributable to the additive effects of genotyped SNPs, `i-LDSC` accurately partitions

₂₄₆ genetic effects into additive versus *cis*-interacting components, which in turn generally leads the ability

₂₄₇ of `i-LDSC` to capture more genetic variance. The mean absolute error between the true generative

heritability and heritability estimates produced by `i-LDSC` and `LDSC` are shown in Supplementary Files 1 and 2, respectively. Generally, the error in heritability estimates is higher for `LDSC` than it is for `i-LDSC` across each of the scenarios that we consider.

Next, we perform an additional set of simulations where we explore other common generative models for complex trait architecture that involve non-additive genetic effects. Specifically, we compare heritability estimates from `LDSC` and `i-LDSC` in the presence of additive effects, *cis*-acting interactions, and a third source of genetic variance stemming from either gene-by-environment (G×E) or or gene-by-ancestry (G×Ancestry) effects. Details on how these components were generated can be found in Materials and Methods. In general, `i-LDSC` underestimates overall heritability when additive effects and *cis*-acting interactions are present alongside G×E (Figure 3 – figure supplement 2) and/or G×Ancestry effects when PCs are included as covariates (Figure 3 – figure supplement 3). Notably, when PCs are not included to correct for residual stratification, both `LDSC` and `i-LDSC` can yield unbounded heritability estimates greater than 1 (Figure 3 – figure supplement 4). Also interestingly, when we omit *cis*-interactions from the generative model (i.e., the genetic architecture of simulated traits is only made up of additive and G×E or G×Ancestry effects), `i-LDSC` will still estimate a nonzero genetic variance component with the *cis*-interaction LD scores (Figure 3 – figure supplement 5-7). Collectively, these results empirically show the important point that *cis*-interaction scores are not enough to recover missing genetic variation for all types of trait architectures; however, they are helpful in recovering phenotypic variation explained by statistical interaction effects. Recall that the linear relationship between (expected) χ^2 test statistics and LD scores proposed by the `LDSC` framework holds when complex traits are generated under the polygenic model where all causal variants have the same expected contribution to phenotypic variation. When *cis*-interactions affect genetic architecture (e.g., in our earlier simulations in Figure 3), these assumptions are violated in `LDSC`, but the inclusion of the additional nonlinear scores in `i-LDSC` help recover the relationship between the expectation of χ^2 test statistics and LD.

As a further demonstration of how `i-LDSC` performs when assumptions of the original LD score model are violated, we also generated synthetic phenotypes with sparse architectures using the spike-and-slab model[20]. Here, traits were simulated with solely additive effects, but this time only variants with the top or bottom $\{1, 5, 10, 25, 50, 100\}$ percentile of LD scores were given nonzero effects (see Material and Methods). Breaking the relationship assumed under the `LDSC` framework between LD scores and chi-squared statistics (i.e., that they are generally positively correlated) led to unbounded estimates of

heritability in all but the (polygenic) scenario when 100% of SNPs contributed to the phenotypic variation

(Figure 3 – figure supplement 8).

Finally, we performed a set of polygenic simulations to assess if `i-LDSC` estimates of non-additive

genetic variance could be spuriously inflated due to either (i) unobserved additive effects (see, for example,

Hemani et al.[39]), (ii) unobserved SNPs that are involved in genetic interactions, or by (iii) nonzero

correlation between the additive and interaction effect sizes in the generative model (i.e., breaking the

independence assumption in Eq. (2)). In the first setting, we observed that, across a range of both minor

allele frequencies and effect sizes, the omission of causal haplotypes had a negligible effect on the estimated

value of the coefficients in `i-LDSC` (Figure 3 – figure supplement 9). We hypothesize this is due to the

fact that the simulations were done for polygenic architectures where all SNPs have at least an additive

effect. As a result, not observing a small subset of SNPs does not hinder the ability of `i-LDSC` to estimate

genetic variance because the effect size of each SNP is small. If these simulations were conducted for sparse

architectures, we would have likely seen a greater impact on `i-LDSC`; although, we have already shown the

LD score regression framework to be uncalibrated for traits with sparse genetic architectures (again see

Figure 3 – figure supplement 8). In the second setting, we observed that the `i-LDSC` framework protects

against the false discovery of non-additive genetic effects and underestimates the variance component ϑ

when causal variants involved in pairwise interactions were unobserved (Figure 3 – figure supplement 10

and 11). As a direct comparison, estimates of the additive variance component τ in `i-LDSC` were not

affected by the unobserved interacting variants. Lastly, in the third setting, we observed that the mean

estimate of the genetic variance in both `LDSC` and `i-LDSC` had a slight upward bias as the correlation

between additive and interaction effect sizes in the generative model increased; however, the median of

these bias estimates was still near zero across all simulated scenarios and their corresponding replicates

(Figure 3 – figure supplement 12 and 13).

Application of `i-LDSC` to the UK Biobank and BioBank Japan

302 To assess whether pairwise interaction genetic effects are significantly affecting estimates of heritability

303 in empirical biobank data, we applied `i-LDSC` to 25 continuous quantitative traits from the UK Biobank

304 and BioBank Japan (Supplementary File 3). Protocols for computing GWAS summary statistics for

305 the UK Biobank are described in the Materials and Methods; while pre-computed summary statistics

306 for BioBank Japan were downloaded directly from the consortium website (see URLs). We release the

cis-acting SNP-by-SNP interaction LD scores used in our analyses on the `i-LDSC` GitHub repository from two reference groups in the 1000 Genomes: 489 individuals from the European superpopulation (EUR) and 504 individuals from the East Asian (EAS) superpopulation (see also Supplementary Files 4 and 5).

In each of the 25 traits we analyzed in the UK Biobank, we detected significant proportions of estimated genetic variation stemming from tagged pairwise cis-interactions (Table 1). This includes many canonical traits of interest in heritability analyses: height, cholesterol levels, urate levels, and both systolic and diastolic blood pressure. Our findings in Table 1 are supported by multiple published studies identifying evidence of non-additive effects playing a role in the architectures of different traits of interest. For example, Li et al.[40] found evidence for genetic interactions that contributed to the pathogenesis of coronary artery disease. It was also recently shown that non-additive genetic effects plays a significant role in body mass index[10]. Generally, we find that the traditional `LDSC` produces lower estimates of trait heritability because it does not consider the additional sources of genetic signal that `i-LDSC` does (Table 1). In BioBank Japan, 23 of the 25 traits analyzed had a significant nonlinear component detected by `i-LDSC` — with HDL and triglyceride levels being the only exceptions.

For each of the 25 traits that we analyzed, we found that the `i-LDSC` heritability estimates are significantly correlated with corresponding estimates from `LDSC` in both the UK Biobank ($r^2 = 0.988$, $P = 5.936 \times 10^{-24}$) and BioBank Japan ($r^2 = 0.849$, $P = 6.061 \times 10^{-11}$) as shown in Figure 4A. Additionally, we found that the heritability estimates for the same traits between the two biobanks are highly correlated according to both `LDSC` ($r^2 = 0.848$, $P = 7.166 \times 10^{-11}$) and `i-LDSC` ($r^2 = 0.666$, $P = 6.551 \times 10^{-7}$) analyses as shown in Figure 4B. After comparing the `i-LDSC` heritability estimates to `LDSC`, we then assessed whether there was significant difference in the amount of phenotypic variation explained by the non-additive genetic effect component in the GWAS summary statistics derived from the the UK Biobank and BioBank Japan (i.e., comparing the estimates of ϑ; see Figure 4 – figure supplement 1A). We show that, while heterogeneous between traits, the phenotypic variation explained by genetic interactions is relatively of the same magnitude for both biobanks ($r^2 = 0.372$, $P = 0.0119$). Notably, the trait with the most significant evidence of tagged cis-interaction effects in GWAS summary statistics is height which is known to have a highly polygenic architecture.

The intercepts estimated by `LDSC` and `i-LDSC` are also highly correlated in both the UK Biobank and the BioBank Japan (Figure 4 – figure supplement 1B). Recall that these intercept estimates represent the confounding factor due to uncontrolled effects. For `LDSC`, this does include phenotypic variation that is

due to unaccounted for pairwise statistical genetic interactions. The `i-LDSC` intercept estimates tend to be correlated with, but are generally different than, those computed with `LDSC` — empirically indicating that non-additive genetic variation is partitioned away and is missed when using the standard LD score alone. This result shows similar patterns in both the UK Biobank ($r^2 = 0.888$, $P = 1.962 \times 10^{-12}$) and BioBank Japan ($r^2 = 0.813$, $P = 7.814 \times 10^{-10}$).

Lastly, we performed an additional analysis in the UK Biobank where the *cis*-interaction scores are included as an annotation alongside 97 other functional categories in the stratified-LD score regression framework and its software `s-LDSC`[41] (Materials and Methods). Here, `s-LDSC` heritability estimates still showed an increase with the interaction scores versus when the publicly available functional categories were analyzed alone, but albeit at a much smaller magnitude (Table 2). The contributions from the pairwise interaction component to the overall estimate of genetic variance ranged from 0.005 for MCHC ($P = 0.373$) to 0.055 for HDL ($P = 0.575$) (Figures 4C and 4D). Furthermore, in this analysis, the estimates of the non-additive components were no longer statistically significant for any of the traits in the UK Biobank (Table 2). Despite this, these results highlight the ability of the `i-LDSC` framework to identify sources of "missing" phenotypic variance explained in heritability estimation. Importantly, moving forward, we suggest using the *cis*-interaction scores with additional annotations whenever they are available as it provides more conservative estimates of the role of non-additive effects on trait architecture.

Discussion

In this paper, we present `i-LDSC`, an extension of the LD score regression framework which aims to recover missing heritability from GWAS summary statistics by incorporating an additional score that measures the non-additive genetic variation that is tagged by genotyped SNPs. Here, we demonstrate how `i-LDSC` builds upon the original `LDSC` model through the development of new "*cis*-interaction" LD scores which help to investigate signals of *cis*-acting SNP-by-SNP interactions (Figure 1 and Figure 1 – figure supplement 1-5). Through extensive simulations, we show that `i-LDSC` is well-calibrated under the null model when polygenic traits are generated only by additive effects (Figure 2 and Figure 2 – figure supplement 1-2), we highlight that `i-LDSC` provides greater heritability estimates over `LDSC` when traits are indeed generated with *cis*-acting SNP-by-SNP interaction effects (Figure 3 and Figure 3 – figure supplement 1, and Supplementary Files 1 and 2), and we tested the robustness of `i-LDSC` on

phenotypes where assumptions of the original LD score model are violated (Figure 3 – figure supplement 2-

13). Finally, in real data, we show examples of many traits with estimated GWAS summary statistics

that tag *cis*-interaction effects in the UK Biobank and BioBank Japan (Figure 4 and Figure 4 – figure

supplement 1, Tables 1 and 2, and Supplementary Files 3-5). We have made `i-LDSC` a publicly available

command line tool that requires minimal updates to the computing environment used to run the original

implementation of LD score regression (see URLs). In addition, we provide pre-computed *cis*-interaction

LD scores calculated from the European (EUR) and East Asian (EAS) reference populations in the 1000

Genomes phase 3 data (see Data and Software Availability under Materials and Methods).

The current implementation of the `i-LDSC` framework offers many directions for future development

and applications. First, an area of future work would be to explore how the relationship between *cis*-

interaction LD scores and interaction effect sizes from the generative model of complex traits might bias

heritability estimates provided by `i-LDSC` (e.g., similar to the relationship we explored between the stan-

dard LD scores and linear effect sizes in Figure 3 – figure supplement 8). Second, as we showed with our

simulation studies (Figure 3 – figure supplement 2-8), the *cis*-interaction LD scores that we propose are

not always enough to recover explainable non-additive genetic effects for all types of trait architectures.

While we focus on pairwise *cis*-acting SNP-by-SNP statistical interactions in this work, the theoretical

concepts underlying `i-LDSC` can easily be adapted to other types of interactions as well. Third, in our

analysis of the UK Biobank and BioBank Japan, we showed that the inclusion of additional categories

via frameworks such as stratified LD score regression[42] can be used to provide more refined heritability

estimates from GWAS summary statistics while accounting for linkage (see results in Table 1 versus Table

2). A key part of our future work is to continue to explore whether considering functional annotation

groups would also improve our ability to identify tagged non-additive genetic effects. Lastly, we have

only focused on analyzing one phenotype at a time in this study. However, many previous studies have

extensively shown that modeling multiple phenotypes can often dramatically increase power[43,44]. There-

fore, it would be interesting to extend the `i-LDSC` framework to multiple traits to study nonlinear genetic

correlations in the same way that `LDSC` was recently extended to uncover additive genetic correlation

maps across traits[45].

URLs

i-LDSC software package for implementing interaction score regression, `https://github.com/lcrawlab/i-LDSC`; LDSC software package for implementing LD score regression, `https://github.com/bulik/ldsc/`; UK Biobank, `https://www.ukbiobank.ac.uk`; BioBank Japan, `http://jenger.riken.jp/en/result`; 1000 Genomes Project genetic map and haplotypes, `http://mathgen.stats.ox.ac.uk/impute/data_download_1000G_phase1_integrated.html`; Database of Genotypes and Phenotypes (dbGaP), `https://www.ncbi.nlm.nih.gov/gap`; NHGRI-EBI GWAS Catalog, `https://www.ebi.ac.uk/gwas/`; GRM-MAF-LD package, `https://github.com/arminschoech/GRM-MAF-LD`; GCTA toolkit, `https://yanglab.westlake.edu.cn/software/gcta/`.

Acknowledgements

We thank Jeffrey P. Spence, Roshni Patel, Matthew Aguirre, Mineto Ota, and our anonymous referees for insightful comments on an earlier version of this manuscript as well as the Harpak, Ramachandran, and Crawford Labs for helpful discussions. This research was conducted in part using computational resources and services at the Center for Computation and Visualization at Brown University. This research was also conducted using the UK Biobank Resource under Application Numbers 14649 (LC) and 22419 (SR). S.P. Smith and D. Udwin were trainees supported under the Brown University Predoctoral Training Program in Biological Data Science (NIH T32 GM128596). S.P. Smith was also supported by NIH RF1AG073593. S.P. Smith and A. Harpak were also supported by NIH R35 GM151108 to A. Harpak. G. Darnell was supported by NSF Grant No. DMS-1439786 while in residence at the Institute for Computational and Experimental Research in Mathematics (ICERM) in Providence, RI. This research was supported in part by an Alfred P. Sloan Research Fellowship and a David & Lucile Packard Fellowship for Science and Engineering awarded to L. Crawford. This research was also partly supported by US National Institutes of Health (NIH) grant R01 GM118652, NIH grant R35 GM139628, and National Science Foundation (NSF) CAREER award DBI-1452622 to S. Ramachandran. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of any of the funders.

Competing Interests

The authors declare no competing interests.

Materials and Methods

Generative statistical model for complex traits

Our goal in this study is to re-analyze summary statistics from genome-wide association studies (GWAS) and estimate heritability while accounting for both additive genetic associations and tagged interaction effects. We begin by assuming the following generative linear model for complex traits which can be seen as an extended view of Eq. (1) in the main text

$$\mathbf{y} = b_0 + \mathbf{X}\boldsymbol{\beta} + \mathbf{X}_D\boldsymbol{\omega} + \mathbf{W}\boldsymbol{\theta} + \boldsymbol{\varepsilon}, \qquad \boldsymbol{\varepsilon} \sim \mathcal{N}(\mathbf{0}, (1 - H^2)\mathbf{I}), \tag{7}$$

where $\mathbf{y}$ denotes an N-dimensional vector of phenotypic states for a quantitative trait of interest measured in N individuals; b_0 is an intercept term; $\mathbf{X}$ is an $N \times J$ matrix of genotypes, with J denoting the number of single nucleotide polymorphism (SNPs) encoded as $\{0, 1, 2\}$ copies of a reference allele at each locus; $\boldsymbol{\beta} = (\beta_1, \ldots, \beta_J)$ is a J-dimensional vector containing the true additive effect sizes for an additional copy of the reference allele at each locus on $\mathbf{y}$; $\mathbf{X}_D$ is an $N \times J$ matrix that represents the dominance for each genotype encoded as $\{0, 1, 1\}$ with corresponding effect sizes $\boldsymbol{\omega}$; $\mathbf{W}$ is an $N \times M$ matrix of genetic interactions; $\boldsymbol{\theta} = (\theta_1, \ldots, \theta_M)$ is an M-dimensional vector containing the interaction effect sizes; $\boldsymbol{\varepsilon}$ is a normally distributed error term with mean zero and variance scaled according to the proportion of phenotypic variation not explained by the broad-sense heritability of the trait, denoted by H^2; and $\mathbf{I}$ denotes an $N \times N$ identity matrix. Note that the encoding for dominance in $\mathbf{X}_D$ was chosen because it imposes orthogonality with the genotype encoding in $\mathbf{X}$[11,46,47].

For convenience, we will assume that the genotype matrix (column-wise), the dominance matrix (also column-wise), and trait of interest have all been standardized[18–20]. Furthermore, while the matrix $\mathbf{W}$ could encode any source of non-additive genetic interactions (e.g., gene-by-environmental effects) in theory, we limit our focus in this study to trait architectures that have been generated with contributions stemming from *cis*-acting statistical SNP-by-SNP (or pairwise) interactions. To that end, we assume that the columns of $\mathbf{W}$ are the Hadamard (element-wise) product between genotypic vectors of the form $\mathbf{x}_j \circ \mathbf{x}_k$ for the j-th and k-th variants. We also want to point out that the generative formulation of Eq. (7) can also be easily extended to accommodate other fixed effects (e.g., age, sex, or genotype principal components), as well as other random effects terms that can be used to account for sample

447 non-independence due to other environmental factors.

448 As a final set of assumptions, we will let the intercept term b_0 be a fixed parameter while allowing

449 the other coefficients to follow independent Gaussian distributions with variances proportional to their

450 individual contributions to the trait heritability [17,20–23,48]

$$\beta_j \sim \mathcal{N}(0, \varphi_\beta^2/J), \qquad \omega_j \sim \mathcal{N}(0, \varphi_\omega^2/J), \qquad \theta_m \sim \mathcal{N}(0, \varphi_\theta^2/M), \tag{8}$$

for $j = 1, \ldots, J$ and $m = 1, \ldots, M$. The broad-sense heritability of the trait is defined as $H^2 = \varphi_\beta^2 +$

$\varphi_\omega^2 + \varphi_\theta^2$. Under the generative model in Eq. (7), we then say that $\mathbb{V}[\mathbf{X}\boldsymbol{\beta}] = \varphi_\beta^2$ is the proportion of

phenotypic variation contributed by additive SNP effects, $\mathbb{V}[\mathbf{X}_D\boldsymbol{\omega}] = \varphi_\omega^2$ is the proportion of phenotypic

variation contributed by dominance effects, and the set of interactions involving some subset of causal

SNPs contribute the remaining proportion to the heritability $\mathbb{V}[\mathbf{W}\boldsymbol{\theta}] = \varphi_\theta^2$. As we mentioned in the main

text, we recognize that the appropriateness of treating genetic effects as random variables in analytical

derivations has been questioned [24], but our simulation studies show that `i-LDSC` accurately recovers

non-additive genetic variance in Eq. (7) under a broad range of conditions.

Orthogonality between additive and non-additive genetic effects

461 Assuming that the effect sizes $\{\boldsymbol{\beta}, \boldsymbol{\omega}, \boldsymbol{\theta}\}$ in Eq. (8) follow independent and zero mean Gaussian distri-

462 butions leads to orthogonality between the additive and non-additive components in Eq. (7). Since the

463 genotypes $\mathbf{X}$ and the dominance values $\mathbf{X}_D$ are fixed orthogonal matrices, it is straightforward to show

464 that $\text{Cov}[\mathbf{X}\boldsymbol{\beta}, \mathbf{X}_D\boldsymbol{\omega}] = 0$ [11,47]. The same relationship can be shown for the additive and the pairwise

465 interaction genetic effects where

$$
\begin{aligned}
\text{Cov}[\mathbf{X}\boldsymbol{\beta}, \mathbf{W}\boldsymbol{\theta}] &= \mathbb{E}[\boldsymbol{\beta}^\mathsf{T}\mathbf{X}^\mathsf{T}\mathbf{W}\boldsymbol{\theta}] - \mathbb{E}[\boldsymbol{\beta}^\mathsf{T}\mathbf{X}^\mathsf{T}]\mathbb{E}[\mathbf{W}\boldsymbol{\theta}] \\
&= \mathbb{E}\left[\sum_{rs} \beta_r \left(\mathbf{X}^\mathsf{T}\mathbf{W}\right)_{rs} \theta_s\right] - \mathbb{E}[\boldsymbol{\beta}^\mathsf{T}]\mathbf{X}^\mathsf{T}\mathbf{W}\mathbb{E}[\boldsymbol{\theta}] \\
&= \sum_{rs} \left(\mathbf{X}^\mathsf{T}\mathbf{W}\right)_{rs} \mathbb{E}[\beta_r\theta_s] - \mathbf{0}^\mathsf{T}\mathbf{X}^\mathsf{T}\mathbf{W}\mathbf{0} \\
&= \sum_{rs} \left(\mathbf{X}^\mathsf{T}\mathbf{W}\right)_{rs} \mathbb{E}[\beta_r]\mathbb{E}[\theta_s] \\
&= 0
\end{aligned}
\tag{9}
$$

467 with $\mathbf{x}_j$ and $\mathbf{w}_m$ denoting the j-th and m-th column of the individual-level genotype matrix $\mathbf{X}$ and the

468 interaction matrix $\mathbf{W}$, respectively. Note that a similar derivation to Eq. (9) can also be done for the

469 dominance and pairwise genetic interaction effects. This concept of orthogonality is important because

470 we want to preserve a unique partitioning of genetic variance when modeling a trait of interest.

Genotypes and their interactions are correlated despite being linearly independent

The design matrices $\mathbf{X}$ and $\mathbf{W}$ in Eq. (7) are not linearly dependent because the pairwise interactions

between two SNPs are encoded as the Hadamard product of two genotypic vectors in the form $\mathbf{x}_j \circ \mathbf{x}_k$

(which is a nonlinear function). Linear dependence would have implied that one could find a transfor-

mation between a SNP and an interaction term in the form $\mathbf{w}_m = c \times \mathbf{x}_j$ for some constant c. However,

despite their linear independence, $\mathbf{X}$ and $\mathbf{W}$ are themselves not orthogonal and still have a nonzero

correlation. This implies that the inner product between genotypes and their interactions is nonzero

$\mathbf{X}^\mathsf{T}\mathbf{W} \neq \mathbf{0}$. To see this, we focus on a focal SNP $\mathbf{x}_j$ and consider three different types of interactions:

• **Scenario I:** Interaction between a focal SNP with itself ($\mathbf{x}_j \circ \mathbf{x}_j$).

• **Scenario II:** Interaction between a focal SNP with a different SNP ($\mathbf{x}_j \circ \mathbf{x}_k$).

• **Scenario III:** Interaction between a focal SNP with a pair of different SNPs ($\mathbf{x}_k \circ \mathbf{x}_l$).

The following derivations rely on the fact that: (1) we assume that genotypes have been mean-centered

and scaled to have unit variance, and (2) under Hardy-Weinberg equilibrium, SNPs marginally follow a

binomial distribution $\mathbf{x}_j \sim \mathrm{Bin}(2, p)$ where p represents the minor allele frequency (MAF)[49,50].

Scenario I. The covariance between a focal SNP and an interaction with itself is $\mathrm{Cov}[\mathbf{x}_j, \mathbf{x}_j\mathbf{x}_j] =$

$\mathbb{E}[\mathbf{x}_j^3] - \mathbb{E}[\mathbf{x}_j]\mathbb{E}[\mathbf{x}_j^2]$. With mean-centered SNPs, this is proportional to $\mathbb{E}[\mathbf{x}_j^3] = (q - p)/\sqrt{2pq}$ which is the

skewness of the binomial distribution where, again, $p = \mathrm{MAF}$ and $q = 1\text{-MAF}$ of the j-th SNP.

Scenario II. Assume that we have two SNPs, $\mathbf{x}_j \sim \mathrm{Bin}(2, p_j)$ and $\mathbf{x}_k \sim \mathrm{Bin}(2, p_k)$ where p_j and p_k

represent their respective minor allele frequencies. We want to compute the correlation between $\mathbf{x}_j$ and

the interaction $\mathbf{x}_j\mathbf{x}_k$ where $\mathrm{Cov}[\mathbf{x}_j, \mathbf{x}_j\mathbf{x}_k] = \mathbb{E}[\mathbf{x}_j^2\mathbf{x}_k] - \mathbb{E}[\mathbf{x}_j]\mathbb{E}[\mathbf{x}_j\mathbf{x}_k]$. Again, with the mean-centered

assumption, the covariance is proportional to the expectation $\mathbb{E}[\mathbf{x}_j^2 \mathbf{x}_k]$. Here, with SNPs taking on values

$\{0, 1, 2\}$, the joint distribution between $\mathbf{x}_j^2$ and $\mathbf{x}_k$ can be written out as the following[51]:

	$\mathbf{x}_j^2 = 0$	$\mathbf{x}_j^2 = 1$	$\mathbf{x}_j^2 = 4$
$\mathbf{x}_k = 0$	u_{jk}^2	$2u_{jk}(1 - p_k - u_{jk})$	$(1 - p_k - u_{jk})^2$
$\mathbf{x}_k = 1$	$2u_{jk}(1 - p_j - u_{jk})$	$2u_{jk}(u_{jk} + p_j + p_k - 1) + 2(1 - p_j - u_{jk})(1 - p_k - u_{jk})$	$2(u_{jk} + p_j + p_k - 1)(1 - p_k - u_{jk})$
$\mathbf{x}_k = 2$	$(1 - p_j - u_{jk})^2$	$2(u_{jk} + p_j + p_k - 1)(1 - p_j - u_{jk})$	$(u_{jk} + p_j + p_k - 1)^2$

where $u_{jk} = (1 - p_j)(1 - p_k) + r_{jk}\sqrt{p_j p_k (1 - p_j)(1 - p_k)}$ and r_{jk} is the Pearson correlation or linkage

disequilibrium (LD) between the j-th and k-th SNPs.

Scenario III. The covariance between a focal SNP and an interaction with a pair of different SNPs

$\mathrm{Cov}[\mathbf{x}_j, \mathbf{x}_k \mathbf{x}_l]$ will be nonzero if the j-th SNP is correlated with either variant (i.e., $r_{jk} \neq 0$ or $r_{jl} \neq 0$).

Traditional estimation of additive GWAS summary statistics

As previously mentioned, the key to this work is that SNP-level GWAS summary statistics can also tag

non-additive genetic effects when there is a nonzero correlation between individual-level genotypes and

their interactions (as defined in Eq. (7)). Throughout the rest of this section, we will use $\mathbf{X}^\intercal \mathbf{X}/N$ to

denote the LD or pairwise correlation matrix between SNPs. We will then let $\mathbf{R}$ represent an LD matrix

empirically estimated from external data (e.g., directly from GWAS study data, or using a pairwise

LD map from a population that is representative of the samples analyzed in the GWAS study). The

important property here is the following

$$\mathbb{E}[\mathbf{X}^\intercal \mathbf{X}] \approx N\mathbf{R}, \qquad \mathbb{E}[\mathbf{x}_j^\intercal \mathbf{x}_j] \approx N, \qquad \mathbb{E}[\mathbf{x}_j^\intercal \mathbf{x}_k] \approx N r_{jk} \tag{10}$$

where the term r_{jk} is again defined as the Pearson correlation coefficient between the j-th and k-th SNPs,

respectively.

In traditional GWAS studies, summary statistics of the true additive effects $\boldsymbol{\beta} = (\mathbf{X}^\intercal \mathbf{X})^{-1} \mathbf{X}^\intercal \mathbf{y}$ in

510 Eq. (7) are typically derived by computing a marginal least squares estimate with the observed data

$$\widehat{\beta}_j = (\mathbf{x}_j^\mathsf{T}\mathbf{x}_j)^{-1}\mathbf{x}_j^\mathsf{T}\mathbf{y} \qquad \Longleftrightarrow \qquad \widehat{\boldsymbol{\beta}} = \text{diag}(\mathbf{X}^\mathsf{T}\mathbf{X})^{-1}\mathbf{X}^\mathsf{T}\mathbf{y}. \tag{11}$$

512 There are two key identities that may be taken from Eq. (11). The first uses Eq. (10) and is the
513 approximate relationship (in expectation) between the moment matrix $\mathbf{X}^\mathsf{T}\mathbf{y}$ and the linear effect size
514 estimates $\widehat{\boldsymbol{\beta}}$:

$$\mathbb{E}[\mathbf{X}^\mathsf{T}\mathbf{y}] = \mathbb{E}[\text{diag}(\mathbf{X}^\mathsf{T}\mathbf{X})\widehat{\boldsymbol{\beta}}] \approx N\widehat{\boldsymbol{\beta}}. \tag{12}$$

516 The second key point combines Eqs. (10) and (12) to describe the asymptotic relationship between the
517 observed marginal GWAS summary statistics $\widehat{\boldsymbol{\beta}}$ and the joint coefficient values $\boldsymbol{\beta}$ where (in expectation)

$$\mathbb{E}[\boldsymbol{\beta}] = \mathbb{E}[(\mathbf{X}^\mathsf{T}\mathbf{X})^{-1}\mathbf{X}^\mathsf{T}\mathbf{y}] \approx (N\mathbf{R})^{-1}N\widehat{\boldsymbol{\beta}} = \mathbf{R}^{-1}\widehat{\boldsymbol{\beta}}. \tag{13}$$

519 After some algebra, the above mirrors a high-dimensional regression model (in expectation) where $\widehat{\boldsymbol{\beta}} = \mathbf{R}\boldsymbol{\beta}$
520 with the estimated summary statistics as the response variables and the empirically estimated LD matrix
521 acting as the design matrix[26,29,31,32,52]. Theoretically, the resulting coefficients output from this high-
522 dimensional model are the desired true effect size estimates used to generate the phenotype of interest.

Additive GWAS summary statistics with tagged interaction effects

When interactions contribute to the architecture of complex traits (i.e., $\boldsymbol{\theta} \neq \mathbf{0}$), the marginal GWAS
summary statistics derived using least squares in Eq. (11) will also explain non-additive variation when
there is a nonzero correlation between genotypes and their interactions. To see this, we use the concept
of "omitted variable bias"[53] where the fitted model aims to estimate the true additive coefficients $\boldsymbol{\beta}$
but does not account for contributions from the non-additive components which also contribute to trait
architecture. In this case, we get the following

$$\begin{aligned} \widehat{\boldsymbol{\beta}} &= \text{diag}(\mathbf{X}^\mathsf{T}\mathbf{X})^{-1}\mathbf{X}^\mathsf{T}\mathbf{y} \\ &= \text{diag}(\mathbf{X}^\mathsf{T}\mathbf{X})^{-1}\mathbf{X}^\mathsf{T}\left[\mathbf{X}\boldsymbol{\beta} + \mathbf{X}_D\boldsymbol{\omega} + \mathbf{W}\boldsymbol{\theta} + \boldsymbol{\varepsilon}\right]. \end{aligned} \tag{14}$$

Since we assume that the genotypes are orthogonal to both the dominance effects in Eq. (7), we know that $\mathbf{X}^\intercal\mathbf{X}_D = \mathbf{0}$. This simplifies the above to be the following

$$\widehat{\boldsymbol{\beta}} = \text{diag}(\mathbf{X}^\intercal\mathbf{X})^{-1}\mathbf{X}^\intercal\mathbf{X}\boldsymbol{\beta} + \text{diag}(\mathbf{X}^\intercal\mathbf{X})^{-1}\mathbf{X}^\intercal\mathbf{W}\boldsymbol{\theta} + \text{diag}(\mathbf{X}^\intercal\mathbf{X})^{-1}\mathbf{X}^\intercal\boldsymbol{\varepsilon} \tag{15}$$

where the matrix $\mathbf{X}^\intercal\mathbf{W}$ (which we showed to be nonzero) can be interpreted as the sample correlation between individual-level genotypes and the *cis*-interactions between causal SNPs. By taking the expectation using Eqs. (10) and (12), we get the following alternative (approximate) relationship between the observed marginal GWAS summary statistics $\widehat{\boldsymbol{\beta}}$ and the true coefficient values $\boldsymbol{\beta}$

$$\mathbb{E}[\widehat{\boldsymbol{\beta}}] = \mathbf{R}\boldsymbol{\beta} + \mathbf{V}\boldsymbol{\theta}, \tag{16}$$

which results from our initial assumption that the residuals are normally distributed with mean zero $\mathbb{E}[\boldsymbol{\varepsilon}] = \mathbf{0}$ in Eq. (7). Here, we define $\mathbf{V}$ to represent a sample estimate of the correlation between the individual-level genotypes and the non-additive genetic interaction matrix such that $\mathbb{E}[\mathbf{X}^\intercal\mathbf{W}] \approx N\mathbf{V}$. Similar to the LD matrix $\mathbf{R}$, the correlation matrix $\mathbf{V}$ is also assumed to be computed from reference panel data. Intuitively, when $\boldsymbol{\theta} \neq \mathbf{0}$ there is additional phenotypic variation contributed by pairwise interactions that can be explained by GWAS effect size estimates. Moreover, when $\mathbf{V}\boldsymbol{\theta} = \mathbf{0}$, then the relationship in Eq. (16) converges onto the conventional asymptotic assumption (in expectation) between GWAS summary statistics and the true additive coefficients in Eq. (13)[26,29,31,32,52].

Connection to quantitative genetics theory

The concept of additive genetic effects partially explaining non-additive variation has also described in classical quantitative genetics[25,35,36]. Consider an individual genotyped at J loci each with major and minor alleles A and B, respectively. Let p_j be the allele frequency of A at the j-th locus, a_j denote the additive effect, and $[aa]_{jk}$ be the additive-by-additive (pairwise) interaction effect between loci j and k, and $[aaa]_{jkl}$ represent a third order interaction between loci j, k, and l. For simplicity in presentation, assume that dominance only makes a small contribution to the genetic variance[11,54,55]. The population

mean is given as the following

$$\mu = 2\sum_{j=1}^{J} p_j a_j + 4\sum_{j=1}^{J}\sum_{k>j}^{J} p_j p_k [aa]_{jk} + 8\sum_{j=1}^{J}\sum_{k>j}^{J}\sum_{l>k>j}^{J} p_j p_k p_l [aaa]_{jkl} + \cdots \tag{17}$$

We follow the assumption that the genetic variation in human complex traits can predominately be explained by additive effects, with the remainder variation being mostly explained by additive-by-additive effects[48,56–58]. As a result, we will ignore the higher-order interaction terms in Eq. (17). Under Hardy-Weinberg equilibrium, we can find the average effect by taking the first derivative of the population mean with respect to the frequency of the increasing allele[35,36]. For the j-th SNP, the average effect (including terms up to second-order interaction) is given by the following

$$\eta_j = \frac{1}{2}\left(\frac{\partial \mu}{\partial p_j}\right) = a_j + 2\sum_{k\neq j}^{J} p_k [aa]_{jk} + O\left([aaa]_{jkl}\right) \tag{18}$$

which notably contains both the additive effect and a summation of additive-by-additive interactions between pairs of loci. The additive genetic variance for the j-th SNP takes on the following form

$$\begin{aligned}\sigma_A^2(j) &= 2p_j(1-p_j)\left[a_j + 2\sum_{k\neq j}^{J} p_k [aa]_{jk}\right]^2 \\ &= 2p_j(1-p_j)\left[a_j^2 + 2a_j\sum_{k\neq j}^{J} p_k [aa]_{jk} + 4\left(\sum_{k\neq j}^{J} p_k [aa]_{jk}\right)^2\right]\end{aligned} \tag{19}$$

which is the product of the square of the average effect in Eq. (18) and the heterozygosity at j-th locus $\mathbb{V}[\mathbf{x}_j] = 2p_j(1-p_j)$ (again assuming that SNPs marginally follow a binomial distribution). The total additive variance is then obtained by summing over the J loci such that $\sigma_A^2 = \sum_j \sigma_A^2(j)$[37].

We can derive a parallel construction for additive genetic variance using the generative random effect model presented in Eq. (7)[35]. Here, we will leverage that with genotype data taken for N individuals, $\sum_i x_{ij}/N = 2p_j$. Ignoring the assumed small contributions from dominance effects, the population mean

for a quantitative trait $\mathbf{y}$ can be written as the following

$$
\begin{aligned}
\mu = \frac{1}{N}\sum_{i=1}^{N} y_i &= \frac{1}{N}\sum_{i=1}^{N}\left[b_0 + \sum_{j=1}^{J} x_{ij}\beta_j + \sum_{j=1}^{J}\sum_{k>j}^{J} x_{ij}x_{ik}\theta_{jk} + \varepsilon_i \right] \\
&= b_0 + 2\sum_{j=1}^{J} p_j\beta_j + 4\sum_{j=1}^{J}\sum_{k>j}^{J} p_j p_k \theta_{jk} + \frac{1}{N}\sum_{i=1}^{N}\varepsilon_i.
\end{aligned}
\tag{20}
$$

To find the average effect for the j-th locus, we this time take the first derivative of the population mean in Eq. (20) with respect to the allele frequency such that

$$
\eta_j = \frac{1}{2}\left(\frac{\partial \mu}{\partial p_j}\right) = \beta_j + 2\sum_{k\neq j}^{J} p_k \theta_{jk}
\tag{21}
$$

which, similar to the theoretical form in quantitative genetics, also contains both the additive effect of the j-th SNP and additional terms encoding the interaction effect between the j-th SNP and all other variants in the data. Once again, under Hardy-Weinberg equilibrium, the additive variance for the j-th SNP is found as taking on the following form

$$
\begin{aligned}
\sigma_A^2(j) &= 2p_j(1-p_j)\left[\beta_j + 2\sum_{k\neq j}^{J} p_k \theta_{jk}\right]^2 \\
&= 2p_j(1-p_j)\left[\beta_j^2 + 2\beta_j\sum_{k\neq j}^{J} p_k\theta_{jk} + 4\left(\sum_{k\neq j}^{J} p_k\theta_{jk}\right)^2\right]
\end{aligned}
\tag{22}
$$

where we can explicitly draw connections between the two frameworks by setting $\beta_j = a_j$ and $\theta_{jk} = [aa]_{jk}$. Note that when there no non-additive effects (such that $\boldsymbol{\theta} = \mathbf{0}$), the above reduces to $\sigma_A^2 = \sum_j 2p_j(1-p_j)\beta_j^2$ which resembles the classical form for the additive genetic variance[58].

Full derivation of interaction LD score regression

In order to derive the interaction LD score (i-LDSC) regression framework, recall that our goal is to recover missing heritability from GWAS summary statistics by incorporating an additional score that measures the non-additive genetic variation that is tagged by genotyped SNPs. To do this, we build upon the LD score regression framework and the LDSC software[17]. Here, we assume nonzero contributions from *cis*-acting pairwise interaction effects in the generative model of complex traits as in Eq. (16), and we use

the observed least squares estimates from Eq. (11) to compute chi-square statistics $\chi_j^2 = N\widehat{\beta}_j^2$ for every

$j = 1, \ldots, J$ variant in the data. Taking the expectation of these statistics yields

$$\mathbb{E}[\chi_j^2] = N\mathbb{E}[\widehat{\beta}_j^2] = N\left[\mathbb{V}[\widehat{\beta}_j] + \left(\mathbb{E}[\widehat{\beta}_j]\right)^2\right]. \tag{23}$$

We can simplify Eq. (23) in two steps. First, by combining the prior assumption in Eq. (8) and the

asymptotic approximation in Eq. (16), we can show that marginal expectation (i.e., when not conditioning

on the true coefficients) $\mathbb{E}[\widehat{\beta}_j] = 0$ for all variants. Second, by conditioning on the generative model from

Eq. (7), we can use the law of total variance to simplify $\mathbb{V}[\widehat{\beta}_j]$ where

$$
\begin{aligned}
\mathbb{V}[\widehat{\beta}_j] &= \mathbb{E}[\mathbb{V}[\widehat{\beta}_j \,|\, \mathbf{X}]] + \mathbb{V}[\mathbb{E}[\widehat{\beta}_j \,|\, \mathbf{X}]] \approx \mathbb{E}[\mathbb{V}[\mathbf{x}_j^\mathsf{T}\mathbf{y}/N \,|\, \mathbf{X}]] + 0 \\
&= \mathbb{E}\left[\frac{1}{N^2}\mathbf{x}_j^\mathsf{T}\left\{\mathbb{V}[\mathbf{y}\,|\,\mathbf{X}]\right\}\mathbf{x}_j\right] \\
&= \mathbb{E}\left[\frac{1}{N^2}\mathbf{x}_j^\mathsf{T}\left\{\frac{\varphi_\beta^2}{J}\mathbf{X}\mathbf{X}^\mathsf{T} + \frac{\varphi_\omega^2}{J}\mathbf{X}_D\mathbf{X}_D^\mathsf{T} + \frac{\varphi_\theta^2}{M}\mathbf{W}\mathbf{W}^\mathsf{T} + (1 - H^2)\right\}\mathbf{x}_j\right] \\
&= \mathbb{E}\left[\frac{1}{N^2}\left\{\frac{\varphi_\beta^2}{J}\mathbf{x}_j^\mathsf{T}\mathbf{X}\mathbf{X}^\mathsf{T}\mathbf{x}_j + \frac{\varphi_\omega^2}{J}\mathbf{x}_j^\mathsf{T}\mathbf{X}_D\mathbf{X}_D^\mathsf{T}\mathbf{x}_j + \frac{\varphi_\theta^2}{M}\mathbf{x}_j^\mathsf{T}\mathbf{W}\mathbf{W}^\mathsf{T}\mathbf{x}_j + N(1 - H^2)\right\}\right] \\
&= \mathbb{E}\left[\frac{1}{N^2}\left\{\frac{\varphi_\beta^2}{J}\mathbf{x}_j^\mathsf{T}\mathbf{X}\mathbf{X}^\mathsf{T}\mathbf{x}_j + \frac{\varphi_\theta^2}{M}\mathbf{x}_j^\mathsf{T}\mathbf{W}\mathbf{W}^\mathsf{T}\mathbf{x}_j + N(1 - H^2)\right\}\right]
\end{aligned}
$$

since $\mathbf{x}_j^\mathsf{T}\mathbf{X}_D = \mathbf{0}$. Using the same logic from the original LDSC regression framework [17], we can use Isserlis'

theorem [59] to write the above in terms of more familiar quantities based on sample correlations

$$\frac{1}{N^2}\mathbf{x}_j^\mathsf{T}\mathbf{X}\mathbf{X}^\mathsf{T}\mathbf{x}_j = \sum_{k=1}^{J}\widetilde{r}_{jk}^2, \qquad \frac{1}{N^2}\mathbf{x}_j^\mathsf{T}\mathbf{W}\mathbf{W}^\mathsf{T}\mathbf{x}_j = \sum_{m=1}^{M}\widetilde{v}_{jm}^2 \tag{24}$$

where $\widetilde{r}_{jk}$ is used to denote the sample correlation between additively-coded genotypes at the j-th and

k-th variants, and $\widetilde{v}_{jm}$ is used to denote the sample correlation between the genotype of the j-th variant

and the m-th genetic interaction on the phenotype of interest (again see Eq. (16)). Furthermore, we can

use the delta method (only displaying terms up to $\mathcal{O}(1/N^2)$) to show that (in expectation)

$$\mathbb{E}[\widetilde{r}_{jk}^2] \approx r_{jk}^2 + (1 - r_{jk}^2)/N, \qquad \mathbb{E}\left[\widetilde{v}_{jm}^2\right] \approx v_{jm}^2 + \left(1 - v_{jm}^2\right)/N. \tag{25}$$

Next, we can then approximate the quantities in Eq. (24) via the following

$$\mathbb{E}\left[\sum_{k=1}^{J}\widetilde{r}_{jk}^2\right] \approx \ell_j + (J - \ell_j)/N, \qquad \mathbb{E}\left[\sum_{m=1}^{M}\widetilde{v}_{jm}^2\right] \approx f_j + (M - f_j)/N \tag{26}$$

where ℓ_j is the corresponding LD score for the additive effect of the j-th variant and f_j represents the "interaction" LD score between the j-th SNP and all other variants in the data set [23], respectively. Altogether, this leads to the specification of the univariate framework with the j-th SNP

$$\mathbb{E}[\chi_j^2] \approx N\left[\left(\frac{\varphi_\beta^2}{J}\right)\ell_j + \left(\frac{\varphi_\theta^2}{M}\right)f_j + \frac{1}{N}(1 - H^2)\right] = \ell_j\tau + f_j\vartheta + 1 \tag{27}$$

where we define $\tau = N\varphi_\beta^2/J$ as estimates of the additive genetic signal, the coefficient $\vartheta = N\varphi_\theta^2/M$ as an estimate of the proportion of phenotypic variation explained by tagged pairwise interaction effects, and $\mathbf{1}$ is the intercept meant to model the misestimation due to uncontrolled confounding effects (e.g., cryptic relatedness and population stratification). Similar to the original LDSC formulation, an intercept greater than one means significant bias. Note that the simplification for many of the terms above such as $(1 - H^2)/N \approx 1/N$ results from our assumption that the number of individuals in our study is large. For example, the sample sizes for each biobank-scale study considered in the analyses of this manuscript are at least on the order of $N \geq 10^4$ observations (see Table 5). Altogether, we can jointly express Eq. (27) in multivariate form as

$$\mathbb{E}[\boldsymbol{\chi}^2] \approx \boldsymbol{\ell}\tau + \boldsymbol{f}\vartheta + \mathbf{1} \tag{28}$$

where $\boldsymbol{\chi}^2 = (\chi_1^2, \ldots, \chi_J^2)$ is a J-dimensional vector of chi-square summary statistics, and $\boldsymbol{\ell} = (\ell_1, \ldots, \ell_J)$ and $\boldsymbol{f} = (f_1, \ldots, f_J)$ are J-dimensional vectors of additive and *cis*-interaction LD scores, respectively. It is important to note that, while $\boldsymbol{\chi}^2$ must be recomputed for each trait of interest, both vectors $\boldsymbol{\ell}$ and $\boldsymbol{f}$ only need to be constructed once per reference panel or individual-level genotypes (see next section for efficient computational strategies).

To identify summary statistics that have significant tagged interaction effects, we test the null hypothesis $H_0 : \vartheta = 0$. The i-LDSC software package implements the same model fitting strategy as LDSC. Here, we use weighted least squares to fit the joint regression in Eq. (28) such that

$$\widehat{\vartheta} = (\boldsymbol{f}^\intercal\boldsymbol{\Psi}\boldsymbol{f})^{-1}\boldsymbol{f}^\intercal\boldsymbol{\Psi}\boldsymbol{\chi}^2, \qquad \psi_{jj} = \left[\ell_j\widehat{\tau} + f_j\widehat{\vartheta} + 1\right]^{-2} \tag{29}$$

where $\boldsymbol{\Psi}$ is a $J \times J$ diagonal weight matrix with nonzero elements set to values inversely proportional to the conditional variance $\mathbb{V}[\chi_j^2 \,|\, \ell_j, f_j] = \psi_{jj}^{-1}$ to adjust for both heteroscedasticity and over-estimation of the summary statistics for each SNP[17]. Standard errors for each coefficient estimate are derived via a jackknife over blocks of SNPs in the data[42], and we then use those standard errors to derive P-values with a two-sided test (i.e., testing the alternative hypothesis $H_A : \vartheta \neq 0$). It is worth noting that the block-jackknife approach tends to be conservative and yield larger standard errors for hypothesis testing[60]. As an alternative, we could first run `i-LDSC` using the block-jackknife procedure over all traits in a study and then use the average of the standard errors to calculate the statistical significance of coefficient estimates; but we do not explore this strategy here and leave that for future work. The quantitative genetics expression for the additive variance σ_A^2 in Eq. (22) is important because it represents the theoretical upper bound on the proportion of phenotypic variance that can be explained from GWAS summary statistics via `i-LDSC`. Using this relationship, we can write the following (approximate) inequality

$$\widehat{\tau} + \widehat{\vartheta} \lesssim \sum_{j=1}^{J} 2 p_j (1 - p_j) \left[\beta_j + 2 \sum_{k \neq j}^{J} p_k \theta_{jk} \right]^2 = \sigma_A^2. \tag{30}$$

For all analyses in this paper, we estimate proportion of phenotypic variance explained by genetic effects using a sum of the coefficients $\widehat{\tau} + \widehat{\vartheta}$ (i.e., the estimated additive component plus the additional genetic variance explained by the tagged pairwise interaction effects).

Efficient computation of *cis*-interaction LD scores

In practice, *cis*-interaction LD scores in `i-LDSC` can be computed efficiently through realizing two key opportunities for optimization. First, given J SNPs, the full matrix of genome-wide interaction effects $\mathbf{W}$ contains on the order of $J(J-1)/2$ total pairwise interactions. However, to compute the *cis*-interaction score for each SNP, we simply can replace the full $\mathbf{W}$ matrix with a subsetted matrix $\mathbf{W}_j$ which includes only interactions involving the j-th SNP. Analogous to the original `LDSC` formulation[17], we consider only interactive SNPs within a *cis*-window proximal to the focal j-th SNP for which we are computing the `i-LDSC` score. In the original `LDSC` model, this is based on the observation that LD decays outside of a window of 1 centimorgan (cM)[17]; therefore, SNPs outside the 1 cM window centered on the j-th SNP j will not significantly contribute to its LD score. The second opportunity for optimization comes from the fact that the matrix of interaction effects for any focal SNP, $\mathbf{W}_j$, does not need to be explicitly

generated. Referencing Eq. (24), the `i-LDSC` scores are defined as $\mathbf{x}_j^\mathsf{T} \mathbf{W}_j \mathbf{W}_j^\mathsf{T} \mathbf{x}_j / N^2$. This can be re-written as $\mathbf{x}_j^\mathsf{T}(\mathbf{D}_j \mathbf{X}^{(j)})(\mathbf{D}_j \mathbf{X}^{(j)})^\mathsf{T} \mathbf{x}_j$, where $\mathbf{D}_j = \mathrm{diag}(\mathbf{x}_j)$ is a diagonal matrix with the j-th genotype as its nonzero elements[23] and $\mathbf{X}^{(j)}$ denotes the subset SNPs within a *cis*-window proximal to the focal j-th SNP. This means that the `i-LDSC` score for the j-th SNP can be simply computed as the following

$$f_j \approx \frac{1}{N^2}(\mathbf{x}_j^\mathsf{T})^2 \mathbf{X}^{(j)} \mathbf{X}^{(j)\mathsf{T}}(\mathbf{x}_j)^2. \tag{31}$$

With these simplifications, the computational complexity of generating `i-LDSC` scores reduces to that of computing LD scores — modulo a vector-by-vector Hadamard product which, for each SNP, is constant factor of N (i.e., the number of genotyped individuals).

Coefficient estimates as determined by *cis*-interaction window size

When computing *cis*-interaction LD scores, the most important de*cis*ion is choosing the number of interacting SNPs to include in $\mathbf{X}^{(j)}$ (or equivalently $\mathbf{W}_j$ for each j-th focal SNP in the calculation of f_j in Eq. (31)). The `i-LDSC` framework considers different estimating windows to account for our lack of *a priori* knowledge about the "correct" non-additive genetic architecture of traits. Theoretically, one could follow previous work[20,28,30,32,33,61] by considering an L-valued grid of possible SNP interaction window sizes. After fitting a series of `i-LDSC` regressions with *cis*-interaction LD scores $\boldsymbol{f}^{(l)}$ generated under the L-different window sizes, we could compute normalized importance weights using their maximized likelihoods via the following

$$\pi^{(l)} = \frac{\mathcal{L}\left(\boldsymbol{\ell}, \boldsymbol{f}^{(l)}; \widehat{\boldsymbol{\beta}}\right)}{\sum_{l'} \mathcal{L}\left(\boldsymbol{\ell}, \boldsymbol{f}^{(l')}; \widehat{\boldsymbol{\beta}}\right)}, \qquad \sum_{l=1}^{L} \pi^{(l)} = 1. \tag{32}$$

As a final step in the model fitting procedure, we could then compute averaged estimates of the coefficients τ and ϑ by marginalizing (or averaging) over the L-different grid combinations of estimating windows

$$\widehat{\tau} = \sum_{l=1}^{L} \pi^{(l)} \widehat{\tau}^{(l)}, \qquad \widehat{\vartheta} = \sum_{l=1}^{L} \pi^{(l)} \widehat{\vartheta}^{(l)}. \tag{33}$$

This final step can be viewed as an analogy to model averaging where marginal estimates are computed via a weighted average using the importance weights[62]. In the current study, we explore the utility of

cis-interaction LD scores generated with different window sizes ± 5, ± 10, ± 25, and ± 50 SNPs around each

j-th focal SNP. In practice, we find that *cis*-interaction LD scores that are calculated using larger windows

lead to the most robust estimates of heritability while also not over representing the total phenotypic

variation explained by tagged non-additive genetic effects (see Figure 3 – figure supplement 1). Therefore,

unless otherwise stated, we use *cis*-interaction LD scores calculated with a ± 50 SNP interaction window

for all simulations and real data analyses conducted in this work. For a direct comparison between

choosing a single window size versus the model averaging strategy described above, see Supplementary

Files 1 and 2.

Relationship between minor allele frequency and effect size

The `LDSC` software computes LD scores using annotations over equally spaced minor allele frequency

(MAF) bins. These annotations enable the per trait relationship between the MAF and the effect size

of each variant in the genome to vary based on the discrete category (or MAF bin) it is placed into.

This additional flexibility is intended to help `LDSC` be more robust when estimating heritability. The

relationship between MAF and effect size is already implicitly encoded in the `LDSC` formulation since we

assume genotypes are normalized. When normalizing by the variance of each SNP (or equivalently its

MAF), we make the assumption that rare variants inherently have larger effect sizes. There exists a true

functional relationship between MAF and effect size which is likely to be somewhere between the two

extremes of (i) normalizing each SNP by its MAF and (ii) allowing the variance per SNP to be dictated

by its MAF.

705   Recent approaches have proposed using a single parameter α to better represent the nonlinear rela-

706 tionship between MAF and variant effect size. The main idea is that this α not only provides the same

707 additional flexibility to `LDSC` as the MAF-based discrete annotations, but it also empirically yields even

708 more precise heritability estimates[63]. Namely, we use

$$\ell_j(c) := \sum_k L_{jk}(\alpha) a_c(k), \qquad L_{jk}(\alpha) = r_{jk}^2 \mathbb{V}[\mathbf{x}_k]^{1-\alpha} \tag{34}$$

710 where $a_c(k)$ is the annotation value for the *c*-th categorical bin. The α parameter is unknown in practice

711 and needs to be estimated for any given trait. While standard ranges for α can be used for heritability es-

712 timates, we use a restricted maximum likelihood (REML) based method which was recently developed[38].

In the `i-LDSC` software, we use this α construction to handle the relationship between MAF and variant effect size for two specific reasons. First, by constructing the LD scores using α, we more accurately capture the variation in chi-square test statistics due to additive effects[63]. Second, we note that there is correlation between MAF and (*i*) LD scores, (*ii*) *cis*-interaction LD scores, and (*iii*) trait architecture. To that end, if we do not properly condition on MAF, there becomes additional bias, and we may falsely attribute some amount of variation in the chi-square test statistics to LD or the tagged interaction effects. Therefore, in our formulation, we include an α term on the LD scores to condition on this effect. We demonstrate in simulations that this removes the bias introduced by the relationship between MAF and trait architecture, and it mitigates potential inflation of type I error rates in the `i-LDSC` test.

Estimation of allele frequency parameters

In the main text, we analyzed 25 complex traits in both the UK Biobank and BioBank Japan data sets. In order to account for minor allele frequency (MAF) dependent trait architecture, we calculated α values for each trait that had not been analyzed by previous studies[38]. The α estimates for each of the 25 traits analyzed in this study are shown in Table 4. Intuitively, α parameterizes the weighting of the effects of each individual variant given its frequency in the study cohort and can take on values in the range of [-1,0]. More negative values of α indicate that lower frequency variants contribute more to the observed variation in a trait of interest, whereas values of α closer to zero indicate that common variants contribute a greater amount of variation to observed trait values.

We took α values for 11 traits (again see Table 4) that had previously been calculated from Schoech et al.[38]. For the remaining 14 traits analyzed in this study, we followed the estimation protocol described in the same manuscript. Specifically, using the variants passing the quality control step in our pipeline for 25,000 randomly selected individuals in the UK Biobank cohort, we constructed MAF-dependent genetic relatedness matrices for values of $\alpha = \{-1, -0.95, -0.9, \ldots, 0\}$ using the `GRM-MAF-LD` software, `https://github.com/arminschoech/GRM-MAF-LD`. We then used the `GCTA` software[64] to obtain heritability and likelihood estimates using REML for each α-trait pairing. We then fit a trait-specific profile likelihood across the range of α values and estimate the maximum likelihood value of α using a natural cubic spline.

Simulation studies

We used a simulation scheme to generate synthetic quantitative traits and SNP-level summary statistics under multiple genetic architectures using real genome-wide data from individuals of self-identified European ancestry in the UK Biobank. Here, we consider phenotypes that have some combination of additive effects, *cis*-acting interactions, and a third source of genetic variance stemming from either gene-by-environment (G×E) or gene-by-ancestry (G×Ancestry) effects. For each scenario, we select some set of SNPs to be causal and assume that complex traits are generated via the following general linear model

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{W}\boldsymbol{\theta} + \mathbf{Z}\boldsymbol{\gamma} + \boldsymbol{\varepsilon}, \qquad \boldsymbol{\varepsilon} \sim \mathcal{N}(\mathbf{0}, \delta^2\mathbf{I}), \tag{35}$$

where $\mathbf{y}$ is an N-dimensional vector containing all the phenotypes; $\mathbf{X}$ is an $N \times J$ matrix of genotypes encoded as 0, 1, or 2 copies of a reference allele; $\boldsymbol{\beta}$ is a J-dimensional vector of additive effect sizes for each SNP; $\mathbf{W}$ is an $N \times M$ matrix which holds all pairwise interactions between the randomly selected subset of the interacting SNPs with corresponding effects $\boldsymbol{\theta}$; $\mathbf{Z}$ is an $N \times K$ matrix of either G×E or G×Ancestry interactions with coefficients $\boldsymbol{\gamma}$; and $\boldsymbol{\varepsilon}$ is an N-dimensional vector of environmental noise. The phenotypic variation is assumed to be $\mathbb{V}[\mathbf{y}] = 1$. All additive and interaction effect sizes for SNPs are randomly drawn from independent standard Gaussian distributions and then rescaled so that they explain a fixed proportion of the phenotypic variance $\mathbb{V}[\mathbf{X}\boldsymbol{\beta}] + \mathbb{V}[\mathbf{W}\boldsymbol{\theta}] + \mathbb{V}[\mathbf{Z}\boldsymbol{\gamma}] = H^2$. Note that we do not assume any specific correlation structure between the effect sizes $\boldsymbol{\beta}$, $\boldsymbol{\theta}$, and $\boldsymbol{\gamma}$. We then rescale the random error term such that $\mathbb{V}[\boldsymbol{\varepsilon}] = (1 - H^2)$. In the main text, we compare the traditional `LDSC` to its direct extension in `i-LDSC`. For each method, GWAS summary statistics are computed by fitting a single-SNP univariate linear model via least squares where $\widehat{\beta}_j = (\mathbf{x}_j^\intercal \mathbf{x}_j)^{-1}\mathbf{x}_j^\intercal\mathbf{y}$ for every $j = 1, \ldots, J$ SNP in the data. These effect size estimates are used to derive the chi-square test statistics $\chi_j^2 = N\widehat{\beta}_j^2$. We implement both `LDSC` and `i-LDSC` with the LD matrix $\mathbf{R} = \mathbf{X}^\intercal\mathbf{X}/N$ and the *cis*-interaction correlation matrix $\mathbf{V} = \mathbf{X}^\intercal\mathbf{W}/N$ being computed using a reference panel of 489 individuals from the European superpopulation (EUR) of the 1000 Genomes Project. The resulting matrices $\mathbf{R}$ and $\mathbf{V}$ are used to compute the additive and *cis*-interaction LD scores, respectively.

Polygenic simulations with *cis*-interactions. In our first set of simulations, we consider phenotypes with polygenic architectures that are made up of only additive and *cis*-acting SNP-by-SNP interactions.

Here, we begin by assuming that every SNP in the genome has at least a small additive effect on the traits of interest. Next, when generating synthetic traits, we assume that the additive effects make up $\rho\%$ of the heritability while the pairwise interactions make up the remaining $(1-\rho)\%$. Alternatively, the proportion of the heritability explained by additivity is said to be $\mathbb{V}[\mathbf{X}\boldsymbol{\beta}] = \rho H^2$, while the proportion detailed by interactions is given as $\mathbb{V}[\mathbf{W}\boldsymbol{\theta}] = (1-\rho)H^2$. The setting of $\rho = 1$ represents the limiting null case for `i-LDSC` where the variation of a trait is driven by solely additive effects. Here, we use the same simulation strategy used in Crawford et al. [23] where we divide the causal *cis*-interaction variants into two groups. One may view the SNPs in group #1 as being the "hubs" of an interaction map. SNPs in group #2 are selected to be variants within some kilobase (kb) window around each SNP in group #1. Given different parameters for the generative model in Eq. (35), we simulate data mirroring a wide range of genetic architectures by toggling the following parameters:

- heritability: $H^2 = 0.3$ and 0.6;

- proportion of phenotypic variation that is generated by additive effects: $\rho = 0.5$, 0.8, and 1;

- percentage of SNPs selected to be in group #1: 1% (sparse), 5%, and 10% (polygenic);

- genomic window used to assign SNPs to group #2: ± 10 and ± 100 kilobase (kb);

- allele frequency parameter: α = -1, -0.5, and 0.

All figures and tables show the mean performances (and standard errors) across 100 simulated replicates.

Polygenic simulations with gene-by-environmental effects.   In our second set of simulations, we continue to consider phenotypes with polygenic architectures that are made up of only additive and *cis*-acting SNP-by-SNP interactions; however, now we also consider each trait to have contributions stemming from nonzero G×E effects. Here, both the additive and *cis*-interaction effects are simulated in the same way as previously described where, for the two groups of interacting variants, 10% of SNPs were selected to be in group #1 and we chose ± 10 kb windows to assign SNPs to group #2. To create G×E effects, we follow a simulation strategy implemented by Zhu et al. [65] and split our sample population in half to emulate two subsets of individuals coming from different environments. We randomly draw the effect sizes for the first environment from a standard Gaussian distribution which we denote as $\boldsymbol{\gamma}_1$. We then selected an amplification coefficient w and set the effect sizes of the G×E interactions in the second

environment to be a scaled version of the first environment effects where $\boldsymbol{\gamma}_2 = w\boldsymbol{\gamma}_1$. In this paper, we generate traits with heritability $H^2 = \{0.3, 0.6\}$ and amplification coefficients set to $w = [1.1, 1.2, \ldots, 2]$. For the first set of simulations, we hold the proportion of phenotypic variation explained by the different genetic components constant by fixing:

- $H^2 = 0.3$: $\mathbb{V}[\mathbf{X}\boldsymbol{\beta}] = 0.15$; $\mathbb{V}[\mathbf{W}\boldsymbol{\theta}] = 0.075$; and $\mathbb{V}[\mathbf{Z}\boldsymbol{\gamma}] = 0.075$;

- $H^2 = 0.6$: $\mathbb{V}[\mathbf{X}\boldsymbol{\beta}] = 0.3$; $\mathbb{V}[\mathbf{W}\boldsymbol{\theta}] = 0.15$; and $\mathbb{V}[\mathbf{Z}\boldsymbol{\gamma}] = 0.15$;

where $\mathbf{Z} = [\mathbf{X}_1, \mathbf{X}_2]$ is the set of genotypes split according to environment and $\boldsymbol{\gamma} = [\boldsymbol{\gamma}_1, \boldsymbol{\gamma}_2]$. To test the sensitivity of the *cis*-interaction LD scores to other sources of non-additive variation, we also repeated the same simulations where there were only additive and G×E effects contributing equally to trait architecture:

- $H^2 = 0.3$: $\mathbb{V}[\mathbf{X}\boldsymbol{\beta}] = 0.15$; $\mathbb{V}[\mathbf{W}\boldsymbol{\theta}] = 0$; and $\mathbb{V}[\mathbf{Z}\boldsymbol{\gamma}] = 0.15$;

- $H^2 = 0.6$: $\mathbb{V}[\mathbf{X}\boldsymbol{\beta}] = 0.3$; $\mathbb{V}[\mathbf{W}\boldsymbol{\theta}] = 0$; and $\mathbb{V}[\mathbf{Z}\boldsymbol{\gamma}] = 0.3$.

Again all figures show the mean performances (and standard errors) across 100 simulated replicates.

Polygenic simulations with gene-by-ancestry effects. In our third set of simulations, we consider phenotypes with polygenic architectures that are made up of additive, *cis*-interactions, and G×Ancestry effects. Here, we follow Sohail et al. [66] and first run a matrix decomposition on the individual-level genotype matrix $\mathbf{X} = \mathbf{U}\mathbf{Q}^{\intercal}$ where $\mathbf{U}$ is a unitary $N \times K$ score matrix, $\mathbf{Q}$ is a $K \times J$ loadings matrix, and K represents the number of (predetermined) principal components (PCs). To generate G×Ancestry interactions, we then create the matrix $\mathbf{Z}_k = \mathbf{X}\mathbf{q}_k$ where $\mathbf{q}_k$ is a J-dimensional vector of SNP loadings for the k-th principal component. In this paper, we generate traits with heritability $H^2 = \{0.3, 0.6\}$ and interaction effects taken over $k = 1, \ldots, 10$ principal components. For the first set of simulations, we hold the proportion of phenotypic variation explained by the different genetic components constant by fixing:

- $H^2 = 0.3$: $\mathbb{V}[\mathbf{X}\boldsymbol{\beta}] = 0.15$; $\mathbb{V}[\mathbf{W}\boldsymbol{\theta}] = 0.075$; and $\mathbb{V}[\mathbf{Z}\boldsymbol{\gamma}] = 0.075$;

- $H^2 = 0.6$: $\mathbb{V}[\mathbf{X}\boldsymbol{\beta}] = 0.3$; $\mathbb{V}[\mathbf{W}\boldsymbol{\theta}] = 0.15$; and $\mathbb{V}[\mathbf{Z}\boldsymbol{\gamma}] = 0.15$;

To test the sensitivity of the *cis*-interaction LD scores to other sources of non-additive variation, we also repeated the same simulations where there were only additive and G×E effects contributing equally to trait architecture:

- $H^2 = 0.3$: $\mathbb{V}[\mathbf{X}\boldsymbol{\beta}] = 0.15$; $\mathbb{V}[\mathbf{W}\boldsymbol{\theta}] = 0$; and $\mathbb{V}[\mathbf{Z}\boldsymbol{\gamma}] = 0.15$;

- $H^2 = 0.6$: $\mathbb{V}[\mathbf{X}\boldsymbol{\beta}] = 0.3$; $\mathbb{V}[\mathbf{W}\boldsymbol{\theta}] = 0$; and $\mathbb{V}[\mathbf{Z}\boldsymbol{\gamma}] = 0.3$.

Note that, for each case, we generate summary statistics in two ways: (i) including the top 10 PCs as covariates in the marginal linear model to correct for population structure and (ii) not correcting for any population structure. Again all figures show the mean performances (and standard errors) across 100 simulated replicates.

Sparse simulation study design with additive effects. In this set of simulations, we consider phenotypes with sparse architectures [20]. Here, traits were simulated with solely additive effects such that $\mathbb{V}[\mathbf{X}\boldsymbol{\beta}] = H^2$, but this time only variants with the top or bottom $\{1, 5, 10, 25, 50, 100\}$ percentile of LD scores were given nonzero coefficients (a similar simulation approach was also previously implemented in both Bulik-Sullivan et al. [1] and Lee et al. [67]). We once again generate traits with heritability $H^2 = \{0.3, 0.6\}$. We also want to note that, in each of these specific analyses, synthetic trait architectures were generated using all UK Biobank genotyped variants that passed initial preprocessing and quality control (see next section). Since not all of these SNPs are HapMap3 SNPs, some variants were omitted from the `LDSC` and `i-LDSC` regression. Overall, as shown in the main text with results taken over 100 replicates, breaking the assumed relationship between LD scores and chi-squared statistics (i.e., that they are generally positively correlated) led to unbounded estimates of heritability in all but the (more polygenic) scenario when 100% of SNPs contributed to phenotypic variation.

Polygenic simulations with unobserved additive effects. In this next set of simulations, we consider another extension of the polygenic case where a portion of the variants with only additive genetic effects are not observed due ascertainment or other quality control procedures. It was found in Hemani et al. [39] that an initial set of signals pointing towards evidence of genetic interactions were actually better explained using linear models of unobserved variants in the same haplotype. Here, we test whether the `i-LDSC` framework is prone to overestimate the non-additive genetic variance when additive effects in the same haplotype are not included in the model. In each simulation, we generated haplotypes that each contain 5,000 variants. Next, we select either a single causal variant with only an additive effect or a set of ten causal variants with only additive effects — each having a MAF that is randomly selected between: (i) (0.01, 0.1), (ii) (0.1, 0.2), (iii) (0.2, 0.3), (iv) (0.3, 0.4), and (v) (0.4, 0.5). The corresponding

additive effect size for each causal variant across the haplotype is simulated inversely proportional with its MAF. For this analysis, we measure the difference between `i-LDSC` coefficient estimates when every variant is included in the model versus when the haplotype causal variants are omitted for two different trait architectures with broad-sense heritability set to $H^2 = 0.3$ and 0.6. Differences in the component estimates between the observed and unobserved single additive variant models are shown in Figure 3 – figure supplement 9A and 9B. Similar estimates when the larger number of ten additive variants are unobserved in each haplotype are shown in Figure 3 – figure supplement 9C and 9D. If `i-LDSC` was prone to overestimating the non-additive effects, then the omission of the variants with only significant additive effects would lead to increased estimates of τ and ϑ. However, across a range of generative broad-sense heritabilities and haplotype architectures we observe that estimates of τ and ϑ are robust. Intuitively, this is likely due to the fact that these simulations were done under polygenic trait architectures where, as a result, the omission of a few causal variants with small marginal effect sizes has little impact on the ability to estimate genetic variance.

Polygenic simulations with unobserved interaction effects. In this set of simulations, we extend the polygenic case to a setting where a portion of the variants involved in genetic interactions are unobserved. Similar to the case with unobserved additive effects, the purpose of these simulations is to assess whether the `i-LDSC` framework is prone to false discovery of non-additive genetic variance when causal interacting SNPs are not included during the estimation of GWAS summary statistics. In each simulation, we generated haplotypes that each contain 5,000 variants. Traits were simulated using the generative model in Eq. (35) with both additive and interaction effects such that $\mathbb{V}[\mathbf{X}\boldsymbol{\beta}] + \mathbb{V}[\mathbf{W}\boldsymbol{\theta}] = H^2$. Here, every SNP in the genome had at least a small additive effect with a corresponding effect size that was drawn to be inversely proportional to its MAF. Only 1% or 5% of variants within each haplotype had causal non-zero interaction effects. However, when running `i-LDSC`, only a percentage of the interacting SNPs $\{1\%, 5\%, 10\%, 25\%, or 50\%\}$ were included in the estimation of $\widehat{\vartheta}$. We once again generate traits with heritability $H^2 = \{0.3, 0.6\}$ such that the proportion of genetic variance explained by additive effects was equal to $\rho = \{0.5, 0.8\}$. As with the other simulation scenarios, all synthetic traits were generated using UK Biobank genotyped variants that passed initial preprocessing and quality control (see next section). Since not all of these SNPs are HapMap3 SNPs, some variants were omitted from the `i-LDSC` regression analyses. Overall, as discussed in the main text with results taken over 100 replicates, `i-LDSC`

underestimated values of $\widehat{\vartheta}$ when there were unobserved interacting variants (see Figure 3 – figure supplement 10 and 11). As expected, estimates of the additive variance component $\widehat{\tau}$, on the other hand, were not affected.

Polygenic simulations with correlated additive and interaction effects. In our last set of simulations, we sought out to better understand how the relationship between the additive (β) and interaction (θ) coefficients in the generative model of complex traits could potentially bias the additive and non-additive variance component estimates in LDSC and i-LDSC. To that end, we performed a set of simulations where we varied the correlation between the set of effects. Specifically, we first drew a set of additive effect sizes for each variant using the MAF-dependent procedure described above (i.e., $\alpha = -1$). We next selected a subset of the causal variants to be in *cis*-interactions. Here, we set the interaction effect sizes to covary with the additive effect size vector in two different ways. In the first, we simply drew the additive and interaction effect sizes from a multivariate normal such that their correlation was equal to $r = \{-1, -0.8, -0.6, \ldots, 0.6, 0.8, 1\}$ (see Figure 3 – figure supplement 12). In the second, we simply amplified the interaction effects to be a linear function $\theta = \beta \times q$ (Figure 3 – figure supplement 13A and 13C) or a squared function $\theta = \beta^{2q}$ (Figure 3 – figure supplement 13B and 13D) of the additive effects where $q = \{0.1, 0.2, \ldots, 0.9, 1\}$. While testing 100 replicates for each value of q, we observed that the mean estimate of genetic variance had a slight upward bias as the correlation between the additive and interaction effect sizes in the generative model increased; however, the distribution of these bias estimates covered zero in the first and third quartiles of all results. We evaluated this behavior for multiple broad-sense heritability levels $H^2 = 0.3$ and 0.6.

Preprocessing for the UK Biobank and BioBank Japan

In order to apply the i-LDSC framework to 25 continuous traits the UK Biobank[68], we first downloaded genotype data for 488,377 individuals in the UK Biobank using the `ukbgene` tool (https://biobank.ctsu.ox.ac.uk/crystal/download.cgi) and converted the genotypes using the provided `ukbconv` tool (https://biobank.ctsu.ox.ac.uk/crystal/refer.cgi?id=149660). Phenotype data for the 25 continuous traits were also downloaded for those same individuals using the `ukbgene` tool. Individuals identified by the UK Biobank as having high heterozygosity, excessive relatedness, or aneuploidy were removed (1,550 individuals). After separating individuals into self-identified ancestral cohorts

using data field `21000`, unrelated individuals were selected by randomly choosing an individual from each pair of related individuals. This resulted in $N = 349,469$ white British individuals to be included in our analysis. We downloaded imputed SNP data from the UK Biobank for all remaining individuals and removed SNPs with an information score below 0.8. Information scores for each SNP are provided by the UK Biobank (`http://biobank.ctsu.ox.ac.uk/crystal/refer.cgi?id=1967`).

Quality control for the remaining genotyped and imputed variants was then performed on each cohort separately using the following steps. All structural variants were first removed, leaving only single nucleotide polymorphisms (SNPs) in the genotype data. Next, all AT/CG SNPs were removed to avoid possible confounding due to sequencing errors. Then, SNPs with minor allele frequency less than 1% were removed using the PLINK 2.0[69] command `--maf 0.01`. We then removed all SNPs found to be out of Hardy-Weinberg equilibrium, using the PLINK `--hwe 0.000001` flag to remove all SNPs with a Fisher's exact test P-value $> 10^{-6}$. Finally, all SNPs with missingness greater than 1% were removed using the PLINK `--mind 0.01` flag.

We then performed a genome-wide association study (GWAS) for each trait in the UK Biobank on the remaining 8,981,412 SNPs. SNP-level GWAS effect sizes were calculated using PLINK and the `--glm` flag[69]. Age, sex, and the first twenty principal components were included as covariates for all traits analyzed[66]. Principal component analysis was performed using FlashPCA 2.0[70] on a set of independent markers derived separately for each ancestry cohort using the PLINK command `--indep-pairwise 100 10 0.1`. Using the parameters `--indep-pairwise` removes all SNPs that have a pairwise correlation above 0.1 within a 100 SNP window, then slides forward in increments of ten SNPs genome-wide.

In order to analyze data from BioBank Japan, we downloaded publicly available GWAS summary statistics for the 25 traits listed in Table 5 from `http://jenger.riken.jp/en/result`. Summary statistics used age, sex, and the first ten principal components as confounders in the initial GWAS study. We then used individuals from the East Asian (EAS) superpopulation from the 1000 Genomes Project Phase 3 to calculate paired `LDSC` and `i-LDSC` scores from a reference panel. We pruned the reference panel using the PLINK command `--indep-pairwise 100 10 0.5` to limit the computational time of calculating scores[69]. This resulted in reference scores for 1,164,666 SNPs that are included on the `i-LDSC` GitHub repository (see URLs). Using summary statistics from BioBank Japan, with scores calculated from the EAS population in the 1000 Genomes, we obtained `i-LDSC` heritability estimates for each of the 25 traits.

Data and software availability

Source code and tutorials for implementing interaction-LD score regression via the `i-LDSC` package are written in Python and are publicly available online at `https://github.com/lcrawlab/i-LDSC`. Files of LD scores, *cis*-interaction LD scores, and GWAS summary statistics used for our analyses of the UK Biobank and BioBank Japan can be downloaded from the Harvard Dataverse (`https://dataverse.harvard.edu/datsset.xhtml?persistentId=doi:10.7910/DVN/W6MA8J&faces-redirect=true`). All software for the traditional and stratified LD score regression framework with `LDSC` and `s-LDSC` were fit using the default settings, unless otherwise stated in the main text. Source code for these approaches was downloaded from `https://github.com/bulik/ldsc`. When applying `s-LDSC`, we used 97 functional annotations from Gazal et al.[41] to estimate heritability. Data from the UK Biobank Resource[68] (`https://www.ukbiobank.ac.uk`) was made available under Application Numbers 14649 and 22419. Data can be accessed by direct application to the UK Biobank.

Figures and Tables

Figure 1. Power of the `i-LDSC` framework to detect tagged pairwise genetic interaction effects on simulated data. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). Next, we randomly select two groups of interacting variants and divide them into two groups. The group #1 SNPs are chosen to be 1%, 5%, and 10% of the total number of SNPs genome-wide (see the x-axis in each panel). These interact with the group #2 SNPs which are selected to be variants within a ±10 kilobase (kb) window around each SNP in group #1. Coefficients for additive and interaction effects were simulated with no minor allele frequency dependency $\alpha = 0$ (see Materials and Methods). Panels **(A)** and **(B)** are results with simulations using a heritability $H^2 = 0.3$, while panels **(C)** and **(D)** were generated with $H^2 = 0.6$. We also varied the proportion of heritability contributed by additive effects to **(A, C)** $\rho = 0.5$ and **(B, D)** $\rho = 0.8$, respectively. Here, we are blind to the parameter settings used in generative model and run `i-LDSC` while computing the *cis*-interaction LD scores using different estimating windows of ±5 (green), ±10 (orange), ±25 (purple), and ±50 (pink) SNPs. Results are based on 100 simulations per parameter combination and the horizontal bars represent standard errors. Generally, the performance of `i-LDSC` increases with larger heritability and lower proportions of additive variation. Note that `LDSC` is not shown here because it does not search for tagged interaction effects in summary statistics.

Figure 1 – figure supplement 1. Power calculations for the `i-LDSC` framework to detect tagged pairwise genetic interaction effects on simulated data using a ±10 kilobase (kb) window to generate *cis*-interactions around a focal SNP with a moderate minor allele frequency dependency $\alpha = -0.5$ for effect sizes. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). Next, we randomly select two groups of interacting variants and divide them into two groups. The group #1 SNPs are chosen to be 1%, 5%, and 10% of the total number of SNPs genome-wide (see the x-axis in each panel). These interact with the group #2 SNPs which are selected to be variants within a ±10 kilobase (kb) window around each SNP in group #1. Coefficients for additive and interaction effects were simulated with minor allele frequency dependency $\alpha = -0.5$ (see Materials and Methods). Panels **(A)** and **(B)** are results of simulations where the total heritability explained by additive SNP effects and *cis*-interaction effects is $H^2 = 0.3$, while panels **(C)** and **(D)** were generated with $H^2 = 0.6$. We also varied the proportion of phenotypic variation explained by additive SNP effects to **(A, C)** $\rho = 0.5$ and **(B, D)** $\rho = 0.8$, respectively. Here, we are blind to the parameter settings used in generative model and run `i-LDSC` while computing the *cis*-interaction LD scores using different estimation windows of ±5 (green), ±10 (orange), ±25 (purple), and ±50 (pink) SNPs. Results are based on 100 simulations per parameter combination and the horizontal black bars represent standard errors.

Figure 1 – figure supplement 2. Power calculations for the `i-LDSC` framework to detect tagged pairwise genetic interaction effects on simulated data using a ±10 kilobase (kb) window to generate *cis*-interactions around a focal SNP with a strong minor allele frequency dependency $\alpha = -1$ for effect sizes. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). Next, we randomly select two groups of interacting variants and divide them into two groups. The group #1 SNPs are chosen to be 1%, 5%, and 10% of the total number of SNPs genome-wide (see the x-axis in each panel). These interact with the group #2 SNPs which are selected to be variants within a ±10 kilobase (kb) window around each SNP in group #1. Coefficients for additive and interaction effects were simulated with minor allele frequency dependency $\alpha = -0.5$ (see Materials and Methods). Panels **(A)** and **(B)** are results of simulations where the total heritability explained by additive SNP effects and *cis*-interaction effects is $H^2 = 0.3$, while panels **(C)** and **(D)** were generated with $H^2 = 0.6$. We also varied the proportion of phenotypic variation explained by additive SNP effects to **(A, C)** $\rho = 0.5$ and **(B, D)** $\rho = 0.8$, respectively. Here, we are blind to the parameter settings used in generative model and run `i-LDSC` while computing the *cis*-interaction LD scores using different estimation windows of ±5 (green), ±10 (orange), ±25 (purple), and ±50 (pink) SNPs. Results are based on 100 simulations per parameter combination and the horizontal black bars represent standard errors.

Figure 1 – figure supplement 3. Power calculations for the `i-LDSC` framework to detect tagged pairwise genetic interaction effects on simulated data using a ±10 kilobase (kb) window to generate *cis*-interactions around a focal SNP with no minor allele frequency dependency $\alpha = 0$ for effect sizes. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). Next, we randomly select two groups of interacting variants and divide them into two groups. The group #1 SNPs are chosen to be 1%, 5%, and 10% of the total number of SNPs genome-wide (see the x-axis in each panel). These interact with the group #2 SNPs which are selected to be variants within a ±10 kilobase (kb) window around each SNP in group #1. Coefficients for additive and interaction effects were simulated with minor allele frequency dependency $\alpha = -0.5$ (see Materials and Methods). Panels **(A)** and **(B)** are results of simulations where the total heritability explained by additive SNP effects and *cis*-interaction effects is $H^2 = 0.3$, while panels **(C)** and **(D)** were generated with $H^2 = 0.6$. We also varied the proportion of phenotypic variation explained by additive SNP effects to **(A, C)** $\rho = 0.5$ and **(B, D)** $\rho = 0.8$, respectively. Here, we are blind to the parameter settings used in generative model and run `i-LDSC` while computing the *cis*-interaction LD scores using different estimation windows of ±5 (green), ±10 (orange), ±25 (purple), and ±50 (pink) SNPs. Results are based on 100 simulations per parameter combination and the horizontal black bars represent standard errors.

Figure 1 – figure supplement 4. Power calculations for the `i-LDSC` framework to detect tagged pairwise genetic interaction effects on simulated data using a ±100 kilobase (kb) window to generate *cis*-interactions around a focal SNP with a moderate minor allele frequency dependency $\alpha = -0.5$ for effect sizes. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). Next, we randomly select two groups of interacting variants and divide them into two groups. The group #1 SNPs are chosen to be 1%, 5%, and 10% of the total number of SNPs genome-wide (see the x-axis in each panel). These interact with the group #2 SNPs which are selected to be variants within a ±10 kilobase (kb) window around each SNP in group #1. Coefficients for additive and interaction effects were simulated with minor allele frequency dependency $\alpha = -0.5$ (see Materials and Methods). Panels **(A)** and **(B)** are results of simulations where the total heritability explained by additive SNP effects and *cis*-interaction effects is $H^2 = 0.3$, while panels **(C)** and **(D)** were generated with $H^2 = 0.6$. We also varied the proportion of phenotypic variation explained by additive SNP effects to **(A, C)** $\rho = 0.5$ and **(B, D)** $\rho = 0.8$, respectively. Here, we are blind to the parameter settings used in generative model and run `i-LDSC` while computing the *cis*-interaction LD scores using different estimation windows of ±5 (green), ±10 (orange), ±25 (purple), and ±50 (pink) SNPs. Results are based on 100 simulations per parameter combination and the horizontal black bars represent standard errors.

Figure 1 – figure supplement 5. Power calculations for the `i-LDSC` framework to detect tagged pairwise genetic interaction effects on simulated data using a ±100 kilobase (kb) window to generate *cis*-interactions around a focal SNP with a strong minor allele frequency dependency $\alpha = -1$ for effect sizes. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). Next, we randomly select two groups of interacting variants and divide them into two groups. The group #1 SNPs are chosen to be 1%, 5%, and 10% of the total number of SNPs genome-wide (see the x-axis in each panel). These interact with the group #2 SNPs which are selected to be variants within a ±10 kilobase (kb) window around each SNP in group #1. Coefficients for additive and interaction effects were simulated with minor allele frequency dependency $\alpha = -0.5$ (see Materials and Methods). Panels **(A)** and **(B)** are results of simulations where the total heritability explained by additive SNP effects and *cis*-interaction effects is $H^2 = 0.3$, while panels **(C)** and **(D)** were generated with $H^2 = 0.6$. We also varied the proportion of phenotypic variation explained by additive SNP effects to **(A, C)** $\rho = 0.5$ and **(B, D)** $\rho = 0.8$, respectively. Here, we are blind to the parameter settings used in generative model and run `i-LDSC` while computing the *cis*-interaction LD scores using different estimation windows of ±5 (green), ±10 (orange), ±25 (purple), and ±50 (pink) SNPs. Results are based on 100 simulations per parameter combination and the horizontal black bars represent standard errors.

Figure 2. The i-LDSC framework is well-calibrated under the null hypothesis and does not identify evidence of tagged non-additive effects when polygenic traits are generated by only additive effects. In these simulations, synthetic trait architecture is made up of only additive genetic variation (i.e., $\rho = 1$). Coefficients for additive and interaction effects were simulated with no minor allele frequency dependency $\alpha = 0$ (see Materials and Methods). Here, we are blind to the parameter settings used in generative model and run i-LDSC while computing the *cis*-interaction LD scores using different estimating windows of ± 5 (green), ± 10 (orange), ± 25 (purple), and ± 50 (pink) SNPs. **(A)** Mean type I error rate using the i-LDSC framework across an array of estimation window sizes for the *cis*-interaction LD scores. This is determined by assessing the P-value of the *cis*-interaction coefficient (ϑ) in the i-LDSC regression model and checking whether $P < 0.05$. **(B)** Estimates of the *cis*-interaction coefficient (ϑ). Since traits were simulated with only additive effects, these estimates should be centered around zero. **(C)** Estimates of the proportions of phenotypic variance explained (PVE) by genetic effects (i.e., estimated heritability) where the true additive variance is set to $H^2\rho = 0.6$. **(D)** QQ-plot of the P-values for the *cis*-interaction coefficient (ϑ) in i-LDSC. Results are based on 100 simulations per parameter combination and the horizontal bars represent standard errors.

Figure 2 – figure supplement 1. The `i-LDSC` framework is well-calibrated under the null hypothesis and does not identify evidence of tagged non-additive effects when polygenic traits are generated by only additive effects and a moderate minor allele frequency dependency $\alpha = -0.5$ for effect sizes. In these simulations, synthetic trait architecture is made up of only additive genetic variation (i.e., $\rho = 1$). Coefficients for additive and interaction effects were simulated with minor allele frequency dependency $\alpha = -0.5$ (see Materials and Methods). Here, we are blind to the parameter settings used in generative model and run `i-LDSC` while computing the *cis*-interaction LD scores using different estimation windows of ± 5 (green), ± 10 (orange), ± 25 (purple), and ± 50 (pink) SNPs. **(A)** Mean type I error rate using the `i-LDSC` framework across an array of estimation window sizes for the *cis*-interaction LD scores. This is determined by assessing the P-value of the *cis*-interaction coefficient (ϑ) in the `i-LDSC` regression model and checking whether $P < 0.05$. **(B)** Estimates of the *cis*-interaction coefficient (ϑ). Since traits were simulated with only additive effects, these estimates should be centered around zero. **(C)** Estimates of the proportions of phenotypic variance explained (PVE) by genetic effects (i.e., estimated heritability) where the true additive variance is set to $H^2\rho = 0.6$. **(D)** QQ-plot of the P-values for the *cis*-interaction coefficient (ϑ) in `i-LDSC`. Results are based on 100 simulations per parameter combination and the horizontal black bars represent standard errors.

Figure 2 – figure supplement 2. The `i`-LDSC framework is well-calibrated under the null hypothesis and does not identify evidence of tagged non-additive effects when polygenic traits are generated by only additive effects and a strong minor allele frequency dependency $\alpha = -1$ for effect sizes. In these simulations, synthetic trait architecture is made up of only additive genetic variation (i.e., $\rho = 1$). Coefficients for additive and interaction effects were simulated with minor allele frequency dependency $\alpha = -0.5$ (see Materials and Methods). Here, we are blind to the parameter settings used in generative model and run `i`-LDSC while computing the *cis*-interaction LD scores using different estimation windows of ± 5 (green), ± 10 (orange), ± 25 (purple), and ± 50 (pink) SNPs. **(A)** Mean type I error rate using the `i`-LDSC framework across an array of estimation window sizes for the *cis*-interaction LD scores. This is determined by assessing the P-value of the *cis*-interaction coefficient (ϑ) in the `i`-LDSC regression model and checking whether $P < 0.05$. **(B)** Estimates of the *cis*-interaction coefficient (ϑ). Since traits were simulated with only additive effects, these estimates should be centered around zero. **(C)** Estimates of the proportions of phenotypic variance explained (PVE) by genetic effects (i.e., estimated heritability) where the true additive variance is set to $H^2\rho = 0.6$. **(D)** QQ-plot of the P-values for the *cis*-interaction coefficient (ϑ) in `i`-LDSC. Results are based on 100 simulations per parameter combination and the horizontal black bars represent standard errors.

Figure 3. `i-LDSC` robustly and accurately estimates the proportions of phenotypic variance explained (PVE) by genetic effects (i.e., estimated heritability) in simulations in polygenic traits, compared to `LDSC`, due to our accounting for interaction effects tagged in additive GWAS summary statistics. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank (Materials and Methods). All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). Next, we randomly select two groups of interacting variants and divide them into two groups. The group #1 SNPs are chosen to be 10% of the total number of SNPs genome-wide. These interact with the group #2 SNPs which are selected to be variants within a ±100 kilobase (kb) window around each SNP in group #1. Coefficients for additive and interaction effects were simulated with no minor allele frequency dependency $\alpha = 0$ (see Materials and Methods). Here, we assume a heritability **(A)** $H^2 = 0.3$ or **(B)** $H^2 = 0.6$ (marked by the black dotted lines, respectively), and we vary the proportion contributed by additive effects with $\rho = \{0.2, 0.4, 0.6, 0.8\}$. The grey dotted lines represent the total contribution of additive effects in the generative model for the synthetic traits ($H^2\rho$). `i-LDSC` outperforms `LDSC` in recovering heritability across each scenario. Results are based on 100 simulations per parameter combination.

Figure 3 – figure supplement 1. i-LDSC robustly and accurately estimates the proportions of phenotypic variance explained (PVE) by genetic effects in polygenic traits by accounting for interaction effects tagged by GWAS summary statistics. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). Next, we randomly select two groups of interacting variants and divide them into two groups. The group #1 SNPs are chosen to be 10% of the total number of SNPs genome-wide. These interact with the group #2 SNPs which are selected to be variants within a ± 100 kilobase (kb) window around each SNP in group #1. Coefficients for additive and *cis*-interaction effects were simulated with no minor allele frequency dependency $\alpha = 0$ (see Materials and Methods). Here, we assume a total heritability explained by additive SNP and *cis*-interaction effects is **(A)** $H^2 = 0.3$ or **(B)** $H^2 = 0.6$ (marked by the black dotted lines, respectively), and we vary the proportion contributed by additive effects with $\rho = \{0.2, 0.4, 0.6, 0.8\}$. The grey dotted line represents the total contribution of additive effects in the generative model for the synthetic traits ($H^2\rho$). We run i-LDSC while computing the *cis*-interaction LD scores using different estimating windows of ± 5, ± 10, ± 25, and ± 50 SNPs, respectively. These results help motivate the selection of scores calculated using a ± 50 SNP window in our empirical analyses. Results are based on 100 simulations per parameter combination.

Figure 3 – figure supplement 2. Performance of `LDSC` and `i-LDSC` on simulated polygenic traits with architectures that are determined by additive, *cis*-interaction, and gene-by-environment (G×E) effects. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). Next, we randomly select two groups of interacting variants and divide them into two groups. The group #1 SNPs are chosen to be 10% of the total number of SNPs genome-wide. These interact with the group #2 SNPs which are selected to be variants within a ±100 kilobase (kb) window around each SNP in group #1. G×E effects were simulated using an amplification model[65] (see Materials and Methods) where we split the sample population in half to emulate two subsets of individuals coming from different environments. We randomly draw variant effect sizes for the first environment from a standard Gaussian distribution. Then effect sizes for the second environment are set to be the product of the effect sizes in from with first environment with an amplifier $w = [1.1, 1.2, \ldots, 2]$ (see the x-axis in each panel). Both the *cis*-interaction and G×E effects were set to explain a quarter of the total phenotypic variation and the remaining half was explained by additive SNP effects. Panels **(A)** and **(B)** show estimates of the proportions of phenotypic variance explained (PVE) by genetic effects (i.e., estimated heritability) from `LDSC` and `i-LDSC`, respectively. Panels **(C)** and **(D)** show `i-LDSC` estimates of the phenotypic variation explained by tagged non-additive genetic effects using the *cis*-interaction LD score (i.e., estimates of ϑ). We assume the total heritability explained by all genetic effects to be **(A, C)** $H^2 = 0.6$ and **(B, D)** $H^2 = 0.3$. Results are based on 100 simulations per parameter combination.

Figure 3 – figure supplement 3. Performance of LDSC and i-LDSC on simulated polygenic traits with architectures that are determined by additive, *cis*-interaction, and gene-by-ancestry (G×Ancestry) effects with principal components (PCs) included in the GWAS model to correct for additional structure. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). Next, we randomly select two groups of interacting variants and divide them into two groups. The group #1 SNPs are chosen to be 10% of the total number of SNPs genome-wide. These interact with the group #2 SNPs which are selected to be variants within a ±100 kilobase (kb) window around each SNP in group #1. G×Ancestry effects were simulated as the product of individual genotypes and the SNP loadings for each of the first 10 PCs (see the x-axis in each panel). Both the *cis*-interaction and G×Ancestry effects were set to explain a quarter of the total phenotypic variation and the remaining half was explained by additive SNP effects. The proportion of genotypic variance explained by each PC is shown in green. Panels **(A)** and **(B)** show estimates of the proportions of phenotypic variance explained (PVE) by genetic effects (i.e., estimated heritability) from LDSC and i-LDSC, respectively. Panels **(C)** and **(D)** show i-LDSC estimates of the phenotypic variation explained by tagged non-additive genetic effects using the *cis*-interaction LD score (i.e., estimates of ϑ). We assume the total heritability explained by all genetic effects to be **(A, C)** $H^2 = 0.6$ and **(B, D)** $H^2 = 0.3$. Results are based on 100 simulations per parameter combination.

Figure 3 – figure supplement 4. Performance of `LDSC` and `i-LDSC` on simulated polygenic traits with architectures that are determined by additive, *cis*-interaction, and gene-by-ancestry (G×Ancestry) effects without correcting for the additional structure in the GWAS analysis. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). Next, we randomly select two groups of interacting variants and divide them into two groups. The group #1 SNPs are chosen to be 10% of the total number of SNPs genome-wide. These interact with the group #2 SNPs which are selected to be variants within a ±100 kilobase (kb) window around each SNP in group #1. G×Ancestry effects were simulated as the product of individual genotypes and the SNP loadings for each of the first 10 PCs (see the x-axis in each panel). Both the *cis*-interaction and G×Ancestry effects were set to explain a quarter of the total phenotypic variation and the remaining half was explained by additive SNP effects. The proportion of genotypic variance explained by each PC is shown in green. Panels **(A)** and **(B)** show estimates of the proportions of phenotypic variance explained (PVE) by genetic effects (i.e., estimated heritability) from `LDSC` and `i-LDSC`, respectively. Panels **(C)** and **(D)** show `i-LDSC` estimates of the phenotypic variation explained by tagged non-additive genetic effects using the *cis*-interaction LD score (i.e., estimates of ϑ). We assume the total heritability explained by all genetic effects to be **(A, C)** $H^2 = 0.6$ and **(B, D)** $H^2 = 0.3$. Results are based on 100 simulations per parameter combination.

Figure 3 – figure supplement 5. Performance of `LDSC` and `i-LDSC` on simulated polygenic traits with architectures that are determined by only additive and gene-by-environment (G×E) effects. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). G×E effects were simulated using an amplification model[65] (see Materials and Methods) where we split the sample population in half to emulate two subsets of individuals coming from different environments. We randomly draw variant effect sizes for the first environment from a standard Gaussian distribution. Then effect sizes for the second environment are set to be the product of the effect sizes in from with first environment with an amplifier $w = [1.1, 1.2, \ldots, 2]$ (see the x-axis in each panel). Additive and G×E effects were set to explain half of the phenotypic variation. Note that unlike results depicted in Figure 3 – figure supplement 2, there are no *cis*-interaction effects that affect trait architecture. Here, panels **(A)** and **(B)** show estimates of the proportions of phenotypic variance explained (PVE) by genetic effects (i.e., estimated heritability) from `LDSC` and `i-LDSC`, respectively. Panels **(C)** and **(D)** show `i-LDSC` estimates of the phenotypic variation explained by tagged non-additive genetic effects using the *cis*-interaction LD score (i.e., estimates of ϑ). We assume the total heritability explained by all genetic effects to be **(A, C)** $H^2 = 0.6$ and **(B, D)** $H^2 = 0.3$. Results are based on 100 simulations per parameter combination.

Figure 3 – figure supplement 6. Performance of `LDSC` and `i-LDSC` on simulated polygenic traits with architectures that are determined by only additive and gene-by-ancestry (G×Ancestry) effects with principal components (PCs) included in the GWAS model to correct for additional structure. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). G×Ancestry effects were simulated as the product of individual genotypes and the SNP loadings for each of the first 10 PCs (see the x-axis in each panel). Additive and G×E effects were set to explain half of the phenotypic variation. The proportion of genotypic variance explained by each PC is shown in green. Note that unlike results depicted in Figure 3 – figure supplement 3, there are no *cis*-interaction effects that affect trait architecture. Here, panels **(A)** and **(B)** show estimates of the proportions of phenotypic variance explained (PVE) by genetic effects (i.e., estimated heritability) from `LDSC` and `i-LDSC`, respectively. Panels **(C)** and **(D)** show `i-LDSC` estimates of the phenotypic variation explained by tagged non-additive genetic effects using the *cis*-interaction LD score (i.e., estimates of ϑ). We assume the total heritability explained by all genetic effects to be **(A, C)** $H^2 = 0.6$ and **(B, D)** $H^2 = 0.3$. Results are based on 100 simulations per parameter combination.

Figure 3 – figure supplement 7. Performance of `LDSC` and `i-LDSC` on simulated polygenic traits with architectures that are determined by only additive and gene-by-ancestry (G×Ancestry) effects without correcting for the additional structure in the GWAS analysis. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). G×Ancestry effects were simulated as the product of individual genotypes and the SNP loadings for each of the first 10 PCs (see the x-axis in each panel). Additive and G×E effects were set to explain half of the phenotypic variation. The proportion of genotypic variance explained by each PC is shown in green. Note that unlike results depicted in Figure 3 – figure supplement 4, there are no *cis*-interaction effects that affect trait architecture. Here, panels **(A)** and **(B)** show estimates of the proportions of phenotypic variance explained (PVE) by genetic effects (i.e., estimated heritability) from `LDSC` and `i-LDSC`, respectively. Panels **(C)** and **(D)** show `i-LDSC` estimates of the phenotypic variation explained by tagged non-additive genetic effects using the *cis*-interaction LD score (i.e., estimates of ϑ). We assume the total heritability explained by all genetic effects to be **(A, C)** $H^2 = 0.6$ and **(B, D)** $H^2 = 0.3$. Results are based on 100 simulations per parameter combination.

Figure 3 – figure supplement 8. Performance of `LDSC` and `i-LDSC` on simulated traits with sparse architectures that are determined by only additive effects. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. Here, traits were generated with solely additive effects where only variants with the top or bottom $\{1, 5, 10, 25, 50, 100\}$ percentile of LD scores were given nonzero coefficients in the generative model (see the x-axis in each panel). Panels **(A)** and **(B)** show estimates of the proportions of phenotypic variance explained (PVE) by genetic effects (i.e., estimated heritability) from `LDSC` and `i-LDSC`, respectively. Panels **(C)** and **(D)** show `i-LDSC` estimates of the phenotypic variation explained by tagged non-additive genetic effects using the *cis*-interaction LD score (i.e., estimates of ϑ). We assume the total heritability explained by all genetic effects to be **(A, C)** $H^2 = 0.6$ and **(B, D)** $H^2 = 0.3$. Results are based on 100 simulations per parameter combination. The overall takeaway is that breaking the assumed relationship between LD scores and chi-squared test statistics (i.e., that they are generally positively correlated) led to unbounded estimates of heritability for both `LDSC` and `i-LDSC` in all but the (polygenic) scenario when 100% of SNPs contributed to phenotypic variation.

Figure 3 – figure supplement 9. The non-additive component estimates in `i-LDSC` are robust to unobserved additive effects in a haplotype. Synthetic trait architectures are simulated such that a substantial proportion of genetic variance is explained by an additive effect that is not directly observed. The goal of these simulations was to assess how these unobserved effects influence the estimation of the non-additive variance component in the `i-LDSC` model. In each simulation, we generated haplotypes that each contain 5,000 variants. Next, we select either **(A, B)** a single causal variant with only an additive effect or **(C, D)** a set of ten causal variants with only additive effects. In each case, the causal variants have a MAF that is randomly selected between: (*i*) (0.01, 0.1), (*ii*) (0.1, 0.2), (*iii*) (0.2, 0.3), (*iv*) (0.3, 0.4), or (*v*) (0.4, 0.5) as depicted on the x-axis. The corresponding additive effect size for each causal variant across the haplotypes is simulated to be inversely proportional to its MAF[38]. On the y-axis, we measure the difference (Δ) between `i-LDSC` coefficient estimates when every variant is included in the model versus when the haplotype causal variants are omitted for two different trait architectures with broad-sense heritability set to **(A, C)** $H^2 = 0.6$ and **(B, D)** $H^2 = 0.3$. Results are based on 100 simulations per parameter combination.

Figure 3 – figure supplement 10. The `i-LDSC` framework protects against the false discovery of non-additive genetic variance when causal interacting SNPs are unobserved and the proportion of genetic variance explained by additive effects is equal to $\rho = 0.5$. Synthetic trait architectures are simulated such that a substantial proportion of genetic variance is explained by pairwise genetic interaction effects that are not directly observed. The goal of these simulations was to assess how these unobserved effects influence the estimation of the non-additive variance component in the `i-LDSC` model. In each simulation, we generated haplotypes that each contain 5,000 variants. Every SNP in the genome had at least a small additive effect. The corresponding additive effect size for each variant across the haplotypes is simulated to be inversely proportional to its MAF[38]. We then set **(A, C)** 1% or **(B, D)** 5% of causal variants in each haplotype to have non-zero interaction effects. On the y-axis, we measure the difference (Δ) between `i-LDSC` coefficient estimates when every variant is included in the model versus when the specified percentage of variants with pairwise genetic interaction effects are omitted for two different trait architectures with broad-sense heritability set to **(A, B)** $H^2 = 0.6$ and **(C, D)** $H^2 = 0.3$. Results are based on 100 simulations per parameter combination.

Figure 3 – figure supplement 11. The `i-LDSC` framework protects against the false discovery of non-additive genetic variance when causal interacting SNPs are unobserved and the proportion of genetic variance explained by additive effects is equal to $\rho = 0.8$. Synthetic trait architectures are simulated such that a substantial proportion of genetic variance is explained by pairwise genetic interaction effects that are not directly observed. The goal of these simulations was to assess how these unobserved effects influence the estimation of the non-additive variance component in the `i-LDSC` model. In each simulation, we generated haplotypes that each contain 5,000 variants. Every SNP in the genome had at least a small additive effect. The corresponding additive effect size for each variant across the haplotypes is simulated to be inversely proportional to its MAF [38]. We then set **(A, C)** 1% or **(B, D)** 5% of causal variants in each haplotype to have non-zero interaction effects. On the y-axis, we measure the difference (Δ) between `i-LDSC` coefficient estimates when every variant is included in the model versus when the specified percentage of variants with pairwise genetic interaction effects are omitted for two different trait architectures with broad-sense heritability set to **(A, B)** $H^2 = 0.6$ and **(C, D)** $H^2 = 0.3$. Results are based on 100 simulations per parameter combination.

Figure 3 – figure supplement 12. Bias in `LDSC` and `i-LDSC` estimates when the additive and interaction effect sizes in the generative model of complex traits are correlated. To simulate synthetic trait architectures, we first simulated additive effects for each variant to be MAF-dependent (i.e., $\alpha = -1$). Here, we set the corresponding interaction effect sizes to have a correlation with the additive effect sizes equal to $r = \{-1, -0.8, -0.6, \ldots, 0.6, 0.8, 1\}$ (labeled across the x-axis). On the y-axis, we measure the bias in the `LDSC` and `i-LDSC` estimates of phenotypic variance explained (PVE) by genetic effects. In each simulation, we generate traits with an equal proportion of variance explained by additive and interaction effects and a total broad-sense heritability set to **(A)** $H^2 = 0.6$ and **(B)** $H^2 = 0.3$. Results are based on 100 simulations for each parameter value.

Figure 3 – figure supplement 13. Bias in `LDSC` and `i-LDSC` estimates when interaction effect sizes in the generative model of complex traits are a linear or squared function of the the additive effects. To simulate synthetic trait architectures, we first simulated additive effects for each variant to be MAF-dependent (i.e., $\alpha = -1$). Here, we set the corresponding interaction effect sizes to be either **(A, C)** a linear function or **(B, D)** a squared function of the additive effects with a scaling factor $q = \{0.1, 0.2, \ldots, 0.8, 1\}$ (labeled across the x-axis). On the y-axis, we measure the bias in the `LDSC` and `i-LDSC` estimates of the phenotypic variance explained (PVE) by genetic effects. In each simulation, we generate traits with an equal proportion of variance explained by additive and interaction effects and a total broad-sense heritability set to **(A, B)** $H^2 = 0.6$ and **(C, D)** $H^2 = 0.3$. Results are based on 100 simulations for each parameter value.

Figure 4. The i-LDSC framework recovers heritability and provides estimates of tagged *cis*-interactions in GWAS summary statistics (ϑ) for 25 quantitiative traits in the UK Biobank and BioBank Japan. **(A)** In both the UK Biobank (green) and BioBank Japan (purple), estimates of phenotypic variance explained (PVE) by genetic effects from i-LDSC and LDSC are highly correlated for 25 different complex traits. The Spearman correlation coefficient between heritability estimates from LDSC and i-LDSC for the UK Biobank and BioBank Japan are $r^2 = 0.989$ and $r^2 = 0.850$, respectively. The $y = x$ dotted line represents the values at which estimates from both approaches are the same. **(B)** PVE estimates from the UK Biobank are better correlated with those from the BioBank Japan across 25 traits using LDSC (Spearman $r^2 = 0.848$) than i-LDSC (Spearman $r^2 = 0.666$). **(C)** Both the original and stratified LDSC models recover the same amount of PVE when the *cis*-interaction LD score is included as an additional component in the UK Biobank analysis (Spearman $r^2 = 0.989$). These models are listed as i-LDSC and s+i-LDSC, respectively. For s+i-LDSC, we included 97 functional annotations from Gazal et al. [41] to estimate heritability. **(D)** Estimates of non-additive variance components in i-LDSC versus s+i-LDSC (Spearmen $r^2 = 0.184$). While not statistically significant in the stratified analysis with the additional annotations, the non-additive component still makes nonzero contributions to the PVE estimation for all 25 traits in the UK Biobank (see Tables 1 and 2).

Figure 4 – figure supplement 1. Additional results from applying `LDSC` and `i-LDSC` for 25 quantitiative traits in the UK Biobank and BioBank Japan. (A) `i-LDSC` estimates of the phenotypic variation explained by tagged non-additive genetic effects using the *cis*-interaction LD score (i.e., estimates of ϑ) between traits in the UK Biobank and BioBank Japan (Spearman $r^2 = 0.372$). **(B)** Estimates of `i-LDSC` and `LDSC` intercept terms for 25 traits analyzed in the UK Biobank and BioBank Japan. Intercept terms using `LDSC` and `i-LDSC` are highly correlated in both the UK Biobank (Spearman $r^2 = 0.888$) and BioBank Japan (Spearman $r^2 = 0.813$). The $x = y$ dotted line represents points for when the two sets of estimates are equal.

Trait	UKB (LDSC)	UKB (i-LDSC)	UKB $\widehat{\vartheta}$	UKB P-value	BBJ (LDSC)	BBJ (i-LDSC)	BBJ $\widehat{\vartheta}$	BBJ P-value
Basophil	0.0250	0.0315	0.0065	1.572×10^{-12}	0.0684	0.1548	0.0864	0.025
BMI	0.1757	0.2349	0.0592	3.083×10^{-84}	0.1667	0.2656	0.0989	2.438×10^{-18}
Cholesterol	0.0954	0.0974	0.0020	1.821×10^{-16}	0.0629	0.1268	0.0639	2.740×10^{-4}
CRP	0.0354	0.0414	0.0060	9.845×10^{-12}	0.0202	0.1625	0.1423	0.020
DBP	0.0940	0.1203	0.0263	1.118×10^{-65}	0.0605	0.1267	0.0662	1.675×10^{-7}
EGFR	0.1521	0.1999	0.0478	1.187×10^{-46}	0.1010	0.1225	0.0215	4.232×10^{-5}
Eosinophil	0.1055	0.1375	0.0320	1.230×10^{-18}	0.0785	0.1973	0.1188	0.001
HBA1C	0.0906	0.1083	0.0177	1.578×10^{-26}	0.1057	0.1308	0.0251	0.031
HDL*	0.1599	0.1768	0.0169	9.636×10^{-37}	0.1590	0.1838	0.0248	0.081
Height	0.3675	0.4815	0.1140	1.038×10^{-64}	0.3941	0.7336	0.3395	7.433×10^{-33}
Hematocrit	0.1078	0.1352	0.0274	2.479×10^{-25}	0.0752	0.0928	0.0176	3.689×10^{-5}
Hemoglobin	0.1177	0.1433	0.0256	4.284×10^{-27}	0.0702	0.0752	0.0050	9.037×10^{-4}
LDL	0.0802	0.0859	0.0057	5.087×10^{-13}	0.0745	0.1438	0.0693	0.018
Lymphocyte	0.0402	0.0501	0.0099	4.906×10^{-19}	0.0844	0.1757	0.0913	5.479×10^{-5}
MCH	0.1361	0.1597	0.0236	1.785×10^{-25}	0.1536	0.2831	0.1295	1.042×10^{-5}
MCHC	0.0317	0.0364	0.0047	3.730×10^{-12}	0.0571	0.0650	0.0079	0.027
MCV	0.1630	0.1902	0.0272	1.180×10^{-29}	0.1530	0.2818	0.1288	1.042×10^{-5}
Monocyte	0.0788	0.0955	0.0167	5.257×10^{-18}	0.0888	0.1549	0.0661	0.004
Neutrophil	0.1102	0.1391	0.0289	1.777×10^{-33}	0.1191	0.2114	0.0923	5.050×10^{-5}
Platelet	0.1992	0.2447	0.0455	2.303×10^{-37}	0.1565	0.2436	0.0871	7.724×10^{-9}
RBC	0.1574	0.1933	0.0359	3.292×10^{-31}	0.1203	0.2068	0.0865	5.972×10^{-8}
SBP	0.0954	0.1201	0.0247	8.660×10^{-75}	0.0769	0.1604	0.0835	9.075×10^{-10}
Triglycerides*	0.1061	0.1204	0.0143	1.410×10^{-26}	0.1171	0.2670	0.1499	0.110
Urate	0.1217	0.1550	0.0333	9.642×10^{-38}	0.1395	0.3462	0.2067	0.015
WBC	0.0962	0.1250	0.0288	9.866×10^{-34}	0.1024	0.2266	0.1242	1.346×10^{-8}

Table 1. i-LDSC heritability estimates and P-values highlighting statistically significant contributions of tagged pairwise genetic interaction effects for 25 traits in the UK Biobank and BioBank Japan. Here, LDSC heritability estimates are included as a baseline. The difference between the approaches is that the i-LDSC heritability estimates include proportions of phenotypic variation that are explained by tagged non-additive variation (see columns with estimates of ϑ). Note that all 25 traits analyzed in the UK Biobank and 23 of the 25 traits analyzed in BioBank Japan have a statistically significant amount of tagged non-additive genetic effects as detected by the *cis*-interaction LD score ($P < 0.05$). The two traits without significant tagged non-additive genetic effects in BioBank Japan were HDL ($P = 0.081$) and Triglyceride ($P = 0.110$). These traits are indicated by *. The i-LDSC P-values are related to the estimates of the ϑ coefficients which are also displayed in Figure 4.

Trait	UKB PVE (`s-LDSC`)	UKB PVE (`s+i-LDSC`)	`s+i-LDSC` P-value
Basophil	0.0363	0.0375	0.4728
BMI	0.2100	0.2482	0.8126
Cholesterol	0.1042	0.1358	0.6202
CRP	0.0452	0.0524	0.6483
DBP	0.1228	0.1441	0.6125
EGFR	0.1826	0.2105	0.8507
Eosinophil	0.1403	0.1578	0.1867
HBA1C	0.1040	0.1275	0.6917
HDL	0.1820	0.2373	0.5754
Height	0.4315	0.4726	0.5224
Hematocrit	0.1416	0.1646	0.3956
Hemoglobin	0.1504	0.1795	0.2299
LDL	0.0858	0.1131	0.8812
Lymphocyte	0.0545	0.0651	0.1453
MCH	0.1497	0.1545	0.0968
MCHC	0.0450	0.0496	0.3728
MCV	0.1814	0.1930	0.1530
Monocyte	0.1085	0.1431	0.5421
Neutrophil	0.1320	0.1599	0.2499
Platelet	0.2317	0.2628	0.7371
RBC	0.1933	0.2223	0.3197
SBP	0.1206	0.1419	0.1100
Triglycerides	0.1335	0.1621	0.5301
Urate	0.1530	0.1736	0.1177
WBC	0.1221	0.1482	0.5155

Table 2. Comparison of `s-LDSC` and `i-LDSC` estimates of phenotypic variance explained (PVE) by genetic effects for 25 complex traits in the UK Biobank. Here, we use stratified LD score regression (`s-LDSC`) to partition heritability across different genomic elements[42]. We used 97 functional annotations from Gazal et al.[41] to estimate heritability in 25 traits. We then appended *cis*-interaction LD scores as an additional annotation to obtain heritability estimates (this method is referred to as `s+i-LDSC` in the table). P-values for the `s+i-LDSC` model detailing the contributions of tagged non-additive genetic effects for 25 traits are provided in the last column. Note that, while not statistically significant in this stratified analysis with the additional annotations, the non-additive component still makes nonzero contributions to the PVE estimation for all 25 traits.

 # Supplementary File Captions

Supplementary File 1. Comparison of `LDSC` and `i-LDSC` estimates of the proportion of phenotypic variance explained (PVE) by genetic effects (i.e., estimated heritability) when the true heritability is set to $H^2 = 0.3$ for polygenic traits. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). Next, we randomly select two groups of interacting variants and divide them into two groups. The group #1 SNPs are chosen to be 10% of the total number of SNPs genome-wide. These interact with the group #2 SNPs which are selected to be variants within a ±100 kilobase (kb) window around each SNP in group #1. Coefficients for additive and interaction effects were simulated with no minor allele frequency dependency $\alpha = 0$ (see Materials and Methods). Here, we assume a heritability $H^2 = 0.3$ and vary the proportion contributed by additive effects with $\rho = \{0.2, 0.4, 0.6, 0.8\}$. We run `i-LDSC` while computing the *cis*-interaction LD scores using different estimating windows of ±5, ±10, ±25, and ±50 SNPs. The "average" column represents results using model averaging over the different estimating windows (see Materials and Methods). We report the mean estimates of heritability (with standard errors in the parentheses) and use mean absolute error (MAE) to quantify the difference between the two methods. Results are based on 100 simulations per parameter combination. As shown in Figure 3 – figure supplement 3 and 1, `LDSC` does not capture the contribution of non-additive genetic effects to trait variation.

Supplementary File 2. Comparison of `LDSC` and `i-LDSC` estimates of the proportion of phenotypic variance explained (PVE) by genetic effects (i.e., estimated heritability) when the true heritability is set to $H^2 = 0.6$. Synthetic trait architecture was simulated using real genotype data from individuals of self-identified European ancestry in the UK Biobank. All SNPs were considered to have at least an additive effect (i.e., creating a polygenic trait architecture). Next, we randomly select two groups of interacting variants and divide them into two interacting groups. The group #1 SNPs are chosen to be 10% of the total number of SNPs genome-wide. These interact with the group #2 SNPs which are selected to be variants within a ±100 kilobase (kb) window around each SNP in group #1. Coefficients for additive and interaction effects were simulated with no minor allele frequency dependency $\alpha = 0$ (see Materials and Methods). Here, we assume a heritability $H^2 = 0.6$ and vary the proportion contributed by additive effects with $\rho = \{0.2, 0.4, 0.6, 0.8\}$. We run `i-LDSC` while computing the *cis*-interaction LD scores using different estimating windows of ±5, ±10, ±25, and ±50 SNPs. The "average" column represents results using model averaging over the different estimating windows (see Materials and Methods). We report the mean estimates of heritability (with standard errors in the parentheses) and use mean absolute error (MAE) to quantify the difference between the two methods. Results are based on 100 simulations per parameter combination. As shown in Figure 3 – figure supplement 3 and 1, `LDSC` does not capture the additional contribution of non-additive genetic effects to trait variation.

Supplementary File 3. Abbreviations used throughout this study for 14 quantitative traits analyzed in this study. The remaining 11 traits analyzed were Basophil count, Cholesterol, Eosinophil count, Height, Hematocrit, Hemoglobin, Lymphocyte count, Monocyte count, Neutrophil count, and Triglyceride levels, respectively. These are not abbreviated in the main text.

Supplementary File 4. Trait-specific α parameters for each of the 25 traits analyzed. Here, α values are used to weight each variant based on its minor allele frequency to account for frequency dependent architectures in each trait. The $*$ indicates α parameters that were taken directly from Schoech et al.[38]. The α parameters for other traits were calculated using the protocol used in that paper. Expansion of trait abbreviations are given in Table 3.

Supplementary File 5. Number of individuals and total SNPs included in the analysis of each trait in BioBank Japan.

References

1. Brendan K Bulik-Sullivan, Po-Ru Loh, Hilary K Finucane, Stephan Ripke, Jian Yang, Nick Patterson, Mark J Daly, Alkes L Price, and Benjamin M Neale. Ld score regression distinguishes confounding from polygenicity in genome-wide association studies. Nature genetics, 47(3):291–295, 2015.

2. Brendan Bulik-Sullivan, Hilary K Finucane, Verneri Anttila, Alexander Gusev, Felix R Day, Po-Ru Loh, Laramie Duncan, John R B Perry, Nick Patterson, Elise B Robinson, Mark J Daly, Alkes L Price, Benjamin M Neale, ReproGen Consortium, Psychiatric Genomics Consortium, and Genetic Consortium for Anorexia Nervosa of the Wellcome Trust Case Control Consortium 3. An atlas of genetic correlations across human diseases and traits. Nature Genetics, 47(11):1236–1241, 2015. doi: 10.1038/ng.3406. URL https://doi.org/10.1038/ng.3406.

3. Huwenbo Shi, Gleb Kichaev, and Bogdan Pasaniuc. Contrasting the genetic architecture of 30 complex traits from summary association data. The American Journal of Human Genetics, 99(1): 139–153, 2016.

4. Noah Zaitlen, Peter Kraft, Nick Patterson, Bogdan Pasaniuc, Gaurav Bhatia, Samuela Pollack, and Alkes L. Price. Using extended genealogy to estimate components of heritability for 23 quantitative and dichotomous traits. PLOS Genetics, 9(5):1–11, 2013. doi: 10.1371/journal.pgen.1003520. URL https://doi.org/10.1371/journal.pgen.1003520.

5. Tinca J C Polderman, Beben Benyamin, Christiaan A de Leeuw, Patrick F Sullivan, Arjen van Bochoven, Peter M Visscher, and Danielle Posthuma. Meta-analysis of the heritability of human traits based on fifty years of twin studies. Nature Genetics, 47(7):702–709, 2015. doi: 10.1038/ng. 3285. URL https://doi.org/10.1038/ng.3285.

6. Kangcheng Hou, Kathryn S. Burch, Arunabha Majumdar, Huwenbo Shi, Nicholas Mancuso, Yue Wu, Sriram Sankararaman, and Bogdan Pasaniuc. Accurate estimation of snp-heritability from biobank-scale data irrespective of genetic architecture. Nature Genetics, 51(8):1244–1251, 2019. doi: 10.1038/s41588-019-0465-0. URL https://doi.org/10.1038/s41588-019-0465-0.

7. Ali Pazokitoroudi, Yue Wu, Kathryn S. Burch, Kangcheng Hou, Aaron Zhou, Bogdan Pasaniuc, and Sriram Sankararaman. Efficient variance components analysis across millions of genomes.

Nature Communications, 11(1):4020, 2020. doi: 10.1038/s41467-020-17576-9. URL https://doi.org/10.1038/s41467-020-17576-9.

8. Loic Yengo, Julia Sidorenko, Kathryn E Kemper, Zhili Zheng, Andrew R Wood, Michael N Weedon, Timothy M Frayling, Joel Hirschhorn, Jian Yang, Peter M Visscher, et al. Meta-analysis of genome-wide association studies for height and body mass index in 700000 individuals of european ancestry. Human molecular genetics, 27(20):3641–3649, 2018.

9. Doug Speed and David J Balding. Sumher better estimates the snp heritability of complex traits from summary statistics. Nature genetics, 51(2):277–284, 2019.

10. Shuang Song, Wei Jiang, Yiliang Zhang, Lin Hou, and Hongyu Zhao. Leveraging ld eigenvalue regression to improve the estimation of snp heritability and confounding inflation. The American Journal of Human Genetics, 2022.

11. Duncan S Palmer, Wei Zhou, Liam Abbott, Emilie M Wigdor, Nikolas Baya, Claire Churchhouse, Cotton Seed, Tim Poterba, Daniel King, Masahiro Kanai, et al. Analysis of genetic dominance in the uk biobank. Science, 379(6639):1341–1348, 2023.

12. Tsz Fung Chan, Xinyue Rui, David V. Conti, Myriam Fornage, Mariaelisa Graff, Jeffrey Haessler, Christopher Haiman, Heather M. Highland, Su Yon Jung, Eimear Kenny, Charles Kooperberg, Loic Le Marchland, Kari E. North, Ran Tao, Genevieve Wojcik, Christopher R. Gignoux, PAGE Consortium, Charleston W. K. Chiang, and Nicholas Mancuso. Estimating heritability explained by local ancestry and evaluating stratification bias in admixture mapping from summary statistics. bioRxiv, page 2023.04.10.536252, 2023. doi: 10.1101/2023.04.10.536252. URL http://biorxiv.org/content/early/2023/04/18/2023.04.10.536252.abstract.

13. Zheng Ning, Yudi Pawitan, and Xia Shen. High-definition likelihood inference of genetic correlations across human complex traits. Nature Genetics, 52(8):859–864, 2020. doi: 10.1038/s41588-020-0653-y. URL https://doi.org/10.1038/s41588-020-0653-y.

14. Yiliang Zhang, Qiongshi Lu, Yixuan Ye, Kunling Huang, Wei Liu, Yuchang Wu, Xiaoyuan Zhong, Boyang Li, Zhaolong Yu, Brittany G Travers, Donna M Werling, James J Li, and Hongyu Zhao. Supergnova: local genetic correlation analysis reveals heterogeneous etiologic sharing of complex

traits. Genome Biol, 22(1):262, 2021. ISSN 1474-760X (Electronic); 1474-7596 (Print); 1474-7596 (Linking). doi: 10.1186/s13059-021-02478-w.

15. Roshni A. Patel, Shaila A. Musharoff, Jeffrey P. Spence, Harold Pimentel, Catherine Tcheandjieu, Hakhamanesh Mostafavi, Nasa Sinnott-Armstrong, Shoa L. Clarke, Courtney J. Smith, Peter P. Durda, Kent D. Taylor, Russell Tracy, Yongmei Liu, W. Craig Johnson, Francois Aguet, Kristin G. Ardlie, Stacey Gabriel, Josh Smith, Deborah A. Nickerson, Stephen S. Rich, Jerome I. Rotter, Philip S. Tsao, Themistocles L. Assimes, and Jonathan K. Pritchard. Genetic interactions drive heterogeneity in causal variant effect sizes for gene expression and complex traits. The American Journal of Human Genetics, 2022. doi: 10.1016/j.ajhg.2022.05.014. URL https://doi.org/10.1016/j.ajhg.2022.05.014.

16. Evan E. Eichler, Jonathan Flint, Greg Gibson, Augustine Kong, Suzanne M. Leal, Jason H. Moore, and Joseph H. Nadeau. Missing heritability and strategies for finding the underlying causes of complex disease. Nature Reviews Genetics, 11(6):446–450, 2010. doi: 10.1038/nrg2809. URL https://doi.org/10.1038/nrg2809.

17. Brendan K Bulik-Sullivan, Po-Ru Loh, Hilary K Finucane, Stephan Ripke, Jian Yang, Schizophrenia Working Group of the Psychiatric Genomics Consortium, Nick Patterson, Mark J Daly, Alkes L Price, and Benjamin M Neale. Ld score regression distinguishes confounding from polygenicity in genome-wide association studies. Nat Genet, 47:291–295, 2015. URL http://dx.doi.org/10.1038/ng.3211.

18. Ismo Strandén and Ole F Christensen. Allele coding in genomic evaluation. Genet Sel Evol, 43(1):25, 2011. ISSN 1297-9686 (Electronic); 0999-193X (Print); 0999-193X (Linking). doi: 10.1186/1297-9686-43-25.

19. Gustavo de Los Campos, Ana I Vazquez, Rohan Fernando, Yann C Klimentidis, and Daniel Sorensen. Prediction of complex human traits using the genomic best linear unbiased predictor. PLoS Genet, 9(7):e1003608, 2013. ISSN 1553-7404 (Electronic); 1553-7390 (Print); 1553-7390 (Linking). doi: 10.1371/journal.pgen.1003608.

20. Xiang Zhou, Peter Carbonetto, and Matthew Stephens. Polygenic modeling with Bayesian sparse linear mixed models. PLoS Genet, 9(2):e1003264, 2013.

21. Jian Yang, Beben Benyamin, Brian P McEvoy, Scott Gordon, Anjali K Henders, Dale R Nyholt, Pamela A Madden, Andrew C Heath, Nicholas G Martin, and Grant W Montgomery. Common snps explain a large proportion of the heritability for human height. Nat Genet, 42(7):565, 2010.

22. Michael C Wu, Seunggeun Lee, Tianxi Cai, Yun Li, Michael Boehnke, and Xihong Lin. Rare-variant association testing for sequencing data with the sequence kernel association test. Am J Hum Genet, 89(1):82–93, 2011. ISSN 1537-6605 (Electronic); 0002-9297 (Linking). doi: 10.1016/ j.ajhg.2011.05.029.

23. Lorin Crawford, Ping Zeng, Sayan Mukherjee, and Xiang Zhou. Detecting epistasis with the marginal epistasis test in genetic mapping studies of quantitative traits. PLoS Genet, 13(7): e1006869, 2017. URL https://doi.org/10.1371/journal.pgen.1006869.

24. Gustavo de los Campos, Daniel Sorensen, and Daniel Gianola. Genomic heritability: What is it? PLOS Genetics, 11(5):e1005048, 2015. URL https://doi.org/10.1371/journal.pgen.1005048.

25. William G. Hill, Michael E. Goddard, and Peter M. Visscher. Data and theory point to mainly additive genetic variance for complex traits. PLOS Genetics, 4(2):1–10, 2008. doi: 10.1371/journal. pgen.1000008. URL https://doi.org/10.1371/journal.pgen.1000008.

26. Farhad Hormozdiari, Emrah Kostem, Eun Yong Kang, Bogdan Pasaniuc, and Eleazar Eskin. Iden-tifying causal variants at loci with multiple signals of association. Genetics, 198(2):497–508, 2014. doi: 10.1534/genetics.114.167908. URL https://pubmed.ncbi.nlm.nih.gov/25104515.

27. Priyanka Nakka, Benjamin J Raphael, and Sohini Ramachandran. Gene and Network Analysis of Common Variants Reveals Novel Associations in Multiple Complex Diseases. Genetics, 204(2): 783–798, 2016. ISSN 1943-2631. doi: 10.1534/genetics.116.188391. URL https://doi.org/10. 1534/genetics.116.188391.

28. Xiang Zhu and Matthew Stephens. Bayesian large-scale multiple regression with summary statistics from genome-wide association studies. Ann Appl Stat, 11(3):1561–1592, 2017. doi: 10.1214/ 17-AOAS1046. URL https://projecteuclid.org:443/euclid.aoas/1507168840.

29. Yan Zhang, Guanghao Qi, Ju-Hyun Park, and Nilanjan Chatterjee. Estimation of complex effect-

size distributions using summary-level statistics from genome-wide association studies across 32 complex traits. Nat Genet, 50(9):1318–1326, 2018.

30. Xiang Zhu and Matthew Stephens. Large-scale genome-wide enrichment analyses identify new trait-associated genes and pathways across 31 human phenotypes. Nat Comm, 9(1):4361, 2018.

31. Wei Cheng, Sohini Ramachandran, and Lorin Crawford. Estimation of non-null snp effect size distributions enables the detection of enriched genes underlying complex traits. PLoS Genet, 16 (6):e1008855, 2020. URL https://doi.org/10.1371/journal.pgen.1008855.

32. Pinar Demetci, Wei Cheng, Gregory Darnell, Xiang Zhou, Sohini Ramachandran, and Lorin Crawford. Multi-scale inference of genetic trait architecture using biologically annotated neural networks. PLOS Genetics, 17(8):1–53, 2021. doi: 10.1371/journal.pgen.1009754. URL https://doi.org/10.1371/journal.pgen.1009754.

33. Yongtao Guan and Matthew Stephens. Bayesian variable selection regression for genome-wide association studies and other large-scale problems. Ann Appl Stat, 5(3):1780–1815, 2011. doi: 10.1214/11-AOAS455. URL https://projecteuclid.org:443/euclid.aoas/1318514285.

34. Ronald Aylmer Fisher. The genetical theory of natural selection: a complete variorum edition. Oxford University Press, 1999. ISBN 0198504403.

35. Valentin Hivert, Julia Sidorenko, Florian Rohart, Michael E Goddard, Jian Yang, Naomi R Wray, Loic Yengo, and Peter M Visscher. Estimation of non-additive genetic variance in human complex traits from a large sample of unrelated individuals. The American Journal of Human Genetics, 108 (5):786–798, 2021.

36. Asko Mäki-Tanila and William G Hill. Influence of gene interaction on complex trait variation with multilocus models. Genetics, 198(1):355–367, 2014. doi: 10.1534/genetics.114.165282. URL https://doi.org/10.1534/genetics.114.165282.

37. Douglas Scott Falconer and Trudy FC Mackay. Quantitative genetics. Longman London, UK, 1983.

38. Armin P Schoech, Daniel M Jordan, Po-Ru Loh, Steven Gazal, Luke J O'Connor, Daniel J Balick, Pier F Palamara, Hilary K Finucane, Shamil R Sunyaev, and Alkes L Price. Quantification

of frequency-dependent genetic architectures in 25 uk biobank traits reveals action of negative selection. Nature communications, 10(1):1–10, 2019.

39. Gibran Hemani, Konstantin Shakhbazov, Harm-Jan Westra, Tonu Esko, Anjali K Henders, Allan F McRae, Jian Yang, Greg Gibson, Nicholas G Martin, Andres Metspalu, et al. Retracted article: Detection and replication of epistasis influencing transcription in humans. Nature, 508(7495): 249–253, 2014.

40. Yabo Li, Hyosuk Cho, Fan Wang, Oriol Canela-Xandri, Chunyan Luo, Konrad Rawlik, Stephen Archacki, Chengqi Xu, Albert Tenesa, Qiuyun Chen, et al. Statistical and functional studies identify epistasis of cardiovascular risk genomic variants from genome-wide association studies. Journal of the American Heart Association, 9(7):e014146, 2020.

41. Steven Gazal, Hilary K Finucane, Nicholas A Furlotte, Po-Ru Loh, Pier Francesco Palamara, Xuanyao Liu, Armin Schoech, Brendan Bulik-Sullivan, Benjamin M Neale, Alexander Gusev, et al. Linkage disequilibrium–dependent architecture of human complex traits shows action of negative selection. Nature genetics, 49(10):1421–1427, 2017.

42. Hilary K Finucane, Brendan Bulik-Sullivan, Alexander Gusev, Gosia Trynka, Yakir Reshef, Po-Ru Loh, Verneri Anttila, Han Xu, Chongzhi Zang, Kyle Farh, Stephan Ripke, Felix R Day, Shaun Purcell, Eli Stahl, Sara Lindstrom, John R B Perry, Yukinori Okada, Soumya Raychaudhuri, Mark J Daly, Nick Patterson, Benjamin M Neale, and Alkes L Price. Partitioning heritability by functional annotation using genome-wide association summary statistics. Nat Genet, 47(11): 1228–1235, 2015. ISSN 1546-1718 (Electronic); 1061-4036 (Print); 1061-4036 (Linking). doi: 10. 1038/ng.3404.

43. Daniel Runcie, Hao Cheng, and Lorin Crawford. Mega-scale linear mixed models for genomic predictions with thousands of traits. bioRxiv, page 2020.05.26.116814, 2020. doi: 10.1101/2020.05.26.116814. URL http://biorxiv.org/content/early/2020/05/29/2020.05. 26.116814.abstract.

44. Julian Stamp, Alan DenAdel, Daniel Weinreich, and Lorin Crawford. Leveraging the genetic correlation between traits improves the detection of epistasis in genome-wide association studies.

bioRxiv, page 2022.11.30.518547, 2022. doi: 10.1101/2022.11.30.518547. URL `http://biorxiv.org/content/early/2022/12/01/2022.11.30.518547.abstract`.

45. Sahin Naqvi, Yoeri Sleyp, Hanne Hoskens, Karlijne Indencleef, Jeffrey P. Spence, Rose Bruffaerts, Ahmed Radwan, Ryan J. Eller, Stephen Richmond, Mark D. Shriver, John R. Shaffer, Seth M. Weinberg, Susan Walsh, James Thompson, Jonathan K. Pritchard, Stefan Sunaert, Hilde Peeters, Joanna Wysocka, and Peter Claes. Shared heritability of human face and brain shape. Nature Genetics, 53(6):830–839, 2021. doi: 10.1038/s41588-021-00827-w. URL `https://doi.org/10.1038/s41588-021-00827-w`.

46. Shaun Purcell, Benjamin Neale, Kathe Todd-Brown, Lori Thomas, Manuel AR Ferreira, David Bender, Julian Maller, Pamela Sklar, Paul IW De Bakker, and Mark J Daly. Plink: a tool set for whole-genome association and population-based linkage analyses. Am J Hum Genet, 81(3): 559–575, 2007.

47. Zulma G Vitezica, Andrés Legarra, Miguel A Toro, and Luis Varona. Orthogonal estimates of variances for additive, dominance, and epistatic effects in populations. Genetics, 206(3):1297–1307, 2017. doi: 10.1534/genetics.116.199406. URL `https://doi.org/10.1534/genetics.116.199406`.

48. Yong Jiang and Jochen C. Reif. Modeling epistasis in genomic selection. Genetics, 201:759–768, 2015.

49. Naomi R Wray, Michael E Goddard, and Peter M Visscher. Prediction of individual genetic risk to disease from genome-wide association studies. Genome Res, 17(10):1520–1528, 2007. ISSN 1088-9051 (Print); 1549-5477 (Electronic); 1088-9051 (Linking). doi: 10.1101/gr.6665407.

50. Christoph Lippert, Gerald Quon, Eun Yong Kang, Carl M Kadie, Jennifer Listgarten, and David Heckerman. The benefits of selecting phenotype-specific variants for applications of mixed models in genomics. Scientific Reports, 3(1):1815, 2013.

51. Seung-Ho Kang and Sin-Ho Jung. Generating correlated binary variables with complete specification of the joint distribution. Biometrical Journal, 43(3):263–269, 2001. doi: https://doi.org/10.1002/1521-4036(200106)43:3⟨263::AID-BIMJ263⟩3.0.CO;2-5. URL `https://doi.org/10.1002/1521-4036(200106)43:3<263::AID-BIMJ263>3.0.CO;2-5`.

52. Farhad Hormozdiari, Martijn van de Bunt, Ayellet V. Segrè, Xiao Li, Jong Wha J. Joo, Michael Bilow, Jae Hoon Sul, Sriram Sankararaman, Bogdan Pasaniuc, and Eleazar Eskin. Colocalization of GWAS and eQTL signals detects target genes. Am J Hum Genet, 99(6):1245–1260, 2016. doi: 10.1016/j.ajhg.2016.10.003. URL https://doi.org/10.1016/j.ajhg.2016.10.003.

53. Humberto Barreto and Frank Howland. Introductory Econometrics: Using Monte Carlo Simulation with Microsoft Excel. Cambridge University Press, Cambridge, 2005. ISBN 9780521843195. doi: DOI:10.1017/CBO9780511809231. URL https://www.cambridge.org/core/books/introductory-econometrics/EA8D7F1057DB5285EFF4E44E6A6A2959.

54. Ali Pazokitoroudi, Alec M. Chiu, Kathryn S. Burch, Bogdan Pasaniuc, and Sriram Sankararaman. Quantifying the contribution of dominance deviation effects to complex trait variation in biobank-scale data. The American Journal of Human Genetics, 108(5):799–808, 2021. ISSN 0002-9297. doi: https://doi.org/10.1016/j.ajhg.2021.03.018. URL https://www.sciencedirect.com/science/article/pii/S0002929721001026.

55. Zhihong Zhu, Andrew Bakshi, Anna A. E. Vinkhuyzen, Gibran Hemani, Sang Hong Lee, Ilja M. Nolte, Jana V. van Vliet-Ostaptchouk, Harold Snieder, Tonu Esko, Lili Milani, Reedik Mägi, Andres Metspalu, William G. Hill, Bruce S. Weir, Michael E. Goddard, Peter M. Visscher, and Jian Yang. Dominance genetic variation contributes little to the missing heritability for human complex traits. The American Journal of Human Genetics, 96(3):377–385, 2024/03/05 2015. doi: 10.1016/j.ajhg.2015.01.001. URL https://doi.org/10.1016/j.ajhg.2015.01.001.

56. Daniel M. Weinreich, Yinghong Lan, Jacob Jaffe, and Robert B. Heckendorn. The influence of higher-order epistasis on biological fitness landscape topography. Journal of Statistical Physics, 172(1):208–225, 2018. doi: 10.1007/s10955-018-1975-3. URL https://doi.org/10.1007/s10955-018-1975-3.

57. Ronald A Fisher. The correlation between relatives on the supposition of mendelian inheritance. Earth and Environmental Science Transactions of the Royal Society of Edinburgh, 52(2):399–433, 1919.

58. Michael Lynch and Bruce Walsh. Genetics and analysis of quantitative traits, volume 1. Sinauer Sunderland, MA, 1998.

59. L. Isserlis. On a formula for the product-moment coefficient of any order of a normal frequency distribution in any number of variables. Biometrika, 12(1-2):134–139, 1918. ISSN 0006-3444. doi: 10.1093/biomet/12.1-2.134. URL https://doi.org/10.1093/biomet/12.1-2.134.

60. Bradley Efron. The jackknife, the bootstrap and other resampling plans. SIAM, 1982. ISBN 0898711797.

61. Peter Carbonetto and Matthew Stephens. Scalable variational inference for bayesian variable selection in regression, and its accuracy in genetic association studies. Bayesian Anal, 7(1):73–108, 2012.

62. Jennifer A. Hoeting, David Madigan, Adrian E. Raftery, and Chris T. Volinsky. Bayesian model averaging: a tutorial (with comments by m. clyde, david draper and e. i. george, and a rejoinder by the authors. Statist Sci, 14(4):382–417, 1999. doi: 10.1214/ss/1009212519. URL https://projecteuclid.org:443/euclid.ss/1009212519.

63. Shadi Zabad, Aaron P Ragsdale, Rosie Sun, Yue Li, and Simon Gravel. Assumptions about frequency-dependent architectures of complex traits bias measures of functional enrichment. Genetic epidemiology, 45(6):621–632, 2021.

64. Jian Yang, S Hong Lee, Michael E Goddard, and Peter M Visscher. Gcta: a tool for genome-wide complex trait analysis. Am J Hum Genet, 88(1):76–82, 2011. ISSN 1537-6605 (Electronic); 0002-9297 (Print); 0002-9297 (Linking). doi: 10.1016/j.ajhg.2010.11.011.

65. Carrie Zhu, Matthew J. Ming, Jared M. Cole, Michael D. Edge, Mark Kirkpatrick, and Arbel Harpak. Amplification is the primary mode of gene-by-sex interaction in complex human traits. Cell Genomics, 3(100297), 2023. doi: 10.1016/j.xgen.2023.100297. URL https://doi.org/10.1016/j.xgen.2023.100297.

66. Mashaal Sohail, Robert M Maier, Andrea Ganna, Alex Bloemendal, Alicia R Martin, Michael C Turchin, Charleston WK Chiang, Joel Hirschhorn, Mark J Daly, Nick Patterson, et al. Polygenic adaptation on height is overestimated due to uncorrected stratification in genome-wide association studies. Elife, 8:e39702, 2019.

67. James J Lee, Matt McGue, William G Iacono, and Carson C Chow. The accuracy of ld score regression as an estimator of confounding and genetic correlations in genome-wide association studies. Genetic epidemiology, 42(8):783–795, 2018.

68. Clare Bycroft, Colin Freeman, Desislava Petkova, Gavin Band, Lloyd T. Elliott, Kevin Sharp, Allan Motyer, Damjan Vukcevic, Olivier Delaneau, Jared O'Connell, Adrian Cortes, Samantha Welsh, Alan Young, Mark Effingham, Gil McVean, Stephen Leslie, Naomi Allen, Peter Donnelly, and Jonathan Marchini. The uk biobank resource with deep phenotyping and genomic data. Nature, 562(7726):203–209, 2018. doi: 10.1038/s41586-018-0579-z. URL `https://doi.org/10.1038/s41586-018-0579-z`.

69. Christopher C Chang, Carson C Chow, Laurent CAM Tellier, Shashaank Vattikuti, Shaun M Purcell, and James J Lee. Second-generation plink: rising to the challenge of larger and richer datasets. Gigascience, 4(1):s13742–015, 2015.

70. Gad Abraham, Yixuan Qiu, and Michael Inouye. Flashpca2: principal component analysis of biobank-scale genotype datasets. Bioinformatics, 2017.