1

Allelic diversity of the pharmacogene *CYP2D6* in New Zealand Māori and Pacific peoples

Leonie M. Hitchman[1], Allamanda Faatoese[2], Tony R. Merriman[3,4], Allison L. Miller[1], Yusmiati Liau[1,5], Oscar E.E. Graham[1], Ping Siu Kee[1], John F. Pearson[1], Tony Fakahau[6], Vicky A. Cameron[2], Martin A. Kennedy[1], Simran D.S. Maggo[1,7]

[1] Department of Pathology and Biomedical Science, University of Otago, Christchurch, New Zealand

[2] Christchurch Heart Institute, Department of Medicine, University of Otago, Christchurch, New Zealand

[3] Biochemistry Department, University of Otago, Dunedin, New Zealand.

[4] Division of Clinical Immunology and Rheumatology, University of Alabama at Birmingham, Alabama, US

[5] Current address: Auckland District Health Board, LabPLUS, Auckland City Hospital, Auckland, New Zealand

[6] Pacific Trust Canterbury, Montreal Street, Christchurch, New Zealand

[7] Department of Pathology, Center for Personalized Medicine, Children's Hospital Los Angeles, Los Angeles, California, USA

Keywords: *CYP2D6*, pharmacogenetics, Polynesian, Māori, Pasifika, drug response, haplotype, ancestry

Joint corresponding authors: Dr Simran Maggo, Prof Martin Kennedy

Abstract

The enzyme cytochrome P450 2D6 (CYP2D6) metabolises approximately 25% of commonly prescribed drugs, including analgesics, anti-hypertensives, and anti-depressants, among many others. Genetic variation in drug metabolising genes can alter how an individual responds to prescribed drugs, including predisposing to adverse drug reactions. The majority of research on the *CYP2D6* gene has been carried out in European and East Asian populations, with Indigenous and minority populations greatly underrepresented. However, genetic variation is often population specific and analysis of diverse ethnic groups can reveal differences in alleles that may be of clinical significance. For this reason, we set out to examine the range and frequency of *CYP2D6* variants in a sample of 202 Māori and Pacific people living in Aotearoa (New Zealand). We carried out a long PCR to isolate the *CYP2D6* region before performing nanopore sequencing to identify all variants and alleles in these samples. We identified eleven novel variants, three of which were exonic missense variations. Six of these occurred in single samples and one was found in 19 samples (9.4% of the cohort). The remaining four novel variants were identified in two samples each. In addition, five new suballeles of *CYP2D6* were identified. One striking finding was that CYP2D6*71, an allele of unknown functional status which has been rarely observed in previous studies, occurs at a relatively high frequency (9.2%) within this cohort. These data will help to ensure that *CYP2D6* genetic analysis for pharmacogenetic purposes can be carried out accurately and effectively in this population group.

Introduction

Pharmacogenetics is the study of genetic variants which impact on an individual's response to drugs, with the aim of guiding prescription practices to improve healthcare quality and outcomes (Bank et al., 2018;Cacabelos et al., 2019). *CYP2D6* is one of the most studied pharmacogenes. This gene encodes an enzyme (CYP2D6) which is expressed in the liver and responsible for metabolising approximately 25% of commonly prescribed drugs or prodrugs (Nofziger et al., 2020).

The *CYP2D6* gene is highly polymorphic, with many recorded single nucleotide variants
(SNVs), structural variants such as small insertions/deletions, larger copy number variants
including whole gene deletions or duplications, as well as hybrid genes formed by
recombination with the closely related pseudogene (*CYP2D7)* that is in close proximity
(Nofziger et al., 2020). Over 150 *CYP2D6* alleles have so far been identified, each of which is
allocated a name using a "star" nomenclature scheme, and tracked within the PharmVar
database. Minor variations are often allocated a "suballele" designation (Gaedigk et al.,
2020;Nofziger et al., 2020).

Variants of the gene may impact CYP2D6 function, ranging from completely inactivating the
enzyme through to elevating its activity, although the impact of many variants is yet to be
quantified. Where the functional impact of alleles is known or can be inferred, individuals
can be categorised into one of four metaboliser phenotypes – ultrarapid metaboliser (UM),
normal metaboliser (NM), intermediate metaboliser (IM), and poor metaboliser (PM), each
of which describes activity of the CYP2D6 enzyme (Gaedigk et al., 2008). Depending on the
type of medication, individuals defined as poor or ultrarapid metabolisers are at particular
risk of experiencing drug toxicity or poor drug response (Cacabelos et al., 2019).

Interethnic differences in allele distribution or frequencies are evident for *CYP2D6* (Gaedigk
et al., 2017;Zhou et al., 2017;Koopmans et al., 2021). For example, *CYP2D6*10* is reported
with a frequency of 45% in East Asian populations, compared to only 1.6% in Europeans
(Zhou et al., 2017). Furthermore, the PharmGKB database of *CYP2D6* variants (Whirl-Carrillo
et al., 2021) includes data from over 64,000 Europeans compared to less than 800
individuals from the Oceanian biogeographic grouping, defined as pre-colonial populations
of the Pacific, including Hawaii, Australia, New Zealand and Papua New Guinea (Huddart et
al., 2019). Many of the existing 'Oceania' studies focus on individuals from Papua New Guinea
and Melanesia, which will not fully represent genetic diversity of Oceanian populations.
Aotearoa (New Zealand) was the last major landmass to be inhabited, with Māori settlers
arriving about 730 years before present (BP)(Gosling and Matisoo-Smith, 2018). Māori and
Pacific Island populations have origins in South-East Asia, before migrations brought them
to remote Oceania around 3000BP (Gosling and Matisoo-Smith, 2018). There have been very
few studies specifically examining pharmacogenetic variability in Māori and Pacific Island

people (Wanwimolruk et al., 1995;Wanwimolruk et al., 1998;Lea et al., 2008). To ensure
equitable application of pharmacogenetic tests that detect clinically relevant alleles in people
of all ancestries, it is important that such pharmacogenetic variants are identified and
quantified by analysis of appropriate population samples.

The choice of technology employed for studies of interethnic diversity in *CYP2D6* is critical.
Many prior studies have used targeted genotyping arrays or allele-specific polymerase chain
reaction (PCR) methods, targeting only known alleles of interest, often derived from analysis
of Europeans (Carvalho Henriques et al., 2021). These analyses are relatively cheap and
straightforward to carry out, but unknown or rare alleles will not be detected (and
potentially reported as *1 if not positive for another allele of interest). A more effective
approach to identify all variants and determine haplotypes is to employ long-read, single
molecule nanopore DNA sequencing on a PCR product encompassing the entire *CYP2D6*
region (Liau et al., 2019). In this paper, we applied these methods to characterise the allelic
landscape of *CYP2D6* in two cohorts of New Zealand Māori or Pacific Island people.

Materials and Methods

Study Cohorts

A total of 202 samples were drawn from two existing studies. Thirty-five of these samples
were from a study called Genetics of Gout, Diabetes, and Kidney Disease in Aotearoa New
Zealand, which recruited individuals aged ≥16 years primarily from the Auckland, Waikato
and Christchurch regions of Aotearoa (New Zealand) (Krishnan et al., 2018). The remaining
167 samples were from the Pasifika Heart Study (Faatoese *et al*. manuscript in preparation).
This study involved 200 Pacific participants aged 20 – 64 years, selected from the patient
register of a Pacific-led primary healthcare clinic (Pacific Trust Canterbury, Christchurch,
NZ), a low-cost health provider largely serving Pacific residents of Christchurch. Screening
clinics for the Pasifika Heart Study were held from May 2015 – June 2016 at the Pacific Trust
Health Clinic and the Nicholls Research Centre, University of Otago, Christchurch. In both
studies, ethnicity was self-reported and the ethnicities of each participant's four
grandparents were also documented. The majority of samples were Polynesian, self-

reporting as Samoan, Tongan, or Māori. Other included ethnicities were Cook Island Māori,

Fijian (Melanesian), Tokelauan, Kiribati, and Niuean.

Ethical approval for the study Genetics of Gout, Diabetes, and Kidney Disease in Aotearoa

New Zealand was given by the NZ Multi-Region Ethics Committee (MEC/05/10/130;

MEC/10/09/092; MEC/11/04/036). The Pasifika Heart Study was approved by the New

Zealand Health and Disability Ethics Committee (14/CEN/72/AM04). Participants in both

studies gave written, informed consent.

121   Table 1 near here

CYP2D6 Long PCR

CYP2D6 was amplified by PCR as a 6.6kb product, including the whole gene as well as

upstream and downstream non-coding regions. Duplication and deletion primers were used

to identify the presence of a *CYP2D6* duplication or deletion via amplification of a secondary

3.5kb fragment if present (Gaedigk et al., 2007;Liau et al., 2019;Maggo et al., 2019b). Primer

sequences are provided in Supplementary Table 1.

Barcoding of PCR amplicons

Oxford Nanopore Technologies (ONT) PCR Barcoding Expansion 1-96 kit was used to add

specific barcodes onto the *CYP2D6* amplicons. This allowed sample pooling in later steps. For

a 50μL reaction, 0.5nM *CYP2D6* 6.6kb tailed amplicon was amplified in the presence of 1x

LongAMP buffer, 0.3mM Kapa dNTPs, 5 units of LongAMP Hot Start *Taq* DNA polymerase,

0.2μM barcode and ultra-pure H_2O to make up the final volume. This followed the ONT

protocol SQK- LSK109, (version PBAC96_9069_v109_revO_14Aug2019). The PCR began

with an incubation at 95°C for 3 minutes, followed by 15 cycles of 95°C for 15 seconds, 62°C

for 15 seconds, 65°C for 7 minutes, and finally 65°C for 7 minutes, as recommended by ONT.

Magnetic Bead Purification

Magnetic beads were used to purify the PCR products both prior to and following the

barcoding step. This removed DNA fragments shorter than 3-4kb (to avoid off-target

amplification) and other impurities. Magbio beads (MagBio Genomics Inc, Gaithersburg,

Maryland, USA) were pelleted and re-dissolved in a buffer with 10mM Tris-HCl, 1.6M NaCl,

1mM EDTA pH 8, 11% (w/v) PEG 8000, 0.20% (v/v) Tween-20, and ultra-pure water to

make a 10mL bead solution (Nagar and Schwessinger, 2018). DNA samples were purified as

described previously (Liau et al., 2019).

Nanopore Sequencing Library Preparation

DNA libraries of barcoded amplicons were prepared according to the 1D PCR barcoding (96)

amplicons ONT protocol, SQK-LSK109. Briefly, after pooling the barcoded amplicons, about

200 fmoles of the DNA pool was subjected to end-repair and ONT adapter ligation. After

purification, approximately 210 ng ($\sim$ 50 fmoles) of the library was introduced to the MinION

flowcell (R9.4.1) and sequenced for up to 48 hours on the GridION X5 nanopore sequencer

(ONT, UK). Some sequencing was also completed on a FLO-FLG-001 flongle (R9.4.1) loaded

with approximately 90 ng ($\sim$ 20 fmoles) of library.

Nanopore Sequencing Data analysis

A previously designed pipeline was used to analyse the data generated (Graham et al., 2020).

The GridION platform conducted real-time filtering, basecalling, and demultiplexing

(separating and binning each sample per barcode) using Guppy version 5.0.12 (ONT, UK). A

quality threshold for sequencing was set, specifying reads between 6 - 8kb in length and a

Qscore of >9. An end-to-end pipeline for data management was designed in a conda

environment using a snakemake workflow (Mölder et al., 2021). The process included

utilising the output of the GridION to initially map the FASTQ files generated against a

reference sequence (CYP2D6_NG008376.3) using MiniMap2 version 2.20-r1061 (Li, 2018).

SAMtools (version 1.7) (Li et al., 2009) was employed to perform indexing at various stages

of this pipeline. Nanopolish (version 0.13.2) (Quick et al., 2016) was used to analyse variant

calls. Whatshap (version 0.17) (Martin et al., 2016) was used to phase the VCF files generated

by Nanopolish. Variants were then matched to *CYP2D6* star alleles using the PharmVar

database (Pharmacogene Variation Consortium) and unmatched variants were identified

and marked as potentially novel to be taken forward for later validation. Finally, Stargazer

v1.0.8 (Lee et al., 2019), an automated tool for pharmacogenetic star allele assignment, was

used to confirm the *CYP2D6* star allele assigned to each sample. Star allele frequencies

between studies were compared with two sided Fisher's exact test and considered
significant at p=0.05.

Sanger sequencing

To validate potential novel alleles, Sanger sequencing was performed following a nested PCR
on the relevant *CYP2D6* gene locations, using the long amplicon as a template as described
(Maggo et al., 2019a; Maggo et al., 2019b). Primer sequences are provided in Supplementary
Table 2.

Results

From an initial 282 samples, 202 were successfully sequenced and analysed at all steps.
Excluded samples included those that failed to amplify the 6.6kb *CYP2D6* PCR product
(n=52), or samples with a low read depth (less than 50 reads; n=28). Genomic DNA samples
which failed to amplify were checked on a TapeStation platform (Agilent, Santa Clara, USA)
and were usually found to be fragmented, with this fragmentation being the likely cause of
long-PCR failure. 120 of the samples were sequenced with an R9.4.1 flow cell, and the
remaining 82 were sequenced with a FLG-001 Flongle, with both flow cell types run on the
GridION X5.

CYP2D6 Genotyping Results

Nine different *CYP2D6* star alleles were identified amongst 202 individuals, including the *5
gene deletion (Table 2). No *CYP2D6* gene duplications were observed. The most common
allele was *1 with an allele frequency of 0.48. Following this, in order of frequency, were *10,
*2, and *71. Four of the identified star alleles, *4, *5, *10, and *41, are known to decrease or
completely inactivate the function of the CYP2D6 liver enzyme, while *1, *2, and *35 have no
impact. *43 and *71 have unknown functional effects on CYP2D6.

Stargazer 1.0.8 was used to call star alleles for the first 120 samples, using variants aligned
to hg19. Of these 120, eleven were called incorrectly. Five of the incorrect haplotype
assignments were due to the presence of a *5 allele, while the remaining six carried *71
variants.

197   Table 2 near here

198 Predicted Metaboliser Phenotypes

The metaboliser status was inferred from the genotype of each sample using the diplotype

activity score as previously defined (Caudle et al., 2020). In this classification, diplotypes

with an activity score of 0 are classified as poor metabolisers, 0.25-1 are intermediate

metabolisers, 1.25-2 are normal metabolisers, and scores above 2 are ultrarapid

metabolisers. Using this definition, the majority (74%) of individuals within the cohort were

classified as normal metabolisers. No individuals were identified as being ultrarapid or poor

metabolisers, and a small subset (8%) of our study cohort were intermediate metabolisers.

Almost 20% of the characterised samples had a metaboliser status that could not be assigned

due to the presence of one or more *71 or *43 alleles (Table 3).

208   Table 3 near here

210 Novel Variants

211   We identified eleven potentially novel variants and validated them using Sanger

212   sequencing (Table 4). Seven variants occur within an intron of the gene, and four are exonic

213   variants, three of which are non-synonymous. These have the potential to form new star

214   alleles or new suballeles.

215   Table 4 near here

Six of the eleven novel variants were singletons (detected in only one individual each), four

were identified in two individuals each, while a variant at position 5709 was found in

nineteen people. In addition, five potentially novel suballeles were also identified (data not

shown). These variants occur within the PharmVar CYP2D6 database but as part of different

star alleles to those observed in this study. The majority exist on the haplotype without any

core variants belonging to star alleles, so are by default *1. Two variants at position 8595

and 8645 are likely a new *10 suballele as all samples have the *10 core variants.

Discussion

Of the nine broad ethnic groups defined by Huddart *et al.,* and reported by PharmGKB,

Oceania is by far the least represented (PharmGKB) (Huddart et al., 2019)**.** We have

successfully used nanopore sequencing to analyse the entire *CYP2D6* gene (including

important upstream and downstream regions) of 202 Māori and Pacific Island individuals,

coupled with analyses to call the star allele diplotypes of each participant. Nine star alleles

were identified amongst this cohort, with frequencies ranging from 0.48 for *1, to 0.002 for

*43. CYP2D6*71, a previously identified rare allele with unknown function, was identified at

a relatively high frequency within this cohort, with an allele frequency of 0.092. From each

individual's diplotype, we were able to categorise the majority of the cohort into metaboliser

phenotypes. We did not find any ultrarapid or poor metabolisers. A large percentage of

participants (74%) were normal metabolisers and 8% had a predicted activity score within

the range of intermediate metabolisers (Caudle et al., 2020)**.** The remaining 18% were

unable to be assigned to a metaboliser status due to carrying one or more copies of the *43

or *71 allele.

Eleven novel variants were also discovered, four of which are exonic. Five known variants

were found within star alleles in which they have not been reported previously, potentially

forming new suballeles.

We did not identify any poor metabolisers in the present study, as all non-functional alleles

were found in diplotypes with a fully functional or partially functional allele. Non-functional

alleles (*4 and *5 gene deletion) were present in ~3% of the cohort, so they do exist at a low

frequency within the population and therefore could give rise to poor metabolisers

phenotypes. In this regard, it is interesting to note that a pharmacological study using the

probe drug debrisoquine in 101 Māori participants identified 5% as poor metabolisers

(Wanwimolruk et al., 1995), and a similar study in 100 Polynesian participants found none

were poor metabolisers (Wanwimolruk et al., 1998). Therefore, our genetic findings are
reasonably congruent with these earlier studies.

However, *CYP2D6* allele frequencies observed in our study do not match allele frequencies
previously reported by PharmGKB for the Oceanian region (PharmGKB). This is almost
certainly because the Oceanian definition encompasses populations with diverse, complex
migration and ancestral histories (Gosling and Matisoo-Smith, 2018). The Oceanian *CYP2D6*
data currently available comes from studies that in total include less than 800 individuals.
The largest study (Gutiérrez Rico et al., 2020) focused on 278 Ni-Vanuatu (Micronesian)
subjects, and found the majority of individuals to carry at least one *1 allele (allele frequency
of 0.804), significantly greater than our identified frequency of 0.478 (P = 4.2 x 10^{-26}). In total,
they identified nine different star alleles amongst their cohort, the majority of which had
allelic frequencies of <0.01. *2 and *10 existed above this frequency (0.068 and 0.029,
respectively) as well as a duplication of *1 at 0.025 (Gutiérrez Rico et al., 2020). Our study
found no duplications and only two alleles had observed frequencies below 0.01 (*35, *43).
Prior reports describing high rates of *CYP2D6* duplications in Oceania have been entirely
based on Papua New Guinea samples (Sistonen et al., 2007;von Ahsen et al., 2010).

The relatively high frequency of the *71 allele we observed is one of the most striking
findings from this study, for several reasons. Because *71 has thus far been considered a rare
allele, it would not be routinely included in most *CYP2D6* genotyping panels. The functional
impact of *71 is unknown, so we were unable to predict metaboliser status for 18% of our
cohort. The first study to identify *71 was performed in a Han Chinese population (Zhou et
al., 2009). This group suggested the allele could be non-functional due to the p.Gly42 >Glu
substitution in the protein N- terminus, a region which is involved in membrane anchoring
of the enzyme (Zhou et al., 2009). However, this is speculative and has yet to be tested. Other
reports, in Micronesian populations, have described allele frequencies of 0.009 in a Ni-
Vanuatu sample (Gutiérrez Rico et al., 2020), significantly less than our 0.092 (P = 3.9x10^{-10})
and 0.058 in a large sample of Solomon Islanders (Charnaud et al., 2022), comparable but
significantly less than our 0.092 (P=0.038).

Conclusion

In conclusion, we have used a nanopore sequencing approach to identify *CYP2D6* star alleles and respective frequencies from 202 individuals of Māori or Pasifika ethnicity. We identified nine known star alleles in this cohort, including CYP2D6*71, a previously identified rare allele of unknown function, which comprised over 9% of the alleles detected. The frequency of this allele prevented us from inferring metabolic function for nearly 20% of this cohort, and clearly, understanding the functional impact of *71 using clinical phenotyping approaches or *in vitro* analyses is a priority area for future research. Should this allele have an altered function, it would be important to ensure it is included in pharmacogenetic testing panels, particularly if to be used in the Oceania region.

Our analysis did not reveal any *CYP2D6* duplications in this cohort, and overall our findings highlight the significant limitations of the Oceania biogeographic grouping for pharmacogenetic research, given the wide allele frequency differences we observe relative to studies on other Oceanian ancestral groups.

Finally, this study has further demonstrated the utility of nanopore sequencing for highly variable genes like *CYP2D6*, with long-read sequencing providing the advantage of analysing the entire gene, including intronic regions, at high throughput and relatively low cost. It is to be hoped this research will contribute to a more equitable uptake of *CYP2D6* pharmacogenetics in people of Māori and Pacific ancestry.

.

References

Bank, P.C.D., Swen, J.J., and Guchelaar, H.J. (2018). Implementation of Pharmacogenomics in Everyday Clinical Settings. *Adv Pharmacol* 83, 219-246.

Cacabelos, R., Cacabelos, N., and Carril, J.C. (2019). The role of pharmacogenomics in adverse drug reactions. *Expert Rev Clin Pharmacol* 12, 407-442.

Carvalho Henriques, B., Buchner, A., Hu, X., Wang, Y., Yavorskyy, V., Wallace, K., Dong, R., Martens, K., Carr, M.S., Behroozi Asl, B., Hague, J., Sivapalan, S., Maier, W., Dernovsek, M.Z., Henigsberg, N., Hauser, J., Souery, D., Cattaneo, A., Mors, O., Rietschel, M., Pfeffer, G., Hume, S., and Aitchison, K.J. (2021). Methodology for clinical genotyping of CYP2D6 and CYP2C19. *Transl Psychiatry* 11, 596.

Caudle, K.E., Sangkuhl, K., Whirl-Carrillo, M., Swen, J.J., Haidar, C.E., Klein, T.E., Gammal, R.S., Relling, M.V., Scott, S.A., Hertz, D.L., Guchelaar, H.-J., and Gaedigk, A. (2020). Standardizing CYP2D6 Genotype to Phenotype Translation: Consensus Recommendations from the Clinical Pharmacogenetics Implementation Consortium and Dutch Pharmacogenetics Working Group. *Clinical and Translational Science* 13, 116-124.

Charnaud, S., Munro, J.E., Semenec, L., Mazhari, R., Brewster, J., Bourke, C., Ruybal-Pesantez, S., James, R., Lautu-Gumal, D., Karunajeewa, H., Mueller, I., and Bahlo, M. (2022). PacBio long-read amplicon sequencing enables scalable high-resolution population allele typing of the complex CYP2D6 locus. *Commun Biol* 5, 168.

Gaedigk, A., Ndjountche, L., Divakaran, K., Dianne Bradford, L., Zineh, I., Oberlander, T., Brousseau, D., Mccarver, D., Johnson, J., and Alander, S. (2007). Cytochrome P4502D6 (CYP2D6) gene locus heterogeneity: characterization of gene duplication events. *Clinical Pharmacology & Therapeutics* 81, 242-251.

Gaedigk, A., Sangkuhl, K., Whirl-Carrillo, M., Klein, T., and Leeder, J.S. (2017). Prediction of CYP2D6 phenotype from genotype across world populations. *Genet Med* 19, 69-76.

Gaedigk, A., Simon, S.D., Pearce, R.E., Bradford, L.D., Kennedy, M.J., and Leeder, J.S. (2008). The CYP2D6 activity score: translating genotype information into a qualitative measure of phenotype. *Clin Pharmacol Ther* 83, 234-242.

Gaedigk, A., Whirl-Carrillo, M., Pratt, V.M., Miller, N.A., and Klein, T.E. (2020). PharmVar and the Landscape of Pharmacogenetic Resources. *Clin Pharmacol Ther* 107, 43-46.

Gosling, A.L., and Matisoo-Smith, E.A. (2018). The evolutionary history and human settlement of Australia and the Pacific. *Curr Opin Genet Dev* 53, 53-59.

Graham, O.E.E., Pitcher, T.L., Liau, Y., Miller, A.L., Dalrymple-Alford, J.C., Anderson, T.J., and Kennedy, M.A. (2020). Nanopore sequencing of the glucocerebrosidase (GBA) gene in a New Zealand Parkinson's disease cohort. *Parkinsonism Relat Disord* 70, 36-41.

Gutiérrez Rico, E.M., Kikuchi, A., Saito, T., Kumondai, M., Hishinuma, E., Kaneko, A., Chan, C.W., Gitaka, J., Nakayoshi, T., Oda, A., Saito, S., Hirasawa, N., and Hiratsuka, M. (2020). CYP2D6 genotyping analysis and functional characterization of novel allelic variants in a Ni-Vanuatu and Kenyan population by assessing dextromethorphan O-demethylation activity. *Drug Metabolism and Pharmacokinetics* 35, 89-101.

Huddart, R., Fohner, A.E., Whirl-Carrillo, M., Wojcik, G.L., Gignoux, C.R., Popejoy, A.B., Bustamante, C.D., Altman, R.B., and Klein, T.E. (2019). Standardized Biogeographic

Grouping System for Annotating Populations in Pharmacogenetic Research. *Clin Pharmacol Ther* 105**,** 1256-1262.

Koopmans, A.B., Braakman, M.H., Vinkers, D.J., Hoek, H.W., and Van Harten, P.N. (2021). Meta-analysis of probability estimates of worldwide variation of CYP2D6 and CYP2C19. *Transl Psychiatry* 11**,** 141.

Krishnan, M., Major, T.J., Topless, R.K., Dewes, O., Yu, L., Thompson, J.M.D., Mccowan, L., De Zoysa, J., Stamp, L.K., Dalbeth, N., Harre Hindmarsh, J., Rapana, N., Deka, R., Eng, W.W.H., Weeks, D.E., Minster, R.L., Mcgarvey, S.T., Viali, S., Naseri, T., Sefuiva Reupena, M., Wilcox, P., Grattan, D., Shepherd, P.R., Shelling, A.N., Murphy, R., and Merriman, T.R. (2018). Discordant association of the CREBRF rs373863828 A allele with increased BMI and protection from type 2 diabetes in Maori and Pacific (Polynesian) people living in Aotearoa/New Zealand. *Diabetologia* 61**,** 1603-1613.

Lea, R., Roberts, R., Green, M., Kennedy, M., and Chambers, G. (2008). Allele frequency differences of cytochrome P450 polymorphisms in a sample of New Zealand Maori. *The New Zealand medical journal* 121**,** 33-37.

Lee, S.-B., Wheeler, M.M., Thummel, K.E., and Nickerson, D.A. (2019). Calling Star Alleles With Stargazer in 28 Pharmacogenes With Whole Genome Sequences. *Clinical Pharmacology & Therapeutics* 106**,** 1328-1337.

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34**,** 3094-3100.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25**,** 2078-2079.

Liau, Y., Maggo, S., Miller, A.L., Pearson, J.F., Kennedy, M.A., and Cree, S.L. (2019). Nanopore sequencing of the pharmacogene CYP2D6 allows simultaneous haplotyping and detection of duplications. *Pharmacogenomics* 20**,** 1033-1047.

Maggo, S., Kennedy, M.A., Barczyk, Z.A., Miller, A.L., Rucklidge, J.J., Mulder, R.T., and Foulds, J.A. (2019a). Common CYP2D6, CYP2C9, and CYP2C19 Gene Variants, Health Anxiety, and Neuroticism Are Not Associated With Self-Reported Antidepressant Side Effects. *Frontiers in Genetics* 10.

Maggo, S.D.S., Sycamore, K.L.V., Miller, A.L., and Kennedy, M.A. (2019b). The Three Ps: Psychiatry, Pharmacy, and Pharmacogenomics, a Brief Report From New Zealand. *Front Psychiatry* 10**,** 690.

Martin, M., Patterson, M., Garg, S., Fischer, S., Pisanti, N., Klau, G., Schöenhuth, A., and Marschall, T. (2016). "WhatsHap: fast and accurate read-based phasing. bioRxiv, 085050".).

Mölder, F., Jablonski, K., Letcher, B., Hall, M., Tomkins-Tinch, C., Sochat, V., Forster, J., Lee, S., Twardziok, S., Kanitz, A., Wilm, A., Holtgrewe, M., Rahmann, S., Nahnsen, S., and Köster, J. (2021). Sustainable data analysis with Snakemake [version 1; peer review: 1 approved, 1 approved with reservations]. *F1000Research* 10.

Nagar, R., and Schwessinger, B. (2018). DNA size selection (> 3–4 kb) and purification of DNA using an improved homemade SPRI beads solution. v. 1. *Protocols. io* 10.

Nofziger, C., Turner, A.J., Sangkuhl, K., Whirl-Carrillo, M., Agundez, J.a.G., Black, J.L., Dunnenberger, H.M., Ruano, G., Kennedy, M.A., Phillips, M.S., Hachad, H., Klein, T.E., and Gaedigk, A. (2020). PharmVar GeneFocus: CYP2D6. *Clin Pharmacol Ther* 107**,** 154-170.

Pharmacogene Variation Consortium *CYP2D6* [Online]. Available: https://www.pharmvar.org/gene/CYP2D6 [Accessed 6 October 2021].

Pharmgkb *CYP2D6 Frequency Table* [Online]. Available: https://api.pharmgkb.org/v1/download/file/attachment/CYP2D6_frequency_table.xlsx [Accessed 30th Sept 2021].

Quick, J., Loman, N.J., Duraffour, S., Simpson, J.T., Severi, E., Cowley, L., Bore, J.A., Koundouno, R., Dudas, G., and Mikhail, A. (2016). Real-time, portable genome sequencing for Ebola surveillance. *Nature* 530**,** 228-232.

Sistonen, J., Sajantila, A., Lao, O., Corander, J., Barbujani, G., and Fuselli, S. (2007). CYP2D6 worldwide genetic variation shows high frequency of altered activity variants and no continental structure. *Pharmacogenet Genomics* 17**,** 93-101.

Steen, V., Andreassen, O., Daly, A., Tefre, T., Borresen, A., Idl, E.J., and Gulbrandsen, A. (1995). Detection of the poor metabolizer-associated CYP2D6(D) gene deletion allele by long-PCR technology. *Pharmacogenetics* 5**,** 214-223.

Von Ahsen, N., Tzvetkov, M., Karunajeewa, H.A., Gomorrai, S., Ura, A., Brockmöller, J., Davis, T.M.E., Mueller, I., Ilett, K.F., and Oellerich, M. (2010). CYP2D6 and CYP2C19 in Papua New Guinea: High frequency of previously uncharacterized CYP2D6 alleles and heterozygote excess. *International journal of molecular epidemiology and genetics* 1**,** 310-319.

Wanwimolruk, S., Bhawan, S., Coville, P.F., and Chalcroft, S.C. (1998). Genetic polymorphism of debrisoquine (CYP2D6) and proguanil (CYP2C19) in South Pacific Polynesian populations. *Eur J Clin Pharmacol* 54**,** 431-435.

Wanwimolruk, S., Pratt, E.L., Denton, J.R., Chalcroft, S.C., Barron, P.A., and Broughton, J.R. (1995). Evidence for the polymorphic oxidation of debrisoquine and proguanil in a New Zealand Maori population. *Pharmacogenetics* 5**,** 193-198.

Whirl-Carrillo, M., Huddart, R., Gong, L., Sangkuhl, K., Thorn, C.F., Whaley, R., and Klein, T.E. (2021). An Evidence-Based Framework for Evaluating Pharmacogenomics Knowledge for Personalized Medicine. *Clin Pharmacol Ther* 110**,** 563-572.

Wright, G.E.B., Niehaus, D.J.H., Drögemöller, B.I., Koen, L., Gaedigk, A., and Warnich, L. (2010). Elucidation of CYP2D6 Genetic Diversity in a Unique African Population: Implications for the Future Application of Pharmacogenetics in the Xhosa Population. *Annals of Human Genetics* 74**,** 340-350.

Zhou, Q., Yu, X.M., Lin, H.B., Wang, L., Yun, Q.Z., Hu, S.N., and Wang, D.M. (2009). Genetic polymorphism, linkage disequilibrium, haplotype structure and novel allele analysis of CYP2C19 and CYP2D6 in Han Chinese. *Pharmacogenomics J* 9**,** 380-394.

Zhou, Y., Ingelman-Sundberg, M., and Lauschke, V.M. (2017). Worldwide Distribution of Cytochrome P450 Alleles: A Meta-analysis of Population-scale Sequencing Projects. *Clin Pharmacol Ther* 102**,** 688-700.

Tables:

Table 1 Ethnicity details of participants

	Primary Ethnicity	Other Ethnicity 1	Other Ethnicity 2	Other Ethnicity 3
Samoan	80	9	2	1
Tongan	53	3		
Fijian	35	6		1
NZ Māori	23	1	1	
Cook Island Māori	5	3		
NZ European	3	11	1	
Kiribati	1			
Tokelauan	1			
Niuean	1			
Grand Total	**202**	**33**	**4**	**2**

Table 2 Star allele frequencies and associated information

Star allele	Allele frequency (n=202)	Number of haplotypes observed (n=404)	Activity score[1]	Impact
*1	0.478	193	1	Normal function
*2	0.141	57	1	Normal function
*4	0.017	7	0	No function
*5	0.012	5	0	No function
*10	0.183	74	0.25	Decreased function
*35	0.007	3	1	Normal function
*41	0.067	27	0.5	Decreased function
*43	0.002	1	?	Uncertain function
*71	0.092	37	?	Uncertain function

[1] Activity score and associated metaboliser phenotype as defined by Caudle et al (2020).

Table 3 Inferred metaboliser frequencies within cohort

Inferred Phenotype (from genotype)	Number	Frequency
Ultrarapid	0	0
Normal	150	0.743
Intermediate	16	0.079
Poor	0	0
Unknown	36	0.178

Table 4 Novel Variants

Variant position (CYP2D6 reference)[1]	SNP rsID	Type of variant	Number of times observed	Allelic designation
5252 (Exon 2)	rs/na	Missense A>G (Gln > Arg)	1	*new
5709 (Intron 2)	rs369508051	G>A	19[2]	*41.new
6128 (Exon 4)	rs755518310	Missense G>A (Gly > Glu)	1	*71.new
6239 (Intron 4)	rs564994275	A>C	1	*4.new
6242 (Intron 4)	rs778008161	G>C	1	*1.new
6419 (Intron 4)	rs/na	G>C	2	*1.new
7557 (Intron 7)	rs376909251	G>A	1	*1.new
7574 (Intron 7)	rs/na	Indel T>TCAGCAC	2	*1.new
8030 (Exon 8)	rs1602566413	Missense G>C (Val > Leu)	2[2]	*new
8068 (Exon 8)	rs1037093492	Synonymous C>T (Leu > Leu)	2[2]	*2.new
8202 (Intron 8)	rs769045995	A>G	1	*41.new

[1]Reference sequence AY545216 (1=sequence start) NG_008376.3

[2]Variant also identified by Charnaud et al. (2022)

Supplementary Table 1 Primer Sequences for CYP2D6 amplification

Primer Name	Primer Sequence (5'-3')	Reference
6.6kb F	ATGGCAGCTGCCATACAATCCACCTG	(Gaedigk et al., 2007)
6.6kb R	CGACTGAGCCCTGGGAGGTAGGTAG	(Gaedigk et al., 2007)
Tailed F	TTTCTGTTGGTGCTGATATTGC	ONT
Tailed R	ACTTGCCTGTCGCTCTATCTTC	ONT
CYP-13 (Del)	ACCGGGCACCTGTACTCCTCA	(Steen et al., 1995)
CYP-24 (Del)	GCATGAGCTAAGGCACCCAGAC	(Steen et al., 1995)
FragB F (Dup)	CCATGGAAGCCCAGGACTGAGC	(Gaedigk et al., 2007)
FragB R (Dup)	CGGCAGTGGTCAGCTAATGAC	(Gaedigk et al., 2007)

Supplementary Table 2. Primer Sequences for novel variant confirmation

Primer Name	Primer Sequence (5'-3')	References
Exon 2 Forward	TCCTCCTTCCACCTGCTCAC	(Wright et al., 2010)
Exon 2 Reverse	CTTTGCCCCACCTCGTCTCT	(Wright et al., 2010)
Exon 3 4 (old) Forward	AGCTGGAATCCGGTGTCGAA	(Wright et al., 2010)
Exon 3 4 (old) Reverse	AGCCATCTCCAGGTAGACCCAG	(Wright et al., 2010)
Exon 3 4 (new) Forward	ATAGGGTTGGAGTGGGTGGT	This publication
Exon 3 4 (new) Reverse	AAATCCTGCTCTTCCGAGGC	This publication
Exon 7 Forward	CCAACATAGGAGGCAAGAAG	(Wright et al., 2010)
Exon 7 Reverse	ACTGGACTCTAGGATGCTGG	(Wright et al., 2010)
Exon 8 Forward	GTCTAGTGGGGAGACAAACCAG	This publication
Exon 8 Reverse	TGCCCTGAGGAGGATGATC	This publication