Climate and Morphology Drive Breeding Periods in Frogs

Running Title: Drivers of amphibian reproduction

Abstract

Aim: Climate can have profound effects on reproductive behavior and physiology, especially in
ectothermic animals. Breeding periods in amphibians have received little attention despite
amphibian reliance on climate and water due to their reproductive biology and ecological
diversity. The aim of this study is to determine how global climate impacts breeding periods in
ectothermic animals through physiology, body size, and microhabitat.
11
12 Location: All continents, except Antarctica.
13
14 Time period: Breeding periods and climate both dating as far back as 1970.
15
16 Major taxa studied: 497 (7%) anuran species of 41 (76%) families.
17
18 Methods: We used phylogenetic comparative methods to analyze a global dataset of frog
19 breeding periods, climate, body size, and microhabitat for 497 species.
20
Results: We found support for a global latitudinal gradient of breeding periods which are longer
in the warmer, wetter tropics while shorter in the colder, dryer temperate zone. Latitudinal and
non-latitudinal global patterns of breeding period were composites of the same patterns in the
temperate and tropical zones. However, the effect of climate and body size in each zone is
unique. Breeding periods displayed weak phylogenetic signal and were similar across
microhabitats.
27
Main conclusions: Breeding periods show a global latitudinal gradient but this concept breaks
down within the tropical zone. Our results are consistent with the importance of physiology in
driving breeding periods and we describe how breeding period and body size may exhibit trade-
offs which make latitudinal gradients context-dependent. Our results set within an
ecophysiological framework have broad implications for understanding reproductive diversity in
other ectothermic organisms.
34
35 **Key Words:** Anura, body size, environment, microhabitat, phenology, reproduction

Introduction

Latitudinal gradients help reveal the origins of biodiversity. Examples include latitudinal gradients of species diversity (Pianka, 1966), geographical ranges (Stevens, 1989), niche breadth (Vázquez & Stevens 2004), biotic interactions (Schemske *et al.*, 2009), and body size (Blackburn & Hawkins, 2004). Latitudinal gradients of species traits are important because they are evidence of evolution by natural selection in response to environmental differences. Behavioral traits responding to temporal changes in climate (phenology) are particularly helpful in understanding ecological adaptation (Wcislo, 1989; Blomberg *et al.,* 2003). The diversification of behavior is often studied by measuring trait differences between one or more species exposed to various conditions in controlled environments or by determining how behavior is related to climate among many species through comparative approaches (Harvey & Purvis, 1991). This includes research on the effects of latitudinal climate on flowering and fruiting in plants (Badeck *et al.*, 2004; Butt *et al.*, 2015; Daru *et al.*, 2019), hatching, emergence, and flight in arthropods (Kaspari *et al.,* 2001; Śniegula *et al.,* 2016), and migration, breeding, and biotic interactions in animals (reviewed in Cohen *et al.,* 2018). Much of this research has benefited from using ectothermic organisms although research on amphibians is relatively limited. Since many amphibian groups are threatened by population declines, understanding how behavioral traits covary with climate fills a gap in our general understanding of phenology and informs future conservation strategies.

Frogs (Anura) are an ecologically diverse group (>7,400 species; AmphibiaWeb, 2022) ideal for studying phenological patterns. In addition to being ectothermic, reproductive cycles and calling behaviors in frogs vary with temperature, precipitation, and latitude (Wells, 2007). Additionally, anurans exhibit clear and rapid responses to climate. For example, warmer winters result in earlier breeding in frogs (Beattie, 1985; Reading, 2003; Neveu, 2009; Todd *et al.,* 2011;

While & Uller, 2014; Benard, 2015). However, relative to breeding start dates, little is known

about how climate impacts breeding periods despite the stunning contrast of tropical species

that breed year-round to explosive-breeding species that limit reproduction to only a few days of

the year (Wells, 1977). Natural selection likely acts on breeding periods because longer periods

of reproduction are associated with the production of more egg clutches (Bull & Shine, 1979;

Morrison & Hero, 2003) and warm and wet climates (Morrison & Hero, 2003; Wells, 2007). One

study on anurans found breeding durations (at the family level) were shorter in the dryer parts of

Brazil (Forti *et al.*, 2022). To our knowledge, no studies have examined climate effects on frog

breeding phenology specifically at a broad taxonomic and global scale.

There are several non-exclusive hypotheses relating climate and ecology to frog

breeding periods. For example, breeding periods could be constrained by cold and dry climates

since frogs generally rely on warmth and water for reproduction. This 'climate hypothesis'

predicts that species at lower latitudes should have longer breeding periods. Additionally,

reproductive behaviors could vary across latitudes due to climatic differences between regions.

This 'zone hypothesis' predicts that phenology may be influenced more by temperature in the

temperate region and precipitation in the tropics (Cohen *et al.,* 2018). Finally, extreme habitat

differences may drive reproductive behaviors, as burrowing frogs living in deserts often have

breeding lasting from only a few days (Wells, 1977). This 'microhabitat hypothesis' predicts that

desert environments constrains breeding compared to other microhabitats.

In addition to the climate and ecology, body size may also be an important factor

influencing phenology (Cohen *et al.,* 2018). Larger body sizes in frogs are associated with larger

clutches (Gomez-Mestre *et al.,* 2012; Furness *et al.,* 2022) and possibly access to larger-bodied

prey (Wilson, 1975) or greater storage capacity of energy from food (as in mammals; Demment

& Van Soest, 1985). The pace of life framework (Hille & Cooper, 2015) connects environmental

variation with life history and body size. For example, it describes how warmer, wetter, and less

seasonal environments (where growth and reproduction are not environmentally constrained)

might select for larger body sizes, slower developmental rates, larger clutches, and many

reproductive events (iteroparity). The converse applies to smaller species under the pace of life

framework. Thus, we predict relatively larger body sizes are associated with longer breeding

periods in warmer, wetter, and less seasonal environments and the converse for smaller

species. We refer to these dual predictions as the 'climate-size hypothesis'.

The goal of this study is to use the comparative approach to understand the climatic and

morphological factors which might influence breeding periods, i.e. breeding durations at the

population level (Wells, 1977), in frogs. Specifically, we test predictions from the climate, zone,

and climate-size hypotheses to determine: 1) whether breeding periods are driven by latitude,

geographical zone, or phylogenetic history, 2) whether longer breeding periods are associated

with warm, wet, and less seasonal climates, 3) whether burrowing lifestyles result in shorter

breeding periods, and 4) whether the relationship between breeding period and climate depends

on body size.

Material and Methods

Coordinate and climate data

We obtained coordinate data for 497 species representing 41 of 54 anuran families from

GBIF.org (GBIF.org, 2021a,b,c,d, accessed 3 October 2021). We describe our data cleaning

procedures in Appendix S1 in Supporting Information. We subsetted the coordinate data to

match the set of species for which we obtained behavior, phylogenetic, microhabitat, and body

size data. This resulted in 822,953 coordinates (mean = 1,656 and range = 3–83,284

observations per species) from which we obtained species mean estimates of absolute latitude

and climate data using WorldClim 2.1 (Fick & Hijmans, 2017), and envirem (Title & Bemmels,

2018) at 30 seconds (~1km^2) resolution. We used an equal-area Mollweide projection to avoid

biases in land area. From these projections, we extracted from each coordinate variables

affecting temperature and water availability covering all possible breeding period values (0–365

days). We reduced the effects of spatial autocorrelation by avoiding repeated sampling of the

same (1km^2) pixels. Specifically, this dataset included annual mean temperature, temperature

seasonality, annual precipitation, precipitation seasonality, annual potential evapotranspiration

(PET), PET seasonality, and topographic wetness. Topographic wetness is the land's ability to

retain water and is obtained from estimates of upslope incurrent areas and local slope

(Sørensen *et al.,* 2005). We calculated species means for each climate variable to analyze the

dataset in a phylogenetic comparative framework.

Breeding period data

We compiled species means of breeding period (sensu Wells, 1977) using data from the

primary literature, AmphibiaWeb (2022), and online databases. We estimated the breeding

period as the difference in calendar days between the breeding start and end dates, assuming

365 days, when the breeding period was not already reported in days. We then averaged these

values across populations (N = 1–6, mean = 1) to obtain species means. Some records were

reported as qualitative data (e.g., "early May"). When breeding periods were reported as starting

or ending on a specific month, we ascribed values of either the first or last day of the month,

respectively. When breeding periods were reported as the "early" or "late" parts of a month, we

matched this to the end of the 1st week and 3rd week, respectively. Rarely, breeding periods

were reported as specific seasons (e.g., "Summer" or "wet season") and we defined these as

the number of days corresponding to each season and country. Since explosive breeding is <30

days (Wells, 1977) and because we did not find any records of explosive breeding which took

longer than 2 weeks, we conservatively assigned a period of 2 weeks to species whose only

description of the breeding period was "explosive breeding".

Phylogenetic, habitat, and body size data

We obtained the consensus tree and 1,000 trees randomly sampled from the

pseudoposterior distribution of trees from Jetz and Pyron (2018) to account for shared

evolutionary history and phylogenetic uncertainty in our analyses. Briefly, Jetz and Pyron (2018)

used PASTIS (Thomas *et al.,* 2013) which combined taxonomic information from AmphibiaWeb

and a molecular supermatrix of 5 mitochondrial and 10 nuclear genes (Pyron, 2014; Jetz &

Pyron, 2018) to estimate a posterior distribution of trees. We collected microhabitat and

maximum male body sizes (snout-vent length) from the primary literature and other datasets

(Moen & Wiens, 2017; Womack & Bell, 2020; AmphibiaWeb, 2022; IUCN, 2022). Microhabitat

categories included aquatic, arboreal, burrowing, terrestrial, torrential, semi-arboreal, semi-

aquatic, and semi-burrowing (Moen & Wiens, 2017). A list of the sources cited in data tables is

found in Appendix 1.

Statistical analysis and visualization

We inspected the latitude and climate data in many ways to understand their

relationships in a biogeographical context. All analyses and visualizations were done in R v4.2.1

(R Core Team, 2022). First, we plotted species means for each climate variable against

absolute mean species latitude to characterize the latitudinal climate gradient in our data. Next,

we centered and scaled each climate variable to perform principal component analysis (PCA)

and understand the covariances among climate variables across species distributions. We also

visualized spatial variation of breeding periods and climate data by estimating per-pixel mean

breeding period and principal component scores at 100 km^2 resolution using letsR v.4.0 (Vilela

& Villalobos, 2015).

To understand how breeding periods and absolute latitudes are distributed across the

phylogeny of frogs, we plotted average species breeding periods and absolute latitude along the

tips of the consensus tree of Jetz and Pyron (2018) using *ggtree* v3.4.0 (Yu *et al.,* 2017; Yu *et*

al., 2018; Yu, 2020). To determine the extent to which phylogenetic history might explain our

data, we estimated the phylogenetic signal (Adams, 2014) of breeding periods and absolute

latitude and using *geomorph* v4.0.1 (Adams *et al.,* 2021; Baken *et al.,* 2021b). Additionally, we

also obtained summary statistics of breeding periods and absolute latitudes for the four largest

families represented in our dataset: Hylidae (N = 98), Myobatrachidae (N = 86), Bufonidae (N =

62), and Ranidae (N = 44). To test for a latitudinal gradient, we regressed breeding periods onto

absolute latitude using phylogenetic generalized least squares (PGLS; Grafen, 1989; Martins &

Hansen, 1997). PGLS regression uses a phylogenetic covariance matrix to account for residual

correlation due to shared evolutionary history. Furthermore, we implemented PGLS regression

through *RRPP* v1.3.1 (Collyer & Adams, 2018, 2019) which uses a residual randomized

permutation procedure to fit linear models, evaluate significance, and calculate empirical effect

sizes for model terms in all models. We used 9,999 iterations to evaluate significance in all

models.

We tested whether colder, dryer, and more seasonal environments are associated with

shorter breeding periods using principal component (PC) regression. PC regression avoids

collinearity issues while helping us understand how breeding periods are linked to environments

defined by covariation among climate variables (PC loadings). We combined principal

component regression with PGLS regression by regressing species means of breeding period

onto all seven principal components. We then compared the fit of this model to the fit of a

phylogenetic ANCOVA model including body size, microhabitat, and interactions between body

size and each principal component to determine how the relationship between breeding period

and climate might depend on body size. We also performed pairwise comparisons of breeding

periods among microhabitat categories to determine whether burrowing lifestyles are associated

with shorter breeding periods. To account for multiple comparisons, we controlled Type I error

rates ($\alpha = 0.05$) using the Sidak-like step-down procedure in *mutoss* v0.1-12 (MuToss Coding

Team *et al.,* 2017). This method is more powerful than both the Sidak and Bonferroni methods

(Holland & Copenhaver, 1987). We used the natural logarithm of body size to account for

evolutionary allometry of body size. To test the zone hypothesis, we repeated the same

modeling procedures above for two subsets containing either temperate or tropical zone

species.

Finally, we evaluated whether the phylogenetic ANCOVA results above were robust to

phylogenetic uncertainty. Following Baken *et al.* (2021a), we fit new PGLS models using each of

the 1,000 trees from Jetz and Pyron (2018) to obtain *Z*-scores for each independent variable.

This procedure resulted in 1,000 *Z*-scores for each model term from which we calculated the

proportion of trees for which the effect size was significant corresponding to $Z > 1.645$ for

empirically generated *Z*-scores (Collyer & Adams, 2013; Collyer *et al.,* 2015).

198   Results

Climate data showed latitudinal and non-latitudinal differences

Relationships between species means of each climate variable and absolute latitude

(Fig. 1) showed climate differences across the tropical and temperate zones. Overall, frogs in

lower (tropical) latitudes live in warmer, wetter, and less seasonal areas, relative to frogs at

higher latitudes. One important exception is that tropical frogs experience the most precipitation

seasonality. Precipitation seasonality also increases with latitude in tropical frog habitats.

Additionally, annual precipitation and precipitation seasonality is more variable than annual

mean temperature and temperature seasonality among tropical frogs. Absolute latitude was

highly correlated ($r = -0.94$; Fig. 2) with PC1 (54% variance; Table 1). However, 46% variance

(PC's 2–7) was independent of latitude (PC1).

Breeding periods exhibit a latitudinal gradient

211
We examined phylogenetic patterns of variation and covariation for mean breeding period

(mean = 148 days, range = 1–365 days, std. error = 5 days) and absolute latitude (mean =

27.17° N, range = 0.03–64.77° N, std. error = 0.71°). Each trait exhibited a weak phylogenetic

signal, including breeding period (K = 0.19, p = 0.0001; Fig. 3) and absolute latitude (K = 0.51, p

= 0.0001). Taxon-specific averages of breeding periods and absolute latitude matched the

prediction of finding longer breeding periods at lower latitudes: hylid treefrogs (1.92° N) bred for

187 days, bufonid toads (18° N) bred for 114 days, and ranid water frogs (44° N) bred for 110

days. However, Australian and southeast Asian myobatrachid frogs bred for 143 days despite

living in the temperate zone (34° S). Summary statistics suggested tropical species bred for

longer periods, on average (mean = 174 days, range = 5–365 days, std. error = 7 days), relative

to temperate species (mean = 128 days, range = 1–365 days, std. error = 5 days). In fact, we

found longer breeding periods were weakly associated with lower latitudes (F = 16.60, r^2 = 0.03,

Z = 3.09, p = 0.0004).

Global scale analyses reveal latitude, climate, and body size drive breeding period diversity

We found breeding periods exhibited a latitudinal gradient (Fig. 4a). However, the effect

of climate on breeding periods depended on body size (Fig. 4b; Fig. S1.1 in Appendix S1), as

we found an interaction between climate PC1 and body size (r^2 = 0.07; Table 2, Rows 1, 8, 10).

Relatively larger frogs exhibited longer breeding periods in the tropics (high values of PC1)

defined by warm and wet environments with less temperature seasonality, compared to

relatively smaller frogs which exhibited longer breeding periods in the temperate zone (low

values of PC1). Notably, the geographical distribution of climate represented as scores on PC1

exhibited a strong latitudinal gradient (Fig. 4c) and this is consistent with the strong negative

correlation with absolute latitude. We also determined the climate-only model ($r^2 = 0.13$) was

significantly improved (F = 2.88, p = 0.0003) by adding body size, microhabitat, and interactions

between body size and climate ($r^2 = 0.21$). Finally, we found burrowing frogs, which tend to

inhabit desert microhabitats, did not exhibit relatively shorter breeding periods after correcting

for multiple comparisons at the global scale (or within the temperate or tropical zones; Tables

S1.1–3, Fig. S1.2).

Three other key findings were independent of latitude (PC1). First, relatively larger frogs

exhibited longer breeding periods in rainy environments with low topographical wetness (high

values of PC2), while relatively smaller frogs had longer breeding periods in dryer environments

with high topographical wetness (low values of PC2; $r^2 = 0.04$; Table 2, Rows 2, 8, 11, Fig.

S1.3). Second, we found a tendency for species living in topographically wet environments with

low rain seasonality (high values of PC3) to breed for longer periods ($r^2 = 0.04$; Table 2, Row 3;

Fig. S1.4). Although PC3 showed an interaction with body size (Table 2, Row 12), this result

was not robust to phylogenetic uncertainty (see below). Third, we found longer breeding periods

were associated with warmer environments with limited potential for evapotranspiration (high

values of PC7; Table 2, Row 8; Fig. S1.5). Importantly, although PC7 explained <1% of climate

variance (Table 1), it explained $r^2 = 0.01$ of differences in breeding periods and had an effect

size ($Z = 2.32$) comparable to other PC's. All significant results described for this global-scale

analysis were supported in 59.7–89.9% of sampled phylogenetic trees (Fig. S1.6).

Temperate and tropical patterns are not identical to global scale patterns

We found three notable differences in the factors affecting breeding periods which

differed between the global scale analysis and the temperate zone-specific analysis. First, we

found a size-independent pattern where temperate species closest to the equator (high values

of PC1) exhibited longer breeding periods ($r^2 = 0.11$; Table S1.4, Row 1, 10; Fig. S1.7). Second,

we found that relatively larger temperate zone species exhibited longer breeding periods in

topographically wet environments with low rain seasonality (high values of PC3), while relatively

smaller species exhibited longer breeding periods in topographically dry environments with high

rain seasonality (low values of PC3; r^2 = 0.08; Table S1.4, Rows 3, 8, 12; Fig. S1.8). Third, we

found a size-independent pattern where temperate zone species living in rainy areas with low

potential evapotranspiration (high values of PC4) exhibited longer breeding periods (r^2 = 0.03;

Table S1.4, Row 4; Fig. 5a,b). Furthermore, as in the global scale analysis, we found a size-

dependent relationship between breeding period and PC2 (r^2 = 0.04; Table S1.4, Rows 2, 8, 11;

Fig. S1.9). Each of these results were supported in 62.1–97.4% of sampled phylogenies (Fig.

S1.10). We also determined the climate-only temperate zone model (r^2 = 0.20) was significantly

improved (F = 3.59, p = 0.0001) by adding body size, microhabitat, and interactions between

body size and climate (r^2 = 0.34).

270   Unlike the global and temperate zone patterns, body size did not influence any

relationships between breeding period and climate in the tropical zone. As found at the global

scale, we found 1) tropical species exhibited longer breeding periods in environments with low

precipitation seasonality and high topographic wetness (high values of PC3; r^2 = 0.04; Table

S1.5 Row 3; Fig. S1.11) and 2) species bred for longer periods in warmer areas with low

potential for evapotranspiration (high values of PC7; r^2 = 0.03; Table S1.5 Row 7; Fig. S1.12).

Opposite to temperate species, tropical species exhibited shorter breeding periods in rainy

environments with low topographical wetness (high values of PC4; r^2 = 0.03; Table S1.5 Row 4;

Fig. 5c,d). Each of these patterns were supported in 53.1–90.6% of sampled phylogenies (Fig.

S1.13).

Discussion

281   We used phylogenetic comparative methods to show how breeding periods in frogs are

related to latitude, climate, and body size. We found longer breeding periods are generally

associated with warmer, wetter, and less seasonal climates and body size can alter these

behavior by environment relationships. However, body size and latitude were not major factors

in the tropics. Overall, we found a low phylogenetic signal for breeding period and our findings

support the climate and climate-size hypotheses. The zone and microhabitat hypotheses were

partially and not supported, respectively. Both temperature and precipitation were important

factors related to breeding periods globally, but the manner through which they impacted

breeding periods differed between the temperate and tropical zones. These findings suggest

that global ecology, but not phylogenetic history, is an important determinant of breeding

periods in frogs.

Support for the climate-size hypothesis might be related to food availability. Warmer,

wetter, and less seasonal environments may create seasonal fluxes of food resulting in

environmentally constrained growth and reproduction, under the pace of life framework (Hille &

Cooper, 2015). As shown previously, greater seasonality in temperature and precipitation is

associated with decreased food availability (Lack, 1948; Dunn *et al.,* 2000). Thus, since frogs

invest more resources into reproduction when there is a greater food supply (Girish & Saidapur,

2000; Lardner & Loman, 2003), we may reasonably expect food resources to affect breeding

periods. However, size-dependent (and food-independent) climate effects might explain why

relatively smaller species breed for longer periods at higher latitudes if we consider the

importance of energetic limitations during and after hibernation. First, frogs use metabolic

energy from fat stores to overwinter and relatively smaller species have smaller fat stores

(Brodeur *et al.,* 2020). If smaller species begin to run low on metabolic energy during

hibernation, this might result in earlier emergence from hibernation and a longer breeding period

relative to larger species. Second, relatively larger species might warm slowly after emergence

from hibernation such that longer and colder winters might restrict their breeding season,

relative to smaller frogs. This prediction arises from the 'heat balance hypothesis' which

describes how thermoconformers may increase heat transfer through smaller body sizes at

higher latitudes (Olalla-Tarraga *et al.,* 2006; Olalla-Tárraga 2011; Castro *et al.,* 2021).

Global breeding periods are composites of temperate and tropical ecology

While latitudinal gradients are typically thought of as global patterns, our study shows

how global patterns of breeding period are composite patterns representing ecological dynamics

that are different in temperate and tropical environments. Importantly, the global latitudinal

gradient seems to reflect a latitudinal gradient in the temperate region (see Figs. 4,S1.7), as this

gradient is lacking in the tropics. We also found the combined effects of body size and latitude

(PC1) on global periods to be an emergent global property since we did not find this interaction

within either the temperate or tropical zone. This may occur because the tropics as a whole are

much warmer and wetter with less temperature seasonality compared to the temperate zone,

such that differences in climate or food availability might result in unique responses of

differently-sized species across latitudes and zones.

Body size and climate effects on breeding periods of tropical or temperate frogs account

for global patterns. For example, we found the effects on breeding periods of regions with

contrasting rain and topographic wetness (PC2) depended on body size. These effects were

identical between the temperate zone and global scale. This result is consistent with the

climate-size hypothesis and shows how breeding periods are determined by the combined

effects of rain, topography, and body size. Additionally, longer breeding periods at the global

scale resulting from less precipitation seasonality and increased topographic wetness (PC3)

seem attributable to the tropics (Fig. S1.11) and large species in the temperate zone (Fig. S1.8).

This effect is consistent with the climate-size hypothesis, but may also reflect the importance of

body size in ephemeral water bodies used for amphibian reproduction in the temperate region.

Flat, rainy, and unephemeral regions may contain fish predators which may select for slower

growth rates and larger body sizes at metamorphosis due to predator-prey size asymmetry

(Hecnar & M'Closkey, 1997; Lardner, 2000; Jara *et al.,* 2019). We suspect predation may

shorten breeding periods in smaller species, relative to larger species which benefit from less

predator-prey size asymmetry. Notably, smaller frogs are predated more often in permanent

water bodies (Kats *et al.,* 1988; Magnusson & Hero, 1991; Summers & McKeon, 2004) and

frogs reach metamorphosis slower and adulthood at a larger size in permanent water bodies

filled with predators (Lardner, 2000). Finally, the global effect of warmer temperatures and

diminished potential evapotranspiration (positive PC7 scores) resulting in longer breeding

periods is due to the same effect in the tropics, but not in the temperate zone and is consistent

with the climate hypothesis.

Biogeographical context is crucial for understanding the drivers of breeding period

Complementing phylogenetic comparative methods with biogeographical approaches

helps us understand how breeding periods (Fig. 4a) covary with climate through space. We

found that frog breeding periods were shorter in the Australian Outback and longer in the

southwestern Himalayas compared to regions at the same latitude. These observations coincide

with dry, monsoonal regions of the northern Outback and monsoons in the southwestern

Himalayas (Jeelani *et al.,* 2017), given that PC2 was a contrast between precipitation and

topographic wetness. These interpretations match known reproductive traits in amphibians,

birds, and fishes in the Outback and Himalayas (Zann *et al.,* 1995; Bentley, 1966; Joshi *et al.,*

2018; Ellepola *et al.,* 2021). Second, breeding periods showed a reverse-latitudinal trend in

Australia and a longitudinal trend in South America and the United States. The contrast of rain

seasonality and topographic wetness (PC3) in association with mountains or plains and river

basins, seems to account for longer breeding periods in southern Australia, eastern United

States, and in the Amazon and Pampas regions of South America. Notably, these features

might explain why myobatrachid frogs (endemic to Australia and South East Asia) exhibited

longer-than-expected breeding periods, relative to their average latitude. Importantly, breeding

activity of frogs in northern Australia depends on heavy rains and local flooding (Tyler, 1998).

Lastly, the positive effect of PC7 on global breeding periods is due to warmer temperatures and

lower potential evapotranspiration. This is seemingly associated with tropical coastlines or

islands and could be related to how increased conductive heat flux from exposed ground

decreases evaporation in areas without canopy cover (see Lister & Garcia, 2018)

Unlike each of the previous examples, we found one variable (PC4) had opposite effects

in the temperate and tropical regions. In the temperate zone, more rain and less potential

evapotranspiration (high values of PC4) was conducive to longer breeding periods. This pattern

is apparent in comparing the eastern and western parts of Europe, North America, and South

America (e.g., the Chaco and Pampas regions). In the tropics, regions with less rain and more

potential evapotranspiration led to longer breeding periods which might account for shorter than

expected (relative to latitude) breeding periods in the Amazon, West Africa, southwestern India,

and in the island of Borneo. These results support the climate hypothesis in the temperate zone

but not in the tropical zone and might be explained if vegetative growth (shelter and breeding

sites), insect growth (food), and breeding periods are restricted by too much rain and too little

sunlight (low potential evapotranspiration). Typically, one might predict that less rain, not more

rain, restricts plant growth. However, Huete et al. (2006) found plant growth in the Amazon

rainforest is enhanced in dryer and more sunlit conditions. Thus, we infer growing seasons are

longer in dryer and more sunlit environments in other tropical rainforests too, which can shorten

breeding seasons in tropical frogs living in rainy environments with little sunlight. These

interpretations represent hypotheses to be tested in future studies since the maps reflect

community-level spatial variation while the comparative analyses summarize this variation into

species means. Nonetheless, we expect both species and communities to show similar

biogeographic trends relating breeding period and climate. Thus, interpreting comparative

results in a biogeographic context is crucial for understanding the origins of trait variation.

Breeding period gradients are context-dependent

In this study, we have shown how breeding periods in temperate and tropical regions are

differentially impacted by climate and body size. Sometimes, the same variable may have

opposite effects in each region and this may depend on how climate affects plant growth and

insect abundance. This study adds to the growing body of evidence describing how the effects

of environmental gradients are sometimes context-dependent (e.g., Gutiérrez-Pinto *et al.,* 2014;

Chaudhary *et al.,* 2016; Cohen *et al.,* 2018; Juarez *et al.,* 2019; Martinez *et al.,* 2021). While

environmental gradients have historically been treated as one dimensional hypotheses, natural

selection is inherently multivariate and acts on combinations of traits (see Adams & Collyer,

2019) such as breeding periods and body sizes in frogs. Longer breeding periods are

associated with more egg clutches (Bull & Shine, 1979; Morrison & Hero, 2003) and larger frogs

exhibit larger clutches and eggs (Prado & Haddad, 2005; Silva *et al.,* 2020). Therefore, we may

have found that larger frogs always exhibited longer breeding periods. Instead, our results

suggest trade-offs between breeding periods and body size relative to how climate and

physiology affect breeding periods across environmental gradients. Similar findings have been

reported for seasonally reproducing crustaceans which show trade-offs between feeding season

length and body size relative to other reproductive traits and environmental seasonality

(Ejsmond *et al.,* 2018).

A few other aspects of ecology seem to play an important role in determining trade-offs

between breeding periods and body size. For example, shorter breeding periods in smaller

species living in rainier environments with lower topographic wetness (high scores on PC2)

might be related to plant diversity or predation. PC2 (Fig. S1.3) is characteristic of high-altitude

rainforests (cloud forests) such as those found in Central and South America, which hold much

of the world's anuran diversity (Duellman, 1988; Giaretta *et al.,* 1999; Cortés *et al.,* 2008).

Therefore, areas with less topographic wetness and correspondingly low vascular plant species

richness (Sørensen *et al.,* 2005) might limit reproductive opportunities for smaller frogs

associated with leaf- or phytotelma-breeding (Donnelly & Guyer, 1994; Schulte *et al.,* 2010;

Poelman *et al.,* 2013). Predation risk may also differ between small and large species, where

larger species are less reliant on canopy cover due to their large size reducing predation

pressure (Nakazawa *et al.,* 2013), resulting in longer breeding periods. Alternatively, smaller

frogs might face shorter breeding periods due to temporal partitioning of breeding times with

other species and this depends on reproductive mode (Gottsberger & Gruber, 2004). Future

research investigating the roles of plant and animal ecology on breeding periods seems likely to

further uncover the roles of climate and topography in ectothermic seasonal breeders.

Further study of the relationships between body size, reproductive traits, and climate

would result in a better understanding of breeding period diversity. For example, body size is

related to various reproductive traits which may exhibit trade-offs themselves, including many

reproductive life history traits (Furness *et al.,* 2022). Additionally, previous work has found an

association between decreased fecundity and warm, wet climates in terrestrial egg-laying

species (Gomez-Mestre *et al.,* 2012). Notably, some species may have evolved larger clutches

in response to seasonal environments (Sheridan, 2009). Accounting for covariation in traits and

environments is crucial for understanding the contexts and mechanisms which govern changes

in seasonal reproduction.

References

Adams, D.C. (2014). A generalized K statistic for estimating phylogenetic signal from shape and
other high-dimensional multivariate data. *Systematic Biology*, **63**, 685–697.

Adams, D.C. & Collyer, M.L. (2019). Phylogenetic comparative methods and the evolution of
multivariate phenotypes. *Annual Review of Ecology, Evolution, and Systematics*, **50**,
405–425.

Adams, D.C., Collyer, M.L., Kaliontzopoulou, A. & Baken E.K. (2021). geomorph: Software for
geometric morphometric analyses. R package version 4.0.1. https://cran.r-
project.org/package=geomorph.

AmphibiaWeb. (2022). *Amphibian Species By the Numbers.* Available at:
https://amphibiaweb.org/amphibian/speciesnums.html. University of California, Berkeley,
CA, USA. Last accessed 12 July 2022.

Badeck, F.-W., Bondeau, A., Bottcher, K., Doktor, D., Lucht, W., Schaber, J. & Sitch, S. (2004).
Responses of spring phenology to climate change. *The New Phytologist*, **162**, 295–309.

Baken, E.K., Mellenthin, L.E. & Adams, D.C. (2021a). Is salamander arboreality limited by
broad-scale climatic conditions? *PloS One*, **16**, e0255393.

Baken, E.K., Collyer, M.L., Kaliontzopoulou, A. & Adams, D.C. (2021b). geomorph v4.0 and
gmShiny: enhanced analytics and a new graphical interface for a comprehensive
morphometric experience. *Methods in Ecology and Evolution*, **12**, 2355–2363.

Beattie, R.C. (1985). The date of spawning in populations of the Common frog (*Rana
temporaria*) from different altitudes in northern England. *Journal of Zoology*, **205**, 137–
154.

Benard, M.F. (2015). Warmer winters reduce frog fecundity and shift breeding phenology, which
consequently alters larval development and metamorphic timing. *Global Change
Biology*, **21**, 1058–1065.

Bentley, P.J. (1966). Adaptations of amphibia to arid environments. *Science*, **152**, 619–623.

Blackburn, T.M. & Hawkins, B.A. (2004). Bergmann's rule and the mammal fauna of northern
North America. *Ecography*, **27**, 715–724.

Blomberg, S.P., Garland, T., Jr & Ives, A.R. (2003). Testing for phylogenetic signal in
comparative data: behavioral traits are more labile. *Evolution*, **57**, 717–745.

Brodeur, J.C., Vera Candioti, J., Damonte, M.J., Bahl, M.F., Poliserpi, M.B. & D'Andrea, M.F.
(2020). Frog somatic indices: Importance of considering allometric scaling, relation with
body condition and seasonal variation in the frog *Leptodactylus latrans*. *Ecological
Indicators*, **116**, 106496.

Bull, J.J. & Shine, R. (1979). Iteroparous animals that skip opportunities for reproduction.
American Naturalist, **114**, 296–303.

Butt, N., Seabrook, L., Maron, M., Law, B.S., Dawson, T.P., Syktus, J. & McAlpine, C.A. (2015).

Cascading effects of climate extremes on vertebrate fauna through changes to low-latitude tree flowering and fruiting phenology. *Global Change Biology*, **21**, 3267–3277.

Castro, K.M.S.A., Amado, T.F., Olalla-Tárraga, M.Á., Gouveia, S.F., Navas, C.A. & Martinez, P.A. (2021). Water constraints drive allometric patterns in the body shape of tree frogs. *Scientific Reports*, **11**, 1218.

Chaudhary, C., Saeedi, H. & Costello, M.J. (2016). Bimodality of latitudinal gradients in marine species richness. *Trends in Ecology & Evolution*, **31**, 670–676.

Cohen, J.M., Lajeunesse, M.J. & Rohr, J.R. (2018). A global synthesis of animal phenological responses to climate change. *Nature Climate Change*, **8**, 224–228.

Collyer, M.L. & Adams, D.C. (2019). RRPP: Linear model evaluation with randomized residuals in a permutation procedure. https://CRAN.R-project.org/package=RRPP.

Collyer, M.L. & Adams, D.C. (2018). RRPP: An R package for fitting linear models to high‐dimensional data using residual randomization. *Methods in Ecology Evolution,* **9**, 1772–1779.

Collyer, M.L. & Adams, D.C. (2013). Phenotypic trajectory analysis: comparison of shape change patterns in evolution and ecology. *Hystrix* , **24**, 75.

Collyer, M.L., Sekora, D.J. & Adams, D.C. (2015). A method for analysis of phenotypic change for phenotypes described by high-dimensional data. *Heredity*, **115**, 357–365.

Cortés, A.M., Ramírez-Pinilla, M.P., Suárez, H.A. & Tovar, E. (2008). Edge effects on richness, abundance and diversity of frogs in Andean cloud forest fragments. *South American Journal of Herpetology*, **3**, 213–222.

Daru, B.H., Kling, M.M., Meineke, E.K. & van Wyk, A.E. (2019). Temperature controls phenology in continuously flowering *Protea* species of subtropical Africa. *Applications in Plant Sciences*, **7**, e01232.

Demment, M.W. & Van Soest, P.J. (1985). A nutritional explanation for body-size patterns of ruminant and nonruminant herbivores. *American Naturalist*, **125**, 641–672.

Donnelly, M.A. & Guyer, C. (1994). Patterns of reproduction and habitat use in an assemblage of Neotropical hylid frogs. *Oecologia*, **98**, 291–302.

Duellman, W.E. (1988). Patterns of species diversity in anuran amphibians in the American tropics. *Annals of the Missouri Botanical Garden*, **75**, 79–104.

Dunn, P.O., Thusius, K.J., Kimber, K. & Winkler, D.W. (2000). Geographic and ecological variation in clutch size of tree swallows. *The Auk*, **117**, 215–221.

Ejsmond, M.J., McNamara, J.M., Søreide, J. & Varpe, Ø. (2018). Gradients of season length and mortality risk cause shifts in body size, reserves and reproductive strategies of determinate growers. *Functional Ecology*, **32**, 2395–2406.

Ellepola, G., Pie, M.R., Pethiyagoda, R., Hanken, J. & Meegaskumbura, M. (2021). Drivers of an epic radiation: the role of climate and islands in species diversification and reproductive-mode evolution of Old-World tree frogs. *bioRxiv*, 2021.09.22.461377.

Fick, S.E. & Hijmans, R.J. (2017). WorldClim 2: new 1‑km spatial resolution climate surfaces
for global land areas. *International Journal of Climatology*, **37**, 4302–4315.

Forti, L.R., Hepp, F., de Souza, J.M., Protazio, A. & Szabo, J.K. (2022). Climate drives anuran
breeding phenology in a continental perspective as revealed by citizen‑collected data.
Diversity & Distributions, **28**, 2094–2109.

Furness, A.I., Venditti, C. & Capellini, I. (2022). Terrestrial reproduction and parental care drive
rapid evolution in the trade-off between offspring size and number across amphibians.
PLoS Biology, **20**, e3001495.

GBIF.org. (2021a). Data from: GBIF Occurrence Download. Available at:
https://doi.org/10.15468/dl.354f8k. Accessed: 03 October 2021.

GBIF.org. (2021b). Data from: GBIF Occurrence Download. Available at:
https://doi.org/10.15468/dl.pdjqte. Accessed: 03 October 2021.

GBIF.org. (2021c). Data from: GBIF Occurrence Download. Available at:
https://doi.org/10.15468/dl.nbku3v. Accessed: 03 October 2021.

GBIF.org. (2021d). Data from: GBIF Occurrence Download. Available at:
https://doi.org/10.15468/dl.uejpg6. Accessed: 03 October 2021.

Giaretta, A.A., Facure, K.G., Sawaya, R.J., Meyer, J.H.D.M. & Chemin, N. (1999). Diversity and
abundance of litter frogs in a montane forest of southeastern Brazil: seasonal and
altitudinal changes. *Biotropica*, **31**, 669–674.

Girish, S. & Saidapur, S.K. (2000). Interrelationship between food availability, fat body, and
ovarian cycles in the frog, *Rana tigrina*, with a discussion on the role of fat body in
anuran reproduction. *Journal of Experimental Zoology*, **286**, 487–493.

Gomez-Mestre, I., Pyron, R.A. & Wiens, J.J. (2012). Phylogenetic analyses reveal unexpected
patterns in the evolution of reproductive modes in frogs. *Evolution*, **66**, 3687–3700.

Gottsberger, B. & Gruber, E. (2004). Temporal partitioning of reproductive activity in a
neotropical anuran community. *Journal of Tropical Ecology*, **20**, 271–280.

Grafen, A. (1989). The phylogenetic regression. *Philosophical Transactions of the Royal Society
of London. Series B, Biological sciences*, **326**, 119–157.

Gutiérrez-Pinto, N., McCracken, K.G., Alza, L., Tubaro, P., Kopuchian, C., Astie, A. & Cadena,
C.D. (2014). The validity of ecogeographical rules is context-dependent: testing for
Bergmann's and Allen's rules by latitude and elevation in a widespread Andean duck.
Biological Journal of the Linnean Society, **111**, 850–862.

Harvey, P.H. & Purvis, A. (1991). Comparative methods for explaining adaptations. *Nature*, **351**,
619–624.

Hecnar, S.J. & M'Closkey, R.T. (1997). The effects of predatory fish on amphibian species
richness and distribution. *Biological Conservation*, **79**, 123–131.

Hille, S.M. & Cooper, C.B. (2015). Elevational trends in life histories: revising the pace-of-life
framework. *Biological Reviews of the Cambridge Philosophical Society*, **90**, 204–213.

Holland, B.S. & Copenhaver, M.D. (1987). An improved sequentially rejective Bonferroni test
procedure. *Biometrics*, **43**, 417–423.

Huete, A.R., Didan, K., Shimabukuro, Y.E., Ratana, P., Saleska, S.R., Hutyra, L.R., Yang, W.,
Nemani, R.R. & Myneni, R. (2006). Amazon rainforests green-up with sunlight in dry
season. *Geophysical Research Letters*, **33**, 1–4.

IUCN. (2022). Data from: The IUCN Red List of Threatened Species, version 2021-3. Available
at: https://www.iucnredlist.org.

Jara, F.G., Thurman, L.L., Montiglio, P.-O., Sih, A. & Garcia, T.S. (2019). Warming-induced
shifts in amphibian phenology and behavior lead to altered predator-prey dynamics.
Oecologia, **189**, 803–813.

Jeelani, G., Deshpande, R.D., Shah, R.A. & Hassan, W. (2017). Influence of southwest
monsoons in the Kashmir Valley, western Himalayas. *Isotopes in Environmental and
Health Studies*, **53**, 400–412.

Jetz, W. & Pyron, R.A. (2018) The interplay of past diversification and evolutionary isolation with
present imperilment across the amphibian tree of life. *Nature Ecology & Evolution*, **2**,
850–858.

Joshi, K.D., Das, S.C.S., Pathak, R.K., Khan, A., Sarkar, U.K. & Roy, K. (2018). Pattern of
reproductive biology of the endangered golden mahseer *Tor putitora* (Hamilton 1822)
with special reference to regional climate change implications on breeding phenology
from lesser Himalayan region, India. *Journal of Applied Animal Research*, **46**, 1289–
1295.

Juarez, B.H., Speiser, D.I. & Oakley, T.H. (2019). Context-dependent evolution of ostracod
morphology along the ecogeographical gradient of ocean depth. *Evolution*, **73**, 1213–
1225.

Kaspari, M., Pickering, J. & Windsor, D. (2001). The reproductive flight phenology of a
neotropical ant assemblage. *Ecological Entomology*, **26**, 245–257.

Kats, L.B., Petranka, J.W. & Sih, A. (1988). Antipredator defenses and the persistence of
amphibian larvae with fishes. *Ecology*, **69**, 1865–1870.

Lack, D. (1948). The significance of clutch-size. Part III.-Some interspecific comparisons. *The
Ibis*, **90**, 25–45.

Lardner, B. (2000). Morphological and life history responses to predators in larvae of seven
anurans. *Oikos*, **88**, 169–180.

Lardner, B. & Loman, J. (2003). Growth or reproduction? Resource allocation by female frogs
Rana temporaria. *Oecologia*, **137**, 541–546.

Lister, B.C. & Garcia, A. (2018). Climate-driven declines in arthropod abundance restructure a
rainforest food web. *Proceedings of the National Academy of Sciences of the United
States of America*, **115**, E10397–E10406.

Magnusson, W.E. & Hero, J.-M. (1991). Predation and the evolution of complex oviposition
behaviour in Amazon rainforest frogs. *Oecologia*, **86**, 310–318.

Martinez, C.M., Friedman, S.T., Corn, K.A., Larouche, O., Price, S.A. & Wainwright, P.C.
(2021). The deep sea is a hot spot of fish body shape evolution. *Ecology Letters*, **24**,
1788–1799.

Martins, E.P. & Hansen, T.F. (1997). Phylogenies and the comparative method: a general
approach to incorporating phylogenetic information into the analysis of interspecific data.
American Naturalist, **149**, 646–667.

Moen, D.S. & Wiens, J.J. (2017). Microhabitat and climatic niche change explain patterns of
diversification among frog families. *American Naturalist*, **190**, 29–44.

Morrison, C. & Hero, J.-M. (2003). Geographic variation in life-history characteristics of
amphibians: a review. *Journal of Animal Ecology*, **72**, 270–279.

MuToss Coding Team, Blanchard, G., Dickhaus, T., Hack, N., Konietschke, F., Rohmeyer, K.,
… *Werft, W.* (2017). mutoss: unified multiple testing procedures. R package version 0.1-
12. https://CRAN.R-project.org/package=mutoss.

Nakazawa, T., Ohba, S.-Y. & Ushio, M. (2013). Predator-prey body size relationships when
predators can consume prey larger than themselves. *Biology Letters*, **9**, 20121193.

Neveu, A. (2009). Incidence of climate on common frog breeding: long-term and short-term
changes. *Acta Oecologica*, **35**, 671–678.

Olalla-Tárraga, M.Á. (2011). "Nullius in Bergmann" or the pluralistic approach to
ecogeographical rules: a reply to Watt et al. (2010). *Oikos*, **120**, 1441–1444.

Olalla-Tarraga, M.A., Rodriguez, M.A. & Hawkins, B.A. (2006). Broad-scale patterns of body
size in squamate reptiles of Europe and North America. *Journal of Biogeography*, **33**,
781–793.

Pianka, E.R. (1966). Latitudinal gradients in species diversity: a review of concepts. *American
Naturalist*, **100**, 33–46.

Poelman, E.H., van Wijngaarden, R.P.A. & Raaijmakers, C.E. (2013). Amazon poison frogs
(*Ranitomeya amazonica*) use different phytotelm characteristics to determine their
suitability for egg and tadpole deposition. *Evolutionary Ecology*, **27**, 661–674.

Prado, C.P.A. & Haddad, C.F.B. (2005). Size-fecundity relationships and reproductive
investment in female frogs in the Pantanal, south-western Brazil. *The Herpetological
Journal*, **15**, 181–189.

Pyron, R.A. (2014). Biogeographic analysis reveals ancient continental vicariance and recent
oceanic dispersal in amphibians. *Systematic Biology*, **63**, 779–797.

R Core Team. (2022). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. Available at: https://www.R-project.org/.

Reading, C.J. (2003). The effects of variation in climatic temperature (1980-2001) on breeding
activity and tadpole stage duration in the common toad, *Bufo bufo. Science of the Total
Environment*, **310**, 231–236.

Schemske, D.W., Mittelbach, G.G., Cornell, H.V., Sobel, J.M. & Roy, K. (2009). Is there a

latitudinal fradient in the importance of biotic interactions? *Annual Review of Ecology, Evolution, and Systematics,* **40**, 245–269.

Schulte, Rödder, Schulte & Lötters (2010). Preference and competition for breeding plants in coexisting *Ranitomeya* species (Dendrobatidae): does height play a role. *Salamandra*, **46**, 180–184.

Sheridan, J.A. (2009). Reproductive variation corresponding to breeding season length in three tropical frog species. *Journal of Tropical Ecology*, **25**, 583–592.

Silva, N.R., Berneck, B.V.M., da Silva, H.R., Haddad, C.F.B., Zamudio, K.R., Mott, T., Nali, R.C. & Prado, C.P.A. (2020). Egg-laying site, fecundity and degree of sexual size dimorphism in frogs. *Biological Journal of the Linnean Society,* **131**, 600–610.

Śniegula, S., Gołąb, M.J. & Johansson, F. (2016). Time constraint effects on phenology and life history synchrony in a damselfly along a latitudinal gradient. *Oikos*, **125**, 414–423.

Sørensen, R., Zinko, U. & Seibert, J. (2005). On the calculation of the topographic wetness index: evaluation of different methods based on field observations. *Hydrology and Earth System Sciences Discussions*, **2**, 1807–1834.

Stevens, G.C. (1989). The latitudinal gradient in geographical range: how so many species coexist in the tropics. *American Naturalist*, **133**, 240–256.

Summers, K. & McKeon, C.S. (2004). The evolutionary ecology of phytotelmata use in Neotropical poison frogs. *Miscellaneous Publications, Museum of Zoology, University of Michigan*, **193**, 55–73.

Thomas, G.H., Hartmann, K., Jetz, W., Joy, J.B., Mimoto, A. & Mooers, A.O. (2013). PASTIS: an R package to facilitate phylogenetic assembly with soft taxonomic inferences. *Methods in Ecology and Evolution*, **4**, 1011–1017.

Title, P.O. & Bemmels, J.B. (2018). ENVIREM: an expanded set of bioclimatic and topographic variables increases flexibility and improves performance of ecological niche modeling. *Ecography*, **41**, 291–307.

Todd, B.D., Scott, D.E., Pechmann, J.H.K. & Gibbons, J.W. (2011). Climate change correlates with rapid delays and advancements in reproductive timing in an amphibian community. *Proceedings of the Royal Society B: Biological Sciences*, **278**, 2191–2197.

Tyler, M. J. (1998). *Australian frogs: a natural history.* Ithaca, New York. Cornell University Press.

Vázquez, D.P. & Stevens, R.D. (2004). The latitudinal gradient in niche breadth: concepts and evidence. *American Naturalist*, **164**, E1–19.

Vilela, B. & Villalobos, F. (2015). letsR: a new R package for data handling and analysis in macroecology. *Methods in Ecology and Evolution*, **6**, 1229–1234.

Wcislo, W.T. (1989). Behavioral environments and evolutionary change. *Annual Review of Ecology and Systematics*, **20**, 137–169.

Wells, K.D. (2007). *The ecology and behavior of amphibians.* University of Chicago Press.

Wells, K.D. (1977). The social behaviour of anuran amphibians. *Animal Behaviour*, **25**, 666–
693.

While, G.M. & Uller, T. (2014). Quo vadis amphibia? Global warming and breeding phenology in
frogs, toads and salamanders. *Ecography*, **37**, 921–929.

Wilson, D.S. (1975). The Adequacy of body size as a niche difference. *American Naturalist*,
109, 769–784.

Womack, M.C. & Bell, R.C. (2020). Two□hundred million years of anuran body□size evolution
in relation to geography, ecology and life history. *Journal of Evolutionary Biology*, **33**,
1417–1432.

Yu, G. (2020). Using ggtree to visualize data on tree-like structures. *Current Protocols in*
Bioinformatics, **69**, e96.

Yu, G., Lam, T.T.-Y., Zhu, H. & Guan, Y. (2018). Two methods for mapping and visualizing
associated data on phylogeny using ggtree. *Molecular Biology and Evolution*, **35**, 3041–
3043.

Yu, G., Smith, D., Zhu, H., Guan, Y. & Lam, T.T-Y. (2017). ggtree: an R package for
visualization and annotation of phylogenetic trees with their covariates and other
associated data. *Methods in Ecology and Evolution,* **8**, 28–36.

Zann, R.A., Morton, S.R., Jones, K.R. & Burley, N.T. (1995). The timing of breeding by zebra
finches in relation to rainfall in central Australia. *The Emu*, **95**, 208–222.
681

682 **Data accessibility:** All data and code are publicly available on datadryad.org (LINK TO BE

683 UPDATED AFTER REVIEW).

684
685 **Tables**
686
687 **Table 1. PCA summary and loadings of climate variables.** PCA based on means for 497 species.

688 Cumulative proportion of variance is given in parentheses next to PC heading. Loadings with a magnitude

689 >=0.30 are in bold.

Variable	PC1 (52)	PC2 (76)	PC3 (90)	PC4 (95)	PC5 (97)	PC6 (99)	PC7 (100)
Annual Mean Temperature	**0.47**	-0.27	0.08	-0.01	0.25	0.15	**0.78**
Temperature Seasonality	**-0.47**	-0.14	-0.09	**0.46**	0.23	**0.70**	0.04

Annual Precipitation	**0.38**	**0.38**	0.14	**0.75**	0.25	-0.21	-0.14
Precipitation Seasonality	0.15	-0.25	**-0.90**	0.23	-0.16	-0.16	0.00
Annual Potential Evapotranspiration (PET)	**0.42**	**-0.40**	0.00	-0.22	**0.47**	0.23	**-0.59**
PET Seasonality	**-0.45**	-0.28	0.01	0.03	**0.60**	**-0.59**	0.10
Topographic Wetness	0.01	**-0.67**	**0.40**	**0.36**	**-0.47**	-0.16	-0.10

691
692
Table 2. Global model summary of effects of climate and body size on breeding periods in 497

frog species. Climate shown as principal components (PC's). Body size is snout-vent length (SVL) in

mm. Df = degrees of freedom, SS = sum of squares, MS = mean square, r^2 = coefficient of determination,

F = F-statistic, Z = standardized effect size (empirical Z-score), and p = p-value. Significant terms ($p <$

0.05), assessed using a consensus tree (see Methods), are in bold. Asterisks indicate significant terms

supported in >50% of 1000 phylogenetic trees.

Row	Term	Df	SS	MS	r^2	F	Z	p
1	**PC1***	**1**	**6258.53**	**6258.53**	**0.05**	**26.87**	**3.71**	**0.0001**
2	**PC2***	**1**	**2391.84**	**2391.84**	**0.02**	**10.27**	**2.6**	**0.0017**
3	**PC3***	**1**	**6197.66**	**6197.66**	**0.04**	**26.61**	**3.75**	**0.0001**
4	PC4	1	78.93	78.93	0	0.34	-0.1	0.5480
5	PC5	1	840.89	840.89	0.01	3.61	1.52	0.0585
6	PC6	1	834.98	834.98	0.01	3.59	1.52	0.0603
7	**PC7***	**1**	**1967.68**	**1967.68**	**0.01**	**8.45**	**2.32**	**0.0045**
8	**SVL**	**1**	**1077.61**	**1077.61**	**0.01**	**4.63**	**1.75**	**0.0351**
9	**Microhabitat***	**7**	**4393.72**	**627.67**	**0.03**	**2.69**	**2.24**	**0.0122**
10	**PC1:SVL***	**1**	**1310.68**	**1310.68**	**0.01**	**5.63**	**1.93**	**0.0186**

11	PC2:SVL*	1	1741.24	1741.24	0.01	7.48	2.24	0.0051
12	PC3:SVL	1	1290.44	1290.44	0.01	5.54	1.94	0.0168
13	PC4:SVL	1	9.07	9.07	0	0.04	-1.06	0.8380
14	PC5:SVL	1	117.52	117.52	0	0.5	0.08	0.4837
15	PC6:SVL	1	77.08	77.08	0	0.33	-0.15	0.5688
16	PC7:SVL	1	41.04	41.04	0	0.18	-0.46	0.6736
17	Residuals	474	110398.01	232.91	0.79			
18	Total	496	139026.91					

699
700
701 **Figures**
702
703
Figure 1. Relationships between species means climate and absolute latitude. Generally, tropical

and temperate climates differ substantially relative to absolute latitude when considering annual mean

temperature, temperature seasonality, annual precipitation, precipitation seasonality, potential

evapotranspiration, and potential evapotranspiration seasonality. Red points indicate tropical species and

blue points indicate temperate zone species. Vertical red line corresponds to 23.43628 decimal degrees.

Figure 2. Relationship between species means of absolute latitude and Climate PC1. Blue points

correspond to temperate zone species and red points are tropical zone species. Tropical and temperate

zone cutoff used is 23.43628 decimal degrees. Absolute latitude is strongly and negatively correlated to

PC1. r^2 = 0.88 and p < 0.0001.

Figure 3. Phylogenetic differences in average breeding period and Absolute (Mean) Latitude

across 497 species of frogs. Phylogeny of 497 species with breeding period and absolute mean latitude

mapped at the tips. Node labels identify the 4 highest-sampled families in our dataset. Inner ring shows

species means of breeding period and outer ring shows species means of absolute latitude. Anurans

breed for a mean of 148 days (range = 1–365 days, std. error = 4.5 days). However, temperate species

breed for an average of 127.8 days (median = 107 days, range = 1–365 days, std. error = 5.87 days) and

tropical species breed for an average of 167.3 days (median = 181 days, range = 5–365 days, std. error =

6.52 days).

Figure 4. Global biogeography of breeding periods, climate PC1, and body size in 497 frog

species. a. Mollweide equal-area projection showing pixels corresponding to 100 km^2 areas. Pixel colors

represent mean breeding periods of all species found within a pixel. **b.** The relationship between breeding

period, climate PC1 (latitude), and body size. Breeding periods and PC1 scores are species means.

Regression lines show the positive relationships between breeding periods and PC1 at body sizes

corresponding to the 10th (24 mm) and 90th (80.48 mm) percentiles of snout-vent length after accounting

for phylogenetic relationships. Larger frogs exhibit a steeper relationship between breeding periods and

PC1 and larger species have longer breeding periods in the tropics while smaller species have longer

breeding periods in the temperate zone. PC1 and body size explain r^2 = 0.07 of breeding period diversity

and their interaction is significant (p = 0.0186). High scores on PC1 indicate tropical latitudes and low

scores indicate temperate latitudes. We used a regression of absolute latitude on PC1 to determine the

PC score which corresponds to 23.43628 decimal degrees. Individual points were removed for clarity but

all data is displayed in Fig. S1.1. **c.** Mollweide equal-area projection showing pixels corresponding to 100

km^2 areas. Pixel colors represent PC1 scores of climate found within a pixel.

Figure 5. Biogeography of breeding periods and climate PC4 in the temperate and tropical zones.

a. The marginal (positive) effect of climate PC4 on breeding periods for 279 temperate species. The red

line is the regression line which accounts for phylogenetic relationships. PC4 is a contrast between

annual precipitation and potential evapostranspiration; high scores are rainy areas with low potential

evapotranspiration. Points are species means. **b.** Mollweide equal-area projection showing pixels

corresponding to 100 km^2 areas. Pixel colors represent PC4 scores of climate found within a pixel in the

temperate zone. **c.** The marginal (negative) effect of climate PC4 on breeding periods for 218 tropical

species. The red line is the regression line which accounts for phylogenetic relationships. PC4 is a

contrast between annual precipitation and potential evapotranspiration; high scores are rainy areas with

low potential evapotranspiration. Points are species means. **d.** Mollweide equal-area projection showing

pixels corresponding to 100 km^2 areas. Pixel colors represent PC4 scores of climate found within a pixel

in the tropics.

Bufonidae

Hylidae

Ranidae

Myobatrachidae

Absolute Latitude
(Decimal degrees)   20 40 60

Breeding
Period
(Days)   100 200 300