1   **TITLE: Unified Tumor Growth Mechanisms from Multimodel Inference and Dataset Integration**

2   **SHORT TITLE: Determining tumor growth mechanisms via model hypothesis exploration**

3   **AUTHORS: Samantha P. Beik[1], Leonard A. Harris[2,3,4], Michael A. Kochen[5], Julien Sage[6,7], Vito**

4   **Quaranta[8,9], Carlos F. Lopez[9,10]***

[1] Medical Scientist Training Program, Vanderbilt University School of Medicine, Nashville, TN, 37232, USA

[2] Department of Biomedical Engineering, University of Arkansas, Fayetteville, AR, USA

[3] Interdisciplinary Graduate Program in Cell & Molecular Biology, University of Arkansas, Fayetteville, AR, USA

[4] Cancer Biology Program, Winthrop P. Rockefeller Cancer Institute, University of Arkansas for Medical Sciences, Little Rock, AR, USA

[5] Department of Bioengineering, University of Washington, Seattle, WA, USA

[6] Departments of Pediatrics, Stanford University, Stanford, CA, USA

[7] Departments of Genetics, Stanford University, Stanford, CA, USA

[8] Program in Chemical and Physical Biology, Vanderbilt University, Nashville, TN, USA

[9] Department of Biochemistry, Vanderbilt University, Nashville, TN, USA

[10] Altos Laboratories, Redwood City, CA, USA

* Correspondence: clopez@altoslabs.com

ABSTRACT

Mechanistic models of biological processes can help explain observed phenomena and predict response to a perturbation. A mathematical model is typically constructed using expert knowledge and informal reasoning to generate a mechanistic explanation for a given observation. Although this approach works well for simple systems with abundant data and well-established principles, quantitative biology is often faced with a dearth of both data and knowledge about a process, thus making it challenging to identify and validate all possible mechanistic hypothesis underlying a system behavior. To overcome these limitations, we introduce a Bayesian multimodel inference (Bayes-MMI) methodology, which quantifies how mechanistic hypotheses can explain a given experimental datasets, and concurrently, how each dataset informs a given model hypothesis, thus enabling hypothesis space exploration in the context of available data. We demonstrate this approach to probe standing questions about heterogeneity, lineage plasticity, and cell-cell interactions in tumor growth mechanisms of small cell lung cancer (SCLC). We integrate three datasets that each formulated different explanations for tumor growth mechanisms in SCLC, apply Bayes-MMI and find that the data supports model predictions for tumor evolution promoted by high lineage plasticity, rather than through expanding rare stem-like populations. In addition, the models predict that in the presence of SCLC-N or SCLC-A2 cells, the transition from SCLC-A to SCLC-Y through an intermediate is decelerated. Together, these predictions provide a testable hypothesis for observed juxtaposed results in SCLC growth and a mechanistic interpretation for tumor recalcitrance.

AUTHOR SUMMARY

To make a mathematical model, an investigator needs to know and incorporate biological relationships present in the system of interest. However, if we don't know the exact relationships, how can we build a model? Building a single model may include spurious relationships or exclude important ones, so model selection enables us to build multiple, incorporating various combinations of biological features and the relationships between them. Each biological feature represents a distinct hypothesis, which can be investigated via model fitting to experimental data. We aim to improve upon the information theoretic

framework of model selection by incorporating Bayesian elements. We apply our approach to small cell

lung cancer (SCLC), using multiple datasets, to address hypotheses about cell-cell interactions, phenotypic

transitions, and tumor makeup across experimental model systems. Incorporating Bayesian inference, we

can add into model selection an assessment of whether these hypotheses are likely or unlikely, or even

whether the data enables assessment of a hypothesis at all. Our analysis finds that SCLC is likely highly

plastic, with cells able to transition phenotypic identities easily. These predictions could help explain why

SCLC is such a difficult disease to treat, and provide the basis for further experiments.

INTRODUCTION

A mechanistic understanding of biological processes that explains causal input-output relationships and predicts population behaviors (1) remains a central challenge to all areas of quantitative biology. Mathematical models have become an established practice to specify precise relationships within a biological system, (2) and thereby hypothesize, and subsequently test, the existence of these relationships. For example, multiple mechanistic models of apoptosis execution have been formulated to explore the nature of biochemical interactions that lead to cellular commitment to death, demonstrating that careful model design and suitable data can lead to important biological insights (3–5). A more challenging situation emerges when models are formulated for biological processes that are poorly defined or understood, leading to multiple competing, and often juxtaposed mechanistic explanations for a given biological process. For example, in Small Cell Lung Cancer (SCLC), a study of circulating tumor cell-derived xenografts showed that Non-NE-related subtypes could act as a stemlike population ("source"), (6) but archetype analysis of a genetically engineered mouse model tumor showed the Non-NE subtype SCLC-Y acts as an end-state ("sink", rather than source) (7). Therefore, continued exploration of the hypotheses generated from these works can help elucidate the differences between these and other potential explanations for tumor growth mechanisms.

This phenomenon where multiple mechanistic hypotheses are concurrently proposed but must be assessed with limited data is not restricted to quantitative biology but common to other fields with similar data availability limitations such as ecology (8) climatology (9), and evolutionary biology (10), to name a few. To address this challenge, methods such as model selection and multimodel inference have been proposed using information theoretic scoring techniques such as Akaike Information Criterion, (AIC) with limited success stemming from difficulties with model averaging (11,12) and the fact that AIC scores do not inherently describe whether a model or features within are informed by the data. More recently, the use of AI and machine learning approaches has given impetus to causal relationship inference (13) but these relationships remain difficult to elucidate, thus underscoring the need for both novel tools for hypothesis

exploration, and tools that can be used with rigor in the face of missing data that may not inform all

hypotheses.

The model selection process involves candidate model evaluation from a superset of plausible models,

relative to a given experimental dataset. A scoring function is then applied to each model usually with the

goal to identify the top-scoring model that best represents the process being investigated. A typical approach

is to consider a "best" model as comprising the most relevant variables that capture important mechanistic

aspects of the explored process, while excluded variables capture process features that are less relevant for

the question being explored. However, variables throughout all candidate models can contribute to

knowledge about the overall system (14). In the cases where data is simply less informative for a given set

of hypotheses, uncertainty will remain about what constitutes a "best" model, (15), necessitating approaches

such as model averaging, where parameter values can be weighted by model probability and then combined

into a distribution of likely values. Unfortunately, for information theoretic applications of model

averaging, this probability must be weighted and summed across all possible models, which are often not

possible to enumerate exhaustively in a biological context.

To address the challenge of employing multimodel inference approaches in the context of biological

processes where models cannot be exhaustively enumerated, data may not inform all model evaluation, and

model averaging across all models is desired to learn about the system of interest, we introduce Bayesian

multimodel inference (Bayes-MMI), a method which combines Bayesian inference with model selection

and model averaging. In our approach, Bayesian model selection enables exploration of mechanistic model

hypotheses by inclusion or exclusion of species and behaviors that could play a role in each biological

process. Application of Bayesian principles in turn reveals the extent to which data informs a given model

and its constitutive parameters.

In our system of interest, small cell lung cancer, (SCLC) we integrate the most suitable available datasets

and published theories of SCLC cellular biology to enumerate mechanistic hypotheses for SCLC tumor

growth. We test the resulting thousands of candidate population dynamics models via nested sampling,

comparing candidate model output to tumor steady-state data, applying the principles of model selection

and model averaging for a principled and comprehensive assessment of SCLC mechanistic hypotheses. We

estimate the probability of each mechanistic hypothesis given the data, generating an interpretation of SCLC

tumor growth: highly likely non-hierarchical phenotypic transitions indicating SCLC subtype plasticity,

and less likely cell-cell interactions that affect the rate of phenotypic transitions across subtypes. We show

how certain aspects of the SCLC model, such as phenotypic transitions and cell-cell interactions related to

these, are well informed by the available data, but other aspects, such as tumor initiation and growth rate

effects, are not informed. Our approach is generalizable to other biological systems, and as such we suggest

a shift away from considering only one "best" model or the variables within and instead propose a move

toward Bayesian principles in multimodel inference and a probabilistic understanding of whether each

cellular behavior in a model contributes to the behavior of the system under study.

RESULTS

Existing datasets yield multiple hypotheses in SCLC tumor growth mechanisms

Small cell lung cancer (SCLC) has been denominated a recalcitrant tumor where relapse after treatment is

commonplace and the survival prognosis is typically poor. SCLC comprises ~15% of all lung cancer cases

worldwide and results in ~200,000 deaths annually with a 5-year survival rate of less than 10% (16).

Intratumoral heterogeneity is hypothesized to be the main contributor to the natural history of this disease

and its morbidity and mortality (16–18). SCLC tumors comprise a mix of functionally distinct subtypes of

interacting cells, (19–21), most notably neuroendocrine (NE) and Non-NE. As shown in **Fig 1A**, SCLC

populations comprise a collection of cellular subtypes within a tumor, identified by differential expression

of transcriptional regulators (17).

Fig 1. Conclusions and hypotheses from literature build mechanistic hypothesis exploration space for tumor growth and development. (A) Synthesis of what is currently known about SCLC subtypes, which have been divided into two overall phenotypes, neuroendocrine (NE) and Non-NE, and then further classified into subtypes based on transcription factor expression. (1) NE SCLC cells, which do not express Hes1, transition into Non-NE cells, which do. (2) Hes1+ cells release unidentified factors (gray circle) that support viability and growth of Hes1- cells, and the two Hes1+ and Hes1- populations grow better together rather than separately. (3) Consensus across the field led to labeling SCLC phenotypic subtypes by the dominant transcription factor expressed in that subtype. (4) Subtype with transcriptional signature intermediate between NE and Non-NE, named SCLC-A2. (5) Phenotypic transitions occur in a hierarchical manner from SCLC-A to SCLC-N to SCLC-Y cells. (B)-(E) Candidate model examples representing SCLC biological hypotheses (Table 1). (B) Model topologies constructed with 2+ subtypes, with number of combinations per number of subtypes. (C) Subtype effect schema, where there are different effectors between candidates and different affected cellular actions. (D) Potential initiation schema, where all subtypes in topology must be accessible either as initiating subtypes or via transitions. (E) Potential transition schema where, unidirectional transitions are those that follow a hierarchy, and bidirectional transitions must be symmetrical when present. HES1, Hes Family BHLH Transcription Factor 1; ASCL1, Achaete-scute homolog 1; NEUROD1, neurogenic differentiation factor 1; POU2F3, POU class 2 homeobox 3; YAP1, yes-associated protein.

The overall goal in this work is to computationally explore tumor growth mechanism hypotheses in SCLC.

Tumor features that emerge as highly supported by data about the growth mechanism could be used to

predict differences in growth across tumors of different genetic backgrounds, responses to *in silico*

treatment, or even predict patient-specific tumor behavior after various treatments. Unfortunately, these

goals are currently hypothetical, because to build one SCLC model that could be used for these purposes,

one would need a unified understanding of the SCLC tumor as a system, and knowledge of SCLC currently

exists as nonoverlapping conclusions and hypotheses. We summarize the current knowledge about SCLC

tumor growth mechanisms, highlight potential knowledge gaps, and refer interested readers to (17) for a

comprehensive review of recent SCLC literature beyond that noted here.

Multiple SCLC subtypes have been identified depending on the experimental model studied, shown in **Fig**

1A, as SCLC-A, N, A2, and Y (Rudin et al., 2019). Other experiments have led to additional proposed

phenotypic subtypes, including canonical subtype SCLC-P, but these were not included in our analysis

(6,21–24). Our previous work aimed to identify whether all subtypes may be present in a tumor or if only

a subset are present, with the result that tumors can be composed of one, multiple, or all subtypes tested

(25). A comprehensive account of initiating SCLC subtype(s) (cell(s) of tumor origin) has not been made,

but multiple have been hypothesized in (25–27).

Studies *in vitro* and *in vivo* have suggested that Non-NE subtype(s) support growth of NE subtypes (20)

(**Fig 1A(2)**), including vasculogenic mimicking SCLC cells having such supportive effects (21). The

presence of NE subtypes has a dampening effect on Non-NE growth (28). Recent work has shown that the

Hes1-positive (Non-NE) cells supporting NE subtype growth (20) may have upregulated YAP1 (29) (**Fig**

1A(2)), and are likely SCLC-Y; otherwise, the referenced studies were completed before the adoption of

the canonical subtypes SCLC-A, N, A2, P, and Y, and so it is unclear which of these exactly contribute

such effects in each case.

NE cells may undergo a transition a toward Hes1+, (likely YAP1+) identity, (**Fig 1A(1)**) which modulates

these Non-NE cells' sensitivity to anticancer drug treatment (20). Other work found that SCLC-A subtype

cells can transition to the SCLC-N subtype and from SCLC-N to SCLC-Y (26) (**Fig 1A(5)**). Without the

ability to undergo a transition toward a more Non-NE phenotype, tumors were smaller and less aggressive;

however, this study did not assess Non-NE or SCLC-Y sensitivity to anticancer drugs (26). These two

landmark studies assessing phenotypic transitions do not assess the same phenotypic transition pathway

and thus we cannot compare intermediates, although we hypothesize that the transitions begin with SCLC-

A and it seems reasonable to assume the hierarchical pathway ends in SCLC-Y. While our investigations

Table 1. Existing data pertaining to SCLC intratumoral heterogeneity and communication.
Tumor composition   • Neuroendocrine-classified (NE) subtypes are A (ASCL1+) and N (NEUROD1+), with non-neuroendocrine-classified (Non-NE) subtype Y (YAP1+) and P (POU2F3+). (17)   • Previous work predicted subtype A2, and SCLC-A, SCLC-A2 and SCLC-N have been seen to express ASCL1 (17,20,25,26). SCLC-A2 expresses Hes1. (25)   • Hes1-positive TKO tumor cells (Non-NE) have YAP1 upregulated. (29)   • It is unclear whether A2 is more NE or Non-NE in character.   • Tumors can be made up of one or more of these subtypes. (17,25)   • Some subtype combinations have been experimentally verified and others have been predicted using CIBERSORT. (17,25,26)   • We do not see SCLC-P in our previous subtype deconvolution. (25)
Evidence of phenotypic transitions   • TKO tumor cells sorted for Hes1-negativity (NE identity) become Hes1 positive (Non-NE identity) when plated with Notch ligand DLL4. (20)   • *Ex vivo* culturing of *in situ* RPM tumors results in histologic and transcriptional phenotypic changes from NE to Non-NE gene expression over time. (26)   • Transitions between A and A2, A2 and N have not been studied.   • No evidence of SCLC-Y transition to NE identity. (20)
Subtype-to-subtype effects   • Cell viability and division are increased when Hes1-negative cells are plated with Hes1-positive cells, compared to Hes1-negative cells only. (20)   • NE cells suppress Non-NE cell division. (30)   • Application of conditioned media or of isolated exosomes results in a morphological change in TKO-derived NE cell line KP3.*
* Personal communication (Alissa Weaver, Vanderbilt University)   ASCL1, Achaete-scute homolog 1; NEUROD1, neurogenic differentiation factor 1; POU2F3, POU class 2 homeobox 3; YAP1, yes-associated protein; HES1, Hes Family BHLH Transcription Factor 1. TKO, $p53^{fl/fl};Rb^{fl/fl};p130^{fl/fl}$ tumors (20); RPM, $Rb1^{fl/fl};Trp53^{fl/fl}$;Lox-Stop-Lox[LSL]-*Myc*T58A tumors (28)

support that SCLC-Y acts as an end state for phenotypic transitions, (7) another study identified that Non-

NE subtypes may have stemlike potential, (6) which contrasts with Non-NE or SCLC-Y acting as the end

of the hierarchical pathway.

Multiple mechanistic hypotheses emerge from existing data

Considering the aspects of SCLC tumor growth observed in the previous section, it is clear that no one

model exists that could easily recapitulate all datasets. To address this challenge, we explored mechanistic

hypotheses in the realm of tumor initiation and composition, phenotypic transitions and their hierarchy, and

subtype-to-subtype effects (**Table 1**). To select which of these multiple hypotheses to include in a

mechanistic model of SCLC without additional findings would introduce bias into the modeling process.

Instead, we can address these questions computationally, by including or excluding these behaviors across

multiple mechanistic models and evaluating whether model behaviors recapitulate SCLC data; that is, we

turn to model selection (14,31).

For our tumor growth mechanism exploration, we interpret tumor topology, initiation, potential subtype

behaviors, etc., as features (model terms) in candidate models (**Table 1**). We define *model variables* as

representations of species in the model (e.g., "subtype A"), and *model terms* as qualitative actions in the

model (e.g., "subtype A cell division"), whose rates are denoted by *kinetic parameters* (e.g., "subtype A

division rate") (**Fig S1A-C**).

To fully account for all possible tumor subtype makeup and an exhaustive set of possible explanations, we

explored models comprising between two and four subtypes per model (**Fig 1B**) and included all possible

cell subtype interactions. Growth supportive effects and transition-inducing effects (**Fig 1C**) (and growth

dampening effects, not shown) are included in some candidate models where, e.g., presence of an effector

(supportive cell subtype) increases the rate of growth of a subtype it affects (supports). Subtype A2 has

expression features of both NE and Non-NE cells (25), including expression of ASCL1 (seen in NE cells)

and HES1 (seen in Non-NE cells) and we therefore assigned A2 NE features in some candidate models and

Non-NE in others (**Fig 1C**).

To compare a hierarchical system, where a cancer stem cell (CSC) can (re)populate a tumor, and non-

hierarchical systems in which phenotypic transitions can occur among multiple or all SCLC subtypes, we

include candidate models with several different potential phenotypic transition schemes. Thus, the set of

candidate models considered include models without phenotypic transitions, models with transitions that

reflect hierarchical transitions observed experimentally (20,26), and models with reversible transitions, i.e.,

high plasticity (**Fig 1D,E**). Unidirectional transitions stemming from one cell subtype indicate a potential

CSC, while bidirectional transitions from multiple subtypes indicate phenotypic plasticity. We additionally

include tumor initiation from one cell of origin *vs.* multiple. Thus, candidate models include different

numbers of initiating subtypes (**Fig 1D**, **Fig S2G**).

To ensure we built a comprehensive set of candidate models that enable exhaustive exploration of

biologically relevant hypothesis space, we combined the potential SCLC behaviors (**Fig 1, Table 1**) with

prior knowledge about mechanistic behavior of tumor populations (32–37). For example, if there is indeed

plasticity in the system, it is likely to be shared among subtypes, leading to symmetrical bidirectional

phenotypic transitions across the model (**Fig 1E**). We therefore expect that all plausible SCLC tumor

growth mechanisms are represented in our candidate model hypothesis space to the best of our knowledge.

Accounting for all these different possibilities led to a set of 5,891 unique candidate models, each

representing a possible SCLC tumor growth mechanistic hypothesis.

Bayesian exploration of candidate population dynamics models using experimental data

We use multiple datasets to identify consensus behavior of SCLC and provide a unifying model of tumor

growth mechanisms broadly supported by available data (**Fig 2A**). These datasets include two genetically-

engineered mouse models (GEMMs), the triple-knockout (TKO) model ($p53^{fl/fl}$;$Rb^{fl/fl}$;$p130^{fl/fl}$ tumors (20);

equivalent to the RPR2 GEMM (26,28,38)), and the RPM model ($Rb1^{fl/fl}$;$Trp53^{fl/fl}$;Lox-Stop-Lox[LSL]-

Myc[T58A] (28)), and cell lines from the Cancer Cell Line Encyclopedia (CCLE) (39) made up largely of the

SCLC-A subtype determined in (25) (**Fig 2A**; **File S1**). This data provided SCLC subtype-assigned

proportions of tumor samples, using the same gene signatures across samples, which had been automatically

determined by CIBERSORT from samples of CCLE SCLC cell lines (39) and their consensus clustering

class labels (25). We consider this preferable to our own *ad hoc* decisions of individual cell subtype identity

necessary to assign the required subtype proportions of tumors had we used newer, available single-cell

RNA sequencing data. The different datasets represented in **Fig 2A** demonstrate differing SCLC tumor

makeup, according to the experimental model employed in the study.

Fig 2. Population composition data and probabilistic representation. (A) CIBERSORT deconvolution of TKO

and RPM genetically engineered mouse model (GEMM) samples (previously published) as well as SCLC-A cell line

samples. CIBERSORT was performed on bulk RNA-sequencing data. (B) Probabilistic representation of tumor

proportion based on mean and standard deviation of proportions across samples within an experimental model; these

distributions were then used for fitting models to data. TKO, p53[fl/fl];Rb[fl/fl];p130[fl/fl] tumors [20]; RPM, Rb1[fl/fl];Trp53[fl/fl];Lox-

Stop-Lox[LSL]-Myc[T58A] tumors [28]; SCLC-A cell lines, a subset of SCLC cell lines from the CCLE [39] that we previously

assigned as representative of tumors made up largely of the SCLC-A subtype [25].

To explore the roles of phenotypic heterogeneity and cellular behaviors (cell-cell interactions, phenotypic

transitions) on SCLC tumor growth dynamics, we used population dynamics modeling, building on our

previous work (40). Population dynamics models employ a mathematical description of the dynamics

within and between heterogeneous subpopulations in an overall population (41,42). With such models,

researchers can mathematically simulate population growth over time and investigate growth dynamics

inherent in the simulations (**Box 1**).

With 5,891 candidate models, (**Fig 1B-E**) we aimed to determine which one(s) could best represent the

SCLC system – in an ideal world, we would determine which model represents reality. However, because

we will never know the "true" model, we must evaluate each model probabilistically – which model is most

likely to explain the data? Before comparing our 5,891 models to data to determine which is best, we

consider each to have the same probability of explaining the data. That is, we assign all models an equal

probability, as this way, we will not bias the exploration of the data toward any particular model or group

of models. This is the reason we built a set of models that enables full exploration of the hypothesis space,

(detailed above) so that with the addition of data, we can ensure we will study all possibilities and find the

best model(s) within that hypothesis space. We compare each candidate model to the data discussed above,

and with model optimization, we assess how well its simulations match what we see in that data. After this,

each model's probability will be updated, with a model that explains the data better achieving a higher

probability than one that does not explain the data as well.

The principles of model selection enable us to assess which candidates are best supported by experimental

data, while prioritizing model simplicity (14,43). Information theoretic approaches for model selection

using the Akaike Information Criterion (AIC) have been used in prior work but do not yield a much-needed

statistical understanding of the data. For a didactic demonstration of this, we refer the reader to **Note S1**.

We therefore employed the marginal likelihood as a more principled means for model ranking and model

averaging.

Box 1. Population dynamics modeling and inter-subtype effects

A population dynamics model represents behaviors of species over time, tracking the size of a cell population, rather than tracking individual cells. However, such a model can also include signaling and dependence between species (Harris et al., 2019). Cell population changes are represented as reactions, where subpopulation abundances increase or decreases due to varying events (division, death, phenotypic transitions), and the rates of increase or decrease can be affected by the presence of other subpopulations. Here, cell types x and w are able to undergo division, death and phenotypic transition at rates k_{div}, k_{die}, and $k_{x\text{-}w}$, respectively (A).

A

$$x \xrightarrow{k_{div}} 2x$$

$$x \xrightarrow{k_{die}} \emptyset$$

$$x \xrightarrow{k_{x \to w}} w$$

B

In our case, where species are cells of an SCLC tumor population, each is assigned a subtype identity (A, N, A2, or Y). The model simulates tumor growth over time, (B, top) calculating the increase in subpopulation amount (B, middle), from which tumor subtype proportion may be calculated (B, bottom) and compared at steady state to tumor proportion data. We define steady state as the composition of a tumor based on the relative abundance of each cell subtype in the tumor, without external perturbations. Though tumor growth may continue exponentially in a steady state, the proportion of subpopulations within remains constant (Harris et al., 2019).

Non-spatial cell-cell interactions may also be modeled, by changing the rates of reactions for one subpopulation according to the amount of another subpopulation present. Here, using the hypothesized biology where subtype w produces a secreted signal f that affects subtype x, we calculate the change in reaction rates. In (C), w secretes the unknown signaling factor f at a rate of k_f while f is degraded at rate k_{-f}. Factor f may affect subtype x at a rate of k_e, where x becomes x^* ("x under effect of f"). Subtype x can then revert back to its unaffected form at a rate of k_{-e}, indicating the rate at which the signaling is completed.

C

$$w \xrightarrow{k_f} w + f$$

$$f \xrightarrow{k_{-f}} \emptyset$$

$$x + f \xrightarrow{k_e} x^* + f$$

$$x^* \xrightarrow{k_{-e}} x$$

D

$$\frac{k_{-f}}{k_f} \equiv K_f^{eq} \qquad \frac{k_{-e}}{k_e} \equiv K_D$$

$$\nu_{fate} = \left(\frac{k_{fate} K_D K_f^{eq} + k_{fate}^*[w]}{K_D K_f^{eq} + [w]}\right)[x]_T$$

With k_f the production rate constant and k_{-f} the degradation rate constant, K_f^{eq} is the equilibrium constant for the amount of factor f in the system (D). Similarly, K_D is the equilibrium constant related to the on-effect rate constant k_e and the off-effect rate constant k_{-e}. Given these, the rate of a cell fate (division, death, or phenotypic transition) for x (ν_{fate}) can be calculated as a function of the population size of the effector cell w (D) (Harris et al., 2019) (see Note S1 for calculations).

By assigning the value of k_{fate}^* as more or less than k_{fate} for a particular cell fate, the presence of the effector cell subpopulation can increase or decrease, respectively, the rate of the cell fate for subpopulation x. In our population dynamics model, typically effector cells increase division and transition rates and decrease death rates, as shown in (E).

E

$$x \xrightarrow{k_{div}^*} 2x$$

$$x \xrightarrow{k_{die}^*} \emptyset$$

$$x \xrightarrow{k_{x \to w}^*} w$$

Prior work has estimated the marginal likelihood for kinetic model fitting using thermodynamic integration

(44). In this work we instead use nested sampling, (45), which is computationally more efficient and has

fewer limitations with regard to the shape of the probability space traversed during evidence calculation

(46–48). The nested sampling method was run once for each of the 5,891 candidate models on each of the

three experimental datasets, amounting to 17,484 potential interpretations of tumor growth mechanisms.

The average fitting time for each model was ~19 wall-clock hours, thus necessitating high-performance

computing for a complete parameter space exploration of the candidate models.

Each model is thus optimized to our datasets via nested sampling, which explores the full volume of the

likely parameter space. Each point in parameter space represents a set of possible parameter values (**Fig**

S3). At each of these points, nested sampling assigns a likelihood value for how well that set of parameter

values fits the data. On completion of the algorithm, the output includes the highest-likelihood parameter

values. Since each tested point in parameter space is a set of parameter values, the highest-likelihood values

for a model are returned by the algorithm as a list of parameter sets (**Fig S4**). Returning a list of parameter

sets rather than one top-scoring set already incorporates Bayesian methodology into the process – each

individual parameter has multiple best-fitting values, which can be interpreted as a distribution of parameter

values (44) – but with nested sampling we add yet more Bayesian methodology. Having assigned a

likelihood to every point in parameter space, nested sampling uses these to calculate one overall likelihood

per model, the marginal likelihood, which takes into account parameter fit as well as model simplicity

(number of parameters). For more detail on how the marginal likelihood is calculated to incorporate both

model fit and size, see **Note S1** and **Methods**. Finally, with marginal likelihood values for each model in

the candidate set, and the candidate model set representing the full hypothesis space with all potential SCLC

population dynamics models, we can calculate a probability. Summing the marginal likelihood values, and

dividing each individual marginal likelihood by these, results in a model posterior probability, representing

a change in probability from pre-model fitting (all models with equal prior probability) to post-model fitting

(see **Note S1**). We are then able to compare model probabilities and additionally perform model averaging

to evaluate kinetic parameter value distributions and probabilities of model variables and terms.

A small subset of candidate tumor growth models is supported by experimental data

In a Bayesian model selection approach, a more likely model comprises a higher proportion of the

probability of the candidate model space (**Fig 3A-C**). After nested sampling, our results indicate the highest-

scoring model for each dataset is $\sim 10^{19}$ times more likely than the lowest-scoring model, and $\sim 10^3$ times

more likely than the median scoring model. For reference, the smallest comparison between models that is

considered significant is $10^{1/2}$ (49) (**Note S1**). Therefore, our results indicate that the data used for model

fitting has informed our knowledge about the system, because before nested sampling, all models are

equally likely.

Performing nested sampling on all candidate models did not yield a unique best-fitting model for any dataset

(**Fig 3A-C**). We therefore leveraged a multi-model inference approach and calculated a confidence interval

(CI) representing a set of best-fitting models per dataset. While a 95% CI is a traditional approach, (**Fig**

3A-C, orange) we also calculate a "relative likelihood confidence interval," as discussed in (14) (see

Methods). For this relative likelihood CI, we calculate the Bayes Factor (BF) between the highest-scoring

model and every other model, using the least strict cutoff of $BF > 10^{1/2}$ (as above, and in **Note S1**). Even

with this permissive cutoff, the relative likelihood CI includes only tens of models per dataset, a large

decrease from the initial number of candidates (~1% or less, **Fig 3D**).

In summary, we can determine a subset of candidate models that adequately represent the data, conditional

on the fitted parameter sets resulting from the model optimization in nested sampling. Investigating these

parameter sets can provide more insight into the similarities and differences between candidate models and

their fits within and between datasets. Moving beyond the parameter values assigned to each model term,

we wanted to investigate how the data available can inform model terms. If data does not inform model

terms and variables and the corresponding fitted parameter rates, it indicates that the mechanistic

conclusions we desire to draw from this data using mathematical modeling may require additional or

different data.

Fig 3. Fitting to data and assigning Bayesian evidence separates candidate models into more and less likely.
(A)-(C). Evidence values (left y-axis) and posterior probability values (right y-axis) from nested sampling, one point
per model, ordered from model with greatest evidence to model with least evidence. Models whose evidence value
are within $10^{1/2}$ of the greatest evidence value, the "relative likelihood confidence interval," are colored in red. Nested
sampling and evidence calculation is performed per dataset. (A) TKO dataset. (B) RPM dataset. (C) SCLC-A cell line
dataset. (D) Numbers and percentages of models in the relative likelihood confidence interval, 95% confidence
interval, and remaining non-confidence interval models. TKO, $p53^{fl/fl};Rb^{fl/fl};p130^{fl/fl}$ tumors [20]; RPM, $Rb1^{fl/fl};Trp53^{fl/fl}$;Lox-
Stop-Lox[LSL]-Myc^{T58A} tumors [28]; SCLC-A cell lines, a subset of SCLC cell lines from the CCLE [39] that we previously
assigned as representative of tumors made up largely of the SCLC-A subtype [25].

High-likelihood model topologies are nonoverlapping between datasets

Given our observation that no one candidate model stands out among other models to explain the experimental data, we employed the multi-model inference technique of Bayesian model averaging (BMA). Briefly, the reasoning behind BMA is that a combination of candidate models will perform better in explaining the data than a single model (15). In BMA, each model is weighted by its posterior model evidence (31) and the model terms within each model receive an averaged likelihood (50) (**Note S1-S3**).

However, before we could investigate model-averaged probabilities, we found that the fitted parameter distribution outcome was dependent on choice of initiating subtype (**Fig S5**). While we aimed to evaluate initiating subtype possibilities with our Bayes-MMI approach, we cannot do this in a unifying fashion as this choice affects fitted parameters and thus model behavior. Therefore, we turned to the literature to impose stricter constraints about initial subtype conditions, to investigate model variable and term probabilities when kinetic parameter posterior distributions were constrained. As mentioned previously, reports link NE SCLC subtypes and long-term tumor propagation (20,30) and, in particular, cells of subtype A (26). We thus used only candidate models with an initiating subtype of A, with or without other initiating subtypes. Since we required that subtype A be an initiating subtype, model structures that do not include subtype A received zero posterior probability (models 3 and 8–10 in **Fig 4A**; model topology probabilities without filtering by initiating subtype are shown in **Fig S6A**).

Fig 4. Likely model topologies vary across datasets; transition rates vary according to subtype presence in similar ways. (A) Hypothesis assessment of model topologies, per dataset. Probability indicates the result of Bayes theorem using equivalent prior probabilities per topology (e.g., 9% probability that one of the topologies in the x-axis best represents a dataset) and evidence values (marginal likelihoods) summed per topology. Model topologies represented by images and corresponding numbers along the x-axis. Posterior probability based on marginal likelihoods of all candidate models that include A as an initiating subtype. (B) Division and phenotypic transition parameters for TKO, RPM, and SCLC-A cell line datasets, comparing between higher-probability topologies (A) and four-subtype topology per dataset. TKO, $p53^{fl/fl}$;$Rb^{fl/fl}$;$p130^{fl/fl}$ tumors [20]; RPM, $Rb1^{fl/fl}$;$Trp53^{fl/fl}$;Lox-Stop-Lox[LSL]-Myc^{T58A} tumors [28]; SCLC-A cell lines, a subset of SCLC cell lines from the CCLE [39] that we previously assigned as representative of tumors made up largely of the SCLC-A subtype [25]. (*) indicates significance between samples from BMA parameter distributions at family-wise error rate (FWER) = 0.01, averaged over ten sampling iterations using one-way ANOVA plus Tukey HSD.

We perform BMA across all models for each dataset. As shown in **Fig 4A**, all datasets (TKO, RPM, and

SCLC-A cell lines) support both two- and three-subtype topologies. Higher probabilities for two-subtype

topologies are expected given that nested sampling prioritizes model simplicity and goodness of fit (45).

Statistically, this result suggests that a two-subtype model could be used to interpret the data reasonably

well, but it also shows that topologies comprising three subtypes cannot be excluded. Approximately 10%

of the probability for the GEMM datasets (TKO and RPM) fall in the three-subtype topology that

encompasses the high-probability two-subtype topologies (model 1 for RPM and model 2 for TKO in **Fig**

4A). For the models fit to SCLC-A cell line data, most of the probability occurs in the topologies with

higher probabilities for the GEMM data. This is reasonable, given that SCLC-A cell line data appears as an

intermediate between the GEMMs (**Fig 2A**). However, the SCLC-A cell line data also has probability that

falls in the A, N, and A2 topology (model 4 in **Fig 4A**) – this is the only topology at all likely to represent

the SCLC-A cell line data but not at all likely to represent the other two datasets.

We interpret the spread of probabilities across multiple topologies, and that most topologies either are

probable as representing either TKO or RPM data but not both, to mean that data coverage from these

datasets is not sufficient to support one unifying topology. Therefore, each dataset supports a different

representation of SCLC tumor growth given its particular (epi)genetic background and environment. This

does not mean that a unifying topology or unifying model of SCLC growth cannot exist, but that the biases

underlying the experimental data result in different explanations for tumor growth mechanisms.

All datasets support alteration of phenotypic transition rates in the presence of N or A2 subtypes

After establishing that multiple model topologies can explain tumor growth mechanisms, we explored

whether the rates of distinct cellular subtype behaviors were characteristic for given model topologies

within each dataset. We therefore compared kinetic parameters across models to learn about dynamic

variation between model topologies. We again used BMA to attain this goal, applying the approach to fitted

kinetic parameter distributions from nested sampling. In this setting, parameter values from more likely

models are assigned higher weights and corresponding parameter distributions are weighted accordingly.

The highest likelihood model topologies (**Fig 4A**, blue) for the TKO GEMM data, along with the four-

subtype topology, are compared in **Fig 4B** (left). Three model terms have significantly different parameter

rates across model topologies, all of which are discussed in **Note S4**. Here we highlight differences in the

A-to-Y and A-to-A2 transitions across model topologies in the TKO dataset: the A-to-Y transition has a

slower rate if A2 is present in the population, and only Y affects the A-to-A2 transition, increasing its rate.

The mechanistic implication of these observations are as such: A2 may represent an intermediate

subpopulation in the tumor that is longer-lived, and will only slowly transition to Y. In the topology with

A, A2, and Y (**Fig 4A**, structure 2; **Fig 4B** left, third-darkest blue), the A-to-A2 transition takes up more of

the flux in the network. Additionally, the N-to-Y transition is faster relative to the A2-to-Y transition (**Fig**

S6B), suggesting that N is a shorter-lived intermediate in the A-to-N-to-Y transition. This result aligns with

previous experiments (Ireland et al., 2020) where N was identified as a short-lived state in the A-to-N-to-

Y transition. We therefore predict that A2 and N are involved in regulating the relative abundance of, and

flux between, A and Y in the tumor.

We also compared the highest likelihood model topologies (**Fig 4A**, red) for RPM-fitted models, as well as

the four-subtype topology (**Fig 4B**, middle). Five parameter rates are significantly different across model

topologies, and we highlight again the A-to-Y transition and here the A-to-N transition (see **Note S4** for

discussion of the remaining significantly different parameter rates). The same A-to-Y transition affected in

the TKO-fitted models is affected in the RPM model in the same way (reduced rate via an intermediate, in

this case N), and the A-to-N transition is affected by Y similarly to the A-to-A2 transition affected by Y in

the TKO data. These similar effects occur despite the experimental data used for BMA being different. We

thus predict that N and A2 are modulating the transition between, and relative abundance of, A and Y.

Unlike in the TKO data, when A2 is present the flux through the system spends more time in the N subtype,

with more frequent transitions to N and less frequent transitions to Y. We predict that while N may be a

shorter-lived intermediate than A2, A2 regulates the flux from A-to-N-to-Y.

Next, we compared the highest likelihood model topologies (**Fig 4A**, green) for the SCLC-A cell line data

and the four-subtype topology (**Fig 4B**, right). Six parameter rates are significantly different across model

topologies, five of which recapitulate rate alterations based on the presence or absence of different subtypes

in TKO or RPM datasets, including the rate alterations discussed above (see **Note S4** for more detail).

In summary, BMA enabled us to determine that the A-to-Y transition is regulated in a similar manner for

the RPM, TKO, and SCLC-A datasets and that the A-to-N and A-to-A2 transitions are regulated similarly

in each dataset. Using the higher likelihood model topologies and model-averaged parameter sets, we can

infer features of the SCLC tumor generally, despite disparate datasets. Finding the same or similar effects

on kinetic parameter rates across independent datasets lends more weight to these predictions about how

the N and A2 subtypes may regulate the system flux from A to Y through intermediates and is an advantage

of our methodology using Bayes-MMI to work toward a unifying model of SCLC tumor growth based on

multiple datasets.

Model analysis supports a non-hierarchical differentiation scheme among SCLC subtypes

We have considered candidate models (**Fig 3**), model topologies (**Fig 4A**), and kinetic parameters (**Fig 4B**)

to explore tumor growth mechanisms in SCLC. There is compelling experimental evidence for multi-

subtype tumor composition, which implies multiple potential growth mechanisms (6,20,26,28,51). We

therefore focused on model topologies 1, 2, and 4, which are three-subtype topologies with detectable

probability ($> 1\%$) (**Fig 4A**), along with the four-subtype model. Using these, we integrate candidate

models, topologies, and kinetic parameters, investigating phenotypic transitions between subtypes, whether

the presence of certain subtypes affects the behaviors of other subtypes, and if so, which subtypes bring

about the effects (**Table 2**). We conclude by proposing a unifying four-subtype model of tumor growth in

SCLC, aiming to represent with one model the varying growth mechanisms accessible across datasets.

Table 2. Probabilities after hypothesis exploration using Bayesian multimodel inference

Informed high posterior probabilities

- Simulated tumors appeared more likely to be made up of fewer than four subtypes, indicating the model selection algorithm's preference for parsimony (fewer subtypes to explain the same data).
- Phenotypic transitions A-to-N, N-to-Y, A-to-Y had posterior probabilities between 60% and 71%
- Posterior odds for these are between 1.5 and 2.5.
- Transitions are bidirectional: phenotypic transitions N-to-A, A2-to-A, Y-to-N, Y-to-A2, and Y-to-A, had posterior probabilities between 61% and 81%
- Posterior odds: ~1.5 to 4.
- Transitions between N and A2 (N-to-A2, A2-to-N) had posterior probability 66%
- Posterior odds: ~2

Informed low posterior probabilities

- Low probability of effects that lead to more/quicker phenotypic transitions from NE to Non-NE subtypes, posterior probabilities between 17% and 43% (average 32%)
- Posterior odds: ~0.5
- In SCLC-A cell line datasets, trophic effects, where Non-NE subtypes increase NE division and decrease NE death, had posterior probability 16%
- Posterior odds: 0.19

Uninformed posterior probabilities

- Initiating / early post-initiation number of subtypes: out of 15 model initiation options, each probability was between 1% and 19%
- Phenotypic transitions A-to-A2 and A2-to-Y had posterior probabilities between 52% and 62%
- Posterior odds for these are between 1.08 and 1.6.
- In TKO and RPM datasets, trophic effects, where Non-NE subtypes increase NE division and decrease NE death, had posterior probabilities between 45% and 46%
- Posterior odds: 0.79 to 0.85.
- In TKO three-subtype models (only three-subtype model with both types of effects) A2 and Y effects are 53% probable *vs.* Y only effects at 47% probable
- Posterior odds for A2 and Y effects is 1.13, posterior odds for Y only effects is 0.89.

We investigate the posterior probabilities, and therefore posterior odds, of each model term (see **Methods**).

Despite different posterior probability values (**Fig 5A**), the probabilities of model terms across datasets

were similar in their trends: across all three-subtype topologies, phenotypic transition probabilities were all

more than ½ (**Fig 5A**, red squares). While some probability values were poorly informed (light red),

(probability between ½ and ⅔), more were informed by the data (deep red) (⅔ or more). Conversely,

probabilities of Non-NE effects on the growth or transitions were all less than ½ (**Fig 5A**, blue squares).

Some probability values were poorly informed, (light blue) (between ⅓ and ½) and others were informed

(deep blue) (⅓ or less) with the addition of data.

Overall, the data suggests that Non-NE effects on transition rates of N-to-Y, or A2-to-Y, are unlikely, (**Fig**

5B-D, deep blue) regardless of whether "Non-NE" defines only the Y subtype, or both A2 and Y are Non-

NE (**Fig 1C**). Inter-subtype effects on SCLC phenotypic transition rates have not previously been studied

and our analysis predicts that at least effects on "late transition" (**Fig 5A**), those interactions affecting N-

to-Y or A2-to-Y, are unlikely to exist. By contrast, transitions involving A-to-N, N-to-A, A2-to-A, N-to-Y,

Y-to-N, A2-to-Y, Y-to-A2, N-to-A2, A2-to-N, A-to-Y, and Y-to-A had posterior probabilities informed by

the data (**Fig 5B-D**, deeper red). We interpret these results as transitions being likely, i.e., our degree of

belief in these transitions has increased. Investigating initiating events via one or multiple cells of origin

across the candidate models, we find that from equal prior probabilities of 6.67% per initiating subtype(s)

(**Fig S2G**) the posterior probabilities are not significantly altered, being between 1% and 15% (not shown).

Thus, initiating subtype events were poorly informed by the data. Additionally, analyzing specific model

terms, inter-subtype effects on NE subtype growth, inter-subtype effects on transition rates between A and

N, or A and A2, and the A-to-A2 transition, were also poorly informed by the data (**Fig 5B-D**, light blue,

light red).

Fig 5. Across datasets, multimodel inference indicates likely bidirectional phentoypic transitions, suggesting high SCLC phenotypic plasticity. (A) Heatmap for high probability three-subtype topologies for each dataset (rows), all models initiated by A +/- other subtypes. Color represents the probability of each cellular behavior (column). Since prior probability starts at 0.5 (white), deeper colors indicate a larger deviation from the prior, with red vs blue indicating more likely or less likely, respectively. (B)-(D). Model schematics with each cellular behavior represented by edges coming from or moving toward each cell subtype, (gray circles) growth rates, (self-arrows) or transitions (arrows between gray circles). Edge colors correspond to colors for that behavior in the heatmap in (A). Top-scoring three-state topology for TKO dataset (B), RPM dataset (C), and SCLC-A cell line dataset (D). (E) Schematic of consolidated model behaviors, drawn from each dataset's high-probability three-subtype topology results ((B)-(D)). When multiple dataset results included different posterior probabilities for a model feature, the one closest to 0.5 was chosen (most conservative). Edge colors correspond to posterior probabilities, with intensity of colors representing information gained from data, as in (A)-(E). (F) Parameter fitting results (part of the nested sampling algorithm) for four-subtype topology models initiated by A +/- other subtypes, across datasets. tsn, transition (e.g., subtype transition). TKO, $p53^{fl/fl}$;$Rb1^{fl/fl}$;$p130^{fl/fl}$ tumors (20); RPM, $Rb1^{fl/fl}$;$Trp53^{fl/fl}$;Lox-Stop-Lox[LSL]-Myc^{T58A} tumors (28); SCLC-A cell lines, a subset of SCLC cell lines from the CCLE (39) that we previously assigned as representative of tumors made up largely of the SCLC-A subtype (25). (*) indicates significance between samples from BMA parameter distributions at family-wise error rate (FWER) = 0.01, averaged over ten sampling iterations using one-way ANOVA plus Tukey HSD.

Finally, to consolidate phenotypic transitions and cell-cell interactions into a unifying mechanism for SCLC

tumor growth, we integrated model probabilities from each of the three-subtype topologies for each dataset

into one model (**Fig 5E**). Briefly, phenotypic transition probabilities were chosen from the models least

informed by the data in an attempt to make conservative predictions (see **Methods**). Model-averaged

parameter rates were visually compared (**Fig 5F**) to ensure that they were within reasonable bounds and

that transition rates relate to each other between datasets similarly to our analyses using high-probability

topologies (**Fig 4B**). Values from the consolidated probabilities (**Fig 5E**) are those reported in **Table 2**,

along with posterior odds to compare one hypothesis to its opposite.

Taken together, these results provide insight not only into what model terms and variables the data is able

to inform, but SCLC tumor behavior as well. Knowledge of trophic effects provided by Non-NE cells to

the benefit of NE cells was not provided by this particular data; therefore, we cannot use it to understand

this behavior. However, we were able to gain knowledge about the likelihood of phenotypic transitions, in

fact indicating that nearly all options for phenotypic transitions are likely to exist. We interpret this as high

SCLC plasticity, supporting a non-hierarchical differentiation scheme where tumor population equilibrium

is achieved through any phenotypic transitions (**Fig 5E**). It is also clear that consolidating the results across

different tumor types is an important step in order to achieve a broader view of the SCLC tumor as a system

rather than as one particular experimental model.

DISCUSSION

The experimental data used for this analysis favors two-subtype topologies as higher-probability candidates.

This is not surprising, because nested sampling prioritizes simpler models. Despite this finding, mounting

experimental evidence supports multi-subtype tumor composition and these data have been interpreted in

the context of multiple existing phenotypes (6,20,26,28,51). In fact, previous work from our labs suggests

that the tumor genetic background dictates possible phenotypic subtypes within a tumor population, and

that phenotypic transitions likely mediate the ability of tumor cells to achieve these phenotypes (40,52,53).

We hypothesize that with additional timecourse and perturbation data, the topology likelihood for tumor

growth mechanism could be shifted toward three- or four-subtype models. This is consistent with studies

of tumor dynamics from cancer broadly, both with regard to phenotypic transitions toward or away from a

rarer subpopulation (34,54–56) and changes in the proportion of phenotypic subpopulations after a

perturbation. Often this perturbation is the application of drug treatment (57–62), but may also be changes

in microenvironment or related factors (26,63).

The results presented here also provide strong evidence for phenotypic plasticity in SCLC tumors, based

on the higher likelihood for most phenotypic transitions tested, regardless of differentiation hierarchy. With

a more plastic and less stem-cell based phenotypic equilibrium, instead of rare remaining stem-like

populations leading to regenerate a tumor after treatment, we hypothesize that any SCLC subtype that

remains post-treatment can lead to tumor regeneration and subsequent treatment resistance, patient

morbidity and mortality. While hierarchical phenotypic heterogeneity *vs.* phenotypic plasticity has not been

experimentally studied in SCLC, we predict that plasticity is highly likely, and extremely important in the

growth and evolution of the SCLC tumor. It is of particular interest to compare phenotypic plasticity and

the prevalence of non-hierarchical transitions in treated *vs.* untreated tumor samples, as treatment is likely

to alter the mechanisms by which tumor population equilibrium is maintained. Time-course experiments

with surface marker labeling or live-reporter imaging can resolve and provide confirmation for bidirectional

phenotypic transitions, which are crucial to understand in order to battle SCLC treatment resistance.

The invasive or metastatic potential of the SCLC tumor is known to be increased by Non-NE subtypes

(21,64). It is unclear whether the conclusions and predictions presented here apply to SCLC in the invasive

or metastatic setting, but future work will include model additions to place the tumor growth in a

physiologic context that includes both the tumor *in situ* and during invasion.

We believe a shift from information theoretic multimodel inference toward a Bayesian approach, enabling

investigation of optimal model(s) with identifiable parameters for a particular dataset, will benefit modeling

in systems biology. The methodology employed herein incorporates model selection and model averaging

into a multimodel inference framework, followed by Bayesian analysis to identify not only whether a

hypothesis investigated via mechanistic modeling is or is not likely, but *how* likely (and thus how informed

by the data) that hypothesis is. Understanding which hypotheses are informed by the data is especially

important given variability between data in investigations of the same systems, such as a particular tumor

type. It is difficult to attain a consensus model since investigators use varying experimental models within

the same physiologic or disease process and thus may draw nonoverlapping conclusions, building parts of

a picture but not a whole. Striving for the whole picture, via principled statistical analysis, to be followed

by experiments based on informed model predictions, will advance cancer research and lead to better

treatments.

METHODS

CIBERSORT deconvolution of RNA sequencing data

Data from two GEMM models provide multiple replicates of tumors from two genetic backgrounds: one from $p53^{fl/fl};Rb^{fl/fl};p130^{fl/fl}$ (triple-knockout, or TKO) GEMM tumors (20), and another from $Rb1^{fl/fl};Trp53^{fl/fl}$;Lox-Stop-Lox[LSL]-$Myc^{T58A}$ (RPM) GEMM tumors (28). We also used publicly available SCLC cell line data. Having been originally derived from human tumors, each cell line has a different genetic background, and therefore we have only one (genetically identical) replicate per cell line sequencing event. To approximate genetic similarity between cell lines, and thus approximate multiple replicates, we expect that cell lines exhibiting similar steady state composition will be more genetically similar than those whose steady state compositions differ. Previously, in (25), we both clustered publicly available SCLC cell line data into clusters that align with the different SCLC subtypes and used CIBERSORT to deconvolute the proportions of cell line data and tumor samples into SCLC subtypes from their RNA sequencing signatures. Results used for this publication can be found as **File S1**.

Population dynamics modeling in PySB

A population dynamics model represents the abundance of species over time, whether increase or decrease due to birth/growth or death. We use ordinary differential equation (ODE) models coded via PySB to generate population dynamics models (65). PySB is a rule-based modeling language, where one will encode

$$A \xrightarrow{k} 2A, k = 0.469 \tag{1}$$

to indicate that A doubles at a rate of 0.469 doublings per day.

Inter-subtype effects are represented by the increase or decrease of the rate of affected reaction. For example, the above division rule has a baseline rate of 0.469 doublings per day, but in the presence of an effector subtype the division rule will have a rate of 0.469*1.05 = 0.493 doublings per day. In this case the effector subtype has increased the division rate by 5%. Thus the rule-based representation is

$$A + Y \xrightarrow{k} 2A + Y, k = 0.493 \tag{2}$$

to indicate that A doubles at a rate of 0.493 doublings per day in the presence of Y.

To simulate the passage of time, the speed at which the division/death/transition reaction occurs–its rate, k,

of cells per unit time–must be assigned as in the equations above. While a literature search reveals

approximate rates of division and death among different SCLC subtypes (**Table S5**), each of these are in a

different context than the system we model here – for example, division rates for the A subtype are

measured *in vitro* in the presence of only that one subtype, whereas our population dynamics model is

meant to simulate this subtype in the presence of others as well as *in vivo* in a mouse tumor. Therefore, we

use the rates in the literature as our prior expectations for division and death, that is, we use these values as

approximate starting values for these parameters during the estimation process. Other rates, such as those

indicating the speed of transitions between subtypes, or any rates including the effects of Non-NE subtypes,

have not previously been noted in the literature and we used much wider ranges for each as our prior

expectations. Rate prior expectations (**Fig S3**) are then provided to the Multinest algorithm to perform

nested sampling.

Multiple hypothesis generation via HypBuilder

Because we perform model selection, we use 5,891 ODE models coded via rule-based modeling in PySB.

Each model is generated to include or exclude from 44 reaction rules. There are eight rules that represent

division and death for each subtype, and with the potential for three different inter-subtype effects

(including none) to have an impact on division or death, each division and death reaction has 3 options,

leading to 24 potential rules relating to division/death in total. There are four rules that represent hierarchical

phenotypic transitions, which likewise have three potential inter-subtype effects, for 12 rules in total

representing hierarchical phenotypic transitions. There are eight rules related to non-hierarchical

phenotypic transitions, for 20 total potential phenotypic transition rules out of the 44 rule options.

We use HypBuilder (https://github.com/LoLab-VU/HypBuilder) to automatically generate the 5,891 PySB

models that we would otherwise have to code by hand. HypBuilder is software for the automatic generation

user-specified collections of mechanistic rule-based models in the PySB format. The input CSV file

contains a global list of all possible model components, and reactions, as well as any instructions regarding

model creation. The instructions dictate which subsets of model components and reactions will be

combinatorially enumerated to create the collection of models. The reactions are parsed via HypBuilder's

molecular interaction library, a library of defined reaction rule sets that is outfitted with common PySB

interactions and is customizable to include more interactions should the user need them. Once parsed and

enumerated each combination of rules is exported as an executable model via PySB.

The instructions for model construction used in this work direct HypBuilder to use a "list" method to

enumerate all candidate models of interest using prior knowledge of likely combinations of model variables

(see https://github.com/LoLab-MSM/Bayes-MMI for code used to enumerate candidate models and create

the list for HypBuilder).

If the candidate model set contains every relevant biologically plausible possibility, we can consider the

entire set of models as representative of 100% of the probability that one of the candidate models explains,

or provides the mathematical basis underlying, the data. This is an assumption that cannot truly be met, and

most model selection literature acknowledges that one cannot find the "true" model (14,43). However, prior

knowledge enables us to determine that all 5,891 models represent all possibilities with regard to

outstanding SCLC hypotheses to the best of our ability.

We visualize the prior expectations for the 44 rate parameters as a probabilistic distribution per parameter

(prior marginal distribution) (**Fig S3**). Correspondingly, a probabilistic representation of best-fitting rates

for each model is returned by the Multinest algorithm (posterior marginal distribution) (e.g., **Fig 4B, 5E**;

Fig S5, S6B).

Parameter estimation and evidence calculation by nested sampling

581   As noted in Eqs. (1) and (2), rate parameters must be set in order to run simulations of a mathematical

582   model. Parameter estimation is the process of determining optimal rates that result in a model simulation

583   recapitulating the data it is meant to represent. Multiple methods exist for parameter fitting or model

584   optimization, (66,67) with Bayesian methods utilizing a prior rate parameter distribution, $P(\boldsymbol{\theta})$, where $\boldsymbol{\theta}$

585   represents the set of n parameters $\{\theta_1, \theta_2, ..., \theta_n\}$, and a likelihood function to assess a parameter set

$$L(D|\boldsymbol{\theta_i}) = P(D|\boldsymbol{\theta_i}), \tag{3}$$

Where θ_i is the ith parameter set and D represents the data being used for fitting. Parameter set θ_i is scored

via the likelihood function $L(D/\theta_i)$ and optimization continues, moving toward better-scoring parameter

sets until an optimal score is reached.

For our likelihood function, we represent SCLC tumor steady-state proportion probabilistically, generating

a Beta distribution (bounded by 0 and 1) to represent the means and standard deviations of sample replicate

subtype proportions, accounting for noise in the proportional space. We test n-1 subtype proportions to

ensure independence of each sampled subtype proportion, to result in the probability of n-1 independent

events (tumor subtype proportions) occurring.

With a prior probability, $P(\boldsymbol{\theta})$, and a likelihood (Eq. 3) the posterior probability can be calculated via Bayes'

Theorem,

$$P(\boldsymbol{\theta_i}|D) = \frac{P(D|\boldsymbol{\theta_i})P(\boldsymbol{\theta_i})}{\int P(D|\boldsymbol{\theta})P(\boldsymbol{\theta})d\boldsymbol{\theta}}. \tag{4}$$

The denominator of Bayes' Theorem represents the likelihood integrated over all parameter sets, called the

marginal likelihood or model evidence. Nested sampling computes this value (Skilling, 2004).

To perform nested sampling, we utilize the Multinest algorithm (46–48). Multinest samples multi-

dimensional parameter space, bounding its search by parameter values along each axis in each of the

multiple dimensions based on prior expectation of parameters, $P(\theta)$ input by the user. It removes the lowest-

probability parameter set and chooses a new one from within the bounded parameter space, subsequently

re-drawing the search space with the bounds incorporating the new parameter set. This continues until all

parameter sets representing the bounds of the search space have approximately equal probability, and the

algorithm estimates that the remaining probability of parameter sets within the bounds is less than a user-

defined tolerance. Each parameter set is evaluated based on a user-defined likelihood function (Eq. 3).

Finally, the likelihood values that correspond to each sampled parameter set are arranged in the order they

were replaced, and the integral over these is taken to approximate the integral over all possible models, that

is, the marginal likelihood or Bayesian model evidence. We used the Multinest-returned importance nested

sampling evidence value, because multiple importance nested sampling runs (multiple 'replicates') for the

same candidate model and prior parameters returned more consistent evidence values than 'vanilla' nested

sampling (data not shown) (48).

For our likelihood function we represented simulation outcomes – proportions of subtypes at steady state –

by a Beta distribution, calculating α and β using the mean μ and variance σ^2 of each dataset (68):

616
$$\nu = \alpha + \beta = \frac{\mu(1-\mu)}{\sigma^2} - 1, where\ \nu = (\alpha + \beta) > 0\ \left(meaning\ \sigma^2 < \mu(1-\mu)\right) \tag{5}$$

617
$$\alpha = \mu\nu = \mu\left(\frac{\mu(1-\mu)}{\sigma^2} - 1\right), if\ \sigma^2 < \mu(1-\mu) \tag{6}$$

618
$$\beta = (1-\mu)\nu = (1-\mu)\left(\frac{\mu(1-\mu)}{\sigma^2} - 1\right), if\ \sigma^2 < \mu(1-\mu). \tag{7}$$

619   We calculated the log-likelihood of each subtype mean from the dataset being fit against the simulated

620   subtype's beta distribution,

621
$$\sum_{s}^{S-1} LL$$

622   where

623
$$LL = \begin{cases} log\left(\frac{x_{s_{data}}^{\alpha_{s_{sim}}-1}\left(1-x_{s_{data}}\right)^{\beta_{s_{sim}}-1}}{B(\alpha,\beta)}\right) if\ s \in\ simulated\ model\ topology \\ log\left(\lambda e^{-\lambda x_{s_{data}}}\right), where\ \lambda = \frac{1}{\sigma_{s_{data}}},\ otherwise \end{cases} . \tag{8}$$

S is the set of subtypes, $\alpha_{s_{sim}}$ and $\beta_{s_{sim}}$ indicate the calculation of α (Eq. 6) and β (Eq. 7) using the

proportion of subtype s from the simulation and the variance of subtype s in the data, $x_{s_{data}}$ is the mean

proportion of subtype s in the dataset, $B(\alpha,\beta)$ is $\frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$ and Γ is the Gamma function, and $\sigma_{s_{data}}$ is the

standard deviation of the data. Using the exponential function $(log(\lambda e^{-\lambda x_{s_{data}}}))$ as part of the log-

likelihood (Eq. 8) enabled us to calculate a likelihood value for subtypes not present in a model's topology,

which should be a poor log-likelihood if the subtype has a high proportion in the data but was not included

in the model topology, or a better log-likelihood if the subtype has a low proportion in the data but was not

included in the model topology (and therefore potentially contributing to overfitting). The Python module

scipy.stats was used to calculate the Beta log likelihood (Eq. 8, above) and the exponential log likelihood

(Eq. 8, below). A simulation would not be scored (return *NaN* and thus be thrown out by the Multinest

fitting algorithm) if the tumor subtype proportions did not reach steady state (calculated by whether a

proportion timecourse had a slope of zero for the last 7.5% of the simulation).

Multinest is run per model per dataset, which equates to performing 5,891 mechanistic interpretations, 3

times each. CPU time for one model fitting was on average 19 hours (~0.80 days), with a range of 5 minutes

to 28 days. If Multinest had not reached its stopping point by 28 days, we assumed that all regions of

parameter space were similarly unlikely and that further running of the algorithm would only continue to

refine the search of the unlikely space; models with this difficulty are very likely to have low marginal

likelihood due to the unlikeliness of the parameter space. We do not include these incompletely-searched

models in our multimodel inference analyses (**Figs 3-5**) and we confirmed that all models that reached 28

days of CPU time without reaching the Multinest stopping point have an extremely low evidence value at

the time they were terminated.

Candidate model prior and posterior probabilities and confidence interval calculation

Each candidate model is considered equally likely prior to fitting by Multinest. That is, every candidate

model has an equal prior probability of being the optimal model to represent the underlying SCLC tumor

system,

$$P(M_k) = \frac{1}{|M|},$$ (9)

Where *M* is the set of all candidate models. With the model evidence, or marginal likelihood, *P(M$_k$)*

estimated by Multinest, (46–48) the posterior probability per model can be calculated as

$$P(M_k|D) = \frac{P(D|M_k)P(M_k)}{\sum_j P(D|M_j)P(M_j)}. \tag{10}$$

With a posterior probability per model, (**Fig 3**) we calculate a 95% confidence interval. This is

accomplished by summing decreasing model posterior probabilities until the sum is 0.95, then considering

those models as our 95% CI (14) (**Fig 3**, orange). Using this confidence interval results in ~1000 models

per dataset, a considerable decrease from the initial 5,891. This is a more traditional approach to

determining a confidence set of models.

We also took an approach discussed in (14). In this approach, a CI is informed by use of the Bayes Factor

between the highest-scoring model and consecutively decreasing scoring models, until the Bayes Factor is

larger than a particular cutoff. The models in this CI would be those models *i* for which $\frac{P(M_{highest})}{P(M_i)} >$

$cutoff$. Burnham and Anderson denote such a method as a "relative likelihood confidence interval" and

discuss its support by statistical theory, noting that it is uncommonly found in the model selection literature.

We used a cutoff of $10^{1/2}$, the lowest Bayes Factor at which a difference may be determined (49). Even with

this permissive cutoff, the relative likelihood CI includes only tens of models, an even greater decrease

from the initial number of candidates.

Prior and posterior probabilities per hypothesis being investigated

Each hypothesis has an assigned prior probability based on our prior expectations. For all hypotheses, we

took an approach where we considered each hypothesis as equally likely compared to competing

hypotheses. For the inclusion of most model terms, this was a prior probability of 50%, where it is 50%

likely the model term is part of a model that is the best representation of the tumor system, and 50% likely

that same term is not part of that model. For the inclusion of effects in the candidate models, the prior

probability for a given effect is 33%, where it is equally likely that an effect is generated by Y, generated

by A2 and Y, or that no effect is present. The comparison between effect types (including none) is included

in **Tables S6-S8**, while the comparison of any effect at all *vs.* no effect (50% *vs.* 50%) is included in the

main text.

For the model topology analysis, we considered it equally likely that any model topology could best

represent the tumor system that generated each dataset, and with 11 possible model topologies this resulted

in a 9% prior probability per model topology (**Fig 4A**). For model initiating subtype hypotheses, (**Fig S2G**)

with 15 potential combinations of initiating subtypes, each initiating subtype combination has a 6.67% prior

probability.

Each candidate model can then be assigned a prior probability conditional on the hypothesis being

considered, $P(M_k|H_i)$, where M_k is the k^{th} candidate model and H_i is the hypothesis being considered. The

calculation of $P(M_k|H_i)$ is based on the number of candidate models that fall under the hypothesis being

considered,

685
$$P(M_k|H_i) = \frac{P(H_i)}{|\{M_j \in M, H_i\}|} ,$$
(11)

686 where $\{M_j \subset M, H_i\}$ is the set of all models assigned to H_i. For example, if H_i is the hypothesis that the

687 model term "A to Y transition" is part of the model that would best represent the SCLC tumor system, then

688 $\{M_j \subset M, H_i\}$ is the set of all candidate models that include the "A to Y transition" model term.

689 Using this prior probability, the posterior probability for an individual model, conditional on the hypothesis

690 being considered, can be calculated as

691
$$P(M_k|D, H_i) = \frac{P(D|M_k,H_i)P(M_k|H_i)}{\sum_j P(D|M_j,H_i)P(M_k|H_i)} ,$$
(12)

692 Where $P(D|M_k,H_i)$ is the Bayesian model evidence (marginal likelihood) for $Model_k$.

693 The posterior probability for an individual model k under hypothesis H_i, $P(M_k|D,H_i)$, is not directly used,

694 as the posterior probability of H_i itself, $P(H_i|D)$ is of principal interest. Under Bayes' Theorem,

695
$$P(H_i|D) = \frac{P(D|H_i)P(H_i)}{\sum_j P(D|H_j)P(H_j)}, \qquad (13)$$

696 Where $P(D|H_i)$ is the marginal likelihood of H_i over all models to which it applies, $\{M_j \subset M, H_i\}$. According

697 to (49), this can be calculated as

698
$$P(D|H_i) = \sum_k P(D|M_k, H_i)P(M_k|H_i), \qquad (14)$$

with a summation instead of an integral because each model has a discrete prior probability as calculated

in Eq. (11).

Using the results of Eq. (14) in Eq. (13), we then calculate the posterior probability for each hypothesis,

pictured in **Fig 5A-D** and noted in **Table 2**. In this way, we can use Bayesian calculation rather than

parameter importance analysis (Galipaud et al., 2014, 2017; see Note S1-S3) to determine the posterior

probability of each model term. This also enables us to avoid bias in considering models with and without

certain model term, if an uneven number of candidate models contain a model term *vs.* do not contain the

term (69).

Posterior odds per hypothesis being investigated

All model terms and variables begin with a prior probability of 0.5. With equal prior probabilities across

all model term hypotheses, the posterior odds represented by $\frac{posterior\ probability}{(1-posterior\ probability)}$ is equivalent to the

Bayes Factor. Therefore, calculation of the posterior odds and the Bayes Factors for each model term are

equivalent.

A posterior probability of model term inclusion of 0.75 or more, or probability of 0.25 or less, would be

considered substantial evidence for inclusion or exclusion of that term, respectively (49). Given the nature

of the posterior odds, where a value of 2 indicates that one hypothesis is twice as likely to be true as the

other, we also consider posterior probabilities of 0.667 or more, or 0.333 or less, to be notable evidence for

inclusion or exclusion of the model term considered. We consider probabilities between 0.333 and 0.667 to

not have been significantly informed by the data.

Bayesian model averaging of parameter sets

Since Multinest returns multiple best-fitting parameter sets, each parameter in a model has a frequency distribution representing the values it takes on over these parameter sets. We thus consider each parameter using a probabilistic representation, per model (posterior marginal distribution) (**Fig 4B, 5E**; **Fig S5, S6B**). Since each candidate model is assigned a posterior probability as in Equation (10), all best-fitting parameter sets for that model can be assigned the same posterior probability. The frequency distribution of one parameter's values across a model's best-fitting parameter sets are thus weighted by its model's posterior probability. Then, the frequency distributions of weighted parameter values per model can be combined, representing the distribution of potential values of a particular parameter, weighted by model posterior probabilities. This way, parameter values in the distribution that come from models with a higher posterior probability (thus higher model evidence) will have more of an effect on the probabilistic representation, since they represent more likely values for the parameter.

To assemble representative fitted parameter sets for each candidate model, we used the first 1000 parameter sets from the Multinest equally weighted posterior samples per model. With up to 44 parameters and up to 5,891 models, the collection has 44 parameter columns and up to 5,891,000 rows representing a parameter vector. The collections were made per dataset.

Comparing parameter distributions

As above, each kinetic parameter has a frequency distribution representing 1000 fitted values per candidate model, meaning up to 5,891,000 fitted values across all models (weighted using Bayesian model averaging, as above). To compare parameter rates across models in the same dataset but with different topologies, we grouped each parameter according to the model topology from which it came. We then sampled 1000 values from the BMA-weighted distribution per kinetic parameter across all models of the same topology. We performed ANOVA followed by Tukey HSD at family-wise error rate (FWER) of 0.01, using the Python module statsmodels. Below an FWER of 0.01, we considered the sampled parameters significantly different

742 across models. We then repeated the sampling, ANOVA, and Tukey HSD for a total of 10 iterations. We

743 then averaged across determinations of significant/non-significant and if a parameter comparison across

744 model topologies was significantly different more often than it was not different, we considered the

745 parameter rates to be different comparing model topologies. The same methodology was used to compare

746 parameter rates across different datasets.

Generating a consolidated model of the SCLC tumor

A hypothesis (model term) whose posterior probability is further from its prior probability indicates more

information gained during the nested sampling process - more knowledge provided by the data. Conversely,

a posterior probability similar to the corresponding prior probability indicates that the data did not inform

our prior knowledge.

To unify the varying models into one view of SCLC biology, we brought together model probabilities from

each three-subtype topology per dataset (**Fig 5E**). To bring together the results for each three-subtype

topology results in the investigation of what appears as a four-subtype topology. In fact, if we are to envision

one model that can represent one system that generated all three datasets, it would need to include all four

subtypes. We consider this a reasonable practice in that all transition posterior probabilities in the three-

topology subtypes either were little informed by the data or had a value indicating that transitions are likely;

in addition, all Non-NE effects were either little informed by the data or had a value indicating that these

effects are unlikely. Posterior probabilities were not the same between three-subtype topologies, but these

trends of likely or unlikely model features generally agreed.

When consolidating models in this way, if model terms were part of multiple topologies (e.g., the A-to-N

transition is part of the A, N, and Y topology, best representing the RPM dataset, and the A, N, and A2

topology, best representing the SCLC-A cell line dataset) we took the posterior probability of the model

feature closer to 0.5. For example, the posterior probability for the A to N transition in the RPM dataset is

0.709 and the posterior probability for this same transition in the SCLC-A dataset is 0.626. Therefore, in

the four-subtype consolidated representation, the posterior probability for the A to N transition is 0.626.

This is the most conservative way to represent the knowledge gained by the data from the perspective of

the entire SCLC system, allowing for the most uncertainty to remain. We consider this practice as avoiding

claiming more certainty about model features than the data may provide.

CONTRIBUTIONS

Conceptualization, S.P.B, L.A.H., and C.F.L; methodology, S.P.B., L.A.H., and M.A.K.; software, S.P.B.,

L.A.H., and M.A.K.; formal analysis and investigation, S.P.B.; resources, C.F.L., J.S. and V.Q.; data

curation, S.P.B.; writing – original draft, S.P.B., C.F.L.; writing – review & editing, S.P.B., L.A.H., M.A.K.,

J.S., V.Q., C.F.L.; visualization, S.P.B.; supervision, V.Q., L.A.H., and C.F.L; funding acquisition, V.Q.

and C.F.L.

ACKNOWLEDGMENTS

The authors would like to thank Sarah Groves, Michael Irvin, and Christine Lovly for insightful

conversations and critical feedback on this work. This work was supported by the following funding

sources: S.P.B. was supported by the National Institutes of Health (NIH) [T32GM007347 and

T32LM012412] and the National Cancer Institute (NCI) [F30CA247078]; C.F.L. was supported by the

National Science Foundation (NSF) [MCB 1411482] and NSF CAREER Award [MCB 1942255]; C.F.L.

and V.Q. were supported by the National Institutes of Health (NIH) [U54-CA217450 and U01-CA215845].

AVAILABILITY

All the code to reproduce the data analysis and the figures is open source and can be found in this GitHub

repository: https://github.com/LoLab-VU/Bayes-MMI. This repository contains all the source code for each

step in the analysis.

Files needed to reproduce the figures without running the data analysis prior can be found in this DropBox

folder: https://www.dropbox.com/sh/4fqzpvu9hgyjicm/AABdfFlCenEuiOPgiH0TT-xqa?dl=0. Download

of these files and placement in the corresponding directories in the GitHub repository enables reproduction

of Figures 3-5.

REFERENCES

1. Baker RE, Peña JM, Jayamohan J, Jérusalem A. Mechanistic models versus machine learning, a fight worth fighting for the biological community? Biol Lett [Internet]. 2018 [cited 2022 Apr 20];14(5). Available from: https://royalsocietypublishing.org/doi/full/10.1098/rsbl.2017.0660

2. Stumpf MPH. Multi-model and network inference based on ensemble estimates: avoiding the madness of crowds. J R Soc Interface [Internet]. 2020 Oct 28 [cited 2021 Nov 3];17(171). Available from: https://royalsocietypublishing.org/doi/abs/10.1098/rsif.2020.0419

3. Kallenberger SM, Beaudouin J, Claus J, Fischer C, Sorger PK, Legewie S, et al. Intra- and interdimeric caspase-8 self-cleavage controls strength and timing of CD95-induced apoptosis. Sci Signal [Internet]. 2014 Mar 11 [cited 2022 Dec 14];7(316). Available from: https://www.science.org/doi/10.1126/scisignal.2004738

4. Kochen MA, Lopez CF. A Probabilistic Approach to Explore Signal Execution Mechanisms With Limited Experimental Data. Front Genet. 2020 Jul 10;11:686.

5. Spencer SL, Sorger PK. Measuring and Modeling Apoptosis in Single Cells. Cell. 2011 Mar 18;144(6):926–39.

6. Gay CM, Stewart CA, Park EM, Diao L, Groves SM, Heeke S, et al. Patterns of transcription factor programs and immune pathway activation define four major subtypes of SCLC with distinct therapeutic vulnerabilities. Cancer Cell. 2021 Mar 8;39(3):346-360.e7.

7. Groves SM, Ildefonso G v, Mcatee CO, Oliver TG, Weaver AM, Quaranta V. Archetype tasks link intratumoral heterogeneity to plasticity and cancer hallmarks in small cell lung cancer. Cell Syst [Internet]. 2022 [cited 2022 Dec 13];13(9):690–710. Available from: https://doi.org/10.1016/j.cels.2022.07.006

8. Tredennick AT, Hooker G, Ellner SP, Adler PB. A practical guide to selecting models for exploration, inference, and prediction in ecology. Ecology [Internet]. 2021 Jun 1 [cited 2022 Dec 13];102(6):e03336. Available from: https://onlinelibrary.wiley.com/doi/full/10.1002/ecy.3336

9. Parker WS. Ensemble modeling, uncertainty and robust predictions. Wiley Interdiscip Rev Clim Change [Internet]. 2013 May 1 [cited 2022 Dec 13];4(3):213–23. Available from: https://onlinelibrary-wiley-com.proxy.library.vanderbilt.edu/doi/full/10.1002/wcc.220

10. Garamszegi LZ, Mundry R. Multimodel-Inference in Comparative Analyses. In: Garamszegi LZ, editor. Modern Phylogenetic Comparative Methods and Their Application in Evolutionary Biology: Concepts and Practice [Internet]. Heidelberg: Springer Berlin; 2014 [cited 2022 Dec 13]. p. 305–31. Available from: http://www.

11. Galipaud M, Gillingham MAF, Dechaume-Moncharmont FX. A farewell to the sum of Akaike weights: The benefits of alternative metrics for variable importance estimations in model selection. Methods Ecol Evol. 2017 Dec 1;8(12):1668–78.

12. Galipaud M, Gillingham MAF, David M, Dechaume-Moncharmont FX. Ecologists overestimate the importance of predictor variables in model averaging: a plea for cautious interpretations. O'Hara RB, editor. Methods Ecol Evol [Internet]. 2014 Oct 1 [cited 2021 Apr 25];5(10):983–91. Available from: http://doi.wiley.com/10.1111/2041-210X.12251

833   13.   Lecca P. Machine Learning for Causal Inference in Biological Networks: Perspectives of This
834   Challenge. Frontiers in Bioinformatics. 2021 Sep 22;1:45.

835   14.   Burnham KP&, Anderson DR. Model Selection and Multimodel Inference: a Practical
836   Information-theoretic Approach. 2nd ed. Springer, New York. 2002.

837   15.   Hinne M, Gronau QF, van den Bergh D, Wagenmakers EJ. A Conceptual Introduction to Bayesian
838   Model Averaging. Adv Methods Pract Psychol Sci. 2020;3(2):200–15.

839   16.   Rudin CM, Brambilla E, Faivre-Finn C, Sage J. Small-cell lung cancer. Nat Rev Dis Primers. 2021
840   Dec 1;7(1).

841   17.   Rudin CM, Poirier JT, Byers LA, Dive C, Dowlati A, George J, et al. Molecular subtypes of small
842   cell lung cancer: a synthesis of human and mouse model data. Nature Reviews Cancer 2019 19:5
843   [Internet]. 2019 Mar 29 [cited 2021 Nov 4];19(5):289–97. Available from:
844   https://www.nature.com/articles/s41568-019-0133-9

845   18.   Zeman KG, Brzezniak CE, Carter CA. Recalcitrant small cell lung cancer: the argument for
846   optimism. J Thorac Dis [Internet]. 2017 Mar 1 [cited 2021 Nov 4];9(3):E295–6. Available from:
847   https://jtd.amegroups.com/article/view/12426/html

848   19.   Calbo J, van Montfort E, Proost N, van Drunen E, Beverloo HB, Meuwissen R, et al. A Functional
849   Role for Tumor Cell Heterogeneity in a Mouse Model of Small Cell Lung Cancer. Cancer Cell.
850   2011 Feb 15;19(2):244–56.

851   20.   Lim JS, Ibaseta A, Fischer MM, Cancilla B, O'Young G, Cristea S, et al. Intratumoural
852   heterogeneity generated by Notch signalling promotes small-cell lung cancer. Nature.
853   2017;545(7654):360–4.

854   21.   Williamson SC, Metcalf RL, Trapani F, Mohan S, Antonello J, Abbott B, et al. Vasculogenic
855   mimicry in small cell lung cancer. Nature Communications 2016 7:1 [Internet]. 2016 Nov 9 [cited
856   2022 May 30];7(1):1–14. Available from: https://www.nature.com/articles/ncomms13322

857   22.   Chan JM, Quintanal-Villalonga Á, Gao VR, Xie Y, Allaj V, Chaudhary O, et al. Signatures of
858   plasticity, metastasis, and immunosuppression in an atlas of human small cell lung cancer. Cancer
859   Cell [Internet]. 2021 Nov 8 [cited 2022 May 8];39(11):1479-1496.e18. Available from:
860   http://www.cell.com/article/S1535610821004979/fulltext

861   23.   Simpson KL, Stoney R, Frese KK, Simms N, Rowe W, Pearce SP, et al. A biobank of small cell
862   lung cancer CDX models elucidates inter- and intratumoral phenotypic heterogeneity. Nature
863   Cancer 2020 1:4 [Internet]. 2020 Apr 13 [cited 2022 May 31];1(4):437–51. Available from:
864   https://www.nature.com/articles/s43018-020-0046-2

865   24.   Huang YH, Klingbeil O, He XY, Wu XS, Arun G, Lu B, et al. POU2F3 is a master regulator of a
866   tuft cell-like variant of small cell lung cancer. Genes Dev [Internet]. 2018 Jul 1 [cited 2022 Dec
867   13];32(13–14):915–28. Available from: http://genesdev.cshlp.org/content/32/13-14/915.full

868   25.   Wooten DJ, Groves SM, Tyson DR, Liu Q, Lim JS, Albert R, et al. Systems-level network
869   modeling of Small Cell Lung Cancer subtypes identifies master regulators and destabilizers. PLoS
870   Comput Biol [Internet]. 2019 Oct 1 [cited 2021 Nov 4];15(10):e1007343. Available from:
871   https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1007343

26.   Ireland AS, Micinski AM, Kastner DW, Guo B, Wait SJ, Spainhower KB, et al. MYC Drives
Temporal Evolution of Small Cell Lung Cancer Subtypes by Reprogramming Neuroendocrine
Fate. Cancer Cell. 2020;38(1):60-78.e12.

27.   Ouadah Y, Rojas ER, Riordan DP, Capostagno S, Kuo CS, Krasnow MA. Rare Pulmonary
Neuroendocrine Cells Are Stem Cells Regulated by Rb, p53, and Notch. Cell [Internet]. 2019 Oct
3 [cited 2021 Nov 4];179(2):403-416.e23. Available from:
http://www.cell.com/article/S0092867419310165/fulltext

28.   Mollaoglu G, Guthrie MR, Böhm S, Brägelmann J, Can I, Ballieu PM, et al. MYC Drives
Progression of Small Cell Lung Cancer to a Variant Neuroendocrine Subtype with Vulnerability to
Aurora Kinase Inhibition. Cancer Cell. 2017 Feb 13;31(2):270–85.

29.   Shue YT, Drainas AP, Li NY, Pearsall SM, Morgan D, Sinnott-Armstrong N, et al. A conserved
YAP/Notch/REST network controls the neuroendocrine cell fate in the lungs. Nat Commun
[Internet]. 2022 [cited 2022 Dec 13];13(2690). Available from: https://doi.org/10.1038/s41467-
022-30416-2

30.   Jahchan NS, Lim JS, Bola B, Morris K, Seitz G, Tran KQ, et al. Identification and Targeting of
Long-Term Tumor-Propagating Cells in Small Cell Lung Cancer. Cell Rep [Internet]. 2016 Jul 19
[cited 2021 Nov 4];16(3):644–56. Available from:
http://www.cell.com/article/S2211124716307616/fulltext

31.   Fragoso TM, Bertoli W, Louzada F. Bayesian Model Averaging: A Systematic Review and
Conceptual Classification. International Statistical Review [Internet]. 2017 Apr 1 [cited 2021 Nov
3];86(1):1–28. Available from: https://onlinelibrary.wiley.com/doi/full/10.1111/insr.12243

32.   Chen X, Wang Y, Feng T, Yi M, Zhang X, Zhou D. The overshoot and phenotypic equilibrium in
characterizing cancer dynamics of reversible phenotypic plasticity. J Theor Biol. 2016 Feb
7;390:40–9.

33.   Fuentes-Garí M, Misener R, García-Munzer D, Velliou E, Georgiadis MC, Kostoglou M, et al. A
mathematical model of subpopulation kinetics for the deconvolution of leukaemia heterogeneity. J
R Soc Interface [Internet]. 2015 Jul 6 [cited 2021 Nov 4];12(108). Available from:
https://royalsocietypublishing.org/doi/abs/10.1098/rsif.2015.0276

34.   Gupta PB, Fillmore CM, Jiang G, Shapira SD, Tao K, Kuperwasser C, et al. Stochastic State
Transitions Give Rise to Phenotypic Equilibrium in Populations of Cancer Cells. Cell [Internet].
2011 Aug 19 [cited 2021 Nov 4];146(4):633–44. Available from:
http://www.cell.com/article/S0092867411008245/fulltext

35.   Greene JM, Levy D, Fung KL, Souza PS, Gottesman MM, Lavi O. Modeling intrinsic
heterogeneity and growth of cancer cells. J Theor Biol. 2015 Feb 21;367:262–77.

36.   Wang W, Quan Y, Fu Q, Liu Y, Liang Y, Wu J, et al. Dynamics between Cancer Cell
Subpopulations Reveals a Model Coordinating with Both Hierarchical and Stochastic Concepts.
PLoS One [Internet]. 2014 Jan 9 [cited 2021 Nov 4];9(1):e84654. Available from:
https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0084654

37.   Zhou JX, Pisco AO, Qian H, Huang S. Nonequilibrium Population Dynamics of Phenotype
Conversion of Cancer Cells. PLoS One [Internet]. 2014 Dec 1 [cited 2021 Nov 4];9(12):e110714.
Available from: https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0110714

38.   Olsen RR, Ireland AS, Kastner DW, Groves SM, Spainhower KB, Pozo K, et al. ASCL1 represses
a SOX9 + neural crest stem-like state in small cell lung cancer. Genes Dev [Internet]. 2021 [cited
2021 Jun 17];35:847–69. Available from:
http://www.genesdev.org/cgi/doi/10.1101/gad.348295.121.

39.   Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, et al. The Cancer Cell
Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. Nature 2012
483:7391 [Internet]. 2012 Mar 28 [cited 2022 May 28];483(7391):603–7. Available from:
https://www.nature.com/articles/nature11003

40.   Harris LA, Beik S, Ozawa PMM, Jimenez L, Weaver AM. Modeling heterogeneous tumor growth
dynamics and cell–cell interactions at single-cell and cell-population resolution. Curr Opin Syst
Biol. 2019 Oct 1;17:24–34.

41.   Charlebois DA, Balázsi G. Modeling cell population dynamics. In Silico Biol [Internet]. 2019 Jan
1 [cited 2021 Nov 10];13(1–2):21–39. Available from: https://github.com/dacharle42/MCPD

42.   Metzcar J, Wang Y, Heiland R, Macklin P. A Review of Cell-Based Computational Modeling in
Cancer Biology. JCO Clin Cancer Inform. 2019 Dec 4;(3):1–13.

43.   Portet S. A primer on model selection using the Akaike Information Criterion. Infect Dis Model
[Internet]. 2020 Jan 1 [cited 2022 Dec 4];5:111. Available from: /pmc/articles/PMC6962709/

44.   Eydgahi H, Chen WW, Muhlich JL, Vitkup D, Tsitsiklis JN, Sorger PK. Properties of cell death
models calibrated and compared using Bayesian approaches. Mol Syst Biol [Internet]. 2013 Jan 1
[cited 2021 Aug 16];9(1):644. Available from:
https://www.embopress.org/doi/full/10.1038/msb.2012.69

45.   Skilling J. Nested Sampling. AIP Conf Proc [Internet]. 2004 Nov 22 [cited 2021 Nov
3];735(1):395. Available from: https://aip.scitation.org/doi/abs/10.1063/1.1835238

46.   Feroz F, Hobson MP. Multimodal nested sampling: an efficient and robust alternative to Markov
Chain Monte Carlo methods for astronomical data analyses. Mon Not R Astron Soc [Internet].
2008 Feb 21 [cited 2021 Nov 3];384(2):449–63. Available from:
https://academic.oup.com/mnras/article/384/2/449/1023701

47.   Feroz F, Hobson MP, Bridges M. MultiNest: an efficient and robust Bayesian inference tool for
cosmology and particle physics. Mon Not R Astron Soc [Internet]. 2009 Oct 1 [cited 2021 Nov
3];398(4):1601–14. Available from: https://academic.oup.com/mnras/article/398/4/1601/981502

48.   Feroz F, Hobson MP, Cameron E, Pettitt AN. Importance Nested Sampling and the MultiNest
Algorithm. The Open Journal of Astrophysics [Internet]. 2019 Nov 27 [cited 2021 Nov
3];2(1):11120. Available from: https://github.com/farhanferoz/MultiNest.

49.   Kass RE, Raftery AE. Bayes Factors. J Am Stat Assoc. 1995;90(430):773–95.

50.   van de Schoot R, Depaoli S, King R, Kramer B, Märtens K, Tadesse MG, et al. Bayesian statistics
and modelling. Nature Reviews Methods Primers 2021 1:1 [Internet]. 2021 Jan 14 [cited 2021
Nov 3];1(1):1–26. Available from: https://www.nature.com/articles/s43586-020-00001-2

51.   Stewart CA, Gay CM, Xi Y, Sivajothi S, Sivakamasundari V, Fujimoto J, et al. Single-cell
analyses reveal increased intratumoral heterogeneity after the onset of therapy resistance in small-

cell lung cancer. Nat Cancer [Internet]. 2020 Apr 17 [cited 2021 Apr 25];1(4):423–36. Available from: https://doi.org/10.1038/s43018-019-0020-z

52. Hayford CE, Tyson DR, Jack Robbins C, Frick PL, Quaranta V, Harris LA. An in vitro model of tumor heterogeneity resolves genetic, epigenetic, and stochastic sources of cell state variability. PLoS Biol. 2021 Jun 1;19(6).

53. Paudel BB, Harris LA, Hardeman KN, Abugable AA, Hayford CE, Tyson DR, et al. A Nonquiescent "Idling" Population State in Drug-Treated, BRAF-Mutated Melanoma. Biophys J. 2018;

54. Chaffer CL, Brueckmann I, Scheel C, Kaestli AJ, Wiggins PA, Rodrigues LO, et al. Normal and neoplastic nonstem cells can spontaneously convert to a stem-like state. Proc Natl Acad Sci U S A [Internet]. 2011 May 10 [cited 2022 Dec 12];108(19):7950–5. Available from: https://www.pnas.org/doi/abs/10.1073/pnas.1102454108

55. Pisco AO, Brock A, Zhou J, Moor A, Mojtahedi M, Jackson D, et al. Non-Darwinian dynamics in therapy-induced cancer drug resistance. Nat Commun [Internet]. 2013 Sep 18 [cited 2022 Dec 12];4(1):1–11. Available from: https://www.nature.com/articles/ncomms3467

56. Yang G, Quan Y, Wang W, Fu Q, Wu J, Mei T, et al. Dynamic equilibrium between cancer stem cells and non-stem cancer cells in human SW620 and MCF-7 cancer cell populations. British Journal of Cancer 2012 106:9 [Internet]. 2012 Apr 3 [cited 2022 Dec 12];106(9):1512–9. Available from: https://www.nature.com/articles/bjc2012126

57. Hata AN, Niederst MJ, Archibald HL, Gomez-Caraballo M, Siddiqui FM, Mulvey HE, et al. Tumor cells can follow distinct evolutionary paths to become resistant to epidermal growth factor receptor inhibition. Nat Med. 2016;22(3):262–9.

58. Ku SY, Rosario S, Wang Y, Mu P, Seshadri M, Goodrich ZW, et al. Rb1 and Trp53 cooperate to suppress prostate cancer lineage plasticity, metastasis, and antiandrogen resistance. Science (1979) [Internet]. 2017 Jan 6 [cited 2022 Dec 12];355(6320):78–83. Available from: https://www.science.org/doi/10.1126/science.aah4199

59. Mu P, Zhang Z, Benelli M, Karthaus WR, Hoover E, Chen CC, et al. SOX2 promotes lineage plasticity and antiandrogen resistance in TP53-and RB1-deficient prostate cancer. Science (1979) [Internet]. 2017 Jan 6 [cited 2022 Jan 19];355(6320). Available from: https://www.science.org/doi/abs/10.1126/science.aah4307

60. Risom T, Langer EM, Chapman MP, Rantala J, Fields AJ, Boniface C, et al. Differentiation-state plasticity is a targetable resistance mechanism in basal-like breast cancer. Nat Commun [Internet]. 2018;9(3815):1–17. Available from: www.nature.com/naturecommunications

61. Shaffer SM, Dunagin MC, Torborg SR, Torre EA, Emert B, Krepler C, et al. Rare cell variability and drug-induced reprogramming as a mode of cancer drug resistance. Nature. 2017;546(7658):431–5.

62. Zou M, Toivanen R, Mitrofanova A, Floch N, Hayati S, Sun Y, et al. Transdifferentiation as a Mechanism of Treatment Resistance in a Mouse Model of Castration-Resistant Prostate Cancer. Cancer Discov [Internet]. 2017;7(7):737–49. Available from: www.aacrjournals.org

63. Watson SS, Dane M, Chin K, Jonas O, Gray JW, Korkola JE. Microenvironment-Mediated
Mechanisms of Resistance to HER2 Inhibitors Differ between HER2+ Breast Cancer Subtypes.
Cell Syst [Internet]. 2018 [cited 2022 Jan 19];6:329–42. Available from:
https://doi.org/10.1016/j.cels.2018.02.001

64. Kwon MC, Proost N, Song JY, Sutherland KD, Zevenhoven J, Berns A. Paracrine signaling
between tumor subclones of mouse SCLC: a critical role of ETS transcription factor Pea3 in
facilitating metastasis. 2015 [cited 2022 May 30]; Available from:
http://www.genesdev.org/cgi/doi/10.1101/gad.262998.115.

65. Lopez CF, Muhlich JL, Bachman JA, Sorger PK. Programming biological models in Python using
PySB. Mol Syst Biol [Internet]. 2013 Jan 1 [cited 2021 Nov 4];9(1):646. Available from:
https://onlinelibrary.wiley.com/doi/full/10.1038/msb.2013.1

66. Mitra ED, Hlavacek WS. Parameter estimation and uncertainty quantification for systems biology
models. Curr Opin Syst Biol [Internet]. 2019 [cited 2022 Mar 10];18:9–18. Available from:
https://doi.org/10.1016/j.coisb.2019.10.006

67. van Ravenzwaaij D, Cassey P, Brown SD. A simple introduction to Markov Chain Monte–Carlo
sampling. Psychon Bull Rev. 2018 Feb 1;25(1):143–54.

68. Kruschke John. Doing Bayesian data analysis : a tutorial introduction with R. 2nd ed. Elsevier
Science & Technology; 2014.

69. Garthwaite PH, Mubwandarikwa E. SELECTION OF WEIGHTS FOR WEIGHTED MODEL
AVERAGING. Aust N Z J Stat. 2010;52(4):363–82.

1011