Thermal cycling-hyperthermia ameliorates cognitive impairment of intracerebroventricular Aβ$_{25-35}$-induced Alzheimer's disease in C57BL/6 mice

Yu-Yi Kuo[1,2], Wei-Ting Chen[1,2], Guan-Bo Lin[1,2], You-Ming Chen[2,3], Hsu-Hsiang Liu[2,3], and Chih-Yu Chao[1,2,3,*]

[1] Department of Physics, Lab for Medical Physics & Biomedical Engineering, National Taiwan University, Taipei 10617, Taiwan

[2] Molecular Imaging Center, National Taiwan University College of Medicine, Taipei 10051, Taiwan

[3] Graduate Institute of Applied Physics, Biophysics Division, National Taiwan University, Taipei 10617, Taiwan

* Corresponding author

E-mail: cychao@phys.ntu.edu.tw (CYC)

Abstract

Despite continuation of some controversies, Alzheimer's disease (AD), the most common cause of dementia nowadays, has been widely believed to derive mainly

from excessive β-amyloid (Aβ) aggregation, that would increase reactive oxygen

species (ROS) and induce neuroinflammation, leading to neuron loss and cognitive

impairment. Existing drugs on Aβ have been ineffective or offer only temporary relief

at best, due to blood-brain barrier or severe side effects. The study employed thermal

cycling-hyperthermia (TC-HT) as an alternative AD therapy and compared its effect

with continuous hyperthermia (HT) *in vivo*. It established an AD mice model via

intracerebroventricular (i.c.v.) injection of $Aβ_{25-35}$, proving that TC-HT is much more

effective in alleviating its performance decline in Y-maze and NOR test, in

comparison with HT. In addition, TC-HT also exhibits a better performance in

decreasing the hippocampal Aβ and BACE1 expressions as well as the

neuroinflammation markers Iba-1 and GFAP levels. Furthermore, the study finds that

TC-HT can elevate more protein expressions of IDE and antioxidative enzyme SOD2

than HT. Besides, after establishment of neuroprotective mechanism, removal of

TC-HT-induced ROS can further augment protection of neural cells against Aβ. In

sum, the study proves the potential of TC-HT in AD treatment, which can be put into

clinical application with the use of focused ultrasound (FUS).

Introduction

Alzheimer's disease (AD) has been the most common cause of dementia nowadays, accounting for over 60% of all the cases [1]. One of the most notable hallmarks of AD pathogenesis is the accumulation in brain of toxic β-amyloid (Aβ) peptides [2], which are cleaved from the amyloid precursor protein (APP) by β- and γ-secretase [3]. Such abnormal Aβ deposits can activate neuroinflammation [4], generate reactive oxygen species (ROS) [5, 6], and cause neuron dysfunction and synapse loss, leading to memory loss and cognitive impairment [7, 8]. Therefore, Aβ appears to be the foremost target in AD therapy [9].

Currently, drug therapy is the mainstream AD treatment, aiming to suppress the generation [10], disrupt the aggregation [11], and promote the clearance [12] of Aβ. Drugs on proteases, such as insulin degrading enzyme (IDE), for alleviating Aβ degradation also have significant potential in AD treatment [13]. However, at present AD drugs have high failure rates, due to difficulty penetrating blood-brain barrier, on top of severe side effects, such as liver toxicity, strong autoimmune response, or carcinogenicity [14]. Hence, there is a need for a safe and effective alternative AD therapy.

Heat has long been regarded as an alternative therapy with significant potential,

given its proven therapeutic effect in physiotherapy, ophthalmology, and cardiology

[15]. Hyperthermia (HT), for instance, has been applied effectively in cancer

treatment, either for directly inhibiting tumor growth [16] or supplementing

radiotherapy and certain chemotherapies to augment their efficacy [17]. In addition,

moderate thermal stress can stimulate the expression of neuroprotective proteins. For

example, HT is capable of increasing the expression of heat shock protein (HSP)

family [18], some of whose members can protect neuron cells against various stresses

to achieve neuroprotection [19, 20]. Besides, HT treatment is conducive to increase

the levels of IDE [21] and antioxidant proteins, such as nuclear factor erythroid

2–related factor 2 (Nrf2) and superoxide dismutase 2 (SOD2) [22, 23].

However, continuous high-temperature exposure can be thermotoxic to human

neural cells, according to some reports [24]. Heat stroke, featuring abnormally high

core body temperature ($> 40°C$), may cause cerebellar injury, hypertension, and

central nervous system derangement [25, 26]. Whole-body HT treatment has also

been reported to cause neurological and systemic complications, such as nausea,

impaired pressure-flow autoregulation, brain edema, and arrhythmias [27]. On the

other hand, although a study showed that long-term heat exposure could reduce the

risk of memory loss among middle-aged men, it was time-consuming without certain

curative effect [28]. Therefore, it is imperative to find a tolerable and effective

thermal treatment for AD.

Our previous study put forth a novel thermal treatment, thermal-cycling

hyperthermia (TC-HT), demonstrating its therapeutic effect in anticancer treatment

[29, 30] and neuroprotection [31] *in vitro*, with minimal side effects. The study

applied TC-HT to treat mice with AD, induced by intracerebroventricular (i.c.v.)

injection of $A\beta_{25-35}$, for comparison with the therapeutic effect of HT. In both cases,

heat was applied to mouse brain specifically, to avoid potential side effect of

whole-body HT.

In this paper, behavioral tests show that TC-HT significantly improves

$A\beta$-induced memory impairment, with an effect much better than HT. In addition,

TC-HT also exhibits a much better performance than HT in decreasing $A\beta$ and

β-secretase (BACE1) expressions, suppressing neuroinflammation markers ionized

calcium-binding adapter molecule 1 (Iba-1) and glial fibrillary acidic protein (GFAP)

levels, and increasing the expression levels of IDE and SOD2 enzymes. Furthermore,

the study shows that removal of TC-HT-induced ROS can further augment the

protective effect of TC-HT in an *in vitro* neural model. In sum, these results

underscore the potential of TC-HT as an alternative AD therapy.

Materials and Methods

Ethics statement

All animal experiments were conducted according to the regulations approved by the

Animal Ethical Committee of National Taiwan University, and the approval protocol

number was NTU-109-EL-174. Mice were monitored every day for signs of distress.

The humane endpoints of the mice included decreased socialization, anorexia, weight

loss of 15%, hunching, or the inability to evade handling. All hydrodynamic

injections and thermal treatments were conducted under anesthesia with 5%

isoflurane (Panion & BF Biotech Inc., Taipei, Taiwan), and all efforts were made to

minimize suffering. For sacrifice, mice were humanely euthanized via carbon dioxide

(CO_2) gas asphyxiation followed by cervical dislocation and were considered

non-survivors.

Animals

C57BL/6 male mice (6-8 weeks old, 22-26 g) were obtained from National

Laboratory Animal Center (Taipei, Taiwan). All mice were housed in groups of 6

mice per cage (32 cm × 16 cm × 14 cm) and maintained in a specific-pathogen-free

(SPF) room under controlled temperature (22±2 °C) and relative humidity (55±10%)

in Animal Resource Center of National Taiwan University. All mice were subjected

to a 12:12 h light/dark cycle (lights on at 7:00 a.m.) with free access to food and water.

All experiments were conducted with mice at age 6-8 weeks. All mice were randomly

categorized into the control group, Aβ group, TC-HT group, and HT group.

β-amyloid administration

Aβ$_{25-35}$ (Sigma-Aldrich; Merck KGaA, Darmstadt, Germany) was dissolved in 0.9%

sterile saline solution to a concentration of 1 mg/mL and then incubated at 37°C for 7

days for fibril aggregation. At the beginning of the experiment (Day 0), 10 μL Aβ

solution was i.c.v.-injected into the mice's brains of the Aβ, TC-HT, and HT groups

according to the protocol described in the previous study [32]. 10 μL of sterile saline

solution was i.c.v.-injected into the mice's brains of the sham-operated control group

HT and TC-HT treatments

To focus on the thermal effect on mice's brain and minimize the potential interference

outside of brains by the conventional whole-body hyperthermia methods [33-35], we

adopted a modified method to heat mice's heads locally. As shown in **Fig 1A**, both

HT and TC-HT thermal treatments were applied by a temperature controlling and

heating system equipped with a DC power supply (9303D, ABM Industries Inc., NY,

USA), a temperature/process controller (PZ900, RKC instrument Inc., Tokyo, Japan),

and a 19 mm × 19 mm polyimide thermofoil for heating (HK5578R4.6L12A, Minco

Products Inc., Fridley, MN, USA). Mice were anesthetized with 5% isoflurane during

thermal treatments. Mice's head fur was shaved and the thermofoil was taped onto the

bare skin with a thin layer of vaseline for better attachment. The thermocouple

measuring tip (K-type) was located under the central region of the thermalfoil, as

shown in **Fig 1B**. The actual temperatures of mice's heads were measured via the

K-type thermocouple and monitored by the temperature/process controller. For

TC-HT treatment, the temperature/process controller was set to implement a 10-cycle

repeated procedure, in which each cycle is consisted of a desired high-temperature

stage for a period of time, followed by a natural thermal dissipating stage for 30 sec at

low-temperature setting 37 °C (**Fig 1C**). In the study, the high-temperature stage in

TC-HT treatment was held at 42 °C for 6 min with a 0.5 °C tolerance. For HT

treatment, the temperature was kept at 42 ± 0.5 °C continuously for 1 h. Monitored

temperatures of mice's heads by HT and TC-HT thermal treatments were shown in

Fig 1D.

Fig 1. Experimental setup and design of TC-HT.

(A) The setup of the temperature controlling and heating system, containing a DC

power supply, a temperature/process controller, and a thermofoil for heating. (B) The

thermal treatment was locally performed on mice's heads by thermofoil, and their

temperatures were measured via a K-type thermocouple, whose measuring tip was

placed in the central circled region under the thermofoil. (C) Schematic diagram of

the TC-HT temperature-controlled programs with parameter settings, and (D) the

temperatures of mice's heads by HT and TC-HT thermal treatments were recorded

every 10 sec throughout the heating period.

Experimental design

At the beginning of the experiment (Day 0), 10 μL Aβ solution was i.c.v.-injected into

the mice's brains of the Aβ, TC-HT, and HT groups. On day 4, 8, and 12, the

respective thermal treatment was applied to the mice of the TC-HT and HT groups.

Y-maze and NOR tests were assessed on day 13 and day 14, respectively. Brain

tissues were collected after the Y-maze test for the following western blot analyses.

The experimental schedule was shown in **Fig 2**.

Fig 2. The experimental schedule.

Aβ solution was i.c.v.-injected at day 0, and the respective thermal treatment was

implemented at day 4, 8, and 12. Behavioral tests of NOR and Y-maze were carried

out at day 13 and 14, respectively. After the Y-maze test, mice were weighted and

then euthanized to collect brain tissues for the following western blot analyses.

Y-maze test

The Y-maze test was used to assess the spatial working memory of mice. The

apparatus is consisted of three identical arms (40 cm × 13 cm × 6.5 cm)

interconnected at a central junction at 120° each. The natural tendency of rodents is to

explore novel environments, which constitutes their spontaneous alteration behavior

to explore a new arm in Y-maze. The assessed mouse was placed at the end of the

start arm, and was allowed to explore all three arms freely for 8 min. Olfactory cues

in the Y-maze were removed by 70% ethanol after each assessment. The behavioral

performance was recorded, and the spontaneous index was calculated as follows:

(number of correct alternations) / (total number of arm entries – 2) × 100. In addition,

the total number of arm entries in the Y-maze was used as a measure to characterize

locomotor activity.

Novel object recognition (NOR) test

We used the NOR test to assess the recognition memory of animals. The NOR test

consists of a habituation phase, a familiarization phase and a test phase. All mice in

the habituation phase were placed in a 40 cm × 40 cm × 30.5 cm empty box for 8 min.

In the familiarization phase, the assessed mouse was placed in the box for 8 min with

2 identical Lego blocks, which were placed close to two diagonal corners of the box.

After the familiarization phase, the mouse was returned to the home cage for 1 h and

then was placed back into the testing box for the test phase. One of the original

objects was replaced with a novel Lego block of a different shape in the test phase,

and the behavioral performance of the mouse was recorded with an overhead video

camera for 8 min. After each assessment in all phases, olfactory cues in the box and

on the Lego blocks were removed by 70% ethanol. Exploration was defined as

sniffing or touching the object with the nose and/or forepaws. The duration of the

exploring time was calculated for the discrimination index. A discrimination index

was defined as follows: (Tn-To) / (Tn+To), where Tn and To stand for the exploration

time of the novel and original objects, respectively.

Brain tissue collection and western blot sample preparation

Mice were euthanized by CO_2 gas. After confirming the death of mice, brains were

taken out quickly and rinsed with phosphate buffered saline (PBS) (HyClone; GE

Healthcare Life Sciences, Chicago, IL, USA) supplemented with 1% fresh protease

inhibitor cocktail (EMD Millipore, Billerica, MA, USA). Whole hippocampi were

rapidly dissected on an ice-cold plate, and then stored in ice-cold RIPA lysis buffer

(EMD Millipore) supplemented with 1% fresh protease inhibitor cocktail.

Hippocampal tissues were homogenized via ultrasound processing (5 sec rectangular

pulse with 5 sec interval for 3 times, amplitude 50%, by a Q125 sonicator; Qsonica,

Newton, CT, USA) and then centrifuged at 23000×g for 30 min at 4°C. The

supernatants were collected and the protein concentration was quantified by the

Bradford protein assay (BioShop Canada Inc., Burlington, Ontario, Canada). Protein

samples were stored at -80°C for subsequent western blotting assessments.

Western blot analysis

Proteins from the homogenized hippocampi were resolved by sodium dodecyl sulfate

polyacrylamide gel electrophoresis (SDS-PAGE) and then electroblotted onto the

polyvinylidene difluoride (PVDF) membranes (EMD Millipore). Blotted membranes

were blocked with 5% skim milk in Tris-buffered saline supplemented with 0.1%

Tween-20 (TBST) (both from BioShop Canada Inc.) for 1 h at room temperature (RT)

and then incubated overnight with diluted primary antibodies in blocking buffer at 4

°C, followed by three rinses with TBST washing buffer. Primary antibodies employed

in this study were listed as following: anti-Aβ (sc-28365, Santa Cruz Biotechnology,

Inc., Dallas, TX, USA), anti-IDE (ab32216, Abcam, Cambridge, UK), anti-BACE1

(#5606), anti-SOD2 (#13141), anti-Iba-1 (#17198) (Cell Signaling Technology,

Danvers, MA, USA), anti-GFAP (GTX100850), and anti-β-actin (GTX110564)

(Gentex, Irvine, CA, USA). After washing with TBST, the membranes were

incubated with horseradish peroxidase-conjugated goat anti-mouse (for probing

anti-Aβ primary antibody) (ab205719, Abcam) or goat anti-rabbit (for probing the

other primary antibodies) secondary antibodies (AB_2313567, Jackson

ImmunoResearch Laboratories, Inc., West Grove, PA, USA) for 1 h at RT. All the

antibodies were diluted according to the manufacturer's instructions to their optimal

concentration. The protein bands were visualized by an enhanced chemiluminescence

substrate (Advansta, San Jose, CA, USA) and detected by an imaging system

Amersham Imager 600 (GE Healthcare Life Sciences). The images were analyzed

with Image Lab software (Bio-Rad Laboratories, Inc.). For normalization of proteins,

β-actin was used as the loading control to determine the relative folds of targeted

proteins.

239   ***In vitro* cell culture and hydrogen gas (H$_2$) treatment**

The human neuroblastoma SH-SY5Y cells were purchased from American Type

Culture Collection (Manassas, VA, USA). The cell culture medium was MEM/F-12

mixture (HyClone; GE Healthcare Life Sciences) containing 10% fetal bovine serum

(HyClone; GE Healthcare Life Sciences) and 1% penicillin-streptomycin (Gibco Life

Technologies, Grand Island, NY, USA), supplemented with 1mM sodium pyruvate

(Sigma-Aldrich; Merck KGaA) and 0.1 mM non-essential amino acids (Invitrogen

Life Technologies, Grand Island, NY, USA). Cells were maintained in a humidified

incubator composed of 5% CO_2 at 37°C. Before the experiment, SH-SY5Y cells were

subcultured and seeded in 96-well plates (Thermo Fisher Scientific Inc., Waltham,

MA, USA) overnight. Cells were treated with 25 μM Aβ for 1 h and then subjected to

the TC-HT treatment for about 2 h. The TC-HT treatment was implemented by the

Thermal Cycler (Applied Biosystems; Thermo Fisher Scientific, Inc.) and the *in vitro*

TC-HT parameter settings were described in our previous study [31]. In the

experiment, H$_2$-mixed gas was composed of 75% H$_2$, 20% oxygen gas (O$_2$), and 5%

CO_2 gas (vol/vol/vol), as described elsewhere [36]. After continuous exposure to the

H$_2$-mixed gas, it has been observed that H$_2$ concentration of the culture medium in a

closed acrylic box gets saturated within 20~30 min at atmospheric pressure [37].

Upon 12~16 h after TC-HT treatment, cultured SH-SY5Y cells were sent into an

acrylic chamber of volume about 450 mL. Mixed gas consisting of 75% H$_2$, 20% O$_2$,

and 5% CO$_2$ (vol/vol/vol) were well mixed and then flowed into the chamber at a rate

of 400 mL/min for 90 min under normal pressure. Upon 4 days after the Aβ

administration, the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide

(MTT) (Sigma-Aldrich; Merck KGaA) assay was used to analyze the cell viability of

SH-SY5Y cells. In brief, the culture medium was replaced with MTT solution (0.5

mg/mL in the MEM/F12 mixture) and incubated at 37°C for 4 h. The formazan

crystals were dissolved by equal volume of the solubilizing buffer containing 10%

sodium dodecyl sulfate (SDS) (Bioshop Canada Inc.) solution in 0.01 N hydrochloric

acid (HCl) (Echo Chemical Co. Ltd., Taiwan) at 37 °C overnight. The absorbance at

570 nm of each well was analyzed by Multiskan GO microplate spectrophotometer

(Thermo Fisher Scientific Inc.), and the background absorbance at 690 nm was

subtracted. The cell viability of each group was expressed as percentage of the

untreated control.

Statistical analysis

All statistical results were shown as mean ± standard deviation (S.D.). Statistical

analysis was conducted with OriginPro 2015 software (OriginLab Corporation,

Northampton, MA, USA) using one-way analysis of variance (ANOVA), followed by

Tukey's post hoc test. P-value < 0.05 was considered to indicate a statistically

significant difference.

Results

TC-HT improves Aβ-induced recognition impairments in mice

The study employed Y-maze test and NOR test to examine the therapeutic effect of

TC-HT and HT on AD mice. In the Y-maze test, the overall percentage of each

mouse's spontaneous index was computed to show its short-term memory function.

As exhibited in **Fig 3A**, the spontaneous index of the Aβ group is inhibited

significantly, ranging from 60.3% to 44.1%, compared with the control group. TC-HT

significantly improves the memory deficiency, elevating the spontaneous index to

61.6%, notably different from HT, which only causes an insignificant improvement

(spontaneous index 46.8%). Besides, there is no significant difference in the total

number of arm entries among the four groups (**Fig 3B**), suggesting that i.c.v. Aβ

injection or both thermal treatments didn't affect the locomotor activities of mice. On

the other hand, NOR test also confirmed that TC-HT is more effective in alleviating

Aβ-induced memory impairment than HT. As shown in **Fig 3C**, along with memory

decline, the discrimination index of the Aβ group plunged from 25.1% down to

-26.2%, which bounces back to 19.2% after TC-HT treatment, significantly

outperforming HT treatment, which only raises the discrimination index to -3.1%.

Thus, TC-HT exhibited a much stronger reversing effect on cognitive impairment

than HT in both Y-maze and NOR tests.

Fig 3. Effect of TC-HT and HT treatments on mice's behavioral performances.

The recovery effects of Aβ-induced recognition impairments after TC-HT and HT

treatments are assessed via (A) the spontaneous index and (B) the total number of arm

entries in the Y-maze test, and (C) the discrimination index in the NOR test. Data are

presented as mean ± S.D. (n = 9 in each group). One-way ANOVA with Tukey's post

hoc test is used to determine statistical significance. (*$P < 0.05$, comparison between

indicated groups).

TC-HT down-regulates Aβ secretion and accumulation in mouse brain

The study first analyzed the accumulated Aβ levels in hippocampi, confirming the

significant effect of TC-HT treatment in repairing Aβ-induced cognitive impairment,

as it reduces the hippocampal Aβ level from 1.34-fold to 0.88-fold compared to the

control group (**Fig 4A**). HT treatment, by comparison, exhibited only slight inhibitive

effect on the hippocampal Aβ level (1.24-fold). On the other hand, it is realized that

Aβ level in mouse brain is partially attributed to the pathological changes of BACE1

[38]. In **Fig 4B**, our results show that i.c.v.-injected Aβ triggers a significant increase

in hippocampal BACE1 expression (1.30-fold) compared with the control group. The

study finds that the elevation of BACE1 expression is significantly suppressed by the

TC-HT treatment (0.89-fold), but only slightly decreased by the HT treatment

(1.19-fold). It's noteworthy that there is no significant difference in BACE1 levels

between the Aβ group and the HT group, suggesting that conventional HT treatment

is ineffective in stopping abnormal Aβ secretion in mouse brain. To the best of our

knowledge, this is the first time that a thermal treatment exhibited an inhibitory ability

on the expression level of BACE1. In sum, the study shows that TC-HT appears to be

much more effective AD treatment than HT, as it's more capable of regulating Aβ

levels actively and inhibiting BACE1 level in hippocampus.

Fig 4. Effect of TC-HT and HT treatments on the expression levels of Aβ and

BACE1 in hippocampi.

Representative western blots and the quantifications of (A) Aβ and (B) BACE1 are

presented for evaluating anti-Aβ effect in hippocampi. β-actin was used as loading

control. Data are presented as mean ± S.D. (n = 9 in each group). One-way ANOVA

with Tukey's post hoc test is used to determine statistical significance. (*$P < 0.05$,

comparison between indicated groups).

TC-HT ameliorates Aβ-induced neuroinflammation in mouse brain

Neuroinflammation is another hallmark of AD, as the risk of cognitive impairment is

highly associated with inflammatory level [39]. The study finds that the expression

levels of both hippocampal Iba-1 and GFAP, the activation markers of microglia cells

and astrocytes, respectively, are significantly enhanced in the Aβ group, compared

with the control group (**Fig 5A** and **5B**). After the TC-HT treatment, both Iba-1 and

GFAP levels drop significantly, while the HT treatment only slightly attenuates their

expression levels. There is no significant difference between the Aβ group and the HT

group in the expressions of both Iba-1 and GFAP. The results indicate that TC-HT

treatment can better ameliorate Aβ-induced neuroinflammation than HT treatment *in*

vivo.

Fig 5. Effect of TC-HT and HT treatments on the expression levels of

neuroinflammatory markers Iba-1 and GFAP in hippocampi.

Representative western blots and the quantifications of (A) Iba-1 and (B) GFAP are

presented for evaluating anti-neuroinflammation effect in hippocampi. β-actin was

used as loading control. Data are presented as mean ± S.D. (n = 9 in each group).

One-way ANOVA with Tukey's post hoc test is used to determine statistical

significance. (*$P < 0.05$, comparison between indicated groups).

TC-HT promotes the expressions of IDE and SOD2 proteins in mouse brain

With IDE long regarded as a major endogenous enzyme attributing to Aβ degradation,

the alteration of IDE activity and expression can cause Aβ accumulation, thereby

increasing the risk of AD [40]. **Fig 6A** shows that while Aβ injection slightly

decreases the hippocampal IDE level (0.88-fold), TC-HT treatment significantly

increases the IDE expression level in hippocampi up to 1.38-fold. By contrast, HT

treatment performs an insignificant altering effect on IDE expression (0.91-fold),

compared with the Aβ group. The findings show that TC-HT can intensify Aβ

clearance via elevating the IDE expression level in hippocampi, while HT is

ineffective in promoting the hippocampal IDE level and thus cannot contribute to the

removal of abnormal hippocampal Aβ in mice.

Antioxidant protein SOD2 also has significant potential in AD treatment, capable

of decreasing hippocampal superoxide and amyloid plaque in AD mice [41]. **Fig 6B**

shows that after i.c.v. Aβ injection, the SOD2 expression level drops significantly

(0.62-fold), which, though, recovers significantly in the TC-HT group (1.41-fold),

much higher than recovery to 0.86-fold after HT treatment. The results show that

TC-HT can activate stronger antioxidant activity in hippocampus against the

Aβ-induced oxidative stress than continuous HT *in vivo*.

Fig 6. Effect of TC-HT and HT treatments on the expression levels of neuroprotective proteins IDE and SOD2 in hippocampi.

Representative western blots and the quantifications of (A) IDE and (B) SOD2 are presented for evaluating Aβ clearance and antioxidant effect in hippocampi, respectively. β-actin was used as loading control. Data are presented as mean ± S.D. (n = 9 in each group). One-way ANOVA with Tukey's post hoc test is used to determine statistical significance. (*$P < 0.05$, comparison between indicated groups).

H_2 treatment produces auxiliary neuroprotective effect against Aβ after TC-HT treatment *in vitro*

To examine whether removal of excessive ROS after TC-HT treatment can intensify the neuroprotective effect against Aβ, H_2 treatment as mentioned in Materials and Methods section was introduced as an exogenous ROS scavenger in an *in vitro* model. **Fig 7A** shows the significant suppression of the viability of SH-SY5Y cells to 60.9% under the insult of 25 μM Aβ, which recovers greatly to 82.5% after TC-HT treatment. Moreover, we find that with a 90 min H_2 treatment at 12~16 h after TC-HT treatment,

the cell viability can bounce back to a near-normal level (96.0%), exhibiting a better

neuroprotective effect than TC-HT treatment alone. However, it should be noted that

administration of H_2 treatment less than 12 h after TC-HT treatment fails to further

boost the viability of SH-SY5Y cells. On the other hand, H_2 treatment alone does not

show any significant protective effect on SH-SY5Y cells against Aβ (**Fig 7B**). These

findings show that while removal of ROS via H_2 treatment alone has no effect against

Aβ, it can further enhance the neuroprotective effect of TC-HT treatment against Aβ,

exhibiting a better neuroprotective effect than TC-HT treatment alone. This could be

attributed to removal of TC-HT-induced ROS with H_2 treatment after establishment

of neuroprotective mechanism at 12~16 h after TC-HT treatment. These results shed

light on the approach in developing AD treatment.

Fig 7. H_2 treatment as a ROS scavenger helps to promote further the

neuroprotective effect of TC-HT against Aβ *in vitro*.

(A) Cell viabilities of SH-SY5Y cells treated with Aβ, Aβ + TC-HT, and Aβ +

TC-HT + H_2 treatment, respectively. The data show that H_2-assisted TC-HT treatment

exhibits a better neuroprotective effect than TC-HT treatment alone. In the other

experiment, SH-SY5Y cells were subjected directly to H_2 treatment after the

administration of 25 µM Aβ. Cell viabilities of SH-SY5Y cells treated with Aβ and

Aβ + H_2 treatment alone are respectively shown in (B). The result indicates that H_2

treatment alone has no effect against Aβ. Data are presented as mean ± S.D. (n = 6 in

each group). One-way ANOVA with Tukey's post hoc test is used to determine

statistical significance. (*P < 0.05, comparison between indicated groups).

Discussion

Given prevalence of AD in our society in recent years, many scientists have taken part

in research on its treatment with various approaches, including Aβ hypothesis, one of

the most noticeable pathogenesis of AD [42]. Aβ peptides could be secreted in brains,

but their abnormal depositions, especially in hippocampus, could induce severe

neurodegeneration and synapse loss, thereby impairing memory and cognitive

abilities [7, 8]. The study employed Aβ$_{25\text{-}35}$, peptide fragment of Aβ$_{1\text{-}42}$ considered the

most pathogenic Aβ species in AD [43, 44], as the model. Development of AD drugs

has been confronted with two major obstacles, namely blood-brain barrier (BBB) and

severe side effects [14, 45]. Hence, the application of heat has potential as an

alternative AD treatment, due to its capability to intensify neuroprotective effect [31]

and sidestep BBB, targeting the diseased brain directly. The study embraced TC-HT,

a modified heat treatment with higher safety and superior protective effect than HT

[31], to investigate the therapeutic potential of thermal treatment on AD *in vivo*.

With memory loss being the first and main symptom of AD [46], AD studies

with mice as model have often employed Y-maze and NOR tests to assess short-term

memory and learning memory [47, 48]. The study probed the therapeutic effect of HT

and TC-HT treatment on Aβ-induced memory impairment, finding that TC-HT

treatment can significantly improve the performance of spontaneous index (**Fig 3A**)

and help memory-impaired mice recognize and remember new objects (**Fig 3C**). The

study demonstrates significant effect of TC-HT treatment in recovering spatial

working memory and recognition memory impaired by Aβ. In comparison, although

HT treatment also enhances behavioral performance, the extent of improvement

significantly lags behind that of TC-HT treatment in both tests. Collectively, TC-HT

treatment exhibits a higher therapeutic potential than HT treatment in improving

Aβ-induced memory deficiency and cognitive dysfunction *in vivo*.

From a molecular perspective, Aβ hypothesis has long been a mainstream

approach in the study on the mechanism of AD pathogenesis and drug development

[49]. Under normal conditions, α- and γ-secretase will hydrolyze APP, preventing

production of Aβ peptides. However, once pathological changes occur, the hydrolysis

of APP is carried out by β- and γ-secretase, leading to secretion of toxic Aβ peptides

and aggregation of insoluble Aβ fibrils [3]. Thus, under the Aβ hypothesis, inhibition

of BACE1 is also considered a major therapeutic target, on top of Aβ clearance, in

AD treatment. The study exhibits the significant effect of TC-HT in decreasing the

hippocampal Aβ level, remarkably outperforming HT which fails to show any

significant Aβ clearance ability (**Fig 4A**). Besides, TC-HT also shows a significantly

better effect in inhibiting the hippocampal BACE1 level than HT, and there is no

significant difference in BACE1 level between the HT and Aβ groups (**Fig 4B**). These

results demonstrate that TC-HT can attenuate the amount of toxic Aβ fragments and

their production in the brains of AD mice, much better than the continuous HT

treatment. To the best of our knowledge, this is the first time that a physical thermal

treatment is proven to be able to modulate the expression of BACE1, shedding light

on alternative approach in AD treatment.

Apart from neurotoxic Aβ in brain, the pathogenesis of AD is also closely linked

to the activation of the brain's immune cells, particularly microglia and astrocytes,

which are triggered by Aβ deposits, according to some reports [4, 50]. Chronic

activation of the immune cells causes neuroinflammation, releasing the inflammatory

mediators which are associated with neurodegeneration and thus exacerbating AD

progression [50, 51]. Consequently, inhibition of neuroinflammation in AD brains is

also important in AD treatment. The study demonstrates extraordinary

anti-neuroinflammatory ability of TC-HT treatment. It is capable of suppressing both

Aβ-induced activation markers of microglial cells (Iba-1) and astrocytes (GFAP)

significantly, as shown in **Fig 5**. By contrast, the continuous HT treatment shows only

little inhibition on both neuroinflammatory markers without significance. Therefore,

the study proves that TC-HT is more capable of ameliorating neuroinflammation in

AD brains than continuous HT *in vivo*.

The clearance of toxic Aβ deposits, a hallmark of Alzheimer's disease, is as

important as curbing its production. First discovered as an enzyme responsible for the

degradation of insulin, IDE has also been found in recent studies to be capable of

degrading several polypeptides including Aβ [40]. In addition, it has been known that

neuron cells are able to regulate the extracellular levels of Aβ via proteolysis by IDE

in vitro [52]. Studies have also shown that the alteration of IDE plays a key role in

AD, as the decrease of IDE activity and expression could result in Aβ accumulation,

thereby raising the risk of AD development [53, 54]. In line with the finding of a

recent study that IDE can be increased in a fashion like HSP [21], our study discovers

that TC-HT treatment can significantly boost the hippocampal IDE level (**Fig 6A**), an

effect unfound in HT treatment. The more pronounced Aβ clearance effect of TC-HT

may result from its higher capability in inhibiting Aβ production (**Fig 4B**) and

clearing Aβ (**Fig 6A**) than continuous HT, indicating its higher potential in AD

therapy than the latter.

Antioxidant enzyme is also an important protective mechanism against AD. It

has been known that Aβ deposition in brain can increase the production of

mitochondrial ROS and intracellular oxidative stress, increasing Aβ formation, which

leads to a vicious cycle and exacerbates AD progression [5, 6]. Besides, excessive

ROS level could be toxic to brain neurons and decrease cognitive performance,

according to some studies [55]. SOD2, a member of the superoxide dismutase family,

can transform superoxide $O_2{}^{\cdot-}$ into hydrogen peroxide and hence regulate excessive

oxidative stress [56]. The reduction of SOD2 can accelerate the onset of

Aβ-dependent memory deficits *in vivo* [57], while the elevation of SOD2 can prevent

the memory deficits [58]. Our study shows that TC-HT can greatly recover

hippocampal SOD2 decreased by Aβ (**Fig 6B**), at a scale far outpacing that of HT.

The result is in line with the behavioral performance of mice in Y-maze and NOR

tests in **Fig 3**, indicating that TC-HT can attenuate the Aβ-induced memory deficits

via removal of excessive ROS in brains by SOD2-related mechanisms.

In addition to the neuroprotective functions of IDE and SOD2, silent mating-type

information regulation 2 homolog 1 (SIRT1) has been reported to attenuate Aβ levels

both *in vitro* and *in vivo* [59]. SIRT1 is a famous deacetylase, known for removing

acetyl groups from substrate proteins to regulate their function. The SIRT1-mediated

deacetylation can not only reduce Aβ production by suppressing the expression of

β-secretase, but also inhibit tau protein aggregation, another pathological hallmark of

AD [59, 60]. Besides, SIRT1 is capable of suppressing inflammation and activating

antioxidative response via deacetylation [61, 62]. Noteworthily, the deacetylation

activity of SIRT1 can be induced by heat [63, 64], which could deacetylate heat shock

factor 1 and lead to gene transcription of heat shock proteins such as HSP70 [64].

HSP70 is a well-known heat-induced chaperone protein that assists proper protein

folding or refold the misfolded proteins in cells [65]. In addition to its role in protein

folding, HSP70 is also capable of blocking the oligomerization of Aβ fibril via

refolding, clearing Aβ through up-regulating IDE expression, inhibiting tau protein

aggregation, and promoting tau protein degradation [66]. Hence, we believe that both

SIRT1 and HSP70 are also involved in the therapeutic effects of TC-HT against AD.

Our results show that i.c.v. Aβ injection significantly reduces hippocampal SIRT1

level to 0.76-fold of the control group, which is significantly recovered by TC-HT

treatment (0.96-fold) (**Fig 8A**). On the other hand, HSP70 expression is not affected

in mice hippocampi by Aβ injection (1.08-fold), but it is significantly elevated in the

TC-HT group (1.32-fold), as shown in **Fig 8B**. Noteworthily, the increase in HSP70

level by TC-HT treatment is consistent with the change of IDE level in **Fig 6A**,

suggesting that TC-HT-induced Aβ clearance might be mediated via

HSP70-dependent signalling.

Fig 8. Effect of TC-HT treatments on the expression levels of SIRT1 and HSP70

in hippocampi.

Representative western blots and the quantifications of (A) SIRT1 and (B) HSP70 are

presented for evaluating anti-Aβ effect in hippocampi. β-actin was used as loading

control. Data are presented as mean ± S.D. (n = 9 in each group). One-way ANOVA

with Tukey's post hoc test is used to determine statistical significance. (*P < 0.05,

comparison between indicated groups).

Excessive ROS accumulation could be detrimental to neurons, damaging

memory and cognitive ability. However, some studies have shown that proper

increase of ROS level is essential for activating the redox signalling and the

associated neuroprotective pathways [67]. Therefore, the study suggests not to

eliminate ROS induced by thermal stress, until setup of relevant protective

mechanisms. To validate the hypothesis, an *in vitro* experiment using H_2 treatment as

an ROS scavenger was introduced in this study. As shown in **Fig 7A**, the result shows

significant recovery of the Aβ-inhibited viability of SH-SY5Y cells after TC-HT

treatment, whose neuroprotective effect could be attributed to activation of Akt and its

downstream signalling pathways, as suggested in our previous study [31]. It has been

reported that early administration of antioxidants may block the construction of

neuroprotection via these pathways [68-70], which is in line with the observation that

administration of H_2 treatment less than 12 h after TC-HT fails to further augment the

viability of SH-SY5Y cells. On the other hand, our study shows that with 90 min H_2

treatment at 12~16 h after TC-HT, removal of TC-HT-induced ROS can further boost

protection of neural cells against Aβ, exhibiting a better neuroprotective effect than

TC-HT treatment alone (**Fig 7A**). Noticeably, the H_2 treatment alone did not show

any protective effect for SH-SY5Y cells against 25 μM Aβ (**Fig 7B**), indicating that

TC-HT treatment is essential for the protective mechanism. The results show that

scavenging TC-HT-induced ROS with H_2 treatment after establishment of

neuroprotective mechanism by TC-HT (12~16 h after TC-HT treatment) can further

enhance the neuroprotective effect of TC-HT treatment against Aβ. Relevant studies

have been in progress *in vivo*.

Although heat could be beneficial for AD, continuous heat exposure may hinder

the function of neuroprotective proteins. Some studies have shown that the

inactivation of proteins may start around 40°C, due to protein denaturation and

unfolding [71]. In the study, continuous HT treatment doesn't exhibit significant

effect in Aβ regulation (**Fig 4 & Fig 6A**) and anti-neuroinflammation activity (**Fig 5**),

and lags behind TC-HT treatment in antioxidative activity (**Fig 6B**). However, with

identical heat exposure time, TC-HT featuring intermittent high-temperature

treatment attains much greater improvement than continuous HT. As a result, despite

its contribution to brain's neuroprotective mechanism, HT treatment may entail

overheating causing protein dysfunction in neural cells and thus interrupting the

pathways for neuroprotection. Therefore, the interspersing low-temperature process is

crucial for TC-HT treatment's neuroprotective effect. Our study demonstrates that

TC-HT can be an effective thermal treatment for AD. In practice, focused ultrasound

(FUS) may be a suitable heating source for TC-HT treatment, as it is capable of

precisely modulating the temperature at the targeted area, via computer-controlled

programmable procedure. The FUS heating area can be set at 4-8 mm [72, 73],

smaller than human brain, thereby quickening heat dissipation to surrounding tissues

and preventing hyperthermic damage. In operation, multi-focal FUS can control the

heating area and temperature by changing the size of scanned spot and intensity in

scanning mode, thereby attaining set temperature and effective thermal dissipation for

TC-HT in AD treatment.

In summary, our study puts forth a novel hyperthermic method, TC-HT, for

application as a potential AD treatment *in vivo*, which demonstrates a better effect in

improving the cognitive performance of mice in Y-maze and NOR tests than

conventional HT treatment, which heats the subjects continuously. Furthermore, the

study shows that TC-HT can suppress the hippocampal Aβ expression to a normal

level, significantly outperforming HT which only exhibits a slight inhibition effect,

perhaps due to TC-HT's higher capability in inhibiting the Aβ productive activity of

BACE1 protein and promoting the Aβ clearance effect of IDE protein. In addition,

TC-HT can significantly inhibit the Aβ-induced overexpression of neuroinflammatory

markers Iba-1 and GFAP in hippocampi, while HT treatment only slightly attenuates

their expression levels without significance. Moreover, TC-HT raises the expression

level of antioxidative protein SOD2 at an extent much higher than HT, indicating that

it can provide a better protection for hippocampal neurons against the i.c.v.-injected

Aβ. Overall, the study validates the therapeutic potential of TC-HT for AD *in vivo*,

shedding light on development of alternative AD treatment with high efficacy.

Funding

This work was supported by grants from Ministry of Science and Technology (MOST

110-2112-M-002-004,   MOST   109-2112-M-002-004,   and   MOST

108-2112-M-002-016 to CYC) of the Republic of China. The funders had no role in

study design, data collection and analysis, decision to publish, or preparation of the

manuscript.

Competing Interests

600   The authors have declared that no competing interests exist.

Ethics Approval

603   This study was approved by the Institutional Animal Care and Use Committee of

604   National Taiwan University. The experiments were conducted with Guide for the

605   Care and Use of Laboratory Animals being strictly followed.

Acknowledgments

608   The authors would like to acknowledge the service provided by Animal Resource

609   Center of National Taiwan University for facility support.

Author Contributions

Conceptualization: Chih-Yu Chao.

Data Curation: Yu-Yi Kuo, Chih-Yu Chao.

Formal analysis: Yu-Yi Kuo, Wei-Ting Chen, Guan-Bo Lin, You-Ming Chen, Hsu-Hsiang Liu, Chih-Yu Chao.

Funding acquisition: Chih-Yu Chao.

Investigation: Yu-Yi Kuo, Wei-Ting Chen, Guan-Bo Lin, Chih-Yu Chao.

Project Administration: Chih-Yu Chao.

Supervision: Chih-Yu Chao.

Validation: Yu-Yi Kuo, Wei-Ting Chen, Guan-Bo Lin, You-Ming Chen, Hsu-Hsiang Liu

Writing – original draft: Yu-Yi Kuo, Chih-Yu Chao.

Writing – review & editing: Chih-Yu Chao.

References

1. 2020 Alzheimer's disease facts and figures. Alzheimer's Dement. 2020; 16(3):391-460.

2. Chen GF, Xu TH, Yan Y, Zhou YR, Jiang Y, Melcher K, et al. Amyloid beta:
structure, biology and structure-based therapeutic development. Acta
Pharmacol Sin. 2017; 38(9):1205-35.

3. Sun X, Chen WD, Wang YD. β-amyloid: the key peptide in the pathogenesis
of Alzheimer's disease. Front Pharmacol. 2015; 6:221.

4. Essa MM, Subash S, Akbar M, Al-Adawi S, Guillemin GJ. Long-term dietary
supplementation of pomegranates, figs and dates alleviate neuroinflammation
in a transgenic mouse model of Alzheimer's disease. PLoS One. 2015;
10(3):e0120964.

5. García-Escudero V, Martín-Maestro P, Perry G, Avila J. Deconstructing
mitochondrial dysfunction in Alzheimer disease. Oxid Med Cell Longev. 2013;
2013:162152.

6. Leuner K, Schütt T, Kurz C, Eckert SH, Schiller C, Occhipinti A, et al.
Mitochondrion-derived reactive oxygen species lead to enhanced amyloid beta
formation. Antioxid Redox Signal. 2012; 16(12):1421-33.

7. Palop JJ, Mucke L. Amyloid-β-induced neuronal dysfunction in Alzheimer's
disease: from synapses toward neural networks. Nat Neurosci. 2010;
13(7):812-8.

8. Kashyap G, Bapat D, Das D, Gowaikar R, Amritkar RE, Rangarajan G, et al.
Synapse loss and progress of Alzheimer's disease - a network model. Sci Rep.
2019; 9(1):6555.

9. Karran E, Mercken M, De Strooper B. The amyloid cascade hypothesis for
Alzheimer's disease: an appraisal for the development of therapeutics. Nat Rev
Drug Discov. 2011; 10(9):698-712.

10. Cao J, Hou J, Ping J, Cai D. Advances in developing novel therapeutic
strategies for Alzheimer's disease. Mol Neurodegener. 2018; 13(1):64.

11. Du Z, Li M, Ren J, Qu X. Current strategies for modulating Aβ aggregation
with multifunctional agents. Acc Chem Res. 2021; 54(9):2172-84.

12. Lee D, Lee G, Yoon DS. Anti-Aβ drug candidates in clinical trials and
plasmonic nanoparticle-based drug-screen for Alzheimer's disease. Analyst.
2018; 143(10):2204-12.

13. Xin SH, Tan L, Cao X, Yu JT, Tan L. Clearance of amyloid beta and tau in
Alzheimer's disease: from mechanisms to therapy. Neurotox Res. 2018;
34(3):733-48.

14. Hung SY, Fu WM. Drug candidates in clinical trials for Alzheimer's disease. J
Biomed Sci. 2017; 24(1):47.

15. Habash RW, Bansal R, Krewski D, Alhafid HT. Thermal therapy, part 1: an
introduction to thermal therapy. Crit Rev Biomed Eng. 2006; 34(6):459-89.

16. Manning MR, Cetas TC, Miller RC, Oleson JR, Connor WG, Gerner EW.
Clinical hyperthermia: results of a phase I trial employing hyperthermia alone
or in combination with external beam or interstitial radiotherapy. Cancer. 1982;
49(2):205-16.

17. Falk MH, Issels RD. Hyperthermia in oncology. Int J Hyperthermia. 2001;
17(1):1-18.

18. Bechtold DA, Brown IR. Heat shock proteins Hsp27 and Hsp32 localize to
synaptic sites in the rat cerebellum following hyperthermia. Brain Res Mol
Brain Res. 2000; 75(2):309-20.

19. Mailhos C, Howard MK, Latchman DS. Heat shock proteins hsp90 and hsp70
protect neuronal cells from thermal stress but not from programmed cell death.
J Neurochem. 1994; 63(5):1787-95.

20. Hatayama T, Yamagishi N, Minobe E, Sakai K. Role of hsp105 in protection
against stress-induced apoptosis in neuronal PC12 cells. Biochem Biophys
Res Commun. 2001; 288(3):528-34.

21. Tundo GR, Sbardella D, Ciaccio C, Bianculli A, Orlandi A, Desimio MG, et al.
Insulin-degrading enzyme (IDE): a novel heat shock-like protein. J Biol Chem.
2013; 288(4):2281-9.

22. Ahmed K, Zaidi SF, Mati-Ur-Rehman, Rehman R, Kondo T. Hyperthermia
and protein homeostasis: cytoprotection and cell death. J Therm Biol. 2020;
91:102615.

23. Banerjee Mustafi S, Chakraborty PK, Dey RS, Raha S. Heat stress upregulates
chaperone heat shock protein 70 and antioxidant manganese superoxide
dismutase through reactive oxygen species (ROS), p38MAPK, and Akt. Cell
Stress Chaperones. 2009; 14(6):579-89.

24. Hashemi M, Abbasiazam A, Oraee-Yazdani S, Lenzer J. Response of human
glioblastoma cells to hyperthermia: cellular apoptosis and molecular events.
Tissue Cell. 2022; 75:101751.

25. Lee JS, Choi JC, Kang SY, Kang JH, Park JK. Heat stroke: increased signal
intensity in the bilateral cerebellar dentate nuclei and splenium on
diffusion-weighted MR imaging. AJNR Am J Neuroradiol. 2009; 30(4):e58.

26. Tabassum S, Raza N, Shah SZ. Outcome of heat stroke patients referred to a
tertiary hospital in Pakistan: a retrospective study. East Mediterr Health J.
2019; 25(7):457-64.

27. Cremer OL, Klakman CJ. Cerebral pathophysiology and clinical neurology of
hyperthermia in humans. Prog Brain Res. 2007; 162:153-69.

28. Laukkanen T, Kunutsor S, Kauhanen J, Laukkanen JA. Sauna bathing is
inversely associated with dementia and Alzheimer's disease in middle-aged
Finnish men. Age Ageing. 2017; 46(2):245-9.

29. Lu CH, Chen WT, Hsieh CH, Kuo YY, Chao CY. Thermal
cycling-hyperthermia in combination with polyphenols, epigallocatechin
gallate and chlorogenic acid, exerts synergistic anticancer effect against
human pancreatic cancer PANC-1 cells. PLoS One. 2019; 14(5):e0217676.

30. Chen WT, Sun YK, Lu CH, Chao CY. Thermal cycling as a novel thermal
therapy to synergistically enhance the anticancer effect of propolis on
PANC-1 cells. Int J Oncol. 2019; 55(3):617-28.

31. Chen WT, Kuo YY, Lin GB, Lu CH, Hsu HP, Sun YK, et al. Thermal cycling
protects SH-SY5Y cells against hydrogen peroxide and β-amyloid-induced
cell injury through stress response mechanisms involving Akt pathway. PLoS
One. 2020; 15(10):e0240022.

32. Kim HY, Lee DK, Chung BR, Kim HV, Kim Y. Intracerebroventricular
injection of amyloid-β peptides in normal mice to acutely induce
Alzheimer-like cognitive deficits. J Vis Exp. 2016; (109):53308.

33. Al Faruque H, Choi ES, Lee HR, Kim JH, Park S, Kim E. Targeted removal of
leukemia cells from the circulating system by whole-body magnetic
hyperthermia in mice. Nanoscale. 2020; 12(4):2773-86.

34. Lin C, Liu Z, Li Y, Chen J. Protocol for fever-range whole-body hyperthermia
(WBH) in mice to study febrile effect on T-cell adhesion and migration.
STAR Protoc. 2021; 2(3):100720.

35. Saim H, Salim MIM, Jemon K. Antitumor effect of fever range whole body
hyperthermia with curcumin in breast cancer-induced mice. Int J Recent
Technol Eng. 2019; 8(2):297-301.

36. Ohsawa I, Ishikawa M, Takahashi K, Watanabe M, Nishimaki K, Yamagata K,
et al. Hydrogen acts as a therapeutic antioxidant by selectively reducing
cytotoxic oxygen radicals. Nat Med. 2007; 13(6):688-94.

37. Murakami Y, Ito M, Ohsawa I. Molecular hydrogen protects against oxidative
stress-induced SH-SY5Y neuroblastoma cell death through the process of
mitohormesis. PLoS One. 2017; 12(5):e0176992.

38. Hampel H, Vassar R, De Strooper B, Hardy J, Willem M, Singh N, et al. The
β-secretase BACE1 in Alzheimer's disease. Biol Psychiatry. 2021;
89(8):745-56.

39. Huang W, Li Z, Zhao L, Zhao W. Simvastatin ameliorate memory deficits and
inflammation in clinical and mouse model of Alzheimer's disease via
modulating the expression of miR-106b. Biomed Pharmacother. 2017;
92:46-57.

40. Tundo GR, Sbardella D, Ciaccio C, Grasso G, Gioia M, Coletta A, et al.
Multiple functions of insulin-degrading enzyme: a metabolic crosslight? Crit
Rev Biochem Mol Biol. 2017; 52(5):554-82.

41. Massaad CA, Washington TM, Pautler RG, Klann E. Overexpression of
SOD-2 reduces hippocampal superoxide and prevents memory deficits in a
mouse model of Alzheimer's disease. Proc Natl Acad Sci USA. 2009;
106(32):13576-81.

42. Selkoe DJ, Hardy J. The amyloid hypothesis of Alzheimer's disease at 25
years. EMBO Mol Med. 2016; 8(6):595-608.

43. Naldi M, Fiori J, Pistolozzi M, Drake AF, Bertucci C, Wu R, et al. Amyloid
β-peptide 25-35 self-assembly and its inhibition: a model undecapeptide
system to gain atomistic and secondary structure details of the Alzheimer's
disease process and treatment. ACS Chem Neurosci. 2012; 3(11):952-62.

44. Xiao Y, Ma B, McElheny D, Parthasarathy S, Long F, Hoshi M, et al.
Aβ(1-42) fibril structure illuminates self-recognition and replication of
amyloid in Alzheimer's disease. Nat Struct Mol Biol. 2015; 22(6):499-505.

45. Jakki SL, Senthil V, Yasam VR, Chandrasekar MJN, Vijayaraghavan C. The
blood brain barrier and its role in Alzheimer's therapy: an overview. Curr
Drug Targets. 2018; 19(2):155-69.

46. Jahn H. Memory loss in Alzheimer's disease. Dialogues Clin Neurosci. 2013;
15(4):445-54.

47. Kraeuter AK, Guest PC, Sarnyai Z. The Y-maze for assessment of spatial
working and reference memory in mice. Methods Mol Biol. 2019;
1916:105-11.

48. Lueptow LM. Novel object recognition test for the investigation of learning
and memory in mice. J Vis Exp. 2017; (126):55718.

49. Qiao O, Ji H, Zhang Y, Zhang X, Zhang X, Liu N, et al. New insights in drug
development for Alzheimer's disease based on microglia function. Biomed
Pharmacother. 2021; 140:111703.

50. Ahmad MH, Fatima M, Mondal AC. Influence of microglia and astrocyte
activation in the neuroinflammatory pathogenesis of Alzheimer's disease:
rational insights for the therapeutic approaches. J Clin Neurosci. 2019;
59:6-11.

51. Heneka MT, Carson MJ, El Khoury J, Landreth GE, Brosseron F, Feinstein
DL, et al. Neuroinflammation in Alzheimer's disease. Lancet Neurol. 2015;
14(4):388-405.

52. Vekrellis K, Ye Z, Qiu WQ, Walsh D, Hartley D, Chesneau V, et al. Neurons
regulate extracellular levels of amyloid β-protein via proteolysis by
insulin-degrading enzyme. J Neurosci. 2000; 20(5):1657-65.

53. Cook DG, Leverenz JB, McMillan PJ, Kulstad JJ, Ericksen S, Roth RA, et al.
Reduced hippocampal insulin-degrading enzyme in late-onset Alzheimer's
disease is associated with the apolipoprotein E-ε4 allele. Am J Pathol. 2003;
162(1):313-9.

54. Eckman EA, Eckman CB. Aβ-degrading enzymes: modulators of Alzheimer's
disease pathogenesis and targets for therapeutic intervention. Biochem Soc
Trans. 2005; 33(5):1101-5.

55. Massaad CA, Klann E. Reactive oxygen species in the regulation of synaptic
plasticity and memory. Antioxid Redox Signal. 2011; 14(10):2013-54.

56. Kim YS, Gupta Vallur P, Phaëton R, Mythreye K, Hempel N. Insights into the
dichotomous regulation of SOD2 in cancer. Antioxidants. 2017; 6(4):86.

57. Esposito L, Raber J, Kekonius L, Yan F, Yu GQ, Bien-Ly N, et al. Reduction
in mitochondrial superoxide dismutase modulates Alzheimer's disease-like
pathology and accelerates the onset of behavioral changes in human amyloid
precursor protein transgenic mice. J Neurosci. 2006; 26(19):5167-79.

58. Tadokoro K, Ohta Y, Inufusa H, Loon AFN, Abe K. Prevention of cognitive
decline in Alzheimer's disease by novel antioxidative supplements. Int J Mol
Sci. 2020; 21(6):1974.

59. Lalla R, Donmez G. The role of sirtuins in Alzheimer's disease. Front Aging
Neurosci. 2013; 5:16.

60. Xu J, Jackson CW, Khoury N, Escobar I, Perez-Pinzon MA. Brain SIRT1
mediates metabolic homeostasis and neuroprotection. Front Endocrinol. 2018;
9:702.

61. Yang H, Zhang W, Pan H, Feldser HG, Lainez E, Miller C, et al. SIRT1
activators suppress inflammatory responses through promotion of p65
deacetylation and inhibition of NF-κB activity. PLoS One. 2012;7(9):e46364.

62. Singh CK, Chhabra G, Ndiaye MA, Garcia-Peterson LM, Mack NJ, Ahmad N.
The role of sirtuins in antioxidant and redox signaling. Antioxid Redox Signal.
2018; 28(8):643-61.

63. Chakraborty S, Singh M, Pandita RK, Singh V, Lo CSC, Leonard F, et al.
Heat-induced SIRT1-mediated H4K16ac deacetylation impairs resection and
SMARCAD1 recruitment to double strand breaks. iScience. 2022;
25(4):104142.

64. Raynes R, Brunquell J, Westerheide SD. Stress inducibility of SIRT1 and its
role in cytoprotection and cancer. Genes Cancer. 2013; 4(3-4):172-82.

65. Mayer MP, Bukau B. Hsp70 chaperones: cellular functions and molecular
mechanism. Cell Mol Life Sci. 2005; 62(6):670-84.

66. Lu RC, Tan MS, Wang H, Xie AM, Yu JT, Tan L. Heat shock protein 70 in
Alzheimer's disease. Biomed Res Int. 2014; 2014:435203.

67. Milkovic L, Cipak Gasparovic A, Cindric M, Mouthuy PA, Zarkovic N. Short
overview of ROS as cell function regulators and their implications in therapy
concepts. Cells. 2019; 8(8):793.

68. Unoki T, Abiko Y, Toyama T, Uehara T, Tsuboi K, Nishida M, et al.
Methylmercury, an environmental electrophile capable of activation and
disruption of the Akt/CREB/Bcl-2 signal transduction pathway in SH-SY5Y
cells. Sci Rep. 2016; 6(1):28944.

69. Jin Y, Wang H. Naringenin inhibit the hydrogen peroxide-induced SH-SY5Y
cells injury through Nrf2/HO-1 pathway. Neurotox Res. 2019; 36(4):796-805.

70. Sun P, Nie X, Chen X, Yin L, Luo J, Sun L, et al. Nrf2 signaling elicits a
neuroprotective role against PFOS-mediated oxidative damage and apoptosis.
Neurochem Res. 2018; 43(12):2446-59.

71. Lepock JR. Protein denaturation during heat shock. Adv Mol Cell Biol. 1997;
19:223-59.

72. Chaplin V, Caskey CF. Multi-focal HIFU reduces cavitation in
mild-hyperthermia. J Ther Ultrasound. 2017; 5(1):12.

73. Partanen A, Tillander M, Yarmolenko PS, Wood BJ, Dreher MR, Kohler MO.
Reduction of peak acoustic pressure and shaping of heated region by use of
multifoci sonications in MR-guided high-intensity focused ultrasound
mediated mild hyperthermia. Med Phys. 2013; 40(1):013301.

Figure 1

Figure 2

Figure 3

Figure 4

A

Aβ   46 kDa

β-actin   45 kDa

B

BACE1   70 kDa

β-actin   45 kDa

Figure 5

A

Iba-1 — 17 kDa

β-actin — 45 kDa

N.S.

Iba-1 / β-actin
(Fold relative to control)

Ctrl   Aβ   TC-HT   HT

B

GFAP — 50 kDa

β-actin — 45 kDa

N.S.

GFAP / β-actin
(Fold relative to control)

Ctrl   Aβ   TC-HT   HT

Figure 6

Figure 7

Figure 8