Thermal cycling-hyperthermia ameliorates cognitive impairment of intracerebroventricular Aβ$_{25\text{-}35}$-induced Alzheimer's disease in C57BL/6 mice

Yu-Yi Kuo[1,2], Wei-Ting Chen[1,2], Guan-Bo Lin[1,2], You-Ming Chen[2,3], Hsu-Hsiang Liu[2,3], and Chih-Yu Chao[1,2,3,*]

[1] Department of Physics, Lab for Medical Physics & Biomedical Engineering, National Taiwan University, Taipei 10617, Taiwan

[2] Molecular Imaging Center, National Taiwan University College of Medicine, Taipei 10051, Taiwan

[3] Graduate Institute of Applied Physics, Biophysics Division, National Taiwan University, Taipei 10617, Taiwan

* Corresponding author

E-mail: cychao@phys.ntu.edu.tw (CYC)

Abstract

Despite continuation of some controversies, Alzheimer's disease (AD), the most common cause of dementia nowadays, has been widely believed to derive mainly from

excessive β-amyloid (Aβ) aggregation, that would increase reactive oxygen species

(ROS) and induce neuroinflammation, leading to neuron loss and cognitive impairment.

Existing drugs on Aβ have been ineffective or offer only temporary relief at best, due

to blood-brain barrier or severe side effects. The study employed thermal cycling-

hyperthermia (TC-HT) as an alternative AD therapy and compared its effect with

continuous hyperthermia (HT) *in vivo*. It established an AD mice model via

intracerebroventricular (i.c.v.) injection of $A\beta_{25-35}$, proving that TC-HT is much more

effective in alleviating its performance decline in Y-maze and NOR test, in comparison

with HT. In addition, TC-HT also exhibits a better performance in decreasing the

hippocampal Aβ and BACE1 expressions as well as the neuroinflammation markers

Iba-1 and GFAP levels. Furthermore, the study finds that TC-HT can elevate more

protein expressions of IDE and antioxidative enzyme SOD2 than HT. Besides, after

establishment of neuroprotective mechanism, removal of TC-HT-induced ROS can

further augment protection of neural cells against Aβ. In sum, the study proves the

potential of TC-HT in AD treatment, which can be put into clinical application with the

use of high-intensity focused ultrasound (HIFU).

Introduction

Alzheimer's disease (AD) has been the most common cause of dementia nowadays, accounting for over 60% of all the cases [1]. One of the most notable hallmarks of AD pathogenesis is the accumulation in brain of toxic β-amyloid (Aβ) peptides [2], which are cleaved from the amyloid precursor protein (APP) by β- and γ-secretase [3]. Such abnormal Aβ deposits can activate neuroinflammation [4], generate reactive oxygen species (ROS) [5, 6], and cause neuron dysfunction and synapse loss, leading to memory loss and cognitive impairment [7, 8]. Therefore, Aβ appears to be the foremost target in AD therapy [9].

Currently, drug therapy is the mainstream AD treatment, aiming to suppress the generation [10], disrupt the aggregation [11], and promote the clearance [12] of Aβ. Drugs on proteases, such as insulin degrading enzyme (IDE), for alleviating Aβ degradation also have significant potential in AD treatment [13]. However, at present AD drugs have high failure rates, due to difficulty penetrating blood-brain barrier, on top of severe side effects, such as liver toxicity, strong autoimmune response, or carcinogenicity [14]. Hence, there is a need for a safe and effective alternative AD therapy.

Heat has long been regarded as an alternative therapy with significant potential,

given its proven therapeutic effect in physiotherapy, ophthalmology, and cardiology

[15]. Hyperthermia (HT), for instance, has been applied effectively in cancer treatment,

either for directly inhibiting tumor growth [16] or supplementing radiotherapy and

certain chemotherapies to augment their efficacy [17]. In addition, HT has also been

reported to be capable of increasing the expression of heat shock protein family [18],

some of whose members can suppress the formation of Aβ fibril and Aβ toxicity [19,

20]. Besides, HT treatment is conducive to increase the levels of IDE [21] and

antioxidant proteins, such as nuclear factor erythroid 2–related factor 2 (Nrf2) and

superoxide dismutase 2 (SOD2) [22, 23].

However, continuous high-temperature exposure can be thermotoxic to human

neural cells, according to some reports [24]. Heat stroke, featuring abnormally high

core body temperature (> 40°C), may cause cerebellar injury and memory deficits [25,

26]. Whole-body HT treatment has also been reported to cause adverse effect on

memory or neural system, such as decreased cognitive performance, impaired pressure-

flow autoregulation, brain edema, intracerebral hemorrhage, and transient central

nervous system dysfunction [27, 28]. Moreover, although a study showed that long-

term heat exposure could reduce the risk of memory loss among middle-aged men, it

was time-consuming without certain curative effect [29]. Therefore, it is imperative to

find a tolerable and effective thermal treatment for AD.

Our previous study put forth a novel thermal treatment, thermal-cycling

hyperthermia (TC-HT), demonstrating its therapeutic effect in anticancer treatment [30,

31] and neuroprotection [32] in vitro, with minimal side effects. The study applied TC-

HT to treat mice with AD, induced by intracerebroventricular (i.c.v.) injection of $A\beta_{25\text{-}35}$,

for comparison with the therapeutic effect of HT. In both cases, heat was applied to

mouse brain specifically, to avoid potential side effect of whole-body HT.

In this paper, behavioral tests show that TC-HT significantly improves $A\beta$-

induced memory impairment, with an effect much better than HT. In addition, TC-HT

also exhibits a much better performance than HT in decreasing $A\beta$ and β-secretase

(BACE1) expressions, suppressing neuroinflammation markers ionized calcium-

binding adapter molecule 1 (Iba-1) and glial fibrillary acidic protein (GFAP) levels,

and increasing the expression levels of IDE and SOD2 enzymes. Furthermore, the study

shows that removal of TC-HT-induced ROS can further augment the protective effect

of TC-HT in an in vitro neural model. In sum, these results underscore the potential of

TC-HT as an alternative AD therapy.

Materials and Methods

Ethics Statement

All animal experiments were conducted according to the regulations approved by

the Animal Ethical Committee of National Taiwan University. Mice were monitored

every day for signs of distress. The humane endpoints of the mice included decreased

socialization, anorexia, weight loss of 15%, hunching, or the inability to evade handling.

All hydrodynamic injections and thermal treatments were conducted under anesthesia

with 5% isoflurane (Panion & BF Biotech Inc., Taipei, Taiwan), and all efforts were

made to minimize suffering. For sacrifice, mice were humanely euthanized via carbon

dioxide (CO_2) gas asphyxiation followed by cervical dislocation and were considered

non-survivors.

Animals

C57BL/6 male mice (6-8 weeks old, 22-26 g) were obtained from National

Laboratory Animal Center (Taipei, Taiwan). All mice were housed in groups of 6 mice

per cage (32 cm × 16 cm × 14 cm) and maintained in a specific-pathogen-free (SPF)

room under controlled temperature (22±2 °C) and relative humidity (55±10%) in

Animal Resource Center of National Taiwan University. All mice were subjected to a

12:12 h light/dark cycle (lights on at 7:00 a.m.) with free access to food and water. All

experiments were conducted with mice at age 6-8 weeks. All mice were randomly

categorized into the control group, Aβ group, TC-HT group, and HT group.

β-amyloid administration

Aβ$_{25-35}$ (Sigma-Aldrich; Merck KGaA, Darmstadt, Germany) was dissolved in

0.9% sterile saline solution to a concentration of 1 mg/mL and then incubated at 37°C

for 7 days for fibril aggregation. At the beginning of the experiment (Day 0), 10 µL Aβ

solution was i.c.v.-injected into the mice's brains of the Aβ, TC-HT, and HT groups

[33]. 10 µL of sterile saline solution was i.c.v.-injected into the mice's brains of the

sham-operated control group

HT and TC-HT treatments

To focus on the thermal effect on mice's brain and minimize the potential interference outside of brains by the conventional whole-body hyperthermia methods [34-36], we adopted a modified method to heat mice's heads locally. As shown in **Fig 1A**, both HT and TC-HT thermal treatments were applied by a temperature controlling and heating system equipped with a DC power supply (9303D, ABM Industries Inc., NY, USA), a temperature/process controller (PZ900, RKC instrument Inc., Tokyo, Japan), and a 19 mm × 19 mm polyimide thermofoil for heating (HK5578R4.6L12A, Minco Products Inc., Fridley, MN, USA). Mice were anesthetized with 5% isoflurane during thermal treatments. Mice's head fur was shaved and the thermofoil was taped onto the bare skin with a thin layer of vaseline for better attachment. The thermocouple measuring tip (K-type) was located under the central region of the thermalfoil, as shown in **Fig 1B**. The actual temperatures of mice's heads were measured via the K-type thermocouple and monitored by the temperature/process controller. For TC-HT treatment, the temperature/process controller was set to implement a 10-cycle repeated procedure, in which each cycle is consisted of a desired high-temperature stage for a period of time, followed by a natural thermal dissipating stage for 30 sec at low-

temperature setting 37 °C (**Fig 1C**). In the study, the high-temperature stage in TC-HT

treatment was held at 42 °C for 6 min with a 0.5 °C tolerance. For HT treatment, the

temperature was kept at 42 ± 0.5 °C continuously for 1 h. Monitored temperatures of

mice's heads by HT and TC-HT thermal treatments were shown in **Fig 1D**.

Fig 1. Experimental setup and design of TC-HT.

(A) The setup of the temperature controlling and heating system, containing a DC

power supply, a temperature/process controller, and a thermofoil for heating. (B) The

thermal treatment was locally performed on mice's heads by thermofoil, and their

temperatures were measured via a K-type thermocouple, whose measuring tip was

placed in the central circled region under the thermofoil. (C) Schematic diagram of the

TC-HT temperature-controlled programs with parameter settings, and (D) the

temperatures of mice's heads by HT and TC-HT thermal treatments were recorded

every 10 sec throughout the heating period.

Experimental design

At the beginning of the experiment (Day 0), 10 μL Aβ solution was i.c.v.-injected

into the mice's brains of the Aβ, TC-HT, and HT groups. On day 4, 8, and 12, the

respective thermal treatment was applied to the mice of the TC-HT and HT groups. Y-

maze and NOR tests were assessed on day 13 and day 14, respectively. Brain tissues

were collected after the Y-maze test for the following western blot analyses. The

experimental schedule was shown in **Fig 2**.

Fig 2. The experimental schedule.

Aβ solution was i.c.v.-injected at day 0, and the respective thermal treatment was

implemented at day 4, 8, and 12. Behavioral tests of NOR and Y-maze were carried out

at day 13 and 14, respectively. After the Y-maze test, mice were weighted and then

euthanized to collect brain tissues for the following western blot analyses.

Y-maze test

The Y-maze test was used to assess the spatial working memory of mice. The

apparatus is consisted of three identical arms (40 cm × 13 cm × 6.5 cm) interconnected

at a central junction at 120° each. The assessed mouse was placed at the end of the start

arm, and was allowed to explore all three arms freely for 8 min. The behavioral

performance was recorded, and the spontaneous index was calculated as follows:

(number of correct alternations) / (total number of arm entries – 2) × 100. In addition,

the total number of arm entries in the Y-maze was used as a measure to characterize

locomotor activity.

Novel object recognition (NOR) test

We used the NOR test to assess the recognition memory of animals. The NOR test

consists of a familiarization phase and a test phase. In the familiarization phase, the

assessed mouse was placed in a 40 cm × 40 cm × 30.5 cm box for 8 min with 2 identical

Lego blocks, which were placed close to two diagonal corners of the box. After the

familiarization phase, the mouse was returned to the home cage for 1 h and then was

placed back into the testing box for the test phase. One of the original objects was

replaced with a novel Lego block of a different shape in the test phase, and the

behavioral performance of the mouse was recorded with an overhead video camera for

8 min. Exploration was defined as sniffing or touching the object with the nose and/or

forepaws. The duration of the exploring time was calculated for the discrimination

index. A discrimination index was defined as follows: (Tn-To) / (Tn+To), where Tn

and To stand for the exploration time of the novel and original objects, respectively.

Brain tissue collection and Western blot sample preparation

Mice were euthanized by CO_2 gas. After confirming the death of mice, brains were

taken out quickly and rinsed with phosphate buffered saline (PBS) (HyClone; GE

Healthcare Life Sciences, Chicago, IL, USA) supplemented with 1% fresh protease

inhibitor cocktail (EMD Millipore, Billerica, MA, USA). Hippocampi were rapidly

dissected on an ice-cold plate, and then stored in ice-cold RIPA lysis buffer (EMD

Millipore) supplemented with 1% fresh protease inhibitor cocktail. Hippocampal

tissues were homogenized via ultrasound processing (5 sec rectangular pulse with 5 sec

interval for 3 times, amplitude 50%, by a Q125 sonicator; Qsonica, Newton, CT, USA)

and then centrifuged at 23000×g for 30 min at 4°C. The supernatants were collected

and the protein concentration was quantified by the Bradford protein assay (BioShop

Canada Inc., Burlington, Ontario, Canada). Protein samples were stored at -80°C for

subsequent western blotting assessments.

Western blot analysis

Proteins from the homogenized hippocampi were resolved by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and then electroblotted onto the polyvinylidene difluoride (PVDF) membranes (EMD Millipore). Blotted membranes were blocked with 5% skim milk in Tris-buffered saline supplemented with 0.1% Tween-20 (TBST) (both from BioShop Canada Inc.) for 1 h at room temperature (RT) and then incubated overnight with diluted primary antibodies in blocking buffer at 4 °C, followed by three rinses with TBST washing buffer. Primary antibodies employed in this study were listed as following: anti-Aβ (Santa Cruz Biotechnology, Inc., Dallas, TX, USA), anti-IDE (Abcam, Cambridge, UK), anti-BACE1, anti-SOD2, anti-Iba-1 (Cell Signaling Technology, Danvers, MA, USA), anti-GFAP, and anti-β-actin (Gentex, Irvine, CA, USA). After washing with TBST, the membranes were incubated with horseradish peroxidase-conjugated goat anti-mouse (for probing anti-Aβ primary antibody) or goat anti-rabbit (for probing the other primary antibodies) secondary antibodies (both from Jackson ImmunoResearch Laboratories, Inc., West Grove, PA, USA) for 1 h at RT. All the antibodies were diluted according to the manufacturer's instructions to their optimal concentration. The protein bands were

visualized by an enhanced chemiluminescence substrate (Advansta, San Jose, CA, USA)

and detected by an imaging system Amersham Imager 600 (GE Healthcare Life

Sciences). The images were analyzed with Image Lab software (Bio-Rad Laboratories,

Inc.). For normalization of proteins, β-actin was used as the loading control to

determine the relative folds of targeted proteins.

In vitro cell culture and hydrogen gas (H$_2$) treatment

The human neuroblastoma SH-SY5Y cells were purchased from American Type

Culture Collection (Manassas, VA, USA). The cell culture medium was MEM/F-12

mixture (HyClone; GE Healthcare Life Sciences) containing 10% fetal bovine serum

(HyClone; GE Healthcare Life Sciences) and 1% penicillin-streptomycin (Gibco Life

Technologies, Grand Island, NY, USA), supplemented with 1mM sodium pyruvate

(Sigma-Aldrich; Merck KGaA) and 0.1 mM non-essential amino acids (Invitrogen Life

Technologies, Grand Island, NY, USA). Cells were maintained in a humidified

incubator composed of 5% CO$_2$ at 37°C. Before the experiment, SH-SY5Y cells were

subcultured and seeded in 96-well plates (Thermo Fisher Scientific Inc., Waltham, MA,

USA) overnight. Cells were treated with 25 μM Aβ for 1 h and then subjected to the

TC-HT treatment for about 2 h. The TC-HT treatment was implemented by the Thermal

Cycler (Applied Biosystems; Thermo Fisher Scientific, Inc.) and the in vitro TC-HT

parameter settings were described in our previous study [32]. In the experiment, H_2-

mixed gas was composed of 75% H_2, 20% oxygen gas (O_2), and 5% CO_2 gas

(vol/vol/vol), as described elsewhere [37]. After continuous exposure to the H_2-mixed

gas, it has been observed that H_2 concentration of the culture medium in a closed acrylic

box gets saturated within 20~30 min at atmospheric pressure [38]. Upon 12~16 h after

TC-HT treatment, cultured SH-SY5Y cells were sent into an acrylic chamber of volume

about 450 mL. Mixed gas consisting of 75% H_2, 20% O_2, and 5% CO_2 (vol/vol/vol)

were well mixed and then flowed into the chamber at a rate of 400 mL/min for 90 min

under normal pressure. Upon 4 days after the Aβ administration, the 3-(4,5-

dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Sigma-Aldrich; Merck

KGaA) assay was used to analyze the cell viability of SH-SY5Y cells. In brief, the

culture medium was replaced with MTT solution (0.5 mg/mL in the MEM/F12 mixture)

and incubated at 37°C for 4 h. The formazan crystals were dissolved by equal volume

of the solubilizing buffer containing 10% sodium dodecyl sulfate (SDS) (Bioshop

Canada Inc.) solution in 0.01 N hydrochloric acid (HCl) (Echo Chemical Co. Ltd.,

Taiwan) at 37 °C overnight. The absorbance at 570 nm of each well was analyzed by

Multiskan GO microplate spectrophotometer (Thermo Fisher Scientific Inc.), and the

background absorbance at 690 nm was subtracted. The cell viability of each group was

expressed as percentage of the untreated control.

Statistical Analysis

All statistical results were shown as mean ± standard deviation (S.D.). Statistical

analysis was conducted with OriginPro 2015 software (OriginLab Corporation,

Northampton, MA, USA) using one-way analysis of variance (ANOVA), followed by

Tukey's post hoc test. P-value < 0.05 was considered to indicate a statistically

significant difference.

Results

TC-HT improves Aβ-induced recognition impairments in mice

The study employed Y-maze test and NOR test to examine the therapeutic effect

of TC-HT and HT on AD mice. In the Y-maze test, the overall percentage of each

mouse's spontaneous index was computed to show its short-term memory function. As

exhibited in **Fig 3A**, the spontaneous index of the Aβ group is inhibited significantly,

ranging from 60.3% to 44.1%, compared with the control group. TC-HT significantly

improves the memory deficiency, elevating the spontaneous index to 61.6%, notably

different from HT, which only causes an insignificant improvement (spontaneous index

46.8%). Besides, there is no significant difference in the total number of arm entries

among the four groups (**Fig 3B**), suggesting that i.c.v. Aβ injection or both thermal

treatments didn't affect the locomotor activities of mice. On the other hand, NOR test

also confirmed that TC-HT is more effective in alleviating Aβ-induced memory

impairment than HT. As shown in **Fig 3C**, along with memory decline, the

discrimination index of the Aβ group plunged from 25.1% down to -26.2%, which

bounces back to 19.2% after TC-HT treatment, significantly outperforming HT

treatment, which only raises the discrimination index to -3.1%. Thus, TC-HT exhibited

a much stronger reversing effect on cognitive impairment than HT in both Y-maze and

NOR tests.

Fig 3. Effect of TC-HT and HT treatments on mice's behavioral performances.

The recovery effects of Aβ-induced recognition impairments after TC-HT and HT

treatments are assessed via (A) the spontaneous index and (B) the total number of arm

entries in the Y-maze test, and (C) the discrimination index in the NOR test. Data are

presented as mean ± S.D. (n = 9 in each group). One-way ANOVA with Tukey's post

hoc test is used to determine statistical significance (*P < 0.05 between indicated

groups).

TC-HT down-regulates Aβ secretion and accumulation in mouse brain

The study first analyzed the accumulated Aβ levels in hippocampi, confirming the

significant effect of TC-HT treatment in repairing Aβ-induced cognitive impairment,

as it reduces the hippocampal Aβ level from 1.34-fold to 0.88-fold compared to the

control group (**Fig 4A**). HT treatment, by comparison, exhibited only slight inhibitive

effect on the hippocampal Aβ level (1.24-fold). On the other hand, it is realized that Aβ

level in mouse brain is partially attributed to the pathological changes of BACE1 [39].

In **Fig 4B**, our results show that i.c.v.-injected Aβ triggers a significant increase in

hippocampal BACE1 expression (1.30-fold) compared with the control group. The

study finds that the elevation of BACE1 expression is significantly suppressed by the

TC-HT treatment (0.89-fold), but only slightly decreased by the HT treatment (1.19-

fold). It's noteworthy that there is no significant difference in BACE1 levels between

the Aβ group and the HT group, suggesting that conventional HT treatment is

ineffective in stopping abnormal Aβ secretion in mouse brain. To the best of our

knowledge, this is the first time that a thermal treatment exhibited an inhibitory ability

on the expression level of BACE1. In sum, the study shows that TC-HT appears to be

much more effective AD treatment than HT, as it's more capable of regulating Aβ

levels actively and inhibiting BACE1 level in hippocampus.

Fig 4. Effect of TC-HT and HT treatments on the expression levels of Aβ and

BACE1 in hippocampi.

Representative western blots and the quantifications of (A) Aβ and (B) BACE1 are

presented for evaluating anti-Aβ effect in hippocampi. β-actin was used as loading

control. Data are presented as mean ± S.D. (n = 9 in each group). One-way ANOVA

with Tukey's post hoc test is used to determine statistical significance (*P < 0.05

between indicated groups).

TC-HT ameliorates Aβ-induced neuroinflammation in mouse brain

Neuroinflammation is another hallmark of AD, as the risk of cognitive impairment

is highly associated with inflammatory level [40]. The study finds that the expression

levels of both hippocampal Iba-1 and GFAP, the activation markers of microglia cells

and astrocytes, respectively, are significantly enhanced in the Aβ group, compared with

the control group (**Fig 5A** and **5B**). After the TC-HT treatment, both Iba-1 and GFAP

levels drop significantly, while the HT treatment only slightly attenuates their

expression levels. There is no significant difference between the Aβ group and the HT

group in the expressions of both Iba-1 and GFAP. The results indicate that TC-HT

treatment can better ameliorate Aβ-induced neuroinflammation than HT treatment *in*

vivo.

Fig 5. Effect of TC-HT and HT treatments on the expression levels of

neuroinflammatory markers Iba-1 and GFAP in hippocampi.

Representative western blots and the quantifications of (A) Iba-1 and (B) GFAP are

presented for evaluating anti-neuroinflammation effect in hippocampi. β-actin was used

as loading control. Data are presented as mean ± S.D. (n = 9 in each group). One-way

ANOVA with Tukey's post hoc test is used to determine statistical significance (*P <

0.05 between indicated groups).

TC-HT promotes the expressions of IDE and SOD2 proteins in mouse brain

With IDE long regarded as a major endogenous enzyme attributing to Aβ

degradation, the alteration of IDE activity and expression can cause Aβ accumulation,

thereby increasing the risk of AD [41]. **Fig 6A** shows that while Aβ injection slightly

decreases the hippocampal IDE level (0.88-fold), TC-HT treatment significantly

increases the IDE expression level in hippocampi up to 1.38-fold. By contrast, HT

treatment performs an insignificant altering effect on IDE expression (0.91-fold),

compared with the Aβ group. The findings show that TC-HT can intensify Aβ clearance

via elevating the IDE expression level in hippocampi, while HT is ineffective in

promoting the hippocampal IDE level and thus cannot contribute to the removal of

abnormal hippocampal Aβ in mice.

Antioxidant protein SOD2 also has significant potential in AD treatment, capable

of decreasing hippocampal superoxide and amyloid plaque in AD mice [42]. **Fig 6B**

shows that after i.c.v. Aβ injection, the SOD2 expression level drops significantly

(0.62-fold), which, though, recovers significantly in the TC-HT group (1.41-fold),

much higher than recovery to 0.86-fold after HT treatment. The results show that TC-

HT can activate stronger antioxidant activity in hippocampus against the Aβ-induced

oxidative stress than continuous HT *in vivo*.

Fig 6. Effect of TC-HT and HT treatments on the expression levels of

neuroprotective proteins IDE and SOD2 in hippocampi.

Representative western blots and the quantifications of (A) IDE and (B) SOD2 are

presented for evaluating Aβ clearance and antioxidant effect in hippocampi,

respectively. β-actin was used as loading control. Data are presented as mean ± S.D. (n

= 9 in each group). One-way ANOVA with Tukey's post hoc test is used to determine

statistical significance (*$P < 0.05$ between indicated groups).

H$_2$ treatment produces auxiliary neuroprotective effect against Aβ after TC-HT

treatment in vitro

To examine whether removal of excessive ROS after TC-HT treatment can

intensify the neuroprotective effect against Aβ, H$_2$ treatment as mentioned in Materials

and Methods section was introduced as an exogenous ROS scavenger in an in vitro

model. **Fig 7A** shows the significant suppression of the viability of SH-SY5Y cells to

60.9% under the insult of 25 μM Aβ, which recovers greatly to 82.5% after TC-HT

treatment. Moreover, we find that with a 90 min H_2 treatment at 12~16 h after TC-HT

treatment, the cell viability can bounce back to a near-normal level (96.0%), exhibiting

a better neuroprotective effect than TC-HT treatment alone. However, it should be

noted that administration of H_2 treatment less than 12 h after TC-HT treatment fails to

further boost the viability of SH-SY5Y cells. On the other hand, H_2 treatment alone

does not show any significant protective effect on SH-SY5Y cells against Aβ (**Fig 7B**).

These findings show that while removal of ROS via H_2 treatment alone has no effect

against Aβ, it can further enhance the neuroprotective effect of TC-HT treatment

against Aβ, exhibiting a better neuroprotective effect than TC-HT treatment alone. This

could be attributed to removal of TC-HT-induced ROS with H_2 treatment after

establishment of neuroprotective mechanism at 12~16 h after TC-HT treatment. These

results shed light on the approach in developing AD treatment.

Fig 7. H₂ treatment as a ROS scavenger helps to promote further the

neuroprotective effect of TC-HT against Aβ in vitro.

(A) Cell viabilities of SH-SY5Y cells treated with Aβ, Aβ + TC-HT, and Aβ + TC-HT

+ H₂ treatment, respectively. The data show that H₂-assisted TC-HT treatment exhibits

a better neuroprotective effect than TC-HT treatment alone. In the other experiment,

SH-SY5Y cells were subjected directly to H₂ treatment after the administration of 25

μM Aβ. Cell viabilities of SH-SY5Y cells treated with Aβ and Aβ + H₂ treatment alone

are respectively shown in (B). The result indicates that H₂ treatment alone has no effect

against Aβ. Data are presented as mean ± S.D. (n = 6 in each group). One-way ANOVA

with Tukey's post hoc test is used to determine statistical significance (*P < 0.05

between indicated groups).

Discussion

Given prevalence of AD in our society in recent years, many scientists have taken

part in research on its treatment with various approaches, including Aβ hypothesis, one

of the most noticeable pathogenesis of AD [43]. Aβ peptides could be secreted in brains,

but their abnormal depositions, especially in hippocampus, could induce severe

neurodegeneration and synapse loss, thereby impairing memory and cognitive abilities

[7, 8]. The study employed $A\beta_{25-35}$, peptide fragment of $A\beta_{1-42}$ considered the most

pathogenic $A\beta$ species in AD [44, 45], as the model. Development of AD drugs has

been confronted with two major obstacles, namely blood-brain barrier (BBB) and

severe side effects [14, 46]. Hence, the application of heat has potential as an alternative

AD treatment, due to its capability to intensify neuroprotective effect [32] and sidestep

BBB, targeting the diseased brain directly. The study embraced TC-HT, a modified

heat treatment with higher safety and superior protective effect than HT [32], to

investigate the therapeutic potential of thermal treatment on AD *in vivo*.

With memory loss being the first and main symptom of AD [47], AD studies with

mice as model have often employed Y-maze and NOR tests to assess short-term

memory and learning memory [48, 49]. The study probed the therapeutic effect of HT

and TC-HT treatment on $A\beta$-induced memory impairment, finding that TC-HT

treatment can significantly improve the performance of spontaneous index (**Fig 3A**)

and help memory-impaired mice recognize and remember new objects (**Fig 3C**). The

study demonstrates significant effect of TC-HT treatment in recovering spatial working

memory and recognition memory impaired by $A\beta$. In comparison, although HT

treatment also enhances behavioral performance, the extent of improvement

significantly lags behind that of TC-HT treatment in both tests. Collectively, TC-HT

treatment exhibits a higher therapeutic potential than HT treatment in improving Aβ-

induced memory deficiency and cognitive dysfunction *in vivo*.

From a molecular perspective, Aβ hypothesis has long been a mainstream

approach in the study on the mechanism of AD pathogenesis and drug development

[50]. Under normal conditions, α- and γ-secretase will hydrolyze APP, preventing

production of Aβ peptides. However, once pathological changes occur, the hydrolysis

of APP is carried out by β- and γ-secretase, leading to secretion of toxic Aβ peptides

and aggregation of insoluble Aβ fibrils [3]. Thus, under the Aβ hypothesis, inhibition

of BACE1 is also considered a major therapeutic target, on top of Aβ clearance, in AD

treatment. The study exhibits the significant effect of TC-HT in decreasing the

hippocampal Aβ level, remarkably outperforming HT which fails to show any

significant Aβ clearance ability (**Fig 4A**). Besides, TC-HT also shows a significantly

better effect in inhibiting the hippocampal BACE1 level than HT, and there is no

significant difference in BACE1 level between the HT and Aβ groups (**Fig 4B**). These

results demonstrate that TC-HT can attenuate the amount of toxic Aβ fragments and

their production in the brains of AD mice, much better than the continuous HT

treatment. To the best of our knowledge, this is the first time that a physical thermal

treatment is proven to be able to modulate the expression of BACE1, shedding light on

alternative approach in AD treatment.

Apart from neurotoxic Aβ in brain, the pathogenesis of AD is also closely linked

to the activation of the brain's immune cells, particularly microglia and astrocytes,

which are triggered by Aβ deposits, according to some reports [4, 51]. Chronic

activation of the immune cells causes neuroinflammation, releasing the inflammatory

mediators which are associated with neurodegeneration and thus exacerbating AD

progression [51, 52]. Consequently, inhibition of neuroinflammation in AD brains is

also important in AD treatment. The study demonstrates extraordinary anti-

neuroinflammatory ability of TC-HT treatment. It is capable of suppressing both Aβ-

induced activation markers of microglial cells (Iba-1) and astrocytes (GFAP)

significantly, as shown in **Fig 5**. By contrast, the continuous HT treatment shows only

little inhibition on both neuroinflammatory markers without significance. Therefore,

the study proves that TC-HT is more capable of ameliorating neuroinflammation in AD

brains than continuous HT *in vivo*.

The clearance of toxic Aβ deposits, a hallmark of Alzheimer's disease, is as

important as curbing its production. First discovered as an enzyme responsible for the

degradation of insulin, IDE has also been found in recent studies to be capable of

degrading several polypeptides including Aβ [41]. In addition, it has been known that

neuron cells are able to regulate the extracellular levels of Aβ via proteolysis by IDE in

vitro [53]. Studies have also shown that the alteration of IDE plays a key role in AD,

as the decrease of IDE activity and expression could result in Aβ accumulation, thereby

raising the risk of AD development [54, 55]. In line with the finding of a recent study

that IDE can be increased in a fashion like heat shock protein (HSP) [21], our study

discovers that TC-HT treatment can significantly boost the hippocampal IDE level (**Fig

6A**), an effect unfound in HT treatment. The more pronounced Aβ clearance effect of

TC-HT may result from its higher capability in inhibiting Aβ production (**Fig 4B**) and

clearing Aβ (**Fig 6A**) than continuous HT, indicating its higher potential in AD therapy

than the latter.

Antioxidant enzyme is also an important protective mechanism against AD. It has

been known that Aβ deposition in brain can increase the production of mitochondrial

ROS and intracellular oxidative stress, increasing Aβ formation, which leads to a

vicious cycle and exacerbates AD progression [5, 6]. Besides, excessive ROS level

could be toxic to brain neurons and decrease cognitive performance, according to some

studies [56]. SOD2, a member of the superoxide dismutase family, can transform

superoxide $O_2^{\cdot-}$ into hydrogen peroxide and hence regulate excessive oxidative stress

[57]. The reduction of SOD2 can accelerate the onset of Aβ-dependent memory deficits

in vivo [58], while the elevation of SOD2 can prevent the memory deficits [59]. Our

study shows that TC-HT can greatly recover hippocampal SOD2 decreased by Aβ (**Fig**

6B), at a scale far outpacing that of HT. The result is in line with the behavioral

performance of mice in Y-maze and NOR tests in **Fig 3**, indicating that TC-HT can

attenuate the Aβ-induced memory deficits via removal of excessive ROS in brains by

SOD2-related mechanisms.

Excessive ROS accumulation could be detrimental to neurons, damaging memory

and cognitive ability. However, some studies have shown that proper increase of ROS

level is essential for activating the redox signalling and the associated neuroprotective

pathways [60]. Therefore, the study suggests not to eliminate ROS induced by thermal

stress, until setup of relevant protective mechanisms. To validate the hypothesis, an in

vitro experiment using H_2 treatment as an ROS scavenger was introduced in this study.

As shown in **Fig 7A**, the result shows significant recovery of the Aβ-inhibited viability

of SH-SY5Y cells after TC-HT treatment, whose neuroprotective effect could be

attributed to activation of Akt and its downstream signalling pathways, as suggested in

our previous study [32]. It has been reported that early administration of antioxidants

may block the construction of neuroprotection via these pathways [61-63], which is in

line with the observation that administration of H_2 treatment less than 12 h after TC-

HT fails to further augment the viability of SH-SY5Y cells. On the other hand, our

study shows that with 90 min H_2 treatment at 12~16 h after TC-HT, removal of TC-

HT-induced ROS can further boost protection of neural cells against Aβ, exhibiting a

better neuroprotective effect than TC-HT treatment alone (**Fig 7A**). Noticeably, the H_2

treatment alone did not show any protective effect for SH-SY5Y cells against 25 μM

Aβ (**Fig 7B**), indicating that TC-HT treatment is essential for the protective mechanism.

The results show that scavenging TC-HT-induced ROS with H_2 treatment after

establishment of neuroprotective mechanism by TC-HT (12~16 h after TC-HT

treatment) can further enhance the neuroprotective effect of TC-HT treatment against

Aβ. Relevant studies have been in progress *in vivo*.

In addition to the inhibitive effect of ROS on viability, continuous heat exposure

may hinder the function of neuroprotective proteins. Some studies have shown that the

inactivation of proteins may start around 40°C, due to protein denaturation and

unfolding [64]. In the study, continuous HT treatment doesn't exhibit significant effect

in Aβ regulation (**Fig 4 & Fig 6A**) and anti-neuroinflammation activity (**Fig 5**), and

lags behind TC-HT treatment in antioxidative activity (**Fig 6B**). However, with

identical heat exposure time, TC-HT featuring intermittent high-temperature treatment

attains much greater improvement than continuous HT. As a result, despite its

contribution to brain's neuroprotective mechanism, HT treatment may entail

overheating causing protein dysfunction in neural cells and thus interrupting the

pathways for neuroprotection. Therefore, the interspersing low-temperature process is

crucial for TC-HT treatment's neuroprotective effect. Our study demonstrates that TC-

HT can be an effective thermal treatment for AD. In practice, high-intensity focused

ultrasound (HIFU) may be a suitable heating source for TC-HT treatment, as it is

capable of precisely modulating the temperature at the targeted area, via computer-

controlled programmable procedure. The HIFU heating area can be set at only about 4-

8 mm [65, 66], smaller than human brain, thereby quickening heat dissipation to

surrounding tissues and preventing hyperthermic damage. In operation, multi-focal

HIFU can control the heating area and temperature by tuning the size of scanned spot

and intensity in scanning mode, thereby attaining set temperature and effective thermal

dissipation for TC-HT in AD treatment.

In summary, our study puts forth a novel hyperthermic method, TC-HT, for

application as a potential AD treatment *in vivo*, which demonstrates a better effect in

improving the cognitive performance of mice in Y-maze and NOR tests than

conventional HT treatment, which heats the subjects continuously. Furthermore, the

study shows that TC-HT can suppress the hippocampal Aβ expression to a normal level,

significantly outperforming HT which only exhibits a slight inhibition effect, perhaps

due to TC-HT's higher capability in inhibiting the Aβ productive activity of BACE1

protein and promoting the Aβ clearance effect of IDE protein. In addition, TC-HT can

significantly inhibit the Aβ-induced overexpression of neuroinflammatory markers Iba-

1 and GFAP in hippocampi, while HT treatment only slightly attenuates their

expression levels without significance. Moreover, TC-HT raises the expression level of

antioxidative protein SOD2 at an extent much higher than HT, indicating that it can

provide a better protection for hippocampal neurons against the i.c.v.-injected Aβ.

Overall, the study validates the therapeutic potential of TC-HT for AD *in vivo*, shedding

light on development of alternative AD treatment with high efficacy.

545 Funding

546   This work was supported by grants from Ministry of Science and Technology

547 (MOST 110-2112-M-002-004, MOST 109-2112-M-002-004, and MOST 108-2112-M-

548 002-016 to CYC) of the Republic of China. The funders had no role in study design,

549 data collection and analysis, decision to publish, or preparation of the manuscript.

551 Competing Interests

552   The authors have declared that no competing interests exist.

554 Acknowledgments

555   The authors would like to acknowledge the service provided by Animal Resource

556 Center of National Taiwan University for facility support.

558 Author Contributions

Conceptualization: Chih-Yu Chao.

Data Curation: Yu-Yi Kuo, Chih-Yu Chao.

Formal analysis: Yu-Yi Kuo, Wei-Ting Chen, Guan-Bo Lin, You-Ming Chen, Hsu-

Hsiang Liu, Chih-Yu Chao.

Funding acquisition: Chih-Yu Chao.

Investigation: Yu-Yi Kuo, Wei-Ting Chen, Guan-Bo Lin, Chih-Yu Chao.

Project Administration: Chih-Yu Chao.

Supervision: Chih-Yu Chao.

Validation: Yu-Yi Kuo, Wei-Ting Chen, Guan-Bo Lin, You-Ming Chen, Hsu-Hsiang

Liu

Writing – original draft: Yu-Yi Kuo, Chih-Yu Chao.

Writing – review & editing: Chih-Yu Chao.

References

1. 2020 Alzheimer's disease facts and figures. Alzheimer's Dement. 2020;
16(3):391-460.
2. Chen GF, Xu TH, Yan Y, Zhou YR, Jiang Y, Melcher K, et al. Amyloid beta:
structure, biology and structure-based therapeutic development. Acta
Pharmacol Sin. 2017; 38(9):1205-35.

3. Sun X, Chen WD, Wang YD. β-amyloid: the key peptide in the pathogenesis of
Alzheimer's disease. Front Pharmacol. 2015; 6:221.

4. Essa MM, Subash S, Akbar M, Al-Adawi S, Guillemin GJ. Long-term dietary
supplementation of pomegranates, figs and dates alleviate neuroinflammation
in a transgenic mouse model of Alzheimer's disease. PLoS One. 2015;
10(3):e0120964.

5. García-Escudero V, Martín-Maestro P, Perry G, Avila J. Deconstructing
mitochondrial dysfunction in Alzheimer disease. Oxid Med Cell Longev. 2013;
2013:162152.

6. Leuner K, Schütt T, Kurz C, Eckert SH, Schiller C, Occhipinti A, et al.
Mitochondrion-derived reactive oxygen species lead to enhanced amyloid beta
formation. Antioxid Redox Signal. 2012; 16(12):1421-33.

7. Palop JJ, Mucke L. Amyloid-β-induced neuronal dysfunction in Alzheimer's
disease: from synapses toward neural networks. Nat Neurosci. 2010; 13(7):812-
8.

8. Kashyap G, Bapat D, Das D, Gowaikar R, Amritkar RE, Rangarajan G, et al.
Synapse loss and progress of Alzheimer's disease - a network model. Sci Rep.
2019; 9(1):6555.

9. Karran E, Mercken M, De Strooper B. The amyloid cascade hypothesis for
Alzheimer's disease: an appraisal for the development of therapeutics. Nat Rev
Drug Discov. 2011; 10(9):698-712.

10. Cao J, Hou J, Ping J, Cai D. Advances in developing novel therapeutic strategies
for Alzheimer's disease. Mol Neurodegener. 2018; 13(1):64.

11. Du Z, Li M, Ren J, Qu X. Current strategies for modulating Aβ aggregation
with multifunctional agents. Acc Chem Res. 2021; 54(9):2172-84.

12. Lee D, Lee G, Yoon DS. Anti-Aβ drug candidates in clinical trials and
plasmonic nanoparticle-based drug-screen for Alzheimer's disease. Analyst.
2018; 143(10):2204-12.

13. Xin SH, Tan L, Cao X, Yu JT, Tan L. Clearance of amyloid beta and tau in
Alzheimer's disease: from mechanisms to therapy. Neurotox Res. 2018;
34(3):733-48.

14. Hung SY, Fu WM. Drug candidates in clinical trials for Alzheimer's disease. J
Biomed Sci. 2017; 24(1):47.

15. Habash RW, Bansal R, Krewski D, Alhafid HT. Thermal therapy, part 1: an
introduction to thermal therapy. Crit Rev Biomed Eng. 2006; 34(6):459-89.

613 16. Manning MR, Cetas TC, Miller RC, Oleson JR, Connor WG, Gerner EW.
614   Clinical hyperthermia: results of a phase I trial employing hyperthermia alone
615   or in combination with external beam or interstitial radiotherapy. Cancer. 1982;
616   49(2):205-16.

617 17. Falk MH, Issels RD. Hyperthermia in oncology. Int J Hyperthermia. 2001;
618   17(1):1-18.

619 18. Tabuchi Y, Kondo T. Targeting heat shock transcription factor 1 for novel
620   hyperthermia therapy (review). Int J Mol Med. 2013; 32(1):3-8.

621 19. Rivera I, Capone R, Cauvi DM, Arispe N, De Maio A. Modulation of
622   Alzheimer's amyloid β peptide oligomerization and toxicity by extracellular
623   Hsp70. Cell Stress Chaperones. 2018; 23(2):269-79.

624 20. Wilhelmus MM, Boelens WC, Otte-Höller I, Kamps B, de Waal RM, Verbeek
625   MM. Small heat shock proteins inhibit amyloid-β protein aggregation and
626   cerebrovascular amyloid-β protein toxicity. Brain Res. 2006; 1089(1):67-78.

627 21. Tundo GR, Sbardella D, Ciaccio C, Bianculli A, Orlandi A, Desimio MG, et al.
628   Insulin-degrading enzyme (IDE): a novel heat shock-like protein. J Biol Chem.
629   2013; 288(4):2281-9.

630 22. Ahmed K, Zaidi SF, Mati-Ur-Rehman, Rehman R, Kondo T. Hyperthermia and
631   protein homeostasis: cytoprotection and cell death. J Therm Biol. 2020;
632   91:102615.

633 23. Banerjee Mustafi S, Chakraborty PK, Dey RS, Raha S. Heat stress upregulates
634   chaperone heat shock protein 70 and antioxidant manganese superoxide
635   dismutase through reactive oxygen species (ROS), p38MAPK, and Akt. Cell
636   Stress Chaperones. 2009; 14(6):579-89.

637 24. Hashemi M, Abbasiazam A, Oraee-Yazdani S, Lenzer J. Response of human
638   glioblastoma cells to hyperthermia: cellular apoptosis and molecular events.
639   Tissue Cell. 2022; 75:101751.

640 25. McLaughlin CT, Kane AG, Auber AE. MR imaging of heat stroke: external
641   capsule and thalamic T1 shortening and cerebellar injury. AJNR Am J
642   Neuroradiol. 2003; 24(7):1372-5.

643 26. Lee JS, Choi JC, Kang SY, Kang JH, Park JK. Heat stroke: increased signal
644   intensity in the bilateral cerebellar dentate nuclei and splenium on diffusion-
645   weighted MR imaging. AJNR Am J Neuroradiol. 2009; 30(4):e58.

646 27. Walter EJ, Carraretto M. The neurological and cognitive consequences of
647   hyperthermia. Crit Care. 2016; 20(1):199.

28. Cremer OL, Diephuis JC, van Soest H, Vaessen PH, Bruens MG, Hennis PJ, et
al. Cerebral oxygen extraction and autoregulation during extracorporeal whole
body hyperthermia in humans. Anesthesiology. 2004; 100(5):1101-7.

29. Laukkanen T, Kunutsor S, Kauhanen J, Laukkanen JA. Sauna bathing is
inversely associated with dementia and Alzheimer's disease in middle-aged
Finnish men. Age Ageing. 2017; 46(2):245-9.

30. Lu CH, Chen WT, Hsieh CH, Kuo YY, Chao CY. Thermal cycling-
hyperthermia in combination with polyphenols, epigallocatechin gallate and
chlorogenic acid, exerts synergistic anticancer effect against human pancreatic
cancer PANC-1 cells. PLoS One. 2019; 14(5):e0217676.

31. Chen WT, Sun YK, Lu CH, Chao CY. Thermal cycling as a novel thermal
therapy to synergistically enhance the anticancer effect of propolis on PANC-1
cells. Int J Oncol. 2019; 55(3):617-28.

32. Chen WT, Kuo YY, Lin GB, Lu CH, Hsu HP, Sun YK, et al. Thermal cycling
protects SH-SY5Y cells against hydrogen peroxide and β-amyloid-induced cell
injury through stress response mechanisms involving Akt pathway. PLoS One.
2020; 15(10):e0240022.

33. Kim HY, Lee DK, Chung BR, Kim HV, Kim Y. Intracerebroventricular
injection of amyloid-β peptides in normal mice to acutely induce Alzheimer-
like cognitive deficits. J Vis Exp. 2016; (109):53308.

34. Al Faruque H, Choi ES, Lee HR, Kim JH, Park S, Kim E. Targeted removal of
leukemia cells from the circulating system by whole-body magnetic
hyperthermia in mice. Nanoscale. 2020; 12(4):2773-86.

35. Lin C, Liu Z, Li Y, Chen J. Protocol for fever-range whole-body hyperthermia
(WBH) in mice to study febrile effect on T-cell adhesion and migration. STAR
Protoc. 2021; 2(3):100720.

36. Saim H, Salim MIM, Jemon K. Antitumor effect of fever range whole body
hyperthermia with curcumin in breast cancer-induced mice. Int J Recent
Technol Eng. 2019; 8(2):297-301.

37. Ohsawa I, Ishikawa M, Takahashi K, Watanabe M, Nishimaki K, Yamagata K,
et al. Hydrogen acts as a therapeutic antioxidant by selectively reducing
cytotoxic oxygen radicals. Nat Med. 2007; 13(6):688-94.

38. Murakami Y, Ito M, Ohsawa I. Molecular hydrogen protects against oxidative
stress-induced SH-SY5Y neuroblastoma cell death through the process of
mitohormesis. PLoS One. 2017; 12(5):e0176992.

39. Hampel H, Vassar R, De Strooper B, Hardy J, Willem M, Singh N, et al. The
β-secretase BACE1 in Alzheimer's disease. Biol Psychiatry. 2021; 89(8):745-
56.

40. Huang W, Li Z, Zhao L, Zhao W. Simvastatin ameliorate memory deficits and
inflammation in clinical and mouse model of Alzheimer's disease via
modulating the expression of miR-106b. Biomed Pharmacother. 2017; 92:46-
57.

41. Tundo GR, Sbardella D, Ciaccio C, Grasso G, Gioia M, Coletta A, et al.
Multiple functions of insulin-degrading enzyme: a metabolic crosslight? Crit
Rev Biochem Mol Biol. 2017; 52(5):554-82.

42. Massaad CA, Washington TM, Pautler RG, Klann E. Overexpression of SOD-
2 reduces hippocampal superoxide and prevents memory deficits in a mouse
model of Alzheimer's disease. Proc Natl Acad Sci USA. 2009; 106(32):13576-
81.

43. Selkoe DJ, Hardy J. The amyloid hypothesis of Alzheimer's disease at 25 years.
EMBO Mol Med. 2016; 8(6):595-608.

44. Naldi M, Fiori J, Pistolozzi M, Drake AF, Bertucci C, Wu R, et al. Amyloid β-
peptide 25-35 self-assembly and its inhibition: a model undecapeptide system
to gain atomistic and secondary structure details of the Alzheimer's disease
process and treatment. ACS Chem Neurosci. 2012; 3(11):952-62.

45. Xiao Y, Ma B, McElheny D, Parthasarathy S, Long F, Hoshi M, et al. Aβ(1-42)
fibril structure illuminates self-recognition and replication of amyloid in
Alzheimer's disease. Nat Struct Mol Biol. 2015; 22(6):499-505.

46. Jakki SL, Senthil V, Yasam VR, Chandrasekar MJN, Vijayaraghavan C. The
blood brain barrier and its role in Alzheimer's therapy: an overview. Curr Drug
Targets. 2018; 19(2):155-69.

47. Jahn H. Memory loss in Alzheimer's disease. Dialogues Clin Neurosci. 2013;
15(4):445-54.

48. Kraeuter AK, Guest PC, Sarnyai Z. The Y-maze for assessment of spatial
working and reference memory in mice. Methods Mol Biol. 2019; 1916:105-11.

49. Lueptow LM. Novel object recognition test for the investigation of learning and
memory in mice. J Vis Exp. 2017; (126):55718.

50. Qiao O, Ji H, Zhang Y, Zhang X, Zhang X, Liu N, et al. New insights in drug
development for Alzheimer's disease based on microglia function. Biomed
Pharmacother. 2021; 140:111703.

51. Ahmad MH, Fatima M, Mondal AC. Influence of microglia and astrocyte activation in the neuroinflammatory pathogenesis of Alzheimer's disease: rational insights for the therapeutic approaches. J Clin Neurosci. 2019; 59:6-11.

52. Heneka MT, Carson MJ, El Khoury J, Landreth GE, Brosseron F, Feinstein DL, et al. Neuroinflammation in Alzheimer's disease. Lancet Neurol. 2015; 14(4):388-405.

53. Vekrellis K, Ye Z, Qiu WQ, Walsh D, Hartley D, Chesneau V, et al. Neurons regulate extracellular levels of amyloid β-protein via proteolysis by insulin-degrading enzyme. J Neurosci. 2000; 20(5):1657-65.

54. Cook DG, Leverenz JB, McMillan PJ, Kulstad JJ, Ericksen S, Roth RA, et al. Reduced hippocampal insulin-degrading enzyme in late-onset Alzheimer's disease is associated with the apolipoprotein E-ε4 allele. Am J Pathol. 2003; 162(1):313-9.

55. Eckman EA, Eckman CB. Aβ-degrading enzymes: modulators of Alzheimer's disease pathogenesis and targets for therapeutic intervention. Biochem Soc Trans. 2005; 33(5):1101-5.

56. Massaad CA, Klann E. Reactive oxygen species in the regulation of synaptic plasticity and memory. Antioxid Redox Signal. 2011; 14(10):2013-54.

57. Kim YS, Gupta Vallur P, Phaëton R, Mythreye K, Hempel N. Insights into the dichotomous regulation of SOD2 in cancer. Antioxidants. 2017; 6(4):86.

58. Esposito L, Raber J, Kekonius L, Yan F, Yu GQ, Bien-Ly N, et al. Reduction in mitochondrial superoxide dismutase modulates Alzheimer's disease-like pathology and accelerates the onset of behavioral changes in human amyloid precursor protein transgenic mice. J Neurosci. 2006; 26(19):5167-79.

59. Tadokoro K, Ohta Y, Inufusa H, Loon AFN, Abe K. Prevention of cognitive decline in Alzheimer's disease by novel antioxidative supplements. Int J Mol Sci. 2020; 21(6):1974.

60. Milkovic L, Cipak Gasparovic A, Cindric M, Mouthuy PA, Zarkovic N. Short overview of ROS as cell function regulators and their implications in therapy concepts. Cells. 2019; 8(8):793.

61. Unoki T, Abiko Y, Toyama T, Uehara T, Tsuboi K, Nishida M, et al. Methylmercury, an environmental electrophile capable of activation and disruption of the Akt/CREB/Bcl-2 signal transduction pathway in SH-SY5Y cells. Sci Rep. 2016; 6(1):28944.

62. Jin Y, Wang H. Naringenin inhibit the hydrogen peroxide-induced SH-SY5Y
cells injury through Nrf2/HO-1 pathway. Neurotox Res. 2019; 36(4):796-805.
63. Sun P, Nie X, Chen X, Yin L, Luo J, Sun L, et al. Nrf2 signaling elicits a
neuroprotective role against PFOS-mediated oxidative damage and apoptosis.
Neurochem Res. 2018; 43(12):2446-59.
64. Lepock JR. Protein denaturation during heat shock. Adv Mol Cell Biol. 1997;
19:223-59.
65. Chaplin V, Caskey CF. Multi-focal HIFU reduces cavitation in mild-
hyperthermia. J Ther Ultrasound. 2017; 5(1):12.
66. Partanen A, Tillander M, Yarmolenko PS, Wood BJ, Dreher MR, Kohler MO.
Reduction of peak acoustic pressure and shaping of heated region by use of
multifoci sonications in MR-guided high-intensity focused ultrasound mediated
mild hyperthermia. Med Phys. 2013; 40(1):013301.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

B

Figure 7