Mining the human gut microbiome identifies mycobacterial D-arabinan degrading enzymes

Omar Al-Jourani[1, ♣], Samuel Benedict[2,♣], Jennifer Ross[1, ♣], Abigail Layton[2], Phillip van der Peet[3], Victoria M. Marando[4], Nicholas P. Bailey[1], Tiaan Heunis[1], Joseph Manion[1], Francesca Mensitieri[1], Aaron Franklin[2], Javier Abellon-Ruiz[1], Sophia L. Oram[1], Lauren Parsons[1], Alan Cartmell[5], Gareth S. A. Wright[6], Arnaud Baslé[1], Matthias Trost[1], Bernard Henrissat[7], Jose Munoz-Munoz[8], Robert P. Hirt[1], Laura L. Kiessling[4], Andrew Lovering[2], Spencer J. Williams[3], Elisabeth C. Lowe[1,*], Patrick J. Moynihan[2]*

Addresses:

[1]Newcastle University Biosciences Institute, Medical School, Newcastle University, Newcastle upon Tyne, U.K., NE2 4HH

[2]Institute of Microbiology and Infection, School of Biosciences, University of Birmingham, Birmingham, U.K., B15 2TT

[3]School of Chemistry and Bio21 Molecular Science and Biotechnology Institute, University of Melbourne, Parkville, Victoria 3010, Australia

[4] Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA, USA

The Broad Institute of Harvard and MIT, Cambridge, MA, USA The Koch Integrative Cancer Research Institute, Massachusetts Institute of Technology, Cambridge, MA, USA

[5] Department of Biochemistry and Systems Biology, Institute of Systems, Molecular and Integrative Biology, University of Liverpool, Liverpool, UK.

[6] School of Life Sciences, University of Essex, Colchester, UK

[7] Department of Biological Sciences, King Abdulaziz University, Jeddah, Saudi

28   Arabia;

Department of Biotechnology and Biomedicine (DTU Bioengineering), Technical

30   University of Denmark, 2800 Kgs. Lyngby, Denmark

[8] Microbial Enzymology Group, Department of Applied Sciences, Northumbria

32   University, Newcastle upon Tyne, United Kingdom.

34
♣ These authors contributed equally to this work

36   * To whom correspondence should be addressed:
Patrick Moynihan, p.j.moynihan@bham.ac.uk
38   Elisabeth Lowe, elisabeth.lowe@ncl.ac.uk

Key words:

40   carbohydrate-active enzymes (CAZymes); glycoside hydrolase (GH); tuberculosis

(TB); GH172; DUF4185

42   **Abbreviations**:

AG, arabinogalactan; DUF, domain of unknown function; GH, glycoside hydrolase;
44   IC-PAD, ion chromatography/ pulsed amperometric detection; LAM,
Lipoarabinomannan; PUL, polysaccharide utilization locus;

Abstract

Division and degradation of bacterial cell walls requires coordinated action of a myriad of enzymes. This particularly applies to the elaborate cell walls of acid-fast organisms such as *Mycobacterium tuberculosis*, which consist of a multi-layered cell wall that contains an unusual glycan called arabinogalactan. Enzymes that cleave the D-arabinan core of this structure have not previously been identified in any organism. We have interrogated the diverse carbohydrate degrading enzymes expressed by the human gut microbiota and uncovered four families of glycoside hydrolases with the capability to degrade the D-arabinan or D-galactan components of arabinogalactan. Using novel exo-D-galactofuranosidases from gut bacteria we generated enriched D-arabinan and used it to identify *D. gadei* as a D-arabinan degrader. This enabled the discovery of endo- and exo- acting enzymes that cleave D-arabinan. We have identified new members of the DUF2961 family (GH172), and a novel family of glycoside hydrolases (DUF4185) that display endo-D-arabinofuranase activity. The DUF4185 enzymes are conserved in mycobacteria and found in many microbes, suggesting that the ability to degrade mycobacterial glycans plays an important role in the biology of diverse organisms. All mycobacteria encode two conserved endo-D-arabinanases that display different preferences for the D-arabinan-containing cell wall components arabinogalactan and lipoarabinomannan, suggesting they are important for cell wall modification and/or degradation. The discovery of these enzymes will support future studies into the structure and function of the mycobacterial cell wall.

Introduction

Growth and division of all bacteria is a carefully orchestrated process requiring the coordinated action of a host of enzymes. Acid-fast organisms such as *Mycobacterium tuberculosis* possess unusual cell wall glycans and lipids which require additional enzymatic machinery during this process. The core of the cell wall structure is conserved amongst mycobacteria and consists of three layers[1,2]. Like other bacteria, peptidoglycan forms the basal layer of the cell wall, though the precise architecture is unknown. At the other extremity of the wall are the mycolic acids which give the organisms their characteristic waxy appearance and are interspersed with a host of species-specific free lipids. Joining these two layers is a complex polysaccharide called arabinogalactan (AG), which has a chemical composition unique to the Mycobacteriales and entirely distinct from the similarly named molecule found in plants that is composed of L-arabinofuranose (**Figure 1A**)[3]. AG is comprised of two domains with a β-D-galactofuranose backbone decorated by large α-D-arabinofuranose branches[4]. The structure and biosynthesis of this molecule has undergone intense scrutiny, due in part to it being a target of the antimycobacterial drug ethambutol[5-8]. Ethambutol targets the arabinosyltransferase proteins in the cell envelope of mycobacteria that are responsible for biosynthesis of the polysaccharide[9]. Similarly, the biogenesis of mycolic acids and peptidoglycan are the subject of much research due to their biochemical complexity, essentiality, and their synthesis being the target of isoniazid and β-lactams respectively[10,11].

The growth of mycobacteria likely requires not only the coordinated synthesis of the cell wall, but also its eventual degradation and turn-over to allow for cellular expansion, division and the insertion of cell envelope spanning secretion systems. Similarly, the control of cell surface structures affords these bacteria the ability to

modulate how they are sensed by the host. As such, hydrolases of the cell wall may

96   play an important role at the mycobacterial cell surface. Hydrolases of peptidoglycan

influence mycobacterial morphology and infection and it was recently demonstrated

98   that at least some peptidoglycan by-products are recycled by the bacteria[12,13].

Similarly, trehalose-mycolates, a dominant lipid component of the cell wall, undergo

100   recycling,[14] and a hydrolase involved in turning over mycolic acids (Rv3451) and a

recycling pathway for trehalose has been described[15]. Cleavage of the acid-fast cell

102   wall is unlikely to be restricted to the Mycobacteriales. Organisms that predate on acid-

fast bacteria such as phage or bacterial predators like the recently reported

104   *Candidatus* Mycosynbacter amalyticus (hereafter M. amalyticus)*,* are also likely to

require D-arabinan degrading enzymes to penetrate the mycobacterial cell wall[16].

106   Only a single enzyme capable of degrading AG has been characterized to date.

GlfH1 (Rv3096) is an exo-β-D-galactofuranosidase from glycoside hydrolase (GH)

108   family GH5_13 that cleaves the galactan backbone of mycobacterial AG at both β-1,5

and β-1,6 linked residues[17]. The precise role of this enzyme in mycobacterial biology

110   remains unclear but its activity is suggestive of it being part of a remodeling pathway

for this cell wall component. Enzymes with activity on simple D-galactofuranosides (D-

112   Gal*f*) have been identified through large-scale screening of orphan GH sequences[18].

Likewise, no enzymes have been characterized with exo- or endo-D-arabinofuranase

114   activity (enzymes that cut within the D-arabinan polymers) against AG, although this

activity was described in protein extracts from soil bacteria dating back to the

116   1970s[19,20]. Endo-arabinan activity was also described in extracts of *Mycobacterium*

smegmatis and was suggested to increase upon treatment of cells with ethambutol,

118   which blocks D-arabinan biosynthesis[21,22]. During our studies an exo-acting difructose-

dianhydride I synthase/hydrolase was discovered that was active on *p*NP-α-D-arabinofuranoside[23]. Whether this enzyme acts on mycobacterial AG is unknown.

The human gut microbiota is responsible for the degradation of dietary plant polysaccharides, host and microbial glycans. Dominated by the Bacteroidetes, this grouping of organisms is collectively amongst the richest known organisms in diversity of complex carbohydrate degrading enzymes[24]. Carbohydrate utilization by the Bacteroidetes is typically mediated by genes which are organized into polysaccharide utilization loci (PULs), which can be induced upon exposure of the bacterium to a given carbohydrate[25-28]. The abundance and diversity of carbohydrate degrading enzymes in the Bacteroidetes provides a rich opportunity for enzyme discovery.

In this study we have mined the glycolytic capacity of the human gut microbiota and discovered a collection of enzymes able to completely degrade mycobacterial arabinogalactan. We report the discovery of new families of glycoside hydrolases active on the mycobacterial cell wall. We also identify new exo-D-arabinofuranosidases from the DUF2961 family (GH172). Furthermore, we demonstrate that D-arabinan degradation is wide-spread amongst Mycobacteriales, and mycobacteria but is also present in phages, other bacteria, and microbial eukaryotes. Our data point to a key role for these enzymes in mycobacterial biology and can enable sophisticated analysis of mycobacterial cell wall components.

Results

Select human gut Bacteroidetes can utilize mycobacterial AG as a carbon source

While endo-D-arabinofuranase activity was first described in soil bacteria more than 50 years ago and has been known in mycobacteria for at least 30 years[19,20], the enzymes responsible for this activity have escaped identification. We reasoned that

these enzymes might be highly regulated, unstable, or poorly soluble in mycobacteria

146   making purification-based approaches unfeasible. Moreover, if they belong to novel

enzyme class(es), bioinformatics approaches would fail. Instead, we isolated

148   arabinogalactan from *M. smegmatis* mc^{2}155 and used it as sole-carbon source for the

growth of a panel of 14 Bacteroidetes species. Of these, 12 strains were able to grow

150   on this material (**Figure 1B**). Ion chromatography with pulsed amperometric detection

(IC-PAD) analysis of selected culture supernatants (**Figure 1C**) demonstrated the

152   production of free galactose in most cultures, and arabinose in one; *Dysgonomonas*

gadei. Together these data indicate that members of the gut microbiota produce

154   enzymes that can depolymerize mycobacterial D-arabinan and D-galactan.

Figure 1. Bacteroidetes growth on mycobacterial arabinogalactan. A) Schematic of the structure of mycobacterial mycolyl-arabinogalactan-peptidoglycan complex. Succinate (‡) and galactosamine (*) modifications are non-stoichiometric. **B)** Growth of selected Bacteroidetes species on arabinogalactan as monitored by change in OD_{600nm}. **C)** IC-PAD analysis of culture supernatants for bacteria grown on mycobacterial AG as a sole-carbon source. Production of arabinose (green star) and galactose (yellow circle) are indicated and compared to standards. **D)** Schematics of PUL47 and 39 from *B. finegoldii*, whose homologs in *B. cellulosilyticus* were identified by RNAseq as being upregulated during growth on AG. **E)** TLC analysis of endpoint reaction products of *B. finegoldii* enzymes with *C. glutamicum* Δ*ubiA* galactan. Filled circles indicate the presence of a given reaction component.

Identification of exo- and endo-galactofuranosidases that degrade galactan

170   The presence of both galactose and arabinose in *D. gadei* culture supernatants

complicated the identification of PULs specific for either galactan or arabinan.

172   Therefore, we developed a method for production of pure D-arabinan by exploiting D-

galactan specific PULs. Based on the analysis of culture supernatants, *B. finegoldii*

174   and *B. cellulosilyticus* appeared to degrade galactan, but not D-arabinan. RNAseq

analysis of *B. cellulosilyticus* revealed the upregulation of PUL35 and PUL36,

176   containing multiple predicted GHs (**Figure S1**). While we could not heterologously

express the *B. cellulosilyticus* enzymes, the homologs of these enzymes from *B.*

178   *finegoldii* DSM17565 (PULDB ID: PUL39 and PUL47, **Figure 1D**) could be expressed

and purified, and when tested on galactan (**Figure 1E**) demonstrated exo- and endo-

180   activities. To determine their galactan degradative capacity, we purified galactan,

comprised alternating β-1,5- and β-1,6-galactofuranose residues from a strain of

182   *Corynebacterium glutamicum* (ΔubiA) that lacks D-arabinan[29]. As shown in **Figure 1E**

and **Figure S1**, combination of the two *B. finegoldii* enzymes comprising a GH43_31

184   (BACFIN_08810) and a new GH-exo-Gal*f* (BACFIN_04787) family completely

hydrolysed this galactan substrate.

186   *Cultivation on D-arabinan identifies D-arabinan-active PULs*

       To generate D-arabinan we digested mycobacterial AG with the two *B. finegoldii*

188   galactofuranosidases (BACFIN_08810 and BACFIN_04787). This resulted in an

enriched D-arabinan fraction with approximately 70% reduction in galactan as

190   determined by acid hydrolysis (**Figure S1**). The resulting D-arabinan was used as a

sole carbon source for *D. gadei* (which was previously shown to produce both D-

192   galactose and D-arabinose from AG). Proteomics of these bacteria at mid-log phase

during growth on enriched D-arabinan identified a predicted fucose isomerase as the

194   most abundant carbohydrate-active enzyme (**Table S1**). This protein maps to PUL42

in the *D. gadei* genome **(Figure 2A**), and an additional nine of the proteins derived

196   from this PUL were in the top 200 most abundant proteins in the total proteome,

including several that lacked annotation (**Figure S2**). Reasoning that mycobacteria

198   may harbor homologs of *D. gadei* arabinanases to process arabinan, we prioritized

the DUF2961 and DUF4185 superfamily enzymes encoded in PUL42 as they

200   possessed homology to predicted mycobacterial proteins of unknown function within

the same superfamilies.

202   *The DUF2961 superfamily (GH172) includes D-arabinofuranosidases*

At the outset of this study, no member of the DUF2961 family had been

204   characterized. Therefore, we cloned, expressed, and purified all three DUF2961 family

members found in PUL42 in *D. gadei*. Upon incubation with purified AG, we observed

206   D-arabinofuranosidase activity for HMPREF9455_02467, HMPREF9455_02471 and

HMPREF9455_02479 (hereafter Dg$_{GH172a}$, Dg$_{GH172b}$ and Dg$_{GH172c}$, respectively)

208   **(Figure S3)**. To probe the activity of these enzymes, we synthesized the chromogenic

substrates   *p*-nitrophenyl   α-D-arabinofuranoside   (pNP-α-D-Ara*f*)   and   β-D-

210   arabinofuranoside (pNP-β-D-Ara*f*). Dg$_{GH172a}$ and Dg$_{GH172c}$ were active against pNP-α-

D-Ara*f*, and no activity was observed using pNP-β-D-Ara*f* (**Table 1**). Although substrate

212   limitations prevented accurate determination of V_{max} we could use pNP-α-D-Ara*f* to

determine k_{cat}/K_M for Dg$_{GH172c}$ (**Figure S4 and Table 2**). These data demonstrate that

214   gut bacteria can use DUF2961 enzymes to generate D-arabinose from mycobacterial

arabinogalactan.

216   *DUF2961 superfamily (GH172) enzymes are present in acid-fast bacteria and their*

predators

Homologues of the DUF2961 encoding genes are present in the genomes of organisms from the Actinomycetota phylum including pathogens such as

Mycobacterium avium subsp. *paratuberculosis* (MAP_0339c) and *Nocardia brasiliensis* (O3I_017420; Noc$_{GH172}$)*.* Noc$_{GH172}$ was soluble; however, despite repeated

attempts, we were unable to produce soluble MAP_0339c. Noc$_{GH172}$ had D-arabinofuranosidase activity on both pNP-α-D-Ara*f* and purified arabinogalactan

(**Table 2, Figure S3**). We also identified a DUF2961 homolog within the genome of the parasitic bacterium M. amalyticus (GII36_05205; Myc$_{GH172}$), which possessed

similar activity (**Figure 2 and S3A**)[16]. Together these data indicate that exo-D-arabinofuranosidase activity is a feature of the DUF2961 super-family. These

enzymes are encoded by both the Mycobacteriales and their predators.

Figure 2. ***D. gadei* DUF2961 genes encode GH172 exo-D-arabinofuranosidases with orthologs in diverse bacteria**. **A**) Schematic of PUL42 as identified by proteomic analysis of *D. gadei* grown on D-arabinan. **B**) DUF2961 enzymes (1 µM) vs 2 mg ml^{-1} AG and **C**) LAM. Samples were analyzed on a Dionex ICS-6000 with CarboPac Pa300 column, 100 mM NaOH, 20 min isocratic elution followed by a 0-60% 500 mM sodium acetate gradient over 60 minutes. Green star = D-arabinose.

GH172 enzymes degrade α-1,5-D-arabinofuranoside linkages within D-arabinan

240   To understand the substrate specificity of these enzymes against other microbial D-arabinan substrates, we assayed our DUF2961 GH172 enzymes

242   (Dg_{GH172a}, Dg_{GH172b}, Dg_{GH172c}, Noc_{GH172}, and Myc_{GH172}) against mycobacterial LAM and D-araF containing pilin oligosaccharides (**Table 1**). The D-arabinan branches of

244   lipoarabinomannan (LAM) are structurally conserved with those in AG. All enzymes could digest LAM from *M. smegmatis* to produce arabinose, although Dg_{GH172b} and

246   Myc_{GH172} displayed lower activity than the others (**Figure 2C and Figure S3B**).

Pili from *Pseudomonas aeruginosa* PA7 are decorated with α-1,5-D-

248   arabinofuranose oligosaccharides. All our GH172, except for Dg_{GH172b} enzymes could cleave the D-arabinofuranose oligosaccharides from digested pili (**Figure S3C**). To

250   rule out D-galactofuranosidase activity, we repeated these experiments using purified D-galactan (**Figure S3D**) and observed no activity. The combination of these

252   experiments reveals that GH172 enzymes are specific for α-D-arabinofuranoside linkages, are active on D-arabinan, and can cleave α-1,5-D-arabinofuranoside

254   linkages.

Family GH172 proteins can adopt diverse oligomeric states

256   To better understand the structure-function relationship of this protein family, we examined the multimeric state of several of our candidates of interest. Size-

258   exclusion chromatography with laser scattering (SEC-LS) analysis of the GH172 family enzymes allowed assessment of molecular weight and assignment of

260   oligomerization states (**Table 3 and Figure S7**). A wide range of oligomeric states were uncovered: Dg_{GH172c} was assigned a hexamer, Dg_{GH172a} a dodecamer, Noc_{GH172}

262   and Myc_{GH172} as trimers, and Dg_{GH172b} as a dimer. This solution data complements

recent reports of hexameric assemblies in protein crystals of two GH172 members:

264   difructose dianhydride I synthase/hydrolase from *Bifidobacterium dentium* (PDB: 7V1V) and BACUNI_00161 from *Bacteroides uniformis* (PDB: 4KQ7) (**Figure S5**)[23].

266   Interestingly, Dg_{GH172b} is predicted to have 1.5 DUF2961 domains, and so dimerization of Dg_{GH172b} is predicted to provide three DUF2961 domains in the final

268   assembly. SEC-LS analysis is supportive of a quaternary structure for each of the GH172 proteins exhibiting multiple DUF2961 domains: Dg_{GH172b}, Myc_{GH172} and

270   Noc_{GH172} each contain three DUF2961 domains, while Dg_{GH172c} contains six and Dg_{GH172a} contains twelve. Diversity in quaternary structure may lead to differences in

272   activity or substrate specificity.

The 3D structure of Dg_{GH172c} reveals a dimer-of-trimers assembly

The 3D structure of Dg_{GH172c} was determined by X-ray crystallography using molecular replacement with the *B. uniformis* GH172 (Bu_{GH172}, PDB: 4KQ7; 64% sequence

identity with Dg_{GH172c}) as the search model. The structure of Dg_{GH172c} was refined to 1.4 Å resolution, data collection and model refinement statistics are shown in **Table**

S2. The Dg_{GH172c} monomer is comprised of two β jelly rolls and a C-terminal alpha helix (**Figure 3A**). Six monomers assemble in a dimer-of-trimers (**Figure 3B and**

Figure S5A), consistent with the solution phase oligomerization state determined by SEC-LS (**Table 3**). The dimer-of-trimers structure is formed by the C-terminal of each

monomer of one trimer interlocking with the C-terminal of a monomer from a second trimer. The lower side of the trimers face each other in the dimer-of-trimers structures.

The interfaces within the trimer form the active site with residues from both monomers contributing to the active site (**Figure 3C**). This gives a total of 3 active sites per trimer,

and six per hexameric assembly. Seven calcium ions are coordinated within each

Dg$_{GH172c}$ trimer: two within each active site/protomer, and one in the 3-fold axes.

Figure 3: 3D structure of Dg$_{GH172c}$ determined by X-ray crystallography. A)
290 Monomer of Dg$_{GH172a}$ with secondary structure shown as cartoon. The N- and C-
termini are labelled and colored blue and red respectively. Bound calcium ions are
292 shown as green spheres. **B)** Overview of hexameric architecture of Dg$_{GH172c}$. Cartoon
representation with bound calcium ions shown as green sphere. Trimer subunits are
294 colored grey, pink and purple for clarity. **C)** Surface representation of the hexamer with
one monomer shown in cartoon. Active site residues are colored gold, and calcium
296 ions shown in green. **D)** Cartoon representation of two monomers of Dg$_{GH172c}$ which

form the active site, with one monomer in pink and the other in purple. The active site residues and coordinated calcium ions are highlighted by a black dashed box. **E)** Proposed active site residues of Dg_{GH172c}. Residues are shown as sticks with carbon atoms colored pink and purple for different subunits. Calcium ions are shown as green spheres and coordinated by D36', D235 and T251.

The proposed active site of Dg_{GH172c} is formed of residues D36, Y152, W230, E233, D235, T251, E254, D255, and W261 (**Figure 3D**). The proposed catalytic residues identified in the characterization of Bd_{GH172} are conserved in $Dg_{GH172c,}$ (Bd/Dg: E270/254, catalytic nucleophile; E291/233, acid/base) and other members of the GH172 family (**Figure S6, S8**). E233 and E254 are 5.4 Å apart in Dg_{GH172c}, which is consistent with a retaining mechanism of enzyme catalysis, as shown for Bd_{GH172}[23]. Ca^{2+} ions are observed in the active sites of Dg_{GH172c} (contains two Ca^{2+}) and Bd_{GH172} (contains one Ca^{2+}). The calcium ions in Dg_{GH172c} are coordinated by D235, which is not conserved in Bd_{GH172}. Instead, at the equivalent position, Bd_{GH172} has N272. A particular feature of characterized GH172 family members is the formation of the active site by two monomers, an uncommon feature of GHs, but on that has been observed in *E. coli* LacZ β-galactosidase[30].

Dg_{GH172c} catalysis is driven by conserved glutamate residues

To gain insight into the functional roles of the residues in the active site of Dg_{GH172c}, the proposed catalytic carboxylate residues were mutagenized to alanine, generating variants E233A, E254A and D225A. Dg_{GH172c}-E233A and Dg_{GH172c}-D225A had no detectable activity and there was a greater than 10^5-fold reduction in activity for Dg_{GH172c}-E254A. Our data support the assignment of E254 and E233 as catalytic residues (acid/base or nucleophile) and highlight an important role for the adjacent D255 residue. None of the variants displayed a change in oligomerization state suggesting they do not possess structural roles (**Figure S9**).

DUF4185 is widely distributed throughout bacterial species

324   In Bacteroidetes, the degradation of a target polysaccharide is typically a multi-

step process whereby oligosaccharides are generated and subsequently cleaved into

326   their monosaccharide constituents, encoded within co-transcribed operons. We

reasoned that the generation of D-arabinofuranose oligosaccharides was likely

328   achieved by proteins that have no annotated function and so focused our attention on

the DUF4185 proteins. To investigate the conservation of these genes across different

330   organisms, we constructed a phylogeny of DUF4185 proteins. This revealed that they

are common within actinobacteria, especially amongst the Actinomycetota (**Figure 4**

332   **and Figure S10**), in *Bacteroides* species, *Myxococcus* and in the lysis cassette of

some actinobacteriophage (**Figure S11**) and the predacious bacterium M. amalyticus*.*

Figure 4: Phylogeny of the DUF4185 enzyme family. Unrooted ML phylogeny (LG

338   model with empirical base frequencies, invariable sites and the discrete gamma

model) of DUF4185 family sequences. Branches with greater than 75% bootstrap

340   support (100 replicates) are highlighted in green. Units for tree scale are inferred

substitutions per amino acid residue. Colored rings indicate phylum (inner) and

342   kingdom (outer) taxonomy information for sequences. Stars highlight sequences of interest and are filled for proteins that have been experimentally characterized in this

344   work.

DUF4185 comprises a novel GH family with endo-D-arabinanase activity

Initially, we cloned, expressed, and purified the DUF4185 homologs

348 (HMPREF9455_02480 and HMPREF9455_02481) from *D. gadei* (herein, referred to

as Dg$_{GH4185a}$ and Dg$_{GH4185b}$, respectively). When incubated with mycobacterial AG and

350 then analyzed by HPAEC, these enzymes produced a banding pattern characteristic

of an endo-acting GH (**Figure S12A**), consistent with endoarabinanase activity. To

352 assess the breadth of activity for DUF4185 enzymes we selected additional candidate

enzymes from each of the major lobes of the global phylogeny (**Figure 4**).

354 Recombinant proteins were produced from several bacterial lineages and a phage

capable of infecting Gram-positive bacteria *Gordonia* of the order Mycobacteriales.

356 These included *Myxococcus xanthus* (Myxo$_{GH4185}$), M. amalyticus*,* and *Gordonia*

Phage GMA6 (Phage$_{GH4185}$)*.* Where the DUF4185 domain sat within a larger gene

358 containing several other large domains we produced truncated variants containing

only the DUF4185 domain due to low solubility of the multi-domain proteins. All

360 DUF4185 constructs except that from M. amalyticus yielded soluble protein. As shown

in **Figure 5** and **Figure S12**, when incubated with mycobacterial AG all these enzymes

362 possessed endo-D-arabinanase activity with varying product profiles suggesting

differences in enzyme specificity. Some of the enzymes were also active against the

364 linear α-1,5-D-arabinofuranose oligosaccharides derived from *P. aeruginosa* PA7 pili,

consistent with activity against the major linkage of mycobacterial D-arabinan (**Figure**

366 **S12**). The discovery of endo-D-arabinanase activity in proteins from organisms outside

of the Actinomycetota is an interesting observation, however its presence in bacterial

368 predators is consistent with the essentiality of this polymer for mycobacterial viability.

This sugar motif has been reported in a small number of LPS structures, providing a

370 possible explanation for the presence of this enzymatic activity in the gut

microbiota[31,32]. Furthermore, some corynebacterial species are found as commensals

372 of the human oral and gut microbiome[33].

Mycobacterial DUF4185s are endo-D-arabinofuranases

Given the importance of D-arabinan to mycobacterial viability and immunology we next sought to understand the biochemical function of DUF4185 proteins from

representative mycobacteria. As shown in **Figure 4** and **Figure S10,** mycobacteria produce at least two DUF4185s that fall into distinct phylogenetic groupings. In *M.*

tuberculosis these are Rv1754c and Rv3707c. Beyond these two conserved DUF4185 genes, some species additional DUF4185 members. For example, many

Mycobacterium abscessus strains encode at least three distinct members whilst *M. smegmatis* mc^{2}155 encodes five (MSMEG_4352, 4360, 4365, 2107 and 6255). Based

on sequence analysis, MSMEG_2107 and MSMEG_6255 are homologs of Rv1754c and Rv3707c, respectively, while the remainder show greater diversity (**Figure S10**).

A distant homolog of Rv3707c from *Mycobacterium abscessus*, Ga0069448_1118 (hereafter referred to as *Mab*$_{4185}$), was readily produced in soluble form and in good

yield[34]. Despite low sequence identity (33.5%) (Figure S13) to the *D. gadei* enzymes, the TLC and HPAEC profiles of mycobacterial AG digested by Mab$_{4185}$ indicate that it

also possesses endo-D-arabinofuranase activity (**Figure 5**).

Figure 5: DUF4185 proteins are endo-D-arabinofuranases that cleave mycobacterial AG and LAM. DUF4185 enzymes were incubated with 2 mg·ml^{-1} AG (**A** and **C**) or 2 mg· ml^{-1} LAM (**B** and **D**) for 16 hours as described in Materials and Methods. Samples were analyzed by High-performance anion-exchange chromatography/pulsed amperometric detection (HPAEC-PAD) on a Dionex ICS-6000w with CarboPac Pa300 column, 100 mM NaOH 20 min isocratic elution followed by a 0-60% 500 mM sodium acetate gradient over 60 minutes. A ladder of α-1,5-L-arabino-oligosaccharides (25 μM) derived from plant arabinogalactans was used as a standard. In panel C the chromatogram of this ladder has been scaled on the *y*-axis by a factor of 0.2 for clarity of presentation.

400   Encouraged by this success with Mab_{4185} but recognizing it has limited

sequence identity with either of the *M. tuberculosis* proteins (17% and 16.4% identical

402   to Rv3707c and Rv1754c respectively over the entire protein length), we sought to

study Rv3707c and Rv1754. However, despite our best efforts we were unable to

404   produce usable amounts of soluble Rv3707c and Rv1754. A previous report

highlighted that Rv3707c is secreted, but it lacks a discernible signal peptide[35]. We

406   reasoned that the instability of Rv3707c may be due to incorrect annotation of the start

site in the *M. tuberculosis* H37Rv genome, given that mycobacteria frequently use

408   TTG and GTG start codons in addition to the canonical ATG. We re-evaluated the

genomic context of the protein and identified several possible in-frame N-terminal

410   extensions of the gene (**Figure S14**). We compared these potential N-terminal

extensions to alignments of DUF4185 proteins that possessed a second, N-terminal

412   domain to identify sequence motifs that are conserved; in many cases a proline-rich

region was observed, which is also found in the potential N-terminal extensions of

414   Rv3707c (**Figure S14**). Furthermore, by extending the N-terminus of the protein, a

putative signal peptide can be predicted by SignalP 6.0 (**Figure S14**). While the

416   precise start-site is uncertain, the likely cleavage point for the signal peptide could be

confidently assigned. Therefore, we designed an expression construct with the

418   putative signal peptide removed, but the remainder of the proline-rich N-terminus

intact. This produced a reasonably stable and soluble Rv3707c protein in good yield

420   that digested AG to give products consistent with a hepta- or hexa-saccharide, and

was also active on LAM (**Figure 5C/D**). While we were unable to produce soluble

422   Rv1754c, we successfully produced the *M. smegmatis* homolog (MSMEG_2107),

which demonstrated activity against LAM, but not AG (**Figure 5C/D and Figure S12**).

424   *DUF4185 family members from different clades have distinct substrate specificities*

To further probe the substrate specificity of these enzymes and elucidate the function of MSMEG_2107, we incubated each of the DUF4185 enzymes (Dg$_{GH4185a}$, Dg$_{GH4185b}$, Mab$_{GH4185}$, Phage$_{GH4185}$, Myxo$_{GH4185}$, MSMEG_2107, and Rv3707c) with: LAM (**Figure S12B**); purified D-galactan (**Figure S12C**); and pilin oligosaccharides from *P. aeruginosa* PA7 (**Figure S12D**). The majority of the enzymes displayed higher activity against AG than LAM. Conversely, MSMEG_2107 was unique in displaying a preference for LAM with no detectable activity when incubated with AG under similar conditions. To confirm that the enzymes only degraded D-arabinan and not D-galactan we tested their activity with D-galactan but did not observe any product formation (**Figure S12C**). Given that branched D-arabinan is the only conserved structure within both AG and LAM, we conclude the DUF4185 family are endo-D-arabinofuranases. We note that only a subset of the DUF4185 enzymes were active against pili from *P. aeruginosa* PA7. As these oligosaccharides are relatively short and linear, this suggests specific enzyme subsite occupancy for activity, which leads to production of arabinose for some enzymes; by contrast arabinose production was not observed upon digestion of branched, polymeric AG.

We next assessed whether DUF4185 endo-D-arabinofuranases can cleave AG in the context of an intact cell wall. We utilized metabolic arabinogalactan labelling with the azide-modified lipid-linked Ara*f* donor 5-AzFPA and then fluorescently labelled intact bacteria with DBCO-conjugated AF647 using click chemistry (**Figure S16**)[36-38]. Both Mab$_{GH4185}$ and Rv3707c released fluorescently labelled material from cell wall, with greater activity and a greater range of products formed by the former enzyme, as observed by HPAEC-PAD. Likewise, treatment with GH172$_{Noc}$ led to release of fluorescent products, supporting the conclusion that both groups of enzymes can cleave AG within the mycobacterial cell wall.

Rv3707c has a highly flexible structure and belongs to the beta-propeller superfamily of glycoside hydrolases

Attempts were made to crystallize all the DUF4185 homologs assayed above. *M. tuberculosis* Rv3707c yielded crystals that diffracted to a resolution of 2.4 Å. Using

the AlphaFold structural prediction as a search model for molecular replacement, we solved a partial structure. As shown in **Figure 6A**, Rv3707c possesses a beta-

propeller structure, similar to several other glycoside hydrolase families,[39] with each blade consisting of 3 antiparallel beta sheets organized radially around a central pore.

Poor or no density was obtained for amino acid residues 1-9, 22-38, 60-69, 85-86, 285-300, and 325-337, which are located on loops surrounding the predicted active

site. The lack of density suggests significant structural flexibility in these regions, consistent with variations in AlphaFold predictions for both Rv3707c and Rv1754c

(**Figure 6, S15, S17**).

    Multi-sequence alignments of DUF4185 homologs (**Figure S13**) identified

three nearly invariant acidic residues in the active site: Asp39, Asp56, and Glu240 (**Figure 6B).** Comparing our experimental structure to the AlphaFold prediction model,

the loops that contained the conserved aspartate residues in the proposed active site deviate in their positions (**Figure 6B and Figure S17**). The conserved D56 points

away from the active site in our experimental structure, likely in a catalytically incompetent position, while the D39 lies in a similar, but distinct position to that of the

AlphaFold model. Together, these observations suggest that the active site may rearrange upon binding of substrate.

The other regions of incomplete density in the experimentally determined 3D structure include loops that are predicted by AlphaFold to be flexible, which is reflected

in the B-factors for the experimental structure (**Figure S17**). These regions are also

predicted by AlphaFold to be variable in other DUF4185 proteins, including Dg$_{GH4185b}$

476   (**Figure 6C**). This apparent flexibility likely enables adaptive binding to a range of

substructures within highly branched mycobacterial D-arabinan, supporting the

478   production of a broad reaction product profile, as seen in enzymatic digests (**Figure**

5).

Figure 6: 3D structure, active site, and AG digestion by DUF4185 endo-D-arabinofuranases. A) 3D structure of Rv3707c determined by X-ray crystallography has a 5-bladed beta-propeller fold. Conserved acidic residues D39, D56 and E240 are indicated in the proposed active site. **B)** Active site details for Rv3707c (X-ray), Rv3707c (AlphaFold) and Dg$_{GH4185b}$ (AlphaFold) with distances between residues indicated in angstroms. **C)** Overall fold of the same enzymes as in **B** with variable surface features highlighted relative to the Rv3707c X-ray structure. Corresponding regions are in the same color in each protein. **D)** Methanolysis of AG catalyzed by Dg$_{GH4185b}$. The production of methyl glucosides in the presence of methanol indicates a retaining mechanism. **E)** Dg$_{GH4185b}$-WT, Dg$_{GH4185b}$-D149A and Dg$_{GH4185b}$-D167A were incubated at 1 µM with 2 mg ml^{-1} AG for 16 hours. No activity was observed for the D188A or D170A mutants.

DUF4185 family are anomer-retaining enzymes

Glycoside hydrolases can hydrolyze the anomeric linkage through either reversion or retention. Inclusion of a simple alcohol, such as methanol, in an enzymatic

digest can be used to identify retaining enzymes, as they may afford methylated glycosides[40]. Addition of methanol to AG digests by $Dg_{GH4185b}$ afforded methyl

arabinoside, thereby demonstrating a retaining mechanism (**Figure 6D**). Three conserved carboxylate residues are present in the active site (**Figure 6B**). Using site-

directed mutagenesis, we varied these carboxylate residues in $Dg_{GH4185b}$ to generate $Dg_{GH4185b}$-D170A and $Dg_{GH4185b}$-D188A and $Dg_{GH4185b}$-E381A derivatives. Activity was

broadly retained for glutamate substitution, but no activity was observed for the two aspartate mutants (**Figure 6E**). These data support a retaining mechanism for the

DUF4185 family of enzymes and assignment of D170/D39 in $Dg_{GH4185b}$ (and D188/D56 in Rv3707c) as the catalytic residues corresponding to acid/base and nucleophile.

Figure 7. Mycobacterial arabinogalactan-degrading enzymes discovered in this study and their substrates. Each enzyme or enzyme family is listed with its identified function and the mycobacterial cell wall it acts upon. Colored in structures are those for which we have presented experimental structural data, black and white are surface representations of AlphaFold models. MA – mycolic acids; L – linker unit; PG – peptidoglycan; PIMs – phosphatidylinostol mannosides.

Discussion

Endo-D-arabinanase activity was first reported more than 50 years ago, but despite the widespread availability of mycobacterial genomic tools and -omics technologies, these enzymes have escaped identification[22,41]. We have mined the human gut microbiome and leveraged evolutionary conservation to identify these enzymes in mycobacteria, along with new exo-D-arabinofuranosidases and exo-D-galactofuranosidases (**Figure 7**). We reasoned that the abundance of Mycobacteriales in environmental niches in addition to the availability of D-arabinofuranose polymers in organisms such as *P. aeruginosa* PA7 and corynebacteria meant that the capacity to degrade this carbohydrate was likely to be encoded in the human gut microbiota[42,43], allowing identification by metabolic methods with arabinogalactan as sole carbon source.

Initially, we identified D-galactofuranase activity in a gut microbe organism, whose presence may reflect the widespread occurrence of D-Gal*f* in the LPS of some

gram-negative bacteria of the human gut microflora[44]. One of the enzymes we identified, BACFIN_04787, is the founding member of a new glycoside hydrolase

family. The galactan degradation pattern exhibited by BACFIN_04787 is consistent with an exo-acting, non-specific enzyme and the product pattern of BACFIN_08810 is

consistent with an exo-acting enzyme that can cleave either the β-1,5 or β-1,6 linkage within D-galactan. Further characterization of these enzymes will enable detailed

analysis of mycobacterial galactan, whose chain length was recently shown to be important in the biology of these organisms[4].

Identifying organisms that can grow on D-arabinan is complicated by the presence of D-galactan in AG. To overcome this, we used the newly discovered D-

galactofuranase to generate enriched D-arabinan, which we used to identify the Bacteroides *D. gadei*, as a potent D-arabinan degrader. The PULs associated with this

ability contained uncharacterized DUFs as well as DU2961 genes corresponding to family GH172. The GH172 proteins were shown to encode exo-D-arabinosidases.

During the preparation of this manuscript, members of family GH172 were independently shown to exhibit exo-D-arabinofuranosidase activity by Kashima *et al.*,

consistent with our data[23]. The GH172 enzyme from *D. gadei* shares a similar structure to the enzyme described by Kashima *et al*; however, it lacks the large C-terminal

extensions observed in Bd$_{GH172}$ which were postulated by the authors to hold the hexameric protein together. SEC-LS analysis of a range of family GH172 proteins

reveals diverse quaternary structures of multimers up to dodecamers. How this structural diversity is connected to function is as-of-yet unexplored. It has been

suggested that family GH172 proteins and phage capsid proteins may be ancestrally related, raising the possibility for diversification of function within this family[45].

The presence of genes encoding DUF4185 and GH172 enzymes in M. amalyticus suggests that they contribute to its epibiotic lifestyle on host *Gordonia* spp. through feeding on arabinogalactan as a carbon source. The $Myc_{DUF4185}$ enzyme is predicted to be a surface located lipoprotein, while Myc_{GH172} is predicted to not be secreted. This localization is consistent with a model whereby M. amalyticus releases oligosaccharides from the surface of *Gordonia*, internalizes them, and then cleaves them to monosaccharides in the cytoplasm. An alternative explanation is that the DUF4185 and GH172 enzymes locally remodel the cell wall of *Gordonia*, enabling access to cellular contents. The biological role of GH172 enzymes in *Bacteroides* remains unclear. While some may be involved in α-fructan degradation, those associated with D-arabinan PULs are more likely targeted at either glycans derived from Actinomycetota or organisms such as *P. aeruginosa* PA7.

566   DUF4185 family of enzymes include the first known endo-D-arabinofuranases, though it is not the first time this activity has been reported. In 1972 Kotani and colleagues reported the isolation of "mixed D-arabinanase activity" in an extract from an unnamed Gram-positive soil microbe, referred to as the "M-2" fraction[19]. This extract possessed endo-activity and released a wide range of products from mycobacterial cell walls, although the specific protein responsible was not identified. Since then, similar impure enzyme cocktails have contributed to numerous studies on AG and LAM[41]. We have now identified broadly conserved endo-D-arabinofuranases from the Mycobacteriales, which we propose to re-name to AraH1 (Rv1754c) and AraH2 (Rv3707c). The availability of well-characterized enzymes with defined activities should support more detailed studies of mycobacterial cell wall polymers.

A review of functional screens of knockout libraries of *M. tuberculosis* highlights an important role for AraH2 in pathogenesis. AraH2 was identified in a screen for proteins with non-canonical signal sequences[35], likely because of mis-annotation of its signal peptide as demonstrated by our work. In that study, Perkowski and colleagues reported that a transposon mutant in AraH2 was severely defective for replication in macrophages. A separate study identified AraH2 as important for control of phagosome acidification, where it was the second most enriched mutant in an acidified phagosome screen[46]. A homologue of AraH2, PEG_1752, was shown to be upregulated in the phylogenetically related *Mycobacterium llatzerense* upon infection of the amoeba *Acanthamoeba castellanii*[47]. These data point to a role for AraH2 in phagosome survival, and by extension of D-arabinan remodeling in mycobacterial pathogenesis.

The role of AraH1 remains more elusive. The gene is broadly conserved amongst mycobacteria, and our biochemical data suggests this enzyme class is active against LAM, but not AG. This genomic locus is a frequent site of IS6110 element insertion[48], and interruption of this gene could cause LAM structural variability amongst circulating strains of *M. tuberculosis*. It is possible that AraH1 and AraH2 have partially overlapping substrate specificity and the former can partially compensate for the loss of the latter. As well, these proteins may have specific roles under narrowly defined conditions. Analogously, peptidoglycan-lytic enzymes with seemingly redundant reaction specificities are encoded in most bacteria, and lack notable phenotypes for their loss under most growth conditions. However, screens at a range of pH values have identified specific functions for these enzymes[49].

To conclude, we have unearthed a new enzymatic toolkit for the degradation of mycobacterial arabinogalactan that should find utility in the study of the structure and

602   function of this important polymer (**Figure 7**). The functional annotation of these genes will support future investigations of the role of D-arabinan structural modulation in

604   mycobacterial biology, mycobacteriophage infection and interbacterial predation.

Tables

606 **Table 1: Activity of GH172 enzymes against synthetic and bacterial substrates**

Protein	pNP-α-D-Ara$f$	AG	LAM	Pili
Dg$_{GH172a}$	✓	✓	✓	✓
Dg$_{GH172b}$	weak	✓	✓	weak
Dg$_{GH172c}$	✓	✓	✓	✓
Noc$_{GH172}$	✓	✓	✓	✓
Myc$_{GH172}$	✓	✓	✓	weak

608 **Table 2: k_{cat}/K_M of GH172 enzymes for pNP-α-D-Ara$f$ and AG**

Protein	pNP-D-Ara$f$
Dg$_{GH172a}$	n/a
Dg$_{GH172c}$	2.9 x 10^4 ± 4 x 10^2 min^{-1} M^{-1}
Noc$_{GH172}$	1.2 x 10^5 ± 2 x 10^3 min^{-1} M^{-1}

Table 3: SEC-LS oligomerization state of GH172 enzymes

Protein	Observed MW (Da)	Oligomerization state	Theoretical MW of oligomer (Da)
Dg$_{GH172c}$	260,829	hexamer	266,952
Noc$_{GH172}$	123,934	trimer	120,507
Dg$_{GH172a}$	515,568	dodecamer	519,438
Dg$_{GH172b}$	148,350	dimer	146,512
Myc$_{GH172}$	192,623	trimer	202,115

Methods

Bacterial strains and growth conditions

Bacteroidetes sp. were grown on a 2x defined minimal media (**Table S3**) under anaerobic conditions at 37 °C over 24 hours to assay growth on various carbon sources, including arabinogalactan. Strains used were *Bacteroides caccae* ATCC 43185, *B. cellulosilyticus* DSM 14838, *B. dorei* DSM 17855, *B. finegoldii* DSM 17565, *B. intestinalis* DSM 17393, *B. nordii* CL02T12C05, *B. ovatus* ATCC 8483, *B. thetaiotaomicron* VPI-5482, *B. vulgatus* ATCC 8482, *B. xylanisolvens* XB1A, *Dysgonomonas gadei* ATCC BAA-286, *D. mossii* DSM 22836, *Parabacteroides gordonii* DSM 23371, *P. johnsonii* DSM 18315. *Escherichia coli* and *Pseudomonas aeruginosa* PA7 ATCC 15692 strains were grown in lysogeny broth at 37 °C (unless otherwise specified). *Mycobacterium smegmatis* mc^{2}155 ATCC 19420 was grown in Tryptic soy Broth at 37 °C with agitation.

RNA sequencing

B. cellulosilyticus was cultured in defined media (supplementary table 3) containing 5 mg ml^{-1} AG or glucose, in triplicate 5 ml cultures. Cells were harvested at mid-log phase and stored in RNA protect (Qiagen). RNA was purified with the RNAeasy Kit. Prior to library preparation, rRNA was depleted using the Pan-Prokaryote riboPOOLs kit (siTOOLs Biotech). In brief, 1 µg of total RNA was incubated for 10 min at 68 °C and 30 min at 37 °C with 100 pmol of rRNA-specific biotinylated DNA probes in 2.5 mM Tris-HCl pH 7.5, 0.25 mM EDTA, and 500 mM NaCl. DNA-rRNA hybrids were depleted from total RNA by two consecutive 15 min incubations

642   with 0.45 mg streptavidin-coated magnetic Dynabeads MyOne C1 (ThermoFisher

Scientific) in 2.5 mM Tris-HCl pH 7.5, 0.25 mM EDTA, and 1 M NaCl at 37 °C. The

644   rRNA-depleted RNA samples were purified using the Zymo RNA Clean &

Concentrator kit combined with DNase treatment on a solid support (Zymo Research).

646   cDNA libraries were prepared using the NEBNext Multiplex Small RNA Library Prep

kit for Illumina (NEB) in accordance with the manufacturers' instructions.

648   Library preparation and sequencing took place at the Earlham Institute, and

were processed by Newcastle University Bioinformatics Support Unit. Briefly, raw

650   sequencing reads were checked using Fast QC, reads were mapped to *Bacteroides*

cellulosilyticus DSM 14838 (GCA_000158035) downloaded from Ensembl (assembly

652   ID ASM15803v1). Reads were quantified against genes contained in the Ensembl

annotation using featureCounts from the Rsubread package[50]. Counts were

654   normalized by Trimmed Median of Means (TMM) as implemented in DESeq2, and

differentially expressed genes determined according to a Negative Binomial model as

656   per DESeq2.

Proteomic analysis of *D. gadei*

658   *Sample preparation*

Dysgonomonas gadei cells were suspended in 5% sodium dodecyl sulfate

660   (SDS) in 50 mM triethylammonium bicarbonate (TEAB) pH 7.5. The samples were

subsequently sonicated using an ultrasonic homogenizer (Hielscher) for 1 minute. The

662   whole-cell lysate was centrifuged at 10,000 x *g* for 5 min to remove cellular debris.

Protein concentration was determined using a bicinchoninic acid (BCA) protein assay

664   (Thermo Scientific). A total of 20 μg protein was used for further processing. Proteins

were reduced by incubation with 20 mM tris(2-carboxyethyl)phosphine for 15 min at

666   47 °C, and subsequently alkylated with 20 mM iodoacetamide for 30 minutes at room

temperature in the dark. Proteomic sample preparation was performed using the

668   suspension trapping (S-Trap) sample preparation method [51], as recommended by the

supplier (ProtiFi™, Huntington NY). Briefly, 2.5 µl of 12% phosphoric acid was added

670   to each sample, followed by the addition of 165 µl S-Trap binding buffer (90%

methanol in 100 mM TEAB pH 7.1). The acidified samples were added, separately, to

672   S-Trap micro-spin columns and centrifuged at 4,000 x g for 1 min until the solution has

passed through the filter. Each S-Trap micro-spin column was washed with 150 µl S-

674   trap binding buffer by centrifugation at 4,000 x g for 1 min. This process was repeated

for a total of five washes. Twenty-five µl of 50 mM TEAB containing trypsin (1:10 ratio

676   of trypsin:protein) was added to each sample, followed by proteolytic digestion for 2 h

at 47 °C using a thermomixer (Eppendorf). Peptides were eluted with 50 mM TEAB

678   pH 8.0 and centrifugation at 1,000 x g for 1 min. Elution steps were repeated using

0.2% formic acid and 0.2% formic acid in 50% acetonitrile, respectively. The three

680   eluates from each sample were combined and dried using a speed-vac before storage

at -80 °C.

682   *Mass Spectrometry*

Peptides were dissolved in 2% acetonitrile containing 0.1% trifluoroacetic acid,

684   and each sample was independently analyzed on an Orbitrap Fusion Lumos Tribrid

mass spectrometer (Thermo Fisher Scientific), connected to an UltiMate 3000

686   RSLCnano System (Thermo Fisher Scientific). Peptides were injected on a PepMap

100 C_{18} LC trap column (300 µm ID × 5 mm, 5 µm, 100 Å) followed by separation on

688   an EASY-Spray nanoLC C_{18} column (75 µm ID × 50 cm, 2 µm, 100 Å) at a flow rate of

250 nl/min. Solvent A was water containing 0.1% formic acid, and solvent B was 80%

690   acetonitrile containing 0.1% formic acid. The gradient used for analysis was as follows:

solvent B was maintained at 2% for 5 min, followed by an increase from 2 to 35% B in

692   120 min, 35-90% B in 0.5 min, maintained at 90% B for 4 min, followed by a decrease

to 3% in 0.5 min and equilibration at 2% for 10 min. The Orbitrap Fusion Lumos was

694   operated in positive-ion data-dependent mode. The precursor ion scan (full scan) was

performed in the Orbitrap in the range of 400-1,600 *m/z* with a resolution of 120,000

696   at 200 *m/z*, an automatic gain control (AGC) target of 4×10^5 and an ion injection time

of 50 ms. MS/MS spectra were acquired in the linear ion trap (IT) using Rapid scan

698   mode after high-energy collisional dissociation (HCD) fragmentation. An HCD collision

energy of 30% was used, the AGC target was set to 1×10^4 and dynamic injection time

700   mode was allowed. The number of MS/MS events between full scans was determined

on-the-fly to maintain a 3 s fixed duty cycle. Dynamic exclusion of ions within

702   a ± 10 ppm *m/z* window was implemented using a 35 s exclusion duration. An

electrospray voltage of 2.0 kV and capillary temperature of 275 °C, with no sheath and

704   auxiliary gas flow, was used.

    All mass spectra were analyzed using MaxQuant 1.6.12.0[52], and searched

706   against the *Dysgonomonas gadei* ATCC BAA-286 proteome database downloaded

from Uniprot (accessed 09.01.2020). Peak list generation was performed within

708   MaxQuant and searches performed using default parameters and the built-in

Andromeda search engine[53]. The enzyme specificity was set to consider fully tryptic

710   peptides, and two missed cleavages were allowed. Oxidation of methionine, N-

terminal acetylation and deamidation of asparagine and glutamine were allowed as

712   variable modifications. Carbamidomethylation of cysteine was allowed as a fixed

modification. A protein and peptide false discovery rate (FDR) of less than 1% was

714   employed in MaxQuant. Proteins that contained similar peptides and could not be

differentiated based on MS/MS analysis alone were grouped to satisfy the principles

716   of parsimony. Reverse hits, contaminants, and proteins only identified by site

modifications were removed before downstream analysis.

718   **Generation of expression constructs**

Unless stated otherwise genes were purchased as codon-optimized constructs

720   from Twist Biosciences. *D. gadei* and *B. finegoldii* genes were cloned from genomic

DNA using standard restriction cloning methods, followed by ligation into pET28a

722   vectors and transformation into TOP10 cells (Novagen) with subsequent sequencing

of selected purified recombinant plasmids by Eurofins for confirmation.

724   **Protein Expression and Purification**

726   *Expression and purification of Dg$_{GH4185a}$, Dg$_{GH4185b}$, Phage$_{GH4185}$, Myxo$_{GH4185}$, Dg$_{GH172a}$, Dg$_{GH172b}$, Dg$_{GH172c}$, and Myc$_{GH172}$*

728
Recombinant proteins were expressed in competent *E. coli* Tuner cells

730   (Novagen) using pET28a vectors generated as above. Cells were grown in LB media

at 37 °C with shaking, until turbidity reached an OD$_{600}$ of ~0.6, wherein expression was

732   induced with 0.2 mM IPTG and cells were further cultured for 16 hours at 16 °C.

Sonication was used to lyse cells in 20 mM Tris, pH 8.0, 200 mM NaCl.

734   Enzymes were purified using immobilized metal affinity chromatography on

cobalt TALON resin. Proteins were dialyzed into 20 mM HEPES, pH 8.0, 150 mM NaCl

736   buffer dialysis (Medicell). For crystallography proteins were purified further via size-

exclusion chromatography (HiLoad 16/600 Superdex 200, GE Healthcare) in 20 mM

738   HEPES, pH 8.0, 150 mM NaCl. Protein purity was ascertained by SDS-PAGE and

protein concentrations were determined using Nanodrop spectroscopy

740   (Thermofisher).

742   *Purification of Rv3707c and MSMEG_2107*

For protein Rv3707c and MSMEG_2107 expression, an aliquot of competent

744   BL21 DE3 *Escherichia coli* was transformed with plasmid and plated on LB agar

supplemented with 50 μg/mL kanamycin. One plate of bacteria was scraped to

746   inoculate 1 L of modified Studier's autoinduction media[54]. The bacteria were incubated

at 37 °C with shaking (180 RPM) until OD_{600}= ~0.6, whereupon the flasks were cooled

748   on ice with agitation to 20 °C and then returned shake overnight at 20 °C. After

induction, the cultures were pelleted at 3990 *x g* for 25 min at 4 °C. Pellets were

750   resuspended in sterile PBS and pelleted at 7000 *x g* for 10 min. The supernatant was

removed, and pellets were snap frozen in liquid nitrogen and stored at -20 °C until

752   preparation.

754   Rv3707c was purified by suspension of one pellet in cold lysis buffer (25 mM

HEPES pH8; 400 mM NaCl; 5% glycerol; 50 mM L-arginine; 50 mM L-glutamic acid; 1

756   mM beta-mercaptoethanol). 1 mg ml^{-1} deoxyribonuclease I from bovine pancreas

(Sigma-Aldrich) was added to cell slurry and incubated for 30 minutes. Cells were

758   lysed by three passages through a French pressure cell. Insoluble debris was then

pelleted by centrifugation at 40,000 *x g* for 40 minutes (4 ºC). The supernatant was

760   then processed by immobilized metal affinity chromatography (IMAC) on a gravity

column containing 2 mL bed volume of cOmplete His Tag purification resin (Roche).

762   After loading the lysate, the column was washed with 80 mL of lysis buffer, then eluted

with an imidazole gradient of 50, 100, 250, and 500 mM. Protein-containing fractions

764   were pooled and dialyzed exhaustively against three liters of dialysis buffer (25 mM

HEPES pH8; 400 mM NaCl; 5% glycerol; 50 mM L-arginine; 50 mM L-glutamic acid; 2

766   mM dithiothreitol). The crude protein was concentrated to a final volume of 0.5 mL on

a 30k Da molecular weight cutoff Pierce protein concentrator (Thermo Scientific). This

768   fraction was then further purified by size exclusion chromatography on an AKTA Prime

system with a SuperDex 26/600 S200 column in the above dialysis buffer before being

770 concentrated. The protein was always used freshly prepared.

772 *Purification of Noc$_{GH172}$ and Mab$_{GH4185}$*
One plate of BL21-DE3 transformed with an appropriate plasmid was used to

774 inoculate 1L Terrific Broth supplemented with kanamycin. The culture was grown to

an OD$_{600}$ of 0.6 and induced with 0.25 mM IPTG and grown overnight at 20 ºC. After

776 harvest of biomass as in purification of Rv3707c, cell pellets were resuspended in a

buffer containing 25 mM HEPES; 40 mM NaCl, lysozyme, and DNAse I. Subsequent

778 purification steps were identical to those in Rv3707c, but in the above, simpler buffer,

omitting lysozyme and DNAse I.

Crystallography

782 *Crystallography of Dg$_{GH172c}$*
Crystallization of Dg$_{GH172c}$ sample at 10 mg ml^{-1} was screened using

784 commercial kits (Molecular Dimensions and Hampton Research) with vapor diffusion

sitting drop method. Crystals formed in a buffer containing 0.1 M MOPS/Sodium

786 HEPES pH 7.5, 0.12 M alcohol and 30% EDO_P8K over a period of 2 weeks. Crystals

were harvested and flash cooled in liquid nitrogen. X-ray diffraction data were collected

788 at the synchrotron beamline I24t Diamond light source (Didcot, UK) at a temperature

of 100 K. The data were integrated using XDS[55] via XIA2 [56] and scaled with Aimless[57].

790 The space group was confirmed with Pointless. The phase problem was solved by

molecular replacement using Phaser and PDB model 4KQ7 as search model from

792 *Bacteroides uniformis*[58]. Subsequently, the structure was auto built with CCP4build

on CCP4cloud[59]. The model was improved by rounds of manual building with COOT

794 and refinement with Refmac[60,61]. The final model was validated with Molprobity[62] and

COOT[60].

Crystallography of Rv3707c

798   Rv3707c was concentrated to 7 mg ml^{-1} and loaded into a sitting drop

crystallization tray (Clover) using JCSG-plus crystallization screen (Molecular

800   Dimensions) at a ratio of 2 µL protein 2 µL reservoir solution. Crystals were formed in

a buffer containing 0.8 M succinic acid pH 7 over a period of 1 month. The crystals

802   were cryoprotected in a solution of 0.8 M succinic acid pH 7 with 30% glycerol and

flash cooled in liquid nitrogen. X-ray diffraction data were collected at beamline I04 of

804   Diamond light source, Oxford. Data were auto-processed using Xia2 and general file

manipulations were performed using the CCP4 suite of programmes[56,59]. Data were

806   phased by molecular replacement using the Alphafold structural prediction of Rv3707c

as a search model using PHASER (TFZ of 21.1)[58]. The structure was then auto built

808   in PHENIX with rounds of refinement carried out by PHENIX-refine and manually in

COOT[60,63].

810   **Summary of methods for phylogenetic analyses**

Two alignments were used to infer respectively a global phylogeny for members

812   of the PF13810 family and a more restricted phylogeny focusing on close relatives to

the functionally characterized proteins. The global alignment was derived from the

814   "Full" Pfam alignment for PF13810 made of 1145 sequences and 1321 aligned sites

(http://pfam.xfam.org/family/PF13810#tabview=tab3). This led to an alignment of 753

816   sequences and 179 aligned sites by: (i) deleting partial sequences that did not include

the catalytic residues or that corresponded to obsolete sequences (nine sequences)

818   and (ii) adding the three sequences from *Mycobacterium abscessus* (strain 4529

available at the Integrated Microbial Genomes & Microbiomes (IMG/M) database:

820   2635695100/Ga0069448_11324, 2635694794/Ga0069448_1118,

2635698039/Ga0069448_113269) and (iii) deleting sites made of a majority of indels.

822   For the restricted alignment 39 complete sequences were aligned, including the seven

proteins investigated in this study - enzymatic characterization and one structure. The

824   sequences were aligned with Clustal Omega using default settings in SEAVIEW

v.4.6.4 [64,65]. Following minor manual adjustments of the alignment the mask function

826   of SEAVIEW was used to selected aligned residues that included conserved blocks of

sequences with no more than two indels leading to 200 aligned sites. The DUF4185

828   alignment is available as Supplementary Dataset 1 for respectively the global and

restricted alignment. IQ-TREE (v.1.6.12) was used to generate maximum likelihood

830   phylogenies using automatic model selection[66]. The selected models were

LG+F+I+G4 for the global alignment and WAG+I+G4 for the restricted alignment using

832   the Bayesian Information Criterion (BIC). Bootstraps (100 replicates) were computed

to assess branch reliability. iTOL (interactive tree of life) was used to generate the

834   figures[67]. The global phylogeny was annotated with taxonomy information derived

from the UniProt database (https://www.uniprot.org/)[68].

836   **Protein structure prediction by AlphaFold 2.0**

    Amino acid sequences were submitted to the Colabfold_advanced.ipynb server

838   for protein structure prediction[69]. Unless otherwise stated the default values were

used, with the max_recycles set to 12. Figure coloring for pLDDT values were

840   generated in ChimeraX 1.4 using the protein structure prediction tool and a custom

palette.

    Timepoint assays

844   To assess enzymatic activity, reactions were initiated containing substrates (in

water) and enzymes (in 20 mM HEPES pH 7.5, 150 mM NaCl, unless otherwise

846   stated) of various concentrations, with 50 mM potassium phosphate buffer pH 7.2 as

a dominant reaction buffer. Reactions were incubated at 37 °C for 16 hours and

848   subsequently boiled to ensure enzymatic cessation. Time point samples were then

analyzed using TLC or IC-PAD.

Porous graphitic carbon chromatography clean-up of Rv3707c and

852   **MSMEG_2107 arabinogalactan hydrolysis assays**

Due to the high concentration of L-arginine and L-glutamic acid in the buffer

854   used for purification of Rv3707c and MSMEG_2107, enzyme assays were unsuitable

for HPAEC-PAD analysis without prior solid phase extraction. To this end, at each

856   timepoint, reaction mixtures were loaded onto a Hypersep Hypercarb SPE cartridge

(Thermo Scientific) which had been washed with acetonitrile and 50% THF in water

858   and exhaustively equilibrated with water prior to loading. Reaction products were then

eluted in 80% acetonitrile in ddH2O with 0.1% trifluoracetic acid (Sigma-Aldrich) and

860   dried by evaporation in a SpeedVac concentrator before being reconstituted in the

original volume of water.

862   **Kinetic analysis of GH172 arabinofuranosidase activity**

Enzymes (100 nM) were incubated with the indicated concentrations of pNP-α-

D-Ara*f* or AG. Assays were performed in technical triplicate at 37 °C in 20 mM HEPES

866   pH 7.5. for pNP, absorbances were measured at 400 nm and graphs were plotted in

GraphPad Prism 9.3.1. For Ag, arabinose concentration was measured by IC-PAD

868   (see below) with reference to a standard curve.

870   **Thin-layer chromatography (TLC)**

Purified proteins were incubated with at 0.1-5 μM (as indicated) with substrates

872   (for methanolysis, methanol was added to the reaction mixture at a final concentration

of 10%) for 16 h at 37 °C to ensure reaction completion (unless otherwise indicated).

874   Using TLC plate aluminum foils (Silicagel 60, 20 x 20, Merck) which were cut to the

desired size (minimum height of 10 cm), these reaction samples were spotted (6 µl,

876   unless otherwise indicated) and allowed to dry. TLC plates were run (twice) in solvent

(1-butanol/acetic acid/ water 2:1:1 (v/v)). Plates were then removed and dried before

878   visualization of sugars was obtained via immersion of TLC plate in Orcinol stain. Plates

were dried and developed through heating between 50 °C and 80 °C.

880   **Ion Chromatography with Pulsed Amperometric Detection (IC-PAD)**

Oligosaccharides from enzymatic polysaccharide digestion were analyzed

882   using a CARBOPAC PA-300 anion exchange column (ThermoFisher) on an ICS-6000

system. Detection enabled by PAD using a gold working electrode and a PdH

884   reference electrode with standard Carbo Quad waveform. Buffer A – 100 mM NaOH,

Buffer B – 100 mM NaOH, 0.5 M Na Acetate. Each sample was run at a constant flow

886   of 0.25 ml·min^{-1} for 100 minutes using the following program after injection: 0-10 min;

isocratic 100% buffer A, 10-70 min; linear gradient to 60% buffer B, 70-80 min; 100%

888   buffer B. The column was then washed with 10 mins of 500 mM NaOH, then 10 min

re-equilibration in 100% buffer A. L-arabino-oligosaccharides (DP = 2-9) obtained

890   commercially (Megazyme) were used as standards at a concentration of 25 µM. Data

were processed using Chromeleon^{TM} Chromatography Management System V.6.8.

892   Final graphs were created using GraphPad Prism 8.0.1.

Purification of mycobacterial arabinogalactan

894   Large scale purification of mycobacterial arabinogalactan was achieved by

established methodologies[70]. In brief, 8 L of mycobacterial culture was grown to mid-

896   exponential phase, cultures were pelleted and resuspended in a minimal volume of

phosphate-buffered saline (VWR) and lysed using an Emulsiflex. The lysate was then

898   boiled in a final concentration of 1% sodium dodecyl sulphate (SDS) and refluxed.

Insoluble material (containing mycolyl-arabinogalactan-peptidoglycan complex) was

900   collected by centrifugation and washed exhaustively with water to remove SDS. The

mycolate layer was removed by saponification by KOH in methanol at 37 °C for 3 days.

902   Cell wall material was then washed repeatedly to remove saponified mycolic acids

with diethyl ether. The phosphodiester linkage between AG and PG was then cleaved

904   by treatment with H_2SO_4 at 95 °C before being neutralized with sodium carbonate. The

resultant solubilized arabinogalactan was collected in the supernatant, dialyzed

906   exhaustively against water and lyophilized (yield= ~22.5 mg·L^{-1}).

Purification of D-arabinan

908   Arabinogalactan (5 mg/mL) was digested in a 10 mL total volume of 25 mM MOPS

buffer pH 7. One µM final concentration of BACFIN_04787 and BACFIN_08810 were

910   added, and the reaction was incubated at 37°C for 48h. An aliquot of the reaction was

analyzed by TLC to verify the hydrolysis of the substrate and galactose release. Then,

912   the sample was dialyzed over-night against 5 L of deionized water using a 1 kDa

membrane, to eliminate residual galactose from the reaction mixture. The sample was

914   then freeze-dried and resuspended in 0.4 mL water. A TLC analysis confirmed the

elimination of the residual galactose from the sample, and this was further confirmed

916   through acid hydrolysis of the resulting D-arabinan to determine total abundance. To

quantify the purity of isolated d-arabinan, an aliquot of both (0.25 mg/mL) d-arabinan

918   and arabinogalactan were treated by acid hydrolysis using 300 mM HCl at 100°C for

1h. Samples were neutralised to pH 7 with NaOH and analysed by HPAEC.

920   Quantitation was based on the migration of standards and the ratio between galactose

and arabinose.

Purification of mycobacterial lipoglycans

924   One liter of mycobacterial culture was grown to mid exponential phase and

pelleted as previously. The pellet was resuspended in 20 ml PBS, 0.1% Tween-80,

926   chilled and lysed by bead-beating. Lysate was transferred to a Teflon-capped glass

tube and vortexed with an equal volume of citrate buffer saturated with phenol (Sigma-

928   Aldrich), and heated to 80 °C for 3 h, vortexing every hour. A biphase was generated

by centrifugation at 2000 RPM at 10 °C, and the upper aqueous phase transferred to

930   a fresh glass tube and hot phenol wash repeated twice more. The resultant protein

free glycan mixture was dialyzed exhaustively against tap water overnight to remove

932   trace phenol and lyophilized, yielding 34 mg of crude lipoglycans (LAM, LM, PIMS) per

liter of culture.

Pseudomonas aeruginosa pilin oligosaccharide extraction.

936   Pilins were purified as described by Burrows and colleagues, with some

modifications[42]. Briefly, _Pseudomonas aeruginosa_ PA7 were streaked out in a grid

938   pattern onto LB agar plates and grown for 24 hours at 37 °C. Cells were then scraped

from all plates using sterile cell scrapers and resuspended in 4 ml of sterile phosphate-

940   buffered saline (pH 7.4) per plate.

Pili were then sheared from the cell wall by vigorous vortexing of resuspended

942   cells for 2 min. This suspension was centrifuged for 5 min at 6000 $x\,g$. The supernatant

was centrifuged for 20 min at 20,000 $x\,g$. Supernatants were transferred to fresh tubes

944   and $MgCl_2$ was added to give a final concentration of 100 mM. Following inversion of

these tubes to ensure mixing, samples were incubated at 4 °C overnight, allowing

946   precipitation of sheared proteins. Samples were then centrifuged for 20 minutes at

20,000 $x\,g$, yielding a precipitate smudge which was resuspended in 50 mM NH_4HCO_3,

948   pH 8.5 and transferred to fresh Eppendorf tubes. This solution was then dialyzed into

the same buffer to eliminate excess $MgCl_2$.

950   Bradford assays were then performed to assay the mass of protein in the

sample, followed by digestion of the intact protein pilins using proteinase K in a 50:1

952   pilin to enzyme ratio by mass for 24 hours in the presence of 2 mM CaCl$_2$. Glycans

were then purified from digested proteins using porous graphitized carbon

954   chromatography, where sugars were eluted from the column using a twofold

increasing concentration series of a butan-1-ol:H$_2$O gradient from 1:32 to 1:1 using 1

956   mL elutions[71]. Thin-layer chromatography (TLC) of eluates showed various oligomers

of arabinan present in all fractions, all of which were subsequently used as substrates

958   for potential arabinanases.

960   **Synthesis of pNP-α-D-arabinofuranoside**

Para-nitrophenol (pNP)-α/β-D-arabinofuranoside synthesis was achieved

962   following the established procedures [23].

964   **SEC-LS**

Molecular weights were determined by size exclusion chromatography coupled

966   light scattering using an Agilent MDS system with either an Agilent BioSEC 5 1000 Å,

4.6 x 300 mm, 5 µm or GE Superdex 200 5 15 mm columns at appropriate flow rates.

968   Detector offsets were calibrated using a BSA standard and concentrations were

determined by refractive index.

Protein sequence alignment and structure visualization

972   Sequence alignments were performed using Clustal Omega and visualized

using ESPript[64,72]. Figures of protein structures were generated with ChimeraX 1.4[73].

974   **Fluorescent-conjugated mAGP hydrolysis assay**

Fluorescently labeled mycolyl-arabinogalactan-peptidoglycan complex

976   (mAGP) was isolated from *Corynebacterium glutamicum ATCC13032* following

previously reported methods[36-38]. In brief, cells were grown in the presence of 5-

978   AzFPA to saturation, reacted with DBCO-conjugated AF647 and then the mAGP was

isolated. The isolated product was suspended in 2% SDS in PBS and split into the

980   outlined treatment groups in Eppendorf tubes. Samples were pelleted by

centrifugation at 15,000 $x\ g$ for 5 min at 4ºC then washed with PBS once (100 µL).

982   Following this wash, the mAGP was resuspended in 90 µL PBS and enzyme stock

added for a final concentration of 5 µM of each enzyme. Samples were incubated at

984   37 ºC with rotation for 16 h. Following incubation, samples were pelleted by

centrifugation at 15,000 $x\ g$ for 5 min at 4 ºC then washed with PBS twice (100 µL).

986   The pellets were then suspended in 2% SDS in PBS, transferred to a 96-well plate

and the fluorescence emission of each well was then measured on a Tecan Infinite

988   M1000 Pro microplate reader. Monitoring of AF647 fluorescence was achieved by

exciting at 648 nm ± 5 nm and detecting at 671 nm ± 5 nm. Z-position was set to 2

990   mm, and the fluorimeter gain was optimized and then kept constant. Data are reported

in relative fluorescence units (RFU) normalized to PBS treated controls.

992   **Acknowledgements:**
We thank members of the Birmingham mycobacteriology group and the
994   Newcastle glycobiology group for support and discussions. We gratefully acknowledge
Diamond synchrotron source for access to X-ray beamtime.
996

Funding:
998   PJM is supported by a BBSRC David Phillips Fellowship (BB/S010122/1) and a
BBSRC Impact Acceleration Award (1544084). ECL is supported by an Academy of
1000   Medical Sciences Springboard Award (SBF006\1048) and a Royal Society Research
Grant (RGS\R2\202228). ALL is supported by a Wellcome Trust Investigator Award in
1002   Science (209437/Z/17/Z). SJW is supported by the Australian Research Council
(DP210100233, DP210100233). STB is supported by a studentship from the
1004   Wellcome Trust. NPB is supported by a BBSRC studentship (BB/M011186/1). OA-J
is funded by a BBSRC studentship (BB/M011186/1). JM-M was supported by an
1006   Advanced Grant from the European Research Council (Grant no. 322820) awarded to
Harry J Gilbert. AC is funded by Academy of Medical Sciences Springboard
1008   (SBF005\1065) and a Royal Society research grant (RGS\R2\212050). This research
was supported by the National Institute of Allergy and Infectious Disease (R01 AI-
1010   126592) and the NIH Common Fund (U01GM125288).

1012

Author Contributions:

Omar Al-Jourani – Methodology, validation, formal analysis, investigation, data curation, visualization. **Samuel Benedict** – Methodology, validation, formal analysis, investigation, data curation, visualization, resources. **Jennifer Ross** – Methodology, validation, formal analysis, investigation, data curation, writing – original draft preparation, writing – review and editing. **Abigail Layton** – Methodology, validation, resources. **Phillip van der Peet** – Methodology, validation, formal analysis, resources. **Victoria Marando** – Methodology, formal analysis, investigation, visualization, writing – review and editing. **Nicholas P Bailey** – Methodology, validation, formal analysis, visualization. **Tiaan Heunis** – Methodology, formal analysis, resources. **Joseph Manion** – Investigation. **Francesca Mensitieri** – Investigation. **Aaron Franklin** – Investigation, resources. **Javier Abellon-Ruiz** – Investigation. **Sophia L Oram,** – Investigation. **Lauren Parsons** – Investigation. **Alan Cartmell** – Resources, supervision. **Gareth SA Wright** – Resources, supervision. **Arnaud Baslé** – Methodology, validation, writing – review and editing. **Matthias Trost** – Resources, supervision. **Bernard Henrissat** – Resources. **Jose Munoz-Munoz** – Investigation. **Robert P Hirt** – Investigation, resources, writing – review and editing, supervision. **Laura L. Kiessling** – Resources, supervision, funding acquisition, writing – review and editing. **Andrew Lovering** – Methodology, validation, formal analysis, writing – review and editing, supervision, funding acquisition. **Spencer J. Williams** – Resources, validation, formal analysis, writing – review and editing, supervision, funding acquisition. **Elisabeth C. Lowe** – Conceptualization, methodology, validation, formal analysis, data curation, writing – original draft preparation, writing – review and editing, visualization, supervision, funding acquisition. **Patrick J. Moynihan** – Conceptualization, methodology, validation, formal analysis, data curation, writing – original draft preparation, writing – review and editing, visualization, supervision, funding acquisition.

Competing Interests

Drs. Moynihan and Lowe are co-inventors on an unpublished patent application pertaining to some of the enzymes described in this manuscript.

References

1048   1   Abrahams, K. A. & Besra, G. S. Synthesis and recycling of the mycobacterial cell envelope. *Curr Opin Microbiol* **60**, 58-65 (2021).
1050   https://doi.org:10.1016/j.mib.2021.01.012

2   Dulberger, C. L., Rubin, E. J. & Boutte, C. C. The mycobacterial cell envelope
- a moving target. *Nat Rev Microbiol* **18**, 47-59 (2020). https://doi.org:10.1038/s41579-019-0273-7

3   Daffe, M., McNeil, M. & Brennan, P. J. Major structural features of the cell wall arabinogalactans of Mycobacterium, Rhodococcus, and Nocardia spp.
Carbohydr Res **249**, 383-398 (1993). https://doi.org:10.1016/0008-6215(93)84102-c

4   Justen, A. M. *et al.* Polysaccharide length affects mycobacterial cell shape and antibiotic susceptibility. *Sci Adv* **6** (2020).
https://doi.org:10.1126/sciadv.aba4015

Safi, H., Sayers, B., Hazbon, M. H. & Alland, D. Transfer of embB codon 306
1062   mutations into clinical Mycobacterium tuberculosis strains alters susceptibility to ethambutol, isoniazid, and rifampin. *Antimicrob Agents Chemother* **52**, 2027-
1064   2034 (2008). https://doi.org:10.1128/AAC.01486-07

6   Sun, Q. *et al.* Mutations within embCAB Are Associated with Variable Level of
1066   Ethambutol Resistance in Mycobacterium tuberculosis Isolates from China. *Antimicrob Agents Chemother* **62** (2018). https://doi.org:10.1128/AAC.01279-
1068   17

7   Zhao, L. L. *et al.* Analysis of embCAB mutations associated with ethambutol
1070   resistance in multidrug-resistant mycobacterium tuberculosis isolates from China. *Antimicrob Agents Chemother* **59**, 2045-2050 (2015).
1072   https://doi.org:10.1128/AAC.04933-14

8   Brossier, F. *et al.* Molecular Analysis of the embCAB Locus and embR Gene
1074   Involved in Ethambutol Resistance in Clinical Isolates of Mycobacterium tuberculosis in France. *Antimicrob Agents Chemother* **59**, 4800-4808 (2015).
1076   https://doi.org:10.1128/AAC.00150-15

9   Zhang, L. *et al.* Structures of cell wall arabinosyltransferases with the anti-
1078   tuberculosis drug ethambutol. *Science* **368**, 1211-1219 (2020). https://doi.org:10.1126/science.aba9102

1080   10   Winder, F. G. & Collins, P. B. Inhibition by isoniazid of synthesis of mycolic acids in Mycobacterium tuberculosis. *J Gen Microbiol* **63**, 41-48 (1970).
1082   https://doi.org:10.1099/00221287-63-1-41

11   Park, J. T. & Strominger, J. L. Mode of action of penicillin. *Science* **125**, 99-101
1084   (1957). https://doi.org:10.1126/science.125.3238.99

12   Healy, C., Gouzy, A. & Ehrt, S. Peptidoglycan Hydrolases RipA and Ami1 Are
1086   Critical for Replication and Persistence of Mycobacterium tuberculosis in the Host. *mBio* **11** (2020). https://doi.org:10.1128/mBio.03315-19

1088   13   Moynihan, P. J. *et al.* The hydrolase LpqI primes mycobacterial peptidoglycan recycling. *Nat Commun* **10**, 2647 (2019). https://doi.org:10.1038/s41467-019-
1090   10586-2

14   Kalscheuer, R., Weinrick, B., Veeraraghavan, U., Besra, G. S. & Jacobs, W. R.,
Jr. Trehalose-recycling ABC transporter LpqY-SugA-SugB-SugC is essential

for virulence of Mycobacterium tuberculosis. *Proc Natl Acad Sci U S A* **107**, 21761-21766 (2010). https://doi.org:10.1073/pnas.1014642108

15   Yang, Y. *et al.* A hydrolase of trehalose dimycolate induces nutrient influx and stress sensitivity to balance intracellular growth of Mycobacterium tuberculosis. *Cell Host Microbe* **15**, 153-163 (2014).
https://doi.org:10.1016/j.chom.2014.01.008

16   Batinovic, S., Rose, J. J. A., Ratcliffe, J., Seviour, R. J. & Petrovski, S. Cocultivation of an ultrasmall environmental parasitic bacterium with lytic ability against bacteria associated with wastewater foams. *Nat Microbiol* **6**, 703-711
(2021). https://doi.org:10.1038/s41564-021-00892-1

17   Shen, L. *et al.* The endogenous galactofuranosidase GlfH1 hydrolyzes mycobacterial arabinogalactan. *J Biol Chem* **295**, 5110-5123 (2020). https://doi.org:10.1074/jbc.RA119.011817

18   Helbert, W. *et al.* Discovery of novel carbohydrate-active enzymes through the rational exploration of the protein sequences space. *Proc Natl Acad Sci U S A*
116, 6063-6068 (2019). https://doi.org:10.1073/pnas.1815791116

19   Kotani, S., Kato, T., Matsubara, T., Sakagoshi, M. & Hirachi, Y. Inducible enzyme degrading serologically active polysaccharides from mycobacterial and corynebacterial cells. *Biken J* **15**, 1-15 (1972).

20   McNeil, M. R., Robuck, K. G., Harter, M. & Brennan, P. J. Enzymatic evidence for the presence of a critical terminal hexa-arabinoside in the cell walls of
Mycobacterium tuberculosis. *Glycobiology* **4**, 165-173 (1994). https://doi.org:10.1093/glycob/4.2.165

21   Xin, Y. *et al.* Characterization of the in vitro synthesized arabinan of mycobacterial cell walls. *Biochim Biophys Acta* **1335**, 231-234 (1997).
https://doi.org:10.1016/s0304-4165(97)00002-0

22   Dong, X., Bhamidi, S., Scherman, M., Xin, Y. & McNeil, M. R. Development of a quantitative assay for mycobacterial endogenous arabinase and ensuing studies of arabinase levels and arabinan metabolism in Mycobacterium
smegmatis. *Appl Environ Microbiol* **72**, 2601-2605 (2006). https://doi.org:10.1128/AEM.72.4.2601-2605.2006

23   Kashima, T. *et al.* Identification of difructose dianhydride I synthase/hydrolase from an oral bacterium establishes a novel glycoside hydrolase family. *J Biol*
Chem **297**, 101324 (2021). https://doi.org:10.1016/j.jbc.2021.101324

      24   El Kaoutari, A., Armougom, F., Gordon, J. I., Raoult, D. & Henrissat, B. The
1128   abundance and variety of carbohydrate-active enzymes in the human gut microbiota. *Nat Rev Microbiol* **11**, 497-504 (2013).
1130   https://doi.org:10.1038/nrmicro3050

      25   Martens, E. C. *et al.* Recognition and degradation of plant cell wall
1132   polysaccharides by two human gut symbionts. *PLoS Biol* **9**, e1001221 (2011). https://doi.org:10.1371/journal.pbio.1001221

1134   26   Cuskin, F. *et al.* Human gut Bacteroidetes can utilize yeast mannan through a selfish mechanism. *Nature* **517**, 165-169 (2015).
1136   https://doi.org:10.1038/nature13995

      27   Cartmell, A. *et al.* A surface endogalactanase in Bacteroides thetaiotaomicron
1138   confers keystone status for arabinogalactan degradation. *Nat Microbiol* **3**, 1314-1326 (2018). https://doi.org:10.1038/s41564-018-0258-8

1140   28   Luis, A. S. *et al.* Dietary pectic glycans are degraded by coordinated enzyme pathways in human colonic Bacteroides. *Nat Microbiol* **3**, 210-219 (2018).
1142   https://doi.org:10.1038/s41564-017-0079-1

Alderwick, L. J. *et al.* Arabinan-deficient mutants of Corynebacterium glutamicum and the consequent flux in decaprenylmonophosphoryl-D-arabinose metabolism. *Glycobiology* **16**, 1073-1081 (2006). https://doi.org:10.1093/glycob/cwl030

Jacobson, R. H., Zhang, X. J., DuBose, R. F. & Matthews, B. W. Three-dimensional structure of beta-galactosidase from E. coli. *Nature* **369**, 761-766 (1994). https://doi.org:10.1038/369761a0

De Castro, C., De Castro, O., Molinaro, A. & Parrilli, M. Structural determination of the O-chain polysaccharide from Agrobacterium tumefaciens, strain DSM 30205. *Eur J Biochem* **269**, 2885-2888 (2002). https://doi.org:10.1046/j.1432-1033.2002.02955.x

Li, D. *et al.* 3beta-Hydroxysteroid dehydrogenase expressed by gut microbes degrades testosterone and is linked to depression in males. *Cell Host Microbe* **30**, 329-339 e325 (2022). https://doi.org:10.1016/j.chom.2022.01.001

Treerat, P. *et al.* Synergism between Corynebacterium and Streptococcus sanguinis reveals new interactions between oral commensals. *ISME J* **14**, 1154-1169 (2020). https://doi.org:10.1038/s41396-020-0598-2

Meng, F., Wang, C. & Kurgan, L. fDETECT webserver: fast predictor of propensity for protein production, purification, and crystallization. *BMC Bioinformatics* **18**, 580 (2018). https://doi.org:10.1186/s12859-017-1995-z

Perkowski, E. F. *et al.* The EXIT Strategy: an Approach for Identifying Bacterial Proteins Exported during Host Infection. *mBio* **8** (2017). https://doi.org:10.1128/mBio.00333-17

Marando, V. M. *et al.* Biosynthetic Glycan Labeling. *J Am Chem Soc* **143**, 16337-16342 (2021). https://doi.org:10.1021/jacs.1c07430

Calabretta, P. J., Hodges, H. L., Kraft, M. B., Marando, V. M. & Kiessling, L. L. Bacterial Cell Wall Modification with a Glycolipid Substrate. *J Am Chem Soc* **141**, 9262-9272 (2019). https://doi.org:10.1021/jacs.9b02290

Besra, G. S. *et al.* A new interpretation of the structure of the mycolyl-arabinogalactan complex of Mycobacterium tuberculosis as revealed through characterization of oligoglycosylalditol fragments by fast-atom bombardment mass spectrometry and 1H nuclear magnetic resonance spectroscopy. *Biochemistry* **34**, 4257-4266 (1995). https://doi.org:10.1021/bi00013a015

Chen, C. K., Chan, N. L. & Wang, A. H. The many blades of the beta-propeller proteins: conserved but versatile. *Trends Biochem Sci* **36**, 553-561 (2011). https://doi.org:10.1016/j.tibs.2011.07.004

Shallom, D. *et al.* Detailed kinetic analysis and identification of the nucleophile in alpha-L-arabinofuranosidase from Geobacillus stearothermophilus T-6, a family 51 glycoside hydrolase. *J Biol Chem* **277**, 43667-43673 (2002). https://doi.org:10.1074/jbc.M208285200

Xin, Y., Huang, Y. & McNeil, M. R. The presence of an endogenous endo-D-arabinase in Mycobacterium smegmatis and characterization of its oligoarabinoside product. *Biochim Biophys Acta* **1473**, 267-271 (1999). https://doi.org:10.1016/s0304-4165(99)00204-4

Voisin, S. *et al.* Glycosylation of Pseudomonas aeruginosa strain Pa5196 type IV pilins with mycobacterium-like alpha-1,5-linked d-Araf oligosaccharides. *J Bacteriol* **189**, 151-159 (2007). https://doi.org:10.1128/JB.01224-06

Villmones, H. C. *et al.* Investigating the human jejunal microbiota. *Sci Rep* **12**, 1682 (2022). https://doi.org:10.1038/s41598-022-05723-9

1192   44   Wesener, D. A. *et al.* Recognition of microbial glycans by human intelectin-1. *Nat Struct Mol Biol* **22**, 603-610 (2015). https://doi.org/10.1038/nsmb.3053

1194   45   Krupovic, M., Makarova, K. S. & Koonin, E. V. Cellular homologs of the double jelly-roll major capsid proteins clarify the origins of an ancient virus kingdom. 1196   *Proc Natl Acad Sci U S A* **119** (2022). https://doi.org:10.1073/pnas.2120620119

46   Stewart, G. R., Patel, J., Robertson, B. D., Rae, A. & Young, D. B. 1198   Mycobacterial mutants with defective control of phagosomal acidification. *PLoS Pathog* **1**, 269-278 (2005). https://doi.org:10.1371/journal.ppat.0010033

1200   47   Delafont, V. *et al.* Mycobacterium llatzerense, a waterborne Mycobacterium, that resists phagocytosis by Acanthamoeba castellanii. *Sci Rep* **7**, 46270 1202   (2017). https://doi.org:10.1038/srep46270

48   Antoine, R., Gaudin, C. & Hartkoorn, R. C. Intragenic Distribution of IS6110 in 1204   Clinical Mycobacterium tuberculosis Strains: Bioinformatic Evidence for Gene Disruption Leading to Underdiagnosed Antibiotic Resistance. *Microbiol Spectr* 1206   **9**, e0001921 (2021). https://doi.org:10.1128/Spectrum.00019-21

49   Mueller, E. A., Egan, A. J., Breukink, E., Vollmer, W. & Levin, P. A. Plasticity of 1208   Escherichia coli cell wall metabolism promotes fitness and antibiotic resistance across environmental conditions. *Elife* **8** (2019). 1210   https://doi.org:10.7554/eLife.40754

50   Liao, Y., Smyth, G. K. & Shi, W. The R package Rsubread is easier, faster, 1212   cheaper and better for alignment and quantification of RNA sequencing reads. *Nucleic Acids Res* **47**, e47 (2019). https://doi.org:10.1093/nar/gkz114

1214   51   Zougman, A., Selby, P. J. & Banks, R. E. Suspension trapping (STrap) sample preparation method for bottom-up proteomics analysis. *Proteomics* **14**, 1006- 1216   1000 (2014). https://doi.org:10.1002/pmic.201300553

52   Cox, J. & Mann, M. MaxQuant enables high peptide identification rates, 1218   individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat Biotechnol* **26**, 1367-1372 (2008). 1220   https://doi.org:10.1038/nbt.1511

53   Cox, J. *et al.* Andromeda: a peptide search engine integrated into the MaxQuant 1222   environment. *J Proteome Res* **10**, 1794-1805 (2011). https://doi.org:10.1021/pr101065j

1224   54   Studier, F. W. Protein production by auto-induction in high density shaking cultures. *Protein Expr Purif* **41**, 207-234 (2005). 1226   https://doi.org:10.1016/j.pep.2005.01.016

55   Sauter, N. K. & Zwart, P. H. Autoindexing the diffraction patterns from crystals 1228   with a pseudotranslation. *Acta Crystallogr D Biol Crystallogr* **65**, 553-559 (2009). https://doi.org:10.1107/S0907444909010725

1230   56   Winter, G., Lobley, C. M. & Prince, S. M. Decision making in xia2. *Acta Crystallogr D Biol Crystallogr* **69**, 1260-1273 (2013). 1232   https://doi.org:10.1107/S0907444913015308

57   Evans, P. R. & Murshudov, G. N. How good are my data and what is the 1234   resolution? *Acta Crystallogr D Biol Crystallogr* **69**, 1204-1214 (2013). https://doi.org:10.1107/S0907444913000061

1236   58   McCoy, A. J. *et al.* Phaser crystallographic software. *J Appl Crystallogr* **40**, 658-674 (2007). https://doi.org:10.1107/S0021889807021206

1238   59   Krissinel, E., Uski, V., Lebedev, A., Winn, M. & Ballard, C. Distributed computing for macromolecular crystallography. *Acta Crystallogr D Struct Biol* 1240   **74**, 143-151 (2018). https://doi.org:10.1107/S2059798317014565

Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501 (2010). https://doi.org:10.1107/S0907444910007493

Murshudov, G. N. *et al.* REFMAC5 for the refinement of macromolecular crystal structures. *Acta Crystallogr D Biol Crystallogr* **67**, 355-367 (2011). https://doi.org:10.1107/S0907444911001314

Williams, C. J. *et al.* MolProbity: More and better reference data for improved all-atom structure validation. *Protein Sci* **27**, 293-315 (2018). https://doi.org:10.1002/pro.3330

Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221 (2010). https://doi.org:10.1107/S0907444909052925

Sievers, F. & Higgins, D. G. Clustal Omega, accurate alignment of very large numbers of sequences. *Methods Mol Biol* **1079**, 105-116 (2014). https://doi.org:10.1007/978-1-62703-646-7_6

Gouy, M., Guindon, S. & Gascuel, O. SeaView version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol* **27**, 221-224 (2010). https://doi.org:10.1093/molbev/msp259

Trifinopoulos, J., Nguyen, L. T., von Haeseler, A. & Minh, B. Q. W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res* **44**, W232-235 (2016). https://doi.org:10.1093/nar/gkw256

Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nucleic Acids Res* **49**, W293-W296 (2021). https://doi.org:10.1093/nar/gkab301

UniProt, C. UniProt: a hub for protein information. *Nucleic Acids Res* **43**, D204-212 (2015). https://doi.org:10.1093/nar/gku989

Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679-682 (2022). https://doi.org:10.1038/s41592-022-01488-1

Shenderov, K. *et al.* Cord factor and peptidoglycan recapitulate the Th17-promoting adjuvant activity of mycobacteria through mincle/CARD9 signaling and the inflammasome. *J Immunol* **190**, 5722-5730 (2013). https://doi.org:10.4049/jimmunol.1203343

Moynihan, P. J. & Clarke, A. J. Mechanism of action of peptidoglycan O-acetyltransferase B involves a Ser-His-Asp catalytic triad. *Biochemistry* **53**, 6243-6251 (2014). https://doi.org:10.1021/bi501002d

Gouet, P., Robert, X. & Courcelle, E. ESPript/ENDscript: Extracting and rendering sequence and 3D information from atomic structures of proteins. *Nucleic Acids Res* **31**, 3320-3323 (2003). https://doi.org:10.1093/nar/gkg556

Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Sci* **30**, 70-82 (2021). https://doi.org:10.1002/pro.3943