Extracellular matrix sensing via modulation of orientational order of integrins and F-actin in focal adhesions.

Valeriia Grudtsyna[1,2], Swathi Packirisamy[1], Tamara Bidone[3], Vinay Swaminathan[1,4*]

[1] Division of Oncology, Department of Clinical Sciences, Lund University, Lund, Sweden
[2] Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
[3] Department of Biomedical Engineering, The University of Utah, Salt Lake City, UT, USA
[4] Wallenberg Centre for Molecular Medicine, Lund University, Sweden
[*] Correspondence to Vinay Swaminathan: vinay.swaminathan@med.lu.se

Abstract

Specificity of cellular responses to distinct cues from the extracellular matrix (ECM) requires precise and sensitive decoding of information from the cell surface. However, how known mechanisms of mechanosensing such as force dependent catch bonds and conformational changes in focal adhesion (FA) proteins can confer this sensitivity is not known. Using a combination of polarization microscopy and computational modeling, here we identify regulation of orientational order or molecular co-alignment of FA proteins as a mechanism able to precisely tune cell sensitivity to the ECM. We find that αV integrins and F-actin in FAs show changes in orientational order in an ECM-mediated integrin activation dependent manner. This magnitude of orientational order is sensitive to changes in ECM density but independent of myosin-II activity though actomyosin contractility can further fine-tune it. A molecular clutch model for integrin binding ECM ligands demonstrates that orientational order of integrin-ECM binding and catch bonds tune cellular sensitivity to ECM density. This mechanism is also able to decouple ECM density changes from changes in ECM stiffness thus also contributing to specificity. Taken together, our results suggest relative geometric organization of FA components as an important regulator of mechanotransduction.

Introduction

Cells sense and respond to a wide range of physical cues from the extracellular matrix (ECM) to regulate processes such as cell migration, proliferation, and transcription[1]–[3]. These cues which include the composition and density of ECM proteins and associated mechanical properties such as stiffness, viscoelasticity and architecture are converted into physical signals such as forces and deformation at focal adhesion (FA) sites which are then transmitted and transduced to intracellular protein networks and downstream pathways via the process of mechanotransduction[4]–[6]. Forces and deformation as signals from FAs comprise information about magnitude, direction as well as rates and frequency [7]–[11]. To decipher these signals with high sensitivity and precision, cells need to be able to sense and transmit small changes in the magnitude, direction or loading rates along mechanotransduction pathways while maintaining fidelity of the signal intracellularly.

Over the years, a number of mechanotransduction mechanisms have been discovered, many of which rely on proteins undergoing force-induced changes in their function[12]–[14]. These mechanosensory proteins can modulate their function by exhibiting force-dependent catch bond behavior with interacting partners or by undergoing conformational changes due to application of forces which results in altered protein-protein interactions, modifications or activation[15]–[20]. However, how these mechanisms which primarily rely on forces of different magnitudes acting across the molecule, confer specificity to different physical signals from the ECM is not known. In addition, how sensitivity to different properties of

53 physical signals, such as direction and loading rate is exactly sensed and transduced
54 remains largely unknown.
55
Recent studies have shown that physical coupling between these different mechanisms at
FA sites can indeed bestow sensitivity to changes in ECM stiffness [21]–[23]. Additionally,
some catch-bonds in FAs are sensitive to the direction of force acting across the interaction
surface of the binding partners[24], [25]. These results imply that instead of acting in
isolation, mechanosensors form physical circuits along the force transmission pathway and
the relative organization of these molecules with respect to each other as well as with
respect to the direction of force can result in changes in sensitivity to ECM cues and
downstream cellular response.
64
We previously showed that the primary ECM receptors, integrins can anisotropically
organize and co-align with each other in FAs of cells upon activation by its ECM ligand[26].
This organization which resembles orientational order of nematic materials is oriented along
the long axis of FA which is also the direction of the force vector at the cell-ECM
interface[27], [28]. Additionally, we observed orientational order of F-actin in FAs correlated
with integrin order. Based on the above, we hypothesized that the relative organization of FA
proteins as well as their degree of alignment relative to each other and the force direction i.e.
the orientational order can underlie ECM sensitivity to different physical cues and
downstream cellular response.
74
75 **ECM ligand density tunes cell spread area and YAP localization in MEFs in an**
76 **integrin-dependent manner**.
77
We first set out to establish the relationship between physical cues from the ECM and
cellular response in our mouse embryonic fibroblasts (MEFs) cell line. We focused on ECM
density which allows for precise control of integrin activation, is physiologically relevant and
is conducive to high-resolution total internal reflection fluorescence microscopy (TIRF)[29]–
[32]. We plated MEFs for 4 hours on a series of glass bottom dishes adsorbed with varying
concentrations of fibronectin (0.1 ug/ml, 1ug/ml, 10ug/ml) and poly-l-lysine (PLL)(Fig S1a,b).
Fibronectin (FN) binds and activates the $\beta1$ and $\beta3$ integrin receptors while PLL binds to
cells in an integrin activation independent manner. To measure local as well as distal to FA
cellular response, we focused on the cell spread area, focal adhesion morphology and
nuclear localization of the mechanosensitive transcription factor YAP/TAZ. To directly
compare ECM density sensing with the better characterized stiffness sensing, we also
plated MEFs on polyacrylamide (PA) gels covalently coupled to fibronectin (10 ug/ml) with
Youngs moduli of 0.4, 6 and 60 kPa as previously described[23], [33].
91
Immunofluorescence staining showed that overall cell morphology on different FN
concentrations on glass were clearly distinct, with cells spreading poorly with many small,
dispersed focal adhesions and diffused actin and YAP staining on the lowest fibronectin
concentration and PLL similar to the 0.4KPa stiffness PA gels (Figure 1a, Figure S1c).
Increasing ligand density had a similar effect as increasing stiffness, with cells spreading
more, showing distinct actin stress fibers and focal adhesions and clearer nuclear YAP
signal (Figure 1a, S1c).
99
Quantification of cell spread area based on the actin staining showed a 2 fold increase per
100-fold increase in ligand density while the nuclear to cytoplasmic YAP ratio showed a 1.54
fold increase over these ligand density range (Figure 1b 1c). Quantification of these
parameters on PA gels showed a slightly higher level of sensitivity to ECM stiffness with cell
area increases by 3.63 from 0.4 to 60KPa and the YAP ratio increasing by 1.85(Figure 1d,
1e). Interestingly, quantification of focal adhesion morphology based on paxillin
immunostaining showed small differences in FA size and eccentricity over the range of

ligand densities with only the number of focal adhesions/cells being significant (Figure S1,
S1d. S1e). We verified the generality of these cellular response in a human epithelial cell
line MCF10A as well (Figure S2).
110
111 These results collectively show that similar to response to changes to ECM stiffness, cells
112 also respond with high sensitivity to changes in ECM densities in an integrin dependent
113 manner.
114
115 **Cellular response to changes in ECM density correlates with changes in**
116 **integrin activation dependent orientational order of αV integrins and f- actin in**
117 **FAs.**
118
Having previously identified a strong correlation between orientational order of F-actin and
αV integrins[26], we first asked if F-actin order was sensitive to changes in integrin
activation on FN coated surface. To test this, we plated MEFs untreated or pre-treated with 1
mM Mn^{2+}, a potent enhancer of integrin activation, on 0.1ug/ml FN for 4 hours. Cells were
then fixed and stained with Alexa Fluor 488-phalloidin and paxillin and imaged using the EA-
TIRF system as previously described[26]. Briefly, a polarized excitation is used to illuminate
fluorescent molecules in the sample and the parallel ($I_{\parallel}$) and perpendicular($I_{\perp}$) components
of the emission are separated using a polarized beam splitter. The relative intensities of
these 2 components are then used to calculate fluorescence anisotropy using the formula, r=
($I_{\parallel}-I_{\perp}$)/ ($I_{\parallel}+ 2I_{\perp}$). For calculating orientational order of the fluorescent molecule in FAs, the
average 'r' in an FA is plotted as a function of the orientation of the FA long axis with respect
to the excitation polarization, across many FAs and cells for a given condition. The
amplitude(A) of the modulation in this relationship is dependent on the fraction of molecules
co-aligned and directly relates to the molecular order parameter determined for actin in
drosophila embryos and the cell membrane[34], [35]. The peak of the modulation in this
relationship gives the average orientation of the dipoles of the fluorescent molecules.
135
EA-TIRFM imaging of the cells showed distinct differences between cells pretreated with
Mn^{2+} compared to untreated cells. Mn^{2+} pretreatment increased cell spread area and
resulted in more distinct FAs and actin stress fibers compared to the untreated cells (Figure
2a, Figure S2c, S2d)). The paxillin channel was used to segment FAs and the average
anisotropy 'r' of actin was then calculated per FA and plotted as a function of the FA
orientation. Fitting this modulation to a cos^2 function and extracting the amplitude of
modulation showed that indeed integrin activation with Mn^{2+} increased actin order by a factor
of 1.78 (A=0.119, R^2=0.851; A=0.212, R^2=0.959) (Figure 2b, Figure S3b). Thus, integrin
activation increases the orientational order of actin in FAs and confirms that actin order is
indeed a consequence as well as a sensitive readout of integrin activation in FAs.
146
Next, we asked if ECM ligand density resulted in changes in αV integrin and F-actin order.
To measure integrin anisotropy, we expressed a constrained-GFP αV integrin probe which
rotationally constrains the GFP to allow measurement of integrin order (Figure 2b,S3b,S4a,
S4b). The cells on lower FN concentrations showed similar levels of actin and integrin
anisotropy with a narrow dynamic range in all the FAs while cells on higher FN concentration
had a higher dynamic range of integrin and actin anisotropy (Figure 2b, Figure S3, S4). The
fit of the actin anisotropy vs FA orientation data of cells on 10 ug/ml FN showed a 1.84-fold
change in amplitude compared to 0.1ug/ml FN and 1.16-fold change on 1ug/ml FN (Figure
2b, S3b), while the amplitude of integrins across this ECM range changed by a factor of
2.30(Figure S4c). It is important to note that the order parameter for integrins were
substantially lower compared to F-actin at the same FN densities.
158
159 Thus, taken together, changes in ECM density results in highly sensitive and robust changes
160 in αV integrins and F-actin orientational order which is dependent on integrin activation.

Myosin II activity is not required but increases F-actin order and enhances cellular sensitivity to changes in ECM density.

Within a cell, FAs are highly heterogenous in their size, composition, and function[36]. One of the key regulators of this heterogeneity is the physical coupling between FAs and the actomyosin network which results in differences in tension acting across a FA[37], [38]. During its lifetime, FAs undergo myosin II mediated changes in protein composition, size, and protein organization, all of which are critical determinants of its function[37], [39]–[41]. This suggests that myosin II mediated contractility is a key regulator of FA function. However, it's role in FA-mediated ECM sensing is controversial as studies have shown that ECM sensing is both dependent [27], [39], [40] and independent [42] of myosin II activity. Here, we next wanted to investigate if orientational order and ECM density sensing was mediated by a specific subpopulation of FAs and thus elucidate the role of myosin II in this organization.

As mentioned above, FA size is strongly correlated with myosin II activity with small adhesions decoupled to contractility and larger FAs coupled to it. So, we re-analyzed our F-actin order measurements under different conditions of FN density by binning all the FAs based on 3 sizes (0.1-$0.25\mu m^2$; 0.25-$1\mu m^2$ and 1-$20\mu m^2$) (Figure 3a). Sizes smaller than $0.1\mu m^2$ precluded accurate orientation assignment and were thus not analyzed. We then plotted the average 'r' as a function of FA orientation for the 3 different size bins and fit the modulations to extract F-actin order at each FN density (Figure 3a). Unsurprisingly, this analysis first revealed that the big FAs (1-$20\mu m^2$) showed significant differences in F-actin order between the lowest FN density (A=0.154; R^2=0.768) compared to medium and highest FN density (A=0.25; R^2=0.962) (Figure 3b). However, to our surprise, we also found that the biggest difference in F-actin order at the different ECM densities was in the smallest adhesions (0.1-0.25 μm^2) with an almost 3-fold change between 0.1ug/ml FN and 1 ug/ml FN and a 5-fold difference between 0.1 to 10 ug/ml FN(Figure 3b). In fact, the cos^2 fit on the lowest FN density was very poor (R^2=0.268) leading to an unreliable measure of amplitude and thus suggesting a very disordered actin structure (Figure 3a). Additionally, at the highest FN density, the difference in order of F-actin in the smallest vs. the biggest FAs was very small (1.28-fold change; A=0.196; A=0.25). The middle size bin showed intermediate differences between the small and big FAs with less differences between 0.1 ug/ml FN and 1 ug/ml FN while a 2-fold change between the small and the biggest FAs. Taken together, these results show that that the smallest FAs are the most sensitive to changes in ECM ligand density and that on the highest density, F-actin exhibits orientational order to a similar extent at all FA sizes.

Since FA size correlates with FA maturation state and myosin II activity, our results suggested that myosin II activity was not required for changes in ECM density dependent F-actin order. To test this, we plated MEFs on the 3 FN densities and treated cells with the myosin II inhibitor blebbistatin. Inhibition of myosin II results in loss of all stress fibers and big FAs and leaves behind small NAs. However, since NAs are often harder to accurately segment and assign orientation, we washed out the blebbistatin for 5 minutes and then fixed and stained the cells with AF-488 phalloidin and paxillin to analyze F-actin order. EA-TIRFM of actin and analysis on the remaining small FAs after blebbistatin washout revealed that myosin II inhibition resulted in lower F-actin order at all FN densities compared to the untreated cells from Fig 2 (Figure 3c,3d). However, loss of myosin II activity did not completely abrogate FN density dependent changes in order as there still was a 2-fold increase in F-actin order between the 0.1 ug/ml and 10ug/ml (Figure S3a). Thus, ECM density dependent F-actin order is independent of myosin II activity, though its activity can increase this order.

The above results suggest that cellular response to changes in ECM ligand density is independent of myosin II activity with potentially a role for myosin II in enhancing sensitivity

to these changes. To test this, we plated MEFs on FN coated cover-glass at the 3 densities
and then pretreated the cells with blebbistatin or with DMSO (Figure 4a). We fixed and
stained the cells for either actin to measure cell spread area or with YAP to measure its
nuclear translocation. Quantification and analysis showed an insignificant change in cell
spread area in blebbistatin treated cells compared to the DMSO control over each
concentration of FN (Figure 4b). This led to overall similar sensitivity in cell area across the
different ECM ligand densities. Myosin II inhibition also led to decrease in YAP nuclear
translocation at 0.1ug/ml FN and 10 ug/ml FN, while there was no statistical change at 1
ug/ml FN (Figure 4c). This led to a slight decrease in ECM ligand sensitivity in overall YAP
nuclear translocation with a 1.26-fold change from the lowest to the highest(Figure 4c).
226
227 Taken together, these results suggest that ECM ligand density dependent changes in
228 orientational order of F-actin and cellular response is independent of myosin II activity,
229 though motor activity can increase orientational order and enhance cellular sensitivity.
230
231 **Orientational order of FA components increases sensitivity for ECM ligand**
232 **binding by modulating directional catch bonds.**
233
We next evaluated what mechanisms underlie the changes in orientational order of F-actin
to enable sensitivity to ECM density independent of myosin II activity. To do so, we extended
our computational model of nascent adhesion assembly based on the molecular-clutch
mechanism [43], [44]. Each clutch in the model was represented as an explicit particle
diffusing on the top surface of two parallel surfaces, with the top surface mimicking the
ventral surface of a cell, and the bottom one mimicking the substrate with randomly
distributed ligands (Figure 5a). Each clutch underwent cycles of diffusion, binding and
unbinding of substrate ligands.
242
Since our experiments showed that αV integrins orientational order increases with increase
in ECM density (Figure S3), we incorporated a probability of integrin activation (P_a) that
varies linearly with ECM density, with a minimum of 0.5 at 100 ligands/um^2 and a maximum
of 1 at 200 ligands/um^2 (Figure 5c). When integrin was active, the clutch could bind a free
ligand; when integrin was inactive it could only diffuse. When bound to a ligand, each clutch
was also considered bound to the actin cytoskeleton through binding of vinculin to actin (via
talin). While the mechanism by which orientation order of F-actin is established in FAs is
unknown, a previous study has shown that long range F-actin order can be established by
formation of directional catch bonds between vinculin and F-actin [25]. Having explicitly
measured orientational order of F-actin here (Figure 2), we incorporated in the model
changes in F-actin order by using two pathways for clutch unbinding of actin: directional,
$k_{off,1}$, and non-directional, $k_{off,2}$ (Figure 5a-b). The directional pathway corresponded to the
one with the highest peak in bond lifetime for the vinculin-actin bond at ~13 s (Figure 5b), in
which vinculin forms a bond with actin pulling towards its pointed end. The non-directional
pathway corresponded to the lifetime-force relation with a maximum of ~ 3s (Figure 5b).
To include changes in orientational order of F-actin, we tuned the probability of clutch
unbinding actin according to the directional unbinding pathway ($P_{v,1}$) and then tested its
importance in mediating ligand sensitivity (Figure 5c). To evaluate how different ECM-
dependent integrin anisotropy could affect adhesion assembly, we tuned the degree by
which P_a varies with ligands concentration (Figure 5d) and evaluated its effect on the fraction
of ligated clutches. Last, to understand how the combined effects providing orientational
order of FA clutches, ECM-dependent integrin activation and F-actin orientation, affect
sensing of ECM density, we tested how simultaneously varying $P_{v,1}$, and P_a, affected the
average number of ligated clutches (Figure 5e). Varying the maximum $P_{v,1}$ ($P_{v,1}$
corresponding to n = 200 ligands/um^2) from 0.5 to 1 (Figure 5c) while maintaining P_a = 0.5,
increased the average fraction of ligated clutches from a minimum of ~0.73 to a maximum
~0.93 using 100 to 200 ligands/um^2(Figure 5d). Varying the maximum P_a (P_a corresponding

to n = 200 ligands/um^2) from 0.5 to 1 (Figure 5c) and varying n from 100 to 200 ligands/um^2,
while maintaining $P_{v,1}$ = 0.5, decreased the minimum fraction of ligated clutches to ~0.71 and
increased their maximum to ~ 0.95 (Figure 5e).

Simultaneously varying P_a and $P_{v,1}$ from 0.5-1 (Figure 5c) in the same range of ligand
densities resulted in variations of the fraction of ligated clutches from 0.73, up to ~ 0.97
(Figure 5f). The fraction of ligated clutches increased about 1.6% using n = 200 ligands/um^2
and varying $P_{v,1}$ from 0.5 to 1, while keeping P_a = 0.5 (Figure 5g). The fraction of ligated
clutches increased by about 4% using n = 200 ligands/um^2 and varying P_a from 0.5 to 1,
while keeping $P_{v,1}$ = 0.5 (Figure 5g). By simultaneously increasing both P_a and $P_{v,1}$ this
increase in the fraction of ligated clutches was more than 5%. The percentage of short-lived
clutches was lower using P_a = $P_{v,1}$ = 1 than using either P_a = 0.5 or $P_{v,1}$ = 0.5 (Figure S6A),
meaning that clutches were ligated for longer time when FA order was maximized.
Collectively, these results demonstrate that ligand sensitivity increases the most by
increasing both in the probabilities of integrin activation and directional catch bond. By
contrast, by increasing either parameter in isolation narrows the range of the fraction of
ligated clutches, resulting in less sensitivity to ECM density. Therefore, the results from the
model demonstrate that the magnitude of orientational order of actin filaments determine the
precise sensitivity to ECM density through FA unbinding kinetics in an integrin-activation
dependent way.

We next wondered if ECM density sensing was distinct and independent of ECM stiffness in our
orientational order-based motor clutch model. To do so, we systematically changed the ECM
stiffness from 0.4kPa to 60kPa and plotted the distribution of the fraction of ligated clutches at 3
different ECM ligand densities (100, 110 and 180 ligands/um^2) (Figure 6a). Using 0.4kPa, the
fraction of ligated clutches increased from ~0.6 to ~0.8; using 6 kPa the fraction of ligated
clutches increased from ~0.7 to ~0.95, shifting the range of ligand sensitivity upward; using 60
kPa, the fraction of ligated clutches increased from only ~0.55 to ~0.85, shifting the range of
ligand sensitivity downward. Surprisingly, ECM density sensing was independent of ECM
stiffness in our model as all stiffnesses shown a response to changes in ligand density. At
different stiffnesses, sensitivity to ECM density also increased the most when P_a and $P_{v,1}$ were
increased simultaneously (Figure S6). Interestingly, our model also showed a biphasic response
across different stiffnesses for the same ligand density (Figure 6a). This occurred because when
substrate forces were too high adhesions disassembled instead of stabilizing.

We tested these predictions of our model by plating MEFs on the 3 different FN densities (0.1-10
ug/ml) on PA gels of the 3 stiffnesses used in the model (0.4 KPa, 6 KPa and 60 KPa). We fixed
and immunostained the cells after 4 hours of plating for YAP, actin and nucleus, just as
previously described. Quantification of the cell spread area (Figure 6b) and YAP N/C ratio (Figure
6c) across all these conditions showed a remarkable match with the prediction from the model. At
each ECM rigidity, cell area and nuclear localization of YAP increased with increasing FN density
with different sensitivities. We also observed a biphasic response at the highest ECM density
(10ug/ml) across the 3 rigidities matching the prediction of the model.

Taken together, these results confirm that ECM dependent changes in orientational order of FA
components and regulation of directional catch bonds provide a robust mechanism for fine tuning
the sensitivity of cells to ECM density, independent of ECM rigidity.

Outlook

Integrin-based FAs are the primary multimolecular structures that mediate sensing of a wide
range of physical cues from the ECM. At the molecular scale, FAs perform this function by
assembling a network of several functionally different proteins that are functionally and physically
linked[37], [45], [46]. In addition to composition, FA molecular architecture is also highly complex
and thought to be a critical regulator of its function[47]–[49]. The role of multimolecular

architecture is even more important if one considers the mechanisms of activation of
mechanosensory molecules essential for FA function. The changes in protein function of
mechanosensors are driven by changes in protein structure or conformation upon application of
forces which expose new binding surfaces or cryptic sites that undergo further modifications.
Owing to physical properties of force such as magnitude and direction, the geometry or
orientation of interactions between forces and proteins is an important factor that can influence
the changes in conformation in protein structure and thus its function[50], [51]. In addition to its
effect on single proteins or cluster of proteins, the relative orientational organization of different
proteins, physically connected across the whole FA or FA-nucleus or FA- other cellular structures
can have an impact on the nature of forces felt by distal proteins and organelles[52]. However,
the link between physical cues from the ECM, molecular organization within the FA and cellular
response has so far been missing.
337
Here, we have shown that FA molecules can exhibit orientational order which most importantly,
changes upon small changes in physical cues from the ECM. Our data on F-actin and integrins in
combination with our previously published data shows that FAs have an anisotropic molecular
architecture that is dynamic and originates at the ECM-integrin interface and extends to the actin
cytoskeleton, which is indirectly connected to integrins. We find that the establishment and
regulation of order does not require myosin II contractility suggesting that forces due to actin
polymerization is sufficient and thus not required for mechanosensing, though contractility can
enhance both order and mechanosensitivity. Currently, the specific mechanism that results in F-
actin order in FAs is unknown. A computational model for directionally asymmetric catch bonds
between vinculin and F-actin has previously shown the establishment long range F-actin
order[25]. In our study, we show that the modulation of orientational order measured here when
linked to the directional catch bond as previously shown can lead to highly sensitive
mechanosensing of ECM cues. However, it also likely that other mechanisms can contribute to
establishment and regulation of orientational order, which include other directional catch bonds
as well as actin crosslinking proteins that have shown to promote actin order in focal
adhesions[24], [53], [54]. Interestingly, long range order of actin in cells has been previously
shown to play a significant role in modulating ECM dependent changes in its mechanical
properties[55]. While it's tempting to speculate that the origin of this long range order is at FAs via
the mechanism or orientational order discovered here, this and other regulatory mechanisms will
be the subject of future research.
358
Modulation of orientational order also offers an elegant and energetically less expensive
mechanism for enhancing sensitivity to ECM cues compared to recruitment of new proteins to the
FAs and is likely applicable to other subcellular structures as well where previously, order of
protein components have been observed[56], [57]. In addition, these results further emphasis
unique material properties of FAs which seem to resemble liquid-crystalline materials rather than
disordered liquid-like. This may have implication in the assembly and growth of these structures
in the cell as well as in-vitro reconstitution efforts which utilize phase separation
methodologies[58], [59]. Finally, orientational order of load bearing proteins have been observed
across different cell types, length scales and the animal kingdom[60]–[62]. In combination with
the results of this study, this suggests a potentially conserved mechanism where orientational
order is not just the consequence of directional forces but may also be a mechanism that fine
tunes response to the forces across different cell and tissues in our body.
371
372 **Methods**
373
374 Cell culture and sample preparation
375
Mouse embryonic fibroblasts (MEFs) were cultured in Dulbecco's high glucose modified
eagle medium (Gibco) supplemented with 10% fetal bovine serum (Gibco) and 100 units/ml
Penicillin/Streptomycin (Gibco). Polyacrylamide gels of varying Youngs modulus were

prepared and functionalized following a previously described protocol [63]. Functionalized
polyacrylamide gels or 35 mm 1.5 glass-bottom dishes (Cellvis) were coated with Poly-L-
lysine (PLL) (Sigma-Aldrich), 0.1, 1 and 10 ug/ml fibronectin (Sigma-Aldrich) for 30 min at
37°C and blocked with 2% Bovine Serum Albumin (Sigma-Aldrich) in PBS for 1hr at 37°C or
at 4°C overnight. Cells were plated on FN/PLL coated glass-bottom dishes or gels for 4h
prior to fixation and immunostaining. The relative change in ECM density was verified by
staining for fibronectin (Figure S1). For blebbistatin washout experiments, cells were allowed
to attach for 2h prior to adding 50μM blebbistatin or DMSO which was washed out after 2h
for 5min. For integrin activation experiments, cells were plated in the presence of 1mM Mn^{2+}
for 4h before fixing and staining.
389
390 Immunostaining and transfection
391
Cells were fixed with 4% paraformaldehyde (Thermo Scientific) in cytoskeletal buffer (CB) (2
mM EGTA, 138 mM KCl, 3 mM MgCl, 10 mM Mes, pH 6.1) for 20 min at 37°C and
permeabilized with 0.5% Triton TX-100 (Sigma-Aldrich) in CB for 5 min at RT. They were
then incubated with 0.1M glycine in CB for 10 min and washed with Tris Buffered Saline
buffer (TBS), once for 5 min and twice for 10 min. Next, the cells were blocked with 2% BSA
in 0.1% tween TBS (TBST) for 1h and immunostained with mouse anti-paxillin IgG1 antibody
(BD Transduction Laboratories 610052) (1:500) overnight at 4°C. The cells were washed
with TBST and incubated with secondary goat anti-mouse IgG Alexa 568 (Invitrogen
A11031) (1:400) and Alexa Fluor 488 Phalloidin (Invitrogen) (1:500) for 2h. Samples were
washed and imaged in TBS or mounted on glass slides with ProLong Glass Antifade
Mountant (Invitrogen).
403
For YAP immunostaining and cell area measurements, cells were fixed with 4%
paraformaldehyde in PBS for 15 min at RT, permeabilized with 0.2 % Triton TX-100 in PBS
for 5 min and blocked with 3% BSA in PBS for 45 min. Cells were immunostained with
primary mouse-anti YAP monoclonal IgG2a antibody (Santa Cruz biotechnology sc-101199)
(1:400) in 1% BSA in PBS and incubated at 4°C overnight. Cells were washed with 0.05%
Tween in PBS and incubated for 2h in secondary goat anti-mouse Alexa 568-conjugated IgG
antibody (Life Technologies Corporation), Alexa Fluor 488 Phalloidin (Invitrogen) (1:500) and
nuclear dye Hoechst 350 (33342 Solution, Thermo Scientific) (1:4000). Samples were
washed and imaged in TBS or mounted on glass slides.
To measure anisotropy and orientational order of αV Integrin, αV Integrin-GFP-constrained
plasmid based on previous work was used[26] . Cells were co-transfected using the Neon
Transfection system (Invitrogen MPK5000) with 2500ng of αV Integrin-GFP-constrained
plasmid and untagged β3 integrin plasmid at 1500V, 30ms, 1pulse. Transfected cells were
cultured in antibiotic free media for 48h before plating on FN/PLL coated 35 mm 1.5 glass-
bottom dishes for 4h. The cells were fixed and stained for paxillin.
419
420 EA-TIRFM, image acquisition
421
Images were acquired using total internal reflection fluorescence microscopy (TIRF)
configuration on a Nikon Eclipse Ti microscope with the following available laser lines: 405
nm, 488 nm, 561 nm and 657 nm and Spectra EX (Lumencor). TIRF APO 100x 1.49 N.A.
objective was used for acquiring images. Emission/excitation filters used were: GFP (mirror:
498-537 nm and 565-624nm; excitation: 450-490 nm and 545-555 nm; emission: peak 525
nm, range 30 nm) and mCherry(mirror: 430-470 nm, 501-539 nm and 567-627 nm;
excitation: 395-415 nm, 475-495 nm and 540-560 nm; emission: peak 605nm, range 15 nm)
or Continuous STORM (mirror: 420-481 nm, 497-553 nm, 575-628 nm and 667-792 nm;
excitation: 387-417 nm, 483-494 nm, 557-570 nm and 636-661 nm; emission: 422-478 nm,
502-549 nm, 581-625 nm and 674-786nm).
432

A polarized evanescent TIRF wave excited the fluorophores. In the emission pathway, the
emitted light was split into p and s polarization components with a polarization beam splitter
(Laser Beamsplitter zt 561 sprdc), placed into an Optosplit III. The orthogonal images (Ipa
and Ipe) were projected onto two separate fields of view, manually aligned and captured with
a Teledyne Photometrics 95B 22mm camera. The polarization of the evanescent field was
verified by measuring the extinction coefficient.
439
440 EA-TIRFM image analysis
441
All image analysis were performed using Julia (v.1.6.0) and the code is available on request.
The Ipa and Ipe images were aligned using QuadDIRECT
(https://github.com/timholy/QuadDIRECT.jl). G-factor, $G = \frac{Ipa}{Ipe}$ was calculated in conjunction
with every imaging session to correct for polarization bias in the detection system and the
optical path. A low concentration solution of fluorescein (Sigma-Aldrich) in water was imaged
with the same camera settings, using 488 nm LED for excitation [64]. The resulting Ipa and
Ipe images were used for G-factor computation.
The fluorescence anisotropy formula,

450
$$r = \frac{Ipa - G.\,Ipe}{Ipa + 2.\,G.\,Ipe}$$

451
was applied to the orthogonal F-actin-phalloidin or const. αV integrin images to create an
anisotropy heatmap. FAs were segmented from the background subtracted paxillin image
using the "Moments" binarization algorithm from the ImageBinarization.jl package. The FA
mean anisotropy and the angle between the FA long axis and the excitation polarization (θ),
were extracted using in-house packages in Julia. FAs with eccentricity less than 0.9 (and
less than 0.7 in the Bb washout experiments were filtered out for reliable estimation of FA
long axis orientation.
459
460 Mean anisotropy values (mean intensities from the anisotropy heatmap) from all FAs,
461 collected from all cell images within the same condition, were plotted against the
462 corresponding θ angles. The data was binned into 15-degree bins (12 bins in total) and the
463 mean 'r' together with SD were calculated. The following trigonometric function was fitted to
464 the binned means using the "curve_fit" function from the LsqFit.jl package:
465
466
$$r = A.\,cos^2(\theta + p) + C$$
467 where r is the fluorescence emission anisotropy; amplitude A is directly related to
468 orientational order; θ is the angle between the FA long axis and the excitation polarization; p
469 is the phase shift, which defines the average orientation of dipoles in the ensemble of
470 fluorophores; C is the anisotropy offset due to background. Negative counterclockwise
471 angles were converted to positive angles.
472
473 Confocal microscopy, calculation of YAP N/C ratio and FA analysis.
474
Images were acquired using a Nikon Confocal A1RHD microscope, with 405 nm, 488 nm,
561 nm, and 640 nm laser lines and equipped with 2 GaAsP PMT (for 488 and 561 lines), 2
PMT (for 405 and 640 lines) and 1 GaAsP PMT for spectral detection. Following objectives
were used for imaging: TIRF APO 100x 1.49 N.A. and Plan Apochromal Lambda 60x 1.42
N.A. To capture the entire cell volume z-stacks with 0.6 um step size for cells on glass and
1 um step size for cells on gels were taken (usually 10-20 steps per cell).
481
The maximum projection of the z-stack in YAP channel was background subtracted using
functions "sigma_clip" and "estimate_background" provided by the Photometry.jl. Median
filtering was applied to images taken with 100x objective. The nucleus was segmented from
the summed up nuclear channel z-stack and eroded (with five pixels for the images taken

with 60x objective and 7 pixels for images taken with 100x objective) to represent the area
with nuclear YAP. The nuclear segment was dilated by 55 (for 60x) or 59 (for 100x) pixels
and subtracted with a nuclear segment dilated by 10 (for 60x) or 14 (for 100x) pixels, to
attain a ring-shaped segment around the nucleus. This way, the transition area between the
nuclear and the cytoplasmic region was excluded. The cell was segmented in the actin
channel to represent the cell spread area. The intersection of the ring-shaped segment and
slightly eroded actin segment represented the area with cytoplasmic YAP. For computation
of N/C YAP ratio the median YAP intensity was divided by the median cytoplasmic YAP
intensity. The entire cytoplasmic area was not used to avoid the contribution of low YAP
counts caused by thin cell edges. The Julia code is available on request.
496
497   Paxillin images, taken as previously described in TIRF mode, were submitted for analysis to
498   the online Focal Adhesion Analysis Server (FAAS)[65].
499
500   Molecular clutch model of adhesion assembly
501
To understand how the orientational order of actin filaments could enable ligand sensitivity,
we extended our computational model of nascent adhesion assembly at the leading cell
edge. Like previous approaches from us and others[44], [66]–[68], our model was based on
the molecular-clutch mechanisms, in which adhesion clutches intermittently transmit
cytoskeletal force to the substrate and substrate rigidity to the actin flow. Cytoskeletal force
was produced by intracellular myosin motors regulating the actin flow velocity and the
number of engaged clutches. Substrate force regulated the strength of the clutch-ligand
binding, mimicking integrin-fibronectin bonds in terms of lifetime versus force relations.
Cytoskeletal force regulated the strength of the clutch-actin binding, mimicking vinculin-actin
bonds in terms of directional versus non-directional lifetime versus force.
512
Each clutch in the model was represented as an explicit single point particle in 3D, that
included integrin and vinculins. Once a clutch bound a substrate ligand, it was considered
also bound to the actin cytoskeleton. The lifetime of the fibronectin and actin bound state of
the clutch depended on the kinetics of the bonds of integrin with substrate ligand and of
vinculins with actin. It has been shown that both integrin–fibronectin bond and vinculin-actin
bond are catch-slip bonds, meaning that their lifetime first increases as a function of force,
then decreases as the force increases further[16], [25].
520
Initially, a given number of fibronectin molecules were randomly distributed on a substrate,
to which integrins could bind. The substrate was represented as an isotropic and elastic
material, consisting of a bundle of ideal springs which mimicked ligands.
The clutches underwent cycles of diffusion along the top surface of the model mimicking the
ventral membrane of cells, followed by binding, and unbinding of substrate ligands
underneath it (Figure 5a). As clutches bound substrate ligands, they experienced
cytoskeletal force, from vinculins binding actin undergoing retrograde flow, as was seen in
the lamellipodium[69]. The number of vinculin-actin bonds depended on force and varied
between 2 and 11[70]. When a diffusing clutch came in proximity of a free fibronectin, it
established a harmonic interaction, which mimicked binding, with stiffness determined by
substrate rigidity. When either integrin or at least one vinculin were bound, the clutch was
considered engaged. When the clutch was engaged, it transmitted the actin flow to the
substrate and built tension on the bonds. This tension was used to determine the unbinding
rates of vinculins and integrin. Therefore, these clutches governed local balances between
cytoskeletal contractile force and substrate stiffness. When both integrin and all vinculins in
a clutch were unbound, the clutch was free to diffuse, mimicking the free diffusion of integrin
receptors on the ventral surface of cells[71]. The actin flow was modulated by the number of
clutches engaged with the substrate, and the number of molecular motors in the system.
The force on the integrin-fibronectin bonds was also directly proportional to substrate
stiffness. All parameters in the model were based upon available experimental data and

previous modeling approaches[72]. We directly incorporated the lifetime vs. force
relationships of integrin–fibronectin and vinculin-actin bonds from atomic force microscopy
and optical trap single-molecule experiments[16], [25]. To account for ligand-dependent
integrin activation we varied the probability of integrin activation, P_a. To account for different
degrees of actin filaments orientation we systematically varied the probabilities of directional
catch bond for vinculin, P_{v1}. (Figure 5c). To quantify the amount ligated clutches, we
measured the average fraction of ligated clutches from simulations of 300s.
548
549　Data and code availability
550
551　All raw data that support the findings and the image analysis codes are available from the
552　corresponding authors upon request.
553
554　Acknowledgements
555

We thank P Nordenfelt, Amin Doostmohammadi, Martijn Gloerich, Guillaume Jacquemet,
Sebastian Wasserstrom and all the members of laboratory of cell and molecular
mechanobiology (LCMM) for their discussion and support. Johannes Kumra Ahnlide is
specially acknowledged for developing the initial analysis pipeline and all the help in
developing code and maintaining image storage servers. Lund University Bioimaging Centre
(LBIC) at Lund University is gratefully acknowledged for providing experimental resources.
This research was funded by the Knut and Alice Wallenberg foundation (VS, Wallenberg
Centre for Molecular Medicine, Lund);Cancerfonden (VS, 19 0445 Pj, Projekt grant) and the
National Science Foundation (TCB, NSF BMMB 2044394).

565
566　Author contributions
567
568　VG and VSS conceived the project. VG and SP performed experiments and analysed data.
569　TCB developed the computational model. VG, SP, TCB and VS wrote the manuscript. VSS
570　supervised the project.

595
596

References

[1]   J. D. Humphrey, E. R. Dufresne, and M. A. Schwartz, "Mechanotransduction and extracellular matrix homeostasis," *Nature Reviews Molecular Cell Biology 2014 15:12*, vol. 15, no. 12, pp. 802–812, Oct. 2014, doi: 10.1038/nrm3896.

[2]   D. A. C. Walma and K. M. Yamada, "The extracellular matrix in development," *Development*, vol. 147, no. 10, May 2020, doi: 10.1242/DEV.175596/224217.

[3]   O. Chaudhuri, J. Cooper-White, P. A. Janmey, D. J. Mooney, and V. B. Shenoy, "Effects of extracellular matrix viscoelasticity on cellular behaviour," *Nature 2020 584:7822*, vol. 584, no. 7822, pp. 535–546, Aug. 2020, doi: 10.1038/s41586-020-2612-2.

[4]   Z. Sun, S. S. Guo, and R. Fässler, "Integrin-mediated mechanotransduction," *Journal of Cell Biology*, vol. 215, no. 4, pp. 445–456, Nov. 2016, doi: 10.1083/JCB.201609037.

[5]   F. Martino, A. R. Perestrelo, V. Vinarský, S. Pagliari, and G. Forte, "Cellular mechanotransduction: From tension to function," *Frontiers in Physiology*, vol. 9, no. JUL, p. 824, Jul. 2018, doi: 10.3389/FPHYS.2018.00824/BIBTEX.

[6]   K. A. Jansen, P. Atherton, and C. Ballestrem, "Mechanotransduction at the cell-matrix interface," *Seminars in Cell & Developmental Biology*, vol. 71, pp. 75–83, Nov. 2017, doi: 10.1016/J.SEMCDB.2017.07.027.

[7]   C. C. DuFort, M. J. Paszek, and V. M. Weaver, "Balancing forces: architectural control of mechanotransduction," *Nature Reviews Molecular Cell Biology*, vol. 12, no. 5, pp. 308–319, May 2011, doi: 10.1038/nrm3112.

[8]   A. Ray and P. P. Provenzano, "Aligned forces: Origins and mechanisms of cancer dissemination guided by extracellular matrix architecture," *Current Opinion in Cell Biology*, vol. 72, pp. 63–71, Oct. 2021, doi: 10.1016/J.CEB.2021.05.004.

[9]   E. Abu Shah and K. Keren, "Mechanical forces and feedbacks in cell motility," *Current Opinion in Cell Biology*, vol. 25, no. 5, pp. 550–557, 2013, doi: 10.1016/j.ceb.2013.06.009.

[10]   P. Roca-Cusachs, R. Sunyer, and X. Trepat, "Mechanical guidance of cell migration: Lessons from chemotaxis," *Current Opinion in Cell Biology*, vol. 25, no. 5, pp. 543–549, 2013, doi: 10.1016/j.ceb.2013.04.010.

[11]   A. Elosegui-Artola, X. Trepat, and P. Roca-Cusachs, "Control of Mechanotransduction by Molecular Clutch Dynamics," *Trends in Cell Biology*, vol. 28, no. 5, pp. 356–367, May 2018, doi: 10.1016/J.TCB.2018.01.008.

[12]   B. D. Hoffman, C. Grashoff, and M. A. Schwartz, "Dynamic molecular processes mediate cellular mechanotransduction.," *Nature*, vol. 475, no. 7356, pp. 316–323, 2011, doi: 10.1038/nature10316.

[13]   Z. Sun, S. S. Guo, and R. Fässler, "Integrin-mediated mechanotransduction," *The Journal of Cell Biology*, vol. 215, no. 4, 2016, doi: 10.1083/jcb.201609037.

[14]   V. Swaminathan and M. Gloerich, "Decoding mechanical cues by molecular mechanotransduction," *Current Opinion in Cell Biology*, vol. 72, pp. 72–80, Oct. 2021, doi: 10.1016/J.CEB.2021.05.006.

[15]   W. Chen, J. Lou, E. A. Evans, and C. Zhu, "Observing force-regulated conformational changes and ligand dissociation from a single integrin on cells.," *J Cell Biol*, vol. 199, no. 3, pp. 497–512, 2012, doi: 10.1083/jcb.201201091.

[16] F. Kong, A. J. García, a. P. Mould, M. J. Humphries, and C. Zhu, "Demonstration of catch bonds between an integrin and its ligand," *Journal of Cell Biology*, vol. 185, no. 7, pp. 1275–1284, 2009, doi: 10.1083/jcb.200810002.

[17] B. T. Marshall, M. Long, J. W. Piper, T. Yago, R. P. McEver, and C. Zhu, "Direct observation of catch bonds involving cell-adhesion molecules," *Nature*, vol. 423, no. 6936, pp. 190–193, 2003, doi: 10.1038/nature01605.

[18] A. del Rio, R. Perez-Jimenez, R. Liu, P. Roca-Cusachs, J. M. Fernandez, and M. P. Sheetz, "Stretching single talin rod molecules activates vinculin binding.," *Science*, vol. 323, no. 5914, pp. 638–641, 2009, doi: 10.1126/science.1162912.

[19] A. Kumar *et al.*, "Talin tension sensor reveals novel features of focal adhesion force transmission and mechanosensitivity," *Journal of Cell Biology*, vol. 213, no. 3, pp. 371–383, May 2016, doi: 10.1083/jcb.201510012.

[20] M. Yao, B. T. Goult, H. Chen, P. Cong, M. P. Sheetz, and J. Yan, "Mechanical activation of vinculin binding to talin locks talin in an unfolded conformation.," *Sci Rep*, vol. 4, p. 4610, 2014, doi: 10.1038/srep04610.

[21] N. C. Gauthier and P. Roca-Cusachs, "Mechanosensing at integrin-mediated cell–matrix adhesions: from molecular to integrated mechanisms," *Current Opinion in Cell Biology*, vol. 50, pp. 20–26, Feb. 2018, doi: 10.1016/J.CEB.2017.12.014.

[22] S. W. Moore, P. Roca-Cusachs, and M. P. Sheetz, "Stretchy Proteins on Stretchy Substrates: The Important Elements of Integrin-Mediated Rigidity Sensing," *Developmental Cell*, vol. 19, pp. 194–206, 2010, doi: 10.1016/j.devcel.2010.07.018.

[23] A. Elosegui-Artola *et al.*, "Mechanical regulation of a molecular clutch defines force transmission and transduction in response to matrix rigidity," *Nature Cell Biology*, vol. 18, no. 5, pp. 540–548, May 2016, doi: 10.1038/ncb3336.

[24] L. M. Owen, N. A. Bax, W. I. Weis, and A. R. Dunn, "The C-terminal actin-binding domain of talin forms an asymmetric catch bond with F-actin," *Proc Natl Acad Sci U S A*, vol. 119, no. 10, Mar. 2022, doi: 10.1073/PNAS.2109329119.

[25] D. L. Huang, N. A. Bax, C. D. Buckley, W. I. Weis, and A. R. Dunn, "Vinculin forms a directionally asymmetric catch bond with F-actin," *Science (1979)*, vol. 357, no. 6352, 2017.

[26] V. Swaminathan *et al.*, "Actin retrograde flow actively aligns and orients ligand-engaged integrins in focal adhesions.," 2017. doi: 10.1073/pnas.1701136114.

[27] S. v. Plotnikov, A. M. Pasapera, B. Sabass, and C. M. Waterman, "Force fluctuations within focal adhesions mediate ECM-rigidity sensing to guide directed cell migration," *Cell*, vol. 151, no. 7, pp. 1513–1527, 2012, doi: 10.1016/j.cell.2012.11.034.

[28] M. L. Gardel, B. Sabass, L. Ji, G. Danuser, U. S. Schwarz, and C. M. Waterman, "Traction stress in focal adhesions correlates biphasically with actin retrograde fl ow speed," *Journal of Cell Biology*, vol. 183, no. 6, pp. 999–1005, 2008, doi: 10.1083/jcb.200810060.

[29] J. Galon *et al.*, "Type, density, and location of immune cells within human colorectal tumors predict clinical outcome," *Science (1979)*, vol. 313, no. 5795, pp. 1960–1964, Sep. 2006, doi: 10.1126/SCIENCE.1129139/SUPPL_FILE/GALON.SOM.PDF.

[30] P. P. Provenzano *et al.*, "Collagen density promotes mammary tumor initiation and progression," *BMC Medicine*, vol. 6, no. 1, pp. 1–15, Apr. 2008, doi: 10.1186/1741-7015-6-11/FIGURES/8.

[31] M. Egeblad, M. G. Rasch, and V. M. Weaver, "Dynamic interplay between the collagen scaffold and tumor evolution," *Curr Opin Cell Biol*, vol. 22, no. 5, pp. 697–706, Oct. 2010, doi: 10.1016/J.CEB.2010.08.015.

[32] M. J. Paszek *et al.*, "Tensional homeostasis and the malignant phenotype," *Cancer Cell*, vol. 8, no. 3, pp. 241–254, 2005, doi: 10.1016/j.ccr.2005.08.010.

[33] P. W. Oakes, Y. Beckham, J. Stricker, and M. L. Gardel, "Tension is required but not sufficient for focal adhesion maturation without a stress fiber template," *Journal of Cell Biology*, vol. 196, no. 3, pp. 363–374, 2012, doi: 10.1083/jcb.201107042.

[34] A. Kress *et al.*, "Mapping the Local Organization of Cell Membranes Using Excitation-Polarization-Resolved Confocal Fluorescence Microscopy," *Biophysical Journal*, vol. 105, no. 1, pp. 127–136, Jul. 2013, doi: 10.1016/J.BPJ.2013.05.043.

[35] M. Mavrakis *et al.*, "Septins promote F-actin ring formation by crosslinking actin filaments into curved bundles," *Nature Cell Biology 2014 16:4*, vol. 16, no. 4, pp. 322–334, Mar. 2014, doi: 10.1038/ncb2921.

[36] M. L. Gardel, I. C. Schneider, Y. Aratyn-Schaus, and C. M. Waterman, "Mechanical integration of actin and adhesion dynamics in cell migration.," *Annu Rev Cell Dev Biol*, vol. 26, pp. 315–333, 2010, doi: 10.1146/annurev.cellbio.011209.122036.

[37] J.-C. Kuo, X. Han, C.-T. Hsiao, J. R. Yates, and C. M. Waterman, "Analysis of the myosin-II-responsive focal adhesion proteome reveals a role for β-Pix in negative regulation of focal adhesion maturation.," *Nat Cell Biol*, vol. 13, no. 4, pp. 383–93, 2011, doi: 10.1038/ncb2216.

[38] G. Giannone *et al.*, "Lamellipodial Actin Mechanically Links Myosin Activity with Adhesion-Site Formation," *Cell*, vol. 128, no. 3, pp. 561–575, 2007, doi: 10.1016/j.cell.2006.12.039.

[39] Cl. M. Pasapera AM, Schneider, IC, Rericha E, Schlaepfer DD, Waterman, A. M. Pasapera, I. C. Schneider, E. Rericha, D. D. Schlaepfer, and C. M. Waterman, "Myosin II activity regulates vinculin recruitment to focal adhesions through FAK-mediated paxillin phosphorylation," *Journal of Cell Biology*, vol. 188, no. 6, pp. 877–890, 2010, doi: 10.1083/jcb.200906012.

[40] A. M. Pasapera, S. v Plotnikov, R. S. Fischer, L. B. Case, T. T. Egelhoff, and C. M. Waterman, "Rac1-dependent phosphorylation and focal adhesion recruitment of myosin IIA regulates migration and mechanosensing.," *Curr Biol*, vol. 25, no. 2, pp. 175–86, Jan. 2015, doi: 10.1016/j.cub.2014.11.043.

[41] L. B. Case *et al.*, "Molecular mechanism of vinculin activation and nanoscale spatial organization in focal adhesions," *Nature Cell Biology*, pp. 1–10, 2015, doi: 10.1038/ncb3180.

[42] P. W. Oakes *et al.*, "Lamellipodium is a myosin-independent mechanosensor," *Proc Natl Acad Sci U S A*, vol. 115, no. 11, pp. 2646–2651, 2018, doi: 10.1073/PNAS.1715869115/SUPPL_FILE/PNAS.1715869115.SM03.MP4.

[43] T. C. Bidone, A. v. Skeeters, P. W. Oakes, and G. A. Voth, "Multiscale model of integrin adhesion assembly," *PLOS Computational Biology*, vol. 15, no. 6, p. e1007077, Jun. 2019, doi: 10.1371/JOURNAL.PCBI.1007077.

[44] P. W. Oakes *et al.*, "Lamellipodium is a myosin-independent mechanosensor," *Proc Natl Acad Sci U S A*, vol. 115, no. 11, pp. 2646–2651, 2018, doi: 10.1073/PNAS.1715869115/SUPPL_FILE/PNAS.1715869115.SM03.MP4.

[45] E. R. Horton *et al.*, "Definition of a consensus integrin adhesome and its dynamics during adhesion complex assembly and disassembly," *Nature Cell Biology*, 2015, doi: 10.1038/ncb3257.

[46] H. B. Schiller and R. Fässler, "Mechanosensitivity and compositional dynamics of cell-matrix adhesions.," *EMBO Rep*, vol. 14, no. 6, pp. 509–19, 2013, doi: 10.1038/embor.2013.49.

[47] I. Patla *et al.*, "Dissecting the molecular architecture of integrin adhesion sites by cryo-electron tomography," *Nature Cell Biology 2010 12:9*, vol. 12, no. 9, pp. 909–915, Aug. 2010, doi: 10.1038/ncb2095.

[48]   H. van Hoorn *et al.*, "The nanoscale architecture of force-bearing focal adhesions," *Nano Letters*, vol. 14, no. 8, pp. 4257–4262, Aug. 2014, doi: 10.1021/NL5008773/SUPPL_FILE/NL5008773_SI_005.AVI.

[49]   P. Kanchanawong *et al.*, "Nanoscale architecture of integrin-based cell adhesions.," *Nature*, vol. 468, no. 7323, pp. 580–584, 2010, doi: 10.1038/nature09621.

[50]   S. Chakrabarti, M. Hinczewski, and D. Thirumalai, "Plasticity of hydrogen bond networks regulates mechanochemistry of cell adhesion complexes," *Proc Natl Acad Sci U S A*, vol. 111, no. 25, pp. 9048–9053, Jun. 2014, doi: 10.1073/PNAS.1405384111.

[51]   S. Adhikari, J. Moran, C. Weddle, and M. Hinczewski, "Unraveling the mechanism of the cadherin-catenin-actin catch bond," *PLOS Computational Biology*, vol. 14, no. 8, p. e1006399, Aug. 2018, doi: 10.1371/JOURNAL.PCBI.1006399.

[52]   A. T. Blanchard and K. Salaita, "Multivalent molecular tension probes as anisotropic mechanosensors: concept and simulation," *Physical Biology*, vol. 18, no. 3, p. 034001, Mar. 2021, doi: 10.1088/1478-3975/ABD333.

[53]   M. A. Juanes, D. Isnardon, A. Badache, S. Brasselet, M. Mavrakis, and B. L. Goode, "The role of APC-mediated actin assembly in microtubule capture and focal adhesion turnover," *Journal of Cell Biology*, vol. 218, no. 10, pp. 3415–3435, Oct. 2019, doi: 10.1083/JCB.201904165/VIDEO-4.

[54]   O. Loison *et al.*, "Polarization-resolved microscopy reveals a muscle myosin motor-independent mechanism of molecular actin ordering during sarcomere maturation," *PLOS Biology*, vol. 16, no. 4, p. e2004718, Apr. 2018, doi: 10.1371/JOURNAL.PBIO.2004718.

[55]   M. Gupta *et al.*, "Adaptive rheology and ordering of cell cytoskeleton govern matrix rigidity sensing," *Nature Communications*, vol. 6, no. 1, p. 7525, Dec. 2015, doi: 10.1038/ncomms8525.

[56]   A. L. Mattheyses, M. Kampmann, C. E. Atkinson, and S. M. Simon, "Fluorescence anisotropy reveals order and disorder of protein domains in the nuclear pore complex," *Biophysical Journal*, vol. 99, no. 6, pp. 1706–1717, Sep. 2010, doi: 10.1016/j.bpj.2010.06.075.

[57]   M. Kampmann, C. E. Atkinson, A. L. Mattheyses, and S. M. Simon, "Mapping the orientation of nuclear pore proteins in living cells with polarized fluorescence microscopy," *Nat Struct Mol Biol*, vol. 18, no. 6, pp. 643–649, 2011, doi: 10.1038/nsmb.2056.

[58]   L. B. Case, M. de Pasquale, L. Henry, and M. K. Rosen, "Synergistic phase separation of two pathways promotes integrin clustering and nascent adhesion formation," *Elife*, vol. 11, Jan. 2022, doi: 10.7554/ELIFE.72588.

[59]   S. F. Banani *et al.*, "Compositional Control of Phase-Separated Cellular Bodies," *Cell*, vol. 166, no. 3, pp. 651–663, Jul. 2016, doi: 10.1016/J.CELL.2016.06.010.

[60]   M. Cetera *et al.*, "Epithelial rotation promotes the global alignment of contractile actin bundles during Drosophila egg chamber elongation," *Nature Communications 2014 5:1*, vol. 5, no. 1, pp. 1–12, Nov. 2014, doi: 10.1038/ncomms6511.

[61]   Y. Maroudas-Sacks, L. Garion, L. Shani-Zerbib, A. Livshits, E. Braun, and K. Keren, "Topological defects in the nematic order of actin fibres as organization centres of Hydra morphogenesis," *Nature Physics 2020 17:2*, vol. 17, no. 2, pp. 251–259, Nov. 2020, doi: 10.1038/s41567-020-01083-1.

[62]   D. Volfson, S. Cookson, J. Hasty, and L. S. Tsimring, "Biomechanical ordering of dense cell populations," *Proc Natl Acad Sci U S A*, vol. 105, no. 40, pp. 15346–15351, Oct. 2008, doi: 10.1073/PNAS.0706805105/SUPPL_FILE/0706805105SI.PDF.

[63] J. N. Lakins, A. R. Chin, and V. M. Weaver, "Exploring the Link Between Human Embryonic Stem Cell Organization and Fate Using Tension-Calibrated Extracellular Matrix Functionalized Polyacrylamide Gels," *Methods in Molecular Biology*, vol. 916, pp. 317–350, 2012, doi: 10.1007/978-1-61779-980-8_24.

[64] S. Ghosh, S. Saha, D. Goswami, S. Bilgrami, and S. Mayor, "Dynamic Imaging of Homo-FRET in Live Cells by Fluorescence Anisotropy Microscopy," 2012, doi: 10.1016/B978-0-12-388448-0.00024-3.

[65] M. E. Berginski, S. M. Gomez, J. Jones, and K. Yamada, "The Focal Adhesion Analysis Server: a web tool for analyzing focal adhesion dynamics," *F1000Research 2013 2:68*, vol. 2, p. 68, Mar. 2013, doi: 10.12688/f1000research.2-68.v1.

[66] A. Macdonald, A. R. Horwitz, and D. A. Lauffenburger, "Kinetic model for lamellipodal actin-integrin 'clutch' dynamics.," *Cell Adh Migr*, vol. 2, no. 2, pp. 95–105, 2008, doi: 10.4161/CAM.2.2.6210/SUPPL_FILE/KCAM_A_10906210_SM0001.ZIP.

[67] C. E. Chan and D. J. Odde, "Traction dynamics of filopodia on compliant substrates," *Science (1979)*, vol. 322, no. 5908, pp. 1687–1691, Dec. 2008, doi: 10.1126/science.1163595.

[68] A. Elosegui-Artola *et al.*, "Rigidity sensing and adaptation through regulation of integrin types," *Nature Materials 2014 13:6*, vol. 13, no. 6, pp. 631–637, May 2014, doi: 10.1038/nmat3960.

[69] A. Ponti, M. Machacek, S. L. Gupton, C. M. Waterman-Storer, and G. Danuser, "Two distinct actin networks drive the protrusion of migrating cells.," *Science*, vol. 305, no. 5691, pp. 1782–1786, 2004, doi: 10.1126/science.1100533.

[70] M. Yao *et al.*, "The mechanical response of talin," *Nature Communications 2016 7:1*, vol. 7, no. 1, pp. 1–11, Jul. 2016, doi: 10.1038/ncomms11966.

[71] O. Rossier *et al.*, "Integrins β1 and β3 exhibit distinct dynamic nanoscale organizations inside focal adhesions," *Nature Cell Biology*, vol. 14, no. 10, pp. 1057–1067, 2012, doi: 10.1038/ncb2588.

[72] A. Y. Alexandrova *et al.*, "Comparative dynamics of retrograde actin flow and focal adhesions: Formation of nascent adhesions triggers transition from fast to slow flow," *PLoS ONE*, vol. 3, no. 9, 2008, doi: 10.1371/journal.pone.0003234.

Figure 1

a)

Figure 1

ECM ligand density tunes cell spread area and YAP localization in MEFs in an Integrin-dependent manner: (a, top) Representative images of MEFs on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml) and PLL, fixed and stained with Hoechst to label nucleus, phalloidin 488 to label actin and Alexa 568 to label YAP. (a, bottom) Representative images of MEFs on gels of different Youngs modulus (0.4, 6, and 60kPa) coated with 10ug/ml of FN, fixed and stained with Hoechst to label nucleus, phalloidin 488 to label actin and Alexa 568 to label YAP. (b) Box plot quantification of cell area from analysis of immunofluorescence images of cells plated on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml) and PLL. Cell area was obtained by segmenting the actin channel. N = 36 for each condition. (c) Box plot quantification of nuclear/cytoplasmic ratio of YAP from analysis of immunofluorescence images of cells plated on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml). N = 36 for each condition. (d) Box plot quantification of cell area from analysis of immunofluorescence images of cells plated on gels of varying Youngs modulus (0.4, 6, and 60kPa), coated with 10ug/ml of FN. Cell area was obtained by segmenting the actin channel. N = 12 for each condition (e) Box plot quantification of nuclear/cytoplasmic ratio of YAP from analysis of immunofluorescence images of cells plated on gels of varying Youngs modulus (0.4, 6, and 60kPa), coated with 10ug/ml of FN. N = 12 for each condition

Figure 2

Figure 2

Cellular response to changes in ECM ligand density correlates with finely tuned changes in integrin activation and co-alignment of actin in FAs: (a) Representative images of MEFs on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml) and on 0.1 ug/ml FN in the presence of Mn^{2+}. Cells were fixed and stained with phalloidin 488 and imaged with TIRF. Emission from the parallel (Ipa) channel, perpendicular (Ipe) channel and a merge (left, Ipa green, Ipe magenta) are shown. Paxillin stained with Alexa 568 (middle). Emission anisotropy (r) of actin in the whole cell and emission anisotropy of actin in segmented FAs (Right). Magnitude of anisotropy color scale (bottom) (b) mean actin anisotropy (r) in FAs vs FA orientation fit to the $\cos^2$ function $r = C + A\cdot\cos^2(\gamma + \theta_d)$ for cells plated on each FN condition. Error bars represent SD. Adhesions from 12 cells were analysed.

Figure 3

Figure 3

Myosin-2 activity is not required for but enhances ECM density dependent F-actin co-alignment and cellular response: (a) mean actin anisotropy (r) in FAs vs FA orientation fit to the $\cos_2$ function r = C + A·$\cos_2$(γ + θ_d) binned by FA size (0.1-0.25μm^2; 0.25-1μm^2 and 1-20μm^2) in cells plated on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml). (b) Graph showing amplitudes of actin anisotropy (r) binned by FA size from three experiments. (c) Representative images of MEFs on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml) following blebbistatin washout for 5 minutes. Cells were fixed and stained with phalloidin 488 and imaged with TIRF. Emission from the parallel (Ipa) channel, perpendicular (Ipe) channel are shown (left) . Paxillin stained with Alexa 568 (middle right). Emission anisotropy (r) of actin stained with phalloidin 488 in the whole cell (right). Magnitude of anisotropy color scale (bottom) (d) mean actin anisotropy (r) in FAs vs FA orientation fit to the $\cos^2$ function for cells plated on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml) with 5 minutes blebbistatin washout. Adhesions from 15, 12 and 10 cells were analysed for the conditions 0.1 ug/ml, 1ug/ml, 10ug/ml respectively.

Figure 4

Figure 4

F-actin anisotropy in FAs is myosin 2 independent (a) Representative images of MEFs on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml) following blebbistatin treatment, fixed and stained with Hoechst to label nucleus, phalloidin 488 to label actin and Alexa 568 to label YAP. (b) Box plot quantification of cell area from analysis of immunofluorescence images of cells plated on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml) following blebbistatin treatment. Cell area was obtained by segmenting the actin channel. N = 36 for each condition. (c) Box plot quantification of nuclear/cytoplasmic ratio of YAP from analysis of immunofluorescence images of cells plated on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml) following blebbistatin treatment. N = 36 for each condition

Figure 5

Figure 5

Ligand sensitivity is governed by molecular order of FA clutches. (a) Schematic representation of the computational model of FA assembly based on the molecular clutch mechanism. Two 2D surfaces are placed 20 nm apart. The bottom surface represents the substrate with a random distribution of ligands, modeled as elastic springs with stiffness k_{sub}, which depends on the substrate young modulus, Y. The top surface mimics the ventral cell membrane, with integrins diffusing with diffusion coefficient D and establishing interactions with substrate ligands. When integrin binds a substrate ligand, filaments exert a force on the clutch with magnitude that depends on their retrograde velocity and myosin motors activity. The direction of this force determines the clutch unbinding rate, $k_{off,1}$ (directional) or $k_{off,2}$ (non-directional). (b) Unbinding rates follow the lifetime ($1/k_{off}$) versus force relation, where the directional pathway of vinculins in the clutch is indicated with v,1, and the non-directional pathway is indicated with v,2. (c) Linear relations between the probability of $k_{off,1}$ (directional pathway for unbinding) $P_{v,1}$, and integrin activation rate, P_a, with varying ligand density, n, between 100-200 ligands/um^2. Legends indicate the value of $P_{v,1}$ or P_a using $n = 200$. The following relations are used: for $P_{v,1}$ or $P_a = 0.5$ at $n = 200$, $P_{v,1}$ or $P_a = 0.0$ x n +0.5; for $P_{v,1}$ or $P_a = 0.6$ at $n = 200$, $P_{v,1}$ or $P_a = 0.01$ x n +0.4; for $P_{v,1}$ or $P_a = 0.7$ at $n = 200$, $P_{v,1}$ or $P_a = 0.02$ x n +0.3; for $P_{v,1}$ or $P_a = 0.8$ at $n = 200$, $P_{v,1}$ or $P_a = 0.03$ x n +0.2; for $P_{v,1}$ or $P_a = 0.9$ at $n = 200$, $P_{v,1}$ or $P_a = 0.04$ x n +0.1; for $P_{v,1}$ or $P_a = 1$ at $n = 200$, $P_{v,1}$ or $P_a = 0.05$ x n; (d) Average fraction of ligated clutches varying $P_{v,1}$ between 0.5-1 and n between 100-200 ligands/um^2, while keeping $P_a = 0.5$. (e) Average fraction of ligated clutches varying P_a between 0.5-1 and n between 100-200 ligands/um^2, while keeping $P_{v,1} = 0.5$. (f) Average fraction of ligated clutches by simultaneously varying $P_{v,1}$ and P_a between 0.5-1 and varying n between 100-200 ligands/um^2. (g) Average fraction of ligand-bound integrins for different values of $P_{v,1}$, P_a or both, varying n between 100-200 ligands/um^2. All data are obtained as averages from 300 s of simulations for each condition, using $Y = 6$ KPa.

Figure 6

Figure 6

ECM ligand sensitivity is ECM stiffness independent: (a) Boxplots of the fraction of ligand bound integrins varying n between 100-180 ligands/um^2 and using Y= 0.4, 6, and 60 kPa. All data are obtained from 300 s of simulations sampling every 1 s. (b) Box plot quantification of cell area from analysis of immunofluorescence images of cells plated on gels of varying Youngs modulus (0.4, 6, and 60kPa) coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml). Cell area was obtained by segmenting the actin channel. N = 12 for each condition. (c) Box plot quantification of nuclear/cytoplasmic ratio of YAP from analysis of immunofluorescence images of cells plated on gels of varying Youngs modulus (0.4, 6, and 60kPa) coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml). N = 12 for each condition

Figure S1

Figure S1

(a) Representative images of glass bottom dishes coated with different concentrations of fibronectin (0.1 ug/ml, 1ug/ml, 10ug/ml) (b) box plot of mean fibronectin intensities at the three different concentrations. (c) Representative images of MEFs plated on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml), fixed, and stained with phalloidin 488 to label actin and Alexa 568 to label paxillin. Violin plots of FA area (d) FA eccentricity (e) and box plot of number of FA per cell (f) in MEFs plated on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml). Quantifications were from analysis of immunofluorescence images of paxillin. N = 10 for each condition.

Figure S2

Figure S2

(a) Representative images of MCF10a cells plated on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml), fixed and stained with Hoechst to stain nucleus, phalloidin 488 to label actin and Alexa 568 to label YAP. (b) Box plot quantification of cell area of MCF10a cells from analysis of immunofluorescence images of cells plated on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml) and PLL. Cell area was obtained by segmenting the actin channel. N = 13 for each condition. (c) Box plot quantification of nuclear/cytoplasmic ratio of YAP of MCF10a cells from analysis of immunofluorescence images of cells plated on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml) and PLL. N = 13 for each condition.

Figure S3

(a) Representative images of MEFs plated on glass coated with PLL. Immunofluorescence images of phalloidin 488 actin with emission from the parallel (Ipa) and perpendicular (Ipe) channels and merge are shown. Paxillin stained with Alexa 568 (middle). Emission anisotropy (r) of actin stained with phalloidin 488 in the whole cell is shown (right). (b) Graph showing amplitudes of actin anisotropy (r) in cells plated on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml) and in the presence of Mn^{2+} on 0.1 ug/ml of FN from 3 experiments. (c) Box plot quantification of cell area from analysis of immunofluorescence images of cells plated on glass coated with 0.1 ug/ml of FN and in presence or absence of Mn^{2+}. Cell area was obtained by segmenting the actin channel. N = 36 for each condition. (d) Violin plot quantification of nuclear/cytoplasmic ratio of YAP from analysis of immunofluorescence images of cells plated on glass coated with 0.1 ug/ml of FN and in presence or absence of Mn^{2+}. N = 12 for each condition.

Figure S4

(a) Representative images of aV-integrin-GFP transfected MEFs plated on glass coated with PLL and different FN concentrations (0.1ug/ml, 1ug/ml 10ug/ml) imaged with TIRF. Emission from the parallel (Ipa) and perpendicular (Ipe) channels and merge are shown (left Ipa magenta, Ipe green). Paxillin stained with Alexa 568 (middle). Emission anisotropy (r) of actin stained with aV-integrin-GFP in the whole cell and in the FAs are shown (right). Magnitude of anisotropy color scale (bottom) (b) mean integrin anisotropy (r) in FAs vs FA orientation fit to the cos2 function r = C + A·cos2(γ + θd) for cells plated on each PLL/FN condition. Error bars represent SD. N = 15 for each experiment. (c) graph of integrin anisotropy amplitudes of cells on different FN concentrations (0.1ug/ml, 1ug/ml 10ug/ml) from two experiments. PLL experiments were done once. N = 19, 80, 35, 26 for the conditions PLL, 0.1ug/ml FN, 1ug/mlFN, 10ug/ml FN respectively.

a)

Figure S5

(a) Graph showing amplitudes of actin anisotropy (r) of cells plated on glass coated with different FN concentrations (0.1 ug/ml, 1ug/ml, 10ug/ml) following blebbistatin washout. N = 15 for each condition per experiment.

Figure S6

(a) Average percentage of short lived clutches (with lifetime < 0.9 s) relative to the total amount of ligated clutches for different conditions of P_a and $P_{v,1}$ using 100 to 200 ligands/um² and $Y = 6$ KPa or (b) $Y = 1$ KPa. (c-e) Average fraction of ligated clutches, using $Y = 1$ KPa and varying n between 100-200 ligands/um² and either (c) varying $P_{v,1}$ between 0.5-1 while keeping $P_a = 0.5$; (d) varying P_a between 0.5-1 while keeping $P_{v,1} = 0.5$; (e) simultaneously varying $P_{v,1}$ and P_a between 0.5-1. All data are obtained as averages from 300 s of simulations for each condition

Supplementary Note on Computational model

We used a Brownian Dynamics approach to simulate the binding and unbinding of molecular adhesion clutches in response to differences in actin filaments orientations, ligand density and substrate stiffness. Actin filaments were considered implicitly in the model, by modulating the amount of force they exerted on the ligand-bound clutches depending on the number of engaged clutches and molecular motors. Therefore, force on the clutches depended on the actin flow rate, which in turn depended on the number of ligand-bound clutches, substrate rigidity and the number of motors. Unbinding of the clutches could follow two pathways: the direction and the non-directional bond kinetics of vinculin. To simulate different degrees of orientation of the actin filaments, we modulated the probability of establishing a directional versus a non-directional catch bond for the interaction of the clutch with actin.

Computational Domain and Boundary Conditions.

Like our previous implementation of the adhesion assembly model[1], [2], the computational domain was 3D and consisted of two parallel surfaces of 1 µm side, separated in the vertical direction by L=20 nm, a dimension typical of integrin headpiece extension[3]. The bottom surface represented the substrate with immobilized ligands; the top substrate represented the ventral cell membrane with the clutches on it (Figure 5b). The clutches were initially randomly distributed on the top surface, and they diffused along x and y, with diffusion coefficient of integrin, D = 0.29 µm^2/s[4]. To avoid finite size effects on the motion of the free clutches, we used periodic boundary conditions in the lateral directions of the domain.

Substrate Model.

The substrate was considered an elastic solid, consisting of a randomly distributed ideal linear springs, mimicking individual fibronectin molecules, with stiffness depending on substrate rigidity, as:

$$k_{sub} = \frac{YA}{L}$$

where Y is the Young's modulus (we tested values in the range 0.4– 60 kPa), A is the integrin/ligand cross-sectional area (corresponding to 80 nm^2, from an ideal bar of radius ~5 nm, corresponding to approximately half the value of the transmembrane leg separation of an integrin in the open conformation), and L is the equilibrium distance separation between substrate and top layer, corresponding to the equilibrium separation between clutch and bound ligand.
Hooke's law for each spring in the bundle can be written as:

$$\boldsymbol{F_{sub}} = k_{sub}\Delta\boldsymbol{L}$$

where $\Delta\boldsymbol{L}$ is the variation from the equilibrium separation between clutch and bound ligand. We used ligand densities between 100-200 ligands/µm^2.

Actin flow.

Actin was considered implicitly, as a force acting on each ligated clutch, as a function of the flow velocity. It was calculated as:

$$F_{flow} = \zeta_i v_{flow}$$

The actin retrograde flow velocity (v_{flow}) was calculated through the linear force-velocity relationship[5], [6],:

$$v_{flow} = v_u(1 - \frac{x_c k_{sub}}{n_m F_m})$$

where v_u = 0.11 µm /s is the unloaded velocity, x_c is the number of bound clutches, n_m=135 is the number of motors, and F_m=2 pN is the motor stall force.

Clutch Representation

Each *i*-th clutch and *j*-th ligand was defined by a 3D position vectors, r_i and r_j, respectively. The vector r_i presented *x, y,* and *z* coordinates of the *i*-th clutch; the vector r_j presented *x, y* and *z* coordinates of the *j*-th ligand. At every timestep of the simulations, *x* and *y* of r_i were updated to track the clutch displacement on the top surface, and binding/unbinding of substrate ligands, while *z* remained fixed at 0; *x, y,* and *z* of r_j remained all fixed over the course of the simulations because ligands were immobilized, with *z* = -0.02 µm.

When a free, diffusive clutch came in proximity of a free ligand (< 21 nm from it), it bound the ligand by establishing a harmonic interaction. This interaction presented spring constant k_{sub} (proportional to the substrate Youngs' modulus, *Y*), and equilibrium distance *L*, corresponding to the separation of the open extracellular integrin headpiece from the membrane [3]. In the ligated state, a number of vinculin-actin bonds were also formed. This number dependent on the force on the integrin-fibronectin bond: 2 for forces below 8pN; 5 for forces between 8 and 15 pN; 9 for forces between 15 and 21 pN; 11 for forces > 21 pN [7]. The harmonic interactions between clutches and ligands, and between vinculins and actin provided several mechanical links between actin and the substrate. Each of these links lasted for a certain lifetime, that depended on two types of catch-slip bonds (integrin-ligand and vinculin-actin bonds). Each clutch became unligated if integrin and all vinculins failed. Once the clutches unbound their ligands, they became diffusive again until they bound a new ligand.

Implementation Algorithm.

Recognizing that inertia is negligible on the length and time scales of integrin motion in the plasma membrane, the displacement of each *i*-th integrin was governed by the Langevin equation of motion in the limit of high friction[8]

$$\frac{dr_i}{dt} = \frac{F_i}{\zeta_i}$$

where r_i was the position vector of integrin; ζ_i was integrin friction coefficient calculated using Einstein relation, as $\zeta_i = \frac{k_B T}{D}$ = 0.0142 pN s/µm, where $k_B T$ = 4.11 pN nm and D = 0.29 µm²/s; *dt* = 10^{-4} s was the simulation timestep; F_i was the total force on the clutch, including a stochastic

contribution from thermal effects and a deterministic contribution from actin flow, governed by amount of ligated clutches, number of motors and their stall force, and substrate mechanics.

Considering all forces acting on the clutches at every time step, their positions were updated iteratively using an explicit Euler integration scheme:

$$r_i(t + dt) = r_i(t) + \frac{dr_i}{dt}dt = r_i(t) + \frac{F_i}{\zeta_i}dt$$

Forces acting on the clutches

The total force acting on each i-th clutch, F_i, resulted from the sum of a stochastic and a deterministic contribution. A stochastic force, F_T, was applied to all clutches in the membrane at every dt, to mimic thermal effects generating diffusion. This force satisfied the fluctuation-dissipation theorem. Deterministic forces originated from actin flow, F_{flow}, and substrate tension, F_{sub}. Thus, the total force acting on each i-th clutch was calculated as:

$$F_i = F_T + F_{flow} + F_{sub}$$

Lifetime of clutch engagements depends on force

The clutch/substrate interaction persisted for a characteristic lifetime which depended upon the tension on the bond. We assumed that integrin and vinculin feel the effect of this force equally on the clutch bond and that the force increases the dissociation rate constants ($k_{off,integrin}$, $k_{off,+}$, $k_{off,-}$) as originally proposed by Bell [9].

Two characteristic catch bonds were considered: integrin-fibronectin and vinculin-actin. The clutch/substrate interaction broke when both bonds were broken (for the case of vinculin, all vinculin bonds had to break for the clutch to become unligated).

A double exponential pathway determined unbinding rates of integrin-fibronectin and of each vinculin-actin. It included a strengthening pathway, with a negative exponent, and a weakening pathway, with a positive exponent.

For integrin, the unbinding rate was:

$$k_{off} = 2 * e^{-0.0640*F} + 0.00005 * e^{0.26*F}$$

For each vinculin, the unbinding rate depended on filaments orientation and therefore it depended on the direction of force application. For forces applied toward the filament pointed end, the catch bond was directional, with the longest lifetimes and corresponding unbinding rate:

$$k_{off,-} = 2 * e^{-0.046*F} + 0.00005 * e^{0.78*F}$$

For forces applied toward the filament barbed end, the catch bond was non-directional, and the unbinding rate was:

$$k_{off,+} = 4 * e^{-0.28*F} + 0.00005 * e^{0.95*F}$$

References

[1]   T. C. Bidone, A. v. Skeeters, P. W. Oakes, and G. A. Voth, "Multiscale model of integrin adhesion assembly," *PLOS Computational Biology*, vol. 15, no. 6, p. e1007077, Jun. 2019, doi: 10.1371/JOURNAL.PCBI.1007077.

[2]   P. W. Oakes *et al.*, "Lamellipodium is a myosin-independent mechanosensor," *Proc Natl Acad Sci U S A*, vol. 115, no. 11, pp. 2646–2651, 2018, doi: 10.1073/PNAS.1715869115/SUPPL_FILE/PNAS.1715869115.SM03.MP4.

[3]   X. P. Xu, E. Kim, M. Swift, J. W. Smith, N. Volkmann, and D. Hanein, "Three-Dimensional Structures of Full-Length, Membrane-Embedded Human α(IIb)β(3) Integrin Complexes," *Biophys J*, vol. 110, no. 4, pp. 798–809, Feb. 2016, doi: 10.1016/J.BPJ.2016.01.016.

[4]   O. Rossier *et al.*, "Integrins β1 and β3 exhibit distinct dynamic nanoscale organizations inside focal adhesions," *Nature Cell Biology*, vol. 14, no. 10, pp. 1057–1067, 2012, doi: 10.1038/ncb2588.

[5]   B. L. Bangasser and D. J. Odde, "Master equation-based analysis of a motor-clutch model for cell traction force," *Cellular and Molecular Bioengineering*, vol. 6, no. 4, pp. 449–459, Dec. 2013, doi: 10.1007/s12195-013-0296-5.

[6]   C. E. Chan and D. J. Odde, "Traction dynamics of filopodia on compliant substrates," *Science (1979)*, vol. 322, no. 5908, pp. 1687–1691, Dec. 2008, doi: 10.1126/science.1163595.

[7]   M. Yao *et al.*, "The mechanical response of talin," *Nature Communications 2016 7:1*, vol. 7, no. 1, pp. 1–11, Jul. 2016, doi: 10.1038/ncomms11966.

[8]   D. Feldman, "The theory of polymer dynamics, by M. Doi and S. F. Edwards, the Clarendon Press, Oxford University Press, New York, 1986, 391 pp. Price: $78.50.," *Journal of Polymer Science Part C: Polymer Letters*, vol. 27, no. 7, pp. 239–240, Jun. 1989, doi: 10.1002/POL.1989.140270706.

[9]   George I. Bell, "Models for the Specific Adhesion of Cells to Cells," *Science (1979)*, vol. 200, no. 4342, pp. 618–627, 1978, doi: 10.1126/science.347575.